

AT LEAST ONE-IN-SIX GALAXIES IS ALWAYS DEAD

L.E. Abramson^{1,*} and D.D. Kelson¹

¹*Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA 91101, USA*

ABSTRACT

Via numerical experiments, we show that the $\sim 10\%$ – 20% passive fraction seen at $z \gtrsim 3$ is consistent with galaxy star formation histories being maximally correlated stochastic processes. If so, this fraction should reflect a time-independent baseline that holds at any epoch or mass regime where mean star formation rates are rising. Data at $\log M_* \leq 10$ and $z \lesssim 0.5$ bear this out, as should future *James Webb Space Telescope* observations.

Keywords: galaxy evolution — quenched galaxies — star formation

Galaxy star formation histories (SFHs) can be modeled as stochastic processes (Kelson 2014; Kelson et al. 2016; Kelson et al. 2020). In this framework, star formation rates (SFRs) are non-negative sums of random numbers. The SFR at the epoch of observation, T , can be independent of values at earlier times, t , or correlated with them, on average, via a weighting factor $(T - t)^{H-1/2}$. Here, H defines the strength of the correlation (Mandelbrot & Van Ness 1968). If $H = 1/2$, the SFH returns to a purely random Brownian walk. If $1/2 < H \leq 1$, past SFR changes bias future changes in the same direction.

Ensembles of stochastic SFHs describe coeval galaxy populations. While the star formation rate of any individual member cannot be predicted, the ensemble’s mean and dispersion evolve deterministically in time as functions of H (Kelson 2014):

$$\langle \text{SFR}(T) \rangle \propto \frac{T^H}{H}, \quad (1)$$

$$\text{Sig}[\text{SFR}(T)] = \sqrt{H} \langle \text{SFR}(T) \rangle. \quad (2)$$

These equations have been shown to hold at redshifts preceding the peak of cosmic star formation (Kelson et al. 2016) and in mass regimes where sSFRs ($\equiv \text{SFR}/M_*$) are mass-independent (Kelson 2014).

While an ensemble’s mean SFR always rises in time, its dispersion grows more slowly for all $H < 1$. So, at fixed epoch, more objects will reach $\text{SFR}(T) = 0$ if SFHs are maximally correlated ($H = 1$) rather than random ($H = 1/2$). Intuitively, this trend reflects the fact that random walks spend less time at $\text{SFR} = 0$ than ones in which past declines bias future changes toward zero. Physically, it means that the passive galaxy fraction constrains SFH stochasticity.

Figure 1, left, illustrates the above trend in 20 numerical ensembles of 1000 SFHs spanning 1000 timesteps. At $H = 1/2$, only 2% of objects have $\text{SFR} = 0$ in the last timestep. By $H = 1$, 18% do. The overplotted measurements of the passive fraction at $z \gtrsim 3$ and $\log M_* \lesssim 10$ at $z \lesssim 0.5$ —where the stochastic framework most directly applies—clearly favor $H \gtrsim 0.8$.

McLeod et al. (2020)’s $\sim 9\%$ implies the lowest H , but simulations show that these authors’ *UVJ* passive selection—based on Williams et al. (2009)’s $z < 0.5$ prescription—may miss 45% of intrinsically non-starforming galaxies at $z \sim 3$.² If so, those data imply a true passive fraction of $\sim 16\%$, suggesting $H \simeq 1$. Such selection effects worsen with increasing redshift: by $z \sim 4$, only about 1/3 of passive galaxies are accurately identified. If so, Santini et al. (2020)’s $z \gtrsim 4$ estimate of a $\sim 7\%$ passive fraction ($\log M_* \geq 10.5$) is consistent with Marsan et al. (2020)’s $z > 3$ measurement of 15%–25% ($\log M_* \geq 11$) and also suggests $H \simeq 1$,³ as the lower-redshift data clearly prefer (Muzzin et al. 2013; Moustakas et al. 2013; Mao et al. 2020).

Measuring the passive fraction is difficult, but we are confident in the above statements because results based on the SFR–stellar mass relation (Kelson 2014), stellar mass functions (Kelson et al. 2016), and the growth of cosmic structure (Kelson et al. 2020) all independently imply that $H \simeq 1$. Intriguingly, so does an analytical derivation based on Newtonian gravitational collapse and

Corresponding author: L.E. Abramson
labramson@carnegiescience.org

* Carnegie Fellow

² Tests assumed a Gaussian $A_V \leq 2$ mag distribution with a mean and dispersion of 0.5 mag, $\log Z/Z_\odot = -0.5$, a Chabrier (2003) initial mass function, and no photometric errors. Using the relevant observed bands yields better results at $z = 3$ (*F140W*, *K_s*, *Spitzer IRAC2*).

³ Technically, the spectral templates Santini et al. (2020) use to identify passive galaxies come from a library based on a non-evolved *UVJ* cut such that 1/2 to 2/3 of truly passive spectral energy distributions may go unmatched.

mass conservation (Kelson et al. 2020). $H \simeq 1$ also fits with our intuition that galaxies’ observed properties should meaningfully reflect their pasts (Caplar & Tacchella 2019; Iyer et al. 2020).

Indeed, the fact that the same passive fraction describes epochs as disparate as those spanned by the above datasets suggests $H \simeq 1$ on its own. The portion of tracks near the minimum of a stochastic process (SFR = 0) is analytically known to evolve as T^{H-1} (Molchan 2000). This trend—confirmed numerically in Figure 1, right—is incompatible with all observations, confining H near unity a priori. If so, the passive fraction should be time-independent as is observed.

Additional processes separate what we have calculated—the fraction of galaxies expected to have zero star formation given normal fluctuations in hospitable conditions—from the “quenched” fraction measured at higher masses or later times (e.g., Peng et al. 2010). The framework discussed here does not yet capture those forces, so our results should be interpreted as a floor or null expectation for the passive fraction in complete ensembles.

As such, we predict that future unbiased JWST surveys will find 16% to 20% of galaxies to be classifiable as passive at all $z \gtrsim 3$, and also at lower redshifts in mass regimes where sSFRs are mass-independent. We look forward to forthcoming tests!




Figure 1. *Left:* data at $z > 3$ and $\log M_* < 10$ at lower redshifts directly suggest that galaxy SFHs are consistent with $H \gtrsim 0.8$ stochastic processes. Accounting for color selection effects, all data suggest $H \gtrsim 0.9$, or maximal temporal correlation. The data describe regimes in which fuel- (high-redshift) or quenching-related SFR constraints (low-redshift) are absent. *Right:* if $H \simeq 1$ —as the SFR– M_* relation and stellar mass function shape also imply—then a null passive fraction of 16%–20% should be constant in time (uppermost long dashes). At all $H < 1$, the passive fraction declines as T^{H-1} contrary to observations. We start the plot at $t/T = 0.25$ as numerical convergence issues affect the early timesteps as $H \rightarrow 1$. Our $H_{\text{input}} = 1$ may thus reflect $H_{\text{eff}} \simeq 0.92$, but the implied passive fraction evolves so slowly—staying above 15% after 4,000 timesteps—that the results at *left* should capture the true null passive fraction. (See also grey line denoting fractions at $T/2$ at *left*.)

REFERENCES

- Caplar, N., & Tacchella, S. 2019, MNRAS, 487, 3845
- Chabrier, G. 2003, PASP, 115, 763
- Iyer, K. G., Tacchella, S., Genel, S., et al. 2020, MNRAS, 498, 430
- Kelson, D. D. 2014, ArXiv e-prints, arXiv:1406.5191
- Kelson, D. D., Benson, A. J., & Abramson, L. E. 2016, ArXiv e-prints, arXiv:1610.06566
- Kelson, D. D., Abramson, L. E., Benson, A. J., et al. 2020, MNRAS, 494, 2628
- Mandelbrot, B. B., & Van Ness, J. W. 1968, SIAM Review, 10, 422
- Mao, Y.-Y., Geha, M., Wechsler, R. H., et al. 2020, arXiv:2008.12783
- Marsan, Z. C., Muzzin, A., Marchesini, D., et al. 2020, arXiv:2010.04725
- McLeod, D. J., McLure, R. J., Dunlop, J. S., et al. 2020, arXiv:2009.03176
- Molchan, G. M. 2000, Theory Probab. Appl., 44(1), 97–102.
- Moustakas, J., Coil, A. L., Aird, J., et al. 2013, ApJ, 767, 50
- Muzzin, A., Marchesini, D., Stefanon, M., et al. 2013, ApJ, 777, 18
- Peng, Y.-jie ., Lilly, S. J., Kovač, K., et al. 2010, ApJ, 721, 193
- Santini, P., Castellano, M., Merlin, E., et al. 2020, arXiv:2011.10584
- Williams, R. J., Quadri, R. F., Franx, M., et al. 2009, ApJ, 691, 1879