

On the minimal sum of edges in a signed edge-dominated graph

Danila Cherkashin^{a,b,c}, Pavel Prozorov^c

a. Chebyshev Laboratory, St. Petersburg State University, 14th Line V.O., 29B, Saint Petersburg 199178 Russia

b. Moscow Institute of Physics and Technology, Laboratory of Combinatorial and Geometric Structures

c. School 533, St. Petersburg, Russia

Abstract

Let G be a simple graph with n vertices and ± 1 -weights on edges. Suppose that for every edge e the sum of edges adjacent to e (including e itself) is positive. Then the sum of weights over edges of G is at least $-\frac{n^2}{25}$. Also we provide an example of a weighted graph with described properties and the sum of weights $-(1 + o(1))\frac{n^2}{8(1+\sqrt{2})^2}$.

The previous best known bounds were $-\frac{n^2}{16}$ and $-(1 + o(1))\frac{n^2}{54}$ respectively. We show that the constant $-1/54$ is optimal under some additional conditions.

10 May 2021

arXiv:2012.09956v2 [math.CO]

1 Introduction

A graph (finite, simple, undirected) is a pair (V, E) , where V stands for a set of vertices, and E denotes a set of unordered pairs of vertices, elements of E are called edges. Let G be a graph; for a given edge $e = (u, v)$ define its *closed edge-neighborhood* as an edge subset $N[e]$ formed by e and all adjacent to e edges of G . A weight function $f : E \rightarrow \{+1, -1\}$ is called a *signed edge domination function* of G if

$$\sum_{e' \in N[e]} f(e') \geq 1$$

for every $e \in E$; in this case we say that (G, f) is a *SED-pair* of order $|V|$. Let $s[(G, f)]$ be the sum of weights over all edges of a graph G equipped by a weight function f .

Denote by E_+ the set $\{(u, v) \in E \mid f(u, v) = 1\}$ and by E_- the set $\{(u, v) \in E \mid f(u, v) = -1\}$. Define

$$s_v = \sum_{e \in N(v)} f(e);$$

for each $v \in V$, where $N(v)$ stands for the set of edges containing v . Let V_+ be $\{v \in V \mid s_v \geq 0\}$ and V_- be $\{v \in V \mid s_v < 0\}$.

The following problem was posed by Xu in [5, 6].

Problem 1. *What is*

$$g(n) := \min\{s[(G, f)] \mid (G, f) \text{ is a SED-pair of order } n\}$$

for each positive integer n ?

Note that for every $g(n) \leq 0$ since an empty graph provides a SED-pair. The only known result was provided by the following theorem.

Theorem 1 (Akbari – Bolouki – Hatami – Siami [2]). *(i) For every n*

$$g(n) \geq -\frac{n^2}{16}.$$

(ii) There is a sequence of SED-pairs of order n that satisfies¹

$$s[G, f] \leq -(1 + o(1))\frac{n^2}{54}.$$

We refine both items as follows.

Theorem 2. *(i) For every n*

$$g(n) \geq -\frac{n^2}{25}.$$

¹In fact the authors claim the bound $-\frac{n^2}{72}$ but the provided example gives the bound $-(1 + o(1))\frac{n^2}{54}$.

(ii) For every n there is a SED-pair of order n that satisfies

$$s[G, f] < -(1 + o(1)) \frac{n^2}{8(1 + \sqrt{2})^2}.$$

Moreover, if $n = 4(p + q)p$, where $p > 1$ and $q > 1$ are positive integers satisfying $p^2 = 2q^2 - 1$, then

$$s[G, f] = \left[-\frac{n^2}{8(1 + \sqrt{2})^2} + \frac{3\sqrt{2} - 4}{4}n \right].$$

Note that there are infinitely many p and q satisfying the condition $p^2 = 2q^2 - 1$, since it is a special case of Pell's equation; it is well known that the positive solutions are

$$p = \frac{\sqrt{2} - 1}{2}(3 + 2\sqrt{2})^k - \frac{1 + \sqrt{2}}{2}(3 - 2\sqrt{2})^k, \quad q = \frac{\sqrt{2} - 1}{2\sqrt{2}}(3 + 2\sqrt{2})^k + \frac{1 + \sqrt{2}}{2\sqrt{2}}(3 - 2\sqrt{2})^k, \quad k \in \mathbb{N}.$$

We show that Theorem 1(ii) is optimal under additional assumptions.

Theorem 3. Let (G, f) be a SED-pair of order n . Suppose that every $e \in E_-$ connects a vertex from V_+ with a vertex from V_- ; and every $e \in E_+$ connects some vertices from V_+ . Then

$$s[(G, f)] \geq -\frac{1}{54}n^2.$$

1.1 Graphons

A *graphon* (also known as a graph limit) is a symmetric measurable function $W : [0, 1]^2 \rightarrow [0, 1]$. Define a *signed graphon* as a symmetric measurable function $W : [0, 1]^2 \rightarrow [-1, 1]$. A signed graphon is *edge-dominated* if $W(x, y) \neq 0$ implies

$$\int_0^1 (W(x, t) + W(y, t))dt \geq 0.$$

Here we consider a continuous analogue of Problem 1. Denote

$$\kappa := \inf \frac{1}{2} \int_0^1 \int_0^1 W(x, y) dx dy \tag{1}$$

where the infimum is taken over all edge-dominated graphons W .

The following theorem is a standard result in the theory of graph limits [3], we include the proof in Appendix A for completeness.

Theorem 4. (i) $g(n) \geq \kappa n^2$, in other words $s(G, f) \geq \kappa n^2$ for any SED-pair (G, f) of order n ;
(ii) $g(n) = (\kappa + o(1))n^2$ for large n .

Theorems 2 and 3 also have natural continuous analogues.

Structure of the paper. Theorem 2(ii) is proved in Section 2. Section 3 is devoted to the proof of Theorem 2(i). Section 4 cites a result, determining the maximal sum of squares of vertex degrees among all graphs with n vertices and e edges; we use it in Section 5, containing the proof of Theorem 3. Appendix A contains the proof of Theorem 4, Appendices B-D contain auxiliary calculations.

2 Examples

In this section we provide a sequence of SED-pairs that achieves the lower bound $-(1 + o(1)) \frac{n^2}{8(1 + \sqrt{2})^2}$.

2.1 A graphon example

The following signed graphon realizes an example for Theorem 2(ii). Put $[0, 1] = A \sqcup B \sqcup C$, where $|A| = 1 - \frac{1}{\sqrt{2}}$, $|B| = \frac{1}{\sqrt{2}} - \frac{1}{2}$, $|C| = \frac{1}{2}$. The function W is defined in Fig. 1. Note that W is edge-dominated: indeed, for $(x, y) \in A \times A$

$$\int_0^1 (W(x, t) + W(t, y))dt = 2 \left(-\frac{1}{\sqrt{2}}|A| + |B| \right) = 0,$$

for $(x, y) \in A \times B$

$$\int_0^1 (W(x, t) + W(t, y)) dt = -\frac{1}{\sqrt{2}}|A| + |B| + |A| + |B| - \frac{1}{\sqrt{2}}|C| = \frac{1}{2} - \frac{1}{2\sqrt{2}} > 0,$$

for $(x, y) \in B \times B$

$$\int_0^1 (W(x, t) + W(t, y)) dt = 2 \left(|A| + |B| - \frac{1}{\sqrt{2}}|C| \right) = 1 - \frac{1}{\sqrt{2}} > 0,$$

and for $(x, y) \in B \times C$

$$\int_0^1 (W(x, t) + W(t, y)) dt = 2 \left(|A| + |B| - \frac{1}{\sqrt{2}}|C| - \frac{1}{\sqrt{2}}|B| \right) = 0.$$

Finally,

$$\frac{1}{2} \int_0^1 \int_0^1 W(x, y) dx dy = \frac{1}{2} \left(-\frac{|A|^2}{\sqrt{2}} + 2|A| \cdot |B| + |B|^2 - \frac{2|B| \cdot |C|}{\sqrt{2}} \right) = \frac{1}{8(1 + \sqrt{2})^2}.$$

1				1
C	0	$-\frac{1}{\sqrt{2}}$	0	
B	1	1	$-\frac{1}{\sqrt{2}}$	
A	$-\frac{1}{\sqrt{2}}$	1	0	
0	A	B	C	1

Figure 1: A graphon example for Theorem 2(ii)

2.2 An explicit graph approximation

Here we provide the best approximation we can do. Fix p and q such that $p^2 = 2q^2 - 1$, and $p, q > 1$.

We need several auxiliary definitions. Define graph $K_{X, Y, \frac{k}{l}} = (X \cup Y, E_{X, Y, \frac{k}{l}})$ for $|X| = al, |Y| = bl$ and integer $a, b, k \leq l$. Split X onto a disjoint sets of size l : $X = X_1 \cup X_2 \cup \dots \cup X_a, |X_i| = l$; also split Y onto the b disjoint sets of the same size: $Y = Y_1 \cup Y_2 \cup \dots \cup Y_b, |Y_j| = l$. For each pair $1 \leq i \leq a, 1 \leq j \leq b$ consider the following bipartite graph $G_{ij} = (X_i \cup Y_j, E_{ij})$ with parts X_i and Y_j (all graphs G_{ij} are isomorphic). Enumerate vertices as follows $X_i = \{v_1, v_2, \dots, v_l\}, Y_j = \{u_1, u_2, \dots, u_l\}$. Define E_{ij} as the set of all pairs (v_g, u_h) , for which $g - h \pmod l$ lies in $\{1, 2, \dots, k\}$. Put

$$E_{X, Y, \frac{k}{l}} = \bigcup_{1 \leq i \leq a, 1 \leq j \leq b} E_{ij}.$$

Obviously the degree of every vertex in G_{ij} equals to k , so the degree of a vertex in $K_{X, Y, \frac{k}{l}}$ is $bk = |Y| \frac{k}{l}$ for vertices in X , and $ak = |X| \frac{k}{l}$ for vertices in Y .

Now define graph $K_X = (X, E_X)$ for $|X| = 2al$ and integer $a, k < l$. Split X onto $2l$ disjoint sets of size a : $X = X_1 \cup X_2 \cup \dots \cup X_{2l}$. The edge between vertices u and v exists if and only if $i - j \pmod{2l}$ lies in

$$\{-k, -(k-1), \dots, -2, -1, 1, 2, \dots, k-1, k\},$$

where $v \in X_i, u \in X_j$. Then the degree of every vertex in K_X equals to $2ak = |X| \frac{k}{l}$.

Let $K_X = (X, E_X)$ be the complete graph (i.e. every pair of vertices forms an edge) on the vertex set X . Degree of each vertex in K_X equals to $|X| - 1$.

Now we are ready to provide the desired construction. Let p and q be a positive solution of $p^2 = 2q^2 - 1$. Put

$$A = \{a_1, a_2, \dots, a_{2p^2}\}, \quad B_1 = \{b_1, b_2, \dots, b_{2p(p-q)}\}, \quad B_2 = \{b_{2p(p-q)+1}, b_{2p(p-q)+2}, \dots, b_{2pq}\},$$

$$C_1 = \{c_1, c_2, \dots, c_{6p(p-q)}\}, \quad C_2 = \{c_{6p(p-q)+1}, c_{6p(p-q)+2}, \dots, c_{2(p+q)p}\}.$$

Define the vertex set

$$V = A \cup B_1 \cup B_2 \cup C_1 \cup C_2$$

(so $n = 4p^2 + 4pq$). The edge set E and weight function f are defined by explicit expressions for E_+ and E_- :

$$E_+ = E_{A, B_1 \cup B_2, \frac{1}{2}} \cup E_{B_1, \frac{p^2 - pq - 1}{p(p-q)}} \cup E_{B_1, B_2, \frac{1}{2}} \cup E_{B_2}; \quad E_- = E_{A, \frac{q}{p}} \cup E_{B_1, C_2, \frac{q}{p}} \cup E_{B_2, C_1, \frac{q}{p}} \cup E_{B_1, C_1, \frac{2pq - 2q^2 - 1}{2p(p-q)}} \cup E_{B_2, C_2, \frac{4q^2 - 2pq - 1}{2p(2q-p)}}.$$

Since p divides all of the cardinalities $|A|, |B_1|, |B_2|, |C_1|, |C_2|$; $2p(p-q)$ divides $|B_1|, |C_1|$, and $2p(2q-p)$ divides $|B_2|, |C_2|$, the definition of f is correct.

Some annoying calculation gives

$$s_{a_i} = 0, \quad s_{b_i} = p^2, \quad s_{c_i} = -p^2$$

for every i .

Note that there is no edge between A and C or inside C . Also all edges inside A of between B and C are negative, so our construction is a SED-pair.

Finally we count

$$s[G, f] = \frac{1}{2} \sum_{v \in V} s_v = \frac{p^2(|B_1| + |B_2| - |C_1| - |C_2|)}{2} = -p^4.$$

Recall that $p^2 = 2q^2 - 1$ and $n = 4p^2 + 4pq = 2p(2p + \sqrt{2}\sqrt{1+p^2})$. So

$$\frac{s[G, f]}{n^2} = \frac{-p^4}{(2p(2p + \sqrt{2}\sqrt{1+p^2}))^2} = -\frac{1}{8(1 + \sqrt{2})^2} + \frac{5\sqrt{2} - 7}{8p^2} + \frac{31\sqrt{2} - 44}{32p^4} + O(p^{-5}).$$

Since $n = (4 + 2\sqrt{2})p^2 + \sqrt{2} - \frac{1}{2\sqrt{2}p^2} + O(p^{-3})$

$$s[G, f] = -\frac{n^2}{8(1 + \sqrt{2})^2} + \frac{3\sqrt{2} - 4}{4}n - \frac{1}{2(2 + \sqrt{2})} + o(1).$$

One can also derive

$$s[G, f] = \left[-\frac{n^2}{8(1 + \sqrt{2})^2} + \frac{3\sqrt{2} - 4}{4}n \right].$$

3 Lower bound on $-n^2/25$

Consider an arbitrary SED-pair (G, f) , $G = (V, E)$.

It is known that for each $v, u \in V$ if $(v, u) \in E_- \cup E_+$, then $s_v + s_u \geq 0$ (check it by hands or see Lemma 1 in [2]). If V_- is empty, then $s[G, f] \geq 0$. Let x be

$$-\min_{v \in V_-} s_v$$

and consider an arbitrary vertex a such that $s_a = -x$. Let $N_-(a)$ be $\{v \in V \mid (a, v) \in E_-\}$. Then $|N_-(a)| \geq x$ and $s_v \geq x$ for each $v \in N_-(a)$, so $N_-(a) \subset V_+$. Then

$$x^2 \leq \sum_{v \in N_-(a)} s_v \leq \sum_{v \in V_+} s_v.$$

Clearly, V_- is an independent set (i.e. has no edges inside) so

$$\begin{aligned} \sum_{v \in V_+} s_v &= \sum_{v \in V_-} s_v + 2 \left(\sum_{(u,v) \in E_+ \mid u,v \in V_+} 1 - \sum_{(u,v) \in E_- \mid u,v \in V_+} 1 \right) \\ &\leq \sum_{v \in V_-} s_v + 2 \frac{|V_+| \cdot (|V_+| - 1)}{2} \leq \sum_{v \in V_-} s_v + |V_+|^2. \end{aligned}$$

So

$$\sum_{v \in V_-} s_v \geq x^2 - |V_+|^2;$$

recall that

$$\sum_{v \in V_+} s_v \geq x|N_-(a)| \geq x^2.$$

On the other hand

$$s[(G, f)] = \sum_{(x,y) \in E_+} 1 - \sum_{(x,y) \in E_-} 1 = \frac{\sum_{v \in V} s_v}{2},$$

and

$$\sum_{v \in V} s_v = \sum_{v \in V_+} s_v + \sum_{v \in V_-} s_v \geq 2x^2 - |V_+|^2.$$

Also

$$\sum_{v \in V} s_v = \sum_{v \in V_+} s_v + \sum_{v \in V_-} s_v \geq x^2 - x|V_-| = -x(|V_-| - x) = -x(|V| - |V_+| - x).$$

Put $y = \frac{x}{|V|}$, $k = \frac{|V_+|}{|V|}$. Then we have the following system of inequalities:

$$\begin{cases} s[(G, f)] \geq (y^2 - \frac{k^2}{2})|V|^2 \\ s[(G, f)] \geq \frac{-y(1-k-y)}{2}|V|^2. \end{cases}$$

So

$$g(n) \geq \min_{0 \leq y \leq 1, 0 \leq k \leq 1} \left(\max \left(y^2 - \frac{k^2}{2}, -\frac{y(1-k-y)}{2} \right) \right) n^2.$$

One may check by computer (or read explicit calculus in Appendix B) that the minimum is $-\frac{1}{25}$ and is reached at $y = \frac{1}{5}$, $k = \frac{2}{5}$.

4 Degree sequences of a graph

Here we display the results from [1], they are required in the proof of Theorem 3; for a survey see [4].

Definition. Let $n, e \leq \binom{n}{2}$ be integer numbers. Consider the unique representation

$$e = \binom{a}{2} + b, \quad 0 \leq b < a.$$

The quasi-complete graph C_n^e with e edges and n vertices v_1, \dots, v_n has edges (v_i, v_j) for $i, j \leq a$ and $i = a+1, j \in \{1 \dots b\}$.

Definition. Let $n, e \leq \binom{n}{2}$ be integer numbers. Consider the unique representation

$$\binom{n}{2} - e = \binom{c}{2} + d, \quad 0 \leq d < c.$$

The quasi-star graph S_n^e with e edges and n vertices v_1, \dots, v_n and the edges connect vertices v_1, \dots, v_{n-c-1} with all vertices and vertex v_{n-c} is connected with vertices v_1, \dots, v_{n-d} .

Let $F(n, e)$ be the maximal value of

$$\sum_{v \in V} (\deg v)^2$$

among the graphs $G = (V, E)$ with n vertices and e edges. We use the following result.

Theorem 5 (Alshwede – Katona, [1]). *For every n and $0 \leq e \leq \binom{n}{2}$ the value $F(n, e)$ is achieved on C_n^e or S_n^e .*

Corollary 1. *Put $\alpha = \frac{2e}{n^2}$. Then*

$$F(n, e) = (1 + o(1)) \max \left(\alpha^{\frac{3}{2}}, (1 - \sqrt{1 - \alpha})(\sqrt{1 - \alpha} + \alpha) \right) n^3.$$

Define

$$G(\alpha) := \alpha^{\frac{3}{2}}, \quad H(\alpha) = (1 - \sqrt{1 - \alpha})(\sqrt{1 - \alpha} + \alpha).$$

We show that $G(\alpha) < H(\alpha)$ for $\alpha \in (0, 1/2)$ and $G(\alpha) > H(\alpha)$ for $\alpha \in (1/2, 1)$. Define $t = \sqrt{1 - \alpha}$. Note that

$$h^2(\alpha) - g^2(\alpha) = (1 - t)^2(1 + t - t^2)^2 - (1 - t^2)^3 = t^2(1 - t)^2(2t^2 - 1) > 0.$$

For $\alpha \in (1/2, 1)$ one has $t \in (0, \frac{\sqrt{2}}{2})$ and $G(\alpha) > H(\alpha)$. For $\alpha \in (0, 1/2)$ one has $t \in (\frac{\sqrt{2}}{2}, 1)$ and $G(\alpha) < H(\alpha)$.

There are several weaker and better-looking bounds on $F(n, e)$, but they do not meet our aims.

5 Proof of Theorem 3

Put $k = |V_+|$. Let the degrees of vertices in $G[V_+]$ be equal to a_1, \dots, a_k ; the degrees of vertices in $G[V_+, V_-]$ be equal to b_1, \dots, b_k for $b_i \in V_+$ and c_1, \dots, c_{n-k} for $c_j \in V_-$. Define

$$a = \frac{1}{k} \sum_{1 \leq i \leq k} a_i; \quad b = \frac{1}{k} \sum_{1 \leq i \leq k} b_i; \quad c = \frac{1}{n-k} \sum_{1 \leq j \leq n-k} c_j;$$

by double-counting in the graph $G[V_+, V_-]$ we have $kb = (n-k)c$.

By the main condition, if we have an edge between (v_i^+, v_j^-) then

$$a_i - b_i \geq c_j.$$

Sum up all these inequalities; then every vertex v_i^+ is counted b_i times, every vertex v_j^- is counted c_j times. Hence

$$\sum_{1 \leq i \leq k} (a_i - b_i)b_i \geq \sum_{1 \leq j \leq n-k} c_j^2.$$

Applying Cauchy – Bunyakovsky – Schwarz inequality, we get

$$\sqrt{\sum_{1 \leq i \leq k} a_i^2 \sum_{1 \leq i \leq k} b_i^2} - \sum_{1 \leq i \leq k} b_i^2 \geq \sum_{1 \leq i \leq k} (a_i - b_i)b_i.$$

AM-GM inequality implies

$$\sum_{1 \leq j \leq n-k} c_j^2 \geq (n-k)c^2 = \frac{k^2}{n-k} b^2.$$

Put

$$\alpha = \frac{a}{k}; \quad B = \sqrt{\frac{1}{k} \sum_{1 \leq i \leq k} b_i^2}; \quad K = \frac{k}{n}.$$

Then we reduce our problem to the following optimization problem:

$$\begin{cases} W(\alpha)B - B^2 \geq b^2 \frac{K}{1-K}; \\ \text{minimize} & \frac{\alpha}{2} K^2 - bK^2; \\ 0 \leq \alpha \leq 1, & 0 \leq K \leq 1, \quad 0 \leq b \leq B \leq \frac{1-K}{K}, \end{cases} \quad (2)$$

where $W(\alpha) = \max(\sqrt{G(\alpha)}, \sqrt{H(\alpha)})$. We show that the desired minimum equals to $-\frac{1}{54}$; it can be reached by the example from Theorem 1(ii). Note that a possible (with respect to conditions of the system (2)) value of (α, b, B, K) may not correspond to a SED-pair.

Case 1. In this case $\alpha \geq \frac{1}{2}$, so $W(\alpha) = \sqrt{G(\alpha)}$. Then we have to solve the following system

$$\begin{cases} \alpha^{\frac{3}{4}} B - B^2 \geq b^2 \frac{K}{1-K}; \\ 0 \leq \alpha \leq \frac{1}{2}, \quad 0 < K < 1, \quad 0 \leq b \leq \frac{1-K}{K}, \quad 0 \leq B \leq \frac{1-K}{K}; \\ \text{minimize} & \frac{\alpha}{2} K^2 - bK^2. \end{cases}$$

In Appendix C we show that the minimum is $-\frac{1}{54}$.

Case 2. In this case $\alpha \leq \frac{1}{2}$. Then $W(\alpha) = \sqrt{H(\alpha)}$. Then we have a deal with the following system

$$\begin{cases} \sqrt{(1 - \sqrt{1 - \alpha})(\sqrt{1 - \alpha} + \alpha)} B - B^2 \geq b^2 \frac{K}{1-K}; \\ 0 \leq \alpha \leq \frac{1}{2}, \quad 0 < K < 1, \quad 0 \leq b \leq \frac{1-K}{K}, \quad 0 \leq B \leq \frac{1-K}{K}; \\ \text{minimize} & \frac{\alpha}{2} K^2 - bK^2. \end{cases}$$

This system is solved in Appendix D; the minimum is bigger than the desired value $-\frac{1}{54}$.

So we prove $s[(G, f)] \geq -(1 + o(1)) \frac{n^2}{54}$; Theorem 4 finishes the proof.

Acknowledgments. The research of Danila Cherkashin is supported by «Native towns», a social investment program of PJSC «Gazprom Neft» (Sections 2 and 3) and by Grant NSh-2540.2020.1 to support leading scientific schools of Russia (Section 5). The authors are grateful to Fedor Petrov for an introduction in graphon theory and to Georgii Strukov for some drawing.

References

- [1] Rudolf Ahlswede and Gyula O. H. Katona. Graphs with maximal number of adjacent pairs of edges. *Acta Mathematica Hungarica*, 32(1-2):97–120, 1978.
- [2] Saeed Akbari, Sadegh Bolouki, Pooya Hatami, and Milad Siami. On the signed edge domination number of graphs. *Discrete mathematics*, 309(3):587–594, 2009.
- [3] László Lovász. *Large networks and graph limits*, volume 60. American Mathematical Soc., 2012.
- [4] Vladimir Nikiforov. The sum of the squares of degrees: Sharp asymptotics. *Discrete Mathematics*, 307(24):3187–3193, 2007.
- [5] Baogen Xu. On signed edge domination numbers of graphs. *Discrete Mathematics*, 239(1-3):179–189, 2001.
- [6] Baogen Xu. On edge domination numbers of graphs. *Discrete Mathematics*, 294(3):311–316, 2005.

Appendix A

Proof of Theorem 4. (i) Let $(G = (V, E), f)$ be an SED-pair of order n . We partition $[0, 1]$ onto n disjoint sets of measure $1/n$ and identify these n sets with n vertices of G . For points $x, y \in [0, 1]$ denote by v, u the vertices which contain them, respectively, and put

$$W(x, y) = \begin{cases} f(v, u), & \text{if } (v, u) \in E \\ 0, & \text{otherwise.} \end{cases}$$

It is easy to see that $\int_0^1 (W(x, t) + W(y, t)) dt = s_v + s_u \geq 0$ whenever $x \in v, y \in u$ and $(v, u) \in E$. Thus signed graphon W is edge-dominated, and $\kappa \leq \frac{1}{2} \int_0^1 \int_0^1 W = \frac{1}{n^2} s(G, f)$ that proves (i).

(ii) Fix $\varepsilon \in (0, 1)$ and an edge-dominated signed graphon W such that $\frac{1}{2} \int_0^1 \int_0^1 W < \kappa + \varepsilon$. Let n be a (large) integer. Denote $k = \lfloor \varepsilon n \rfloor$, $m = n - k$. Since $\varepsilon > 0$ is arbitrary, and the lower bound $g(n) \geq \kappa n^2$ is already established in (i), for proving (ii) it suffices to prove that

$$g(n) \leq 2kn + m^2(\kappa + \varepsilon) \quad (3)$$

for all large enough n .

Choose m points $v_1, \dots, v_m \in [0, 1]$ uniformly and independently at random. Denote $V = \{1, 2, \dots, n\}$, and define the signed graph $G = (V, E)$ as follows:

- 1) if $i > m$, the vertex i is joined with all other vertices and $f(i, j) = 1$ for all $j \in V \setminus \{i\}$;
- 2) if $i, j \leq m$, we join i and j by an edge with probability $|W(v_i, v_j)|$ and put $f(i, j) = \text{sign}W(v_i, v_j)$ if i and j become joined (the above events are independent).

If we define

$$\tilde{f}(i, j) = \begin{cases} f(i, j), & \text{if } (i, j) \in E \\ 0, & \text{otherwise,} \end{cases}$$

then the expectation of $\tilde{f}(i, j)$ equals $W(v_i, v_j)$. If v_1, \dots, v_m are fixed, the Chernoff bound guarantees that:

a) the probability that $s_i - k = \sum_{j \leq m} \tilde{f}(i, j)$ differs from $\sum_j W(v_i, v_j)$ by a value greater than $k/5$ is exponentially small, and this holds true even if v_1, \dots, v_m are fixed;

b) the probability that $\sum_j W(v_i, v_j)$ differs from $m \int_0^1 W(v_i, t) dt$ by a value greater than $k/5$ is also exponentially small, and this holds true even if v_i is fixed;

c) the probability that $\sum_i \int_0^1 W(v_i, t) dt$ differs from $m \int_0^1 \int_0^1 W(x, y) dx dy$ by more than $k/5$ is also exponentially small.

Therefore with high probability none of the above $2m + 1$ events happens, and we get

$$\left| s_i - k - \int_0^1 W(v_i, t) dt \right| \leq \frac{2k}{5}$$

for all $i = 1, \dots, m$, and

$$\left| \sum_{i=1}^m s_i - km - m^2 \int_0^1 \int_0^1 W \right| \leq \frac{3km}{5}.$$

These bounds yield that (G, f) is an SED-pair, and

$$g(n) \leq s[G, f] = \frac{1}{2} \sum_{j=1}^n s_j \leq k(n-1) + \frac{km}{2} + \frac{3km}{10} + \frac{1}{2} m^2 \int_0^1 \int_0^1 W \leq 2kn + m^2(\kappa + \varepsilon)$$

that is (3). □

Appendix B

We have to calculate

$$\min \left(\max \left(y^2 - \frac{k^2}{2}, -\frac{y(1-k-y)}{2} \right) \right) = -\frac{1}{2} \max \left(\min(k^2 - 2y^2, y - y^2 - ky) \right).$$

Let $k_1, y_1 \in [0, 1]$ be any values representing this maximum (the maximum is reached by compactness).

First, we show that $y_1^2 - \frac{k_1^2}{2} = -\frac{y_1(1-k_1-y_1)}{2}$. Indeed, this equality means that $k_1 = \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2}$. Suppose the contrary; if $k_1 > \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2}$ then

$$\begin{aligned} \min(k_1^2 - 2y_1^2, y_1 - y_1^2 - k_1 y_1) &\leq y_1 - y_1^2 - k_1 y_1 < \\ &< y_1 - y_1^2 - y_1 \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2} = \left(\frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2} \right)^2 - 2y_1^2 = \\ \min \left(y_1 - y_1^2 - y_1 \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2}, \left(\frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2} \right)^2 - 2y_1^2 \right) \end{aligned}$$

and if $k_1 < \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2}$ then

$$\begin{aligned} \min(k_1^2 - 2y_1^2, y_1 - y_1^2 - k_1 y_1) &\leq k_1^2 - 2y_1^2 < \\ &< \left(\frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2} \right)^2 - 2y_1^2 = y_1 - y_1^2 - y_1 \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2} = \\ \min \left(y_1 - y_1^2 - y_1 \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2}, \left(\frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2} \right)^2 - 2y_1^2 \right). \end{aligned}$$

In both cases

$$\min(k_1^2 - 2y_1^2, y_1 - y_1^2 - k_1 y_1) < \min \left(y_1 - y_1^2 - y_1 \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2}, \left(\frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2} \right)^2 - 2y_1^2 \right),$$

and $0 < \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2} < 1$ (because $y_1 < \sqrt{5y_1^2 + 4y_1} < y_1 + 2$), so (k_1, y_1) doesn't represent the maximum, a contradiction.

Since $y_1^2 - \frac{k_1^2}{2} = -\frac{y_1(1-k_1-y_1)}{2}$ for $k_1 = \frac{-y_1 + \sqrt{5y_1^2 + 4y_1}}{2}$, one may search for $\max T(y)$ with $0 \leq y \leq 1$, where

$$T(y) = y - y^2 - y \frac{-y + \sqrt{5y^2 + 4y}}{2} = y - y \frac{y + \sqrt{5y^2 + 4y}}{2}.$$

Consider the derivative of T

$$\begin{aligned} T'(y) &= \left(y - y \frac{y + \sqrt{5y^2 + 4y}}{2} \right)' = 1 - y - \frac{\sqrt{5y^2 + 4y}}{2} - y \frac{10y + 4}{4\sqrt{5y^2 + 4y}} = \\ &= \frac{(y + \sqrt{5y^2 + 4y})(5y - 1)(y + 1)}{\sqrt{5y^2 + 4y}(\sqrt{5y^2 + 4y} + 1)}. \end{aligned}$$

For $y > \frac{1}{5}$ one has $T'(y) < 0$, so $T(y) < T(\frac{1}{5})$ for each $y > \frac{1}{5}$. Analogously $y < \frac{1}{5}$ one has $T'(y) > 0$, so $T(y) < T(\frac{1}{5})$ for each $y < \frac{1}{5}$. Then $T(y) \leq T(\frac{1}{5}) = \frac{2}{25}$ for each $y \in [0, 1]$. So

$$\min \left(\max \left(y^2 - \frac{k^2}{2}, -\frac{y(1-k-y)}{2} \right) \right) = -\frac{1}{2} \max \left(\min(k^2 - 2y^2, y - y^2 - ky) \right) = -\frac{1}{2} \max T(y) = -\frac{1}{25}.$$

6 Appendix C

Here we solve the system

$$\begin{cases} \alpha^{\frac{3}{4}} B - B^2 \geq b^2 \frac{K}{1-K}; \\ \frac{1}{2} \leq \alpha \leq 1, \quad 0 < K < 1, \quad 0 \leq b \leq \frac{1-K}{K}, \quad 0 \leq B \leq \frac{1-K}{K}; \\ \text{minimize} \quad \frac{\alpha}{2} K^2 - bK^2. \end{cases}$$

Case 1: $K > \frac{1}{2}$. Then by AM-GM inequality $\sqrt{G(\alpha)} \geq 2b\sqrt{\frac{K}{1-K}}$ and equality holds for $B = b\sqrt{\frac{K}{1-K}}$. Then

$$b \leq \frac{\sqrt{G(\alpha)}}{2\sqrt{\frac{K}{1-K}}} = \frac{\sqrt{G(\alpha)}}{2} \sqrt{\frac{1-K}{K}}.$$

Hence

$$\frac{\alpha}{2}K^2 - bK^2 \geq \frac{\alpha}{2}K^2 - \frac{\sqrt{G(\alpha)}}{2}\sqrt{1-K}K^{3/2} =: q(\alpha, K);$$

we are going to minimize $q(\alpha, K)$. Derive with respect to K :

$$\frac{dq(\alpha, K)}{dK} = K\alpha - \frac{\sqrt{G(\alpha)}}{2} \frac{3-4K}{2\sqrt{\frac{1-K}{K}}}.$$

Find the roots of the derivative. We may multiply by $\sqrt{\frac{1-K}{K}}$

$$\sqrt{(1-K)K}\alpha = \sqrt{G(\alpha)} \left(\frac{3}{4} - K \right).$$

Then $K < \frac{3}{4}$. Square the equation

$$(1-K)K\alpha^2 = G(\alpha) \left(\frac{3}{4} - K \right)^2.$$

It is quadratic in K

$$(\alpha^2 + G(\alpha))K^2 - \left(\frac{3}{2}G(\alpha) + \alpha^2 \right) K + \frac{9}{16}G(\alpha) = 0.$$

Then $D = \frac{3}{4}G(\alpha)\alpha^2 + \alpha^4$ and the roots are

$$K_1 = \frac{(\frac{3}{2}G(\alpha) + \alpha^2) + \sqrt{\frac{3}{4}G(\alpha)\alpha^2 + \alpha^4}}{2(\alpha^2 + G(\alpha))}; \quad K_2 = \frac{(\frac{3}{2}G(\alpha) + \alpha^2) - \sqrt{\frac{3}{4}G(\alpha)\alpha^2 + \alpha^4}}{2(\alpha^2 + G(\alpha))}.$$

Obviously, the first root is always bigger than $3/4$. Note that

$$K_2 = \frac{1}{2} + \frac{\frac{G(\alpha)}{2} - \sqrt{\frac{3}{4}G(\alpha)\alpha^2 + \alpha^4}}{2(\alpha^2 + G(\alpha))}.$$

Easily

$$\sqrt{\frac{3}{4}G(\alpha)\alpha^2 + \alpha^4} > \alpha^2 > \frac{1}{2}\alpha^{1.5} = \frac{G(\alpha)}{2}$$

since $\alpha \geq \frac{1}{2}$ so the second root is smaller than $1/2$. Hence we should check only $K = 1/2$ and $K = 1$. Clearly $q(\alpha, 1)$ is non-negative; one may check (see Fig. 2 that $q(\alpha, \frac{1}{2})$ is bigger than $-\frac{1}{54}$).




Figure 2: The plot of $q(\alpha, \frac{1}{2}) + \frac{1}{54}$




Figure 3: The plot of $q(\alpha, K_0(\alpha)) + \frac{1}{54}$

Case 2: $K < \frac{1}{2}$. Consider

$$\sqrt{G(\alpha)} \geq B + \frac{b^2}{B} \frac{K}{1-K}.$$

It also implies that $B \geq b\sqrt{\frac{K}{1-K}}$ but the condition $B \geq b$ is stronger since $K < \frac{1}{2}$. Then the optimal B is equal to b and hence $b = \sqrt{G(\alpha)}(1-K)$ and we minimize

$$q(\alpha, K) := \frac{\alpha}{2}K^2 - \sqrt{G(\alpha)}(1-K)K^2.$$

The derivative with respect to K is

$$\alpha K - 2\sqrt{G(\alpha)}K + 3\sqrt{G(\alpha)}K^2.$$

It has zeros at 0 and $\frac{2\sqrt{G(\alpha)} - \alpha}{3\sqrt{G(\alpha)}}$. The derivative is negative on $(0, \frac{2\sqrt{G(\alpha)} - \alpha}{3\sqrt{G(\alpha)}})$, so $q(\alpha, K)$ is decreasing. After $\frac{2\sqrt{G(\alpha)} - \alpha}{3\sqrt{G(\alpha)}}$ the derivative is positive, so the function increases. Hence $q(\alpha, K)$ has local minimum in K at

$$K_0(\alpha) = \frac{2\sqrt{G(\alpha)} - \alpha}{3\sqrt{G(\alpha)}} = \frac{2}{3} - \frac{\alpha}{3\sqrt{G(\alpha)}}.$$

Substitution gives

$$q(\alpha, K_0(\alpha)) = \frac{\alpha}{2} \left(\frac{2}{3} - \frac{\alpha}{3\sqrt{G(\alpha)}} \right)^2 - \sqrt{G(\alpha)} \left(\frac{1}{3} + \frac{\alpha}{3\sqrt{G(\alpha)}} \right) \left(\frac{2}{3} - \frac{\alpha}{3\sqrt{G(\alpha)}} \right)^2 = \frac{\sqrt{h(a)}}{54} \left(\frac{a}{\sqrt{h(a)}} - 2 \right)^3.$$

One may check (see Fig. 3) that $q(\alpha, K_0(\alpha)) > -\frac{1}{54}$.

7 Appendix D

Now we solve the system

$$\begin{cases} \sqrt{(1-\sqrt{1-\alpha})(\sqrt{1-\alpha}+\alpha)}B - B^2 \geq b^2 \frac{K}{1-K}; \\ 0 \leq \alpha \leq \frac{1}{2}, \quad 0 < K < 1, \quad 0 \leq b \leq \frac{1-K}{K}, \quad 0 \leq B \leq \frac{1-K}{K}; \\ \text{minimize} \quad \frac{\alpha}{2}K^2 - bK^2. \end{cases}$$

First, consider $H(\alpha)$. Since it is positive, $\sqrt{H(\alpha)}$ and $H(\alpha)$ have the same intervals of monotonicity. Change the variable $t = \sqrt{1-\alpha}$. Note that $\alpha \in [0; 1/2)$ implies $t \in (\frac{1}{\sqrt{2}}; 1]$. Then

$$H(\alpha) = (1-t)(t+1-t^2) = t^3 - 2t^2 + 1.$$

Since $H'(t) = 3t^2 - 4t = 3t(t - \frac{4}{3}) < 0$ for all t , $H(t)$ is decreasing function. Note that $t(\alpha)$ is decreasing, so $\sqrt{H(\alpha)}$ and $H(\alpha)$ are increasing functions.

Consider two cases.

Case 1: $K > \frac{1}{2}$. Then by AM-GM inequality $\sqrt{H(\alpha)} \geq 2b\sqrt{\frac{K}{1-K}}$ and equality holds for $B = b\sqrt{\frac{K}{1-K}}$. Then

$$b \leq \frac{\sqrt{H(\alpha)}}{2\sqrt{\frac{K}{1-K}}} = \frac{\sqrt{H(\alpha)}}{2} \sqrt{\frac{1-K}{K}}.$$

Analogously to Appendix C we reduce to finding the minimum of

$$q(\alpha, K) := \frac{\alpha}{2}K^2 - \frac{\sqrt{H(\alpha)}}{2}\sqrt{1-K}K^{3/2}.$$

Again derive with respect to K and find the roots

$$K_1 = \frac{(\frac{3}{2}H(\alpha) + \alpha^2) + \sqrt{\frac{3}{4}H(\alpha)\alpha^2 + \alpha^4}}{2(\alpha^2 + H(\alpha))}; \quad K_2 = \frac{(\frac{3}{2}H(\alpha) + \alpha^2) - \sqrt{\frac{3}{4}H(\alpha)\alpha^2 + \alpha^4}}{2(\alpha^2 + H(\alpha))}.$$

Obviously, $K_1 > 3/4$. So the only possible root is K_2 . We should examine $K = K_2$ (in the case when it is bigger than $1/2$), $K = \frac{1}{2}$ and $K = 1$. One can see (for example by compare the plots on Fig. 4 and Fig. 5) that $K_2(\alpha) > \frac{1}{2}$ implies that $q(\alpha, K_2(\alpha))$ is bigger than $-\frac{1}{54}$.




Figure 4: The plot of $q(\alpha, K_2(\alpha)) + \frac{1}{54}$




Figure 5: The plot of $K_2(\alpha) - \frac{1}{2}$

Finally, note that for $K = 1$ function q is positive. For $K = 1/2$ one may see the plot on Fig. 6 to check that $q(\alpha, \frac{1}{2}) > -\frac{1}{54}$.




Figure 6: The plot of $q\left(\alpha, \frac{1}{2}\right) + \frac{1}{54}$




Figure 7: The plot of $q\left(\alpha, K_0(\alpha)\right) + \frac{1}{54}$

Case 2: One can repeat step-by-step the second case of Appendix C. We minimize

$$q(\alpha, K) := \frac{\alpha}{2}K^2 - \sqrt{H(\alpha)}(1 - K)K^2.$$

Derivation and substitution gives

$$q(\alpha, K_0(\alpha)) = \frac{\sqrt{H(a)}}{54} \left(\frac{a}{\sqrt{H(a)}} - 2 \right)^3.$$

One may check (see Fig. 7) that $q(\alpha, K_0(\alpha)) > -\frac{1}{54}$.