

Parallelisms and Translations of (Affine) $SL(2, q)$ -Unitals

Verena Möhler

December 17, 2020

Abstract

Unitals can be obtained as closures of affine unitals via parallelisms. The isomorphism type of the closure depends on the chosen parallelism, which need not be unique. For affine $SL(2, q)$ -unitals, we introduce a class of parallelisms for odd order and one for square order.

Translations are automorphisms of unitals, fixing each block through a given center. For each of the known parallelisms of affine $SL(2, q)$ -unitals, we compute all possible translations with centers on the block at infinity.

2020 MSC: 51A15, 51A10, 05E18

Keywords: design, unital, affine unital, parallelism, automorphism, translation

Most of the results in the present paper have been obtained in the author's Ph.D. thesis [7], where detailed arguments can be found for some statements that we leave to the reader here.

1 Preliminaries

Projective planes arise from affine planes via a unique parallelism. The strategy of building the affine part of a geometry first and then completing it by using a parallelism on the blocks can successfully be applied to other incidence structures than affine and projective planes. We apply this approach to a special construction of unitals. A crucial difference to affine planes is, that for a given affine unital, there may be various parallelisms, leading to non-isomorphic closures.

A **unital of order n** is a $2-(n^3 + 1, n + 1, 1)$ design, i. e. an incidence structure with $n^3 + 1$ points, $n + 1$ points on each block and unique joining blocks for any two points. We also consider *affine unitals*, which arise from unitals by removing one block (and all the points on it) and can be completed to unitals via a parallelism on the short blocks. We give an axiomatic description:

Definition 1.1. Let $n \in \mathbb{N}$, $n \geq 2$. An incidence structure $\mathbb{U} = (\mathcal{P}, \mathcal{B}, I)$ is called an **affine unital of order n** if:

- (AU1) There are $n^3 - n$ points.
- (AU2) Each block is incident with either n or $n + 1$ points. The blocks incident with n points will be called **short blocks** and the blocks incident with $n + 1$ points will be called **long blocks**.
- (AU3) Each point is incident with n^2 blocks.
- (AU4) For any two points there is exactly one block incident with both of them.
- (AU5) There exists a **parallelism** on the short blocks, meaning a partition of the set of all short blocks into $n + 1$ parallel classes of size $n^2 - 1$ such that the blocks of each parallel class are pairwise non-intersecting.

The existence of a parallelism as in (AU5) must explicitly be required (see [7, Example 3.10]). An affine unital \mathbb{U} of order n with parallelism π can be completed to a unital \mathbb{U}^π of order n as follows: For each parallel class, add a new point that is incident with each short block of that class. Then add a single new block $[\infty]^\pi$, incident with the $n + 1$ new points (see [7, Proposition 3.9]). We call \mathbb{U}^π the **π -closure** of \mathbb{U} .

Note that though we must require the existence of a parallelism in the definition of an affine unital, this parallelism need not be unique. It is therefore not convenient to require that isomorphisms of affine unitals respect certain parallelisms and we will only ask them to be isomorphisms of incidence structures. We call two parallelisms π and π' of an affine unital \mathbb{U} **equivalent** if there is an automorphism of \mathbb{U} which maps π to π' .

Given an affine unital \mathbb{U} with parallelisms π and π' , the closures \mathbb{U}^π and $\mathbb{U}^{\pi'}$ are isomorphic with $[\infty]^\pi \mapsto [\infty]^{\pi'}$ exactly if π and π' are equivalent (see [7, Proposition 3.12]).

2 Parallelisms of Affine $\text{SL}(2, q)$ -Unitals

From now on let p be a prime and $q := p^e$ a p -power. We are interested in a special kind of affine unitals, namely affine $\text{SL}(2, q)$ -unitals. The construction of those affine unitals is due to Grundhöfer, Stroppel and Van Maldeghem [3]. They consider translations of unitals, i. e. automorphisms fixing each block through a given point (the so-called center). Of special interest are unitals of order q where two points are centers of translation groups of order q . In the classical (Hermitian) unital of order q , any two such translation groups generate a group isomorphic to $\text{SL}(2, q)$; see [2, Main Theorem] for further possibilities. The construction of (affine) $\text{SL}(2, q)$ -unitals is motivated by this action of $\text{SL}(2, q)$ on the classical unital.

Let $S \leq \mathrm{SL}(2, q)$ be a subgroup of order $q+1$ and let $T \leq \mathrm{SL}(2, q)$ be a Sylow p -subgroup. Recall that T has order q (and thus trivial intersection with S), that any two conjugates $T^h := h^{-1}Th$, $h \in \mathrm{SL}(2, q)$, have trivial intersection unless they coincide and that there are $q+1$ conjugates of T .

Consider a collection \mathcal{D} of subsets of $\mathrm{SL}(2, q)$ such that each set $D \in \mathcal{D}$ contains $\mathbf{1} := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, that $\#D = q+1$ for each $D \in \mathcal{D}$, and the following properties hold:

(Q) For each $D \in \mathcal{D}$, the map

$$(D \times D) \setminus \{(x, x) \mid x \in D\} \rightarrow \mathrm{SL}(2, q), \quad (x, y) \mapsto xy^{-1},$$

is injective, i. e. the set $D^* := \{xy^{-1} \mid x, y \in D, x \neq y\}$ contains $q(q+1)$ elements.

(P) The system consisting of $S \setminus \{\mathbf{1}\}$, all conjugates of $T \setminus \{\mathbf{1}\}$ and all sets D^* with $D \in \mathcal{D}$ forms a partition of $\mathrm{SL}(2, q) \setminus \{\mathbf{1}\}$.

Set

$$\mathcal{P} := \mathrm{SL}(2, q),$$

$$\mathcal{B} := \{Sg \mid g \in \mathrm{SL}(2, q)\} \cup \{T^h g \mid h, g \in \mathrm{SL}(2, q)\} \cup \{Dg \mid D \in \mathcal{D}, g \in \mathrm{SL}(2, q)\}$$

and let the incidence relation $I \subseteq \mathcal{P} \times \mathcal{B}$ be containment.

Then we call the incidence structure $\mathbb{U}_{S, \mathcal{D}} := (\mathcal{P}, \mathcal{B}, I)$ an **affine $\mathrm{SL}(2, q)$ -unital**. Each affine $\mathrm{SL}(2, q)$ -unital is indeed an affine unital of order q , see [7, Prop. 3.15]. If π is a parallelism on the short blocks of an affine $\mathrm{SL}(2, q)$ -unital $\mathbb{U}_{S, \mathcal{D}}$, we call the π -closure $\mathbb{U}_{S, \mathcal{D}}^\pi$ an **$\mathrm{SL}(2, q)$ -(π -)unital**.

For the investigation of parallelisms of affine unitals, we only need to know how the short blocks look like. In any affine $\mathrm{SL}(2, q)$ -unital, the set of short blocks is the set of all right cosets of the Sylow p -subgroups. A parallelism as in (AU5) means a partition of the set of short blocks into $q+1$ sets of q^2-1 pairwise non-intersecting blocks.

Definition 2.1. Let \mathfrak{P} denote the set of all Sylow p -subgroups of $\mathrm{SL}(2, q)$.

(a) We denote by $\mathfrak{S} := \{Tg \mid T \in \mathfrak{P}, g \in \mathrm{SL}(2, q)\}$ the set of short blocks of any affine $\mathrm{SL}(2, q)$ -unital.

(b) A **parallelism on \mathfrak{S}** is a partition of \mathfrak{S} into $q+1$ sets of q^2-1 pairwise non-intersecting cosets.

Note that each right coset Tg is a left coset gT^g of a conjugate of T . For each prime power q , there are hence two obvious parallelisms on \mathfrak{S} , namely partitioning the set of short blocks into the sets of *right* cosets or into the sets of *left* cosets of the Sylow p -subgroups. We name those two parallelisms “flat” and “natural”, respectively, and denote them by the corresponding musical signs

$$\flat := \{\{Tg \mid g \in \mathrm{SL}(2, q)\} \mid T \in \mathfrak{P}\} \quad \text{and} \quad \natural := \{\{gT \mid g \in \mathrm{SL}(2, q)\} \mid T \in \mathfrak{P}\}.$$

Remark 2.2. *Inspired by the construction of $\mathrm{SL}(2, q)$ -unitals from affine $\mathrm{SL}(2, q)$ -unitals via different parallelisms, Nagy and Mezőfi created a method of constructing new incidence structures from old ones by removing a block and attaching it again in a different way, see [5]. They call this method paramodification and used it to construct plenty new unitals of orders 3 and 4, respectively. Their new unitals also comprise the Leonidis unitals (see Section 2.3), since these are obtained from the classical unital of order 4 by paramodification.*

On any affine $\mathrm{SL}(2, q)$ -unital $\mathbb{U}_{S, \mathcal{D}}$, the group $\mathrm{SL}(2, q)$ acts as a group of automorphisms by multiplication from the right. We denote this permutation group by $R := \{\rho_h \mid h \in \mathrm{SL}(2, q)\}$, where ρ_h acts by right multiplication with $h \in \mathrm{SL}(2, q)$. Any automorphism of $\mathrm{SL}(2, q)$ induces a permutation on the point set of $\mathbb{U}_{S, \mathcal{D}}$; we call the corresponding permutation group $\mathfrak{A} \cong \mathrm{Aut}(\mathrm{SL}(2, q))$. For each $a \in \mathrm{GL}(2, q)$, conjugation by a induces a permutation $\gamma_a \in \mathfrak{A}$. Although not every $\alpha \in \mathfrak{A}$ acts as automorphism on $\mathbb{U}_{S, \mathcal{D}}$, we know that for $q \geq 3$, the full automorphism group of $\mathbb{U}_{S, \mathcal{D}}$ is a subgroup of $\mathfrak{A} \times R$, more precisely a subgroup of $\mathfrak{A}_S \times R$ (see [6, Theorem 3.3]). We are therefore also interested in the stabilizers of our parallelisms under the action of $\mathfrak{A} \times R$.

Regarding the parallelisms \mathfrak{b} and \mathfrak{c} , we see that both are stabilized by $\mathfrak{A} \times R$. Note that right multiplication with $h \in \mathrm{SL}(2, q)$ fixes each parallel class of \mathfrak{b} while it acts on the parallel classes of \mathfrak{c} via conjugation on the Sylow p -subgroups $T \in \mathfrak{P}$ (since $gTh = ghT^h$).

Example 2.3.

- (a) For each prime power q we may choose $S = C$ to be cyclic and \mathcal{H} a set of blocks through $\mathbb{1}$ such that $\mathbb{U}_{C, \mathcal{H}}$ is isomorphic to the affine part of the classical unital and the closure $\mathbb{U}_{C, \mathcal{H}}^{\mathfrak{b}}$ is isomorphic to the classical unital. We call $\mathbb{U}_{C, \mathcal{H}}$ the **classical affine $\mathrm{SL}(2, q)$ -unital** and $\mathbb{U}_{C, \mathcal{H}}^{\mathfrak{b}}$ the **classical $\mathrm{SL}(2, q)$ -unital**. See [3, Example 3.1] or [7, Section 3.2.2] for details.
- (b) Several non-classical affine $\mathrm{SL}(2, q)$ -unitals are known, namely one of order 4, described in [3], and three of order 8, described in [8, Section 3].

2.1 A Class of Parallelisms for Odd Order

For each odd prime power q , there is at least one class of parallelisms on \mathfrak{S} apart from \mathfrak{b} and \mathfrak{c} . Let q be odd throughout this section.

Let T be a fixed Sylow p -subgroup of $\mathrm{SL}(2, q)$, namely

$$T := \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{F}_q \right\}.$$

The normalizer of T in $\mathrm{GL}(2, q)$ is the set of upper triangular matrices. Let \mathbb{F}_q^{\square} denote the set of all squares in \mathbb{F}_q , let $\mathbb{F}_q^{\times, \square}$ denote the set of all squares in \mathbb{F}_q^{\times} and $\mathbb{F}_q^{\times, \square^c}$ the set of all non-squares in \mathbb{F}_q^{\times} . Note that $\#\mathbb{F}_q^{\times, \square} = \#\mathbb{F}_q^{\times, \square^c} = \frac{1}{2}(q - 1)$ since q is odd. Let

$$\Omega := \left\{ g \in \mathrm{SL}(2, q) \mid (0, 1) \cdot g \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} \in \mathbb{F}_q^{\square} \right\} \quad \text{and} \quad \Omega^c := \mathrm{SL}(2, q) \setminus \Omega.$$

Note that Ω is a union of T -cosets (left as well as right cosets) since

$$(0, 1) \cdot Tg \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \{(0, 1) \cdot g \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix}\} = (0, 1) \cdot gT \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Let further

$$\begin{aligned} A &:= \{Tg \mid g \in \Omega\} \cup \{gT \mid g \in \Omega^c\}, \\ A' &:= \{Tg \mid g \in \Omega^c\} \cup \{gT \mid g \in \Omega\} \end{aligned}$$

and¹

$$\pi_{\text{odd}} := \{A^h \mid h \in \text{SL}(2, q)\}, \quad \pi'_{\text{odd}} := \{A'^h \mid h \in \text{SL}(2, q)\}.$$

Theorem 2.4. *For odd q , the sets π_{odd} and π'_{odd} are parallelisms on \mathfrak{S} . With $v \in \mathbb{F}_q^{\times, \square}$, conjugation by $\begin{pmatrix} 1 & 0 \\ 0 & v \end{pmatrix}$ maps π_{odd} to π'_{odd} .*

Proof. Since Ω is a union of T -cosets, the cosets in A are pairwise non-intersecting and we get

$$\#A = \frac{\#\Omega}{\#T} + \frac{\#\Omega^c}{\#T} = \frac{\#\text{SL}(2, q)}{\#T} = q^2 - 1.$$

Hence, A is a set of $q^2 - 1$ pairwise non-intersecting short blocks.

Let $h \in \text{N} := \text{N}_{\text{SL}(2, q)}(T)$. Then h is upper triangular, say $h := \begin{pmatrix} a & * \\ 0 & a^{-1} \end{pmatrix}$ with $a \in \mathbb{F}_q^{\times}$. For any $g \in \text{SL}(2, q)$, we have

$$(0, 1) \cdot g^h \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = (0, a) \cdot g \cdot \begin{pmatrix} a \\ 0 \end{pmatrix} = a^2(0, 1) \cdot g \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Since $a^2 \in \mathbb{F}_q^{\times, \square}$, we get $g^h \in \Omega$ exactly if $g \in \Omega$. Hence, conjugation by h stabilizes A .

Now let $h \in \text{SL}(2, q) \setminus \text{N}$ and $Tg \in A$. Then $(Tg)^h = T^h g^h$ is not a right coset of T . Compute $T^h g^h = g^h (T^h)^{g^h} = g^h T^{g^h}$ and consider $gh \in \text{N}$. Then $h = g^{-1}n$ for an $n \in \text{N}$ and we have $g^h = g^n$. But since $g^n \in \Omega$ exactly if $g \in \Omega$ (as shown above), the coset $(Tg)^h$ is not contained in A . A similar consideration shows that $(gT)^h$ is not contained in A if $gT \in A$ and $h \notin \text{N}$.

Let $g, h \in \text{SL}(2, q)$ and assume $A^h \cap A^g \neq \emptyset$. Then $A^{hg^{-1}} \cap A \neq \emptyset$ and we get $hg^{-1} \in \text{N}$, $A^{hg^{-1}} = A$ and hence $A^h = A^g$. Thus, $\pi_{\text{odd}} = \{A^h \mid h \in \text{SL}(2, q)\}$ is indeed a partition of \mathfrak{S} into $\#\text{SL}(2, q)/\#\text{N} = q + 1$ sets of $q^2 - 1$ pairwise non-intersecting cosets, hence a parallelism on \mathfrak{S} .

Now let $v \in \mathbb{F}_q^{\times, \square}$ and $f := \begin{pmatrix} 1 & 0 \\ 0 & v \end{pmatrix}$. Then $f \in \text{N}_{\text{GL}(2, q)}(T)$ and $g^f \in \text{N}$ exactly if $g \in \text{N}$. Further, for any $g \in \text{SL}(2, q) \setminus \text{N}$, we have

$$(0, 1) \cdot g^f \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = (0, v^{-1}) \cdot g \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = v^{-1}(0, 1) \cdot g \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

¹ Note that in [7], the parallelism π_{odd} is called π^{\square} and π'_{odd} is called ${}^{\square}\pi$.

Hence, $g^f \in \Omega \setminus N$ exactly if $g \in \Omega^c$ and we get $A^f = A'$ (note that $N \subseteq \Omega$). For each $A^h \in \pi_{\text{odd}}$, we get

$$(A^h)^f = A^{hf} = (A^f)^{(h^f)} = A'^{(h^f)} \in \pi'_{\text{odd}}.$$

Hence, we obtain π'_{odd} from π_{odd} via conjugation by f , and π'_{odd} is a parallelism on \mathfrak{S} since π_{odd} is a parallelism on \mathfrak{S} . \square

As mentioned above and according to [6, Theorem 3.3], the full automorphism group of any affine $\text{SL}(2, q)$ -unital of order $q \geq 3$ is a subgroup of $\mathfrak{A} \times R$. We compute the stabilizer of the parallelism π_{odd} in $\mathfrak{A} \times R \cong \text{Aut}(\text{SL}(2, q)) \times \text{SL}(2, q)$ (note that q is odd and hence $q \geq 3$). Recall that $\text{Aut}(\text{SL}(2, q)) = \text{PGL}(2, q) := \text{PGL}(2, q) \rtimes \text{Aut}(\mathbb{F}_q)$, where automorphisms of \mathbb{F}_q act entrywise on matrices. We denote by φ the Frobenius automorphism $x \mapsto x^p$ on \mathbb{F}_q .

Theorem 2.5. *Let $c := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. The stabilizer of π_{odd} in the group $\mathfrak{A} \times R$ equals*

(a) $\text{PSL}(2, q) \times \langle \rho_{-1} \rangle$ if $q \equiv 1 \pmod{4}$ and

(b) $\text{PSL}(2, q) \rtimes \langle \gamma_c \cdot \rho_{-1} \rangle$ if $q \equiv 3 \pmod{4}$,

where $\text{PSL}(2, q) := \text{PSL}(2, q) \rtimes \text{Aut}(\mathbb{F}_q)$.

Proof. Note first that the Frobenius automorphism φ stabilizes Ω and T and hence A and A' . Thus, for each $A^h \in \pi_{\text{odd}}$, we have

$$A^h \cdot \varphi = (A \cdot \varphi)^{h \cdot \varphi} = A^{h \cdot \varphi} \in \pi_{\text{odd}}$$

and $\langle \varphi \rangle = \text{Aut}(\mathbb{F}_q)$ stabilizes π_{odd} and equally π'_{odd} .

The action of $\text{PSL}(2, q)$ obviously stabilizes π_{odd} and π'_{odd} by construction. Since the index of $\text{PSL}(2, q)$ in $\text{PGL}(2, q)$ equals 2, the orbit of π_{odd} under the action of $\text{PGL}(2, q)$ has length 1 or 2. From Theorem 2.4, we know that conjugation by $\begin{pmatrix} 1 & 0 \\ 0 & v \end{pmatrix}$ ($v \in \mathbb{F}_q^{\times, \square}$) maps π_{odd} to π'_{odd} , and hence the stabilizer of π_{odd} in the action of $\text{PGL}(2, q)$ equals $\text{PSL}(2, q)$ and conjugation by any element in $\text{PGL}(2, q) \setminus \text{PSL}(2, q)$ interchanges π_{odd} and π'_{odd} .

Since we know now that $\text{PGL}(2, q)$ stabilizes $\{\pi_{\text{odd}}, \pi'_{\text{odd}}\}$, we need to find those elements in R which also stabilize the set of these two parallelisms. Let $\rho_g \in R$ and assume $\pi_{\text{odd}} \cdot \rho_g \in \{\pi_{\text{odd}}, \pi'_{\text{odd}}\}$. Since for each $A^h \in \pi_{\text{odd}}$ the automorphism ρ_g maps the set of right cosets of T^h in A^h on a set of right cosets of T^h , it must then also map the set of left cosets of T^h in A^h on a set of left cosets of T^h . Hence, g is contained in the normalizer of every Sylow p -subgroup of $\text{SL}(2, q)$ and thus $g \in \{\pm 1\}$. We see immediately that

$$\pi_{\text{odd}} \cdot \rho_{-1} = \begin{cases} \pi_{\text{odd}} & \text{if } -1 \in \mathbb{F}_q^{\times, \square}, \\ \pi'_{\text{odd}} & \text{if } -1 \in \mathbb{F}_q^{\times, \square}. \end{cases}$$

If $q \equiv 1 \pmod{4}$, then $-1 \in \mathbb{F}_q^{\times, \square}$ and ρ_{-1} stabilizes π_{odd} . Since ρ_{-1} does not fix $\mathbb{1}$ – while every automorphism in \mathfrak{A} does – and since ρ_{-1} commutes with every automorphism in \mathfrak{A} , statement (a) follows.

If $q \equiv 3 \pmod{4}$, then both ρ_{-1} and conjugation by c interchange π_{odd} and π'_{odd} and hence the product stabilizes π_{odd} . Again, $\gamma_c \cdot \rho_{-1}$ does not fix $\mathbb{1}$ – while every automorphism in \mathfrak{A} does – and hence the product $\text{P}\Sigma\text{L}(2, q) \cdot \langle \gamma_c \cdot \rho_{-1} \rangle$ is semidirect, since ρ_{-1} commutes with every automorphism in \mathfrak{A} and γ_c normalizes $\text{P}\Sigma\text{L}(2, q) \leq \mathfrak{A}$. \square

Remark 2.6. *Knowing the full automorphism group of $\mathbb{U}_{C, \mathcal{H}}$, we use Theorem 2.5 to compute that the stabilizer of $[\infty]^{\pi_{\text{odd}}}$ in the full automorphism group of the closure $\mathbb{U}_{C, \mathcal{H}}^{\pi_{\text{odd}}}$ has order $2e(q+1)$ (see [7, Corollary 5.3]).*

2.2 A Class of Parallelisms for Square Order

For each square order, there also is at least one class of parallelisms on \mathfrak{S} apart from \mathfrak{b} and \mathfrak{t} . Consider the quadratic field extension $\mathbb{F}_{q^2}/\mathbb{F}_q$ and the unique involutory field automorphism

$$\bar{\cdot}: \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}, x \mapsto \bar{x} := x^q,$$

with fixed field \mathbb{F}_q . We let this automorphism act entrywise on matrices over \mathbb{F}_{q^2} . Let again $T := \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{F}_{q^2} \right\}$ be a fixed Sylow p -subgroup of $\text{SL}(2, q^2)$. Let further

$$\Omega := \left\{ g \in \text{SL}(2, q^2) \mid (0, 1) \cdot g \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} \in \mathbb{F}_q \right\} \quad \text{and} \quad \Omega^c := \text{SL}(2, q^2) \setminus \Omega.$$

As in Section 2.1, Ω is a union of T -cosets. Consider the action

$$\vartheta: \mathbb{F}_{q^2}^{2 \times 2} \times \text{GL}(2, q^2) \rightarrow \mathbb{F}_{q^2}^{2 \times 2}, (x, h) \mapsto x \cdot \vartheta_h := h^{-1} x \bar{h},$$

and note that ϑ_h equals conjugation with h if $h \in \text{GL}(2, q)$.

Lemma 2.7. *Let $h \in \text{N}_{\text{SL}(2, q^2)}(T)$. Then $\Omega \cdot \vartheta_h = \Omega$ and $\Omega^c \cdot \vartheta_h = \Omega^c$.*

Proof. Since h normalizes T , we have h upper triangular, say $h := \begin{pmatrix} a & * \\ 0 & a^{-1} \end{pmatrix}$ with $a \in \mathbb{F}_{q^2}^{\times}$. For any $g \in \text{SL}(2, q^2)$, we have

$$(0, 1) \cdot (g \cdot \vartheta_h) \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = (0, a) \cdot g \cdot \begin{pmatrix} \bar{0} \\ 1 \end{pmatrix} = a\bar{a} \cdot (0, 1) \cdot g \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Since $a\bar{a} \in \mathbb{F}_q^{\times}$, we get $g \cdot \vartheta_h \in \Omega$ exactly if $g \in \Omega$. \square

Theorem 2.8. *Let*

$$A := \{Tg \mid g \in \Omega\} \cup \{gT \mid g \in \Omega^c\} \quad \text{and} \quad \pi_{\text{sq}} := \{A \cdot \vartheta_h \mid h \in \text{SL}(2, q^2)\}.$$

Then π_{sq} is a parallelism on \mathfrak{S} .

Proof. Note first that the cosets in A are pairwise non-intersecting and that

$$\#A = \frac{\#\Omega}{\#T} + \frac{\#\Omega^c}{\#T} = \frac{\#\mathrm{SL}(2, q^2)}{\#T} = q^4 - 1.$$

Hence, A is indeed a set of $(q^2)^2 - 1$ non-intersecting short blocks.

Let $h \in \mathbf{N} := \mathbf{N}_{\mathrm{SL}(2, q^2)}(T)$ and $Tg \in A$. Then

$$(Tg) \cdot \vartheta_h = h^{-1}Tg\bar{h} = h^{-1}Thh^{-1}g\bar{h} = T^h(g \cdot \vartheta_h) = T(g \cdot \vartheta_h) \in A,$$

since ϑ_h leaves Ω invariant (Lemma 2.7). Analogously, $(gT) \cdot \vartheta_h \in A$ if $gT \in A$ and hence $A \cdot \vartheta_h = A$ for $h \in \mathbf{N}$.

Now let $h \in \mathrm{SL}(2, q^2) \setminus \mathbf{N}$ and $Tg \in A$. Then $(Tg) \cdot \vartheta_h = T^h(g \cdot \vartheta_h)$ is not a right coset of T . Assume $(Tg) \cdot \vartheta_h = (g \cdot \vartheta_h)T^{g\bar{h}}$ is a left coset of T , i. e. $n := g\bar{h} \in \mathbf{N}$. Hence,

$$g \cdot \vartheta_h = h^{-1}g\bar{h} = \bar{n}^{-1}\bar{g}n = \bar{g} \cdot \vartheta_{\bar{n}},$$

which is in Ω exactly if $\bar{g} \in \Omega$. But since obviously $\bar{\Omega} = \Omega$ and $g \in \Omega$ since $Tg \in A$, we have $g \cdot \vartheta_h \in \Omega$ and hence $(Tg) \cdot \vartheta_h \notin A$. A similar consideration shows that $(gT) \cdot \vartheta_h \notin A$ if $gT \in A$. Thus, $(A \cdot \vartheta_h) \cap A = \emptyset$ if $h \in \mathrm{SL}(2, q^2) \setminus \mathbf{N}$.

Finally, let $g, h \in \mathrm{SL}(2, q^2)$ and assume $(A \cdot \vartheta_g) \cap (A \cdot \vartheta_h) \neq \emptyset$. Then $(A \cdot \vartheta_{gh^{-1}}) \cap A \neq \emptyset$ and hence $gh^{-1} \in \mathbf{N}$, $A \cdot \vartheta_{gh^{-1}} = A$ and $A \cdot \vartheta_g = A \cdot \vartheta_h$. Thus, $\pi_{\mathrm{sq}} = \{A \cdot \vartheta_h \mid h \in \mathrm{SL}(2, q^2)\}$ is indeed a partition of \mathfrak{S} into $\#\mathrm{SL}(2, q^2)/\#\mathbf{N} = q^2 + 1$ sets of $(q^2)^2 - 1$ pairwise non-intersecting cosets, hence a parallelism on \mathfrak{S} . □

Remark 2.9.

- (a) For each parallel class X of π_{sq} , there exists a unique Sylow p -subgroup $P \in \mathfrak{P}$ such that X consists of $q^3 - 1$ right cosets of P (of which $q^2 - 1$ are left cosets of \bar{P}) and of $q^4 - q^3$ left cosets of \bar{P} , none of which is a right coset of P .
- (b) Applying inversion on $\mathrm{SL}(2, q^2)$ to the parallelism π_{sq} yields another parallelism π_{sq}^{-1} . Since no element of $\mathfrak{A} \rtimes R$ maps π_{sq} to π_{sq}^{-1} , the two parallelisms are not equivalent in any affine $\mathrm{SL}(2, q^2)$ -unital.
- (c) If q^2 is odd, neither π_{sq} nor π_{sq}^{-1} are equivalent to π_{odd} .

As for π_{odd} , we compute the stabilizer of π_{sq} in $\mathfrak{A} \rtimes R$. We abbreviate $\Gamma := \mathrm{Stab}_{\mathfrak{A} \rtimes R}(\pi_{\mathrm{sq}})$.

Lemma 2.10. *Let $x \in \mathbb{F}_{q^2}^\times$ and $c \in \mathbb{F}_q^\times$ such that $x\bar{x} = c^2$. Then $x \in \mathbb{F}_{q^2}^{\times, \square}$.*

Proof. If q is even, then every element in \mathbb{F}_{q^2} is a square. Let q be odd and let z be a generator of $\mathbb{F}_{q^2}^\times$. Note that $\mathbb{F}_q^\times = \{z^{l(q+1)} \mid l \in \mathbb{Z}\}$ and let $k, l \in \mathbb{Z}$ such that $x = z^k$ and $c = z^{l(q+1)}$. Then

$$z^{k(q+1)} = x\bar{x} = c^2 = z^{2l(q+1)},$$

and we get $k - 2l \equiv 0 \pmod{q^2 - 1}$. Since $q^2 - 1$ is even, we get that k is even and hence x is a square. \square

Lemma 2.11.

- (a) $\text{Aut}(\mathbb{F}_{q^2}) \leq \Gamma$.
- (b) Let $r \in \text{SL}(2, q^2)$. We have $\rho_r \in \Gamma$ exactly if $r \in \{\pm 1\}$.
- (c) Let $d \in \mathbb{F}_{q^2}^\times$ and $a := \begin{pmatrix} d & 0 \\ 0 & 1 \end{pmatrix}$. Then $d \in \mathbb{F}_{q^2}^{\times, \square}$ exactly if there exists $r \in \text{SL}(2, q^2)$ with $\gamma_a \rho_r \in \Gamma$.
- (d) Let $a \in \text{GL}(2, q^2)$. Then $\det(a) \in \mathbb{F}_{q^2}^{\times, \square}$ exactly if there exists $r \in \text{SL}(2, q^2)$ with $\gamma_a \rho_r \in \Gamma$.
- (e) Let $a \in \text{GL}(2, q^2)$, let $r, s \in \text{SL}(2, q^2)$ and assume $\gamma_a \rho_r, \gamma_a \rho_s \in \Gamma$. Then $s \in \{\pm r\}$.

Proof. (a) As in the proof of Theorem 2.5, we have that the Frobenius automorphism $\varphi \in \text{Aut}(\mathbb{F}_{q^2})$ stabilizes Ω and T and hence A . Further, φ and $\bar{\cdot}$ commute and we thus get $\pi_{\text{sq}} \cdot \varphi = \pi_{\text{sq}}$ and hence $\text{Aut}(\mathbb{F}_{q^2}) \leq \Gamma$.

(b) Note first that $\Omega \cdot \rho_{-1} = \Omega$ and hence $A \cdot \rho_{-1} = A$. Since for any h , the permutation ρ_{-1} commutes with ϑ_h , we get $\pi_{\text{sq}} \cdot \rho_{-1} = \pi_{\text{sq}}$.

Let $r \in \text{SL}(2, q^2)$ and $P \in \mathfrak{P}$. Recall Remark 2.9 (a). The permutation ρ_r maps each set of right cosets of P to a set of right cosets of P and each set of left cosets of \bar{P} to a set of left cosets of \bar{P}^r . Hence, if $\rho_r \in \Gamma$, then $r \in \text{N}_{\text{SL}(2, q^2)}(P)$ for any $P \in \mathfrak{P}$ and thus $r \in \{\pm 1\}$.

(c) Let $d = c^2 \in \mathbb{F}_{q^2}^{\times, \square}$ and set $a' := c^{-1} \cdot a \in \text{SL}(2, q^2)$. Set $r := a'^{-1} \bar{a}$. Then $\gamma_a \rho_r = \gamma_{a'} \rho_r = \vartheta_{a'} \in \Gamma$.

Conversely, let $d \in \mathbb{F}_{q^2}^\times$ and assume that there exists $r \in \text{SL}(2, q^2)$ such that $\gamma_a \rho_r \in \Gamma$. As above, $\gamma_a \rho_r$ maps each set of right cosets of P to a set of right cosets of P^a and each set of left cosets of \bar{P} to a set of left cosets of \bar{P}^{ar} . Hence, we get $ar\bar{a}^{-1} \in \bigcap_{P \in \mathfrak{P}} \text{N}_{\text{GL}(2, q^2)}(P) = \text{Z}(\text{GL}(2, q^2))$ and there exists $\lambda \in \mathbb{F}_{q^2}^\times$ such that $r = \lambda a^{-1} \bar{a}$. Since $r \in \text{SL}(2, q^2)$, we get $1 = \det(r) = \lambda^2 d^{-1} \bar{d}$. Since a normalizes T and $\gamma_a \rho_r$ stabilizes π_{sq} , we get that $\gamma_a \rho_r$ stabilizes A and hence Ω . Compute

$$(0, 1) \cdot g^a r \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = (0, 1) \cdot \begin{pmatrix} d^{-1} & 0 \\ 0 & 1 \end{pmatrix} g \lambda \begin{pmatrix} \bar{d} & 0 \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \lambda \bar{d} \cdot (0, 1) g \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Hence, $\lambda \bar{d} \in \mathbb{F}_q^\times$. But then $d \bar{d} = (d \bar{d}^{-1}) \bar{d}^2 = \lambda^2 \bar{d}^2 = (\lambda \bar{d})^2$ and we get $d \in \mathbb{F}_{q^2}^{\times, \square}$, according to Lemma 2.10.

- (d) Let $d := \begin{pmatrix} \det a & 0 \\ 0 & 1 \end{pmatrix}$ and $a' := d^{-1}a \in \mathrm{SL}(2, q^2)$. Set $s := a'^{-1}\overline{a'}$. Then $\gamma_{a'}\rho_s = \vartheta_{a'} \in \Gamma$. Assume $\det a \in \mathbb{F}_{q^2}^{\times, \square}$. According to (c), there exists $x \in \mathrm{SL}(2, q^2)$ such that $\gamma_d\rho_x \in \Gamma$. Set $r := x^{a'}s \in \mathrm{SL}(2, q^2)$. Then

$$\gamma_a\rho_r = \gamma_d\gamma_{a'}\rho_{x^{a'}s} = \gamma_d\gamma_{a'}(\gamma_{a'})^{-1}\rho_x\gamma_{a'}\rho_s = \gamma_d\rho_x \cdot \gamma_{a'}\rho_s \in \Gamma.$$

Analogously, if there exists $r \in \mathrm{SL}(2, q^2)$ such that $\gamma_a\rho_r \in \Gamma$, then we have $\gamma_d\rho_{(rs^{-1})^{a'^{-1}}} \in \Gamma$ and the statement follows with (c).

- (e) If $\gamma_a\rho_r \in \Gamma$ and $\gamma_a\rho_s \in \Gamma$ then $\rho_{(rs^{-1})^{a'^{-1}}} = \gamma_a\rho_r(\gamma_a\rho_s)^{-1} \in \Gamma$. According to (b), we thus have $ars^{-1}a^{-1} \in \{\pm 1\}$ and hence $s \in \{\pm r\}$. \square

Theorem 2.12. *Let $q := p^e$ and $\Gamma := \mathrm{Stab}_{\mathfrak{A} \times R}(\pi_{\mathrm{sq}})$. Then*

- (a) $\Gamma = (\langle \vartheta_h \mid h \in \mathrm{SL}(2, q^2) \rangle \rtimes \mathrm{Aut}(\mathbb{F}_{q^2})) \times \langle \rho_{-1} \rangle \cong \mathrm{P}\Sigma\mathrm{L}(2, q^2) \times \langle \rho_{-1} \rangle$.
- (b) $\#\Gamma = 2e(q^2 - 1)q^2(q^2 + 1)$.

Proof. Directly from Lemma 2.11. \square

Remark 2.13.

- (a) *Let π be a parallelism on \mathfrak{S} and π^{-1} the parallelism on \mathfrak{S} obtained by inversion. The stabilizers $\mathrm{Stab}_{\mathfrak{A} \times R}(\pi)$ and $\mathrm{Stab}_{\mathfrak{A} \times R}(\pi^{-1})$ are isomorphic via $\psi: \alpha\rho_r \mapsto \alpha\gamma_r\rho_{r^{-1}}$.*
- (b) *We have $\Gamma \cdot \psi = \Gamma$ and thus $\mathrm{Stab}_{\mathfrak{A} \times R}(\pi_{\mathrm{sq}}^{-1}) = \mathrm{Stab}_{\mathfrak{A} \times R}(\pi_{\mathrm{sq}})$.*

Remark 2.14. *For the classical affine $\mathrm{SL}(2, q^2)$ -unital $\mathbb{U}_{C, \mathcal{H}}$, the stabilizer $\mathrm{Aut}(\mathbb{U}_{C, \mathcal{H}}^{\pi_{\mathrm{sq}}})_{[\infty]} = \mathrm{Aut}(\mathbb{U}_{C, \mathcal{H}}^{\pi_{\mathrm{sq}}^{-1}})_{[\infty]}$ is isomorphic to $(\mathrm{P}\Sigma\mathrm{L}(2, q^2)_C) \times \langle \rho_{-1} \rangle$. The order of $\mathrm{Aut}(\mathbb{U}_{C, \mathcal{H}}^{\pi_{\mathrm{sq}}})_{[\infty]}$ equals $4e(q^2+1)$. For each affine $\mathrm{SL}(2, q^2)$ -unital $\mathbb{U}_{S, \mathcal{D}}$, the order of $\mathrm{Aut}(\mathbb{U}_{S, \mathcal{D}}^{\pi_{\mathrm{sq}}})_{[\infty]} = \mathrm{Aut}(\mathbb{U}_{S, \mathcal{D}}^{\pi_{\mathrm{sq}}^{-1}})_{[\infty]}$ equals the order of $\mathrm{Aut}(\mathbb{U}_{S, \mathcal{D}})_1$.*

2.3 The Leonids Unitals of Order Four

For small orders, parallelisms on \mathfrak{S} can be found using a computer. For $q \in \{3, 5\}$, the classical affine $\mathrm{SL}(2, q)$ -unital $\mathbb{U}_{C, \mathcal{H}}$ represents the only isomorphism type of affine $\mathrm{SL}(2, q)$ -unitals (see [3, Theorem 3.3] and [7, Theorem 6.1]). An exhaustive search using GAP [1] shows that up to equivalence, the three known parallelisms \flat , \natural and π_{odd} are the only ones existing for $q \in \{3, 5\}$ (see [7, Section 6.2.1] for details).

For $q = 4$, there exist two isomorphism types of affine $\mathrm{SL}(2, q)$ -unitals, represented by the classical affine $\mathrm{SL}(2, 4)$ -unital $\mathbb{U}_{C, \mathcal{H}}$ and the non-classical affine $\mathrm{SL}(2, 4)$ -unital described in [3], which we denote by $\mathbb{U}_{C, \mathcal{E}}$. The full automorphism group of $\mathbb{U}_{C, \mathcal{H}}$ is $\mathfrak{A}_C \times R \cong (C_5 \rtimes C_4) \times \mathrm{SL}(2, 4)$ and the full automorphism group of $\mathbb{U}_{C, \mathcal{E}}$ is of type $C_4 \times \mathrm{SL}(2, 4)$ and a subgroup of $\mathrm{Aut}(\mathbb{U}_{C, \mathcal{H}})$.

Table 1: Orbit lengths on the set of 182 parallelisms for order 4

$\text{Aut}(\mathbb{U}_{C,\mathcal{H}})$	$\text{Aut}(\mathbb{U}_{C,\mathcal{E}})$
1	1
1	1
30	24
	6
25	20
	5
5	5
60	60
60	60

For order 4, an exhaustive computer search yields 182 parallelisms on \mathfrak{S} . We consider the actions of the automorphism groups $\text{Aut}(\mathbb{U}_{C,\mathcal{H}})$ and $\text{Aut}(\mathbb{U}_{C,\mathcal{E}})$ on the set of 182 parallelisms found by GAP. The orbit lengths of these actions are listed in Table 1. Apart from \mathfrak{b} and \mathfrak{h} , which are invariant under the action of $\mathfrak{A} \times R$, there are hence seven pairwise non-equivalent parallelisms in $\mathbb{U}_{C,\mathcal{E}}$. We name representatives of those parallelisms π_1, \dots, π_7 such that π_1 and π_2 as well as π_3 and π_4 are equivalent in $\mathbb{U}_{C,\mathcal{H}}$. Following the numbering in [7, Table 6.2], the parallelism π_7 is equivalent to π_{sq} and π_6 is equivalent to π_{sq}^{-1} . In fact, the construction of the parallelism π_{sq} was inspired by the discovery of those parallelisms and partly answers one of the open problems stated in the author's Ph. D. thesis ([7, Question 6 in Chapter 7]). Completing the two affine unitals with the seven parallelisms π_1, \dots, π_7 , we obtain twelve pairwise non-isomorphic $\text{SL}(2, q)$ -unitals, namely

$$\mathbb{U}_{C,\mathcal{H}}^{\pi_2}, \mathbb{U}_{C,\mathcal{H}}^{\pi_4}, \mathbb{U}_{C,\mathcal{H}}^{\pi_5}, \mathbb{U}_{C,\mathcal{H}}^{\pi_6}, \mathbb{U}_{C,\mathcal{H}}^{\pi_7}, \mathbb{U}_{C,\mathcal{H}}^{\pi_1}, \mathbb{U}_{C,\mathcal{E}}^{\pi_2}, \mathbb{U}_{C,\mathcal{E}}^{\pi_3}, \mathbb{U}_{C,\mathcal{E}}^{\pi_4}, \mathbb{U}_{C,\mathcal{E}}^{\pi_5}, \mathbb{U}_{C,\mathcal{E}}^{\pi_6}, \mathbb{U}_{C,\mathcal{E}}^{\pi_7}.$$

Since those parallelisms for order 4 were found during the Leonid meteor shower in November 2018, we call the twelve resulting unitals **Leonids unitals**.

Knowing all affine $\text{SL}(2, 4)$ -unitals and all parallelisms on \mathfrak{S} for order 4, we state the following

Theorem 2.15 (by exhaustive computer search). *There are exactly 16 isomorphism types of $\text{SL}(2, 4)$ -unitals, represented by $\mathbb{U}_{C,\mathcal{H}}^{\mathfrak{h}}, \mathbb{U}_{C,\mathcal{H}}^{\mathfrak{b}}, \mathbb{U}_{C,\mathcal{E}}^{\mathfrak{h}}, \mathbb{U}_{C,\mathcal{E}}^{\mathfrak{b}}$ and the twelve Leonids unitals.* \square

Recall that the stabilizer of the block at infinity in any $\text{SL}(2, q)$ -unital $\mathbb{U}_{S,\mathcal{D}}^{\pi}$ equals the group of those automorphisms of the affine unital $\mathbb{U}_{S,\mathcal{D}}$ which stabilize the parallelism π . We compute with GAP that in each Leonids unital there are indeed no automorphisms moving the block at infinity, and we are thus able to compute their full automorphism groups as subgroups of $\text{Aut}(\mathbb{U}_{C,\mathcal{H}})$ or $\text{Aut}(\mathbb{U}_{C,\mathcal{E}})$, respectively. See Table 2 for the isomorphism types of the full automorphism groups of the Leonids unitals.

Table 2: Isomorphism types of the full automorphism groups of the Leonids unitals

\mathbb{U}	$\text{Aut}(\mathbb{U})$	$\# \text{Aut}(\mathbb{U})$
$\mathbb{U}_{C,\mathcal{H}}^{\pi_2}$	$C_4 \times D_5$	40
$\mathbb{U}_{C,\mathcal{H}}^{\pi_4}$	$C_4 \times A_4$	48
$\mathbb{U}_{C,\mathcal{H}}^{\pi_5}$	$C_4 \times (A_4 \times C_5)$	240
$\mathbb{U}_{C,\mathcal{H}}^{\pi_6}$	$C_5 \times C_4$	20
$\mathbb{U}_{C,\mathcal{H}}^{\pi_7}$	$C_5 \times C_4$	20
$\mathbb{U}_{C,\mathcal{E}}^{\pi_1}$	D_5	10
$\mathbb{U}_{C,\mathcal{E}}^{\pi_2}$	$C_4 \times D_5$	40
$\mathbb{U}_{C,\mathcal{E}}^{\pi_3}$	A_4	12
$\mathbb{U}_{C,\mathcal{E}}^{\pi_4}$	$C_4 \times A_4$	48
$\mathbb{U}_{C,\mathcal{E}}^{\pi_5}$	$C_4 \times A_4$	48
$\mathbb{U}_{C,\mathcal{E}}^{\pi_6}$	C_4	4
$\mathbb{U}_{C,\mathcal{E}}^{\pi_7}$	C_4	4

Remark 2.16. *Regarding the orders of the full automorphism groups of the Leonids unitals in Table 2, we see that the order of $\text{Aut}(\mathbb{U}_{C,\mathcal{H}}^{\pi_5})$ is notably greater than the other orders. Indeed, although R is not contained in its full automorphism group, the unital $\mathbb{U}_{C,\mathcal{H}}^{\pi_5}$ admits a group of automorphisms (of isomorphism type $A_4 \times C_5$) which acts regularly on the affine points.*

3 Translations

We consider a special kind of automorphisms of unitals, namely translations. Since in any closure of an affine $\text{SL}(2, q)$ -unital, the respective parallelism determines the possible translations with centers on the block at infinity, the study of translations is closely related to the study of parallelisms.

Definition 3.1. A **translation** with **center** c of a unital \mathbb{U} is an automorphism of \mathbb{U} that fixes the point c and each block through c . The group of all translations with center c will be denoted by $G_{[c]}$.

We call c a **translation center** if $G_{[c]}$ acts transitively on the set of points different from c on any block through c .

Remark 3.2. *Let \mathbb{U} be a unital of order q and c a point of \mathbb{U} . Then the group $G_{[c]}$ of all translations of \mathbb{U} with center c acts semiregularly on the points of \mathbb{U} different from c and hence semiregularly on the blocks of \mathbb{U} that are not incident with c (see [4, Theorem 1.3]). Thus, $\#G_{[c]} \leq q$ and c is a translation center exactly if $\#G_{[c]} = q$. Further, if a block B of \mathbb{U} is fixed by $\text{Aut}(\mathbb{U})$, then the center of any translation of \mathbb{U} lies on B .*

In the classical unital, each point is a translation center. In any non-classical $\mathrm{SL}(2, q)$ - \natural -unital and in any $\mathrm{SL}(2, q)$ - \flat -unital of order $q \geq 3$, the block $[\infty]$ is fixed by the full automorphism group (see [6, Proposition 3.11 and Theorem 3.16]) and hence the center of every translation lies on $[\infty]$.

We label the points at infinity of any $\mathrm{SL}(2, q)$ - π -unital with the Sylow p -subgroups, such that each (affine short) block $P \in \mathfrak{P}$ through $\mathbf{1}$ is incident with the point $P \in [\infty]$. We first compute translations of $\mathrm{SL}(2, q)$ - \natural -unitals.

Lemma 3.3. *Let $\mathbb{U}_{S, \mathcal{D}}^{\natural}$ be an $\mathrm{SL}(2, q)$ - \natural -unital. For each $P \in \mathfrak{P}$, the point $P \in [\infty]$ is a translation center with $G_{[P]} = R_P := \{\rho_x \mid x \in P\}$.*

Proof. For each $P \in \mathfrak{P}$, the set of blocks through the point P in $\mathbb{U}_{S, \mathcal{D}}^{\natural}$ equals $\{[\infty]\} \cup \{gP \mid g \in \mathrm{SL}(2, q)\}$. Hence, R_P obviously is a group of translations of $\mathbb{U}_{S, \mathcal{D}}^{\natural}$ with center P . Since $\#R_P = q$, the statement follows (recall Remark 3.2). \square

If $\mathbb{U}_{S, \mathcal{D}}^{\natural}$ is not classical, then the translations given in Lemma 3.3 are all translations of $\mathbb{U}_{S, \mathcal{D}}^{\natural}$. We need some preparation to compute all possible translations of $\mathrm{SL}(2, q)$ - π -unitals with center on $[\infty]$ and $\pi \in \{b, \pi_{\mathrm{odd}}, \pi_{\mathrm{sq}}, \pi_{\mathrm{sq}}^{-1}\}$ (Theorem 3.7).

Lemma 3.4. *Let $P \in \mathfrak{P}$ and let $\tau \in \mathfrak{A} \times R$ such that $P \cdot \tau = P$. Then $\tau = \alpha\rho_x$ with $\alpha \in \mathfrak{A}_P$ and $x \in P$.*

Proof. Let $\tau = \alpha \cdot \rho_h \in \mathfrak{A} \times R$. Then $P = P \cdot \tau = (P \cdot \alpha)h$ and thus α stabilizes P and we have $h \in P$. \square

Let in the following $T := \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{F}_q \right\}$ be a fixed Sylow p -subgroup of $\mathrm{SL}(2, q)$.

Lemma 3.5. *Let $M := \{Tg \mid g \in N(T)\}$ and $f := \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix}$ with $c \in \mathbb{F}_q^{\times}$.*

- (a) *Let $\tau \in \mathfrak{A} \times R$ such that $X \cdot \tau = X$ for all $X \in M$. Then $\tau = \gamma_a \rho_t$ for some $a \in N_{\mathrm{GL}(2, q)}(T)$ and $t \in T$.*
- (b) *Let $\tau \in \mathfrak{A} \times R$ such that $X \cdot \tau = X$ for all $X \in M \cup \{Tf\}$. Then $\tau = \gamma_{t^{-1}} \rho_t$ for some $t \in T$.*

Proof. (a) Since $T \in M$, we have $\tau = \alpha\rho_t$ with $\alpha \in \mathfrak{A}_T \cong \mathrm{P}(N_{\mathrm{GL}(2, q)}(T)) \rtimes \mathrm{Aut}(\mathbb{F}_q)$ and $t \in T$, according to Lemma 3.4. Note first that right multiplication with any element $t \in T$ fixes each coset $Tg = gT \in M$. If $g = \begin{pmatrix} b & * \\ 0 & b^{-1} \end{pmatrix} \in N(T)$, the coset Tg equals $\left\{ \begin{pmatrix} b & x \\ 0 & b^{-1} \end{pmatrix} \mid x \in \mathbb{F}_q \right\}$. Let $[a] \in \mathrm{P}(N_{\mathrm{GL}(2, q)}(T))$. Then we may choose $a = \begin{pmatrix} 1 & y \\ 0 & a \end{pmatrix}$ and for each Tg with $g \in N(T)$, we have $(Tg) \cdot \gamma_a = (Tg)^a = Tg$. Applying a power φ^l of the Frobenius automorphism φ stabilizes the block $Tg = \left\{ \begin{pmatrix} b & x \\ 0 & b^{-1} \end{pmatrix} \mid x \in \mathbb{F}_q \right\}$ exactly if $b^{(p^l)} = b$. Thus, if $\tau = \alpha\rho_t$ stabilizes each block in M , then $\alpha = \gamma_a$ with $a \in N_{\mathrm{GL}(2, q)}(T)$.

- (b) We already know $\tau = \gamma_a \rho_t$ for some $a \in \mathrm{N}_{\mathrm{GL}(2,q)}(T)$ and $t \in T$. We need to show $\gamma_a = \gamma_{t^{-1}}$. Since a normalizes T , we have $(Tf) \cdot \gamma_a \rho_t = (Tf)^a t = Tf^a t$. Hence, $(Tf) \cdot \tau = Tf$ exactly if $f^a t f^{-1} \in T$. Let $t := \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$ and choose without restriction $a = \begin{pmatrix} 1 & y \\ 0 & d \end{pmatrix}$. Then

$$(0, 1) \cdot f^a t f^{-1} = (c(\frac{1-c(x+y)}{d} - 1), 1 + \frac{c(x+y)}{d}).$$

Since $f^a t f^{-1}$ is in T and $c \neq 0$, we get $y = -x$ and $d = 1$ and hence $a = t^{-1}$. \square

If π is a parallelism on \mathfrak{S} and $A \in \pi$ a parallel class, we denote

$$\mathcal{T}_A^\pi := \{\tau \in \mathfrak{A} \times R \mid \forall X \in A : X \cdot \tau = X\}.$$

If $B \in \mathfrak{S}$ is a short block, we denote by $[B]$ the parallel class of π containing B .

Lemma 3.6. *For any $P \in \mathfrak{P}$, we have:*

- (a) $\mathcal{T}_{[P]}^\flat = \{\gamma_{x^{-1}} \rho_x \mid x \in P\}$.
(b) $\mathcal{T}_{[P]}^\pi$ is trivial for $\pi \in \{\pi_{\mathrm{odd}}, \pi_{\mathrm{sq}}, \pi_{\mathrm{sq}}^{-1}\}$.

Proof. Let $\pi \in \{b, \pi_{\mathrm{odd}}, \pi_{\mathrm{sq}}, \pi_{\mathrm{sq}}^{-1}\}$. Following the notation in Lemma 3.5, the parallel class $[T] \in \pi$ contains M and at least one coset Tf with $f = \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix}$ and $c \in \mathbb{F}_q^\times$. According to Lemma 3.5, we thus get $\mathcal{T}_{[T]}^\pi \subseteq \{\gamma_{t^{-1}} \rho_t \mid t \in T\}$. Note that $\gamma_{t^{-1}} \rho_t$ equals left multiplication with t and that $tgT = gT$ exactly if $g \in \mathrm{N}(T)$ or $t = \mathbf{1}$. The parallel class $[T] \in b$ equals the set of right cosets of T , while in π_{odd} , π_{sq} , and π_{sq}^{-1} , the parallel class $[T]$ contains at least one left coset gT , respectively, with $g \notin \mathrm{N}(T)$. Hence, we get $\mathcal{T}_{[T]}^\flat = \{\gamma_{t^{-1}} \rho_t \mid t \in T\}$ and $\mathcal{T}_{[T]}^\pi$ is trivial if $\pi \in \{\pi_{\mathrm{odd}}, \pi_{\mathrm{sq}}, \pi_{\mathrm{sq}}^{-1}\}$.

For any $\pi \in \{b, \pi_{\mathrm{odd}}, \pi_{\mathrm{sq}}, \pi_{\mathrm{sq}}^{-1}\}$, the stabilizer of π in $\mathfrak{A} \times R$ acts transitively on the parallel classes of π . Hence, statement (b) follows. If $\pi = b$, then each parallel class $[T^h]$ equals $[T]^h$ and we have $(Tg)^h \cdot \tau = (Tg)^h$ exactly if $\tau = \gamma_{t^{-h}} \rho_{t^h}$ for some $t \in T$. \square

Theorem 3.7. *Let $\mathbb{U} := \mathbb{U}_{\mathfrak{S}, \mathcal{D}}^\pi$ be an $\mathrm{SL}(2, q)$ - π -unital.*

- (a) *If $\pi = b$ and $q \geq 3$, then:*
- (i) *If $p = 2$, then every non-trivial translation of \mathbb{U} is given by left multiplication with an involution contained in $\mathrm{N}(S)$. For each Sylow 2-subgroup $P \in \mathfrak{P}$, the normalizer $\mathrm{N}(S)$ contains exactly one non-trivial element of P .*
 - (ii) *If q is odd, then \mathbb{U} does not admit any non-trivial translation.*
- (b) *If $\pi \in \{\pi_{\mathrm{odd}}, \pi_{\mathrm{sq}}, \pi_{\mathrm{sq}}^{-1}\}$, then \mathbb{U} admits no non-trivial translation with center on $[\infty]$.*

Proof. The translations of \mathbb{U} with center on the block $[\infty]$ are automorphisms of \mathbb{U} stabilizing $[\infty]$ and are hence contained in $\mathfrak{A}_S \times R$. Thus, statement (b) follows directly from Lemma 3.6.

Now let $\pi = b$ and $q \geq 3$. Then the block $[\infty]$ is fixed by the full automorphism group of \mathbb{U} (see [6, Theorem 3.16]) and hence the center of every translation of \mathbb{U} lies on $[\infty]$ (recall Remark 3.2). Fix $P \in \mathfrak{P}$. According to Lemma 3.6, we have

$$\Gamma_{[P]} \leq \mathcal{T}_{[P]}^\flat \cap (\mathfrak{A}_S \times R) = \{\gamma_{x^{-1}\rho_x} \mid x \in P \text{ such that } \gamma_x \in \mathfrak{A}_S\}.$$

Thus, every possible translation of \mathbb{U} with center P is given by left multiplication with $x \in P \cap N(S)$. The possible isomorphism types of S and its stabilizer in $\text{Aut}(\text{SL}(2, q))$ are given in [7, Proposition 2.5 and Theorem 2.11]. According to those, we get that $x = 1$ or $\text{ord}(x) = p$ divides $2(q+1)$ and hence $p = 2$.

If $p = 2$, then S is cyclic (see e. g. [8, Proposition 2.2] or [7, Proposition 2.5]) and the normalizer $N(S)$ is a dihedral group of order $2(q+1)$ containing S as normal subgroup of order $q+1$. There are thus $q+1$ involutions in $N(S)$, contained in one coset of S . For any $P \in \mathfrak{P}$, the intersection of S and P is trivial and hence no two non-trivial elements of P are contained in $N(S)$. \square

Corollary 3.8. *For q even, the unital $\mathbb{U}_{C, \mathcal{H}}^\flat$ admits exactly $q+1$ non-trivial translations, each of order 2. These translations generate a dihedral group of order $2(q+1)$. \square*

For the Leonids unitals, all translations can explicitly be computed (e. g. with GAP).

Proposition 3.9 ([7], Proposition 6.17). *All translations of the Leonids unitals are known:*

- (i) $\mathbb{U}_{C, \mathcal{H}}^{\pi_2}$ and $\mathbb{U}_{C, \mathcal{E}}^{\pi_2}$ admit exactly one non-trivial translation each, of order 2.
- (ii) $\mathbb{U}_{C, \mathcal{H}}^{\pi_4}$, $\mathbb{U}_{C, \mathcal{H}}^{\pi_5}$, $\mathbb{U}_{C, \mathcal{E}}^{\pi_3}$, $\mathbb{U}_{C, \mathcal{E}}^{\pi_4}$ and $\mathbb{U}_{C, \mathcal{E}}^{\pi_5}$ admit exactly three non-trivial translations each. These translations, respectively, have a common center (which is thus a translation center) and generate an elementary abelian group of order 4. \square

Acknowledgment. The author wishes to warmly thank her thesis advisor Markus J. Stroppel for his highly valuable support in each phase of this research.

References

- [1] *GAP – Groups, Algorithms, and Programming, Version 4.8.10.* The GAP Group, Jan. 2018. URL: <https://www.gap-system.org>.
- [2] Theo Grundhöfer, Markus J. Stroppel, and Hendrik Van Maldeghem. *Moufang sets generated by translations in unitals.* 2020. arXiv: [2008.11445](https://arxiv.org/abs/2008.11445) [[math.GR](https://arxiv.org/abs/2008.11445)].

- [3] Theo Grundhöfer, Markus J. Stroppel, and Hendrik Van Maldeghem. “A non-classical unital of order four with many translations”. In: *Discrete Math.* 339.12 (2016), pp. 2987–2993. DOI: [10/dftk](https://doi.org/10/dftk).
- [4] Theo Grundhöfer, Markus J. Stroppel, and Hendrik Van Maldeghem. “Unitals admitting all translations”. In: *J. Combin. Des.* 21.10 (2013), pp. 419–431. DOI: [10/dftm](https://doi.org/10/dftm).
- [5] Dávid Mezőfi and Gábor P. Nagy. *New Steiner systems from old ones by paramodifications*. 2020. arXiv: [2003.09233](https://arxiv.org/abs/2003.09233) [[math.CO](https://arxiv.org/abs/2003.09233)].
- [6] Verena Möhler. *Automorphisms of (Affine) $SL(2,q)$ -Unitals*. 2020. arXiv: [2012.10116](https://arxiv.org/abs/2012.10116) [[math.CO](https://arxiv.org/abs/2012.10116)].
- [7] Verena Möhler. “ $SL(2,q)$ -Unitals”. PhD thesis. Karlsruher Institut für Technologie (KIT), 2020. DOI: [10/d27c](https://doi.org/10/d27c).
- [8] Verena Möhler. *Three Affine $SL(2,8)$ -Unitals*. 2020. arXiv: [2012.10134](https://arxiv.org/abs/2012.10134) [[math.CO](https://arxiv.org/abs/2012.10134)].