

# Three Affine $SL(2, 8)$ -Unitals

Verena Möhler

December 17, 2020

## Abstract

$SL(2, q)$ -unitals are unitals of order  $q$  admitting a regular action of  $SL(2, q)$  on the complement of some block. We introduce three non-classical affine  $SL(2, 8)$ -unitals and their full automorphism groups. Each of those three affine unitals can be completed to at least two non-isomorphic unitals, leading to six pairwise non-isomorphic unitals of order 8.

**2020 MSC:** 51E26, 51A10, 05B30

**Keywords:** design, unital, affine unital, non-classical unital, automorphism

Most of the results in the present paper have been obtained in the author's Ph.D. thesis [6], where detailed arguments can be found for some statements that we leave to the reader here.

## 1 Preliminaries

One strategy to construct projective planes is to build an affine plane first and then to add *points at infinity*, namely a new point for each parallel class and a line containing all these new points. This strategy of constructing an affine part of a geometry first and then completing it by adding some objects at infinity can successfully be applied to other incidence structures than affine and projective planes. We apply such an approach to unitals.

A **unital of order  $n$**  is a  $2-(n^3 + 1, n + 1, 1)$  design, i. e. an incidence structure with  $n^3 + 1$  points,  $n + 1$  points on each block and unique joining blocks for any two points. We consider *affine unitals*, which arise from unitals by removing one block (and all the points on it) and can be completed to unitals via a parallelism on the short blocks. We give an axiomatic description:

**Definition 1.1.** Let  $n \in \mathbb{N}$ ,  $n \geq 2$ . An incidence structure  $\mathbb{U} = (\mathcal{P}, \mathcal{B}, I)$  is called an **affine unital of order  $n$**  if:

- (AU1) There are  $n^3 - n$  points.
- (AU2) Each block is incident with either  $n$  or  $n + 1$  points. The blocks incident with  $n$  points will be called **short blocks** and the blocks incident with  $n + 1$  points will be called **long blocks**.
- (AU3) Each point is incident with  $n^2$  blocks.
- (AU4) For any two points there is exactly one block incident with both of them.
- (AU5) There exists a **parallelism** on the short blocks, meaning a partition of the set of all short blocks into  $n + 1$  parallel classes of size  $n^2 - 1$  such that the blocks of each parallel class are pairwise non-intersecting.

The existence of a parallelism as in (AU5) must explicitly be required (see [6, Example 3.10]). An affine unital  $\mathbb{U}$  of order  $n$  with parallelism  $\pi$  can be completed to a unital  $\mathbb{U}^\pi$  of order  $n$  as follows: For each parallel class, add a new point that is incident with each short block of that class. Then add a single new block  $[\infty]^\pi$ , incident with the  $n + 1$  new points (see [6, Proposition 3.9]). We call  $\mathbb{U}^\pi$  the  **$\pi$ -closure** of  $\mathbb{U}$ . Note that the closure depends on the parallelism, which need not be unique. Given an affine unital  $\mathbb{U}$  with parallelisms  $\pi$  and  $\pi'$ , the closures  $\mathbb{U}^\pi$  and  $\mathbb{U}^{\pi'}$  are isomorphic with  $[\infty]^\pi \mapsto [\infty]^{\pi'}$  exactly if there is an automorphism of  $\mathbb{U}$  which maps  $\pi$  to  $\pi'$  (see [6, Proposition 3.12]).

## 2 Affine $\mathrm{SL}(2, q)$ -Unitals

From now on let  $p$  be a prime and  $q := p^e$  a  $p$ -power. We are interested in a special kind of affine unitals, namely affine  $\mathrm{SL}(2, q)$ -unitals. The construction of those affine unitals is due to Grundhöfer, Stroppel and Van Maldeghem [2]. They consider translations of unitals, i. e. automorphisms fixing each block through a given point (the so-called center). Of special interest are unitals of order  $q$  where two points are centers of translation groups of order  $q$ . In the classical (Hermitian) unital of order  $q$ , any two such translation groups generate a group isomorphic to  $\mathrm{SL}(2, q)$ ; see [1, Main Theorem] for further possibilities. The construction of (affine)  $\mathrm{SL}(2, q)$ -unitals is motivated by this action of  $\mathrm{SL}(2, q)$  on the classical unital.

Let  $S \leq \mathrm{SL}(2, q)$  be a subgroup of order  $q + 1$  and let  $T \leq \mathrm{SL}(2, q)$  be a Sylow  $p$ -subgroup. Recall that  $T$  has order  $q$  (and thus trivial intersection with  $S$ ), that any two conjugates  $T^h := h^{-1}Th$ ,  $h \in \mathrm{SL}(2, q)$ , have trivial intersection unless they coincide and that there are  $q + 1$  conjugates of  $T$ .

Consider a collection  $\mathcal{D}$  of subsets of  $\mathrm{SL}(2, q)$  such that each set  $D \in \mathcal{D}$  contains  $\mathbf{1} := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ , that  $\#D = q + 1$  for each  $D \in \mathcal{D}$ , and the following properties hold:

(Q) For each  $D \in \mathcal{D}$ , the map

$$(D \times D) \setminus \{(x, x) \mid x \in D\} \rightarrow \mathrm{SL}(2, q), \quad (x, y) \mapsto xy^{-1},$$

is injective, i. e. the set  $D^* := \{xy^{-1} \mid x, y \in D, x \neq y\}$  contains  $q(q+1)$  elements.

(P) The system consisting of  $S \setminus \{1\}$ , all conjugates of  $T \setminus \{1\}$  and all sets  $D^*$  with  $D \in \mathcal{D}$  forms a partition of  $\mathrm{SL}(2, q) \setminus \{1\}$ .

Set

$$\mathcal{P} := \mathrm{SL}(2, q),$$

$$\mathcal{B} := \{Sg \mid g \in \mathrm{SL}(2, q)\} \cup \{T^h g \mid h, g \in \mathrm{SL}(2, q)\} \cup \{Dg \mid D \in \mathcal{D}, g \in \mathrm{SL}(2, q)\}$$

and let the incidence relation  $I \subseteq \mathcal{P} \times \mathcal{B}$  be containment.

Then we call the incidence structure  $\mathbb{U}_{S, \mathcal{D}} := (\mathcal{P}, \mathcal{B}, I)$  an **affine  $\mathrm{SL}(2, q)$ -unital**. Each affine  $\mathrm{SL}(2, q)$ -unital is indeed an affine unital of order  $q$ , see [6, Prop. 3.15]. We call the sets  $\hat{D} := \{Dd^{-1} \mid d \in D\}$ ,  $D \in \mathcal{D}$ , the **hats** of  $\mathbb{U}_{S, \mathcal{D}}$  and the blocks  $Dg$ ,  $D \in \mathcal{D}$  and  $g \in \mathrm{SL}(2, q)$ , the **arcuate blocks** of  $\mathbb{U}_{S, \mathcal{D}}$ .

For the construction of an affine  $\mathrm{SL}(2, q)$ -unital, we have to choose a subgroup  $S \leq \mathrm{SL}(2, q)$  of order  $q+1$  and find a set  $\mathcal{D}$  of arcuate blocks through  $1$  such that (Q) and (P) hold.

**Example 2.1.**

- (a) For each prime power  $q$  we may choose  $S = C$  to be cyclic and  $\mathcal{H}$  a set of arcuate blocks through  $1$  such that  $\mathbb{U}_{C, \mathcal{H}}$  is isomorphic to the affine part of the classical unital. We call  $\mathbb{U}_{C, \mathcal{H}}$  the **classical affine  $\mathrm{SL}(2, q)$ -unital**. See [2, Example 3.1] or [6, Section 3.2.2] for details.
- (b) In [2], Grundhöfer, Stroppel and Van Maldeghem introduce a non-classical affine  $\mathrm{SL}(2, 4)$ -unital.

**Proposition 2.2.** *Let  $p = 2$  and let  $S \leq \mathrm{SL}(2, q)$  be a subgroup of order  $q+1$ . Then  $S$  is cyclic and unique up to conjugation.*

*Proof.* For  $p = 2$ , we have  $\mathrm{SL}(2, q) \cong \mathrm{PSL}(2, q)$ . Using Dickson's list of subgroups of  $\mathrm{PSL}(2, q)$  (see e. g. [3, Hauptsatz II.8.27]), we see that each subgroup of order  $q+1$  is cyclic. From [3, Satz II.8.5], we get that there is exactly one conjugacy class of cyclic subgroups of  $\mathrm{PSL}(2, q)$  of order  $q+1$ .  $\square$

**Remark 2.3.** *In [6, Proposition 2.5], we give a complete list of possible subgroups  $S \leq \mathrm{SL}(2, q)$  of order  $q+1$ . For  $p \not\equiv 3 \pmod{4}$ , the group  $S$  is cyclic. For  $p \equiv 3 \pmod{4}$ ,  $S$  is cyclic or generalized quaternion and there is one exceptional case for  $q = 23$  and one for  $q = 47$ .*

For each prime power  $q$ , we may choose a cyclic subgroup  $C \leq \mathrm{SL}(2, q)$  of order  $q + 1$  as given in the following

**Remark 2.4.** *Let  $d \in \mathbb{F}_q^\times$  such that  $X^2 - tX + d$  has no root in  $\mathbb{F}_q$ , where  $t = 1$  if  $q$  is even and  $t = 0$  if  $q$  is odd. Then*

$$C := \left\{ \begin{pmatrix} a & b \\ -db & a+tb \end{pmatrix} \mid a^2 + tab + db^2 = 1 \right\}$$

*is a cyclic subgroup of  $\mathrm{SL}(2, q)$  of order  $q + 1$ . Note that  $C$  is the norm 1 group of the quadratic extension field*

$$\mathbb{F}_{q^2} := \left\{ \begin{pmatrix} a & b \\ -db & a+tb \end{pmatrix} \mid a, b \in \mathbb{F}_q \right\}.$$

We take a brief look on automorphisms of affine  $\mathrm{SL}(2, q)$ -unitals, i. e. bijections of the point set such that the block set is invariant. On any affine  $\mathrm{SL}(2, q)$ -unital  $\mathbb{U}_{S, \mathcal{D}}$ , right multiplication with elements of  $\mathrm{SL}(2, q)$  obviously induces automorphisms. Let  $R := \{\rho_h \mid h \in \mathrm{SL}(2, q)\} \leq \mathrm{Aut}(\mathbb{U}_{S, \mathcal{D}})$ , where  $\rho_h \in R$  acts on  $\mathbb{U}_{S, \mathcal{D}}$  by right multiplication with  $h \in \mathrm{SL}(2, q)$ . Every automorphism of  $\mathrm{SL}(2, q)$  obviously induces a bijection of the point set of  $\mathbb{U}_{S, \mathcal{D}}$ , but it need not leave the block set invariant. Let  $\mathfrak{A}$  denote the permutation group given by all automorphisms of  $\mathrm{SL}(2, q)$ .

We import a useful statement from [4]:

**Theorem 2.5** ([4], Theorem 3.3). *Let  $q \geq 3$  and let  $\mathbb{U}_{S, \mathcal{D}}$  and  $\mathbb{U}_{S', \mathcal{D}'}$  be affine  $\mathrm{SL}(2, q)$ -unitals.*

(a) *Let  $\psi: \mathbb{U}_{S, \mathcal{D}} \rightarrow \mathbb{U}_{S', \mathcal{D}'}$  be an isomorphism. Then  $\psi = \alpha\rho_h$  with  $\rho_h \in R$  and  $\alpha \in \mathfrak{A}$  such that  $S \cdot \alpha = S'$ .*

(b)  $\mathrm{Aut}(\mathbb{U}_{S, \mathcal{D}}) \leq \mathfrak{A}_S \times R$ . □

**Remark 2.6.** *The classical affine  $\mathrm{SL}(2, q)$ -unital  $\mathbb{U}_{C, \mathcal{H}}$  admits the whole group  $\mathfrak{A}_C \times R$  as automorphism group (see [6, Proposition 4.6]). Hence,  $\mathfrak{A}_S \times R$  is a sharp upper bound for the automorphism group of any affine  $\mathrm{SL}(2, q)$ -unital  $\mathbb{U}_{S, \mathcal{D}}$  of order  $q \geq 3$ .*

### 3 Three Affine $\mathrm{SL}(2, 8)$ -Unitals

Let  $q = 8$  and  $\mathbb{F}_8^\times = \langle z \rangle$ , with  $z^3 = z + 1$ . The polynomial  $X^2 + X + 1$  has no root in  $\mathbb{F}_8$  and the Frobenius automorphism

$$\varphi: \mathbb{F}_8 \rightarrow \mathbb{F}_8, \quad x \mapsto x^2,$$

has order 3. Since  $q = 8$  is even, any subgroup  $S \leq \mathrm{SL}(2, 8)$  of order 9 is cyclic and we may hence choose

$$S := C = \left\{ \begin{pmatrix} a & b \\ b & a+b \end{pmatrix} \mid a, b \in \mathbb{F}_8, a^2 + ab + b^2 = 1 \right\}.$$

A generator of  $C$  is given by  $g := \begin{pmatrix} z^2 & z^4 \\ z^4 & z \end{pmatrix}$ . Let  $f := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ . Then

$$\mathfrak{A}_C = \text{Aut}(\text{SL}(2, 8))_C = \langle \gamma_g \rangle \rtimes \langle \gamma_f \cdot \varphi \rangle \cong C_9 \rtimes C_6,$$

where  $\varphi$  acts entrywise on a matrix and  $\gamma_x$  describes conjugation with  $x$ . Representatives of the conjugacy classes of minimal subgroups of  $\mathfrak{A}_C$  are

$$F := \langle \gamma_f \rangle \cong \langle \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \rangle \cong C_2,$$

$$U := \langle \gamma_{g^3} \rangle \cong \langle \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \rangle \cong C_3 \text{ and}$$

$$L := \langle \varphi \rangle \cong C_3.$$

**Example 3.1** (The classical affine unital of order 8). Let

$$H_1 := \left\{ \mathbf{1}, \begin{pmatrix} z^5 & 1 \\ z^5 & z^6 \end{pmatrix}, \begin{pmatrix} z^4 & z^2 \\ 1 & z^2 \end{pmatrix}, \begin{pmatrix} 0 & z \\ z^6 & z^5 \end{pmatrix}, \begin{pmatrix} z^3 & z^6 \\ z^4 & z^5 \end{pmatrix}, \begin{pmatrix} z^3 & z \\ z^6 & 0 \end{pmatrix}, \begin{pmatrix} 1 & z^2 \\ 1 & z^6 \end{pmatrix}, \begin{pmatrix} z^4 & 1 \\ z^5 & 1 \end{pmatrix}, \begin{pmatrix} z^5 & 0 \\ 0 & z^2 \end{pmatrix} \right\},$$

$$H_2 := H_1 \cdot \varphi, H_3 := H_1 \cdot \varphi^2,$$

$$H_4 := \left\{ \mathbf{1}, \begin{pmatrix} z^5 & 0 \\ z^6 & z^2 \end{pmatrix}, \begin{pmatrix} z & z^6 \\ z^4 & 1 \end{pmatrix}, \begin{pmatrix} 1 & z^5 \\ z^6 & z^5 \end{pmatrix}, \begin{pmatrix} z & z^4 \\ 0 & z^6 \end{pmatrix}, \begin{pmatrix} z^5 & z^2 \\ z^4 & z^4 \end{pmatrix}, \begin{pmatrix} 0 & z^2 \\ z^5 & z^5 \end{pmatrix}, \begin{pmatrix} 0 & z^4 \\ z^3 & z^4 \end{pmatrix}, \begin{pmatrix} z^2 & z^5 \\ z^5 & z^6 \end{pmatrix} \right\},$$

$$H_5 := H_4 \cdot \varphi, H_6 := H_4 \cdot \varphi^2$$

and  $\mathcal{H} := \{H_1, \dots, H_6\}$ . Then  $\mathbb{U}_{C, \mathcal{H}}$  is the classical affine unital of order 8. Recall that for  $H \in \mathcal{H}$ , we denote by  $\hat{H}$  the set of arcuate blocks  $\{Hh^{-1} \mid h \in H\}$ . As indicated,  $\varphi$  acts on the set of hats  $\{\hat{H} \mid H \in \mathcal{H}\}$  in two orbits of length 3. Conjugation by  $g$  stabilizes each  $\hat{H}$  and acts transitively on the blocks of each  $\hat{H}$ . Conjugation by  $f$  also stabilizes each  $\hat{H}$  but fixes exactly one block per  $\hat{H}$ .

**Theorem 3.2** (Weihnachtsunital). *Let  $C := \langle g \rangle = \langle \begin{pmatrix} z^2 & z^4 \\ z^4 & z \end{pmatrix} \rangle$  as above and let*

$$D_1 := \left\{ \mathbf{1}, \begin{pmatrix} z^5 & 1 \\ z^5 & z^6 \end{pmatrix}, \begin{pmatrix} z^4 & z^2 \\ 1 & z^2 \end{pmatrix}, \begin{pmatrix} 0 & z \\ z^6 & z^2 \end{pmatrix}, \begin{pmatrix} 1 & z^4 \\ z^2 & z^2 \end{pmatrix}, \begin{pmatrix} 1 & z \\ z^6 & 0 \end{pmatrix}, \begin{pmatrix} 1 & z^2 \\ 1 & z^6 \end{pmatrix}, \begin{pmatrix} z^4 & 1 \\ z^5 & 1 \end{pmatrix}, \begin{pmatrix} z^5 & 0 \\ 0 & z^2 \end{pmatrix} \right\},$$

$$D_2 := D_1 \cdot \varphi, D_3 := D_1 \cdot \varphi^2,$$

$$D_4 := \left\{ \mathbf{1}, \begin{pmatrix} z^5 & 0 \\ z^6 & z^2 \end{pmatrix}, \begin{pmatrix} z & z^6 \\ z^4 & 1 \end{pmatrix}, \begin{pmatrix} 0 & z \\ z^6 & z^5 \end{pmatrix}, \begin{pmatrix} z^4 & 0 \\ z^2 & z^3 \end{pmatrix}, \begin{pmatrix} z^5 & z^2 \\ z^3 & z^6 \end{pmatrix}, \begin{pmatrix} 0 & z^2 \\ z^5 & z^5 \end{pmatrix}, \begin{pmatrix} 0 & z^4 \\ z^3 & z^4 \end{pmatrix}, \begin{pmatrix} z^2 & z^5 \\ z^5 & z^6 \end{pmatrix} \right\},$$

$$D_5 := D_4 \cdot \varphi, D_6 := D_4 \cdot \varphi^2$$

and  $\mathcal{D} := \{D_1, \dots, D_6\}$ . Then  $\mathbb{WU} := \mathbb{U}_{C, \mathcal{D}}$  is an affine  $\text{SL}(2, 8)$ -unital and we call it **Weihnachtsunital**<sup>1</sup>. The stabilizer of  $\mathbf{1}$  in  $\text{Aut}(\mathbb{WU})$  is

$$\text{Aut}(\mathbb{WU})_{\mathbf{1}} = U \rtimes (F \times L) = \langle \gamma_{g^3} \rangle \rtimes \langle \gamma_f \cdot \varphi \rangle \cong C_3 \rtimes C_6$$

and the full automorphism group


$$\text{Aut}(\mathbb{WU}) = \text{Aut}(\mathbb{WU})_{\mathbf{1}} \rtimes R$$

has index 3 in  $\text{Aut}(\mathbb{U}_{C, \mathcal{H}}) = \mathfrak{A}_C \rtimes R$ .

<sup>1</sup> The Weihnachtsunital was discovered around Christmas 2017, whence the name.

*Proof.* The proof is basically computation (recall Theorem 2.5). Note that the given description already uses the automorphism  $\varphi \in \text{Aut}(\mathbb{WU})_{\mathbb{1}}$ . Conjugation by  $f$  stabilizes each hat with exactly one fixed block per hat. Conjugation by the generator  $g$  of  $C$  does not induce an automorphism of  $\mathbb{WU}$ , but conjugation by  $g^3$  yields an automorphism of  $\mathbb{WU}$  such that each hat is fixed.  $\square$


Having computed the full automorphism group of  $\text{Aut}(\mathbb{WU})$ , we know in particular that the Weihnachtsunital is not isomorphic to the classical affine  $\text{SL}(2, 8)$ -unital  $\mathbb{U}_{C, \mathcal{H}}$ . Another way to see that  $\mathbb{WU}$  is not isomorphic to  $\mathbb{U}_{C, \mathcal{H}}$  is via O’Nan configurations. An O’Nan configuration consists of four distinct blocks meeting in six distinct points:



O’Nan observed that classical unitals do not contain such configurations (see [7, p. 507]).

**Remark 3.3.** *In  $\mathbb{WU}$ , there are lots of O’Nan configurations, e. g.*

$$\begin{aligned}
 C &= \{\mathbb{1}, g, g^2, g^3, g^4, g^5, g^6, g^7, g^8\}, \\
 T &:= \{\mathbb{1}, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & z^2 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & z^3 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & z^4 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & z^5 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & z^6 \\ 0 & 1 \end{pmatrix}\}, \\
 D_2 \cdot \begin{pmatrix} z^4 & 1 \\ 1 & 0 \end{pmatrix} &= \left\{ \begin{pmatrix} z^4 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & z^3 \\ z^4 & z^3 \end{pmatrix}, \begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} z^2 & 0 \\ z & z^5 \end{pmatrix}, \begin{pmatrix} z^2 & 1 \\ z^2 & z^4 \end{pmatrix}, \begin{pmatrix} z & 1 \\ z^2 & z^5 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} z^4 & z \\ 0 & z^3 \end{pmatrix}, \begin{pmatrix} 1 & z^3 \\ z^4 & 0 \end{pmatrix} \right\}, \\
 D_3 \cdot \begin{pmatrix} z & z^3 \\ 0 & z^6 \end{pmatrix} &= \left\{ \begin{pmatrix} z & z^3 \\ 0 & z^6 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} z^3 & z^4 \\ z & z \end{pmatrix}, \begin{pmatrix} 0 & z^3 \\ z^4 & z^2 \end{pmatrix}, \begin{pmatrix} z & 1 \\ z^2 & z^5 \end{pmatrix}, \begin{pmatrix} z & 0 \\ z^4 & z^6 \end{pmatrix}, \begin{pmatrix} z & z \\ z & z^5 \end{pmatrix}, \begin{pmatrix} z^3 & z \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & z^2 \\ 0 & 1 \end{pmatrix} \right\}.
 \end{aligned}$$



**Theorem 3.4** (Osterunital and Pfingstunital<sup>2</sup>). *Let  $C := \langle g \rangle = \langle \begin{pmatrix} z^2 & z^4 \\ z^4 & z \end{pmatrix} \rangle$  as above.*

(a) *Let*

$$\begin{aligned} D_1 &:= \left\{ \mathbf{1}, \begin{pmatrix} z^5 & 1 \\ z^5 & z^6 \end{pmatrix}, \begin{pmatrix} z^4 & z^2 \\ 1 & z^2 \end{pmatrix}, \begin{pmatrix} 1 & z \\ z^6 & 0 \end{pmatrix}, \begin{pmatrix} 0 & z \\ z^6 & z^2 \end{pmatrix}, \begin{pmatrix} 1 & z^4 \\ z^2 & z^2 \end{pmatrix}, \begin{pmatrix} z^3 & z^5 \\ z^3 & 1 \end{pmatrix}, \begin{pmatrix} z^5 & z^4 \\ z^2 & z^4 \end{pmatrix}, \begin{pmatrix} z^2 & 0 \\ 0 & z^5 \end{pmatrix} \right\}, \\ D_2 &:= D_1^g, D_3 := D_1^{g^2}, \\ D_4 &:= \left\{ \mathbf{1}, \begin{pmatrix} z^5 & 0 \\ z^6 & z^2 \end{pmatrix}, \begin{pmatrix} z & z^6 \\ z^4 & 1 \end{pmatrix}, \begin{pmatrix} z^5 & z^2 \\ z^5 & 0 \end{pmatrix}, \begin{pmatrix} z^3 & z^4 \\ z^6 & z^5 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ z^3 & z \end{pmatrix}, \begin{pmatrix} 1 & z \\ 1 & z^3 \end{pmatrix}, \begin{pmatrix} z & z^2 \\ 1 & z^5 \end{pmatrix}, \begin{pmatrix} z & 0 \\ z & z^6 \end{pmatrix} \right\}, \\ D_5 &:= D_4^g, D_6 := D_4^{g^2} \end{aligned}$$

and  $\mathcal{D} := \{D_1, \dots, D_6\}$ . Then  $\mathbb{OU} := \mathbb{U}_{C, \mathcal{D}}$  is an affine  $\mathrm{SL}(2, 8)$ -unital and we call it **Osterunital**.

(b) *Let  $f = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  as above and let*

$$\begin{aligned} D'_1 &:= D_1, D'_2 := D_2, D'_3 := D_3, \\ D'_4 &:= D_4^f, D'_5 := (D'_4)^g, D'_6 := (D'_4)^{g^2} \end{aligned}$$

and  $\mathcal{D}' := \{D'_1, \dots, D'_6\}$ . Then  $\mathbb{PU} := \mathbb{U}_{C, \mathcal{D}'}$  is an affine  $\mathrm{SL}(2, 8)$ -unital and we call it **Pfingstunital**.

We denote by  $C$  also the automorphism group  $C := \langle \gamma_g \rangle \leq \mathfrak{A}_C$ . The full stabilizers of  $\mathbf{1}$  in  $\mathrm{Aut}(\mathbb{OU})$  and  $\mathrm{Aut}(\mathbb{PU})$ , respectively, are

$$\mathrm{Aut}(\mathbb{OU})_{\mathbf{1}} = \mathrm{Aut}(\mathbb{PU})_{\mathbf{1}} = C \rtimes L = \langle \gamma_g \rangle \rtimes \langle \varphi \rangle \cong C_9 \rtimes C_3$$

and the full automorphism groups

$$\mathrm{Aut}(\mathbb{OU}) = \mathrm{Aut}(\mathbb{PU}) = (C \rtimes L) \rtimes R$$

have index 2 in  $\mathrm{Aut}(\mathbb{U}_{C, \mathcal{H}})$ .


*Proof.* Again this is basically computation. The given description already uses the automorphism  $\gamma_g$  in both  $\mathrm{Aut}(\mathbb{OU})_{\mathbf{1}}$  and  $\mathrm{Aut}(\mathbb{PU})_{\mathbf{1}}$ . The Frobenius automorphism  $\varphi$  acts as automorphism on  $\mathbb{OU}$  as well as on  $\mathbb{PU}$  in the same way as it does on  $\mathbb{U}_{C, \mathcal{H}}$  and on  $\mathbb{WU}$ . The orbits of  $\varphi$  in  $\mathcal{D}$  are  $\{D_1, D_2, D_3\}$  and  $\{D_4, D_5, D_6\}$  and its orbits in  $\mathcal{D}'$  are  $\{D'_1, D'_2, D'_3\}$  and  $\{D'_4, D'_5, D'_6\}$ . Conjugation by  $f$  induces no automorphism on neither  $\mathbb{OU}$  nor  $\mathbb{PU}$ .  $\square$

**Remark 3.5.** *Other than in the Weihnachtsunital, there is a difference between the action of  $\mathrm{Aut}(\mathbb{OU})_{\mathbf{1}} = \mathrm{Aut}(\mathbb{PU})_{\mathbf{1}} \leq \mathfrak{A}_C$  on the set of hats of the Oster- and Pfingstunital, respectively, and its action on the set of hats of the classical affine  $\mathrm{SL}(2, 8)$ -unital  $\mathbb{U}_{C, \mathcal{H}}$ . In  $\mathbb{U}_{C, \mathcal{H}}$ , conjugation by  $g$  fixes every hat, while on  $\mathbb{OU}$  and  $\mathbb{PU}$  it acts on the set of hats in two orbits of length 3.*

<sup>2</sup> The Osterunital and the Pfingstunital were discovered in 2018, you might guess the dates.

**Remark 3.6.** *As in the Weihnachtsunital, there are also many O’Nan configurations in  $\mathbb{O}U$  and  $\mathbb{P}U$ , e. g.*

$$\begin{aligned}
C &= \{1, g, g^2, g^3, g^4, g^5, g^6, g^7, g^8\}, \\
D_1 &= \left\{1, \begin{pmatrix} z^5 & 1 \\ z^5 & z^6 \end{pmatrix}, \begin{pmatrix} z^4 & z^2 \\ 1 & z^2 \end{pmatrix}, \begin{pmatrix} 1 & z \\ z^6 & 0 \end{pmatrix}, \begin{pmatrix} 0 & z \\ z^6 & z^2 \end{pmatrix}, \begin{pmatrix} 1 & z^4 \\ z^2 & z^2 \end{pmatrix}, \begin{pmatrix} z^3 & z^5 \\ z^3 & 1 \end{pmatrix}, \begin{pmatrix} z^5 & z^4 \\ z^2 & z^4 \end{pmatrix}, \begin{pmatrix} z^2 & 0 \\ 0 & z^5 \end{pmatrix}\right\}, \\
D_2 \cdot g &= \left\{g, \begin{pmatrix} z^4 & z^2 \\ 1 & z^2 \end{pmatrix}, \begin{pmatrix} z^3 & 0 \\ z^3 & z^4 \end{pmatrix}, \begin{pmatrix} z^5 & z^5 \\ z^3 & z^5 \end{pmatrix}, \begin{pmatrix} z^6 & 0 \\ 1 & z \end{pmatrix}, \begin{pmatrix} z^6 & z \\ z^3 & z^6 \end{pmatrix}, \begin{pmatrix} z^5 & z^2 \\ z^6 & z^5 \end{pmatrix}, \begin{pmatrix} 1 & z^3 \\ z^4 & 0 \end{pmatrix}, \begin{pmatrix} z^2 & z^3 \\ 0 & z^5 \end{pmatrix}\right\}, \\
D_3 \cdot \begin{pmatrix} z^5 & z^2 \\ z^6 & z^5 \end{pmatrix} &= \left\{\begin{pmatrix} z^5 & z^2 \\ z^6 & z^5 \end{pmatrix}, \begin{pmatrix} z^2 & 0 \\ 0 & z^5 \end{pmatrix}, \begin{pmatrix} z & z^2 \\ z^3 & z^3 \end{pmatrix}, \begin{pmatrix} z^2 & 1 \\ z^6 & 1 \end{pmatrix}, \begin{pmatrix} z^6 & z \\ z^5 & z^3 \end{pmatrix}, \begin{pmatrix} z^4 & z^3 \\ z^4 & 0 \end{pmatrix}, \begin{pmatrix} z & z^4 \\ z^4 & z^2 \end{pmatrix}, \begin{pmatrix} z^3 & z^2 \\ z^5 & 0 \end{pmatrix}, \begin{pmatrix} z^3 & z \\ z^4 & z \end{pmatrix}\right\}.
\end{aligned}$$



Although they look quite similar, the Osterunital and the Pfingstunital are not isomorphic, as is shown in the following

**Proposition 3.7.** *There is no isomorphism between  $\mathbb{O}U$  and  $\mathbb{P}U$ .*

*Proof.* According to Theorem 2.5, any isomorphism between  $\mathbb{O}U$  and  $\mathbb{P}U$  must be contained in  $\mathfrak{A}_C \times R$ . But since the index of  $\text{Aut}(\mathbb{O}U)$  in  $\mathfrak{A}_C \times R$  equals 2 and computation shows that  $D_1^f$  is no block of  $\mathbb{P}U$ , the statement follows.  $\square$

In particular, the Oster- and Pfingstunital are two non-isomorphic affine  $\text{SL}(2, q)$ -unitals with the same full automorphism group.

**Remark 3.8.** *The Weihnachts-, Oster- and Pfingstunital were found by a computer search. In fact, we did an exhaustive search for affine  $\text{SL}(2, 8)$ -unitals, where the groups  $F$ ,  $U$  and  $L$  act in the same way as on the classical affine  $\text{SL}(2, 8)$ -unital. Those three affine unitals were the only ones appearing through the search. See [6, Chapter 6] for details about the search.*



## 4 Completion to Unitals

Any affine unital can be completed to a unital by each of its parallelisms. In any affine  $\mathrm{SL}(2, q)$ -unital, the set of short blocks is the set of all right cosets of the  $q + 1$  Sylow  $p$ -subgroups of  $\mathrm{SL}(2, q)$ . Note that each right coset  $Tg$  is a left coset  $gT^g$  of a conjugate of  $T$ . A parallelism as in (AU5) means a partition of the set of short blocks into  $q + 1$  sets of  $q^2 - 1$  pairwise non-intersecting cosets. For each prime power  $q$ , there are hence two obvious parallelisms, namely partitioning the set of short blocks into the sets of *right* cosets or into the sets of *left* cosets of the Sylow  $p$ -subgroups. We name those two parallelisms “flat” and “natural”, respectively, and denote them by the corresponding musical signs

$$\flat := \{\{Tg \mid g \in \mathrm{SL}(2, q)\} \mid T \in \mathfrak{P}\} \quad \text{and} \quad \natural := \{\{gT \mid g \in \mathrm{SL}(2, q)\} \mid T \in \mathfrak{P}\},$$

where  $\mathfrak{P}$  denotes the set of Sylow  $p$ -subgroups of  $\mathrm{SL}(2, q)$ .

Given an affine  $\mathrm{SL}(2, q)$ -unital  $\mathbb{U}_{S, \mathcal{D}}$  with parallelism  $\pi$ , we call the  $\pi$ -closure an  **$\mathrm{SL}(2, q)$ - $(\pi)$ -unital**. Completing  $\mathbb{WU}$ ,  $\mathbb{OU}$  and  $\mathbb{PU}$  with  $\flat$  and  $\natural$  each, we obtain six pairwise non-isomorphic  $\mathrm{SL}(2, q)$ -unitals of order 8. Since they are all  $\flat$ - or  $\natural$ -closures of non-classical affine  $\mathrm{SL}(2, q)$ -unitals of order  $8 \geq 3$ , we know from [4, Proposition 3.11 and Theorem 3.16] that their full automorphism groups fix the block  $[\infty]$ . Since the parallelisms  $\flat$  and  $\natural$ , respectively, are preserved under the action of  $\mathfrak{A} \times R$ , we get

$$\mathrm{Aut}(\mathbb{U}^\pi) = \mathrm{Aut}(\mathbb{U}^\pi)_{[\infty]} = \mathrm{Aut}(\mathbb{U})$$

for any  $\mathbb{U} \in \{\mathbb{WU}, \mathbb{OU}, \mathbb{PU}\}$  and  $\pi \in \{\flat, \natural\}$ .

**Remark 4.1.** *In any  $\mathrm{SL}(2, q)$ - $\natural$ -unital, the Sylow  $p$ -subgroups act (via right multiplication) as translation groups of order  $q$  with centers on the block  $[\infty]$ . Hence,  $\mathbb{WU}^\natural$ ,  $\mathbb{OU}^\natural$  and  $\mathbb{PU}^\natural$  are examples of non-classical unitals of order  $q$  where the translations generate  $\mathrm{SL}(2, q)$ .*

**Remark 4.2.** *There might be more parallelisms on the short blocks of  $\mathrm{SL}(2, 8)$ -unitals, leading to further closures. We already know a class of parallelisms for each odd order and one for square order (described in [5, Sections 2.1 and 2.2]) and some parallelisms for order 4, leading to 12 new  $\mathrm{SL}(2, 4)$ -unitals, the so-called Leonids unitals (see [5, Section 2.3] and [6, Section 6.2.2]).*

**Acknowledgment.** The author wishes to warmly thank her thesis advisor Markus J. Stroppel for his highly valuable support in each phase of this research.

## References

- [1] Theo Grundhöfer, Markus J. Stroppel, and Hendrik Van Maldeghem. *Moufang sets generated by translations in unitals*. 2020. arXiv: [2008.11445](https://arxiv.org/abs/2008.11445) [[math.GR](#)].

- [2] Theo Grundhöfer, Markus J. Stroppel, and Hendrik Van Maldeghem. “A non-classical unital of order four with many translations”. In: *Discrete Math.* 339.12 (2016), pp. 2987–2993. DOI: [10/dftk](https://doi.org/10/dftk).
- [3] Bertram Huppert. *Endliche Gruppen. I.* Die Grundlehren der Mathematischen Wissenschaften, Band 134. Springer-Verlag, Berlin-New York, 1967.
- [4] Verena Möhler. *Automorphisms of (Affine)  $SL(2,q)$ -Unitals*. 2020. arXiv: [2012.10116](https://arxiv.org/abs/2012.10116) [[math.CO](https://arxiv.org/abs/2012.10116)].
- [5] Verena Möhler. *Parallelisms and Translations of (Affine)  $SL(2,q)$ -Unitals*. 2020. arXiv: [2012.10129](https://arxiv.org/abs/2012.10129) [[math.CO](https://arxiv.org/abs/2012.10129)].
- [6] Verena Möhler. “ $SL(2,q)$ -Unitals”. PhD thesis. Karlsruher Institut für Technologie (KIT), 2020. DOI: [10/d27c](https://doi.org/10/d27c).
- [7] Michael E. O’Nan. “Automorphisms of unitary block designs”. In: *J. Algebra* 20 (1972), pp. 495–511. DOI: [10/br4k6c](https://doi.org/10/br4k6c).