

Determinant of a Sum of Certain Kronecker Products

Dwight Nwaigwe

March 3, 2022

Abstract

We compute the determinant of $\sum_{n=1}^N \mathbf{A}^{(n)} \otimes \mathbf{B}^{(n)}$, where $\mathbf{A}^{(n)}$ is square and $\mathbf{B}^{(n)} = \mathbf{x}^{(n)} \mathbf{y}^{(n)T}$ where $\mathbf{x}^{(n)}$ and $\mathbf{y}^{(n)}$ have length N .

Introduction

Kronecker products appear in a variety of applications [1], [2]. For square matrices \mathbf{A}, \mathbf{B} the determinant of the Kronecker product $\mathbf{A} \otimes \mathbf{B}$ is given by $\det(\mathbf{A})^N \det(\mathbf{B})^F$, where F is the size of \mathbf{A} and N the size of \mathbf{B} [3]. Typically there are no general formulas for more complicated expressions involving Kronecker products, such as sums. In this article we compute the determinant of a sum of certain Kronecker products as described in the next section.

Result

lemma. Let $\{\mathbf{A}^{(n)}\}$ be F by F matrices and $\mathbf{B}^{(n)} = \mathbf{x}^{(n)} \mathbf{y}^{(n)T}$ where $\mathbf{x}^{(n)}$ is the n^{th} column of some N by N matrix \mathbf{X} , and $\mathbf{y}^{(n)}$ is the n^{th} column of some N by N matrix \mathbf{Y} . Also, let

$$\mathbf{G} = \sum_{n=1}^N \mathbf{A}^{(n)} \otimes \mathbf{B}^{(n)}. \quad (1)$$

Then, the following equation holds:

$$\det(\mathbf{G}) = \left(\prod_{n=1}^N \det(\mathbf{A}^{(n)}) \right) \det(\mathbf{X})^F \det(\mathbf{Y})^F. \quad (2)$$

Proof. $\det(\mathbf{G})$ is a multi-linear function of its columns, therefore it can be expressed a sum of determinants where each determinant corresponds to a particular arrangement of columns belonging to the various $\mathbf{A}^{(n)} \otimes \mathbf{B}^{(n)}$. Denote the set of such corresponding matrices as \mathcal{O} . Then $\det(\mathbf{G}) = \sum_{\mathbf{C} \in \mathcal{O}} \det(\mathbf{C})$. Because $\mathbf{B}^{(n)}$ is rank 1, only certain matrices contribute to the sum and as a result we can redefine \mathcal{O} to consist of these matrices. Let $\mathbf{C} \in \mathcal{O}$. The column entries of \mathbf{C} are such that: each of the first N columns come from a different $\mathbf{A}^{(n)} \otimes \mathbf{B}^{(n)}$; each of the columns $N + 1$ to $2N$ come from a different $\mathbf{A}^{(n)} \otimes \mathbf{B}^{(n)}$; and so forth for the $(N - 1)F + 1$ to NF columns. For this reason, let S_N be the permutation group on N elements, and let $(\gamma^1, \gamma^2, \dots, \gamma^F) \in (S_N)^F$. Every $\mathbf{C} \in \mathcal{O}$ can be identified by some $(\gamma^1, \gamma^2, \dots, \gamma^F)$. By definition of γ^k , we have that $\mathbf{A}^{\gamma^k(i)} = \mathbf{A}^{(n)}$ for some i , and n , and $\mathbf{B}^{\gamma^k(i)} = \mathbf{B}^{(n)}$ for some i and n . It is possible that $(\gamma^1, \gamma^2, \dots, \gamma^F)$ may not have distinct elements. Also, we now write $\mathbf{C}^{(\gamma^1, \gamma^2, \dots, \gamma^F)}$ instead of \mathbf{C} to emphasize the identification of $\mathbf{C} \in \mathcal{O}$ with $(S_N)^F$.

For a given $\mathbf{C}^{(\gamma^1, \gamma^2, \dots, \gamma^F)} \in \mathcal{O}$, it is sufficient to only consider the case of a block diagonal structure since contributions from the non-block diagonal terms in $\mathbf{C}^{(\gamma^1, \gamma^2, \dots, \gamma^F)}$ can be expressed as contributions from a block diagonal matrix up to sign. For the purpose of the proof, we thus only calculate contributions from the

Clearly,

$$\det(\mathbf{Z}) = \det(\mathbf{X}) \operatorname{sgn}(\gamma^i) \quad (9)$$

which combined with (8) gives

$$\prod_{i=1}^F \det[\mathbf{B}_1^{\gamma^i(1)} \mathbf{B}_2^{\gamma^i(2)} \dots \mathbf{B}_N^{\gamma^i(N)}] = (\det(\mathbf{X}))^F \prod_{i=1}^F (\mathbf{Y}_{1\gamma^i(1)} \mathbf{Y}_{2\gamma^i(2)} \dots \mathbf{Y}_{N\gamma^i(N)}) \operatorname{sgn}(\gamma^i). \quad (10)$$

Plugging in (10) into (3) gives

$$\det(\mathbf{C}_{diag}^{(\gamma^1, \gamma^2, \dots, \gamma^F)}) \quad (11)$$

$$= (\mathbf{A}_{11}^1 \mathbf{A}_{11}^2 \dots \mathbf{A}_{11}^F) (\mathbf{A}_{22}^1 \mathbf{A}_{22}^2 \dots \mathbf{A}_{22}^F) \dots (\mathbf{A}_{FF}^1 \mathbf{A}_{FF}^2 \dots \mathbf{A}_{FF}^F) (\det(\mathbf{X}))^F.$$

$$\prod_{i=1}^F (\mathbf{Y}_{1\gamma^i(1)} \mathbf{Y}_{2\gamma^i(2)} \dots \mathbf{Y}_{N\gamma^i(N)}) \operatorname{sgn}(\gamma^i). \quad (12)$$

To get the contributions from all block-diagonal matrices, we must sum over all possible sequences of $(\gamma^1, \gamma^2, \dots, \gamma^F) \in (S_N)^F$. Doing this we have

$$\begin{aligned} \sum_{(\gamma^1, \gamma^2, \dots, \gamma^F) \in (S_N)^F} \det(\mathbf{C}_{diag}^{(\gamma^1, \gamma^2, \dots, \gamma^F)}) \quad (13) \\ = (\mathbf{A}_{11}^1 \mathbf{A}_{11}^2 \dots \mathbf{A}_{11}^F) (\mathbf{A}_{22}^1 \mathbf{A}_{22}^2 \dots \mathbf{A}_{22}^F) \dots (\mathbf{A}_{FF}^1 \mathbf{A}_{FF}^2 \dots \mathbf{A}_{FF}^F) (\det(\mathbf{X}))^F \cdot \\ \left(\sum_{(\gamma^1, \gamma^2, \dots, \gamma^F) \in (S_N)^F} \prod_{i=1}^F (\mathbf{Y}_{1\gamma^i(1)} \mathbf{Y}_{2\gamma^i(2)} \dots \mathbf{Y}_{N\gamma^i(N)}) \operatorname{sgn}(\gamma^i) \right). \end{aligned}$$

However, the term $\left(\sum_{(\gamma^1, \gamma^2, \dots, \gamma^F) \in (S_N)^F} \prod_{i=1}^F (\mathbf{Y}_{1\gamma^i(1)} \mathbf{Y}_{2\gamma^i(2)} \dots \mathbf{Y}_{N\gamma^i(N)}) \operatorname{sgn}(\gamma^i) \right)$ is equal to $(\det(\mathbf{Y}))^F$, so that (13) becomes

$$\begin{aligned} \sum_{(\gamma^1, \gamma^2, \dots, \gamma^F) \in (S_N)^F} \det(\mathbf{C}_{diag}^{(\gamma^1, \gamma^2, \dots, \gamma^F)}) \quad (14) \\ = (\mathbf{A}_{11}^1 \mathbf{A}_{11}^2 \dots \mathbf{A}_{11}^F) (\mathbf{A}_{22}^1 \mathbf{A}_{22}^2 \dots \mathbf{A}_{22}^F) \dots (\mathbf{A}_{FF}^1 \mathbf{A}_{FF}^2 \dots \mathbf{A}_{FF}^F) (\det(\mathbf{X}))^F (\det(\mathbf{Y}))^F \end{aligned}$$

Adding the contributions from the non-block diagonal terms to (14), (2) is proved. \square

References

- [1] Peter Benner, Pawan Goyal, Boris Kramer, Benjamin Peherstorfe, and Karen Willcox. Operator inference for non-intrusive model reduction of systems with non-polynomial nonlinear terms. *Computational Methods in Applied Mechancis and Engineering*, 2020.
- [2] Frederik Benzing, Marcelo Matheus Gauy, Asier Mujika, Anders Martinsson, and Angelika Steger. Optimal kronecker-sum approximation of real time recurrent learning. *Proceedings of Machine Learning Research*, 97, 2019.
- [3] Roger Horn and Charles Johnson. *Topics in Matrix Analysis*. 2001.