

# KDV SOLVES BKP

ALEXANDER ALEXANDROV

ABSTRACT. In this note, we prove that any tau-function of the KdV hierarchy also solves the BKP hierarchy after a simple rescaling of times.

Keywords: tau-functions, BKP hierarchy, KdV hierarchy, Hirota bilinear identity

In this paper we answer the question about the relation between the KdV and BKP hierarchies, raised in [Ale20]. Namely, in [Ale20] it was observed that several families of the KdV tau-functions important in enumerative geometry and theoretical physics also solve the BKP hierarchy after a simple rescaling of times. Here we prove that *any* tau-function of KdV solves the BKP hierarchy.

In terms of tau-function  $\tau_{KP}(\mathbf{t})$  the KP hierarchy, introduced in [SS83], is described by the Hirota bilinear identity

$$(1) \quad \oint_{\infty} e^{\xi(\mathbf{t}-\mathbf{t}',z)} \tau_{KP}(\mathbf{t} - [z^{-1}]) \tau_{KP}(\mathbf{t}' + [z^{-1}]) dz = 0.$$

This bilinear identity encodes all nonlinear equations of the KP hierarchy. Here we use the standard short-hand notations

$$(2) \quad \mathbf{t} \pm [z^{-1}] := \{t_1 \pm z^{-1}, t_2 \pm \frac{1}{2}z^{-2}, t_3 \pm \frac{1}{3}z^{-3}, \dots\}$$

and

$$(3) \quad \xi(\mathbf{t}, z) = \sum_{k>0} t_k z^k.$$

If a tau-function of the KP hierarchy does not depend on even time variables,

$$(4) \quad \frac{\partial}{\partial t_{2k}} \tau_{KdV}(\mathbf{t}) = 0 \quad \forall k > 0,$$

than it is a tau-function of the KdV hierarchy. Since we still have arbitrary  $t_{2k}$  and  $t'_{2k}$  in the first factor of the integrand in (1), for the KdV hierarchy the Hirota bilinear identity is given by

$$(5) \quad \oint_{\infty} e^{\xi^o(\mathbf{t}-\mathbf{t}',z)} f(z) \tau_{KdV}(\mathbf{t} - [z^{-1}]) \tau_{KdV}(\mathbf{t}' + [z^{-1}]) dz = 0$$

for arbitrary series  $f(z) \in \mathbb{C}[[z^2]]$ . Here we introduce

$$(6) \quad \xi^o(\mathbf{t}, z) = \sum_{k \in \mathbb{Z}_{\text{odd}}^+} t_k z^k.$$

In terms of the *Hirota derivatives*

$$(7) \quad P(D_1, D_2, \dots) f \cdot g = P\left(\frac{\partial}{\partial y_1}, \frac{\partial}{\partial y_2}, \dots\right) f(\mathbf{t} + \mathbf{y}) g(\mathbf{t} - \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{0}},$$

the first few equations of the KdV hierarchy can be represented as

$$(8) \quad \begin{aligned} (D_1^4 - 4D_1 D_3) \tau_{KdV} \cdot \tau_{KdV} &= 0, \\ (D_1^6 + 4D_1^3 D_3 - 32D_3^2) \tau_{KdV} \cdot \tau_{KdV} &= 0, \\ (D_1^6 - 20D_1^3 D_3 - 80D_3^2 + 144D_1 D_5) \tau_{KdV} \cdot \tau_{KdV} &= 0. \end{aligned}$$

The BKP hierarchy was introduced by Date, Jimbo, Kashiwara and Miwa in [DJKM82\*]. It can be represented in terms of tau-function  $\tau_{BKP}(\mathbf{t})$  by the Hirota bilinear identity similar to (1):

$$(9) \quad \frac{1}{2\pi i} \oint_{\infty} e^{\xi^o(\mathbf{t}-\mathbf{t}',z)} \tau_{BKP}(\mathbf{t} - 2[z^{-1}]) \tau_{BKP}(\mathbf{t}' + 2[z^{-1}]) \frac{dz}{z} = \tau_{BKP}(\mathbf{t}) \tau_{BKP}(\mathbf{t}').$$

The first equation of the BKP hierarchy in terms of the Hirota derivatives is given by

$$(10) \quad (D_1^6 - 5D_1^3 D_3 - 5D_3^2 + 9D_1 D_5) \tau_{BKP} \cdot \tau_{BKP} = 0.$$

Note, that if we map  $D_k \mapsto D_k/2$  then this equation coincides with the last equation in (8). Let us show that this is more than just a coincidence, and all equations of the BKP hierarchy follow from the KdV hierarchy. The main result of this note is

**Theorem 1.** *For any KdV tau-function*

$$(11) \quad \tau_{BKP}(\mathbf{t}) = \tau_{KdV}(\mathbf{t}/2)$$

*is a tau-function of the BKP hierarchy.*

*Proof.* Let us consider

$$(12) \quad f(z) = \frac{e^{\xi^\circ(\mathbf{t}-\mathbf{t}',z)} - e^{-\xi^\circ(\mathbf{t}-\mathbf{t}',z)}}{z} \in \mathbb{C}[[z^2]],$$

then the Hirota bilinear identity (5) for the KdV hierarchy leads to

$$(13) \quad \frac{1}{2\pi i} \oint_{\infty} e^{2\xi^\circ(\mathbf{t}-\mathbf{t}',z)} \tau_{KdV}(\mathbf{t} - [z^{-1}]) \tau_{KdV}(\mathbf{t}' + [z^{-1}]) \frac{dz}{z} = \tau_{KdV}(\mathbf{t}) \tau_{KdV}(\mathbf{t}').$$

Now we substitute the times  $\mathbf{t} \mapsto \mathbf{t}/2$ ,  $\mathbf{t}' \mapsto \mathbf{t}'/2$ :

$$(14) \quad \frac{1}{2\pi i} \oint_{\infty} e^{\xi^\circ(\mathbf{t}-\mathbf{t}',z)} \tau_{KdV}(\mathbf{t}/2 - [z^{-1}]) \tau_{KdV}(\mathbf{t}'/2 + [z^{-1}]) \frac{dz}{z} = \tau_{KdV}(\mathbf{t}/2) \tau_{KdV}(\mathbf{t}'/2).$$

This equation coincides with the Hirota bilinear identity for the BKP hierarchy (9) if we identify tau-functions by (11). This completes the proof.  $\square$

It is easy to see that the converse statement is false, that is, the KdV hierarchy is a certain reduction of the BKP hierarchy. This reduction, as well as similar description of the higher Gelfand–Dickey hierarchies, will be discussed elsewhere.

This theorem partially explains the results of [Ale20, MM20]. Indeed, in [MM20] it was noted that the Kontsevich–Witten tau-function - one of the most important solutions of the KdV hierarchy - has a simple expansion in terms of the Schur Q-functions. This result was generalized in [Ale20] to a family of KdV tau-functions related to the Brézin–Gross–Witten model. On the basis of these expansions the author has conjectured (for the Kontsevich–Witten tau-function) and proved (for the Brézin–Gross–Witten tau-function) that these KdV tau-functions also solve the BKP hierarchy. From Theorem 1 it follows, that this part of the conjectures in [Ale20] trivially follows from a general relation between the KdV and BKP hierarchies. Moreover, this theorem makes the Schur Q-functions a natural basis for expansion of the KdV tau-functions.

There are several different ways to relate KdV and BKP hierarchies. In particular, in [DJKM82] the authors describe an identification of KdV hierarchy with the 4-reduction of BKP, see Remark on page 1098. Another reduction from BKP to KdV (so-called 1-constrained BKP) was described in [Orl91, Orl93, Che92]. To the best of our understanding, these reductions do not coincide with the one considered in this paper.

#### ACKNOWLEDGMENTS

The author is grateful to J. van de Leur, A. Mironov, A. Morozov, and A. Orlov for useful discussions. This work was supported by IBS-R003-D1.

#### REFERENCES

- [Ale20] Alexander Alexandrov. Intersection numbers on  $\overline{\mathcal{M}}_{g,n}$  and BKP hierarchy. *arXiv e-prints*, page arXiv:2012.07573, December 2020.
- [Che92] Yi Cheng. Constraints of the Kadomtsev–Petviashvili hierarchy. *J. Math. Phys.*, 33(11):3774–3782, 1992.
- [DJKM82] Etsuro Date, Michio Jimbo, Masaki Kashiwara, and Tetsuji Miwa. Transformation groups for soliton equations. Euclidean Lie algebras and reduction of the KP hierarchy. *Publ. Res. Inst. Math. Sci.*, 18(3):1077–1110, 1982.
- [DJKM82\*] Etsuro Date, Michio Jimbo, Masaki Kashiwara, and Tetsuji Miwa. Transformation groups for soliton equations. IV. A new hierarchy of soliton equations of KP-type. *Phys. D*, 4(3):343–365, 1981/82.
- [MM20] A. Mironov and A. Morozov. Superintegrability of Kontsevich matrix model. arXiv:2011.12917.
- [Orl91] A. Yu. Orlov. Symmetries for unifying different soliton systems into a single integrable hierarchy. preprint IINS/Oce-04/03, 1991.
- [Orl93] A. Yu. Orlov. Volterra Operator Algebra for Zero Curvature Representation. University of KP. In A. S. Fokas, D. J. Kaup, A. C. Newell, and V. E. Zakharov, editors, *Nonlinear Processes in Physics*, page 126, January 1993.
- [SS83] Mikio Sato and Yasuko Sato. Soliton equations as dynamical systems on infinite-dimensional Grassmann manifold, 1983.

IBS CENTER FOR GEOMETRY AND PHYSICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY (POSTECH), 77 CHEONGAM-RO, NAM-GU, POHANG, GYEONGBUK, 37673, KOREA

*Email address:* alexandrovsash at gmail.com