

KDV SOLVES BKP

ALEXANDER ALEXANDROV

ABSTRACT. In this note, we prove that any tau-function of the KdV hierarchy also solves the BKP hierarchy after a simple rescaling of times.

Keywords: tau-functions, BKP hierarchy, KdV hierarchy, Hirota bilinear identity

In this paper we answer the question about the relation between the KdV and BKP hierarchies, raised in [Ale20]. Namely, in [Ale20] it was observed that several families of the KdV tau-functions important in enumerative geometry and theoretical physics also solve the BKP hierarchy after a simple rescaling of times. Here we prove that *any* tau-function of KdV solves the BKP hierarchy.

In terms of tau-function $\tau_{KP}(\mathbf{t})$ the KP hierarchy, introduced in [SS83], is described by the Hirota bilinear identity

$$(1) \quad \oint_{\infty} e^{\xi(\mathbf{t}-\mathbf{t}',z)} \tau_{KP}(\mathbf{t} - [z^{-1}]) \tau_{KP}(\mathbf{t}' + [z^{-1}]) dz = 0.$$

This bilinear identity encodes all nonlinear equations of the KP hierarchy. Here we use the standard short-hand notations

$$(2) \quad \mathbf{t} \pm [z^{-1}] := \{t_1 \pm z^{-1}, t_2 \pm \frac{1}{2}z^{-2}, t_3 \pm \frac{1}{3}z^{-3}, \dots\}$$

and

$$(3) \quad \xi(\mathbf{t}, z) = \sum_{k>0} t_k z^k.$$

If a tau-function of the KP hierarchy does not depend on even time variables,

$$(4) \quad \frac{\partial}{\partial t_{2k}} \tau_{KdV}(\mathbf{t}) = 0 \quad \forall k > 0,$$

than it is a tau-function of the KdV hierarchy. Since we still have arbitrary t_{2k} and t'_{2k} in the first factor of the integrand in (1), for the KdV hierarchy the Hirota bilinear identity is given by

$$(5) \quad \oint_{\infty} e^{\xi^o(\mathbf{t}-\mathbf{t}',z)} f(z) \tau_{KdV}(\mathbf{t} - [z^{-1}]) \tau_{KdV}(\mathbf{t}' + [z^{-1}]) dz = 0$$

for arbitrary series $f(z) \in \mathbb{C}[[z^2]]$. Here we introduce

$$(6) \quad \xi^o(\mathbf{t}, z) = \sum_{k \in \mathbb{Z}_{\text{odd}}^+} t_k z^k.$$

In terms of the *Hirota derivatives*

$$(7) \quad P(D_1, D_2, \dots) f \cdot g = P\left(\frac{\partial}{\partial y_1}, \frac{\partial}{\partial y_2}, \dots\right) f(\mathbf{t} + \mathbf{y}) g(\mathbf{t} - \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{0}},$$

the first few equations of the KdV hierarchy can be represented as

$$(8) \quad \begin{aligned} (D_1^4 - 4D_1D_3) \tau_{KdV} \cdot \tau_{KdV} &= 0, \\ (D_1^6 + 4D_1^3D_3 - 32D_3^2) \tau_{KdV} \cdot \tau_{KdV} &= 0, \\ (D_1^6 - 20D_1^3D_3 - 80D_3^2 + 144D_1D_5) \tau_{KdV} \cdot \tau_{KdV} &= 0. \end{aligned}$$

The BKP hierarchy was introduced by Date, Jimbo, Kashiwara and Miwa in [DJKM82*]. It can be represented in terms of tau-function $\tau_{BKP}(\mathbf{t})$ by the Hirota bilinear identity similar to (1):

$$(9) \quad \frac{1}{2\pi i} \oint_{\infty} e^{\xi^o(\mathbf{t}-\mathbf{t}',z)} \tau_{BKP}(\mathbf{t} - 2[z^{-1}]) \tau_{BKP}(\mathbf{t}' + 2[z^{-1}]) \frac{dz}{z} = \tau_{BKP}(\mathbf{t}) \tau_{BKP}(\mathbf{t}').$$

The first equation of the BKP hierarchy in terms of the Hirota derivatives is given by

$$(10) \quad (D_1^6 - 5D_1^3D_3 - 5D_3^2 + 9D_1D_5) \tau_{BKP} \cdot \tau_{BKP} = 0.$$

Note, that if we map $D_k \mapsto D_k/2$ then this equation coincides with the last equation in (8). Let us show that this is more than just a coincidence, and all equations of the BKP hierarchy follow from the KdV hierarchy. The main result of this note is

Theorem 1. *For any KdV tau-function*

$$(11) \quad \tau_{BKP}(\mathbf{t}) = \tau_{KdV}(\mathbf{t}/2)$$

is a tau-function of the BKP hierarchy.

Proof. Let us consider

$$(12) \quad f(z) = \frac{e^{\xi^\circ(\mathbf{t}-\mathbf{t}',z)} - e^{-\xi^\circ(\mathbf{t}-\mathbf{t}',z)}}{z} \in \mathbb{C}[[z^2]],$$

then the Hirota bilinear identity (5) for the KdV hierarchy leads to

$$(13) \quad \frac{1}{2\pi i} \oint_{\infty} e^{2\xi^\circ(\mathbf{t}-\mathbf{t}',z)} \tau_{KdV}(\mathbf{t} - [z^{-1}]) \tau_{KdV}(\mathbf{t}' + [z^{-1}]) \frac{dz}{z} = \tau_{KdV}(\mathbf{t}) \tau_{KdV}(\mathbf{t}').$$

Now we substitute the times $\mathbf{t} \mapsto \mathbf{t}/2$, $\mathbf{t}' \mapsto \mathbf{t}'/2$:

$$(14) \quad \frac{1}{2\pi i} \oint_{\infty} e^{\xi^\circ(\mathbf{t}-\mathbf{t}',z)} \tau_{KdV}(\mathbf{t}/2 - [z^{-1}]) \tau_{KdV}(\mathbf{t}'/2 + [z^{-1}]) \frac{dz}{z} = \tau_{KdV}(\mathbf{t}/2) \tau_{KdV}(\mathbf{t}'/2).$$

This equation coincides with the Hirota bilinear identity for the BKP hierarchy (9) if we identify tau-functions by (11). This completes the proof. \square

It is easy to see that the converse statement is false, that is, the KdV hierarchy is a certain reduction of the BKP hierarchy. This reduction, as well as similar description of the higher Gelfand–Dickey hierarchies, will be discussed elsewhere.

This theorem partially explains the results of [Ale20, MM20]. Indeed, in [MM20] it was noted that the Kontsevich–Witten tau-function - one of the most important solutions of the KdV hierarchy - has a simple expansion in terms of the Schur Q-functions. This result was generalized in [Ale20] to a family of KdV tau-functions related to the Brézin–Gross–Witten model. On the basis of these expansions the author has conjectured (for the Kontsevich–Witten tau-function) and proved (for the Brézin–Gross–Witten tau-function) that these KdV tau-functions also solve the BKP hierarchy. From Theorem 1 it follows, that this part of the conjectures in [Ale20] trivially follows from a general relation between the KdV and BKP hierarchies. Moreover, this theorem makes the Schur Q-functions a natural basis for expansion of the KdV tau-functions.

There are several different ways to relate KdV and BKP hierarchies. In particular, in [DJKM82] the authors describe an identification of KdV hierarchy with the 4-reduction of BKP, see Remark on page 1098. Another reduction from BKP to KdV (so-called 1-constrained BKP) was described in [Orl91, Orl93, Che92]. To the best of our understanding, these reductions do not coincide with the one considered in this paper.

ACKNOWLEDGMENTS

The author is grateful to J. van de Leur, A. Mironov, A. Morozov, and A. Orlov for useful discussions. This work was supported by IBS-R003-D1.

REFERENCES

- [Ale20] Alexander Alexandrov. Intersection numbers on $\overline{\mathcal{M}}_{g,n}$ and BKP hierarchy. *arXiv e-prints*, page arXiv:2012.07573, December 2020.
- [Che92] Yi Cheng. Constraints of the Kadomtsev–Petviashvili hierarchy. *J. Math. Phys.*, 33(11):3774–3782, 1992.
- [DJKM82] Etsuro Date, Michio Jimbo, Masaki Kashiwara, and Tetsuji Miwa. Transformation groups for soliton equations. Euclidean Lie algebras and reduction of the KP hierarchy. *Publ. Res. Inst. Math. Sci.*, 18(3):1077–1110, 1982.
- [DJKM82*] Etsuro Date, Michio Jimbo, Masaki Kashiwara, and Tetsuji Miwa. Transformation groups for soliton equations. IV. A new hierarchy of soliton equations of KP-type. *Phys. D*, 4(3):343–365, 1981/82.
- [MM20] A. Mironov and A. Morozov. Superintegrability of Kontsevich matrix model. arXiv:2011.12917.
- [Orl91] A. Yu. Orlov. Symmetries for unifying different soliton systems into a single integrable hierarchy. preprint IINS/Oce-04/03, 1991.
- [Orl93] A. Yu. Orlov. Volterra Operator Algebra for Zero Curvature Representation. University of KP. In A. S. Fokas, D. J. Kaup, A. C. Newell, and V. E. Zakharov, editors, *Nonlinear Processes in Physics*, page 126, January 1993.
- [SS83] Mikio Sato and Yasuko Sato. Soliton equations as dynamical systems on infinite-dimensional Grassmann manifold, 1983.

IBS CENTER FOR GEOMETRY AND PHYSICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY (POSTECH), 77 CHEONGAM-RO, NAM-GU, POHANG, GYEONGBUK, 37673, KOREA

Email address: alexandrovsash at gmail.com