

Graphs cospectral with $\text{NU}(n + 1, q^2)$, $n \neq 3$

Ferdinand Ihringer

Francesco Pavese

Valentino Smaldore

Abstract

Let $\mathcal{H}(n, q^2)$ be a non-degenerate Hermitian variety of $\text{PG}(n, q^2)$, $n \geq 2$. Let $\text{NU}(n+1, q^2)$ be the graph whose vertices are the points of $\text{PG}(n, q^2) \setminus \mathcal{H}(n, q^2)$ and two vertices P_1, P_2 are adjacent if the line joining P_1 and P_2 is tangent to $\mathcal{H}(n, q^2)$. Then $\text{NU}(n+1, q^2)$ is a strongly regular graph. In this paper we show that $\text{NU}(n+1, q^2)$, $n \neq 3$, is not determined by its spectrum.

Keywords: strongly regular graph; unital; Hermitian variety.

1 Introduction

A *strongly regular graph* with parameters (v, k, λ, μ) is a (simple, undirected, connected) graph with v vertices such that each vertex lies on k edges, any two adjacent vertices have exactly λ common neighbours, and any two non-adjacent vertices have exactly μ common neighbours. Let $\mathcal{H}(n, q^2)$ be a non-degenerate Hermitian variety of $\text{PG}(n, q^2)$, $n \geq 2$. Let $\text{NU}(n+1, q^2)$ be the graph whose vertices are the points of $\text{PG}(n, q^2) \setminus \mathcal{H}(n, q^2)$ and two vertices P_1, P_2 are adjacent if the line joining P_1 and P_2 contains exactly one point of $\mathcal{H}(n, q^2)$ (i.e., it is a line tangent to $\mathcal{H}(n, q^2)$). The graph $\text{NU}(n+1, q^2)$ is a strongly regular graph for any q [3, Table 9.9, p. 145]. A variation of Godsil–McKay switching was described by Wang, Qiu, and Hu in [17]. In [13], by using a simplified version of the Wang–Qiu–Hu switching, the authors exhibited a strongly regular graph having the same parameters as $\text{NU}(n+1, 4)$, $n \geq 5$, but not isomorphic to it. Moreover they asked whether or not the graph $\text{NU}(n+1, q^2)$, is determined by its spectrum. In this paper we show that the graph $\text{NU}(n+1, q^2)$, $n \neq 3$, is not determined by its spectrum.

A *unital* \mathcal{U} of a finite projective plane Π of order q^2 is a set of $q^3 + 1$ points such that every line of Π meets \mathcal{U} in either 1 or $q + 1$ points. A Hermitian curve is a unital of $\text{PG}(2, q^2)$, called *classical unital*. Besides the classical one, there are other known unitals in $\text{PG}(2, q^2)$ and they arise from a construction due to Buekenhout, see [1, 2, 14]. Let \mathcal{U} be a unital of $\text{PG}(2, q^2)$ and

F. Ihringer: Department of Mathematics: Analysis, Logic and Discrete Mathematics, Ghent University, Belgium *e-mail:* ferdinand.ihringer@ugent.be

F. Pavese: Dipartimento di Meccanica, Matematica e Management, Politecnico di Bari, Via Orabona 4, 70125 Bari, Italy *e-mail:* francesco.pavese@poliba.it

V. Smaldore: Dipartimento di Matematica, Informatica ed Economia, Università degli Studi della Basilicata, Contrada Macchia Romana, 85100, Potenza, Italy; *e-mail:* valentino.smaldore@unibas.it

Mathematics Subject Classification (2020): Primary 51E20, 05E30; Secondary 05C50.

let $\Gamma_{\mathcal{U}}$ be the graph whose vertices are the points of $\text{PG}(2, q^2) \setminus \mathcal{U}$ and two vertices P_1, P_2 are adjacent if the line joining P_1 and P_2 contains exactly one point of \mathcal{U} . The graph $\Gamma_{\mathcal{U}}$ is strongly regular with the same parameters as $\text{NU}(3, q^2)$. Here we observe the (well-known) fact that if \mathcal{U} is a non-classical unital, then $\Gamma_{\mathcal{U}}$ is not isomorphic to $\text{NU}(3, q^2)$.

In $\text{PG}(n, q^2)$, $n \geq 4$, by applying a simplified version of the Wang–Qiu–Hu switching to the graph $\text{NU}(n+1, q^2)$, we obtain two graphs \mathcal{G}'_n and \mathcal{G}''_n , which are strongly regular and have the same parameters as $\text{NU}(n+1, q^2)$. Moreover we show that \mathcal{G}'_n (if $q > 2$) and \mathcal{G}''_n are not isomorphic to $\text{NU}(n+1, q^2)$.

2 Preliminary results

Let $\text{PG}(n, q^2)$ be the n -dimensional projective space over the finite $\text{GF}(q^2)$ equipped with homogeneous projective coordinates (X_1, \dots, X_{n+1}) . A unitary polarity of $\text{PG}(n, q^2)$ is induced by a non-degenerate Hermitian form on the underlying vector space. A non-degenerate Hermitian variety $\mathcal{H}(n, q^2)$ consists of the absolute points of a unitary polarity of $\text{PG}(n, q^2)$. A non-degenerate Hermitian variety $\mathcal{H}(n, q^2)$ has $(q^{n+1} + (-1)^n)(q^n - (-1)^n) / (q^2 - 1)$ points. Subspaces of maximal dimension contained in $\mathcal{H}(n, q^2)$ are $\lfloor \frac{n-1}{2} \rfloor$ -dimensional projective spaces and are called *generators*. There are either $(q+1)(q^3+1)\dots(q^n+1)$ or $(q^3+1)(q^5+1)\dots(q^{n+1}+1)$ generators in $\mathcal{H}(n, q^2)$, depending on whether n is odd or even, respectively. A line of $\text{PG}(n, q^2)$ meets $\mathcal{H}(n, q^2)$ in 1, $q+1$ or, if $n \geq 3$, in q^2+1 points. The latter lines are the generators of $\mathcal{H}(n, q^2)$ if $n \in \{3, 4\}$; lines meeting $\mathcal{H}(n, q^2)$ in one or $q+1$ points are called *tangent lines* or *secant lines*, respectively. Through a point P of $\mathcal{H}(n, q^2)$ there pass $(q^{n-1} + (-1)^{n-2})(q^{n-2} - (-1)^{n-2}) / (q^2 - 1)$ generators and these generators are contained in a hyperplane. The hyperplane containing these generators is the polar hyperplane of P with respect to the unitary polarity of $\text{PG}(n, q^2)$ defining $\mathcal{H}(n, q^2)$ and it is also called the *tangent hyperplane* to $\mathcal{H}(n, q^2)$ at P . The tangent lines through P are precisely the remaining lines of its polar hyperplane that are incident with P . If $P \notin \mathcal{H}(n, q^2)$ then the polar hyperplane of P is a hyperplane of $\text{PG}(n, q^2)$ meeting $\mathcal{H}(n, q^2)$ in a non-degenerate Hermitian variety $\mathcal{H}(n-1, q^2)$ and it is said to be *secant* to $\mathcal{H}(n, q^2)$. The stabilizer of $\mathcal{H}(n, q^2)$ in $\text{PGL}(n+1, q^2)$ is the group $\text{PGU}(n+1, q^2)$.

For a point P of $\text{PG}(2, q^2)$, its polar line meets $\mathcal{H}(2, q^2)$ exactly in P or in the $q+1$ points of a Baer subline, according as P belongs to $\mathcal{H}(2, q^2)$ or not. It is well known that $\mathcal{H}(2, q^2)$ contains no *O'Nan configuration* [15, p. 507], which is a configuration consisting of four distinct lines intersecting in six distinct points of $\mathcal{H}(2, q^2)$. Dually, $\mathcal{H}(2, q^2)$ cannot contain a *dual O'Nan configuration*, that is a configuration formed by four points of $\text{PG}(2, q^2) \setminus \mathcal{H}(2, q^2)$ no three on a line, such that the six lines connecting two of them are tangent lines. In $\text{PG}(2, q^2)$, a *Hermitian pencil of lines* is a cone having as vertex a point V and as base a Baer subline of a line ℓ , where $V \notin \ell$. See [1] for more details.

A plane of $\text{PG}(4, q^2)$ meets $\mathcal{H}(4, q^2)$ in a line, a Hermitian pencil of lines or a non-degenerate Hermitian curve, according as its polar line has q^2+1 , 1 or $q+1$ points in common with $\mathcal{H}(4, q^2)$, respectively.

Lemma 2.1. *In $\text{PG}(2, q^2)$, let P_1, P_2, P_3 be three points on a line γ tangent to a non-degenerate Hermitian curve $\mathcal{H}(2, q^2)$ at the point T , where $T \neq P_i$, $i = 1, 2, 3$. Let s be the unique Baer subline of γ containing P_1, P_2, P_3 .*

- *If $T \in s$, then there is no point $P \in \text{PG}(2, q^2) \setminus (\mathcal{H}(2, q^2) \cup \gamma)$ such that the lines PP_i , $i = 1, 2, 3$, are tangent to $\mathcal{H}(2, q^2)$.*
- *If $T \notin s$, then there are q points $P \in \text{PG}(2, q^2) \setminus (\mathcal{H}(2, q^2) \cup \gamma)$ such that the lines PP_i , $i = 1, 2, 3$, are tangent to $\mathcal{H}(2, q^2)$.*

Proof. Let \perp be the unitary polarity of $\text{PG}(2, q^2)$ defining $\mathcal{H}(2, q^2)$. Assume first that $T \in s$. Suppose by contradiction that there is a point $P \in \text{PG}(2, q^2) \setminus (\mathcal{H}(2, q^2) \cup \gamma)$ such that the lines PP_i , $i = 1, 2, 3$, are tangent to $\mathcal{H}(2, q^2)$. Then by projecting the $q + 1$ points of $P^\perp \cap \mathcal{H}(2, q^2)$ from P onto γ we obtain a Baer subline of γ containing P_1, P_2, P_3 , i.e., the Baer subline s . Since $T \in s$, we have that the line PT has to be tangent to $\mathcal{H}(2, q^2)$ at T . Hence $PT = \gamma$, contradicting the fact that $P \notin \gamma$.

Assume now that $T \notin s$. By using [9, Theorem 15.3.11] it can be seen that there are $q^2(q - 1)$ Baer sublines in γ such that $T \notin s$, and these are permuted in a single orbit by the stabilizer $\text{PGU}(3, q^2)_\gamma$ of γ in $\text{PGU}(3, q^2)$. On the other hand, $\text{PGU}(3, q^2)_\gamma$ permutes in a unique orbit the $q^3(q - 1)$ points of $\text{PG}(2, q^2) \setminus (\mathcal{H}(2, q^2) \cup \gamma)$, see for instance [15, Lemma 2.7]. Consider the incidence structure whose point set \mathcal{P} consists of the $q^4 - q^3$ points of $\text{PG}(2, q^2) \setminus (\mathcal{H}(2, q^2) \cup \gamma)$ and whose block set \mathcal{L} of the $q^3 - q^2$ Baer sublines of γ , where a point $P \in \mathcal{P}$ is incident with a block $r \in \mathcal{L}$ if r is obtained by projecting the $q + 1$ points of $P^\perp \cap \mathcal{H}(2, q^2)$ from P onto γ . Since both, \mathcal{P} and \mathcal{L} , are orbits of the same group, the considered incidence structure is a tactical configuration [11, Lemma 1.45] and hence we have that a block of \mathcal{L} is incident with q points of \mathcal{P} . In other words, there are q points of $\text{PG}(2, q^2) \setminus (\mathcal{H}(2, q^2) \cup \gamma)$ that are projected onto the same Baer subline of γ , as required. \square

Lemma 2.2. *In $\text{PG}(2, q^2)$, let t, t' be two lines tangent to a non-degenerate Hermitian curve $\mathcal{H}(2, q^2)$ and let $R = t \cap t'$. There is a bijection between the points P of $t' \setminus (\mathcal{H}(2, q^2) \cup \{R\})$ and the Baer sublines s_P of t passing through R , not containing $\mathcal{H}(2, q^2) \cap t$ and such that a line joining P with a point of s_P is tangent to $\mathcal{H}(2, q^2)$.*

Proof. It is enough to note that the stabilizer of t and t' in $\text{PGU}(3, q^2)$ acts transitively on the $q^2 - 1$ points of $t' \setminus (\mathcal{H}(2, q^2) \cup \{R\})$ and the $q^2 - 1$ Baer sublines of t through R not containing $\mathcal{H}(2, q^2) \cap t$. \square

Lemma 2.3. *In $\text{PG}(2, q^2)$, let t be a line tangent to a non-degenerate Hermitian curve $\mathcal{H}(2, q^2)$ at T and let u, u_1, u_2 be three distinct points of t such that the Baer subline of t containing them does not pass through T . For a point $u' \in \text{PG}(2, q^2) \setminus \mathcal{H}(2, q^2)$ such that $\langle u, u' \rangle$ is secant to $\mathcal{H}(2, q^2)$ and $\langle u', u_1 \rangle, \langle u', u_2 \rangle$ are tangents, there are exactly two points R_i , $i = 1, 2$, of $\text{PG}(2, q^2) \setminus (\mathcal{H}(2, q^2) \cup t)$ such that the lines $\langle R_i, u \rangle, \langle R_i, u_1 \rangle, \langle R_i, u_2 \rangle, \langle R_i, u' \rangle$ are tangent to $\mathcal{H}(2, q^2)$. Moreover $\langle R_i, u' \rangle \cap t = u_i$.*

Proof. We may assume without loss of generality that $\mathcal{H}(2, q^2) : X_1^q X_2 + X_1 X_2^q + X_3^{q+1} = 0$, $T = (1, 0, 0)$, $u = (0, 0, 1)$, $u_i = (x_i, 0, 1)$, with $x_1^q x_2 - x_1 x_2^q \neq 0$, otherwise T would belong to the Baer subline of t containing u, u_1, u_2 . Let $R = (c, 1, d)$. By [10, Lemma 2.3], the lines $\langle R, u \rangle, \langle R, u_i \rangle$, $i = 1, 2$, are tangents if and only if $c + c^q = 0$ and $d = \frac{x_1^{q+1} x_2^q - x_1^q x_2^{q+1}}{x_1^q x_2 - x_1 x_2^q}$. Let $u' = (a, 1, b)$, where $a + a^q \neq 0$, since $\langle u, u' \rangle$ is secant, $a + a^q + b^{q+1} \neq 0$, since $u' \notin \mathcal{H}(2, q^2)$ and $b - d = \frac{(a+a^q)(x_1^q - x_2^q)}{x_1^q x_2 - x_1 x_2^q}$, since $|\langle u', u_i \rangle \cap \mathcal{H}(2, q^2)| = 1$, $i = 1, 2$. Thus the line $\langle R, u' \rangle$ is tangent if and only if c satisfies $X^2 + X(a^q - a + b^q d - b d^q) + a d(b-d)^q + a^q d^q (b-d) - a^{q+1} = 0$. The two solutions of the previous quadratic equation are $\frac{a x_1^q (x_2 - x_1) + a^q x_1 (x_2^q - x_1^q)}{x_1^q x_2 - x_1 x_2^q}$ and $\frac{a x_2^q (x_2 - x_1) + a^q x_2 (x_2^q - x_1^q)}{x_1^q x_2 - x_1 x_2^q}$. Moreover it can be easily checked that $\langle u', R \rangle \cap t \in \{u_1, u_2\}$. \square

Lemma 2.4. *Let ℓ be a line of $\text{PG}(2, q^2)$ and let P_1, P_2, P_3 be three non-collinear points of $\text{PG}(2, q^2) \setminus \ell$. Then there are $q^2 + 2q$ Hermitian pencils of lines meeting ℓ in one point and containing P_1, P_2, P_3 .*

Proof. Let \mathcal{C} be a Hermitian pencil of lines consisting of $q + 1$ lines through a point of ℓ and containing the points P_1, P_2, P_3 . If the lines $P_1 P_2, P_1 P_3, P_2 P_3$ are not lines of \mathcal{C} , then there are $q^2 - 2$ of such curves. Note that by projecting the Baer subline of $P_2 P_3$ containing $\ell \cap (P_2 P_3)$, P_2, P_3 from P_1 onto ℓ and by projecting the Baer subline of $P_1 P_3$ containing $\ell \cap (P_1 P_3)$, P_1, P_3 from P_2 onto ℓ and by projecting the Baer subline of $P_1 P_2$ containing $\ell \cap (P_1 P_2)$, P_1, P_2 from P_3 onto ℓ , we obtain the same Baer subline s of ℓ . If a Hermitian pencil of lines contains ℓ and the points P_1, P_2, P_3 , but does not contain the lines $P_1 P_2, P_1 P_3, P_2 P_3$, then its vertex is a point of $s \setminus \{\ell \cap (P_1 P_2), \ell \cap (P_1 P_3), \ell \cap (P_2 P_3)\}$. Hence, among the $q^2 - 2$ Hermitian pencils of lines described above there are exactly $q - 2$ that contain ℓ . There are $q + 1$ Hermitian pencils of lines consisting of $q + 1$ lines through the point of $\ell \cap (P_1 P_2)$, containing the points P_1, P_2, P_3 and the line $P_1 P_2$. Exactly one of these contains ℓ . Similarly for $P_1 P_3$ and $P_2 P_3$. Therefore there are $q^2 - 2 - (q - 2) + 3(q + 1) - 3 = q^2 + 2q$ Hermitian pencils of lines meeting ℓ in one point and containing P_1, P_2, P_3 . \square

Lemma 2.5. *In $\text{PG}(2, q^2)$, let $\mathcal{H}(2, q^2)$ be a non-degenerate Hermitian curve and let P_1, P_2, P_3 be three non-collinear points of $\text{PG}(2, q^2) \setminus \mathcal{H}(2, q^2)$ such that the line $P_i P_j$ is tangent to $\mathcal{H}(2, q^2)$, $1 \leq i < j \leq 3$. Then there are exactly $3q$ Hermitian pencils of lines meeting $\mathcal{H}(2, q^2)$ in $q + 1$ points and containing P_1, P_2, P_3 .*

Proof. Let \perp be the unitary polarity of $\text{PG}(2, q^2)$ defining $\mathcal{H}(2, q^2)$. Let \mathcal{C} be a Hermitian pencil of lines such that $|\mathcal{C} \cap \mathcal{H}(2, q^2)| = q + 1$ and $P_i \in \mathcal{C}$, $i = 1, 2, 3$. Let V be the vertex of \mathcal{C} . Then necessarily $V^\perp = \langle \mathcal{C} \cap \mathcal{H}(2, q^2) \rangle$ and each of the $q + 1$ lines of \mathcal{C} is tangent to $\mathcal{H}(2, q^2)$. If V is one among P_1, P_2, P_3 , then \mathcal{C} is uniquely determined. If $V \notin \{P_1, P_2, P_3\}$, then V must lie on one of the lines $P_1 P_2, P_1 P_3, P_2 P_3$, otherwise $\mathcal{H}(2, q^2)$ would contain the dual of an O'Nan configuration, a contradiction. On the other hand the point V can be chosen in $q - 1$ ways on each of the three lines $P_1 P_2, P_1 P_3, P_2 P_3$ and for every choice of V the curve \mathcal{C} is uniquely determined. \square

Lemma 2.6. *In $\text{PG}(2, q^2)$, let \mathcal{H} and \mathcal{H}' be two non-degenerate Hermitian curves with associated polarity \perp and \perp' , respectively. Let $|\mathcal{H} \cap \mathcal{H}'| \in \{1, q+1\}$ and let ℓ be the (unique) line of $\text{PG}(2, q^2)$ such that $\ell \cap \mathcal{H} = \ell \cap \mathcal{H}' = \mathcal{H} \cap \mathcal{H}'$. Thus $\ell^\perp = \ell^{\perp'}$ and if $P \in \mathcal{H} \setminus \mathcal{H}'$, then $P^{\perp'} \cap P^\perp \in \ell$.*

Proof. We may assume without loss of generality that \mathcal{H} is given by $X_1^q X_2 + X_1 X_2^q + X_3^{q+1} = 0$ and that \mathcal{H}' is given by $X_1^q X_2 + X_1 X_2^q + \lambda X_3^{q+1} = 0$, $\lambda \in \text{GF}(q) \setminus \{0, 1\}$, if $|\mathcal{H} \cap \mathcal{H}'| = q+1$, or that \mathcal{H}' is given by $\lambda X_1^{q+1} + X_1^q X_2 + X_1 X_2^q + X_3^{q+1} = 0$, $\lambda \in \text{GF}(q) \setminus \{0\}$, if $|\mathcal{H} \cap \mathcal{H}'| = 1$. For more details on the intersection of two non-degenerate Hermitian curves see [1, 7, 12]. In the former case ℓ is the line $X_3 = 0$ and $\ell^\perp = \ell^{\perp'} = (0, 0, 1)$. Let $P = (a, b, 1) \in \mathcal{H} \setminus \mathcal{H}'$, then $P^\perp \cap P^{\perp'} = (a^q, -b^q, 0) \in \ell$. In the latter case ℓ is the line $X_1 = 0$ and $\ell^\perp = \ell^{\perp'} = (0, 1, 0)$. Let $P = (1, a, b) \in \mathcal{H} \setminus \mathcal{H}'$, then $P^\perp \cap P^{\perp'} = (0, -b^q, 1) \in \ell$. \square

Proposition 2.7. *In $\text{PG}(2, q^2)$, let $\mathcal{H}(2, q^2)$ be a non-degenerate Hermitian curve and let P_1, P_2, P_3 be three non-collinear points of $\text{PG}(2, q^2) \setminus \mathcal{H}(2, q^2)$ such that the line $P_i P_j$ is tangent to $\mathcal{H}(2, q^2)$, $1 \leq i < j \leq 3$. Then there are exactly $q^2 - q + 1$ non-degenerate Hermitian curves meeting $\mathcal{H}(2, q^2)$ in one or $q+1$ points and containing P_1, P_2, P_3 .*

Proof. Assume that $P_1 = (1, 0, 0)$, $P_2 = (0, 1, 0)$, $P_3 = (0, 0, 1)$. If \mathcal{H} is a non-degenerate Hermitian curve containing P_1, P_2, P_3 , then \mathcal{H} is given by

$$a_{12} X_1 X_2^q + a_{12}^q X_1^q X_2 + a_{13} X_1 X_3^q + a_{13}^q X_1^q X_3 + a_{23} X_2 X_3^q + a_{23}^q X_2^q X_3 = 0,$$

for some $a_{12}, a_{13}, a_{23} \in \text{GF}(q^2)$, with

$$a_{12} a_{13}^q a_{23} + a_{12}^q a_{13} a_{23}^q \neq 0. \quad (2.1)$$

On the other hand, if $\mathcal{H}(2, q^2)$ is a non-degenerate Hermitian curve such that $P_1, P_2, P_3 \in \text{PG}(2, q^2) \setminus \mathcal{H}(2, q^2)$ and the line $P_i P_j$ is tangent to $\mathcal{H}(2, q^2)$, $1 \leq i < j \leq 3$, then $\mathcal{H}(2, q^2)$ is given by

$$y^{q+1} X_1^{q+1} + z^{q+1} X_2^{q+1} + X_3^{q+1} - xz^{q+1} X_1 X_2^q - x^q z^{q+1} X_1^q X_2 - yX_1 X_3^q - y^q X_1^q X_3 - zX_2 X_3^q - z^q X_2^q X_3 = 0,$$

for some $x, y, z \in \text{GF}(q^2)$ such that $y^{q+1} = x^{q+1} z^{q+1}$ and $y \neq -xz$. By using the projectivity $X'_1 = yX_1, X'_2 = zX_2, X'_3 = X_3$, we may assume that $\mathcal{H}(2, q^2)$ is given by

$$X_1^{q+1} + X_2^{q+1} + X_3^{q+1} - \alpha X_1 X_2^q - \alpha^q X_1^q X_2 - X_1 X_3^q - X_1^q X_3 - X_2 X_3^q - X_2^q X_3 = 0,$$

for some $\alpha \in \text{GF}(q^2)$ such that $\alpha^{q+1} = 1$ and $\alpha \neq -1$. Then the lines $P_1 P_2$, $P_1 P_3$ and $P_2 P_3$ are tangent to $\mathcal{H}(2, q^2)$ at the points $L_3 = (1, \alpha, 0)$, $L_2 = (1, 0, 1)$ and $L_1 = (0, 1, 1)$, respectively. Let \perp and \perp' be the polarities defining $\mathcal{H}(2, q^2)$ and \mathcal{H} , respectively. If $|\mathcal{H}(2, q^2) \cap \mathcal{H}| \in \{1, q+1\}$, according to Lemma 2.6, we have that $L_i^\perp \cap L_i^{\perp'} \in \ell$, $i = 1, 2, 3$, where ℓ is the unique line of $\text{PG}(2, q^2)$ such that $\ell \cap \mathcal{H}(2, q^2) = \ell \cap \mathcal{H} = \mathcal{H}(2, q^2) \cap \mathcal{H}$. Denote by Z_i the point $L_i^\perp \cap L_i^{\perp'}$, $i = 1, 2, 3$ and let L be the point ℓ^\perp . Then

$$Z_1 = (0, -a_{23}^q, a_{23}), Z_2 = (-a_{13}^q, 0, a_{13}), Z_3 = (-a_{12}^q, \alpha^q a_{12}, 0)$$

are on a line if and only if

$$a_{12}^q a_{13} a_{23}^q - \alpha^q a_{12} a_{13}^q a_{23} = 0. \quad (2.2)$$

Note that $L = Z_1^\perp \cap Z_2^\perp = (a_{13}(a_{23} + a_{23}^q), \alpha a_{23}(a_{13} + a_{13}^q), \alpha a_{13}^q a_{23} + a_{13} a_{23}^q)$ and since $\ell^\perp = \ell^{\perp'}$, we have that $Z_1, Z_2 \in L^{\perp'}$. Some calculations show that $Z_1, Z_2 \in L^{\perp'}$ if and only if

$$(a_{23} + a_{23}^q)(a_{23} a_{13}^q - a_{23}^q a_{12}^q) + a_{23}^{q+1}(\alpha^q a_{23}^q - a_{23}) = 0, \quad (2.3)$$

$$(a_{13} + a_{13}^q)(a_{23}^q a_{13} - a_{13}^q a_{12}) + a_{13}^{q+1}(\alpha a_{13}^q - a_{13}) = 0. \quad (2.4)$$

Since $\alpha \neq -1$, it follows from (2.4) that

$$a_{13} + a_{13}^q \neq 0. \quad (2.5)$$

Hence from (2.4), we have that

$$a_{12} = \frac{a_{13} a_{23}^q (a_{13} + a_{13}^q) + a_{13}^{q+1} (\alpha a_{13}^q - a_{13})}{a_{13}^q (a_{13} + a_{13}^q)}. \quad (2.6)$$

By substituting (2.6) in (2.2), we get that one of the following has to be satisfied:

$$a_{13}^q - \alpha^q a_{13} = 0, \quad (2.7)$$

$$a_{23}^{q+1}(a_{13} + a_{13}^q) - a_{13}^{q+1}(a_{23} + a_{23}^q) = 0. \quad (2.8)$$

Assume that (2.8) is satisfied. Since $a_{23} + a_{23}^q \neq 0$, then

$$a_{13} = \frac{a_{23}^{q+1}}{a_{23} + a_{23}^q} (\xi + 1), \quad (2.9)$$

where $\xi \in \text{GF}(q^2)$, $\xi^{q+1} = 1$, with $\xi \neq -1$, otherwise $a_{13} = 0$, contradicting (2.1). By taking into account (2.9), equation (2.6) becomes

$$a_{12} = \frac{a_{23}^q (\xi a_{23}^q + \alpha a_{23})}{a_{23} + a_{23}^q}. \quad (2.10)$$

Moreover, if (2.9) and (2.10) hold true, then (2.3) is satisfied, whereas some calculations show that condition (2.1) is satisfied whenever $\xi a_{23}^q + \alpha a_{23} \neq 0$, i.e,

$$a_{23}^{q-1} \neq -\xi^q \alpha. \quad (2.11)$$

The equation $X^{q-1} = -\xi^q \alpha$ has $q-1$ solutions in $\text{GF}(q^2)$ since $(-\xi^q \alpha)^{q+1} = 1$.

Assume that (2.7) holds true. Thus it turns out that necessarily $a_{12} = \alpha a_{23}^q$ and $a_{13} = a_{23}^{q+1}(\alpha + 1)/(a_{23} + a_{23}^q)$ and hence we retrieve a particular solution of the previous case (the one with $\xi = \alpha$).

Observe that for a fixed $\xi \in \text{GF}(q^2)$, with $\xi^{q+1} = 1$, $\xi \neq -1$, we have that

$$|\{a_{23} \in \text{GF}(q^2) \mid a_{23} + a_{23}^q \neq 0, a_{23}^{q-1} \neq -\xi^q \alpha\}| = \begin{cases} q(q-1) & \text{if } \xi = \alpha, \\ (q-1)^2 & \text{if } \xi \neq \alpha. \end{cases}$$

Furthermore for a fixed $a_{23} \in \text{GF}(q^2)$ satisfying (2.5) and (2.11), then a_{12} and a_{13} are uniquely determined by (2.10) and (2.9), respectively, and, if $\rho \in \text{GF}(q) \setminus \{0\}$, the values $a_{23}, \rho a_{23}$ give rise to the same Hermitian curve. \square

Lemma 2.8. *In $\text{PG}(3, q^2)$, let $\mathcal{H}(3, q^2)$ be a non-degenerate Hermitian surface with associated polarity \perp and let $\gamma = P^\perp$ be a plane secant to $\mathcal{H}(3, q^2)$. If P' is a point of $\text{PG}(3, q^2) \setminus (\mathcal{H}(3, q^2) \cup \gamma \cup \{P\})$, then the points of γ lying on a line passing through P' and tangent to $\mathcal{H}(3, q^2)$ form a non-degenerate Hermitian curve \mathcal{H}' of γ such that $\gamma \cap \mathcal{H}' \cap \mathcal{H}(3, q^2) \subset \langle PP' \rangle^\perp$. Viceversa, if \mathcal{H}' is a non-degenerate Hermitian curve of γ meeting $\gamma \cap \mathcal{H}(3, q^2)$ in one or $q+1$ points and ℓ is the (unique) line of γ such that $\ell \cap \mathcal{H}' = \ell \cap \mathcal{H}(3, q^2) = \gamma \cap \mathcal{H}' \cap \mathcal{H}(3, q^2)$, then there are exactly $q+1$ points P' of $\ell^\perp \setminus (\mathcal{H}(3, q^2) \cup \gamma \cup \{P\})$ such that the lines joining P' with a point of \mathcal{H}' are tangent to $\mathcal{H}(3, q^2)$.*

Proof. Let $\mathcal{H}(3, q^2)$ be the Hermitian surface of $\text{PG}(3, q^2)$ given by $X_1^{q+1} + X_2^{q+1} + X_3^{q+1} + X_4^{q+1} = 0$ and let γ be the plane $X_4 = 0$. Let P' be a point of $\text{PG}(3, q^2) \setminus (\mathcal{H}(3, q^2) \cup \gamma \cup \{P\})$. If PP' is tangent to $\mathcal{H}(3, q^2)$, then we may assume that $PP' \cap \gamma = (1, \xi, 0, 0)$, for some $\xi \in \text{GF}(q^2)$ such that $\xi^{q+1} = -1$. Then $P' = (1, \xi, 0, \lambda)$ where $\lambda \in \text{GF}(q^2) \setminus \{0\}$. Similarly if PP' is secant to $\mathcal{H}(3, q^2)$, then we may assume that $PP' \cap \gamma = (1, 0, 0, 0)$. In this case $P' = (1, 0, 0, \lambda)$ where $\lambda \in \text{GF}(q^2) \setminus \{0\}$ and $\lambda^{q+1} \neq -1$. From [10, Lemma 2.3], the line joining P' and $Q = (x_1, x_2, x_3, 0) \in \gamma$ is tangent to $\mathcal{H}(3, q^2)$ if and only if $Q \in \mathcal{H}'$, where \mathcal{H}' is the non-degenerate Hermitian curve of γ given by

$$(\lambda^{q+1} - 1)x_1^{q+1} + (\lambda^{q+1} + 1)x_2^{q+1} + \lambda^{q+1}x_3^{q+1} - \xi^q x_1^q x_2 - \xi x_1 x_2^q = 0$$

if $|PP' \cap \mathcal{H}(3, q^2)| = 1$ or by

$$\lambda^{q+1}x_1^{q+1} + (\lambda^{q+1} + 1)(x_2^{q+1} + x_3^{q+1}) = 0$$

if $|PP' \cap \mathcal{H}(3, q^2)| = q+1$. Note that $\gamma \cap \mathcal{H}' \cap \mathcal{H}(3, q^2) \subset \langle PP' \rangle^\perp$ and by replacing λ with λ' , where $\lambda^{q+1} = \lambda'^{q+1}$ we retrieve the same Hermitian curve \mathcal{H}' . \square

3 Graphs cospectral with $\text{NU}(3, q^2)$

Let Π be a finite projective plane of order q^2 and let \mathcal{U} be a unital of Π . A line of Π is said to be *tangent* or *secant* to \mathcal{U} according as it meets \mathcal{U} in 1 or $q+1$ points, respectively. There are $q^3 + 1$ tangent lines and $q^4 - q^3 + q^2$ secant lines. In particular, through each point of \mathcal{U} there pass q^2 secant lines and one tangent line, while through each point of $\Pi \setminus \mathcal{U}$ there pass $q+1$ tangent lines and $q^2 - q$ secant lines. Starting from \mathcal{U} , it is possible to define a unital \mathcal{U}^* of the dual plane Π^* of Π . This can be done by taking the tangent lines to \mathcal{U} as the points of \mathcal{U}^* and the points of $\Pi \setminus \mathcal{U}$ as the secant lines to \mathcal{U}^* , where incidence is given by reverse containment. Thus \mathcal{U}^* is a unital of Π^* , called the *dual unital* to \mathcal{U} .

Let $\Gamma_{\mathcal{U}}$ be the graph whose vertices are the points of $\Pi \setminus \mathcal{U}$ and two vertices P_1, P_2 are adjacent if the line joining P_1 and P_2 is tangent to \mathcal{U} .

Proposition 3.1. *The graph $\Gamma_{\mathcal{U}}$ is strongly regular with parameters $(q^2(q^2 - q + 1), (q+1)(q^2 - 1), 2(q^2 - 1), (q+1)^2)$.*

Proof. Since every point of $\Pi \setminus \mathcal{U}$ lies on $q+1$ lines that are tangent to \mathcal{U} , the graph $\Gamma_{\mathcal{U}}$ is $(q+1)(q^2 - 1)$ -regular. Consider two distinct vertices u_1, u_2 of $\Gamma_{\mathcal{U}}$ corresponding to the points

P_1, P_2 of $\Pi \setminus \mathcal{U}$. If u_1, u_2 are adjacent, then the line P_1P_2 is tangent to \mathcal{U} and u_1, u_2 have $2(q^2 - 1)$ common neighbours. If u_1, u_2 are not adjacent, then the line P_1P_2 is secant to \mathcal{U} and u_1, u_2 have $(q + 1)^2$ common neighbours. \square

A *clique* of a graph is a subset of pairwise adjacent vertices. A clique is said to be *maximal* if it is maximal with respect to set theoretical inclusion.

Proposition 3.2. *The graph $\Gamma_{\mathcal{U}}$ contains a class of $q^3 + 1$ maximal cliques of size q^2 and a class of $q^2(q^3 + 1)(q^2 - q + 1)$ maximal cliques of size $q + 2$.*

Proof. In the first case, a clique corresponds to the points on a tangent line with the tangent point excluded. In the second case, it corresponds to $q + 1$ points on a tangent line ℓ , and the other point off ℓ . In both cases these cliques are maximals. \square

Let $\Pi = \text{PG}(2, q^2)$ and let \mathcal{U} be the classical unital $\mathcal{H}(2, q^2)$. In this case $\Gamma_{\mathcal{U}}$ is denoted by $\text{NU}(3, q^2)$. Since a dual O’Nan configuration cannot be embedded in $\mathcal{H}(2, q^2)$, we have the following (see also [4, Corollary 3]).

Corollary 3.3. *The maximal cliques of the graph $\text{NU}(3, q^2)$ are exactly those described in Proposition 3.2.*

In what follows we recall few known properties of some classes unitals of $\text{PG}(2, q^2)$. Besides the classical one, there are other known unitals in $\text{PG}(2, q^2)$ and all the known unitals of $\text{PG}(2, q^2)$ arise from a construction due to Buekenhout, see [1, 2, 14]. A *Buekenhout unital* of $\text{PG}(2, q^2)$ is either an orthogonal Buekenhout–Metz unital or a Buekenhout–Tits unital. An *orthogonal Buekenhout–Metz unital* of $\text{PG}(2, q^2)$, $q > 2$, is projectively equivalent to one of the following form:

$$\mathcal{U}_{\alpha, \beta} = \{(x, \alpha x^2 + \beta x^{q+1} + z, 1) \mid z \in \text{GF}(q), x \in \text{GF}(q^2)\} \cup \{(0, 1, 0)\},$$

where α, β are elements in $\text{GF}(q^2)$ such that $(\beta - \beta^q)^2 + 4\alpha^{q+1}$ is a non-square in $\text{GF}(q)$ if q is odd, or $\beta \notin \text{GF}(q)$ and $\alpha^{q+1}/(\beta + \beta^q)^2$ has absolute trace 0 if q is even and $q > 2$. The unital $\mathcal{U}_{\alpha, \beta}$ is classical if and only if $\alpha = 0$. If q is odd, $\beta \in \text{GF}(q)$ and $\alpha \neq 0$, then $\mathcal{U}_{\alpha, \beta}$ can be obtained glueing together q conics of $\text{PG}(2, q^2)$ having in common the point $(0, 1, 0)$. Let $m > 1$ be an odd integer and let $q = 2^m$. A *Buekenhout–Tits unital* of $\text{PG}(2, q^2)$ is projectively equivalent to

$$\mathcal{U}_T = \{(x_0 + x_1\beta, (x_0^{\delta+2} + x_0x_1 + x_1^\delta)\beta + z, 1) \mid x_0, x_1, z \in \text{GF}(q)\} \cup \{(0, 1, 0)\},$$

where β is an element of $\text{GF}(q^2) \setminus \text{GF}(q)$ and $\delta = 2^{\frac{m+1}{2}}$. See [6] for more details.

As the Desarguesian plane is self-dual, that is, it is isomorphic to its dual plane, the dual of a unital in $\text{PG}(2, q^2)$ is a unital in $\text{PG}(2, q^2)$. The non-classical orthogonal Buekenhout–Metz unital $\mathcal{U}_{\alpha, \beta}$ as well as the Buekenhout–Tits unital \mathcal{U}_T of $\text{PG}(2, q^2)$ are self-dual [1, Theorem 4.17, Theorem 4.28, Corollary 4.35]. Moreover the O’Nan configuration can be embedded in each non-classical Buekenhout unital of $\text{PG}(2, q^2)$ [6]. Therefore it follows that there exists a dual O’Nan configuration embedded in both unitals $\mathcal{U}_{\alpha, \beta}$ and \mathcal{U}_T^\dagger .

[†]We thank A. Aguglia for pointing this out.

Proposition 3.4. *There exists a dual O’Nan configuration embedded in a non-classical Buekenhout unital of $\text{PG}(2, q^2)$.*

For the convenience of the reader we illustrate the previous statement with an example. Let q be odd and let us fix α, β in $\text{GF}(q^2)$, $\alpha \neq 0$, such that $(\beta - \beta^q)^2 + 4\alpha^{q+1}$ is a non-square in $\text{GF}(q)$. Set $\mathcal{U} = \{(-2\alpha x + (\beta^q - \beta)x^q, 1, \alpha x^2 - \beta^q x^{q+1} - z) \mid x \in \text{GF}(q^2), z \in \text{GF}(q)\} \cup \{(0, 0, 1)\}$. From the proof of [1, Theorem 4.17] \mathcal{U} is projectively equivalent to $\mathcal{U}_{\alpha, \beta}$. Consider the six lines given by: $[0, 0, 1]$, $[0, -2x_{\lambda_1}x_{\lambda_2}, x_{\lambda_1} + x_{\lambda_2}]$, $[x_{\lambda_1}, -x_{\lambda_1}, 1]$, $[x_{\lambda_2}, -x_{\lambda_2}, 1]$, $[-x_{\lambda_1}, -x_{\lambda_1}, 1]$, $[-x_{\lambda_2}, -x_{\lambda_2}, 1]$, where $x_{\lambda_i} = -\frac{\alpha^q + \lambda_i - \beta}{\alpha^{q+1} - (\lambda_i - \beta)^{q+1}}$, $i = 1, 2$, and $\lambda_1, \lambda_2 \in \text{GF}(q)$ are such that $(\lambda_1 + \alpha^q - \beta)^{q+1}(\alpha^{q+1} - \beta^{q+1} - \lambda_2^2) + (\lambda_2 + \alpha^q - \beta)^{q+1}(\alpha^{q+1} - \beta^{q+1} - \lambda_1^2) = 0$. By [1, Theorem 4.17] and [6, Corollary 3.6] these lines are tangent to \mathcal{U} and there are four non-collinear points of $\text{PG}(2, q^2) \setminus \mathcal{U}$ such that the lines joining two of these four points are exactly the six lines given above.

An easy consequence of the previous proposition is the following result.

Corollary 3.5. *If \mathcal{U} is a non-classical Buekenhout unital of $\text{PG}(2, q^2)$, then the graph $\Gamma_{\mathcal{U}}$ has at least a further class of cliques than those described in Proposition 3.2. A clique in this class contains four points no three on a line.*

Comparing Corollary 3.3 with Corollary 3.5, we see that $\Gamma_{\mathcal{U}}$ has more maximal cliques than $\Gamma_{\mathcal{U}}$ and we obtain the desired result for $q > 2$.

Theorem 3.6. *Let $q > 2$. If \mathcal{U} is a non-classical Buekenhout unital of $\text{PG}(2, q^2)$, then the graph $\Gamma_{\mathcal{U}}$ is not isomorphic to $\text{NU}(3, q^2)$.*

4 Graphs cospectral with $\text{NU}(n + 1, q^2)$, $n \geq 4$

In this section we introduce two classes of graphs cospectral with $\text{NU}(n + 1, q^2)$, $n \geq 4$. These graphs are obtained by modifying few adjacencies in $\text{NU}(n + 1, q^2)$ following an idea proposed by Wang, Qiu, and Hu. For a graph \mathcal{G} and for a vertex u of \mathcal{G} , let $N_{\mathcal{G}}(u)$ be the set of neighbours of u in \mathcal{G} . Let us recall the Wang–Qiu–Hu switching in a simplified version as stated in [13].

Theorem 4.1 (WQH switching). *Let \mathcal{G} be a graph whose vertex set is partitioned as $\ell_1 \cup \ell_2 \cup \mathcal{D}$. Assume that the induced subgraphs on ℓ_1, ℓ_2 , and $\ell_1 \cup \ell_2$ are regular, and that the induced subgraphs on ℓ_1 and ℓ_2 have the same size and degree. Suppose that each $x \in \mathcal{D}$ either has the same number of neighbours in ℓ_1 and ℓ_2 , or $N_{\mathcal{G}}(x) \cap (\ell_1 \cup \ell_2) \in \{\ell_1, \ell_2\}$. Construct a new graph \mathcal{G}' by switching adjacency and non-adjacency between $x \in \mathcal{D}$ and $\ell_1 \cup \ell_2$ when $N_{\mathcal{G}}(x) \cap (\ell_1 \cup \ell_2) \in \{\ell_1, \ell_2\}$. Then \mathcal{G} and \mathcal{G}' are cospectral.*

4.1 The graphs \mathcal{G}'_n and \mathcal{G}''_n

Let $\mathcal{H}(n, q^2)$ be a non-degenerate Hermitian variety of $\text{PG}(n, q^2)$, $n \geq 4$, with associated polarity \perp . Let us denote by \mathcal{G}_n the graph $\text{NU}(n + 1, q^2)$. Recall that the vertices of \mathcal{G}_n are the points of

$\text{PG}(n, q^2) \setminus \mathcal{H}(n, q^2)$ and that two vertices are adjacent if their span is a line tangent to $\mathcal{H}(n, q^2)$. The graph \mathcal{G}_n is strongly regular with parameters

$$v = \frac{q^n(q^{n+1} - \epsilon)}{q + 1}, \quad k = (q^n + \epsilon)(q^{n-1} - \epsilon),$$

$$\lambda = q^{2n-3}(q + 1) - \epsilon q^{n-1}(q - 1) - 2, \quad \mu = q^{n-2}(q + 1)(q^{n-1} - \epsilon),$$

where $\epsilon = (-1)^{n+1}$. Fix a point $P \in \mathcal{H}(n, q^2)$ and let $\bar{\ell}_1, \bar{\ell}_2$ be two lines of $\text{PG}(n, q^2)$ such that $\bar{\ell}_1 \cap \mathcal{H}(n, q^2) = \bar{\ell}_2 \cap \mathcal{H}(n, q^2) = P$. Let π be the plane spanned by $\bar{\ell}_1$ and $\bar{\ell}_2$ and let $\ell_i = \bar{\ell}_i \setminus \{P\}$, $i = 1, 2$. Thus ℓ_i consists of q^2 vertices of \mathcal{G}_n . There are two possibilities: either $\pi \cap \mathcal{H}(n, q^2)$ is a Hermitian pencil of lines or is a line. Let us define the following sets:

$$\begin{aligned} \mathcal{A} &= (\cap_{u \in \ell_1} N_{\mathcal{G}_n}(u)) \cap (\cap_{u \in \ell_2} N_{\mathcal{G}_n}(u)), \\ \mathcal{A}_1 &= (\cap_{u \in \ell_1} N_{\mathcal{G}_n}(u)) \setminus (\mathcal{A} \cup \ell_2), \\ \mathcal{A}_2 &= (\cap_{u \in \ell_2} N_{\mathcal{G}_n}(u)) \setminus (\mathcal{A} \cup \ell_1). \end{aligned}$$

Lemma 4.2. *Let $n = 4$.*

- *If $\pi \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines, then the set \mathcal{A} or \mathcal{A}_i , $i = 1, 2$, consists of the points lying on $(q + 1)^2$ or $(q + 1)(q^2 - q - 2)$ lines tangent to $\mathcal{H}(4, q^2)$ at P , minus P itself. In particular $|\mathcal{A}| = q^2(q + 1)^2$ and $|\mathcal{A}_1| = |\mathcal{A}_2| = q^2(q + 1)(q^2 - q - 2)$.*
- *If $\pi \cap \mathcal{H}(4, q^2)$ is a line, then the set \mathcal{A} or \mathcal{A}_i , $i = 1, 2$, consists of the points lying on $2(q^2 - 1)$ or $q(q^2 - q - 1)$ lines tangent to $\mathcal{H}(4, q^2)$ at P , minus P itself. In particular $|\mathcal{A}| = 2q^2(q^2 - 1)$ and $|\mathcal{A}_1| = |\mathcal{A}_2| = q^3(q^2 - q - 1)$.*

Proof. Assume that $\pi \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines. Let u' be a vertex of \mathcal{G}_4 such that it is adjacent to any vertex of ℓ_1 . Then $u' \notin \ell$, otherwise $u' \in N_{\mathcal{G}_4}(u')$, a contradiction. If $u \in \ell_1$, the line determined by u and u' contains a unique point of $\mathcal{H}(4, q^2)$ and hence the plane σ spanned by $\bar{\ell}_1$ and u' has exactly $q^2 + 1$ points of $\mathcal{H}(4, q^2)$. Therefore $\sigma \cap \mathcal{H}(4, q^2)$ is a line and σ contains $q^4 - q^2$ vertices of \mathcal{G}_4 adjacent to every vertex of ℓ_1 . Note that there are $q + 1$ planes through ℓ_1 meeting $\mathcal{H}(4, q^2)$ in a line. Hence $|\mathcal{A}_1 \cup \mathcal{A}| = (q + 1)(q^4 - q^2)$. A similar argument holds for ℓ_2 . Moreover if σ_i is a plane through ℓ_i having $q^2 + 1$ points in common with $\mathcal{H}(4, q^2)$, $i = 1, 2$, then $\sigma_1 \cap \sigma_2$ is a line tangent to $\mathcal{H}(4, q^2)$ at P . Therefore $\mathcal{A} = q^2(q + 1)^2$ and the first part of the statement follows. If $\pi \cap \mathcal{H}(n, q^2)$ is a line, one can argue as before. \square

It can be easily deduced that $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2 \subseteq P^\perp$. If $\pi \cap \mathcal{H}(n, q^2)$ is a Hermitian pencil of lines and $(n, q) = (4, 2)$, then $|\mathcal{A}_1| = |\mathcal{A}_2| = 0$.

Construction 4.3. Define a graph \mathcal{G}'_n or \mathcal{G}''_n , according as $\pi \cap \mathcal{H}(n, q^2)$ is a Hermitian pencil of lines or a line. The vertices of \mathcal{G}'_n and \mathcal{G}''_n are the vertices of \mathcal{G}_n and the edges are as follows:

$$\begin{cases} N_{\mathcal{G}_n}(u) & \text{if } u \notin \mathcal{A}_1 \cup \mathcal{A}_2 \cup \ell_1 \cup \ell_2, \\ (N_{\mathcal{G}_n}(u) \setminus \mathcal{A}_1) \cup \mathcal{A}_2 & \text{if } u \in \ell_1, \\ (N_{\mathcal{G}_n}(u) \setminus \mathcal{A}_2) \cup \mathcal{A}_1 & \text{if } u \in \ell_2, \\ (N_{\mathcal{G}_n}(u) \setminus \ell_1) \cup \ell_2 & \text{if } u \in \mathcal{A}_1, \\ (N_{\mathcal{G}_n}(u) \setminus \ell_2) \cup \ell_1 & \text{if } u \in \mathcal{A}_2. \end{cases}$$

Note that the graphs \mathcal{G}'_n and \mathcal{G}''_n are obtained from \mathcal{G}_n by applying the WQH switching as described in Theorem 4.1. Indeed, the vertex set of \mathcal{G}_n consists of $\ell_1 \cup \ell_2 \cup \mathcal{D}$, where $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2 \subset \mathcal{D}$. Furthermore, if $x \in \mathcal{D} \setminus (\mathcal{A}_1 \cup \mathcal{A}_2)$, then $N_{\mathcal{G}_n}(x) \cap \ell_1 = N_{\mathcal{G}_n}(x) \cap \ell_2 \in \{0, 1, q^2\}$, whereas if $x \in \mathcal{A}_i$, then $N_{\mathcal{G}_n}(x) \cap \ell_i = q^2$ and $N_{\mathcal{G}_n}(x) \cap \ell_j = 0$, where $\{i, j\} = \{1, 2\}$.

Theorem 4.4. *The graphs \mathcal{G}'_n and \mathcal{G}''_n are strongly regular and have the same parameters as \mathcal{G}_n .*

Proof. It is sufficient to show that the hypotheses of Theorem 4.1 are satisfied. Consider the graph \mathcal{G}'_n . The induced subgraph on ℓ_i is the complete graph on q^2 vertices; as there is no edge between a vertex of ℓ_1 and a vertex of ℓ_2 , the induced subgraph on $\ell_1 \cup \ell_2$ is the union of the two complete graphs. Let u be a vertex of \mathcal{G}_n with $u \notin \ell_1 \cup \ell_2$. If u is not in π , let σ_i be the span of u and ℓ_i for $i = 1, 2$. If $\sigma_i \cap \mathcal{H}(n, q^2)$ is a line, then $N_{\mathcal{G}_n}(u) \cap \ell_i = \ell_i$. If $\sigma_i \cap \mathcal{H}(n, q^2)$ is a Hermitian pencil of lines, then all lines of σ_i that are tangent to $\mathcal{H}(n, q^2)$ go through P , so $|N_{\mathcal{G}_n}(u) \cap \ell_i| = 0$. If $\sigma_i \cap \mathcal{H}(n, q^2)$ is a non-degenerate Hermitian curve, then ℓ_i is the unique line of σ_i that is tangent to $\mathcal{H}(n, q^2)$ at P . In this case, as u lies on $q+1$ lines of σ_i that are tangent to $\mathcal{H}(n, q^2)$, we have that $|N_{\mathcal{G}_n}(u) \cap \ell_i| = q+1$. The curve $\sigma_1 \cap \mathcal{H}(n, q^2)$ is non-degenerate if and only if the line $\sigma_1 \cap \sigma_2$ is secant to $\mathcal{H}(n, q^2)$, i.e., if and only if the curve $\sigma_2 \cap \mathcal{H}(n, q^2)$ is non-degenerate. Assume that $u \in \pi$. Thus $|N_{\mathcal{G}_n}(u) \cap \ell_i| = 0$, $i = 1, 2$. Hence, we have shown that we can apply Theorem 4.1 and obtain a strongly regular graph with the same parameters as \mathcal{G}_n . By using the same arguments used for \mathcal{G}' , it is possible to see that the graph \mathcal{G}''_n is strongly regular and has the same parameters as \mathcal{G}_n . \square

4.2 The vertices adjacent to three pairwise adjacent vertices of \mathcal{G}_4

For a point R of $\text{PG}(4, q^2) \setminus \mathcal{H}(4, q^2)$ let \mathcal{T}_R be the set of points lying on a line tangent to $\mathcal{H}(4, q^2)$ at R . Thus $\mathcal{T}_R = N_{\mathcal{G}_4}(R) \cup (R^\perp \cap \mathcal{H}(4, q^2)) \cup \{R\}$. Let P_1, P_2, P_3 be three points of $\text{PG}(4, q^2) \setminus \mathcal{H}(4, q^2)$ such that the line $P_i P_j$ is tangent to $\mathcal{H}(4, q^2)$, $1 \leq i < j \leq 3$, or, in other words, P_1, P_2, P_3 are three pairwise adjacent vertices of \mathcal{G}_4 . Let $\gamma = \langle P_1, P_2, P_3 \rangle$. Note that if γ is a plane, then $\gamma \cap \mathcal{H}(4, q^2)$ is not a degenerate Hermitian curve of rank two. Indeed every tangent line contained in a plane meeting $\mathcal{H}(4, q^2)$ in a degenerate Hermitian curve of rank two pass through the same point. Hence there are three possibilities:

- 1) γ is a line tangent to $\mathcal{H}(4, q^2)$ and hence $\gamma \cap \mathcal{H}(4, q^2)$ is a point;
- 2) γ is a plane and $\gamma \cap \mathcal{H}(4, q^2)$ is a line;
- 3) γ is a plane and $\gamma \cap \mathcal{H}(4, q^2)$ is a non-degenerate Hermitian curve.

Lemma 4.5. *Let P_1, P_2, P_3 be three points of $\text{PG}(4, q^2) \setminus \mathcal{H}(4, q^2)$ such that the line $P_i P_j$ is tangent to $\mathcal{H}(4, q^2)$, $1 \leq i < j \leq 3$, $\gamma = \langle P_1, P_2, P_3 \rangle$ is a plane and $\gamma \cap \mathcal{H}(4, q^2)$ is a line ℓ .*

- *If R is a point of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma)$, then $\mathcal{T}_R \cap \gamma$ is a Hermitian pencil of lines meeting ℓ in one point.*
- *If \mathcal{C} is a Hermitian pencil of lines of γ meeting ℓ in one point, then there are q^3 points $R \in \text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma)$ such that $\mathcal{T}_R \cap \gamma = \mathcal{C}$.*

Proof. Let R be a point of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma)$. The plane $\langle R, \ell \rangle$ meets $\mathcal{H}(4, q^2)$ in a Hermitian pencil of lines and hence the line $t = R^\perp \cap \gamma = \langle R, \gamma^\perp \rangle^\perp = \langle R, \ell \rangle^\perp$ is tangent to $\mathcal{H}(4, q^2)$. By projecting the plane γ from R onto R^\perp , we get a plane γ' , where $\gamma \cap \gamma' = t$. Moreover, since $R^\perp \cap \mathcal{H}(4, q^2)$ is a non-degenerate Hermitian surface and $\gamma' \subset R^\perp$, it follows that $|\gamma' \cap \mathcal{H}(4, q^2)| \in \{q^3 + q^2 + 1, q^3 + 1\}$. Note that if L is a point of ℓ , then the line RL shares with $\mathcal{H}(4, q^2)$ either one or $q + 1$ points according as $L \in R^\perp$ or $L \notin R^\perp$, respectively, and $\ell \cap R^\perp = \ell \cap t$, since $\ell \subset \gamma$. Hence $|\ell \cap t| = 1$. By projecting the line ℓ from R onto R^\perp , we obtain a line ℓ' that is tangent to $\mathcal{H}(4, q^2)$, with $\ell' \subset \gamma'$ and $\ell' \cap \mathcal{H}(4, q^2) = t \cap \mathcal{H}(4, q^2)$. In particular in γ' , there are two distinct tangent lines, namely t and ℓ' , such that $t \cap \ell' \in \ell$ and hence $\gamma' \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines. Therefore, we deduce that there are $q^3 + q^2$ points of $\gamma \setminus \mathcal{H}(4, q^2)$ belonging to $N_{\mathcal{G}_4}(R)$. This shows the first part of the claim.

In the plane γ , there are $q^2(q-1)(q^2+1)$ Hermitian pencils of lines meeting ℓ in one point, and these are permuted in a single orbit by the group $\text{PGU}(5, q^2)_\gamma$, the stabilizer of γ in $\text{PGU}(5, q^2)$. On the other hand, $\text{PGU}(5, q^2)_\gamma$ permutes in a unique orbit the $q^5(q-1)(q^2+1)$ points of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma)$. Consider the incidence structure whose point set \mathcal{P} is formed by the points of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma)$ and whose block set \mathcal{L} consists of the $q^2(q-1)(q^2+1)$ degenerate Hermitian curves described above, where a point $R \in \mathcal{P}$ is incident with a block $\mathcal{C} \in \mathcal{L}$ if $\mathcal{T}_R \cap \gamma = \mathcal{C}$. Since both, \mathcal{P} and \mathcal{L} , are orbits of the same group, the considered incidence structure is a tactical configuration and hence we have that a block of \mathcal{L} is incident with q^3 points of \mathcal{P} , as required. \square

Lemma 4.6. *Let P_1, P_2, P_3 be three points of $\text{PG}(4, q^2) \setminus \mathcal{H}(4, q^2)$ such that the line $P_i P_j$ is tangent to $\mathcal{H}(4, q^2)$, $1 \leq i < j \leq 3$, $\gamma = \langle P_1, P_2, P_3 \rangle$ is a plane and $\gamma \cap \mathcal{H}(4, q^2)$ is a non-degenerate Hermitian curve.*

- If R is a point of $\gamma^\perp \setminus \mathcal{H}(4, q^2)$, then $\mathcal{T}_R \cap \gamma \subset \mathcal{H}(4, q^2)$.
- If R is a point of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma^\perp)$ and $\langle R, \gamma \rangle \cap \gamma^\perp \in \mathcal{H}(4, q^2)$, then $\mathcal{T}_R \cap \gamma$ is a Hermitian pencil of lines, say \mathcal{C} , meeting $\mathcal{H}(4, q^2)$ in $q + 1$ points. In this case there are $(q + 1)(q^2 - 1)$ points $R \in \text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma^\perp)$ such that $\mathcal{T}_R \cap \gamma = \mathcal{C}$.
- If R is a point of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma^\perp)$ and $\langle R, \gamma \rangle \cap \gamma^\perp \notin \mathcal{H}(4, q^2)$, then $\mathcal{T}_R \cap \gamma$ is a non-degenerate Hermitian curve, say \mathcal{C} , meeting $\mathcal{H}(4, q^2)$ in 1 or $q + 1$ points. In this case there are $(q + 1)(q^2 - q)$ points $R \in \text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma^\perp)$ such that $\mathcal{T}_R \cap \gamma = \mathcal{C}$.

Proof. Let R be a point of $\gamma^\perp \setminus \mathcal{H}(4, q^2)$, then a line t through R that is tangent to $\mathcal{H}(4, q^2)$ meets γ in a point of $\mathcal{H}(4, q^2)$. Hence $\mathcal{T}_R \cap \gamma \subset \mathcal{H}(4, q^2)$.

Let R be a point of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma^\perp)$. The solid $\Gamma = \langle R, \gamma \rangle$ meets γ^\perp in a point, say R' , and let $R'' = \Gamma^\perp = R^\perp \cap \gamma^\perp$. Let us define γ' as the plane $R^\perp \cap \Gamma$ and r as the line $\gamma \cap \gamma'$. Note that γ' is the plane obtained by projecting γ from R onto R^\perp .

If $R' \in \mathcal{H}(4, q^2)$, then $R' = R''$, $R'^\perp = \Gamma$, the line RR' is tangent to $\mathcal{H}(4, q^2)$ at R' and the plane $\langle \gamma^\perp, RR' \rangle$ meets $\mathcal{H}(4, q^2)$ in $q^3 + 1$ points. Hence $\gamma' = R^\perp \cap \Gamma = R^\perp \cap R'^\perp = \langle RR' \rangle^\perp$ is a plane meeting $\mathcal{H}(4, q^2)$ in $q^3 + q^2 + 1$ points. Moreover, $r = \langle \gamma^\perp, RR' \rangle^\perp$ has $q + 1$ points

in common with $\mathcal{H}(4, q^2)$, since $|\langle \gamma^\perp, RR' \rangle \cap \mathcal{H}(4, q^2)| = q^3 + 1$. In this case by projecting the $q^3 + q^2 + 1$ points of $\gamma' \cap \mathcal{H}(4, q^2)$ from R onto γ , we get a Hermitian pencil of lines of γ , say \mathcal{C} , such that $\mathcal{C} \cap r = \mathcal{C} \cap \mathcal{H}(4, q^2) = r \cap \mathcal{H}(4, q^2)$. Further the vertex of \mathcal{C} is the point $V = RR' \cap \gamma$. Varying the point R' in $\gamma^\perp \cap \mathcal{H}(4, q^2)$ and by replacing R with one of the $q^2 - 1$ points of $VR' \setminus \{V, R'\}$, we obtain the same curve \mathcal{C} .

If $R' \notin \mathcal{H}(4, q^2)$, then $R' \neq R''$, $R''^\perp = \Gamma$ and the line RR'' is secant to $\mathcal{H}(4, q^2)$, since $R \in R''^\perp$, with $R \notin \mathcal{H}(4, q^2)$. Hence $\gamma' = R^\perp \cap \Gamma = R^\perp \cap R''^\perp = \langle RR'' \rangle^\perp$ is a plane meeting $\mathcal{H}(4, q^2)$ in $q^3 + 1$ points. Note that $r = \gamma \cap \gamma' = \langle R', R'' \rangle^\perp \cap \langle R, R'' \rangle^\perp = \langle R, R', R'' \rangle^\perp$. Two possibilities arise: either RR' is a line tangent to $\mathcal{H}(4, q^2)$ at the point $RR' \cap \gamma$ or RR' is secant to $\mathcal{H}(4, q^2)$ and $RR' \cap \gamma \notin \mathcal{H}(4, q^2)$. In the former case $|r \cap \mathcal{H}(4, q^2)| = 1$, since $|\langle R, R', R'' \rangle \cap \mathcal{H}(4, q^2)| = q^3 + q^2 + 1$, whereas in the latter case $|r \cap \mathcal{H}(4, q^2)| = q + 1$, since $|\langle R, R', R'' \rangle \cap \mathcal{H}(4, q^2)| = q^3 + 1$. By projecting the $q^3 + 1$ points of $\gamma' \cap \mathcal{H}(4, q^2)$ from R onto γ , we get a non-degenerate Hermitian curve of γ , say \mathcal{C} , such that $\mathcal{C} \cap r = \mathcal{C} \cap \mathcal{H}(4, q^2) = r \cap \mathcal{H}(4, q^2)$. By Lemma 2.8 there are exactly $q + 1$ points of $RR' \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma')$ which give rise to the same curve \mathcal{C} . Since $|\gamma^\perp \setminus \mathcal{H}(4, q^2)| = q^2 - q$, it follows that there are $(q + 1)(q^2 - q)$ points $R \in \text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma^\perp)$ such that $\mathcal{T}_R \cap \gamma = \mathcal{C}$. \square

Theorem 4.7. *Let P_1, P_2, P_3 be three pairwise adjacent vertices of \mathcal{G}_4 and let $\gamma = \langle P_1, P_2, P_3 \rangle$.*

1. *If γ is a line tangent to $\mathcal{H}(4, q^2)$, let s be the unique Baer subline of γ containing P_1, P_2, P_3 and let $T = \gamma \cap \mathcal{H}(4, q^2)$. Then*

$$|N_{\mathcal{G}_4}(P_1) \cap N_{\mathcal{G}_4}(P_2) \cap N_{\mathcal{G}_4}(P_3)| = \begin{cases} q^5 + q^4 - q^3 - 3 & \text{if } T \in s, \\ 2q^5 + q^4 - q^3 - 3 & \text{if } T \notin s. \end{cases}$$

2. *If γ is a plane and $\gamma \cap \mathcal{H}(4, q^2)$ is a line, then $|N_{\mathcal{G}_4}(P_1) \cap N_{\mathcal{G}_4}(P_2) \cap N_{\mathcal{G}_4}(P_3)| = q^5 + 3q^4 - 3$.*
3. *If γ is a plane and $\gamma \cap \mathcal{H}(4, q^2)$ is a non-degenerate Hermitian curve, then $|N_{\mathcal{G}_4}(P_1) \cap N_{\mathcal{G}_4}(P_2) \cap N_{\mathcal{G}_4}(P_3)| = q^5 + 2q^4 + 3q^3 - 2q^2 - q - 3$.*

Proof. Assume that γ is a line tangent to $\mathcal{H}(4, q^2)$ and let σ be a plane through γ . If $\sigma \cap \mathcal{H}(4, q^2)$ is a line, then σ contains $q^4 - q^2$ points R not in γ such that the lines RP_i , $i = 1, 2, 3$, are tangent to $\mathcal{H}(4, q^2)$. If $\sigma \cap \mathcal{H}(4, q^2)$ is Hermitian pencil of lines, then σ contains no point R not in γ such that the lines RP_i , $i = 1, 2, 3$, are tangent to $\mathcal{H}(4, q^2)$, whereas if $\sigma \cap \mathcal{H}(4, q^2)$ is a non-degenerate Hermitian curve, then from Lemma 2.1, σ contains either 0 or q points R not in γ such that the lines RP_i , $i = 1, 2, 3$, are tangent to $\mathcal{H}(4, q^2)$ according as $T \in s$ or $T \notin s$, respectively. Since there are $q + 1$ planes meeting $\mathcal{H}(4, q^2)$ in a line, $q^2 - q$ planes intersecting $\mathcal{H}(4, q^2)$ in a Hermitian pencil of lines and q^4 planes having $q^3 + 1$ points in common with $\mathcal{H}(4, q^2)$, we have that $|N_{\mathcal{G}_4}(P_1) \cap N_{\mathcal{G}_4}(P_2) \cap N_{\mathcal{G}_4}(P_3)| = (q + 1)(q^4 - q^2) + |\gamma \setminus \{T, P_1, P_2, P_3\}| = q^5 + q^4 - q^3 - 3$, if $T \in s$, and $|N_{\mathcal{G}_4}(P_1) \cap N_{\mathcal{G}_4}(P_2) \cap N_{\mathcal{G}_4}(P_3)| = (q + 1)(q^4 - q^2) + q^4 \cdot q + |\gamma \setminus \{T, P_1, P_2, P_3\}| = 2q^5 + q^4 - q^3 - 3$, if $T \notin s$.

Assume that γ is a plane and that $\gamma \cap \mathcal{H}(4, q^2)$ is a line, say ℓ . If R is a point of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma)$ such that the lines RP_i , $i = 1, 2, 3$, are tangent to $\mathcal{H}(4, q^2)$, then from Lemma 4.5,

$\mathcal{T}_R \cap \gamma$ is a Hermitian pencil of lines meeting ℓ in one point. From Lemma 2.4, in γ there are $q^2 + 2q$ Hermitian pencils of lines meeting ℓ in one point and containing P_1, P_2, P_3 . From Lemma 4.5, for each of these $q^2 + 2q$ Hermitian pencils of lines, there are q^3 points R of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma)$ such that the lines $RP_i, i = 1, 2, 3$, are tangent to $\mathcal{H}(4, q^2)$. Therefore in this case $|N_{\mathcal{G}_4}(P_1) \cap N_{\mathcal{G}_4}(P_2) \cap N_{\mathcal{G}_4}(P_3)| = q^3(q^2 + 2q) + |\gamma \setminus (\ell \cup \{P_1, P_2, P_3\})| = q^5 + 3q^4 - 3$.

Assume that γ is a plane and that $\gamma \cap \mathcal{H}(4, q^2)$ is a non-degenerate Hermitian curve. If R is a point of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma)$ such that the lines $RP_i, i = 1, 2, 3$, are tangent to $\mathcal{H}(4, q^2)$, then from Lemma 4.6, we have that $R \in \text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma^\perp)$ and either $\mathcal{T}_R \cap \gamma$ is a Hermitian pencil of lines meeting $\mathcal{H}(4, q^2)$ in $q+1$ points or $\mathcal{T}_R \cap \gamma$ is a non-degenerate Hermitian curve meeting $\mathcal{H}(4, q^2)$ in one or $q+1$ points. From Lemma 2.5, in γ there are $3q$ Hermitian pencils of lines meeting $\gamma \cap \mathcal{H}(4, q^2)$ in $q+1$ points and containing P_1, P_2, P_3 . Similarly from Proposition 2.7, in γ there are $q^2 - q + 1$ non-degenerate Hermitian curves meeting $\gamma \cap \mathcal{H}(4, q^2)$ in one or $q+1$ points and containing P_1, P_2, P_3 . From Lemma 4.6, for each of these Hermitian curves, there are either $(q+1)(q^2 - 1)$ or $(q+1)(q^2 - q)$ points R of $\text{PG}(4, q^2) \setminus (\mathcal{H}(4, q^2) \cup \gamma \cup \gamma^\perp)$ such that the lines $RP_i, i = 1, 2, 3$ are tangent to $\mathcal{H}(4, q^2)$ according as the Hermitian curve is degenerate or not, respectively. Observe that if R belongs to γ then necessarily R has to lie on a line joining two of the three points P_1, P_2, P_3 , otherwise $\gamma \cap \mathcal{H}(4, q^2)$ would contain the dual of an O'Nan configuration, a contradiction. Therefore in this case $|N_{\mathcal{G}_4}(P_1) \cap N_{\mathcal{G}_4}(P_2) \cap N_{\mathcal{G}_4}(P_3)| = 3q \cdot (q+1)(q^2 - 1) + (q^2 - q + 1) \cdot (q+1)(q^2 - q) + 3 \cdot (q-1) = q^5 + 2q^4 + 3q^3 - 2q^2 - q - 3$. \square

4.3 The isomorphism issue

Set $n = 4$. Let u be a point of ℓ_1 and let t be a line through u tangent to $\mathcal{H}(4, q^2)$ at T and contained in P^\perp , with $t \neq \bar{\ell}_1$. If $\pi \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines, then $|t \cap \mathcal{A}_1| = q^2 - q - 2$, $|t \cap \mathcal{A}_2| = 0$ and t shares with \mathcal{A} the $q+1$ points of a Baer subline. If $\pi \cap \mathcal{H}(4, q^2)$ is a line, then $|t \cap \mathcal{A}_1| = q^2 - q - 1$, $|t \cap \mathcal{A}_2| = 0$, $|t \cap \mathcal{A}| = q$ and $(t \cap \mathcal{A}) \cup \{u\}$ is Baer subline. Let u_1 and u_2 be two distinct points of \mathcal{A} and let b be the Baer subline of t containing u, u_1, u_2 . In the case when $|\pi \cap \mathcal{H}(4, q^2)| = q^3 + q^2 + 1$, assume that $T \notin b$. Observe that u, u_1, u_2 are three pairwise adjacent vertices of $\mathcal{G}_4, \mathcal{G}'_4$ and \mathcal{G}''_4 .

Proposition 4.8. $|N_{\mathcal{G}'_4}(u) \cap N_{\mathcal{G}'_4}(u_1) \cap N_{\mathcal{G}'_4}(u_2)| = |N_{\mathcal{G}''_4}(u) \cap N_{\mathcal{G}''_4}(u_1) \cap N_{\mathcal{G}''_4}(u_2)| = 2q^5 + q^3 - 3$.

Proof. From the definition of \mathcal{G}'_4 and \mathcal{G}''_4 we have that

$$\begin{aligned} N_{\mathcal{G}'_4}(u) \cap N_{\mathcal{G}'_4}(u_1) \cap N_{\mathcal{G}'_4}(u_2) &= N_{\mathcal{G}''_4}(u) \cap N_{\mathcal{G}''_4}(u_1) \cap N_{\mathcal{G}''_4}(u_2) \\ &= ((N_{\mathcal{G}_4}(u) \setminus \mathcal{A}_1) \cup \mathcal{A}_2) \cap N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2) \\ &= ((N_{\mathcal{G}_4}(u) \cap N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2)) \setminus \mathcal{A}_1) \cup (N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2) \cap \mathcal{A}_2). \end{aligned}$$

Let σ be the plane spanned by t and P . Since $\sigma \cap \mathcal{H}(4, q^2)$ is a line, it follows that every point of $\sigma \cap \mathcal{A}_1$ spans with both u_1 and u_2 a line that is tangent to $\mathcal{H}(4, q^2)$. Hence $\mathcal{A}_1 \cap \sigma \subseteq N_{\mathcal{G}_4}(u) \cap N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2)$. Note that $|\mathcal{A}_2 \cap \sigma| = 0$, whereas $|\mathcal{A}_1 \cap \sigma|$ equals $q^2(q^2 - q - 2)$ or $q^2(q^2 - q - 1)$ according as $\pi \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines or a line.

Denote by σ' a plane through t contained in P^\perp , with $\sigma' \neq \sigma$, and let $u' = \sigma' \cap \ell_2$. Then $\sigma' \cap \mathcal{H}(4, q^2)$ is a non-degenerate Hermitian curve.

If $|\pi \cap \mathcal{H}(4, q^2)|$ equals $q^3 + q^2 + 1$ (resp. $q^2 + 1$), the points of \mathcal{A}_1 contained in $\sigma' \setminus \sigma$ are on the q (resp. $q - 1$) lines of σ' tangent to $\sigma' \cap \mathcal{H}(4, q^2)$ passing through u and distinct from t (resp. t and $\sigma' \cap \pi$). By Lemma 2.2, there is a unique point R on each of these tangent lines such that $R \in N_{\mathcal{G}_4}(u) \cap N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2)$ and if $R \in N_{\mathcal{G}_4}(u) \cap N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2)$, then $R \in (\mathcal{A} \cup \mathcal{A}_1) \cap (\sigma' \setminus \sigma)$. In particular, if the case $\langle R, u' \rangle \cap t \notin \{u_1, u_2\}$ occurs, then $R \notin \mathcal{A}$, otherwise R, u_1, u_2, u' would be a dual O'Nan configuration of $\sigma' \cap \mathcal{H}(4, q^2)$, which is impossible. Hence $R \in \mathcal{A}_1$. Assume that $\langle R, u' \rangle \cap t \in \{u_1, u_2\}$. If $\pi \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines, then $R \in \mathcal{A}$, by Lemma 2.3. If $\pi \cap \mathcal{H}(4, q^2)$ is a line, then $R \in \mathcal{A}_1$, otherwise R, u_1, u, u' or R, u_2, u, u' would be a dual O'Nan configuration of $\sigma' \cap \mathcal{H}(4, q^2)$. Therefore $|N_{\mathcal{G}_4}(u) \cap N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2) \cap \mathcal{A}_1|$ equals $q^2(q^2 - q - 2) + q^2(q - 2) = q^4 - 4q^2$ or $q^2(q^2 - q - 1) + q^2(q - 1) = q^4 - 2q^2$, according as $\pi \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines or a line, respectively.

In a similar way, if $|\pi \cap \mathcal{H}(4, q^2)|$ equals $q^3 + q^2 + 1$ (resp. $q^2 + 1$), the points of \mathcal{A}_2 contained in $\sigma' \setminus \sigma$ are on the $q + 1$ (resp. q) lines of σ' tangent to $\sigma' \cap \mathcal{H}(4, q^2)$ passing through u' (and distinct from $\sigma' \cap \pi$). Each of these lines meets t in a point of \mathcal{A} . Let R be a point on one of these $q + 1$ (resp. q) lines, with $R \in N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2) \setminus (t \cup \{u'\})$. Hence $R \in (\mathcal{A} \cup \mathcal{A}_2) \cap (\sigma' \setminus \sigma)$. If $\langle R, u' \rangle \cap t \notin \{u_1, u_2\}$, then $R \notin \mathcal{A}_2$, otherwise R, u_1, u_2, u' would be a dual O'Nan configuration of $\sigma' \cap \mathcal{H}(4, q^2)$, a contradiction. Assume that $\langle R, u' \rangle \cap t \in \{u_1, u_2\}$. There are $q - 1$ possibilities for R , since the line $\langle u', u_i \rangle$ contains $q - 1$ points distinct from u' belonging to $N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2)$, by Lemma 2.2. If $\pi \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines, then exactly one of these $q - 1$ points is in \mathcal{A} by Lemma 2.3. If $\pi \cap \mathcal{H}(4, q^2)$ is a line, then for all the $q - 1$ possibilities we have that $R \in \mathcal{A}_2$, otherwise R, u_1, u, u' or R, u_2, u, u' would be a dual O'Nan configuration of $\sigma' \cap \mathcal{H}(4, q^2)$. Therefore $|N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2) \cap \mathcal{A}_2 \cap \sigma'|$ is $2(q - 2)$ or $2(q - 1)$ and hence $|N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2) \cap \mathcal{A}_2|$ equals $2(q - 2)q^2$ or $2(q - 1)q^2$, according as $\pi \cap \mathcal{H}(4, q^2)$ is a Hermitian pencil of lines or a line, respectively.

It follows that $|N_{\mathcal{G}'_4}(u) \cap N_{\mathcal{G}'_4}(u_1) \cap N_{\mathcal{G}'_4}(u_2)| = 2q^5 + q^4 - q^3 - 3 - (q^4 - 4q^2) + 2(q^3 - 2q^2) = 2q^5 + q^3 - 3$ and $|N_{\mathcal{G}''_4}(u) \cap N_{\mathcal{G}''_4}(u_1) \cap N_{\mathcal{G}''_4}(u_2)| = 2q^5 + q^4 - q^3 - 3 - (q^4 - 2q^2) + 2(q^3 - q^2) = 2q^5 + q^3 - 3$. \square

As a consequence of Theorem 4.7 and Proposition 4.8, we have the following.

Theorem 4.9. *Let $n \geq 4$. The graph \mathcal{G}_n is not isomorphic neither to \mathcal{G}'_n (if $q > 2$), nor to \mathcal{G}''_n .*

Proof. Let Π be a four-space of $\text{PG}(n, q)$ such that $\Pi \cap \mathcal{H}(n, q^2) = \mathcal{H}(4, q^2)$ and $\bar{\ell}_i \subset \Pi$, $i = 1, 2$. Then the induced subgraph by \mathcal{G}_n or \mathcal{G}'_n or \mathcal{G}''_n on $\Pi \setminus \mathcal{H}(4, q^2)$ is \mathcal{G}_4 or \mathcal{G}'_4 or \mathcal{G}''_4 . Let u, u_1, u_2 be three pairwise adjacent vertices of \mathcal{G}_4 , \mathcal{G}'_4 and \mathcal{G}''_4 . By Theorem 4.7, $|N_{\mathcal{G}_4}(u) \cap N_{\mathcal{G}_4}(u_1) \cap N_{\mathcal{G}_4}(u_2)|$ takes four different values. By Proposition 4.8, both $|N_{\mathcal{G}'_4}(u) \cap N_{\mathcal{G}'_4}(u_1) \cap N_{\mathcal{G}'_4}(u_2)|$ and $|N_{\mathcal{G}''_4}(u) \cap N_{\mathcal{G}''_4}(u_1) \cap N_{\mathcal{G}''_4}(u_2)|$ take at least five different values. \square

5 Conclusions

We have seen that if $n \neq 3$, then $\text{NU}(n + 1, q^2)$ is not determined by its spectrum. Regarding the case $n = 3$, there are exactly 28 strongly regular graphs cospectral with $\text{NU}(4, 4)$, see [16]. A strongly regular graph having the same parameters as $\text{NU}(4, 9)$ and admitting $2 \times \text{U}(4, 4)$

as a full automorphism group has been constructed in [5]. We ask whether or not the graph $\text{NU}(4, q^2)$, $q \geq 4$, is determined by its spectrum.

Acknowledgments. The first author is supported by a postdoctoral fellowship of the Research Foundation - Flanders (FWO).

References

- [1] S.G. Barwick, G.L. Ebert, *Unitals in projective planes*, Springer Monographs in Mathematics. Springer, New York, 2008.
- [2] F. Buekenhout, Existence of unitals in finite translation planes of order q^2 with a kernel of order q , *Geom. Dedicata*, 5 (2), (1976), 189–194.
- [3] A.E. Brouwer, W.H. Haemers, *Spectra of graphs*, Universitext, Springer, New York, 2012.
- [4] A.A. Bruen, J.C. Fisher, An observation on certain point-line configurations in classical planes, *Discrete Math.*, 106/107, (1992), 93–96.
- [5] D. Crnković, S. Rukavina, A. Švob, New strongly regular graphs from orthogonal groups $O^+(6, 2)$ and $O^-(6, 2)$, *Discrete Math.*, 341, (2018), no. 10, 2723–2728.
- [6] T. Feng, W. Li, On the existence of O’Nan configurations in ovoidal Buekenhout–Metz unitals in $\text{PG}(2, q^2)$, *Discrete Math.*, 342 (8), (2019), 2324–2332.
- [7] L. Giuzzi, Collineation groups of the intersection of two classical unitals, *J. Combin. Des.*, 9, (2001), 445–459.
- [8] J.W.P. Hirschfeld, *Projective Geometries over Finite Fields*, Oxford Mathematical Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1998.
- [9] J.W.P. Hirschfeld, *Finite projective spaces of three dimensions*, Oxford Mathematical Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1985.
- [10] J.W.P. Hirschfeld, J.A. Thas, *General Galois Geometries*, Oxford Mathematical Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1991.
- [11] D.R. Hughes, F. Piper, *Design theory*, Cambridge University Press, Cambridge, 1985.
- [12] B.C. Kestenband, Unital intersections in finite projective planes, *Geom. Dedicata*, 11, (1981), 107–117.
- [13] F. Ihringer, A. Munemasa, New strongly regular graphs from finite geometries via switching, *Linear Algebra Appl.*, 580, (2019), 464–474.

- [14] R. Metz, On a class of unitals, *Geom. Dedicata*, 8 (1), (1979), 125–126.
- [15] M.E. O’Nan, Automorphisms of unitary block designs, *J. Algebra*, 20, (1972), 495–511.
- [16] E. Spence, The strongly regular $(40, 12, 2, 4)$ graphs, *Electron. J. Combin.*, 7, (2000), R22, 4p.
- [17] W. Wang, L. Qiu, Y. Hu, Cospectral graphs, GM-switching and regular rational orthogonal matrices of level p , *Linear Algebra Appl.*, 563, (2019), 154–177.