

**PHASE RETRIEVAL BY PROJECTIONS IN \mathbb{R}^n REQUIRES
 $2n - 2$ PROJECTIONS
 (THIS PAPER HAS A GAP)**

PETER G. CASAZZA AND DORSA GHOREISHI

ABSTRACT. We will answer the most significant open problem in real phase retrieval by projections by showing it requires at least $2n - 2$ projections to do phase retrieval in \mathbb{R}^n .

1. INTRODUCTION

Phase retrieval is one of the most active and applied subjects in mathematics today, with applications to pure mathematics, applied mathematics, engineering, medicine, computer science, and more. Frame theory is fundamental to the digitalizing of information for processing and storing. Phase retrieval has applications to X-ray crystallography, electron microscopy, astronomical imaging, optics, quantum physics, and much more. Phase retrieval will even be needed to align the mirrors in the new James Webb Space Telescope.

Definition 1.1. *A family of vectors $\{\phi_i\}_{i=1}^m$ in \mathbb{R}^n or \mathbb{C}^n does **phase retrieval** if for any non-zero $x, y \in \mathbb{R}^n$ satisfying $|\langle x, \phi_i \rangle| = |\langle y, \phi_i \rangle|$ for all $i \in [m]$ this will imply $x = cy$ with $|c| = 1$.*

A fundamental idea here is:

Definition 1.2. *A family of vectors $\{\phi_i\}_{i=1}^m$ in \mathbb{R}^n has the **complement property** if for all subsets $I \subset [m]$, either $\text{span}\{\phi_i\}_{i \in I} = \mathbb{R}^n$ or $\text{span}\{\phi_i\}_{i \in I^c} = \mathbb{R}^n$.*

The most important classification of phase retrieval in \mathbb{R}^n appeared in [1].

Theorem 1.3. *A family of vectors $\{\phi_i\}_{i=1}^m$ does phase retrieval in \mathbb{R}^n if and only if it has the complement property.*

Corollary 1.4. *If a family of vectors $\{\phi_i\}_{i=1}^m$ satisfies phase retrieval in \mathbb{R}^n , then $m \geq 2n - 1$.*

In some applications a signal must be reconstructed from the norms of higher dimensional components. In X-ray crystallography such a problem

2020 *Mathematics Subject Classification.* 42C15.

The authors were supported by NSF DMS 1609760 and 1906725.

arises in crystal twinning [7]. In this scenario, there exists a similar phase retrieval problem.

Definition 1.5. A family of subspaces $\{W_i\}_{i=1}^m$ in \mathbb{R}^n or \mathbb{C}^n with respective orthogonal projections $\{P_i\}_{i=1}^m$ satisfy **phase retrieval** if for all $x, y \in \mathbb{R}^n$, $\|P_i x\| = \|P_i y\|$ for all $i \in [m]$, implies $x = cy$ with $|c| = 1$.

A deep study of phase retrieval by projections appears in [4]. This includes the result:

Theorem 1.6. For any $0 < r_i < n$, $i \in [2n-1]$ there are subspaces $\{W_i\}_{i=1}^{2n-1}$ in \mathbb{R}^n doing phase retrieval with $\dim W_i = r_i$.

A fundamental result in this area from [4] is:

Theorem 1.7. Let $\{W_i\}_{i=1}^m$ be subspaces of \mathbb{R}^n . The following are equivalent:

- (1) $\{W_i\}_{i=1}^m$ does phase retrieval.
- (2) Whenever we choose orthonormal bases $\{u_{ij}\}_{j=1}^{I_i}$ for W_i , the family of vectors $\{u_{ij}\}_{i=1, j=1}^m$ does phase retrieval.

Eddin [8] proved a fundamental result concerning phase retrieval by projections (See [5] for an elementary proof of this result.)

Theorem 1.8. A family of projections $\{P_i\}_{i=1}^m$ does phase retrieval in \mathbb{R}^n if and only if for every $0 \neq x \in \mathbb{R}^n$, $\text{span}\{P_i x\}_{i=1}^m = \mathbb{R}^n$.

The following theorem in [11] shows a particular case of \mathbb{R}^4 in which phase retrieval is possible with $2n - 2 = 2(4) - 2 = 6$ subspaces.

Theorem 1.9. There are 6 two-dimensional subspaces of \mathbb{R}^4 satisfying phase retrieval.

In [3] it was shown that there are 6 hyperplanes in \mathbb{R}^4 doing phase retrieval.

The following result also appears in [3]. Since this result is central to this paper, we include its short proof.

Theorem 1.10. It takes at least $2n - 2$ hyperplanes in \mathbb{R}^n to do phase retrieval.

Proof. Let $\{W_i\}_{i=1}^m$ be a family of hyperplanes in \mathbb{R}^n doing phase retrieval with respective projections $\{P_i\}_{i=1}^m$.

By way of contradiction, assume $m \leq 2n - 3$. Choose a vector

$$0 \neq x \in \bigcap_{i=1}^{n-1} W_i$$

so $P_i x = x$ for $i \in [n-1]$. Therefore we can write, $\{P_i x\}_{i=1}^m = \{x\} \cup \{P_i x\}_{i=n}^m$. Notice that $2n - 3 - (n - 1) + 1 = n - 1$ which shows $\{P_i x\}_{i=1}^m$ has at most

$n - 1$ non-zero vectors and cannot span the space. This is contradicting Theorem 1.8.

□

The main open question in this area has been about finding the least number of subspaces satisfying phase retrieval. We will show in this paper that it takes at least $2n - 2$ subspaces to satisfy phase retrieval in \mathbb{R}^n .

Another open problem in this area is motivated by this theorem of Edidin [8]:

Theorem 1.11. *If $n = 2^k - 1$, for some k , then it takes $2n - 1$ subspaces of \mathbb{R}^n to do phase retrieval.*

So now the main open problem in this area is:

Problem 1.12. *Can we do phase retrieval in \mathbb{R}^n with $2n - 2$ subspaces whenever $n \neq 2^k - 1$ for any k .*

The complex case is significantly more complicated. It is known [2] that phase retrieval can be done in \mathbb{C}^n with $4n - 4$ vectors. It was believed this result was best possible. But Vinzant [9] showed that phase retrieval can be done in \mathbb{C}^4 with 11 vectors (in other words $4n - 5$ vectors). So the major open problem here is:

Problem 1.13. *What is the minimum number of vectors required to do phase retrieval in \mathbb{C}^n ?*

Phase retrieval by projections in \mathbb{C}^n is still unknown.

Problem 1.14. *What is the minimum number of projections required to do phase retrieval in \mathbb{C}^n*

Notice that Theorem 1.8 (Edidin's theorem) fails in \mathbb{C}^n . A complex version of this theorem appears in [11], Theorem 2.1 (3). A completely different approach to a complex version of Edidin's theorem was proved in [6]. In this paper they associate vectors in \mathbb{C}^n with rank 2 projections in \mathbb{R}^{2n} .

2. MAIN RESULT

The following is immediate by Theorem 1.8.

Lemma 2.1. *If a family of projections $\{P_i\}_{i=1}^m$ does phase retrieval in \mathbb{R}^n , then for every $0 \neq x \in \mathbb{R}^n$, either $\text{span}\{P_i x\}_{i=2}^m = \mathbb{R}^n$ or $\text{span}\{P_i x\}_{i=2}^m$ spans a hyperplane.*

Moreover, if this is a hyperplane, $P_1 x \notin \text{span}\{P_i x\}_{i=2}^m$.

The next result is fundamental to our proof.

Theorem 2.2. *Let a family of projections $\{P_i\}_{i=1}^m$ do phase retrieval on \mathbb{R}^n and let*

$$F = \{x : \|x\| = 1 \text{ and } H_x = \text{span}\{P_i x\}_{i=2}^m \neq \mathbb{R}^n\}.$$

If $x_1, x_2 \in F$ and $H_{x_1} = H_{x_2}$, then $x_1 = \pm x_2$.

Proof. Since $H_{x_1} = H_{x_2}$ is a hyperplane, there is a vector $\|z\| = 1$ so that $z \perp P_i x_1$ and $z \perp P_i x_2$ for all $i = 2, 3, \dots, m$. Hence

$$0 = \langle z, P_i x_1 \rangle = \langle P_i z, x_1 \rangle = \langle z, P_i x_2 \rangle = \langle P_i z, x_2 \rangle, \text{ for all } i = 2, 3, \dots, m.$$

Since this family spans a hyperplane, x_1, x_2 are both orthogonal to this hyperplane and so $x_1 = \pm x_2$. \square

The main result will be based on the following theorem:

Theorem 2.3. *Let a family of subspaces $\{W_i\}_{i=1}^m$ do phase retrieval in \mathbb{R}^n . If $\dim W_1^\perp \geq 2$, there is a $z \in W_1^\perp$ with $\|z\| = 1$ so that $\{W_1 \cup z\} \cup \{W_2, \dots, W_m\}$ does phase retrieval.*

Proof. We will do this in steps.

(1) Choose orthonormal bases $\{u_{ij}\}_{j=1}^{I_i}$ for W_i when $i \in [m]$ so that these vectors satisfy phase retrieval. For $z \in W_1^\perp$ let $W_z = \text{span}\{W_1, z\}$ and let P_z be the projection onto W_z .

We need to show there exists a $z \in W_1^\perp$ so that $\text{span}\{P_z, P_2, \dots, P_m\}$ does phase retrieval in \mathbb{R}^n .

(2) Choose $x \in \mathbb{R}^n$ with $\|x\| = 1$. If $\text{span}\{P_i x\}_{i=2}^m = \mathbb{R}^n$, this shows that if we add a vector z which is in W_1^\perp , the subspaces still satisfy phase retrieval and we are done for this x .

Now we need to consider the cases when $\text{span}\{P_i x\}_{i=2}^m \neq \mathbb{R}^n$,

(3) By Theorem 2.2, if $\text{span}\{P_i x\}_{i=2}^m = \text{span}\{P_i y\}_{i=2}^m \neq \mathbb{R}^n$ and $\|x\| = \|y\|$, then $x = \pm y$.

(4) Note

$$\text{span}\{P_i x\}_{i=2}^m = \text{span}\{u_{kj} : \langle P_i x, u_{kj} \rangle \neq 0 \text{ for some } i = 1, 2, \dots, m\}.$$

It follows from (3) that there are at most a finite number of x 's for which $\text{span}\{P_i x\}_{i=2}^m \neq \mathbb{R}^n$. Call these $\{x_j\}_{j=1}^k$, and let $V_{x_j} = \text{span}\{P_i x_j\}_{i=2}^m$ and choose $\|y_j\| = 1$ with $y_j \perp V_{x_j}$.

(5) Now we need to find a $z \in W_1^\perp$ with $\|z\| = 1$ and $P_z x_j \notin V_{x_j}$ for all $j \in [k]$.

Since there are a finite number of V_{x_j} , it suffices to show that for all $j \in [k]$ if

$$(1) \quad A = \{z \in W_1^\perp : \|z\| = 1, P_z x_j \in V_{x_j} = y_j^\perp, \}$$

the family of lines in the unit ball of A going through the origin is a set of measure zero relative to Lebesgue measure on W_1^\perp .

$$(6) \text{ We have } A \cap y_j^\perp = A \cap x_j^\perp = \emptyset.$$

- If $z \in A \cap x_j^\perp$ then:

$$\begin{aligned} P_z x_j &= P_1 x_j + \langle x_j, z \rangle z \\ &= P_1 x_j \in y_j^\perp. \end{aligned}$$

It follows that $\text{span}\{P_i x_j\}_{i=1}^m = y_j^\perp \neq \mathbb{R}^n$. So by Theorem 1.8, $\{P_i\}_{i=1}^m$ fails phase retrieval - a contradiction.

- If $z \in A \cap y_j^\perp$ then

$$P_z x_j = P_1 x_j + \langle x_j, z \rangle z \in y_j^\perp.$$

Since y_j^\perp is a subspace and $z \in y_j^\perp$ it follows again that $P_1 x_j \in y_j^\perp$ and so $\{P_i\}_{i=1}^m$ fails phase retrieval - again a contradiction.

(7) **Claim:** There is a constant $0 < c \leq 1$ so that for every $z \in A$ we have

$$\langle x_j, z \rangle \langle y_j, z \rangle = c.$$

Proof of the claim: Let $z_1, z_2 \in A$. Then if $P_{z_1} x_j, P_{z_2} x_j \in y_j^\perp$, since y_j^\perp is a subspace,

$$P_{z_1} x_j - P_{z_2} x_j = (P_1 x_j + \langle x_j, z_1 \rangle z_1) - (P_1 x_j + \langle x_j, z_2 \rangle z_2) \in y_j^\perp.$$

Hence

$$\langle x_j, z_1 \rangle \langle y_j, z_1 \rangle - \langle x_j, z_2 \rangle \langle y_j, z_2 \rangle = 0.$$

Notice that $c \neq 0$, since if $c = 0$, then for every $z \in A$ either $z \perp x_j$ or $z \perp y_j$ which is not possible by (6).

(8) After a rotation, we may assume $X = \text{span}\{x_j, y_j\} = \mathbb{R}^2 \subset \mathbb{R}^n$ with canonical basis $\{e_1, e_2\}$ and $x_j = e_1$.

Let $y_j = (e, f) \in \mathbb{R}^2$ and let P be the projection onto X . We want to find the points $v \in X$ so that $\langle x_j, v \rangle \langle y_j, v \rangle = c$.

The points on the line ℓ_1 orthogonal to $x_j = e_1$ and passing through $(x, 0)$ represents all the points in \mathbb{R}^2 whose inner product with x_j is x . Let m be the slope of the line orthogonal to y_j . The points on the line ℓ_2 orthogonal

to y_j and passing through $\frac{c}{x}(e, f)$ are the points in \mathbb{R}^2 whose inner product with y_j is $\frac{c}{x}$ since $e^2 + f^2 = 1$. This line has equation:

$$y - \frac{c}{x}f = m(x - \frac{c}{x}e).$$

That is:

$$y = mx + \frac{c}{x}(f - em).$$

So if $v \in \ell_1 \cap \ell_2$ for some $c \leq x \leq 1$ then

$$v = (x, mx + \frac{c}{x}(f - em)).$$

Therefore, the points $v \in X$ so that $\langle x_j, v \rangle \langle y_j, v \rangle = c$ are:

$$\{(x, mx + \frac{c}{x}(f - em)) : c \leq x \leq 1\} \quad (*)$$

(9) We will answer the case where $\dim W_1^\perp = 2$. We will examine 3 cases.

Case 1: P maps W_1^\perp onto X .

Since P is onto, there is a $v \in W_1^\perp$ with $\|v\| = 1$ and $Pv = re_1$. Let v^\perp be the unit vector in W_1^\perp orthogonal to v and $y_j = (e, f) \in \mathbb{R}^2$. Let $Pv^\perp = (g, h)$. If $z \in A$, $z = av^\perp + bv$ and $a^2 + b^2 = 1$. By (8) and the argument above, $Pz = (x, mx + \frac{c}{x}(f - em))$. Also,

$$\begin{aligned} Pz &= aPv^\perp + bPv \\ &= (ag, ah) + (br, 0) \\ &= (ag + br, ah). \end{aligned}$$

So in order for $\langle x, Pz \rangle \langle y, Pz \rangle = c$ we must have for some x :

$$ag + \sqrt{1 - a^2}r = x \text{ and } ah = mx + \frac{c}{x}(f - em).$$

Substituting the first equation into the second equation for x , since all constants except a are fixed, we have a quadratic equation in a . So A is of cardinality at most two and the lines through A are a set of measure zero in W_1^\perp .

Case 2: $PW_1^\perp = \emptyset$.

In this case, if $z \in A$ then $Pz = 0$. So $\langle z, x_j \rangle = \langle Pz, x_j \rangle = 0$. By (6), this is a contradiction unless $A = \emptyset$.

Case 3: PW_1^\perp is dimension one in X .

In this case, $x_j^\perp \cap W_1^\perp = y_j^\perp \cap W_1^\perp$. So $x_j^\perp \cap W_1^\perp$ is a hyperplane in W_1^\perp . Let $y_j = (e, f) \in \mathbb{R}^2$ and choose $\|v\| = 1$ in W_1^\perp with $Pv^\perp = 0$. Let $Pv = (g, h)$.

Then, if $z \in A$ with $z = av + bv^\perp$ by (8) and (*),

$$Pz = (ag, ah) = (x, mx + \frac{c}{x}(f - em))$$

Hence,

$$ag = x \text{ and } ah = mx + \frac{c}{x}(f - em).$$

Substituting the first equation in the second equation for x , all constants except a are fixed so this is a quadratic equation for a and hence has at most two solutions. That is, A consists of at most two lines and so is a set of measure zero in W_1^\perp .

(10) We check the case $\dim W_1^\perp \geq 3$.

Since $x_j^\perp \cap y_j^\perp$ is a subset of \mathbb{R}^n of codimension 1 or 2, we will examine these two cases.

Case 1: $(x_j^\perp \cap y_j^\perp) \cap W_1^\perp$ is a set of codimension 2 in W_1^\perp .

Choose a two dimensional subspace V of W_1^\perp so that $W_1^\perp = V \oplus (x_j^\perp \cap y_j^\perp)$.

Claim: $PV = X$.

Proof of claim: If not, then there is a $v \in V$ with $Pv = 0$. That is,

$$\langle v, x_j \rangle = \langle Pv, x_j \rangle = 0 = \langle v, y_j \rangle.$$

But $v \perp (x_j^\perp \cap y_j^\perp)$, a contradiction.

Choose $v \in V$, $\|v\| = 1$ and $Pv = re_1$. Let v^\perp be in V and orthogonal to v and $Pv^\perp = (g, h)$. It follows that every $z \in W_1^\perp$ is of the form

$$z = bv + av^\perp + u$$

when $a^2 + b^2 + \|u\|^2 = 1$, and $P(u) = 0$. By (8) and (*),

$$Pz = (x, mx + \frac{c}{x}(f - em))$$

But

$$\begin{aligned} Pz &= bPv + aPv^\perp \\ &= (br, 0) + (ag, ah) \\ &= (ag + br, ah). \end{aligned}$$

So we must have for some x :

$$ag + br = x \text{ and } ah = mx + \frac{c}{x}(f - em).$$

Now, from the second equation

$$ah = m(ag + bj) + \frac{c}{ag + br}(f - em),$$

All the constants here are fixed except a,b. This equation, represents a quadratic equation in a,b. Hence,

$$\{z = av^\perp + bv : \langle x_j, Pz \rangle \langle y_j, Pz \rangle = c\}$$

is the union of two continuous quadratic graphs G_1, G_2 in V . For $z \in A$, $z = av + bv^\perp + u$, $\|z\| = 1$. So

$$A = \{x + u : x \in G_1, \|u\|^2 = 1 - \|x\|^2\}$$

or

$$A = \{x + u : x \in G_2, \|u\|^2 = 1 - \|x\|^2\}.$$

In each case, since the unit ball of $(x_j^\perp \cap y_j^\perp) \cap W_1^\perp$ is a set of measure zero in W_1^\perp and A is a curve on the sphere of W_1^\perp and since $\dim W_1^\perp \geq 3$, the lines through A are a set of measure zero in W_1^\perp .

Case 2: $(x_j^\perp \cap y_j^\perp) \cap W_1^\perp$ is a set of codimension 1 in W_1^\perp .

Choose $v \in W_1^\perp$ with $v \perp x_j$. Then every $z \in W_1^\perp$, $\|z\| = 1$, is of the form $z = av + u$ and $u \in x_j^\perp$. Let $Pv = (g, h)$. Now, $Pz = Pav$ so

$$Pz = aPv = (ag, ah).$$

Hence, by (8) and (*),

$$ag = x \text{ and } ah = mx + \frac{c}{x}(f - em).$$

This yields

$$ah = mag + \frac{c}{ag}(f - em).$$

This quadratic equation in a has at most two solutions a_1, a_2 . So A is of the form

$$\{a_i v + u : \|u\|^2 = 1 - a_i^2\}.$$

But these sets are two translates of a circle in $(x_j^\perp \cap y_j^\perp) \cap W_1^\perp$ and the lines through these sets form a set of measure zero in W_1^\perp .

(11) Now, for any x_j the set of all $z \in W_1^\perp$ with $P_z x_j \in y_1^\perp$ is a set of measure zero in W_1^\perp . So choose $z \in W_1^\perp$ which is in none of these k subsets of W_1^\perp of measure zero. Now, for any $0 \neq x \in \mathbb{R}^n$, if $x \neq x_j$ for any $j \in [k]$, then $\{P_i x\}_{i=2}^m$ spans \mathbb{R}^n . And if $x = x_j$ for some $j \in [k]$ then $P_z x_j \notin y_j^\perp$ and so $\{P_z x_j, P_i x_j\}_{i=2}^m$ spans \mathbb{R}^n . That is, $\{P_z, P_i\}_{i=2}^m$ does phase retrieval in \mathbb{R}^n .

This completes the proof of the theorem. \square

As a consequence, applying the theorem over and over, we have

Corollary 2.4. *Assume a family of subspaces $\{W_i\}_{i=1}^m$ with $\dim W_i = k_i$, $1 \leq k_i \leq n - 1$, for all $i = 1, 2, \dots$, does phase retrieval in \mathbb{R}^n . The following hold:*

- (1) *For any $1 \leq k_i \leq r_i \leq n - 1$, there are subspaces W'_i of \mathbb{R}^n so that $W_i \subset W'_i$, $\dim W'_i = r_i$ and $\{W'_i\}_{i=1}^m$ does phase retrieval.*
- (2) *There are hyperplanes $\{W'_i\}_{i=1}^m$ with $W_i \subset W'_i$ and $\{W'_i\}_{i=1}^m$ does phase retrieval. So by Theorem 1.10, $m \geq 2n - 2$.*

REFERENCES

- [1] R. Balan, P.G. Casazza, and D. Edidin, *On Signal Reconstruction Without Phase*, Appl. Comput. Harmon. Anal., **20** (3) (2006) 345-356.
- [2] A.S. Bandeira, J. Cahill, D. Mixon and A.A. Nelson, *Saving phase: injectivity and stability for phase retrieval*, Appl. and Comput. Harmonic Anal., **37** (1) (2014) 106-125.
- [3] Sara Botelho-Andrade, Peter G. Casazza, Desai Cheng, John I. Haas, Tin T. Tran, Janet C. Tremain, and Zhiqiang Xu, *Real Phase Retrieval by orthogonal complements and hyperplanes*, Proceedings of SPIE (2017).
- [4] J. Cahill, P.G. Casazza, J. Peterson and L. Woodland, *Phase retrieval by projections*, Houston Journal of Mathematics **42** No. 2 (2016) 537-558.
- [5] P.G. Casazza, *An elementary proof of a fundamental result in frame theory*, Arxiv 2010.07058.
- [6] D. Cheng and P.G. Casazza, *Associating vectors in C^n with rank-2 projections in \mathbb{R}^{2n} : with applications*. arxiv: 1703.02657.
- [7] J. Drenth, *Principles of protein x-ray crystallography*, Springer, 2010.
- [8] D. Edidin, *Projections and phase retrieval*, Appl. Comput. Harmon. Anal., **42** (2)(2017) 350-359.
- [9] C. Vinzant, *A small frame and a certificate of its injectivity*, Proceedings of the International conference on Sampling Theory and Applications, 7148879 (2015), 197-200.
- [10] Y. Wang and Z. Zu, *Generalized phase retrieval: measurement number, matrix recovery and beyond*, ArXiv 1605.08034.
- [11] Z. Xu, *The minimal measurement number for low rank matrix recovery*, Applied and Computational Harmonic Analysis (2018).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211-4100

Email address: Casazzap@missouri.edu

Email address: dorsa.ghoreishi@slu.edu