

# APPROXIMATE OBLIQUE DUAL FRAMES

JORGE P. DÍAZ<sup>1</sup>, SIGRID B. HEINEKEN<sup>2</sup>, AND PATRICIA M. MORILLAS<sup>1,\*</sup>

<sup>1</sup>*Instituto de Matemática Aplicada San Luis, UNSL-CONICET, Ejército de los Andes 950, 5700 San Luis, Argentina*

<sup>2</sup>*Departamento de Matemática, FCEyN, Universidad de Buenos Aires, Pabellón I, Ciudad Universitaria, IMAS, UBA-CONICET, C1428EGA C.A.B.A., Argentina*

**ABSTRACT.** In representations using frames, oblique duality appears in situations where the analysis and the synthesis has to be done in different subspaces. In some cases, we cannot obtain an explicit expression for the oblique duals and in others there exists only one oblique dual frame which has not the properties we need. Also, in practice the computations are not exact. To give a solution to these problems, in this work we introduce and investigate the notion of approximate oblique dual frames first in the setting of separable Hilbert spaces. We focus then on approximate oblique dual frames in shift-invariant subspaces of  $L^2(\mathbb{R})$ . We give different conditions on the generators that assure the existence of approximate oblique dual frames. We present an example that illustrates the results.

**Key words:** Frames, Oblique dual frames, Approximate dual frames, Oblique projections, Shift-invariant spaces.

**AMS subject classification:** Primary 42C15; Secondary 42C40, 46C05, 41A30.

## 1. INTRODUCTION

A frame is a sequence of vectors in a separable Hilbert space. Frames generalize bases [3, 6, 7, 10, 17, 21]. The main difference to bases is that the reconstruction of each vector is not necessarily unique, i.e. there exists more than one collection of coefficients to represent each vector as a combination of the elements of a frame. These frame coefficients are associated to other sequences called dual frames.

In practice, the computation of duals is not exact and sometimes it is even not possible to give in theory an analytic expression of it. Another difficulty is that in applications we need to truncate the frame representations in order to work with finite sequences of numbers. Also the frame coefficients can in general only be computed approximately. Approximate dual frames appear as an answer to these problems [5, 9, 15, 22]. They are also easier to build and can be adapted to our needs.

For frames in a subspace, the reconstruction can also be done with coefficients that depend on duals that do not necessarily belong to the same subspace. These are known as oblique dual frames [1, 13, 4, 11, 12, 14, 18]. In this situation we again

---

\* Corresponding author. E-mail address: morillas.unsl@gmail.com  
*E-mail addresses:* jpdiaz1179@gmail.com (J. P. Díaz), sheinek@dm.uba.ar (S. B. Heineken), morillas.unsl@gmail.com (P. M. Morillas).

have the limitations and the computational difficulties described before. Moreover, sometimes the subspaces are given by the problem and there exists a unique oblique dual frame. This unique oblique dual could need to be improved in some aspects. In order to respond to these different problems, in this work we introduce the concept of approximate oblique dual frames and study its properties. Since in general there exists more than one approximate oblique dual, we also gain freedom in its construction.

In Section 2 we give a brief review of existing definitions and results.

In Section 3 we introduce the notion of approximate oblique dual frames and investigate some of its properties. We also give its interpretation in the setting of signal processing theory and the conditions of uniqueness and consistency in order to have a good approximation of a signal.

In Section 4 we obtain characterizations of approximate oblique dual frames, in terms of series expansions as well as in terms of operators.

In Section 5 we extend to the oblique setting an important property of approximate dual frames, that is to obtain a reconstruction of the vectors as close as we desire starting from any pair of approximate oblique dual frames. We also show how to construct oblique dual frames from approximate oblique dual frames.

In Section 6 we obtain approximate oblique dual frames from the perturbation of oblique duals. We prove that if two frames are “close”, each oblique dual frame of any of them is an approximate oblique dual frame of the other, expressing its bounds in terms of the original ones.

In Section 7 we consider shift-invariant subspaces of  $L^2(\mathbb{R})$  and obtain conditions on their generators to provide an approximate reconstruction in one of them. The technique introduced in the proof of the main result of this section (Theorem 7.2) is the key for the obtention of the central results of section 8 about approximate oblique dual frames.

We begin Section 8 stating sufficient conditions for approximate oblique duality, that are derived from results of section 7. We then give an expression for the Fourier transform of the oblique projection when the spaces are shift-invariant. We use it to provide a more manageable sufficient condition, and also a necessary condition on the generators of shift-invariant subspaces for the existence of approximate oblique dual frames. We finally consider the necessity of working in the applications with approximate oblique dual frames. To illustrate their behavior, we present an example with frame sequences of translates generated by  $B$ -splines.

## 2. PRELIMINARIES

We consider  $\mathcal{H}, \mathcal{K}$  separable Hilbert spaces. The space of bounded operators from  $\mathcal{H}$  to  $\mathcal{K}$  will be denoted by  $L(\mathcal{H}, \mathcal{K})$ . For  $T \in L(\mathcal{H}, \mathcal{K})$  denote the image, the null space and the adjoint of  $T$  by  $\mathcal{R}(T)$ ,  $\mathcal{N}(T)$  and  $T^*$ , respectively. If  $T$  has closed range we also consider the Moore-Penrose pseudo-inverse of  $T$  denoted by  $T^\dagger$ . The inner product and the norm in  $\mathcal{H}$  will be denoted by  $\langle \cdot, \cdot \rangle$  and  $\| \cdot \|$ , respectively.

The Lebesgue measure of a measurable subset  $A$  of  $\mathbb{R}$  will be denoted by  $|A|$ . We denote the characteristic function of  $A$  with  $\chi_A$ .

We now review briefly definitions and present properties that we use later. Unless we say the contrary, in the sequel  $\mathcal{V}$  and  $\mathcal{W}$  will be two closed subspaces of  $\mathcal{H}$  such that  $\mathcal{H} = \mathcal{V} \oplus \mathcal{W}^\perp$ .

**2.1. Oblique projections.** The oblique projection onto  $\mathcal{V}$  along  $\mathcal{W}^\perp$ , is the unique operator that satisfies

$$\begin{aligned}\pi_{\mathcal{V}\mathcal{W}^\perp} f &= f \text{ for all } f \in \mathcal{V}, \\ \pi_{\mathcal{V}\mathcal{W}^\perp} f &= 0 \text{ for all } f \in \mathcal{W}^\perp.\end{aligned}$$

Equivalently,  $\mathcal{R}(\pi_{\mathcal{V}\mathcal{W}^\perp}) = \mathcal{V}$  and  $\mathcal{N}(\pi_{\mathcal{V}\mathcal{W}^\perp}) = \mathcal{W}^\perp$ . Observe that  $(\pi_{\mathcal{V}\mathcal{W}^\perp})^* = \pi_{\mathcal{V}\mathcal{W}^\perp}$ . If  $\mathcal{W} = \mathcal{V}$  we obtain the orthogonal projection onto  $\mathcal{W}$ , which we denote by  $P_{\mathcal{W}}$ . The properties involving oblique projections presented in this section will be used throughout the paper.

The concept of angle between two closed subspaces that we will use is the following:

**Definition 2.1.** Given  $\mathcal{W}$  y  $\mathcal{V}$  two closed subspaces of  $\mathcal{H}$ , we define *angle from  $\mathcal{V}$  to  $\mathcal{W}$*  as the unique real number  $\theta(\mathcal{V}, \mathcal{W}) \in [0, \frac{\pi}{2}]$  such that

$$(2.1) \quad \cos \theta(\mathcal{V}, \mathcal{W}) = \inf_{v \in \mathcal{V}, \|v\|=1} \|P_{\mathcal{W}} v\|.$$

**Theorem 2.2.** [19] *Let  $\mathcal{V}, \mathcal{W}$  be closed subspaces of a separable Hilbert space  $\mathcal{H}$ . Then the following assertions are equivalent:*

- (i)  $\mathcal{H} = \mathcal{W} \oplus \mathcal{V}^\perp$ .
- (ii)  $\mathcal{H} = \mathcal{V} \oplus \mathcal{W}^\perp$ .
- (iii)  $\cos(\mathcal{V}, \mathcal{W}) > 0$  and  $\cos(\mathcal{W}, \mathcal{V}) > 0$ .

We have  $\|\pi_{\mathcal{W}\mathcal{V}^\perp}\| = \frac{1}{\cos(\mathcal{W}, \mathcal{V})} \geq 1$  [2].

**2.2. Frames.** The concept of frame has been introduced by Duffin and Schaeffer in [10]. Using a frame, each element of a Hilbert space has a representation which in general is not unique. This flexibility makes them attractive for many applications involving signal expansions.

We will now recall the concept of frame for a closed subspace of  $\mathcal{H}$ .

**Definition 2.3.** Let  $\mathcal{W}$  be a closed subspace of  $\mathcal{H}$  and  $\{f_k\}_{k=1}^\infty \subset \mathcal{W}$ . Then  $\{f_k\}_{k=1}^\infty$  is a *frame* for  $\mathcal{W}$ , if there exist constants  $0 < \alpha \leq \beta < \infty$  such that

$$(2.2) \quad \alpha \|f\|^2 \leq \sum_{k=1}^{\infty} |\langle f, f_k \rangle|^2 \leq \beta \|f\|^2 \text{ for all } f \in \mathcal{W}.$$

If the right inequality in (2.2) is satisfied,  $\{f_k\}_{k=1}^\infty$  is a *Bessel sequence* for  $\mathcal{W}$ . The constants  $\alpha$  and  $\beta$  are the *frame bounds*. In case  $\alpha = \beta$ , we call  $\{f_k\}_{k=1}^\infty$  an  *$\alpha$ -tight frame*, and if  $\alpha = \beta = 1$  it is a *Parseval frame* for  $\mathcal{W}$ .

To a Bessel sequence  $\mathcal{F} = \{f_k\}_{k=1}^\infty$  for  $\mathcal{W}$  we associate the *synthesis operator*

$$T_{\mathcal{F}} : \ell^2 \rightarrow \mathcal{H}, T_{\mathcal{F}} \{c_k\}_{k=1}^\infty = \sum_{k=1}^{\infty} c_k f_k,$$

the *analysis operator*

$$T_{\mathcal{F}}^* : \mathcal{H} \rightarrow \ell^2, T_{\mathcal{F}}^* f = \{\langle f, f_k \rangle\}_{k=1}^\infty,$$

and the *frame operator*

$$S_{\mathcal{F}} = T_{\mathcal{F}} T_{\mathcal{F}}^*.$$

A Bessel sequence  $\mathcal{F} = \{f_k\}_{k=1}^\infty$  for  $\mathcal{W}$  is a frame for  $\mathcal{W}$  if and only  $\mathcal{R}(T_{\mathcal{F}}) = \mathcal{W}$ , or equivalently,  $S_{\mathcal{F}}$  is invertible when restricted to  $\mathcal{W}$ . Furthermore,  $\mathcal{F}$  is an  $\alpha$ -tight frame for  $\mathcal{W}$  if and only if  $S_{\mathcal{F}} = \alpha P_{\mathcal{W}}$ . A *Riesz basis* for  $\mathcal{W}$  is a frame for  $\mathcal{W}$  which is also a basis. If  $\{f_k\}_{k=1}^\infty$  is a frame for  $\overline{\text{span}}\{f_k\}_{k=1}^\infty$  we say that  $\{f_k\}_{k=1}^\infty$  is a *frame sequence*.

The *optimal upper frame bound*, i.e., the infimum over all upper frame bounds is  $\|S_{\mathcal{F}}\| = \|T_{\mathcal{F}}\|^2$ , and the *optimal lower frame bound*, i.e., the supremum over all lower frame bounds is  $\|S_{\mathcal{F}}^{-1}\|^{-1} = \|T_{\mathcal{F}}^{\dagger}\|^{-2}$ .

**Definition 2.4.** Let  $\mathcal{W}$  be a closed subspace of  $\mathcal{H}$ . Let  $\mathcal{F} = \{f_k\}_{k=1}^{\infty}$  and  $\mathcal{G} = \{g_k\}_{k=1}^{\infty}$  be frames for  $\mathcal{W}$ . If

$$T_{\mathcal{G}}T_{\mathcal{F}}^* = \pi_{\mathcal{W}},$$

we say that  $\mathcal{G}$  is a *dual frame* of  $\mathcal{F}$  in  $\mathcal{W}$ .

The sequence  $\{S_{\mathcal{F}}^{\dagger}f_k\}_{k=1}^{\infty}$  is the *canonical dual frame* of  $\{f_k\}_{k=1}^{\infty}$  in  $\mathcal{W}$ .

For more details about frames we refer the reader to [3, 6, 21].

**2.3. Shift-invariant frame sequences.** For  $f \in L^1(\mathbb{R})$ , its Fourier transform, that we denote by  $\widehat{f}$  or  $\mathcal{F}f$ , is defined by

$$\widehat{f}(\gamma) = \int_{-\infty}^{\infty} f(x)e^{-i2\pi x\gamma} dx$$

for  $\gamma \in \mathbb{R}$ . The Fourier transform can be extended as a unitary operator in  $L^2(\mathbb{R})$ .

The discrete-time Fourier transform is given as the function in  $L^2(0, 1)$  defined by

$$\widehat{\mathbf{c}}(\gamma) = \sum_{k \in \mathbb{Z}} c_k e^{-i2\pi k\gamma}$$

for  $\mathbf{c} = \{c_k\}_{k \in \mathbb{Z}} \in \ell^2(\mathbb{Z})$ . Given  $k \in \mathbb{Z}$ , the *translation operator*  $T_k : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  is given by

$$T_k f(x) = f(x - k),$$

and the *modulation operator*  $E_k : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  by

$$E_k f(x) = e^{i2\pi kx} f(x).$$

Let  $\phi \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  is a Bessel sequence. Let  $f \in L^2(\mathbb{R})$ . Then  $f = \sum_{k \in \mathbb{Z}} c_k T_k\phi$  with  $\{c_k\}_{k \in \mathbb{Z}} \in \ell^2(\mathbb{Z})$  if and only if its Fourier transform is  $\widehat{f} = H\widehat{\phi}$  where  $H = \sum_{k \in \mathbb{Z}} c_k E_{-k} \in L^2(0, 1)$  [3, Lemma 9.2.2].

Let  $\mathcal{W} := \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$ . A space of this type is called *shift-invariant*. If  $\{T_k\phi\}_{k \in \mathbb{Z}}$  is a frame sequence, then

$$\mathcal{W} = \{\sum_{k \in \mathbb{Z}} c_k T_k\phi : \{c_k\}_{k \in \mathbb{Z}} \in \ell^2(\mathbb{Z})\}.$$

We will note

$$\Phi : [0, 1] \rightarrow \mathbb{R}, \quad \Phi(\gamma) = \sum_{k \in \mathbb{Z}} |\widehat{\phi}(\gamma + k)|^2,$$

The function  $\Phi$  belongs to  $L^1(0, 1)$ .

**Theorem 2.5.** *Let  $\phi \in L^2(\mathbb{R})$ . Then*

- (i)  $\{T_k\phi\}_{k \in \mathbb{Z}}$  is a Bessel sequence with bound  $\beta$  if and only if  $\Phi(\gamma) \leq \beta$ , a.e.  $\gamma \in [0, 1]$ .
- (ii)  $\{T_k\phi\}_{k \in \mathbb{Z}}$  is a frame sequence with bounds  $\alpha, \beta$  if and only if  $\alpha \leq \Phi(\gamma) \leq \beta$  a.e.  $\gamma \in \{\Phi \neq 0\}$ .

Let  $\phi, \widetilde{\phi} \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\widetilde{\phi}\}_{k \in \mathbb{Z}}$  are Bessel sequences with bounds  $\beta$  and  $\widetilde{\beta}$ , respectively. By Theorem 2.5,

$$(2.3) \quad \sum_{n \in \mathbb{Z}} |\widehat{\phi}(\gamma + n)\overline{\widehat{\widetilde{\phi}}(\gamma + n)}| \leq \sqrt{\beta\widetilde{\beta}} \quad \text{a.e. } \gamma \in [0, 1].$$

**Proposition 2.6.** [4] *Let  $\phi, \widetilde{\phi} \in L^2(\mathbb{R})$ , and assume that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\widetilde{\phi}\}_{k \in \mathbb{Z}}$  are frame sequences. Let  $\mathcal{W} := \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\mathcal{V} := \overline{\text{span}}\{T_k\widetilde{\phi}\}_{k \in \mathbb{Z}}$ . Then the following are equivalent:*

- (i)  $L^2(\mathbb{R}) = \mathcal{V} \oplus \mathcal{W}^{\perp}$ .

(ii)  $\{\Phi = 0\} = \{\tilde{\Phi} = 0\}$  and there exists a constant  $c > 0$  such that

$$\left| \sum_{k \in \mathbb{Z}} \widehat{\phi}(\gamma + k) \overline{\widehat{\phi}(\gamma + k)} \right| \geq c \text{ a.e. } \gamma \in \{\Phi \neq 0\}.$$

Let  $\mathcal{W}$  and  $\mathcal{V}$  be two closed subspaces of  $L^2(\mathbb{R})$  such that  $L^2(\mathbb{R}) = \mathcal{V} \oplus \mathcal{W}^\perp$ . In [4] it is shown that if  $\{T_k \phi\}_{k \in \mathbb{Z}}$  is a frame for  $\mathcal{W}$  and  $\{T_k \tilde{\phi}\}_{k \in \mathbb{Z}}$  is a frame for  $\mathcal{V}$  then

$$(2.4) \quad \cos(\mathcal{W}, \mathcal{V}) \geq \text{ess inf}_{w \in \{\Phi \neq 0\}} \frac{|\sum_{k \in \mathbb{Z}} \widehat{\phi}(w - k) \overline{\widehat{\phi}(w - k)}|}{\sqrt{\sum_{k \in \mathbb{Z}} |\widehat{\phi}(w - k)|^2} \sqrt{\sum_{k \in \mathbb{Z}} |\widehat{\phi}(w - k)|^2}}.$$

### 3. DEFINITION AND FUNDAMENTAL PROPERTIES

In this section we present the concept of approximate oblique duality and explore some of its properties. First we state the next result that can be deduced from [16, Lemma 2.1] and the definitions of orthogonal and oblique projections:

**Lemma 3.1.** *The following holds:*

- (i)  $(P_{\mathcal{W}})_{|\mathcal{V}} : \mathcal{V} \rightarrow \mathcal{W}$  and  $(P_{\mathcal{V}})_{|\mathcal{W}} : \mathcal{W} \rightarrow \mathcal{V}$  are isomorphisms,  $((P_{\mathcal{W}})_{|\mathcal{V}})^{-1} = (\pi_{\mathcal{V}\mathcal{W}^\perp})_{|\mathcal{W}}$  and  $((P_{\mathcal{V}})_{|\mathcal{W}})^{-1} = (\pi_{\mathcal{W}\mathcal{V}^\perp})_{|\mathcal{V}}$ .
- (ii)  $((P_{\mathcal{W}})_{|\mathcal{V}})^* = (P_{\mathcal{V}})_{|\mathcal{W}}$  and  $((\pi_{\mathcal{W}\mathcal{V}^\perp})_{|\mathcal{V}})^* = (\pi_{\mathcal{V}\mathcal{W}^\perp})_{|\mathcal{W}}$ .

We introduce the following definition of approximate oblique dual frames:

**Definition 3.2.** Let  $\mathcal{F} = \{f_k\}_{k=1}^\infty$  and  $\mathcal{G} = \{g_k\}_{k=1}^\infty$  be frames for  $\mathcal{W}$  and  $\mathcal{V}$ , respectively. Let  $\epsilon \geq 0$ . We say that  $\mathcal{F}$  and  $\mathcal{G}$  are  $\epsilon$ -approximate oblique dual frames if

$$(3.1) \quad \|\pi_{\mathcal{W}\mathcal{V}^\perp} - T_{\mathcal{F}} T_{\mathcal{G}}^*\| \leq \epsilon \text{ or } \|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}} T_{\mathcal{F}}^*\| \leq \epsilon.$$

In this case we also say that  $\{g_k\}_{k=1}^\infty$  is an  $\epsilon$ -approximate oblique dual frame of  $\{f_k\}_{k=1}^\infty$  in  $\mathcal{V}$  and that  $\{f_k\}_{k=1}^\infty$  is an  $\epsilon$ -approximate oblique dual frame of  $\{g_k\}_{k=1}^\infty$  in  $\mathcal{W}$ . If  $\mathcal{V} = \mathcal{W}$ , then  $\mathcal{F}$  and  $\mathcal{G}$  will be called  $\epsilon$ -approximate dual frames in  $\mathcal{W}$ .

Note that the conditions of the definition can be written as

$$(3.2) \quad \left\| \pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k=1}^{\infty} \langle f, g_k \rangle f_k \right\| \leq \epsilon \|f\| \text{ for all } f \in \mathcal{H}$$

and

$$(3.3) \quad \left\| \pi_{\mathcal{V}\mathcal{W}^\perp} f - \sum_{k=1}^{\infty} \langle f, f_k \rangle g_k \right\| \leq \epsilon \|f\| \text{ for all } f \in \mathcal{H},$$

respectively.

*Remark 3.3.* (i) If  $\epsilon = 0$ ,  $\mathcal{F}$  and  $\mathcal{G}$  are oblique dual frames [11, 4]. If  $\mathcal{V} = \mathcal{W} = \mathcal{H}$  and  $\epsilon < 1$ , then  $\mathcal{F}$  and  $\mathcal{G}$  are approximate dual frames as defined in [5].

(ii) If  $\epsilon < 1$ , by Neumann's Theorem  $(T_{\mathcal{F}} T_{\mathcal{G}}^*)_{|\mathcal{W}}$  is an invertible operator. Note that  $(T_{\mathcal{F}} T_{\mathcal{G}}^*)_{|\mathcal{V}} = (T_{\mathcal{F}} T_{\mathcal{G}}^*)_{|\mathcal{W}} (P_{\mathcal{W}})_{|\mathcal{V}}$ . So, by (i) of Lemma 3.1,  $(T_{\mathcal{F}} T_{\mathcal{G}}^*)_{|\mathcal{V}}$  is an invertible operator from  $\mathcal{V}$  to  $\mathcal{W}$ .

(iii) If  $\epsilon < 1$ , it is sufficient that  $\{f_k\}_{k=1}^\infty$  and  $\{g_k\}_{k=1}^\infty$  are Bessel sequences, since  $\mathcal{R}(T_{\mathcal{F}}) = \mathcal{W}$  by (3.3). Hence  $\{f_k\}_{k=1}^\infty$  is a frame for  $\mathcal{W}$ . Analogously  $\{g_k\}_{k=1}^\infty$  is a frame for  $\mathcal{V}$ .

(iv) Assume that  $\epsilon < 1$ . If  $f \in \mathcal{W}$ , by (3.3),

$$f = (T_{\mathcal{F}}T_{\mathcal{G}|\mathcal{W}}^*)^{-1}T_{\mathcal{F}}T_{\mathcal{G}}^*f = \sum \langle f, g_k \rangle (T_{\mathcal{F}}T_{\mathcal{G}|\mathcal{W}}^*)^{-1}f_k.$$

Similarly, for  $f \in \mathcal{V}$  we have

$$f = (T_{\mathcal{G}}T_{\mathcal{F}|\mathcal{V}}^*)^{-1}T_{\mathcal{G}}T_{\mathcal{F}}^*f = \sum \langle f, f_k \rangle (T_{\mathcal{G}}T_{\mathcal{F}|\mathcal{V}}^*)^{-1}g_k.$$

Hence  $\{(T_{\mathcal{F}}T_{\mathcal{G}|\mathcal{W}}^*)^{-1}f_k\}_{k=1}^{\infty}$  and  $\{g_k\}_{k=1}^{\infty}$  ( $\{f_k\}_{k=1}^{\infty}$  and  $\{(T_{\mathcal{G}}T_{\mathcal{F}|\mathcal{V}}^*)^{-1}g_k\}_{k=1}^{\infty}$ ) are oblique dual frames by [4, lema 3.1(i)].

The next proposition tells us that approximate oblique duality is preserved under the action of unitary operators.

**Proposition 3.4.** *Let  $\mathcal{F} = \{f_k\}_{k=1}^{\infty} \subset \mathcal{W}$  and  $\mathcal{G} = \{g_k\}_{k=1}^{\infty} \subset \mathcal{V}$  be  $\epsilon$ -approximate oblique dual frames and  $U : \mathcal{H} \rightarrow \mathcal{H}$  a unitary operator. Then  $U\mathcal{F} = \{Uf_k\}_{k=1}^{\infty}$  and  $U\mathcal{G} = \{Ug_k\}_{k=1}^{\infty}$  are  $\epsilon$ -approximate oblique dual frames.*

*Proof.* By [3, Corollary 5.3.4], the sequences  $U\mathcal{F}$  and  $U\mathcal{G}$  are frames for  $U\mathcal{W}$  and  $U\mathcal{V}$ , respectively. We have  $T_{U\mathcal{F}} = UT_{\mathcal{F}}$  and  $T_{U\mathcal{G}} = UT_{\mathcal{G}}$ .

To see that  $\mathcal{H}$  can be decomposed as  $\mathcal{H} = U\mathcal{W} \oplus (U\mathcal{V})^{\perp}$ , observe first that  $\mathcal{H} = \mathcal{W} \oplus \mathcal{V}^{\perp}$  and  $I = UU^*$ . So

$$(3.4) \quad I = U\pi_{\mathcal{W}\mathcal{V}^{\perp}}U^* + U\pi_{\mathcal{V}^{\perp}\mathcal{W}}U^*.$$

This shows that  $\mathcal{H} = U\mathcal{W} + (U\mathcal{V}^{\perp})$ .

Let  $f \in U\mathcal{W} \cap (U\mathcal{V}^{\perp})$ . Then there exist  $g \in \mathcal{W}$  and  $h \in \mathcal{V}^{\perp}$  such that  $f = Ug = Uh$ . Since  $U$  is injective  $g = h$ . But  $\mathcal{W} \cap \mathcal{V}^{\perp} = \{0\}$ . Hence  $g = h = 0$  and so  $f = 0$ . This shows that  $U\mathcal{W} \cap (U\mathcal{V}^{\perp}) = \{0\}$  and so  $\mathcal{H} = U\mathcal{W} \oplus (U\mathcal{V}^{\perp})$ .

From the definition of orthogonal complement and unitary operator it follows that  $(U\mathcal{V})^{\perp} = U\mathcal{V}^{\perp}$ .

Therefore,

$$(3.5) \quad \mathcal{H} = U\mathcal{W} \oplus (U\mathcal{V})^{\perp}.$$

From (3.5) and (3.4), we have

$$(3.6) \quad \pi_{U\mathcal{W}(U\mathcal{V})^{\perp}} = U\pi_{\mathcal{W}\mathcal{V}^{\perp}}U^*.$$

But then

$$\begin{aligned} \|\pi_{U\mathcal{W}(U\mathcal{V})^{\perp}} - T_{U\mathcal{F}}T_{U\mathcal{G}}^*\| &= \|U\pi_{\mathcal{W}\mathcal{V}^{\perp}}U^* - UT_{\mathcal{F}}T_{\mathcal{G}}^*U^*\| = \|U(\pi_{\mathcal{W}\mathcal{V}^{\perp}} - T_{\mathcal{F}}T_{\mathcal{G}}^*)U^*\| \\ &= \|\pi_{\mathcal{W}\mathcal{V}^{\perp}} - T_{\mathcal{F}}T_{\mathcal{G}}^*\| \leq \epsilon, \end{aligned}$$

and so  $U\mathcal{F}$  and  $U\mathcal{G}$  are  $\epsilon$ -approximate oblique dual frames.  $\square$

Now we briefly discuss the interpretation of the approximate oblique duality in the context of signal processing theory. Let  $\mathcal{F}$  be a frame for  $\mathcal{W}$ . Assume that the samples  $T_{\mathcal{F}}^*f = (\langle f, f_k \rangle)_{k=1}^{\infty}$  of an unknown signal  $f \in \mathcal{H}$  are given. Our goal is the reconstruction of  $f$  from these samples using a frame  $\mathcal{G}$  for  $\mathcal{V}$  in such a way that the reconstruction  $f_r \in \mathcal{V}$  is a good approximation of  $f$ . This means on one hand that there doesn't exist two different signals in  $\mathcal{V}$  with the same samples. On the other, that the samples of the reconstruction  $f_r$  and those of the original signal  $f$  are close. Specifically the following two conditions are required. Let  $\epsilon > 0$ ,

- (i) *Uniqueness of the reconstruction:* If  $f, g \in \mathcal{V}$  and  $T_{\mathcal{F}}^*f = T_{\mathcal{F}}^*g$ , then  $f = g$ .
- (ii)  *$\epsilon$ -consistent reconstruction:* For every  $f \in \mathcal{H}$ ,  $\|T_{\mathcal{F}}^*f_r - T_{\mathcal{F}}^*f\| \leq \epsilon \|f\|$ .

If (ii) holds, we say that  $f_r$  is an  $\epsilon$ -consistent reconstruction of  $f$  in  $\mathcal{V}$ . Note that a 0-consistent reconstruction is a consistent reconstruction as defined in [11, 12, 14]. The condition (i) is equivalent to  $\mathcal{V} \cap \mathcal{W}^\perp = \{0\}$ .

Using an oblique dual frame of  $\mathcal{F}$  in  $\mathcal{V}$ , we can obtain the following bound for  $\|f_r - \pi_{\mathcal{V}\mathcal{W}^\perp} f\|$ :

**Proposition 3.5.** *Let  $\mathcal{F} = \{f_k\}_{k=1}^\infty$  be a frame for  $\mathcal{W}$  and  $\mathcal{G}$  be an oblique dual frame of  $\mathcal{F}$  in  $\mathcal{V}$ . Let  $\epsilon \geq 0$ . If  $f \in \mathcal{H}$  and  $f_r$  is an  $\epsilon$ -consistent reconstruction of  $f$  in  $\mathcal{V}$ , then*

$$\|f_r - \pi_{\mathcal{V}\mathcal{W}^\perp} f\| \leq \epsilon \|T_{\mathcal{G}}\| \|f\|.$$

*Proof.* Using that  $T_{\mathcal{G}}T_{\mathcal{F}}^* = \pi_{\mathcal{V}\mathcal{W}^\perp}$  and  $f_r \in \mathcal{V}$ , we have

$$\|f_r - \pi_{\mathcal{V}\mathcal{W}^\perp} f\| = \|T_{\mathcal{G}}T_{\mathcal{F}}^*f_r - T_{\mathcal{G}}T_{\mathcal{F}}^*\pi_{\mathcal{V}\mathcal{W}^\perp}f\| \leq \|T_{\mathcal{G}}\| \|T_{\mathcal{F}}^*f_r - T_{\mathcal{F}}^*f\| \leq \epsilon \|T_{\mathcal{G}}\| \|f\|. \quad \square$$

The following result relates an  $\epsilon$ -consistent reconstruction  $f_r$  to the oblique projection  $\pi_{\mathcal{V}\mathcal{W}^\perp} f$  of  $f$ .

**Theorem 3.6.** *Let  $\mathcal{F}$  be a frame for  $\mathcal{W}$ ,  $f \in \mathcal{H}$  and  $\epsilon \geq 0$ . The following hold:*

- (i) *If  $f_r$  is an  $\epsilon$ -consistent reconstruction of  $f$  in  $\mathcal{V}$ , then  $\|f_r - \pi_{\mathcal{V}\mathcal{W}^\perp} f\| \leq \epsilon \|\pi_{\mathcal{V}\mathcal{W}^\perp}\| \|T_{\mathcal{F}}^\dagger\| \|f\|$ .*
- (ii) *If  $\|f_r - \pi_{\mathcal{V}\mathcal{W}^\perp} f\| \leq \frac{\epsilon}{\|T_{\mathcal{F}}^\dagger\|} \|f\|$ , then  $f_r$  is an  $\epsilon$ -consistent reconstruction of  $f$  in  $\mathcal{V}$ .*

*Proof.* If  $\mathcal{G}$  is the canonical oblique dual frame of  $\mathcal{F}$  then  $T_{\mathcal{G}} = \pi_{\mathcal{V}\mathcal{W}^\perp} S_{\mathcal{F}}^\dagger T_{\mathcal{F}}$ . We also have,  $T_{\mathcal{F}}^\dagger = T_{\mathcal{F}}^* S_{\mathcal{F}}^\dagger$ . So (i) follows from Proposition 3.5.

To prove (ii), assume that  $\|f_r - \pi_{\mathcal{V}\mathcal{W}^\perp} f\| \leq \frac{\epsilon}{\|T_{\mathcal{F}}^\dagger\|} \|f\|$ . Then we have

$$\|T_{\mathcal{F}}^* f_r - T_{\mathcal{F}}^* f\| = \|T_{\mathcal{F}}^* f_r - T_{\mathcal{F}}^* \pi_{\mathcal{V}\mathcal{W}^\perp} f\| \leq \epsilon \|f\|. \quad \square$$

The previous theorem allows to link  $\epsilon$ -consistent reconstruction with approximate oblique duality via the next corollary.

**Corollary 3.7.** *Let  $\mathcal{F}$  be a frame for  $\mathcal{W}$ ,  $\mathcal{G}$  a frame for  $\mathcal{V}$  and  $\epsilon \geq 0$ . Then*

- (i) *If  $f_r = T_{\mathcal{G}}T_{\mathcal{F}}^*f$  is an  $\frac{\epsilon}{\|\pi_{\mathcal{V}\mathcal{W}^\perp}\| \|T_{\mathcal{F}}^\dagger\|}$ -consistent reconstruction of  $f$  in  $\mathcal{V}$ , for each  $f \in \mathbb{H}$ , then  $\mathcal{F}$  and  $\mathcal{G}$  are  $\epsilon$ -approximate oblique dual frames.*
- (ii) *If  $\mathcal{F}$  and  $\mathcal{G}$  are  $\epsilon$ -approximate oblique dual frames, then  $f_r = T_{\mathcal{G}}T_{\mathcal{F}}^*f$  is an  $\epsilon \|T_{\mathcal{F}}^\dagger\|$ -consistent reconstruction of  $f$  in  $\mathcal{V}$ , for each  $f \in \mathbb{H}$ .*

#### 4. CHARACTERIZATION OF APPROXIMATE OBLIQUE DUAL FRAMES

The following lemma gives equivalent conditions for approximate oblique duality.

**Lemma 4.1.** *Let  $\mathcal{F} = \{f_k\}_{k=1}^\infty$ ,  $\mathcal{G} = \{g_k\}_{k=1}^\infty$  be Bessel sequences in  $\mathcal{H}$  such that  $\overline{\text{span}}\{f_k\}_{k=1}^\infty = \mathcal{W}$  and  $\overline{\text{span}}\{g_k\}_{k=1}^\infty = \mathcal{V}$ . Let  $\epsilon \geq 0$ . Then the following assertions are equivalent:*

- (i)  *$\{f_k\}_{k=1}^\infty$  and  $\{g_k\}_{k=1}^\infty$  are  $\epsilon$ -approximate oblique dual frames.*
- (ii)  *$\|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k=1}^\infty \langle f, g_k \rangle f_k\| \leq \epsilon \|f\|$  for all  $f \in \mathcal{H}$ .*
- (iii)  *$\|\pi_{\mathcal{V}\mathcal{W}^\perp} f - \sum_{k=1}^\infty \langle f, f_k \rangle g_k\| \leq \epsilon \|f\|$  for all  $f \in \mathcal{H}$ .*
- (iv)  *$|\langle \pi_{\mathcal{W}\mathcal{V}^\perp} f, g \rangle - \sum_{k=1}^\infty \langle f, g_k \rangle \langle f_k, g \rangle| \leq \epsilon \|f\| \|g\|$  for all  $f, g \in \mathcal{H}$ .*
- (v)  *$|\langle \pi_{\mathcal{V}\mathcal{W}^\perp} f, g \rangle - \sum_{k=1}^\infty \langle f, f_k \rangle \langle g_k, g \rangle| \leq \epsilon \|f\| \|g\|$  for all  $f, g \in \mathcal{H}$ .*

*Proof.* Recall that (ii) and (iii) are the equivalent conditions (3.2) and (3.3) of Definition 3.2.

Let  $f, g \in \mathcal{H}$ . Assume that (ii) holds. Using that

$$|\langle \pi_{\mathcal{W}\mathcal{V}^\perp} f, g \rangle - \sum_{k=1}^{\infty} \langle f, g_k \rangle \langle f_k, g \rangle| = |\langle \pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k=1}^{\infty} \langle f, g_k \rangle f_k, g \rangle|,$$

and Schwartz's inequality, we obtain (iv).

Assume now that (iv) holds. We have

$$\left\| \pi_{\mathcal{W}\mathcal{V}^\perp} g - \sum_{k=1}^{\infty} \langle g, g_k \rangle f_k \right\| = \sup_{\|f\|=1} |\langle \pi_{\mathcal{W}\mathcal{V}^\perp} g - \sum_{k=1}^{\infty} \langle g, g_k \rangle f_k, f \rangle| \leq \epsilon \|g\|.$$

So (ii) holds.

The proof of (iii)  $\Leftrightarrow$  (v) is similar to (ii)  $\Leftrightarrow$  (iv).  $\square$

Condition (ii) of Lemma 4.1 implies

$$(4.1) \quad \left\| f - \sum_{k=1}^{\infty} \langle f, g_k \rangle f_k \right\| \leq \epsilon \|f\| \quad \text{for } f \in \mathcal{W},$$

whereas (iii) implies  $\|g - \sum_{k=1}^{\infty} \langle g, f_k \rangle g_k\| \leq \epsilon \|g\|$  for  $g \in \mathcal{V}$ . We conclude that the elements of each subspace can be reconstructed approximately, with a uniform relative error.

Let  $\{f_k\}_{k=1}^{\infty}$  be a Bessel sequence in  $\mathcal{W}$  and  $\{g_k\}_{k=1}^{\infty}$  be a Bessel sequence in  $\mathcal{V}$ . Observe that for  $f \in \mathcal{H}$ ,

$$\begin{aligned} \left\| \pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k=1}^{\infty} \langle f, g_k \rangle f_k \right\| &= \left\| \pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k=1}^{\infty} \langle f, \pi_{\mathcal{V}\mathcal{W}^\perp} g_k \rangle f_k \right\| \\ &= \left\| \pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k=1}^{\infty} \langle \pi_{\mathcal{W}\mathcal{V}^\perp} f, g_k \rangle f_k \right\| \end{aligned}$$

So, we have the following result:

**Proposition 4.2.** *Let  $\{f_k\}_{k=1}^{\infty}$  be a frame for  $\mathcal{W}$  and  $\{g_k\}_{k=1}^{\infty}$  be a frame for  $\mathcal{V}$  such that (4.1) is satisfied. Then  $\{f_k\}_{k=1}^{\infty}$  and  $\{g_k\}_{k=1}^{\infty}$  are  $\epsilon \|\pi_{\mathcal{W}\mathcal{V}^\perp}\|$ -approximate oblique dual frames.*

Similarly we have:

**Proposition 4.3.** *Let  $\{f_k\}_{k=1}^{\infty}$  be a Bessel sequence in  $\mathcal{W}$  and  $\{g_k\}_{k=1}^{\infty}$  be a Bessel sequence in  $\mathcal{H}$  such that (4.1) holds. Let  $\mathcal{U}$  be a closed subspace such that  $\mathcal{H} = \mathcal{W} \oplus \mathcal{U}^\perp$ . If  $\epsilon \|\pi_{\mathcal{W}\mathcal{U}^\perp}\| < 1$ , then  $\{f_k\}_{k=1}^{\infty}$  is a frame for  $\mathcal{W}$ ,  $\{\pi_{\mathcal{U}\mathcal{W}^\perp} g_k\}_{k=1}^{\infty}$  is a frame for  $\mathcal{U}$  and  $\{f_k\}_{k=1}^{\infty}$  and  $\{\pi_{\mathcal{U}\mathcal{W}^\perp} g_k\}_{k=1}^{\infty}$  are  $\epsilon \|\pi_{\mathcal{W}\mathcal{U}^\perp}\|$ -approximate oblique dual frames.*

In [13, Lemma B.1] a characterization of oblique dual frames is given by oblique inverses. We have a similar characterization for approximate oblique dual frames using *approximate oblique left inverses*.

**Definition 4.4.** Let  $\mathcal{F} = \{f_k\}_{k=1}^{\infty}$  be a frame for  $\mathcal{W}$ . We say that a linear bounded operator  $A : \ell^2(\mathbb{N}) \rightarrow \mathcal{V}$  is an  $\epsilon$ -approximate oblique left inverse of  $T_{\mathcal{F}}^*$  in  $\mathcal{V}$  along  $\mathcal{W}^\perp$  if

$$(4.2) \quad \|\pi_{\mathcal{V}\mathcal{W}^\perp} - AT_{\mathcal{F}}^*\| \leq \epsilon.$$

We now give the following characterization.



**Lemma 4.5.** *Let  $\mathcal{F} = \{f_k\}_{k=1}^\infty$  be a frame for  $\mathcal{W}$ . Let  $\{\delta_k\}_{k=1}^\infty$  be the canonical basis of  $\ell^2(\mathbb{N})$ . Let  $\epsilon < 1$ . The  $\epsilon$ -approximate oblique dual frames of  $\mathcal{F}$  in  $\mathcal{V}$  are the sequences  $\{g_k\}_{k=1}^\infty = \{A\delta_k\}_{k=1}^\infty$ , where  $A : \ell^2(\mathbb{N}) \rightarrow \mathcal{V}$  is an  $\epsilon$ -approximate oblique left inverse of  $T_{\mathcal{F}}^*$  in  $\mathcal{V}$  along  $\mathcal{W}^\perp$ .*

*Proof.* Clearly, if  $\mathcal{G} = \{g_k\}_{k=1}^\infty$  is an  $\epsilon$ -approximate oblique dual frame of  $\{f_k\}_{k=1}^\infty$  in  $\mathcal{V}$ , then  $T_{\mathcal{G}}$  is an  $\epsilon$ -approximate oblique left inverse of  $T_{\mathcal{F}}^*$  in  $\mathcal{V}$  along  $\mathcal{W}^\perp$  and  $\{g_k\}_{k=1}^\infty = \{T_{\mathcal{G}}\delta_k\}_{k=1}^\infty$ .

Let now  $A : \ell^2(\mathbb{N}) \rightarrow \mathcal{V}$  be an  $\epsilon$ -approximate oblique left inverse. Let  $\mathcal{G} = \{g_k\}_{k=1}^\infty = \{A\delta_k\}_{k=1}^\infty$ . By [3, Theorem 3.2.3],  $\mathcal{G}$  is a Bessel sequence in  $\mathcal{V}$  with  $T_{\mathcal{G}} = A$ . We have  $\|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*\| \leq \epsilon$ . By Remark 3.3,  $\mathcal{G}$  is a frame for  $\mathcal{V}$ . Thus,  $\mathcal{G}$  is an  $\epsilon$ -approximate oblique dual frame of  $\{f_k\}_{k=1}^\infty$  in  $\mathcal{V}$ .  $\square$

Using Lemma 4.5 and an expression of the approximate oblique left inverses we obtain a description of the approximate oblique dual frames:

**Theorem 4.6.** *Let  $\mathcal{F} = \{f_k\}_{k=1}^\infty$  be a frame for  $\mathcal{W}$ . Let  $\epsilon < 1$ . Then any  $\epsilon$ -approximate oblique dual frame  $\mathcal{G} = \{g_k\}_{k=1}^\infty \subset \mathcal{V}$  of  $\mathcal{F}$  in  $\mathcal{V}$  is of the form*

$$\{g_k\}_{k=1}^\infty = \{\pi_{\mathcal{V}\mathcal{W}^\perp}S_{\mathcal{F}}^\dagger f_k + h_k - \sum_{j=1}^\infty \langle S_{\mathcal{F}}^\dagger f_k, f_j \rangle h_j + r_k\}_{k=1}^\infty$$

where  $\{h_k\}_{k=1}^\infty$  and  $\{r_k\}_{k=1}^\infty$  are Bessel sequences in  $\mathcal{V}$  and  $\|\sum \langle f, f_k \rangle r_k\| \leq \epsilon \|f\|$  for all  $f \in \mathcal{H}$ .

*Proof.* We note that  $A : \ell^2(\mathbb{N}) \rightarrow \mathcal{V}$  is an  $\epsilon$ -approximate oblique left inverse of  $T_{\mathcal{F}}^*$  in  $\mathcal{V}$  if and only if  $A = L + R$  where  $L : \ell^2(\mathbb{N}) \rightarrow \mathcal{V}$  is an oblique left inverse of  $T_{\mathcal{F}}^*$  in  $\mathcal{V}$  and  $R : \ell^2(\mathbb{N}) \rightarrow \mathcal{V}$  is a bounded operator such that  $\|RT_{\mathcal{F}}^*\| \leq \epsilon$ . So, from Lemma 4.5 and [4, Lemma B.2], any  $\epsilon$ -approximate oblique dual frame  $\mathcal{G} = \{g_k\}_{k=1}^\infty \subset \mathcal{V}$  of the frame  $\mathcal{F}$  in  $\mathcal{V}$  is of the form

$$\{g_k\}_{k=1}^\infty = \left\{ \left( \pi_{\mathcal{V}\mathcal{W}^\perp} S_{\mathcal{F}}^\dagger T_{\mathcal{F}} + W(I - T_{\mathcal{F}}^* S_{\mathcal{F}}^\dagger T_{\mathcal{F}}) + R \right) \delta_k \right\}_{k=1}^\infty$$

where  $W : \ell^2(\mathbb{N}) \rightarrow \mathcal{V}$  and  $R : \ell^2(\mathbb{N}) \rightarrow \mathcal{V}$  are bounded operators such that  $\|RT_{\mathcal{F}}^*\| \leq \epsilon$ . Setting  $h_k = W\delta_k$  and  $r_k = R\delta_k$  for each  $k$ , we obtain the conclusion.  $\square$

## 5. IMPROVING APPROXIMATION

Given a frame  $\mathcal{F} = \{f_k\}_{k=1}^\infty$  in  $\mathcal{W}$  and an approximate oblique dual frame  $\mathcal{G} = \{g_k\}_{k=1}^\infty \subset \mathcal{V}$  of it, the following proposition says that it is possible to construct from  $\mathcal{G}$  other approximate oblique dual frames for  $\mathcal{F}$  in  $\mathcal{V}$  that provide a reconstruction as “good” as needed. This is analogous to a result for approximate dual frames [5, Proposition 3.2].

**Proposition 5.1.** *Let  $\mathcal{F} = \{f_k\}_{k=1}^\infty \subset \mathcal{W}$  and  $\mathcal{G} = \{g_k\}_{k=1}^\infty \subset \mathcal{V}$  be  $\epsilon$ -approximate oblique dual frames with  $0 \leq \epsilon < 1$ . For a fixed  $N \in \mathbb{N}$ , let*

$$(5.1) \quad \tilde{g}_k^{(N)} = \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n g_k$$

and  $\tilde{\mathcal{G}}_N = \{\tilde{g}_k^{(N)}\}_{k=1}^\infty$ . Then  $\tilde{\mathcal{G}}_N$  is an  $\epsilon^{N+1}$ -approximate oblique dual frame of  $\mathcal{F}$  in  $\mathcal{V}$  and

$$(5.2) \quad \left\| \pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}_N}T_{\mathcal{F}}^* \right\| \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

*Proof.* Let  $\tilde{g}_k^{(N)} = \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n g_k$  for  $N \in \mathbb{N}$  and  $\tilde{\mathcal{G}}_N = \{\tilde{g}_k^{(N)}\}_{k=1}^\infty$ .

Since  $\{g_k\}_{k=1}^\infty$  is a Bessel sequence and  $\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*$  is a bounded operator, it follows that  $\{\tilde{g}_k^{(N)}\}_{k=1}^\infty$  is a Bessel sequence. Hence, by [3, Theorem 3.2.3] its synthesis operator  $T_{\tilde{\mathcal{G}}_N}$  is well defined. Let  $f \in \mathcal{H}$ , then

$$\begin{aligned}
T_{\tilde{\mathcal{G}}_N}T_{\mathcal{F}}^*f &= \sum_{k=1}^\infty \langle f, f_k \rangle \tilde{g}_k^{(N)} = \sum_{k=1}^\infty \langle f, f_k \rangle \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n g_k \\
&= \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n T_{\mathcal{G}}T_{\mathcal{F}}^*f = \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n T_{\mathcal{G}}T_{\mathcal{F}}^* \pi_{\mathcal{V}\mathcal{W}^\perp} f \\
&= \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n (\pi_{\mathcal{V}\mathcal{W}^\perp} - (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)) \pi_{\mathcal{V}\mathcal{W}^\perp} f \\
&= \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n \pi_{\mathcal{V}\mathcal{W}^\perp} f - \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^{n+1} \pi_{\mathcal{V}\mathcal{W}^\perp} f \\
&= \pi_{\mathcal{V}\mathcal{W}^\perp} f - (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^{N+1} \pi_{\mathcal{V}\mathcal{W}^\perp} f
\end{aligned}$$

Now, using that  $\mathcal{W}^\perp \subseteq \mathcal{N}(\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)$ ,

$$\pi_{\mathcal{V}\mathcal{W}^\perp} f - T_{\tilde{\mathcal{G}}_N}T_{\mathcal{F}}^*f = (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^{N+1} \pi_{\mathcal{V}\mathcal{W}^\perp} f = (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^{N+1} f$$

Therefore,  $\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}_N}T_{\mathcal{F}}^* = (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^{N+1}$  and

$$(5.3) \quad \left\| \pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}_N}T_{\mathcal{F}}^* \right\| \leq \left\| \pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^* \right\|^{N+1} \leq \epsilon^{N+1} \rightarrow 0$$

as  $N \rightarrow \infty$ . □

*Remark 5.2.* Note that in the previous proof we obtain

$$\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}_N}T_{\mathcal{F}}^* = (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^{N+1}.$$

Considering

$$(5.4) \quad L_N = \sum_{n=0}^N (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n,$$

the operator  $T_{\tilde{\mathcal{G}}_N}$  can be expressed as  $T_{\tilde{\mathcal{G}}_N} = L_N T_{\mathcal{G}}$ .

Setting  $\tilde{f}_k^{(N)} = \sum_{n=0}^N (\pi_{\mathcal{W}\mathcal{V}^\perp} - T_{\mathcal{F}}T_{\mathcal{G}}^*)^n f_k$  and  $\tilde{\mathcal{F}}_N = \{\tilde{f}_k^{(N)}\}_{k=1}^\infty$ , we have  $T_{\tilde{\mathcal{F}}_N} = L_N^* T_{\mathcal{F}}$ ,  $\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\tilde{\mathcal{F}}_N}^* = (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^{N+1}$  and

$$(5.5) \quad \left\| \pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\tilde{\mathcal{F}}_N}^* \right\| \leq \left\| \pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^* \right\|^{N+1} \leq \epsilon^{N+1} \rightarrow 0.$$

From (5.3) and (5.5), we observe that we can improve the approximation given by the approximate oblique dual frames  $\mathcal{F}$  and  $\mathcal{G}$ , modifying either of them.

The following proposition shows that the frames  $\tilde{\mathcal{F}}_N$  and  $\tilde{\mathcal{G}}_N$  defined previously are related with oblique dual frames of  $\mathcal{F}$  and  $\mathcal{G}$ .

**Proposition 5.3.** *Let  $\mathcal{F} = \{f_k\}_{k=1}^\infty \subset \mathcal{W}$  and  $\mathcal{G} = \{g_k\}_{k=1}^\infty \subset \mathcal{V}$  be  $\epsilon$ -approximate oblique dual frames with  $0 \leq \epsilon < 1$ . Then  $I - (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)$  is invertible and*

$$\begin{aligned}
(5.6) \quad L &:= (I - (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*))^{-1} \\
&= I + \sum_{n=1}^\infty (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n.
\end{aligned}$$

Let  $\tilde{\mathcal{G}} = \{\tilde{g}_k\}_{k=1}^\infty$  with  $\tilde{g}_k = LT_{\mathcal{G}}\delta_k$  and  $\tilde{\mathcal{F}} = \{\tilde{f}_k\}_{k=1}^\infty$  with  $\tilde{f}_k = L^*T_{\mathcal{F}}\delta_k$ . Then:

- (i)  $\tilde{\mathcal{G}}$  is an oblique dual frame of  $\mathcal{F}$  in  $\mathcal{V}$ .
- (ii)  $\tilde{\mathcal{F}}$  is an oblique dual frame of  $\mathcal{G}$  in  $\mathcal{W}$ .
- (iii)  $\tilde{\mathcal{G}}$  is an  $\frac{\epsilon}{1-\epsilon}$ -approximate oblique dual frame of  $\tilde{\mathcal{F}}$  in  $\mathcal{V}$ .

*Proof.* The assertions about  $I - (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*)$  follow from Neumann's Theorem.

If  $L_N$  is as in (5.4), then  $L_N \rightarrow L$  as  $N \rightarrow \infty$ . So,  $T_{\tilde{\mathcal{G}}_N} = L_N T_{\mathcal{G}} \rightarrow T_{\tilde{\mathcal{G}}} = LT_{\mathcal{G}}$  as  $N \rightarrow \infty$ . Using (5.3), we have

$$\begin{aligned} \|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*\| &= \|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}_N}T_{\tilde{\mathcal{F}}}^*\| + \|T_{\tilde{\mathcal{G}}_N}T_{\tilde{\mathcal{F}}}^* - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*\| \\ &\leq \epsilon^{N+1} + \|T_{\tilde{\mathcal{G}}_N} - T_{\tilde{\mathcal{G}}}\| \|T_{\tilde{\mathcal{F}}}\| \end{aligned}$$

Since the right hand side tends to 0 as  $N \rightarrow \infty$ ,  $T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^* = \pi_{\mathcal{V}\mathcal{W}^\perp}$ . This shows that  $\tilde{\mathcal{G}}$  is an oblique dual frame of  $\mathcal{F}$  in  $\mathcal{V}$ . Similarly, it can be proved that  $\tilde{\mathcal{F}}$  is an oblique dual frame of  $\mathcal{G}$  in  $\mathcal{W}$ . Since  $\mathcal{R}(I - L) \subseteq \mathcal{V}$ , by (i),

$$\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^* = \pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*L = \pi_{\mathcal{V}\mathcal{W}^\perp}(I - L) = \sum_{n=1}^\infty (\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*)^n.$$

Then  $\|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*\| \leq \frac{\epsilon}{1-\epsilon}$ .  $\square$

*Remark 5.4.* We note that  $\tilde{\mathcal{G}}$  is not an oblique dual frame of  $\tilde{\mathcal{F}}$  in  $\mathcal{V}$ . In effect, assume that  $\tilde{\mathcal{G}}$  is an oblique dual frame of  $\tilde{\mathcal{F}}$  in  $\mathcal{V}$ . Then, from the equalities above  $L = I$ , and so  $\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^* = 2I$ . But this implies  $\|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*\| = 2$ , which contradicts  $\|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*\| \leq \epsilon < 1$ .

*Remark 5.5.* It is natural to ask if we can have an approximation as close as wanted to  $\pi_{\mathcal{V}\mathcal{W}^\perp}(f)$  using simultaneously the frames  $\tilde{\mathcal{F}}_N$  and  $\tilde{\mathcal{G}}_N$ . The answer is no. More precisely, since

$$\|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*\| \leq \|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}_N}T_{\tilde{\mathcal{F}}_N}^*\| + \|T_{\tilde{\mathcal{G}}_N}T_{\tilde{\mathcal{F}}_N}^* - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*\|$$

and

$$\lim_{N \rightarrow \infty} \|T_{\tilde{\mathcal{G}}_N}T_{\tilde{\mathcal{F}}_N}^* - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*\| = 0,$$

we obtain, using Remark 5.4,

$$\liminf_{N \rightarrow \infty} \|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}_N}T_{\tilde{\mathcal{F}}_N}^*\| \geq \|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{F}}}^*\| > 0.$$

## 6. APPROXIMATE OBLIQUE DUAL FRAMES BY PERTURBATION

The following propositions say that if two frames are ‘‘close’’, each oblique dual frame of one of them is an approximate oblique dual frame of the other. They are extensions of [5, Proposition 4.1 and Proposition 4.3], which are results about approximate dual frames.

**Proposition 6.1.** *Let  $0 \leq \epsilon < 1$ . Let  $\tilde{\mathcal{G}} = \{\tilde{g}_k\}_{k=1}^\infty$  be a frame for  $\mathcal{V}$ ,  $\mathcal{G} = \{g_k\}_{k=1}^\infty \subset \mathcal{V}$  a sequence such that  $\sum_{k=1}^\infty |\langle f, g_k - \tilde{g}_k \rangle|^2 \leq r \|f\|^2$  for each  $f \in \mathcal{V}$ , and  $\mathcal{F} = \{f_k\}_{k=1}^\infty$  an oblique dual frame of  $\tilde{\mathcal{G}}$  in  $\mathcal{W}$  which upper frame bound  $\beta$  satisfies  $\sqrt{\beta r} \leq \epsilon$ . Then,  $\mathcal{G}$  is an  $\epsilon$ -approximate dual frame of  $\mathcal{F}$  in  $\mathcal{V}$ .*

*Proof.* We will first see that  $\{g_k\}_{k=1}^\infty$  is a Bessel sequence. Let  $f \in \mathcal{V}$ . Applying the Cauchy-Schwartz inequality,

$$\sum_{k=1}^\infty |\langle f, g_k \rangle|^2 \leq \sum_{k=1}^\infty (|\langle f, g_k - \tilde{g} \rangle| + |\langle f, \tilde{g} \rangle|)^2$$

$$\begin{aligned}
&= \sum_{k=1}^{\infty} |\langle f, g_k - \tilde{g} \rangle|^2 + 2 \sum_{k=1}^{\infty} |\langle f, g_k - \tilde{g} \rangle| |\langle f, \tilde{g} \rangle| + \sum_{k=1}^{\infty} |\langle f, \tilde{g} \rangle|^2 \\
(6.1) \quad &\leq (r + 2\sqrt{r\tilde{\beta}} + \tilde{\beta}) \|f\|^2,
\end{aligned}$$

where  $\tilde{\beta}$  is the upper frame bound for  $\{\tilde{g}_k\}_{k=1}^{\infty}$ . Hence,  $\{g_k\}_{k=1}^{\infty}$  is a Bessel sequence with bound  $(\sqrt{r} + \sqrt{\tilde{\beta}})^2$ .

Note that the condition given for  $\{g_k\}_{k=1}^{\infty}$  can be written as  $\|T_{\mathcal{G}}^* - T_{\tilde{\mathcal{G}}}^*\| \leq \sqrt{r}$ .

We have

$$\|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{F}}T_{\mathcal{G}}^*\| = \|T_{\mathcal{F}}T_{\tilde{\mathcal{G}}}^* - T_{\mathcal{F}}T_{\mathcal{G}}^*\| \leq \|T_{\mathcal{F}}\| \|T_{\tilde{\mathcal{G}}}^* - T_{\mathcal{G}}^*\| \leq \sqrt{\beta r} \leq \epsilon.$$

Therefore,  $\{f_k\}_{k=1}^{\infty}$  and  $\{g_k\}_{k=1}^{\infty}$  are  $\epsilon$ -approximate oblique dual frames.  $\square$

**Proposition 6.2.** *Let  $\mathcal{F} = \{f_k\}_{k=1}^{\infty}$  be a frame for  $\mathcal{W}$  with bounds  $\alpha$  and  $\beta$ ,  $\epsilon \geq 0$  and  $\tilde{\mathcal{G}} = \{\tilde{g}_k\}_{k=1}^{\infty} \subset \mathcal{W}$  such that  $\sum_{k=1}^{\infty} |\langle f, f_k - \tilde{g} \rangle|^2 \leq r \|f\|^2$ , for  $f \in \mathcal{W}$ , with  $r \leq \frac{\alpha\epsilon^2}{(\|\pi_{\mathcal{V}\mathcal{W}^\perp}\| + \epsilon)^2}$ . Then,*

- (i)  $\tilde{\mathcal{G}}$  is a frame for  $\mathcal{W}$ .
- (ii) If  $\mathcal{G} = \{g_k\}_{k=1}^{\infty} \subset \mathcal{V}$  is given by  $g_k = \pi_{\mathcal{V}\mathcal{W}^\perp}(T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{G}}}^*)^\dagger \tilde{g}_k$ , then  $\mathcal{G}$  is an  $\epsilon$ -approximate oblique dual frame of  $\mathcal{F}$  in  $\mathcal{V}$ .

*Proof.* (i) Since  $r < \alpha$ , by [3, Corollary 22.1.5] it follows that  $\tilde{\mathcal{G}}$  is a frame for  $\mathcal{W}$  with bounds  $(\sqrt{\alpha} - \sqrt{r})^2$  and  $(\sqrt{r} + \sqrt{\beta})^2$ .

To see (ii), observe that, by [3, Lemma 5.1.5], the canonical dual frame  $\{(T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{G}}}^*)^\dagger \tilde{g}_k\}_{k=1}^{\infty}$  of  $\{\tilde{g}_k\}_{k=1}^{\infty}$  in  $\mathcal{W}$  has bounds  $\frac{1}{(\sqrt{r} + \sqrt{\beta})^2}$  and  $\frac{1}{(\sqrt{\alpha} - \sqrt{r})^2}$ .

By [4, Theorem 3.2], the sequence  $\mathcal{G} = \{g_k\}_{k=1}^{\infty} \subset \mathcal{V}$  given by  $g_k = \pi_{\mathcal{V}\mathcal{W}^\perp}(T_{\tilde{\mathcal{G}}}T_{\tilde{\mathcal{G}}}^*)^\dagger \tilde{g}_k$  is an oblique dual frame of  $\tilde{\mathcal{G}}$  in  $\mathcal{V}$ . Its synthesis operator satisfies  $\|T_{\mathcal{G}}\| \leq \frac{\|\pi_{\mathcal{V}\mathcal{W}^\perp}\|}{\sqrt{\alpha - \sqrt{r}}}$ . Hence,

$$\|\pi_{\mathcal{V}\mathcal{W}^\perp} - T_{\mathcal{G}}T_{\mathcal{F}}^*\| = \|T_{\mathcal{G}}T_{\tilde{\mathcal{G}}}^* - T_{\mathcal{G}}T_{\mathcal{F}}^*\| \leq \|T_{\mathcal{G}}\| \|T_{\tilde{\mathcal{G}}}^* - T_{\mathcal{F}}^*\| \leq \frac{\|\pi_{\mathcal{V}\mathcal{W}^\perp}\|}{\sqrt{\alpha - \sqrt{r}}} \sqrt{r} \leq \epsilon.$$

So  $\{f_k\}_{k=1}^{\infty}$  and  $\{g_k\}_{k=1}^{\infty}$  are  $\epsilon$ -approximate oblique dual frames.  $\square$

## 7. APPROXIMATE RECONSTRUCTION IN SHIFT-INVARIANT SUBSPACES

In this section we give conditions on the generators of shift-invariant subspaces  $\mathcal{W}$  and  $\mathcal{V}$  of  $\mathcal{H} = L^2(\mathbb{R})$ , in order to obtain an approximate reconstruction in one of them. We emphasize that for these results the subspaces don't necessarily decompose  $L^2(\mathbb{R})$  in direct sum, as we assumed before. As a consequence we obtain in the next section sufficient conditions on the generators for approximate oblique duality (see Corollaries 8.1 and 8.2).

In order to prove the results we will use this lemma:

**Lemma 7.1.** *Let  $\phi, \tilde{\phi} \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\tilde{\phi}\}_{k \in \mathbb{Z}}$  are Bessel sequences. The following holds:*

- (i) If  $f \in L^2(\mathbb{R})$ , then  $\sum_{n \in \mathbb{Z}} \widehat{f}(\cdot + n) \overline{\widehat{\tilde{\phi}}(\cdot + n)} \in L^2(0, 1)$  and
$$\sum_{k \in \mathbb{Z}} \langle \widehat{f}, \widehat{T_k\tilde{\phi}} \rangle \widehat{T_k\phi}(\gamma) = \widehat{\phi}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{f}(\gamma + n) \overline{\widehat{\tilde{\phi}}(\gamma + n)}.$$

(ii) If  $f \in \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$ , then

$$(7.1) \quad \sum_{k \in \mathbb{Z}} \langle \widehat{f}, \widehat{T_k\tilde{\phi}} \rangle \widehat{T_k\phi} = \widehat{f} \sum_{n \in \mathbb{Z}} \widehat{\phi}(\cdot + n) \overline{\widehat{\phi}(\cdot + n)}.$$

*Proof.* By the Cauchy-Schwarz inequality and Theorem 2.5, if  $f \in L^2(\mathbb{R})$  then

$$\int_0^1 |\sum_{n \in \mathbb{Z}} \widehat{f}(\cdot + n) \overline{\widehat{\phi}(\cdot + n)}|^2 \leq \beta \int_0^1 \sum_{n \in \mathbb{Z}} |\widehat{f}(\cdot + n)|^2 \leq \beta \|f\|^2,$$

where  $\beta$  is an upper frame bound of  $\{T_k\phi\}_{k \in \mathbb{Z}}$ . So, the equality in (i), which coincides with [4, Equality (12)], holds.

From (i), if  $f = \sum_{k \in \mathbb{Z}} c_k T_k\phi$ , where  $\{c_k\}_{k \in \mathbb{Z}} \in \ell^2(\mathbb{Z})$  has a finite number of nonzero elements,

$$\begin{aligned} \sum_{k \in \mathbb{Z}} \langle \widehat{f}, \widehat{T_k\tilde{\phi}} \rangle \widehat{T_k\phi}(\gamma) &= \widehat{\phi}(\gamma) \sum_{n \in \mathbb{Z}} \left( \sum_{k \in \mathbb{Z}} c_k E_{-k}(\gamma) \right) \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} \\ &= \left( \sum_{k \in \mathbb{Z}} c_k E_{-k}(\gamma) \right) \widehat{\phi}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} \\ &= \widehat{f}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)}. \end{aligned}$$

It follows that (7.1) holds for  $f \in \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$ . By continuity of the operator  $f \mapsto \sum_{k \in \mathbb{Z}} \langle f, T_k\tilde{\phi} \rangle T_k\phi$  and since  $\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} \in L^\infty(0, 1)$ , (7.1) holds for  $f \in \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$ .  $\square$

The key result of this section is the following:

**Theorem 7.2.** *Let  $\phi, \tilde{\phi} \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\tilde{\phi}\}_{k \in \mathbb{Z}}$  are Bessel sequences. Let  $\mathcal{W} = \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$ . Then, given  $\epsilon \geq 0$ , the following statements are equivalent:*

- (i)  $\|f - \sum_{k \in \mathbb{Z}} \langle f, T_k\tilde{\phi} \rangle T_k\phi\| \leq \epsilon \|f\|$ , for  $f \in \mathcal{W}$ .
- (ii)  $\left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} - 1 \right| \leq \epsilon$  for a.e.  $\gamma \in \{\gamma : \Phi(\gamma) \neq 0\}$ .

*Proof.* (ii)  $\Rightarrow$  (i): Let  $f = \sum_{k \in \mathbb{Z}} c_k T_k\phi$ , where  $\{c_k\}_{k \in \mathbb{Z}} \in \ell^2(\mathbb{Z})$  has a finite number of nonzero elements. Note that if  $\Phi(\gamma) = 0$  then  $\widehat{f}(\gamma) = 0$ . By Lemma 7.1 and (ii), we have that

$$\|f - \sum_{k \in \mathbb{Z}} \langle f, T_k\tilde{\phi} \rangle T_k\phi\| = \left\| \chi_{\{\Phi \neq 0\}} \widehat{f} \left[ 1 - \sum_{n \in \mathbb{Z}} \widehat{\phi}(\cdot + n) \overline{\widehat{\phi}(\cdot + n)} \right] \right\| \leq \epsilon \|\widehat{f}\| = \epsilon \|f\|.$$

So, as in the proof of Lemma 7.1, (i) holds for  $f \in \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$ .

(i)  $\Rightarrow$  (ii): Assume there exists  $E \subseteq \{\Phi \neq 0\}$  such that  $|E| > 0$  and

$$|\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} - 1| > \epsilon \text{ for } \gamma \in E.$$

We will see that there exists  $f \in \mathcal{W}$  such that

$$\|f - \sum_{k \in \mathbb{Z}} \langle f, T_k\tilde{\phi} \rangle T_k\phi\| > \epsilon \|f\|.$$

We can write  $E = \bigcup_{k \in \mathbb{N}} E_k$ , where

$$E_k = \left\{ \gamma \in \{\Phi \neq 0\} : \left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} - 1 \right| \geq \frac{1}{k} + \epsilon \right\}$$

for  $k \in \mathbb{N}$ . If  $|E_k| = 0$  for all  $k$ , then  $|E| = 0$ , which is a contradiction. So, there exist  $\epsilon' > \epsilon$  and  $E' \subseteq E$  such that  $|E'| > 0$  and

$$(7.2) \quad \left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} - 1 \right| \geq \epsilon' \text{ for } \gamma \in E'.$$

Let  $F \in L^2(0, 1)$  such that  $\text{supp}(F) \subseteq E'$ . Let  $f \in \mathcal{W}$  that verifies  $\widehat{f} = F\widehat{\phi}$ . By Lemma 7.1 and (7.2),

$$\begin{aligned} \|f - \sum_{k \in \mathbb{Z}} \langle f, T_k \tilde{\phi} \rangle T_k \phi\|^2 &= \left\| \widehat{f} \left[ 1 - \sum_{n \in \mathbb{Z}} \widehat{\phi}(\cdot + n) \overline{\widehat{\phi}(\cdot + n)} \right] \right\|^2 \\ &= \left\| \widehat{f} \left[ 1 - \sum_{n \in \mathbb{Z}} \widehat{\phi}(\cdot + n) \overline{\widehat{\phi}(\cdot + n)} \right] \right\|_{L^2(E')}^2 \\ &\geq (\epsilon')^2 \|\widehat{f}\|_{L^2(E')}^2 = (\epsilon')^2 \|f\|^2 > \epsilon^2 \|f\|^2, \end{aligned}$$

which contradicts (i).  $\square$

*Remark 7.3.* If  $\epsilon < 1$ , from condition (ii),  $\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} \neq 0$  for a.e.  $\gamma \in \{\Phi \neq 0\}$ . So  $\{\tilde{\Phi} = 0\} \subseteq \{\Phi = 0\}$  a.e..

Using the previous theorem we obtain a result about approximate dual frames in subspaces.

**Corollary 7.4.** *Let  $\phi, \tilde{\phi} \in L^2(\mathbb{R})$  be such that  $\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\{T_k \tilde{\phi}\}_{k \in \mathbb{Z}}$  are Bessel sequences. Let  $\mathcal{W} = \overline{\text{span}}\{T_k \phi\}_{k \in \mathbb{Z}}$ . Then, given  $0 \leq \epsilon < 1$ , the following assertions are equivalent:*

- (i)  $\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\{\underline{T_k P_{\mathcal{W}} \tilde{\phi}}\}_{k \in \mathbb{Z}}$  are  $\epsilon$ -approximate dual frames in  $\mathcal{W}$ .
- (ii)  $\left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}(\gamma + n)} - 1 \right| \leq \epsilon$  for a.e.  $\gamma \in \{\gamma : \Phi(\gamma) \neq 0\}$ .

*Proof.* The operators  $P_{\mathcal{W}}$  and  $T_k$  commute. Since  $\{T_k \tilde{\phi}\}_{k \in \mathbb{Z}}$  is a Bessel sequence, then  $\{T_k P_{\mathcal{W}} \tilde{\phi}\}_{k \in \mathbb{Z}} = \{P_{\mathcal{W}} T_k \tilde{\phi}\}_{k \in \mathbb{Z}}$  is too. Given  $f \in \mathcal{W}$ ,

$$\|f - \sum_{k \in \mathbb{Z}} \langle f, T_k P_{\mathcal{W}} \tilde{\phi} \rangle T_k \phi\| = \|f - \sum_{k \in \mathbb{Z}} \langle f, T_k \tilde{\phi} \rangle T_k \phi\|.$$

So the conclusion follows from Theorem 7.2 and Remark 3.3(iii).  $\square$

Let  $\{T_k \phi\}_{k \in \mathbb{Z}}$  be a Bessel sequence,  $\mathcal{W} = \overline{\text{span}}\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\phi_1 \in L^2(\mathbb{R})$ . The theorem below gives conditions so that there exists  $\tilde{\phi} \in \overline{\text{span}}\{T_k \phi_1\}_{k \in \mathbb{Z}}$  which translates allow approximate reconstruction in  $\mathcal{W}$ .

**Theorem 7.5.** *Let  $\phi, \phi_1 \in L^2(\mathbb{R})$  be such that  $\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\{T_k \phi_1\}_{k \in \mathbb{Z}}$  are Bessel sequences. Let  $\mathcal{W} = \overline{\text{span}}\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\mathcal{V} = \overline{\text{span}}\{T_k \phi_1\}_{k \in \mathbb{Z}}$ . Let  $\tilde{\phi} \in \mathcal{V}$  such that  $\widehat{\tilde{\phi}} = \widehat{H} \widehat{\phi_1}$  with  $\widehat{H} \in L^\infty(0, 1)$ . Assume  $\epsilon \geq 0$ . Then  $\{T_k \tilde{\phi}\}_{k \in \mathbb{Z}}$  is a Bessel sequence and the following are equivalent:*

- (1)  $\|f - \sum_{k \in \mathbb{Z}} \langle f, T_k \tilde{\phi} \rangle T_k \phi\| \leq \epsilon \|f\|$  for  $f \in \mathcal{W}$ .
- (2)  $\left| \widehat{H}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi_1}(\gamma + n)} - 1 \right| \leq \epsilon$  for a.e.  $\gamma \in \{\gamma : \Phi(\gamma) \neq 0\}$ .

*Proof.* Observe that

$$\tilde{\Phi}(\gamma) = \sum_{k \in \mathbb{Z}} |\widehat{H}(\gamma) \widehat{\phi_1}(\gamma + k)|^2 = |\widehat{H}(\gamma)|^2 \Phi_1(\gamma)$$

$$\leq \|\tilde{H}\|_{L^\infty(0,1)}^2 \Phi_1(\gamma) \leq \|\tilde{H}\|_{L^\infty(0,1)}^2 \beta_1,$$

where  $\beta_1$  is the Bessel bound of  $\{T_k\phi_1\}_{k \in \mathbb{Z}}$ . So, by Theorem 2.5,  $\{T_k\tilde{\phi}\}_{k \in \mathbb{Z}}$  is a Bessel sequence.

Since  $\left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi}(\gamma+n)} - 1 \right| = \left| \overline{\tilde{H}(\gamma)} \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi}_1(\gamma+n)} - 1 \right|$ , the conclusion follows from Theorem 7.2.  $\square$

If in the previous statements  $\epsilon = 0$ , the results of [4] for oblique duality are recovered.

From Theorem 7.5 and (2.3) we obtain the following sufficient condition for approximate reconstruction:

**Corollary 7.6.** *Let  $\phi, \phi_1 \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\phi_1\}_{k \in \mathbb{Z}}$  are Bessel sequences with  $\{\Phi_1 = 0\} \subseteq \{\Phi = 0\}$  a.e. and Bessel bounds  $\beta$  and  $\beta_1$ . Let  $\mathcal{W} = \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$ ,  $\mathcal{V} = \overline{\text{span}}\{T_k\phi_1\}_{k \in \mathbb{Z}}$  and  $\tilde{\phi} \in \mathcal{V}$  be such that  $\widehat{\tilde{\phi}} = \tilde{H}\widehat{\phi_1}$  with  $\tilde{H} \in L^\infty(0,1)$ . Assume  $\epsilon \geq 0$ . Then  $\{T_k\tilde{\phi}\}_{k \in \mathbb{Z}}$  is a Bessel sequence and if*

$$(7.3) \quad \left| \overline{\tilde{H}(\gamma)} - \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi}_1(\gamma+n)}} \right| \leq \frac{\epsilon}{\sqrt{\beta\beta_1}}$$

for a.e.  $\gamma \in \{\Phi \neq 0\}$ , then

$$\|f - \sum_{k \in \mathbb{Z}} \langle f, T_k\tilde{\phi} \rangle T_k\phi\| \leq \epsilon \|f\| \quad \text{for } f \in \mathcal{W}.$$

The next result about approximate dual frames in shift-invariant subspaces is a consequence of Corollary 7.4, with a proof similar to the one of Theorem 7.5.

**Corollary 7.7.** *Let  $\phi, \phi_1 \in L^2(\mathbb{R})$  such that  $\{T_k\phi\}_{k \in \mathbb{Z}}$ ,  $\{T_k\phi_1\}_{k \in \mathbb{Z}}$  are Bessel sequences with  $\{\Phi_1 = 0\} = \{\Phi = 0\}$  a.e. Set  $\mathcal{W} = \overline{\text{span}}\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\mathcal{V} = \overline{\text{span}}\{T_k\phi_1\}_{k \in \mathbb{Z}}$ . Assume that  $\tilde{\phi} \in \mathcal{V}$  and  $\tilde{\phi}_1 \in \mathcal{W}$  are such that  $\widehat{\tilde{\phi}} = \tilde{H}\widehat{\phi_1}$  and  $\widehat{\tilde{\phi}_1} = \tilde{H}\widehat{\phi}$  with  $\tilde{H} \in L^\infty(0,1)$ . Then  $\{T_k\tilde{\phi}\}_{k \in \mathbb{Z}}$ ,  $\{T_k\tilde{\phi}_1\}_{k \in \mathbb{Z}}$  are Bessel sequences and if*

$$\left| \overline{\tilde{H}(\gamma)} \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1(\gamma+n)}} - 1 \right| \leq \epsilon$$

a.e.  $\gamma \in \{\Phi \neq 0\}$  with  $0 \leq \epsilon < 1$ , then  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k P_{\mathcal{W}}\tilde{\phi}\}_{k \in \mathbb{Z}}$  are  $\epsilon$ -approximate dual frames in  $\mathcal{W}$ , whereas  $\{T_k\phi_1\}_{k \in \mathbb{Z}}$  and  $\{T_k P_{\mathcal{V}}\tilde{\phi}_1\}_{k \in \mathbb{Z}}$  are  $\epsilon$ -approximate dual frames in  $\mathcal{V}$ .

## 8. APPROXIMATE OBLIQUE DUAL FRAMES IN SHIFT-INVARIANT SPACES

In this section we study the concept of approximate oblique duality when  $\mathcal{H} = L^2(\mathbb{R})$  and  $\mathcal{W}$  and  $\mathcal{V}$  are shift-invariant subspaces of  $L^2(\mathbb{R})$ . In what follows we assume that  $L^2(\mathbb{R}) = \mathcal{W} \oplus \mathcal{V}^\perp$ .

The following corollaries give conditions on the generators of two shift-invariant subspaces for approximate oblique duality. The first is a consequence of Theorem 7.2 and Proposition 4.2.

**Corollary 8.1.** *Let  $\phi, \tilde{\phi} \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\tilde{\phi}\}_{k \in \mathbb{Z}}$  are frames for  $\mathcal{W}$  and  $\mathcal{V}$  respectively. Assume  $\epsilon \geq 0$ . If*

$$\left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\tilde{\phi}}(\gamma+n)} - 1 \right| \leq \frac{\epsilon}{\|\pi_{\mathcal{W}\mathcal{V}^\perp}\|} \quad \text{for a.e. } \gamma \in \{\gamma : \Phi(\gamma) \neq 0\};$$

then  $\{T_k\phi\}_{k\in\mathbb{Z}}$  and  $\{T_k\tilde{\phi}\}_{k\in\mathbb{Z}}$  are  $\epsilon$ -approximate oblique dual frames.

**Corollary 8.2.** *Let  $\phi, \phi_1 \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k\in\mathbb{Z}}$  and  $\{T_k\phi_1\}_{k\in\mathbb{Z}}$  are frames for  $\mathcal{W}$  and  $\mathcal{V}$  respectively. Let  $\tilde{\phi} \in \mathcal{V}$  be such that  $\tilde{\phi} = \tilde{H}\widehat{\phi_1}$  with  $\tilde{H} \in L^2(0, 1)$ . Assume  $0 \leq \epsilon < 1$ . If*

$$\left| \tilde{H}(\gamma) \sum_{n\in\mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)} - 1 \right| \leq \frac{\epsilon}{\|\pi_{\mathcal{W}\mathcal{V}^\perp}\|} \text{ for a.e. } \gamma \in \{\gamma : \Phi(\gamma) \neq 0\}$$

then  $\{T_k\tilde{\phi}\}_{k\in\mathbb{Z}}$  is a frame for  $\mathcal{V}$  such that  $\{T_k\phi\}_{k\in\mathbb{Z}}$  and  $\{T_k\tilde{\phi}\}_{k\in\mathbb{Z}}$  are  $\epsilon$ -approximate oblique dual frames.

*Proof.* By Proposition 2.6,  $\{\Phi = 0\} = \{\Phi_1 = 0\}$  and there exists  $c > 0$  such that

$$\left| \tilde{H}(\gamma) \right| \leq \frac{1}{|\sum_{n\in\mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}|} \left( 1 + \frac{\epsilon}{\|\pi_{\mathcal{W}\mathcal{V}^\perp}\|} \right) \leq \frac{1}{c} \left( 1 + \frac{\epsilon}{\|\pi_{\mathcal{W}\mathcal{V}^\perp}\|} \right).$$

for a.e.  $\gamma \in \{\gamma : \Phi(\gamma) \neq 0\}$ . So, by Theorem 2.5,

$$\tilde{\Phi}(\gamma) = |\tilde{H}(\gamma)|^2 \Phi_1(\gamma) \leq \frac{1}{c^2} \left( 1 + \frac{\epsilon}{\|\pi_{\mathcal{W}\mathcal{V}^\perp}\|} \right)^2 \beta_1,$$

where  $\beta_1$  is an upper frame bound of  $\{T_k\phi_1\}_{k\in\mathbb{Z}}$ . Applying again Theorem 2.5 we conclude that  $\{T_k\tilde{\phi}\}_{k\in\mathbb{Z}}$  is a Bessel sequence.

Now the conclusion follows from Theorem 7.5, Proposition 4.2 and Remark 3.3(iii).  $\square$

The following theorem gives an expression for the Fourier transform of the oblique projection when the subspaces are shift-invariant, in terms of the corresponding generators.

**Theorem 8.3.** *Let  $\phi, \phi_1 \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k\in\mathbb{Z}}$  and  $\{T_k\phi_1\}_{k\in\mathbb{Z}}$  are frames for  $\mathcal{W}$  and  $\mathcal{V}$  respectively. Then*

$$\widehat{\pi_{\mathcal{W}\mathcal{V}^\perp} f}(\gamma) = \begin{cases} \frac{\sum_{n\in\mathbb{Z}} \widehat{f}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}}{\sum_{n\in\mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}} \widehat{\phi}(\gamma) & \gamma \in \mathcal{N}^c(\Phi) \\ 0 & \gamma \in \mathcal{N}(\Phi) \end{cases}$$

for  $f \in L^2(\mathbb{R})$ .

*Proof.* Let  $f \in L^2(\mathbb{R})$ . Since  $\{T_k\phi\}_{k\in\mathbb{Z}}$  is a frame for  $\mathcal{W}$  there exists  $\tau \in L^2(0, 1)$  such that  $\widehat{\pi_{\mathcal{W}\mathcal{V}^\perp} f} = \tau \widehat{\phi}$ . So, if  $\gamma \in \mathcal{N}(\Phi)$ ,  $\widehat{\pi_{\mathcal{W}\mathcal{V}^\perp} f}(\gamma) = 0$ . By [8, Lemma 2.8],

$$\begin{aligned} \sum_{n\in\mathbb{Z}} \widehat{f}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)} &= \sum_{n\in\mathbb{Z}} \widehat{\pi_{\mathcal{W}\mathcal{V}^\perp} f}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)} \\ &= \tau(\gamma) \sum_{n\in\mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}. \end{aligned}$$

If  $\gamma \in \{\Phi \neq 0\}$ , by Proposition 2.6,  $\sum_{n\in\mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)} \neq 0$ , hence  $\tau(\gamma) = \frac{\sum_{n\in\mathbb{Z}} \widehat{f}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}}{\sum_{n\in\mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}}$  and  $\widehat{\pi_{\mathcal{W}\mathcal{V}^\perp} f}(\gamma) = \frac{\sum_{n\in\mathbb{Z}} \widehat{f}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}}{\sum_{n\in\mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}} \widehat{\phi}(\gamma)$ .  $\square$

If  $\mathcal{W} = \mathcal{V}$  the previous theorem reduces to [8, Theorem 2.9].

Using the expression for the Fourier transform of the oblique projection given in Theorem 8.3, we obtain the following sufficient condition for approximate duality which is different to the one of Corollary 8.2. If  $\{T_k\phi\}_{k\in\mathbb{Z}}$  and  $\{T_k\phi_1\}_{k\in\mathbb{Z}}$  are Riesz bases for  $\mathcal{W}$  and  $\mathcal{V}$  respectively, we have an equality in (2.4) [20]. So, in this case the sufficient condition of the next theorem is weaker than the one of Corollary 8.2.



**Theorem 8.4.** *Let  $\phi, \phi_1 \in L^2(\mathbb{R})$  be such that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\phi_1\}_{k \in \mathbb{Z}}$  are frames for  $\mathcal{W}$  and  $\mathcal{V}$  respectively. Let  $0 \leq \epsilon < 1$ . If  $\psi \in \mathcal{V}$  verifies  $\widehat{\psi} = H\widehat{\phi_1}$ , where  $H \in L^2(0, 1)$  is such that*

$$(8.1) \quad \left| \overline{H(\gamma)} - \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)}} \right| \leq \frac{\epsilon}{\sqrt{\Phi(\gamma)\Phi_1(\gamma)}}$$

for a.e.  $\gamma \in \{\Phi \neq 0\}$ , then  $\{T_k\phi\}_{k \in \mathbb{Z}}$ ,  $\{T_k\psi\}_{k \in \mathbb{Z}}$  are  $\epsilon$ -approximate oblique dual frames.

*Proof.* Let  $f \in L^2(\mathbb{R})$ . Let  $\epsilon \geq 0$  and  $H \in L^2(0, 1)$  that verify (8.1). Let  $\psi \in \mathcal{V}$  be such that  $\widehat{\psi} = H\widehat{\phi_1}$ . Then, using Lemma 7.1 and Theorem 8.3,

$$\begin{aligned} & \|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k \in \mathbb{Z}} \langle f, T_k\psi \rangle T_k\phi\|^2 = \|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k \in \mathbb{Z}} \langle \widehat{f}, E_k\widehat{\psi} \rangle E_k\widehat{\phi}\|^2 \\ &= \int_{\{\Phi \neq 0\}} \left| \frac{\sum_{n \in \mathbb{Z}} \widehat{f}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)}}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)}} \widehat{\phi}(\gamma) - \widehat{\phi}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{f}(\gamma+n)\overline{\widehat{\psi}(\gamma+n)} \right|^2 d\gamma \\ &= \int_{\{\Phi \neq 0\}} \left| \widehat{\phi}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{f}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)} \left( \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)}} - \overline{H(\gamma)} \right) \right|^2 d\gamma \\ &\leq \int_{\{\Phi \neq 0\}} \Phi(\gamma) \left| \sum_{n \in \mathbb{Z}} \widehat{f}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)} \left( \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)}} - \overline{H(\gamma)} \right) \right|^2 d\gamma \\ &\leq \int_{\{\Phi \neq 0\}} \Phi(\gamma)\Phi_1(\gamma) \sum_{n \in \mathbb{Z}} |\widehat{f}(\gamma+n)|^2 \left| \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)}} - \overline{H(\gamma)} \right|^2 d\gamma \\ &\leq \epsilon^2 \int_{\{\Phi \neq 0\}} \sum_{n \in \mathbb{Z}} |\widehat{f}(\gamma+n)|^2 d\gamma = \epsilon^2 \int_{\{\Phi \neq 0\}} |\widehat{f}(\gamma)|^2 d\gamma \leq \epsilon^2 \|\widehat{f}\|^2 dx = \epsilon^2 \|f\|^2. \end{aligned}$$

Hence  $\|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k \in \mathbb{Z}} \langle f, T_k\psi \rangle T_k\phi\| \leq \epsilon \|f\|$  for  $f \in \mathcal{H}$ . So,  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\psi\}_{k \in \mathbb{Z}}$  are  $\epsilon$ -approximate oblique dual frames.  $\square$

*Remark 8.5.* Assume that there exists  $E \subseteq \{\Phi \neq 0\}$  with  $|E| > 0$  and that

$$\left| \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)}} - \overline{H(\gamma)} \right| > \frac{\epsilon}{\sqrt{\Phi(\gamma)\Phi_1(\gamma)}}$$

for  $\gamma \in E$ . Let  $\epsilon' > 0$  with  $\epsilon' < \frac{c\epsilon}{\sqrt{\beta\beta_1}}$  (where  $c > 0$  is the constant of Proposition 2.6). Note that  $\epsilon' < \frac{c\epsilon}{\sqrt{\Phi(\gamma)\Phi_1(\gamma)}} \leq \frac{c\epsilon}{|\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n)\overline{\widehat{\phi_1}(\gamma+n)}|} \leq \epsilon$ .

We are going to see that  $\{T_k\phi\}_{k \in \mathbb{Z}}$  and  $\{T_k\psi\}_{k \in \mathbb{Z}}$  are not  $\epsilon'$ -approximate oblique dual frames, i.e., there exists  $f \in L^2(\mathbb{R})$  such that

$$\|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k \in \mathbb{Z}} \langle f, T_k\phi \rangle T_k\phi\| > \epsilon' \|f\|.$$

Let  $F \in L^2(0, 1)$  such that  $\text{supp}(F) \subseteq E$  and  $f \in L^2(\mathbb{R})$  given by  $\widehat{f} = F\widehat{\phi}$ . Then  $f \in \mathcal{W}$ . Therefore

$$\|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k \in \mathbb{Z}} \langle f, T_k\psi \rangle T_k\phi\|^2 =$$

$$\begin{aligned}
&= \int_{\{\Phi \neq 0\}} \sum_{n \in \mathbb{Z}} |\widehat{f}(\gamma + n)|^2 \left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)} \left( \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)}} - \overline{H}(\gamma) \right) \right|^2 d\gamma \\
&\geq \epsilon^2 \int_{\{\Phi \neq 0\}} \sum_{n \in \mathbb{Z}} |\widehat{f}(\gamma + n)|^2 \frac{\left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)} \right|^2}{\Phi(\gamma) \Phi_1(\gamma)} d\gamma \\
&\geq \frac{\epsilon^2 \epsilon^2}{\beta \beta_1} \int_{\{\Phi \neq 0\}} \sum_{n \in \mathbb{Z}} |\widehat{f}(\gamma + n)|^2 d\gamma > \epsilon'^2 \int_{\{\Phi \neq 0\}} \sum_{n \in \mathbb{Z}} |\widehat{f}(\gamma + n)|^2 d\gamma \\
&= \epsilon'^2 \int |\widehat{f}(\gamma)|^2 d\gamma = \epsilon'^2 \|f\|^2
\end{aligned}$$

So,

$$\|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k \in \mathbb{Z}} \langle f, T_k \widehat{\phi} \rangle T_k \phi\| > \epsilon' \|f\|.$$

We have the following necessary condition for approximate oblique duality.

**Theorem 8.6.** *Let  $\phi, \phi_1 \in L^2(\mathbb{R})$  be such that  $\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\{T_k \phi_1\}_{k \in \mathbb{Z}}$  are frames for  $\mathcal{W}$  and  $\mathcal{V}$  respectively. Let  $0 \leq \epsilon < 1$ . Set  $\psi \in \mathcal{V}$  such that  $\widehat{\psi} = H \widehat{\phi}_1$ , with  $H \in L^2(0, 1)$ . If  $\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\{T_k \psi\}_{k \in \mathbb{Z}}$  are  $\epsilon$ -approximate oblique dual frames, then*

$$(8.2) \quad \left| \overline{H}(\gamma) - \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)}} \right| \leq \frac{\epsilon}{\left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)} \right|}$$

for a.e.  $\gamma \in \{\Phi \neq 0\}$ .

*Proof.* We proceed in a similar way to the proof of Theorem 7.2. Assume there exists  $E \subseteq \{\Phi \neq 0\}$  with  $|E| > 0$  such that

$$\left| \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)}} - \overline{H}(\gamma) \right| > \frac{\epsilon}{\left| \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)} \right|} \text{ for } \gamma \in E.$$

Then

$$\left| 1 - \overline{H}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)} \right| > \epsilon \text{ for } \gamma \in E.$$

There exist  $\epsilon' > \epsilon$  and  $E' \subseteq E$  such that  $|E'| > 0$  and

$$\left| 1 - \overline{H}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)} \right| \geq \epsilon' \text{ for } \gamma \in E'.$$

Consider  $F \in L^2(0, 1)$  such that  $\text{supp}(F) \subseteq E'$  and  $f \in \mathcal{W}$  with  $\widehat{f} = F \widehat{\phi}$ . Hence,

$$\begin{aligned}
&\|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k \in \mathbb{Z}} \langle f, T_k \psi \rangle T_k \phi\|^2 \\
&= \int_{\{\Phi \neq 0\}} \left| \frac{\sum_{n \in \mathbb{Z}} \widehat{f}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)}}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)}} \widehat{\phi}(\gamma) - \widehat{\phi}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{f}(\gamma + n) \overline{\widehat{\psi}(\gamma + n)} \right|^2 d\gamma \\
&= \int_{\{\Phi \neq 0\}} |\widehat{f}(\gamma)|^2 \left| 1 - \overline{H}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)} \right|^2 d\gamma \\
&= \int_{\{\Phi \neq 0\}} \sum_{n \in \mathbb{Z}} |\widehat{f}(\gamma + n)|^2 \left| 1 - \overline{H}(\gamma) \sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma + n) \overline{\widehat{\phi}_1(\gamma + n)} \right|^2 d\gamma
\end{aligned}$$

$$\geq \epsilon'^2 \int_{\{\Phi \neq 0\}} \sum_{n \in \mathbb{Z}} |\widehat{f}(\gamma + n)|^2 d\gamma = \epsilon'^2 \int_{\{\Phi \neq 0\}} |\widehat{f}(\gamma)|^2 d\gamma > \epsilon^2 \|f\|^2.$$

Consequently,

$$\|\pi_{\mathcal{W}\mathcal{V}^\perp} f - \sum_{k \in \mathbb{Z}} \langle f, T_k \tilde{\phi} \rangle T_k \phi\| > \epsilon \|f\|.$$

This contradicts the fact that  $\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\{T_k \psi\}_{k \in \mathbb{Z}}$  are  $\epsilon$ -approximate oblique dual frames.  $\square$

Let  $\phi, \phi_1 \in L^2(\mathbb{R})$  be such that  $\{T_k \phi\}_{k \in \mathbb{Z}}$  and  $\{T_k \phi_1\}_{k \in \mathbb{Z}}$  are frames for  $\mathcal{W}$  and  $\mathcal{V}$  respectively. The unique oblique dual frame of  $\{T_k \phi\}_{k \in \mathbb{Z}}$  in  $\mathcal{V}$  is  $\{T_k \tilde{\phi}\}_{k \in \mathbb{Z}}$  where  $\tilde{\phi} = \sum_{k \in \mathbb{Z}} c_k T_k \phi_1$  and  $\{c_k\}_{k \in \mathbb{Z}}$  are the Fourier coefficients of  $\tilde{H}(\gamma) = \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}}$ .

In practice, the generator of the oblique dual frame can only be obtained approximately since the computation of the previous series and the calculus of the Fourier coefficients are not exact. So, in fact, in the applications we always have to work with an approximate oblique dual frame. In the following example we illustrate this situation. We consider here the effect of the truncation required for the series that defines  $\tilde{\phi}$  when  $\tilde{H}$  is not a trigonometric polynomial. In this case, the used  $\epsilon$ -approximate dual frame is  $\{T_k \psi_K\}_{k \in \mathbb{Z}}$  where  $\psi_K = \sum_{|k| \leq K} c_k T_k \phi_1$  for some  $K \in \mathbb{N}$ . Taking into account (8.1) we have the error

$$\epsilon = \sup_{\gamma \in [0,1]} \left| \sqrt{\Phi(\gamma)\overline{\Phi_1(\gamma)}} \left| \tilde{H}(\gamma) - \frac{1}{\sum_{n \in \mathbb{Z}} \widehat{\phi}(\gamma+n) \overline{\widehat{\phi_1}(\gamma+n)}} \right| \right|$$

where  $H(\gamma) = \sum_{|k| \leq K} c_k e^{-2i\pi k\gamma}$ .

**Example 8.7.** We consider shift-invariant subspaces generated by  $B$ -splines (see [4, Example 4.5]). Let  $\phi = B_1$  and  $\phi_1 = B_3$  where  $B_1$  and  $B_3$  are the  $B$ -splines of order 1 and 3 respectively. In this case

$$\tilde{H}(\gamma) = \frac{1}{\sum_{n \in \mathbb{Z}} \left( \frac{\sin \pi(\gamma+n)}{\pi(\gamma+n)} \right)^4}$$

is not a trigonometric polynomial. As a consequence, the generator of the oblique dual frame of  $\{T_k B_1\}_{k \in \mathbb{Z}}$  in  $\mathcal{V}$  has not compact support whereas for the generator  $\psi_K$  of the approximate oblique dual frame we have  $\text{supp}(\psi_K) = [-\frac{3+2K}{2}, \frac{3+2K}{2}]$ . The next table shows the supports and the estimated errors for different values of  $K$ .

$K$	$\text{supp}(\psi_K)$	$\epsilon$
0	$[-1.5, 1.5]$	0.732050720745298
1	$[-2.5, 2.5]$	0.196152304804580
2	$[-3.5, 3.5]$	0.052558805558347
3	$[-4.5, 4.5]$	0.014083071590247
4	$[-5.5, 5.5]$	0.003773541369707

TABLE 1. The supports and the estimated errors for different values of  $K$ .

In Figure 1 we plot at the left the generator  $\psi_4$  of the approximate oblique dual frame which support is  $[-5.5, 5.5]$ , and at the right the generators  $\tilde{\phi}$  and  $\psi_4$  in the interval  $[3.5, 8.5]$ .




FIGURE 1. (a) The generator  $\psi_4$  of the approximate oblique dual frame; (b) The generators  $\tilde{\phi}$  (dashed line) and  $\psi_4$  (solid line).

#### ACKNOWLEDGEMENT

This research has been supported by Grants PIP 112-201501-00589-CO (CONICET), PROICO 03-1618 (UNSL), PICT-2014-1480 and UBACyT 20020130100422BA.

#### REFERENCES

- [1] M. A. Akinlar and J. P. Gabardo, *Oblique duals associated with rational subspace Gabor frames*, J. Integral Equations Appl. **20** (2008), no. 3, 283–309.
- [2] P. Berger, K. Gröchenig and G. Matz; *Sampling and reconstruction in distinct subspaces using oblique projections*, J. Fourier Anal. Appl. **25** (2019), 1080–1112.
- [3] O. Christensen, *An introduction to frames and Riesz bases*, 2nd edn., Birkhäuser, Boston, 2016.
- [4] O. Christensen and Y. C. Eldar, *Oblique dual frames and shift-invariant spaces*, Appl. Comput. Harmon. Anal. **17** (2004), 48–68.
- [5] O. Christensen and R. S. Laugesen, *Approximately dual frames in Hilbert spaces and applications to Gabor frames*, Sampl. Theory in Signal and Image Process. **9** (2011), 77–90.
- [6] I. Daubechies, *Ten Lectures on Wavelets*, SIAM, Philadelphia, PA, 1992.
- [7] I. Daubechies, A. Grossmann and Y. Meyer, *Painless nonorthogonal expansions*, J. Math. Phys. **27** (1986), 1271–1283.
- [8] C. de Boor, R. DeVore and A. Ron, *Approximation from shift-invariant subspaces of  $L^2(\mathbb{R}^d)$* , Trans. Amer. Math. Soc. **341** (1994), 787–806.
- [9] M. Dörfler and E. Matusiak; *Nonstationary Gabor frames - approximately dual frames and reconstruction errors*, Adv. Comput. Math. **41** (2015), 293–316.
- [10] R. J. Duffin and A. C. Schaeffer, *A class of nonharmonic Fourier series*, Trans. Amer. Math. Soc. **72** (1952), 341–366.
- [11] Y. C. Eldar, *Sampling with arbitrary sampling and reconstruction spaces and oblique dual frame vectors*, J. Fourier Anal. Appl. **9** (2003), no. 1, 77–96.
- [12] Y. C. Eldar, *Sampling without input constrains: consistent reconstruction in arbitrary spaces*. In Sampling, Wavelets and Tomography, A. Zayed and J. J. Benedetto Eds., pp. 33–60, Birkhäuser Boston, 2003.

- [13] Y. C. Eldar and O. Christensen, *Characterization of oblique dual frame pairs*, EURASIP J. Appl. Signal Process. (2006), 1–11.
- [14] Y. C. Eldar and T. Werther, *General framework for consistent sampling in Hilbert spaces*, Int. J. Wavelets Multiresolut. Inf. Process. **3** (2005), no. 4, 497–509.
- [15] H. G. Feichtinger, D. M. Onchis and C. Wiesmeyr, *Construction of approximate dual wavelet frames*, Adv. Comput. Math. **40** (2014), 273–282.
- [16] S. B. Heineken and P. M. Morillas, *Oblique dual fusion frames*, Numer. Funct. Anal. Optim. **39** (2018), no. 7, 800–824.
- [17] C. Heil (Ed.): Harmonic Analysis and Applications. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston, 2006.
- [18] A. A. Hemmat and J. P. Gabardo, *Properties of oblique dual frames in shift-invariant systems*, J. Math. Anal. Appl. **356** (2009), no. 1, 346–354.
- [19] W. S. Tang, *Oblique projections, biorthogonal Riesz bases and multiwavelets in Hilbert spaces*, Proc. Amer. Math. Soc. **128** (2000), no. 2, 463–473.
- [20] M. Unser and A. Aldroubi, *A general sampling theory for non-ideal acquisition devices*, IEEE Trans. Signal Process. **42** (1994), 2915–2925.
- [21] R. Young, *An Introduction to Nonharmonic Fourier Series*, Academic, New York (1980) (revised first edition 2001).
- [22] W. Zhang, *Dual and approximately dual Hilbert-Schmidt frames in Hilbert spaces*, Results Math. **73** (2018), no. 4.