

# Beating Random Assignment for Approximating Quantum 2-Local Hamiltonian Problems

Ojas Parekh\* and Kevin Thompson†

## Abstract

The quantum  $k$ -Local Hamiltonian problem is a natural generalization of classical constraint satisfaction problems ( $k$ -CSP) and is complete for QMA, a quantum analog of NP. Although the complexity of  $k$ -Local Hamiltonian problems has been well studied, only a handful of approximation results are known. For Max 2-Local Hamiltonian where each term is a rank 3 projector, a natural quantum generalization of classical Max 2-SAT, the best known approximation algorithm was the trivial random assignment, yielding a 0.75-approximation. We present the first approximation algorithm beating this bound, a classical polynomial-time 0.764-approximation. For strictly quadratic instances, which are maximally entangled instances, we provide a 0.801 approximation algorithm, and numerically demonstrate that our algorithm is likely a 0.821-approximation. We conjecture these are the hardest instances to approximate. We also give improved approximations for quantum generalizations of other related classical 2-CSPs. Finally, we exploit quantum connections to a generalization of the Grothendieck problem to obtain a classical constant-factor approximation for the physically relevant special case of strictly quadratic traceless 2-Local Hamiltonians on bipartite interaction graphs, where a inverse logarithmic approximation was the best previously known (for general interaction graphs). Our work employs recently developed techniques for analyzing classical approximations of CSPs and is intended to be accessible to both quantum information scientists and classical computer scientists.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Previous Work . . . . .	4
1.2	Overview of Our Work . . . . .	5
1.3	Outlook and Conjectures . . . . .	9
<b>2</b>	<b>Semidefinite Relaxation and Rounding Approach</b>	<b>10</b>
2.1	Quantum Information Notation . . . . .	10
2.2	Formal Problem Statement . . . . .	10
2.3	Semidefinite Relaxation . . . . .	11
2.4	Rounding Approach and Formal Statement of Results . . . . .	14
2.5	Analysis Overview . . . . .	16
<b>3</b>	<b>Notations</b>	<b>18</b>

---

\*Sandia National Laboratories, *email*: odparek@sandia.gov

†Sandia National Laboratories, *email*: kevthom@sandia.gov

<b>4</b>	<b>Standard Form for the Expectation of an Edge</b>	<b>20</b>
4.1	Moment Matrix Description of Two Qubit Quantum States . . . . .	20
4.2	SVD Argument . . . . .	21
<b>5</b>	<b>Proof of the Main Technical Lemmas</b>	<b>24</b>
5.1	Hermite Polynomials . . . . .	25
5.1.1	Definitions and Expansions . . . . .	25
5.1.2	Hermite Expansions of Interest . . . . .	26
5.1.3	Bounds on Hermite Expansions . . . . .	28
5.2	Optimizing Rational Functions over Polytopes . . . . .	30
5.3	Technical Estimates for Functions of Interest . . . . .	31
5.4	Proof of Lemma 17 . . . . .	32
5.5	Proof of Lemma 18 . . . . .	33
<b>6</b>	<b>Proof of Main Theorems</b>	<b>34</b>
6.1	Proof of Theorem 11 (strictly quadratic case) . . . . .	34
6.2	Proof of Theorem 10 . . . . .	35
<b>7</b>	<b>Conclusion</b>	<b>37</b>
7.1	Acknowledgments . . . . .	37
<b>A</b>	<b>Connecting 2-Local Hamiltonians and Classical 2-CSPs</b>	<b>41</b>
<b>B</b>	<b>Identities</b>	<b>42</b>
<b>C</b>	<b>Calculation of Hermite Coefficients</b>	<b>44</b>
<b>D</b>	<b>Lower Bounds for the Expectation of an Edge</b>	<b>48</b>
<b>E</b>	<b>Technical Quantum Facts</b>	<b>53</b>
<b>F</b>	<b>Traceless, Bipartite, and Strictly Quadratic Hamiltonians</b>	<b>57</b>

## 1 Introduction

The design and analysis of approximation algorithms [WS11, Vaz13] is an extensively studied area in theoretical computer science. In this setting, we are given some (generally NP-hard) optimization problem, and we are tasked with producing a valid (feasible) solution with objective within some provable factor of the optimal objective value. To understand this formally imagine we are given some optimization problem  $\mathcal{P}$ , which corresponds to an infinite set of problem instances  $\{P_i\}$ . Each problem instance corresponds to a triple  $P_i = (f_i, \mathcal{T}_i, i_n)$  where  $\mathcal{T}_i \subseteq \{0, 1\}^{i_n}$  and each triple corresponds to an optimization problem of the form:

$$OPT_i = \max_{v \in \mathcal{T}_i} f_i(v).$$

An approximation algorithm  $\mathcal{A}$  acts on an efficient description of the instance to produce some feasible solution to the problem:  $\mathcal{A}(f_i, \mathcal{T}_i, i_n) = v_i^* \in \mathcal{T}_i$  in time polynomial in the instance size (polynomial in  $i_n$ ). It is said that algorithm has approximation factor  $\alpha$  for  $0 < \alpha \leq 1$  if in the

worst-case (over all instances), the solution produced by the algorithm is a factor of  $\alpha$  off of the optimal answer:

$$\min_i \frac{f_i(v_i^*)}{OPT_i} \geq \alpha.$$

Since we should not expect to solve NP-hard problems, the interesting question is then the approximability of NP-hard optimization problems, or the study of which approximation factors  $\alpha$  are obtainable for different problems. As one might expect, approximability is highly problem sensitive and there are many classes of natural problems with very different attainable approximation factors [Hås99, DS05, Vaz13]. So, in general approximation algorithms are developed in the context of specific problems, and many such algorithms are known [Vaz13, WS11].

**2-Local Hamiltonian.** In stark contrast, although QMA-hard quantum optimization problems arise naturally through well-known physically motivated problems [BS07, SV09], they have very few known approximation algorithms with provable approximation factors [BBT09, GK12, BH16, HM17, BGKT19, GP19, HLP20, AGM20]<sup>1</sup>. The QMA-hard optimization studied in these works, as well as the problem we will study here, is the *2-Local Hamiltonian problem* [KKR06, KSVV02]. An instance of this problem is specified by a problem size,  $n$ , as well as a set of 2-local interactions,  $\{H_e\}$ . Each  $H_e$  is some *local* Hamiltonian which can be written as the tensor product of  $n-2$  identity terms with some nontrivial operator that acts on at most 2 qubits, i.e.  $H_e = \mathcal{O}_{ij} \otimes (\mathbb{I}_2)^{\otimes(n-2)}$ . The optimization problem corresponding to a particular instance is to find the smallest or largest eigenvalue,  $\lambda_{min}$  or  $\lambda_{max}$ , of  $H = \sum_e H_e$ . Ideally, an algorithm solving this problem would also produce a description of or access to a corresponding eigenvector. An approximation algorithm,  $\mathcal{A}$ , acts on the size of the problem ( $n$ ) and a description of the local Hamiltonians  $\{H_e\}$  to produce a classical description of a valid quantum state. Once again we say that the algorithm achieves approximation factor  $\alpha$  if:

$$\frac{\text{Tr}[(\sum_e H_e) \mathcal{A}(n, \{H_e\})]}{\lambda_{max}(\sum_e H_e)} \geq \alpha \text{ for all instances.}$$

Generally we assume some property of the Hamiltonian which forces  $\lambda_{max}(\sum_e H_e) > 0$  so that this is a sensible definition. A common assumption [KKR06, GK12, HLP20] is that the terms  $H_e$  are positive semi-definite (PSD) and nonzero. We note that when all of the terms  $H_e$  are taken to be diagonal projectors (in say, the standard computational basis), the corresponding instance of 2-Local Hamiltonian corresponds precisely to an instance of the classical 2-Constraint-Satisfaction problem (2-CSP). In this case, the 4 diagonal entries of  $\mathcal{O}_{ij}$  correspond to the  $\{0, 1\}$  output values of a Boolean function on Boolean variables  $x_i$  and  $x_j$  corresponding to  $i$  and  $j$ . See Appendix A for more details as well as a classical motivation for 2-Local Hamiltonian. In addition Table 1 highlights classical 2-CSP specializations of quantum 2-local Hamiltonian problems for which approximation algorithms are known.

The 2-Local Hamiltonian problem is interesting in many different contexts of physics and quantum information [KKR06, KSVV02, Osb12]. This problem is manifestly interesting to physicists because the 2-local nature of the problem matches the local nature of many physical systems (spin chains, Ising model, etc.). Hence, the study of eigenstates and energies is of utmost importance, and has been since the beginnings of quantum mechanics itself [Bet31]. From a theoretical computer science perspective, the 2-Local Hamiltonian problem is interesting for the same reasons that classical approximation algorithms are interesting. Under standard complexity theoretic assumptions,

<sup>1</sup>Here and throughout this paper we mean a *classical* algorithm which takes as input a classical description of a quantum problem and produces a classical description of a quantum state. An approximation algorithm for a QMA-hard problem can have several natural meanings distinct from this (quantum input, quantum algorithm which produces classical output, etc.)

we should not expect to be able to solve the problem, so the interesting direction is the study of the approximability of the problem. Can we find rigorous approximation algorithms, and how well can we expect to be able to approximate the answer? Moreover, which classes of instances admit constant-factor approximation algorithms?

The generic 2-Local Hamiltonian problem is a generalization of several classical optimization problems [WB03, BGKT19] with very different approximability. Maximum independent set is one such example [WB03], and it is well known that such a problem cannot be approximated to within a constant factor unless  $P=NP$ , so we should not expect the 2-Local Hamiltonian problem to have a constant-factor approximation algorithm which holds uniformly for all instances. In light of this fact researchers make specific assumptions on the terms  $H_e$ , and attempt to find approximation algorithms under these assumptions. The 2-Local Hamiltonian instances we consider generalize a variety of classical optimization problems, including Max 2-SAT, Max Cut, general Max 2-CSP, and the Grothendieck problem (see Table 1 for our results).

## 1.1 Previous Work

In the interest of describing classical approximation algorithms for 2-Local Hamiltonian, let  $OPT = \lambda_{\max}(\sum_{e \in E} H_e)$  be the largest eigenvalue of an instance of 2-Local Hamiltonian,  $H = \sum_{e \in E} H_e$ , and let

$$OPT_{prod} = \max_{\substack{|\phi_1\rangle, \dots, |\phi_n\rangle \\ \in \mathbb{C}^2}} \langle \phi_1 | \otimes \dots \otimes \langle \phi_n | H | \phi_1 \rangle \otimes \dots \otimes | \phi_n \rangle$$

be the product state<sup>2</sup> with the largest objective value or *energy*.

One common assumption is on the *geometry* of the interactions in  $E$ . Bansal, Bravyi, and Terhal show that 2-Local Hamiltonian on bounded-degree planar graphs admits a polynomial-time approximation scheme<sup>3</sup> (PTAS) [BBT09], and Brandão and Harrow generalize this to arbitrary planar graphs [BH16]. On the other end, for  $k$ -Local Hamiltonian on dense graphs, Gharibian and Kempe give a PTAS with respect to  $OPT_{prod}$  [GK12], and Brandão and Harrow extend this result to obtain a PTAS for dense graphs with respect to  $OPT$  [BH16]. Brandão and Harrow also show the existence of good product-states or give product-state approximations for a variety of graph classes [BH16].

Many authors make assumptions on the form of the terms  $H_e$ . One common assumption is that each  $H_e$  is traceless [BGKT19, HM17], or equivalently that they each can be written as a linear combination of tensor products of Pauli operators, excluding the identity operator. Unfortunately, this case is still general enough to capture problems with no constant factor approximation algorithms [ABE<sup>+</sup>05] (under complexity theoretic assumptions), although Bravyi, Gosset, König, and Temme give an approximation algorithm for traceless 2-Local Hamiltonian with guarantee that depends inverse logarithmically on the problem size [BGKT19]. Harrow and Montanaro give an approximation algorithm for traceless  $k$ -Local Hamiltonian with respect to the maximum degree and size of the interaction hypergraph [HM17]. Other authors make assumptions which force particular physically relevant forms for the terms  $H_e$  [GP19], or on the rank of the terms  $H_e$  [HLP20].

It is these latter works which our work is most easily compared to, so we provide descriptions of them here (the approximation guarantees are specified in Table 1). A unifying theme among them is that they rely upon a semi-definite program (SDP) to provide an upper bound on  $OPT$  and then

<sup>2</sup>As is clear from the expression, a product state is a quantum state which factors according to tensor product of individual quantum states. Such states have no entanglement and are considered “classical” states.

<sup>3</sup>This is an approximation algorithm that allows an arbitrarily good, but constant, approximation factor at the expense of an increase in runtime.

use generalization of a classical randomized rounding scheme to produce a product state [BGKT19, GP19, HLP20]. Such an approach was first carried out by Brandão and Harrow [BH16].

The work of Hallgren, Lee, and Parekh [HLP20] assumes that the terms  $H_e$  are all PSD, and that each one is some projector of fixed rank, which is also our primary problem of interest. They solve an SDP relaxation of  $OPT$  and use the solution they get to “round” to a valid quantum product state, say  $|\psi\rangle\langle\psi|$ . Employing results used in classical SDP rounding algorithms [BdOFV10, GW95] in a black-box fashion, along with a rounding scheme designed to handle 1-local terms, the authors are able to show that the quantum state from the rounding algorithm is within some fraction of the optimal state, or that  $\langle\psi|\sum_e H_e|\psi\rangle \geq \alpha OPT$ . The approximation factors they obtain are worse than the maximally mixed state (random assignment) for all cases except when  $rank(H_e) = 1$ , and here they are able to get the first approximation factor beating random assignment. They also provide an approximation algorithm when each  $H_e$  is a product state, which is a QMA-hard class of 2-Local Hamiltonian.

Another result which is particularly relevant to our work is that of Gharibian and Parekh [GP19], who consider a QMA-hard 2-Local Hamiltonian generalization of the classical Max Cut problem. Here the authors assume a particular form for the  $H_e$  by assuming that there are only specific terms in the Pauli decomposition. They use a simpler SDP relaxation than [HLP20], and they round a solution of this SDP to a product state using a result of Briët, de Oliveira Filho, and Vallentin [BdOFV14] in a black box fashion to prove the approximation factor. Improved approximation results for this problem, that go beyond using product states, have been recently obtained by Anshu, Gosset, and Morenz [AGM20].

For traceless Hamiltonians, the main result of interest is that of Bravyi, Gosset, König, and Temme [BGKT19]. This work builds off [CW04, HM17] and provides an inverse logarithmic approximation factor for generic traceless Hamiltonians. Such Hamiltonians necessarily have a non-negative and a non-positive eigenvalue. Note that such a class generalizes all problems considered in this paper, since adding copies of the identity can only make the approximation factor better, however there is no reason to expect their analysis could be used to prove constant factor approximations for the classes we study. The SDP relaxation is the same as [GP19]; however, the rounding scheme and analysis is a generalization of a classical approximation by Charikar and Wirth [CW04].

## 1.2 Overview of Our Work

There are two problems of interest to us. The first is the general problem of finding the largest eigenvalue of a 2-local traceless Hamiltonian on a bipartite interaction graph, and the second problem is finding the largest eigenvalue of a 2-local Hamiltonian where all the local terms are projectors (eigenvalues are 0 or 1).

**Traceless Hamiltonians.** For the traceless case, we consider Hamiltonians on bipartite interaction graphs that are “strictly quadratic.” Informally, the latter means that the 2-local terms of the Hamiltonian do not contain any implicit 1-local terms (see Definition 5 in Section 2.2). The classical analog of a strictly quadratic traceless Hamiltonian is a multilinear quadratic polynomial that does not contain linear terms (see Appendix A for connections between Hamiltonians and multilinear polynomials).

A natural classical analog of this problem is the symmetric Grothendieck problem [BdOFV14, FL20]. In this problem the objective is to maximize a quadratic form of a set of variables,  $\mathbf{z}^T A \mathbf{z}$ , subject to the constraint that each of the variables  $\mathbf{z}_i \in \{\pm 1\}$ , and where we assume diagonal elements of  $A$  are 0. The strictly quadratic nature of the problem is apparent, since the objective is a quadratic form. It is also apparent that  $A$  is traceless, but this is not the real reason the analogy

Table 1: **Summary of our and related results.** The number of qubits or Boolean variables is  $n$ . For readability, we omit weights  $w_{ij} \geq 0$  that may be present in both 2-local Hamiltonian (2-LH) and 2-CSP problems. Appendix A provides more details on the relationship between 2-LH and 2-CSP, as well as definitions for  $X_i, Y_i, Z_i$ . An “N” denotes a numerical result, and the classical results are implicitly numerical. The abbreviation “quad.” refers to strictly quadratic instances.

Max 2-LH problem (QMA-hard)	Max 2-CSP specialization (NP-hard)	Classical approx. for 2-CSP	Classical approx. for 2-LH problem (product state)
<b>Traceless</b> $\sum_{ij \in E} H_{ij} \otimes \mathbb{I}_{[n] \setminus \{ij\}}$ $H_{ij}$ has no $\mathbb{I}$ terms	<b>Classical Ising</b> $\max -\sum_{ij \in E} z_i z_j$ $z_i \in \{\pm 1\}$	$\Omega(\frac{1}{\log n})$ [CW04]	$\Omega(\frac{1}{\log n})^\dagger$ [BGKT19]
<b>Bipartite Traceless</b> $\sum_{e \in E} H_{ij} \otimes \mathbb{I}_{[n] \setminus \{ij\}}$ $H_{ij}$ has no $\mathbb{I}$ terms $E$ bipartite	<b>Grothendieck</b> $\max -\sum_{ij \in E} z_i z_j$ $z_i \in \{\pm 1\}$ $E$ bipartite	$0.561 + \epsilon$ [BMMN13]	$0.187^\dagger$ (quad.)
<b>Positive/Rank 1</b> $\sum_{ij \in E} H_{ij} \otimes \mathbb{I}_{[n] \setminus \{ij\}}$ $\mathbb{I} \succeq H_{ij} \succeq 0$ ( $\equiv H_{ij}$ rank 1 projector)	<b>Max 2-CSP</b> ( $\equiv 1$ satisfying assignment per clause)	$0.874$ [LLZ02]	$0.25$ (random) $0.328$ [HLP20] <b>0.387</b> <b>0.467 (quad.)</b> <b>0.498 (quad., N)</b> $0.5$ (upper bound)
<b>Max Heisenberg</b> $\sum_{ij \in E} \mathbb{I} - X_i X_j - Y_i Y_j - Z_i Z_j$ (special case of above)	<b>Max Cut</b> $\max \sum_{ij \in E} 1 - z_i z_j$ $z_i \in \{\pm 1\}$	$0.878$ [GW95]	$0.25$ (random) $0.498$ [GP19] $0.5$ (upper bound) $0.53^*$ [AGM20]
<b>Rank 2</b> $\sum_{ij \in E} H_{ij} \otimes \mathbb{I}_{[n] \setminus \{ij\}}$ $H_{ij}$ rank 2 projector	<b>Max 2-CSP</b> with 2 satisfying assignments/clause	$0.874$ [LLZ02]	$0.5$ (random) <b>0.565</b> <b>0.639 (quad.)</b> <b>0.653 (quad., N)</b> $0.667$ (upper bound)
<b>2-QSAT</b> $\sum_{ij \in E} H_{ij} \otimes \mathbb{I}_{[n] \setminus \{ij\}}$ $H_{ij}$ rank 3 projector	<b>Max 2-SAT</b> ( $\equiv 3$ satisfying assignments/clause)	$0.940$ [LLZ02]	$0.75$ (random) <b>0.764</b> <b>0.805 (quad.)</b> <b>0.821 (quad., N)</b> $0.834$ (upper bound)

\*This exceeds the product-state upper bound because it is achieved by a classical approximation algorithm that rounds to a non-product state.

$\dagger$ For any traceless 2-LH problem, we obtain a product-state approximation ratio that is  $\frac{1}{3}$  of an approximation ratio for a related classical CSP, using the appropriate classical approximation algorithm as a black box (see Appendix F).

is appropriate. Note that, since  $\mathbf{z}_i \in \{\pm 1\}$ , we will always pick up the diagonal elements of  $A$ :  $\mathbf{z}_i^2 A_{ii} = A_{ii}$ . Hence, finding the exact solution to the case where  $\text{Tr}[A] \neq 0$  is equivalent to finding the exact solution when  $\text{Tr}[A] = 0$ . On the quantum side, there is a standard decomposition for local Hamiltonians such that  $H = \alpha \mathbb{I} + \beta \mathcal{O}$ , where  $\mathcal{O}$  is some traceless operator. Since we are trying to find  $\max \langle \psi | (\alpha \mathbb{I} + \beta \mathcal{O}) | \psi \rangle$  for normalized  $|\psi\rangle$ , we will always pick up the constant  $\alpha$  independent of our choice of  $|\psi\rangle$ , hence we get an equivalent optimization problem for any value of  $\alpha$ . This is why the analogy is appropriate, we are optimizing over a quadratic form with the extra identity contribution subtracted off.

The rounding algorithm we use is a simple modification of a known algorithm [BdOFV14] for solving a variant of the symmetric Grothendieck problem (which also uses the bipartite assumption). Since we are able to use [BdOFV14] in a black-box fashion, the technical details of that algorithm are not needed for this paper. Essentially, we use this result to obtain  $\{\pm 1\}$  variables such that the objective upper bounds the optimal quantum objective. Rounding to a quantum state is then easily accomplished by dividing these variables by a large enough constant that they can be taken to be Bloch vectors for a valid quantum state. In the end, we obtain a  $\frac{2 \ln(1+\sqrt{2})}{3\pi}$ -approximation algorithm. Note that the best previously known result is a  $\Omega(\frac{1}{\log(n)})$ -approximation by Bravyi, Gosset, König, and Temme [BGKT19] on general graphs. More generally, our approach allows one to obtain a product-state approximation algorithm for traceless instances by using an approximation algorithm for a related classical CSP as a black box, losing a factor of 3 in the approximation ratio. This also gives a more direct means of obtaining the result of Bravyi, Gosset, König, and Temme [BGKT19]. Since this result is disparate from our main results, we present both the formal statements and analysis in Appendix F.

**2-Local projectors.** The main contribution of this work is the second problem we mentioned, finding the largest eigenvalue of a 2-local Hamiltonian where each of the local terms are projectors. In order to understand our rounding algorithm, we must first understand the local nature of the 2-local Hamiltonian problem. Let  $H = \sum_e H_e$  be the 2-local Hamiltonian where each  $H_e$  is a local term affecting only two qubits, say qubits  $i$  and  $j$ . Let  $\rho \in \mathbb{C}^{2^n \times 2^n}$  be the optimal density matrix. The value of the objective can be calculated as  $\text{Tr}[\rho H] = \sum_e \text{Tr}[\rho H_e] = \sum_e \text{Tr}[\rho_{ij} \widetilde{H}_e]$ , where  $\rho_{ij} \in \mathbb{C}^{4 \times 4}$  is the marginal density matrix on qubits  $i, j$ , and  $\widetilde{H}_e \in \mathbb{C}^{4 \times 4}$  is the 2-local part of the term  $H_e \in \mathbb{C}^{2^n \times 2^n}$  that acts on  $i, j$ . Here  $\widetilde{H}_e$  plays a role analogous to a classical Boolean constraint on 2 variables. Hence, if we were given the set of marginals  $\{\rho_{ij}\}$ , and descriptions of the  $\widetilde{H}_e$ , we could calculate the objective on a polynomial sized classical computer since  $\rho_{ij}, \widetilde{H}_e \in \mathbb{C}^{4 \times 4}$ . This implies that if we were able to optimize the set  $\{\rho_{ij}\}$  subject to the constraint that the  $\rho_{ij}$  were valid marginals of a global density matrix, we would be able to solve the local Hamiltonian problem. Indeed, the issue here is that deciding if a set of marginal density matrices is globally consistent is itself a QMA-complete problem [Liu06, BG19].

A natural question is then whether or not, in polynomial time, we can impose some global constraint which is weaker than consistency. Optimizing with this constraint would provide a relaxation, hence a polynomial-time-computable upper bound on the optimal objective. Then one might use generalizations of standard classical recipes [GW95] for deriving an approximation algorithm from the relaxation. In the classical case, a problem of this form can be constrained using a semidefinite constraint on a “moment matrix” (e.g. [RT12]). A moment matrix “tracks” low order statistics of a global probability distribution. These statistics have the property that they are only a function of marginal distributions and that the moment matrix is guaranteed to be PSD if the marginals are consistent (although the converse does hold in general). Hence, defining the moment matrix and forcing it to be PSD gives a weaker condition than global consistency, which can easily

be checked. We will adopt the same approach, except we will be tracking local *quantum* statistics. Our approach is related to existing hierarchies of quantum moment matrices [DLTW08, PNA10]. We will have, as variables in our optimization problem, a set of marginal distributions  $\{\rho_{ij}\}$ , as well as an overall moment matrix  $M$ , where each entry of  $M$  can be evaluated using at most one specific  $\rho_{ij}$ . Just as in the classical case, the matrix  $M$  will have the property that  $M \succeq 0$  for a consistent set of marginals.

Once we have the relaxation, we can efficiently solve it (it will be a polynomially-large SDP) to obtain an optimal moment matrix  $M^*$  and optimal marginals  $\{\rho_{ij}^*\}$  for the relaxation. The solution will have the property that  $\{\rho_{ij}^*\}$  likely represent a globally inconsistent set of marginals, and the objective will be larger than the objective for the optimal quantum state (recall we are applying a condition that is weaker than consistency). The task for us is to then generate a set of consistent density matrices  $\{\rho_{ij}\}$  with quantifiable loss in objective. The loss of objective will correspond to an approximation factor, as is the theme in many works. The approach we take is to (randomly) generate single qubit marginals  $\{\rho_i\}$  from the set of two qubit marginals  $\{\rho_{ij}^*\}$  and output the consistent density matrix  $\bigotimes_i \rho_i$ . Note that, just as in the classical case, non-overlapping marginals can always be assumed consistent. Understanding the loss in objective then reduces to understanding the loss of objective due to the random rounding procedure.

**Analysis.** The analysis proceeds as one might expect, by linearity we can reduce the expectation  $\mathbb{E}[\text{Tr}[\bigotimes_i \rho_i \sum_e H_e]] = \sum_e \mathbb{E}[\text{Tr}[\bigotimes_i \rho_i H_e]]$ . Then, a bound on the “worst-case” edge provides a bound on the expectation overall. The issue with accomplishing this directly is the number of parameters involved. Arguments concerning classical problems do not have to contend with this. For example, Max Cut has all terms proportional to  $(1 - z_i z_j)$  for  $z_i, z_j$  scalar variables. Max 2-SAT has 4 kinds of clauses depending on negation of the variables; it has nowhere near the variability of a generic 2-local projector. Naively, it is determined by 16 parameters, since  $\widetilde{H}_e$  is a  $4 \times 4$  matrix. Therefore the first task in our analysis is to reduce the terms  $\widetilde{H}_e$  to a standard form, specified by a small number of parameters. This is accomplished with some singular value decompositions, as well as exploiting the rotational invariance of standard multivariate normal distributions, and applying some results concerning 2-qubit density matrices or projectors [Gam16]. Given the standard form, calculating the expected objective reduces to calculating the expectations of certain functions of multivariate normal variables. This is analyzed using expansions in Hermite polynomials, and the resulting expressions are bounded or given in terms of special functions. The analysis itself is partially inspired by [BdOFV14, BGG<sup>+</sup>18], and we are able to obtain a result of [BdOFV14] as a special case of our analysis. The final results we obtain are summarized here informally:

**Theorem 1** (Informal). *Given a 2-local Hamiltonian problem  $\{H_e\}$  where all  $H_e$  are proportional to 2-local projectors with  $\widetilde{H}_e$  of rank  $k \in \{1, 2, 3\}$ , we give a classical randomized polynomial-time algorithm with approximation ratio  $\alpha(k)$  where*

$$\alpha(k) = \begin{cases} 0.387 & \text{if } k = 1 \\ 0.565 & \text{if } k = 2 \\ 0.764 & \text{if } k = 3. \end{cases}$$

**Theorem 2** (Informal). *If in addition to the assumptions of Theorem 1, the terms  $H_e$  are strictly quadratic, we give a classical randomized polynomial-time algorithm with approximation ratio  $\alpha(k)$ ,*



where

$$\alpha(k) = \begin{cases} 0.467 & \text{if } k = 1 \\ 0.639 & \text{if } k = 2 \\ 0.805 & \text{if } k = 3. \end{cases}$$

Projectors on 2 qubits have rank at most 4, so there are three cases of interest for this problem, rank 1, rank 2, and rank 3. We obtain novel approximation factors for each case. The decision version of the problem we consider is known as Quantum-SAT and was introduced in 2006 by Bravyi [Bra11], and the approximability of Max Quantum-SAT was first considered in 2011 by Gharibian and Kempe [GK12], who observed that the maximally mixed state trivially achieves an approximation ratio of  $k/4$ , where  $k \in \{1, 2, 3\}$  is the rank of the projectors  $\widetilde{H}_e$ . The only nontrivial result previously known is a 0.328-approximation for the  $k = 1$  case by Hallgren, Lee, and Parekh [HLP20]. The main difference between our work and the previous works [GP19, HLP20] is that we are able to directly analyze the expectation rather than appealing to other works in a black-box fashion. Both of these works appealed to approximation results of Briët, de Oliveira Filho, and Vallentin [BdOFV10, BdOFV14]. Our analysis may be seen as a generalization of the result of Briët, de Oliveira Filho, and Vallentin [BdOFV14] employed in [GP19].

### 1.3 Outlook and Conjectures

**Strictly quadratic instances.** We believe the strictly quadratic case is an interesting special case for several reasons. As noted, one of the difficulties in analyzing rounding schemes for 2-Local Hamiltonian is the sheer number of parameters involved. The quadratic case reduces the number of parameters to consider, while still including physically relevant QMA-hard instances such as the Max Heisenberg model that serves as a quantum generalization of Max Cut [GP19]. Indeed we believe that quadratic instances allow one to glean insights and develop techniques that might otherwise be obscured in more general instances. One of the first rigorous approximation algorithms for a 2-local Hamiltonian that goes beyond product states was developed for a quadratic instance, and one that has considerably fewer parameters than those we consider here [AGM20]. Moreover, maximally entangled instances are quadratic, and we conjecture these are the hardest cases to approximate.

Additionally, we believe that the analysis we provide is tighter for this case. Since there are no linear terms in the objective, we can focus on one particular Hermite expansion and carefully bound it. For the general case we must contend with understanding the linear and quadratic terms simultaneously, which makes the analysis much more difficult. We conjecture that the true performance of our algorithm for the general case is:

**Conjecture 3.** *(Informal) Our rounding algorithm achieves approximation ratio:*

$$\alpha(k) = \begin{cases} 0.498 & \text{if } k = 1 \\ 0.653 & \text{if } k = 2 \\ 0.821 & \text{if } k = 3. \end{cases}$$

We have indeed attempted to verify the approximation ratios in Conjecture 3 numerically, and via both sampling and integration methods, we observe that the worst-case instances for our algorithm match the values stated in the conjecture. The difficulty in taking these encouraging results as fact is that sampling is of course not exhaustive, and the integrals exhibit poor convergence. We do give upper bounds on  $\alpha(k)$  in Theorem 40 in the Appendix. These are derived by furnishing instances on 2 qubits that demonstrate a gap between the maximum eigenvalue and the maximum

objective value achieved by a product state. These bounds are  $\frac{1}{2}$  ( $k = 1$ ),  $\frac{2}{3}$  ( $k = 2$ ), and  $\frac{5}{6}$  ( $k = 3$ ), which are fairly close the values in Conjecture 3. We suspect that one may be able to prove that the approximation ratios in Conjecture 3 are best possible under the unique games conjecture.

**Significance of our work.** We give the first approximation algorithm beating random assignment for Max 2-QSAT (and related problems). We show how to move beyond numerical evaluation of approximation ratios for 2-local Hamiltonian problems, which is not as critical in the classical case that enjoys only a handful of parameters. This is accomplished by bringing analysis employing Hermite polynomials to bear on 2-local Hamiltonian problems. Moreover, our rounding scheme is a simple and natural generalization of hyperplane rounding.

## 2 Semidefinite Relaxation and Rounding Approach

In this section we present a rigorous but high-level overview of our approach, with technical lemmas deferred to later sections. We define the main problems considered and our semidefinite relaxation and rounding algorithm. We conclude by motivating the analysis that will occur in subsequent sections.

### 2.1 Quantum Information Notation

We adopt some standard notations used in quantum information [NC10]. The *kets*  $|0\rangle := [0, 1]$  and  $|1\rangle := [1, 0]$  represent the standard basis vectors for  $\mathbb{C}^2$ , while the *bras*  $\langle 0|$  and  $\langle 1|$  represent their conjugate transposes. The  $d \times d$  identity matrix is denoted by  $\mathbb{I}_d$ , and the subscript will be omitted when redundant. We obtain the standard bases for  $\mathbb{C}^{2^n}$  as  $|b_1 b_2 \dots b_n\rangle := |b_1\rangle |b_2\rangle \dots |b_n\rangle := |b_1\rangle \otimes |b_2\rangle \otimes \dots \otimes |b_n\rangle$ , with  $b_i \in \{0, 1\}$ . The Pauli matrices will have the usual definition:

$$\sigma^0 = \mathbb{I} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \sigma^1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma^2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \text{and} \quad \sigma^3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (1)$$

We will generally use subscripts to indicate quantum subsystems. If  $\rho$  is a density matrix on  $n$  qubits, for instance,  $\rho_{ij}$  will correspond to the marginal density matrix on qubits  $i$  and  $j$ , i.e. the partial trace  $\rho_{ij} = \text{Tr}_{[n] \setminus \{i, j\}}[\rho]$  (e.g. [NC10], Section 2.4.3). Similarly,  $\sigma_i^j$  corresponds to Pauli  $j$  on qubit  $i$ . Subscripts will supercede position in many cases in the paper, for instance  $\sigma_i^1 \otimes \sigma_j^2 \otimes \mathbb{I}_{[n] \setminus \{i, j\}}$  is meant as  $\mathbb{I} \otimes \mathbb{I} \otimes \dots \otimes \sigma^1 \otimes \dots \otimes \mathbb{I} \otimes \sigma^2 \otimes \dots \otimes \mathbb{I}$  where  $\sigma^1$  is at the  $i$ th position and  $\sigma^2$  is at the  $j$ th position. We encourage readers familiar with classical constraint satisfaction problems to consult Appendix A, which casts such problems as quantum local Hamiltonian problems.

### 2.2 Formal Problem Statement

In the 2-Local Hamiltonian problem we are given as input the problem size,  $n$ , and a list of classical descriptions of 2-local terms,  $\{H_e\}_{e \in E}$ . We allow multiedges, i.e. distinct edges  $e$  and  $e'$  on the same pair of qubits  $i, j$ ; we will use the notation  $e_1$  and  $e_2$  to refer to the qubits on which  $e$  acts. In this context a term  $H_e$  is 2-local if it can be written in the form  $H_e = \mathcal{O}_e \otimes \mathbb{I}_{[n] \setminus \{e_1, e_2\}}$ , using the subscript notation from Section 2.1. Local Hamiltonians have polynomially-sized descriptions which can be given in terms of the local operators  $\mathcal{O}_e$ , but for our purposes the details of the description will not be important. It is important to note at this point that we will use  $\text{rank}(H_e)$  to mean  $\text{rank}(\mathcal{O}_e)$ . The actual rank of  $H_e$  is  $\text{rank}(\mathcal{O}_e)2^{n-2}$ , but for ease of exposition we will say that the “rank” of a 2-local term is equal to the rank of its non-trivial part. We are tasked with determining the largest eigenvalue of the Hamiltonian  $H = \sum_{e \in E} H_e$ :

**Problem 4** (QLH( $n, \{H_e\}_{e \in E}$ )). *Given a problem size,  $n$ , as well as a classical description of a set of 2-local terms  $\{H_e\}$  with  $H_e \in \mathbb{C}^{2^n \times 2^n}$  Hermitian, find:*

$$\lambda_{\max}(H) := \max_{|\phi\rangle \in [\mathbb{C}^2]^{\otimes n}} \text{Tr} \left[ \sum_{e \in E} H_e |\phi\rangle \langle \phi| \right] = \max_{\substack{\rho \in \mathbb{C}^{2^n} \times \mathbb{C}^{2^n} \\ \text{Tr}(\rho)=1, \rho \succeq 0}} \text{Tr} \left[ \sum_{e \in E} H_e \rho \right].$$

The main problems of interest to us are instances of QLH where each term is a projector and the special case where each projector is strictly quadratic. The strictly quadratic case precludes non-identity 1-local terms (i.e.  $\mathcal{O}_i \otimes \mathbb{I}_{[n] \setminus \{i\}}$  with  $\mathcal{O}_i \neq \mathbb{I}$ ) that may be implicit in a 2-local term.

**Definition 5** (Strictly Quadratic). *Let  $H_e$  be a 2-local term on  $n$  qubits. Write  $H_e = \mathcal{O}_e \otimes \mathbb{I}_{[n] \setminus \{e_1, e_2\}}$  for some nontrivial operator  $\mathcal{O}_e$ . Express  $\mathcal{O}_e$  in the Pauli basis as:*

$$\mathcal{O}_e = \sum_{k,l=0}^3 \alpha_{k,l} \sigma^k \otimes \sigma^l. \quad (2)$$

We say that  $H_e$  is a strictly quadratic if  $\alpha_{k,0} = 0$  for all  $k \neq 0$ , and  $\alpha_{0,l} = 0$  for all  $l \neq 0$ .

Note that the coefficients in Equation (2) may be obtained as  $\alpha_{k,l} = \text{Tr}[\sigma^k \otimes \sigma^l \mathcal{O}_e]/4$  and are real since  $\mathcal{O}_e$  is Hermitian.

We focus our attention on QLH restricted to projectors and strictly quadratic projectors, both of which remain QMA-hard [PM15]. In this case  $\mathcal{O}_e = w_e P_e$ , where  $P_e$  is a 2-qubit projector, and  $w_e \geq 0$  is a weight. There are three interesting cases, depending on the rank of  $P_e$ . We will obtain approximation factors for each.

**Problem 6** (QLHP( $n, k, \{H_e\}_{e \in E}$ )). *Given a problem size,  $n$ , as well as a classical description of a set of 2-local terms  $\{H_e = w_e P_e \otimes \mathbb{I}_{[n] \setminus \{e_1, e_2\}}\}$  with  $w_e \geq 0$  and  $P_e \in \mathbb{C}^{4 \times 4}$  a 2-qubit projector of rank at least  $k$ , find  $\lambda_{\max}(\sum_{e \in E} H_e)$ .*

It is worth mentioning that since any  $\mathcal{O}_e \succeq 0$  can be written as a positive combination of rank 1 projectors, QLHP with  $k = 1$  captures instances of QLH where each  $H_e \succeq 0$ .

### 2.3 Semidefinite Relaxation

We employ a semidefinite programming relaxation for QLH (Problem 4) that is a refinement of the now standard SDP relaxation that has been used in designing approximation algorithms [BH16, BGKT19, GP19]. Our relaxation is related to one used by Hallgren, Lee, and Parekh [HLP20] and may be viewed as a specialization of noncommutative Lasserre hierarchies proposed for quantum information applications [DLTW08, PNA10].

In this section we assume, for the sake of exposition, that there is a single edge  $ij$  on any pair of qubits  $i, j \in [n]$ ; however, the relaxation and rounding algorithm may be readily extended to handle general instances of QLH with multiedges. Suppose we have an instance of QLH on  $n$  qubits. As previously stated, the first set of variables in our SDPs will be marginal density matrices  $\{\rho_{ij}\}$ . Since there are  $n$  qubits, there are  $\binom{n}{2}$  many of these, and each of them is a  $4 \times 4$  Hermitian matrix. While we cannot impose global consistency, we can force each  $\rho_{ij}$  to be a valid density matrix on its own:  $\text{Tr}[\rho_{ij}] = 1$  and  $\rho_{ij} \succeq 0$  for all  $i, j \in [n]$ . We could also explicitly force overlapping marginals to be consistent on single qubit density matrices, however this will be implicit through our use of moment matrices.

**Moment matrices.** Suppose we have a quantum state on  $n$  qubits  $|\psi\rangle \in \mathbb{C}^{2^n}$ . Consider the 1-local Pauli operators,  $\mathcal{M} = \{\mathbb{I}\} \cup \{\sigma_i^k \otimes \mathbb{I}_{[n]\setminus\{i\}} \mid k \in [3], i \in [n]\}$ . We apply each of the  $3n + 1$  Pauli operators  $\mathcal{O} \in \mathcal{M}$  on  $|\psi\rangle$  to obtain columns of a matrix  $V = \{\mathcal{O}|\psi\rangle\}_{\mathcal{O} \in \mathcal{M}} \in \mathbb{C}^{2^n \times (3n+1)}$ . We call  $M := V^\dagger V \in \mathbb{C}^{(3n+1) \times (3n+1)}$  the *moment matrix* of  $|\psi\rangle$  with respect to  $\mathcal{M}$ ; note that  $M$  is Hermitian and  $M \succeq 0$  by construction. The notation  $M(\mathcal{O}, \mathcal{P})$  refers to the entry of  $M$  at the row and column corresponding to  $\mathcal{O}, \mathcal{P} \in \mathcal{M}$  respectively. We have  $M(\mathcal{O}, \mathcal{P}) = \langle \psi | \mathcal{O} \mathcal{P} | \psi \rangle$ , for all  $\mathcal{O}, \mathcal{P} \in \mathcal{M}$ , so that  $M$  captures all the 2-local Pauli statistics of  $|\psi\rangle$ . In particular the quantity  $\langle \psi | H | \psi \rangle$  is a linear function of the entries of  $M$  for a 2-local Hamiltonian  $H$ . If we let  $\mathcal{M}_k$  consist of all the  $k$ -local tensor products of Paulis, instead of just the 1-local ones, the corresponding moment matrix  $M_k$ , of size  $O(n^k)$  by  $O(n^k)$ , includes all the  $2k$ -local Pauli statistics. We may obtain SDP relaxations for QLH problems by constructing a relaxed  $M_k \succeq 0$  that satisfies linear constraints of the form  $\text{Tr}[AM_k] = b$ , that a true moment matrix would satisfy. This corresponds to the  $k$ th level of noncommutative Lasserre hierarchies introduced for quantum information [DLTW08, PNA10]. Our approach relaxes  $M_1$  and adds additional constraints enforcing positivity of 2-local marginals; the relaxation we obtain sits between the  $k = 1$  and  $k = 2$  levels of the noncommutative Lasserre hierarchy.

**SDP Relaxation.** We define a (relaxed) moment matrix  $M$ , which will track local statistics of the set of marginals  $\{\rho_{ij}\}$ . Let  $M$  be a symmetric,  $(3n + 1) \times (3n + 1)$  real matrix whose rows and columns correspond to operators in  $\mathcal{M}$ . Entries of  $M$  will correspond to coefficients of the marginal density matrices  $\{\rho_{ij}\}$  in the Pauli basis. We use the notation  $M(\sigma_i^k, \sigma_j^l)$  to refer to entries of  $M$  for  $i, j \in [n]$  and  $k, l \in [3]$ ; in addition we have a row and column of  $M$  indexed by  $\mathbb{I}$ . We set  $M(\sigma_i^k, \sigma_j^l) = \text{Tr}[\sigma^k \otimes \sigma^l \rho_{ij}]$  for  $(i, k), (j, l)$  in  $[n] \times [3]$ . In addition we set  $M(\mathbb{I}, \mathbb{I}) = 1$ , and  $M(\sigma_i^k, \mathbb{I}) = \text{Tr}[\sigma^k \otimes \mathbb{I} \rho_{ij}]$  for all  $(i, k) \in [n] \times [3]$  and  $j \in [n]$ . Note that this constraint forces consistent single-qubit marginals since

$$\text{Tr}_u[\rho_{iu}] = \text{Tr}_v[\rho_{iv}] \Leftrightarrow \text{Tr}[\sigma^l \otimes \mathbb{I} \rho_{iu}] = \text{Tr}[\sigma^l \otimes \mathbb{I} \rho_{iv}] \quad \forall l.$$

Since  $M$  contains all local information of  $\{\rho_{ij}\}$ , we can use  $M$  to evaluate the objective of our SDP. In this direction, we will define a weight matrix for each edge  $H_{ij} = w_{ij} \mathcal{O}_{ij} \otimes \mathbb{I}_{n \setminus \{i, j\}}$ , where  $\mathcal{O}_{ij} \in \mathbb{C}^{4 \times 4}$ , and  $w_{ij}$  is a scalar weight. We define the  $(3n + 1) \times (3n + 1)$  matrix  $C_{ij}$ , which contains the coefficients of  $\mathcal{O}_{ij}$  in the Pauli basis:

$$\begin{aligned} C_{ij}(\sigma_i^k, \sigma_j^l) &= C_{ij}(\sigma_j^l, \sigma_i^k) = \text{Tr}[\sigma^k \otimes \sigma^l \mathcal{O}_{ij}] / 8 \quad \forall k, l \in [3], \\ C_{ij}(\sigma_i^k, \mathbb{I}) &= C_{ij}(\mathbb{I}, \sigma_i^k) = \text{Tr}[\sigma^k \otimes \mathbb{I} \mathcal{O}_{ij}] / 8 \quad \forall k \in [3], \\ C_{ij}(\sigma_j^l, \mathbb{I}) &= C_{ij}(\mathbb{I}, \sigma_j^l) = \text{Tr}[\mathbb{I} \otimes \sigma^l \mathcal{O}_{ij}] / 8 \quad \forall l \in [3], \end{aligned} \tag{3}$$

and all other entries of  $C_{ij}$  are 0. To illustrate application of the matrix  $C_{ij}$ , suppose  $\mathcal{O}_{ij}$  and the marginal density matrix  $\rho_{ij}$  have Pauli decompositions:

$$\mathcal{O}_{ij} = \sum_{k, l=0}^3 \alpha_{kl} \sigma^k \otimes \sigma^l \quad \text{and} \quad \rho_{ij} = \frac{1}{4} \sum_{k, l=0}^3 \beta_{kl} \sigma^k \otimes \sigma^l.$$

Since, for  $k, l \geq 0$ ,  $(\sigma^k)^2 = \mathbb{I}$  and  $\text{Tr}[\sigma^k \sigma^l] = 0$  when  $k \neq l$ , the value we gain from edge  $ij$ , ignoring the weight  $w_{ij}$ , is written as:

$$\text{Tr}[\mathcal{O}_{ij} \rho_{ij}] = \alpha_{00} \beta_{00} + \sum_{\substack{k, l: \\ (k \neq 0) \vee (l \neq 0)}} \alpha_{kl} \beta_{kl} = \frac{\text{Tr}[\mathcal{O}_{ij}]}{4} + \text{Tr}[C_{ij} M]. \tag{4}$$

With these facts in hand, we may finally give the main SDP relaxation in this work:

**Problem 7.** *Given an instance of QLH (Problem 4) on  $n$  qubits with local terms  $\{H_{ij} = w_{ij} \mathcal{O}_{ij} \otimes \mathbb{I}_{[n] \setminus \{i,j\}}\}$ , let  $C_{ij}$  be defined according to Equation (3) for each  $ij \in E$ . Solve the following SDP:*

$$\max \sum_{ij \in E} w_{ij} \left( \frac{\text{Tr}[\mathcal{O}_{ij}]}{4} + \text{Tr}[C_{ij}M] \right) \quad (5)$$

$$\text{s.t.} \quad M(\mathbb{I}, \mathbb{I}) = 1, \quad (6)$$

$$M(\sigma_i^k, \sigma_i^k) = 1 \quad \forall i \in [n] \text{ and } k \in [3], \quad (7)$$

$$M(\sigma_i^k, \sigma_i^l) = 0 \quad \forall i \in [n] \text{ and } k \neq l \in [3], \quad (8)$$

$$M(\sigma_i^k, \sigma_j^l) = \text{Tr}[\sigma^k \otimes \sigma^l \rho_{ij}] \quad \forall ij \in E \text{ and } k, l \in [3], \quad (9)$$

$$M(\sigma_i^k, \mathbb{I}) = \text{Tr}[\sigma^k \otimes \mathbb{I} \rho_{ij}] \quad \forall ij \in E \text{ and } k \in [3], \quad (10)$$

$$M(\sigma_j^l, \mathbb{I}) = \text{Tr}[\mathbb{I} \otimes \sigma^l \rho_{ij}] \quad \forall ij \in E \text{ and } l \in [3], \quad (11)$$

$$\text{Tr}[\rho_{ij}] = 1 \quad \forall ij \in E, \quad (12)$$

$$\mathcal{H}(\mathbb{C}^{4 \times 4}) \ni \rho_{ij} \succeq 0 \quad \forall ij \in E, \quad (13)$$

$$\mathcal{S}(\mathbb{R}^{(3n+1) \times (3n+1)}) \ni M \succeq 0, \quad (14)$$

where  $\mathcal{S}(\cdot)$  and  $\mathcal{H}(\cdot)$  refer to the symmetric and Hermitian matrices, respectively.

**Theorem 8.** *The mathematical program of Problem 7 is an efficiently computable semidefinite program that provides an upper bound on  $\lambda_{\max}(\sum_{ij \in E} H_{ij})$ .*

*Proof.* Constraints (6)–(12) are linear equalities on the entries of PSD matrices  $M$  and  $\rho_{ij} \forall ij \in E$ , hence we do indeed have an SDP. Since there are polynomially many variables of polynomial size, the usual considerations show computational efficiency, i.e. the program can be solved to arbitrary additive precision in polynomial time [BV04].

A larger matrix  $X \succeq 0$ , consisting of  $M$  and the  $\rho_{ij}$  as its diagonal blocks may be used to put the SDP into a more standard form (e.g. [BV04], Section 4.6.2). Although the  $\rho_{ij}$  are complex, the SDP may be solved as a real SDP by appealing to the standard approach of tracking the real and imaginary parts separately and observing  $X \succeq 0$  if and only if

$$\begin{bmatrix} \text{Re}(X) & -\text{Im}(X) \\ \text{Im}(X) & \text{Re}(X) \end{bmatrix} \succeq 0.$$

Let  $|\psi\rangle$  be an eigenvector corresponding to  $\lambda_{\max}(\sum_{ij \in E} H_{ij})$ , and let  $\rho_{ij}^*, \forall ij \in E$ , be the 2-qubit marginal density matrices of  $\rho = |\psi\rangle \langle \psi|$ , so that Constraints (12) and (13) are satisfied for the  $\rho_{ij}^*$ . In addition consider the moment matrix  $M$  for  $|\psi\rangle$  with respect to  $\mathcal{M}$ , as described above. The matrix  $M$  satisfies Constraints (6), (7), (9)–(11), and (14) by the definition of a moment matrix, since

$$\langle \psi | \sigma_i^k \otimes \sigma_j^l \otimes \mathbb{I}_{[n] \setminus \{i,j\}} | \psi \rangle = \text{Tr}[\sigma^k \otimes \sigma^l \rho_{ij}^*], \text{ for } 0 \leq k, l \leq 3. \quad (15)$$

Constraint (8) is the only one that remains. Note that the real part of  $M$ ,  $M^* := \text{Re}(M) \succeq 0$  since  $M \succeq 0$ . By Equation (15), for any  $j \in [n]$  and  $k \neq l \in [3]$ ,  $M(\sigma_i^k, \sigma_i^l) = \pm i \text{Tr}[\sigma^m \otimes \mathbb{I} \rho_{ij}^*]$ , where  $m \in [3] \setminus \{k, l\}$ . The quantities in Equation (15) are real since tensor products of Pauli operators are Hermitian. This implies that  $M(\sigma_i^k, \sigma_i^l)$  for  $k \neq l \in [3]$  is imaginary and more generally that  $M^*$  and the  $\rho_{ij}^*$  satisfy all the constraints.

Consider the objective value for this solution,  $\sum_{ij \in E} w_{ij} (\text{Tr}[\mathcal{O}_{ij}]/4 + \text{Tr}[C_{ij}M^*]) =$

$$\sum_{ij \in E} w_{ij} \text{Tr}[\mathcal{O}_{ij} \rho_{ij}^*] = \text{Tr} \left[ \sum_{ij \in E} H_{ij} \rho \right] = \lambda_{max} \left( \sum_{ij \in E} H_{ij} \right),$$

where the first equality follows from Equation (4). It follows that the optimal solution to Problem 7 has value at least that of the optimal solution of QLH.  $\square$

## 2.4 Rounding Approach and Formal Statement of Results

**Overview.** In classical SDP-based rounding schemes, one typically seeks to randomly “round” unit vectors  $\mathbf{v}_i \in \mathbb{R}^d$  to scalars  $z_i \in \{\pm 1\}$  so that the expected value of  $z_i z_j$  approximates  $\mathbf{v}_i \cdot \mathbf{v}_j$ . The seminal hyperplane rounding scheme of Goemans and Williamson [GW95] achieves this by selecting a random unit vector  $\mathbf{r} \in \mathbb{R}^d$  and setting  $z_i = \mathbf{r} \cdot \mathbf{v}_i / |\mathbf{r} \cdot \mathbf{v}_i|$ .

Rounding solutions from SDP relaxations for QLH to product states generalizes this approach. Recall that a product state has the form  $|\psi\rangle = |\psi_1\rangle \otimes \dots \otimes |\psi_n\rangle$  where each  $|\psi_i\rangle \in \mathbb{C}^2$  is a local state on qubit  $i$ . We obtain a density matrix  $\rho = |\psi\rangle\langle\psi| = |\psi_1\rangle\langle\psi_1| \otimes \dots \otimes |\psi_n\rangle\langle\psi_n|$ , which is a tensor product of single-qubit density matrices  $\rho_i := |\psi_i\rangle\langle\psi_i|$ . Any such  $\rho_i$  may be expressed in the Pauli basis as  $\rho_i = \frac{1}{2}(\mathbb{I} + \theta_{i1}\sigma^1 + \theta_{i2}\sigma^2 + \theta_{i3}\sigma^3)$ , where  $\theta_{ik} = \text{Tr}[\sigma^k \rho_i] \in \mathbb{R}$  and  $\sum_{k \in [3]} \theta_{ik}^2 = 1$ . In particular, product states with  $\theta_{i1} = \theta_{i2} = 0$  and  $\theta_{i3}^2 = 1$  correspond precisely to the classical setting (see Appendix A for an explicit connection between the two). Product states exhibit no entanglement, and we may specify  $\theta_{ik}$  independently for each qubit  $i$ . However, instead of producing a single  $z_i^2 = 1$  as in the classical case, we must produce a unit vector  $\theta_i = [\theta_{i1}, \theta_{i2}, \theta_{i3}] \in \mathbb{R}^3$  for each  $i \in [n]$ . Briët, de Oliveira Filho, and Vallentin were the first to consider such generalizations of scalars to unit vectors, in the context of the Grothendieck problem [BdOFV10, BdOFV14], and their analysis has fueled recent approximation algorithms for QLH [GP19, HLP20].

The classical Goemans-Williamson rounding scheme obtains the unit vectors  $\mathbf{v}_i$  from a Cholesky decomposition of a PSD matrix  $V^T V = R \succeq 0$ . Taking the  $\mathbf{v}_i$  to be columns of  $V$  yields  $R_{i,j} = \mathbf{v}_i \cdot \mathbf{v}_j$ . Recent approximation algorithms [BGKT19, GP19, HLP20] for QLH have mimicked this approach, as do we. Let  $M^* \succeq 0$  be an optimal solution to Problem 7 (the  $\rho_{ij}^*$  are not necessary to describe the rounding algorithm). We find a Cholesky decomposition  $V^T V = M^*$ , and let  $\mathbf{v}_{ik} \in \mathbb{R}^d$  be the column of  $V$  associated with  $\sigma_i^k$  for  $i \in [n], k \in [3]$ ; we may assume  $d \leq 3n + 1$ . In addition we let  $\mathbf{v}_0$  be the column of  $V$  corresponding to  $\mathbb{I}$ . These are unit vectors as a consequence of Constraints (6) and (7).

We will employ the same rounding algorithm for both the general and strictly quadratic cases. While previous related works [BGKT19, GP19, HLP20] have in some cases had to rely on more sophisticated rounding schemes because they have been amenable to analysis, we are able to shed light on what is arguably the most natural generalization of the Goemans-Williamson approach. We draw  $\mathbf{r} \sim \mathcal{N}(0, \mathbb{I}_d)$ , i.e. a multivariate distribution over  $d$  independent and standard Gaussian variables. For each qubit, we obtain the desired vector  $\theta_i = [\theta_{i1}, \theta_{i2}, \theta_{i3}]$  as:

$$[\mathbf{v}_{i1} \cdot \mathbf{r}, \mathbf{v}_{i2} \cdot \mathbf{r}, \mathbf{v}_{i3} \cdot \mathbf{r}] / Q_i,$$

where  $Q_i := \sqrt{(\mathbf{v}_{i1} \cdot \mathbf{r})^2 + (\mathbf{v}_{i2} \cdot \mathbf{r})^2 + (\mathbf{v}_{i3} \cdot \mathbf{r})^2}$  is a normalization.

The classical Max Cut problem corresponds to a strictly quadratic Hamiltonian (see Appendix A for justification); however, classical Max 2-SAT and more general Max 2-CSP have 1-local terms (i.e., linear terms in  $\{\pm 1\}$  variables). In contrast, strictly quadratic instances of QLHP serve as a quantum generalization of Max 2-SAT and Max 2-CSP that have no 1-local terms. In order to

obtain effective classical approximations in the presence of 1-local terms, an additional vector  $\mathbf{v}_0$  is necessary, representing (scalar) identity. Generally, the vector  $\mathbf{v}_0$  is used in conjunction with more sophisticated rounding schemes (e.g. [LLZ02]) to obtain positive expectation from the 1-local terms. For the quantum case, relatively simple approaches suffice to get good approximations [HLP20]. Using the vector  $\mathbf{v}_0$  is not necessary for the strictly quadratic case, and including it does not affect its approximation.

**Rounding algorithm.** The rounding approach described above produces single-qubit density matrices:

$$\rho_i = \frac{1}{2} \left( \mathbb{I} + \frac{\mathbf{v}_{i1} \cdot \mathbf{r}}{Q_i} \sigma^1 + \frac{\mathbf{v}_{i2} \cdot \mathbf{r}}{Q_i} \sigma^2 + \frac{\mathbf{v}_{i3} \cdot \mathbf{r}}{Q_i} \sigma^3 \right).$$

Hence, on any 1-local term,  $\mathbb{E}[\text{Tr}[\sigma^k \rho_i]] = \mathbb{E}[\mathbf{v}_{ik} \cdot \mathbf{r}/Q_i] = 0$  since  $Q_i$  is an even function and  $\mathbf{v}_{ik} \cdot \mathbf{r}$  is an odd function in each entry of  $\mathbf{r}$ . Thus, in order to get a nontrivial approximation on 1-local terms, we will use the vector  $\mathbf{v}_0$  to globally flip the sign of the  $\theta_i$  vectors of all qubits, i.e.  $\mathbf{v}_{ik} \cdot \mathbf{r}/Q_i \rightarrow \text{sign}(\mathbf{v}_0 \cdot \mathbf{r})(\mathbf{v}_{ik} \cdot \mathbf{r}/Q_i)$ . Since  $\text{sign}(\mathbf{v}_0 \cdot \mathbf{r}) \in \{\pm 1\}$ , for quadratic objective terms this factor will cancel out, but for 1-local terms we will gain additional objective from the correlation of  $\mathbf{v}_0 \cdot \mathbf{r}$  and  $\mathbf{v}_{ik} \cdot \mathbf{r}$ . Formally, we can state the rounding algorithm, which applies to any instance of QLH, as follows.

**Algorithm 9.** *Hyperplane rounding for 2-Local Hamiltonian:*

1. Given some instance of Problem 4 formulate and solve the corresponding instance of Problem 7. Let  $M^*$  be the optimal moment matrix obtained from Problem 7.
2. Find the Cholesky decomposition of  $M^*$ , obtaining Cholesky vectors  $\mathbf{v}_0 \in \mathbb{R}^d$  and  $\{\mathbf{v}_{ik} \in \mathbb{R}^d\}$  such that  $M^*(\sigma_i^k, \sigma_j^l) = \mathbf{v}_{ik} \cdot \mathbf{v}_{jl}$  and  $M^*(\mathbb{I}, \sigma_i^k) = \mathbf{v}_0 \cdot \mathbf{v}_{ik}$  for  $i, j \in [n]$  and  $k, l \in [3]$ .
3. Let  $\mathbf{r}$  be a random vector with  $\mathbf{r} \sim \mathcal{N}(0, \mathbb{I}_d)$ .
4. For each qubit  $i$ , set  $Q_i = \sqrt{(\mathbf{v}_{i1} \cdot \mathbf{r})^2 + (\mathbf{v}_{i2} \cdot \mathbf{r})^2 + (\mathbf{v}_{i3} \cdot \mathbf{r})^2}$ , and set  $\theta_{ik} = \text{sign}(\mathbf{v}_0 \cdot \mathbf{r})(\mathbf{v}_{ik} \cdot \mathbf{r}/Q_i)$  for  $k \in [3]$ .
5. Output the (pure) state:

$$\rho = \bigotimes_{i=1}^n \frac{1}{2} (\mathbb{I} + \theta_{i1} \sigma^1 + \theta_{i2} \sigma^2 + \theta_{i3} \sigma^3).$$

We will give the following approximation guarantees for QLHP:

**Theorem 10.** *Fix  $k \in \{1, 2, 3\}$ . Suppose we are given an instance of QLHP (Problem 6),  $\{H_e\}$  where  $H_e = w_e P_e \otimes \mathbb{I}_{[n] \setminus \{e_1, e_2\}}$  for  $w_e \geq 0$  and  $P_e$  a projector of rank at least  $k$ , for all  $e \in E$ . Let  $M^*$  be the optimal moment matrix for the corresponding SDP relaxation, Problem 7, and let  $\rho$  be the random output of Algorithm 9. Then,*

$$\mathbb{E} \left[ \text{Tr} \left[ \sum_{e \in E} H_e \rho \right] \right] \geq \alpha(k) \left( \sum_{e \in E} w_e \left( \frac{\text{rank}(P_e)}{4} + \text{Tr}[C_e M^*] \right) \right) \geq \alpha(k) \lambda_{\max} \left( \sum_{e \in E} H_e \right),$$

where

$$\alpha(k) = \begin{cases} 2/\pi - 1/4 \approx 0.387 & \text{if } k = 1 \\ 16/(9\pi) \approx 0.565 & \text{if } k = 2 \\ 3/8 + 11/(9\pi) \approx 0.764 & \text{if } k = 3. \end{cases}$$

**Theorem 11.** *If, in addition to the assumptions of Theorem 10, the  $P_e$  are strictly quadratic projectors, then the random output of Algorithm 9 satisfies:*

$$\mathbb{E} \left[ \text{Tr} \left[ \sum_{ij \in E} H_{ij} \rho \right] \right] \geq \alpha(k) \lambda_{\max} \left( \sum_{ij \in E} H_{ij} \right),$$

where

$$\alpha(k) = \begin{cases} 22/(15\pi) \approx 0.467 & \text{if } k = 1 \\ 1/3 + 24/(25\pi) \approx 0.639 & \text{if } k = 2 \\ 1/2 + 388/(405\pi) \approx 0.804 & \text{if } k = 3. \end{cases}$$

The above results are rigorous, but non-optimal. The quadratic analysis depends crucially on an expansion of a particular expectation in Hermite polynomials. One can consider a higher order Hermite series to more accurately capture the expectation and achieve a better approximation factor. We have such results, but opt to not include them in the interest of the reader. Higher orders bring increased tedium, and our technique should be clear enough at the end of the paper that an interested reader could do the higher order calculation.

One can ask, why not include a high enough order that the result becomes essentially optimal? The issue is that polynomial expansions often converge slowly in the presence of discontinuities [PMS19]. Indeed, computationally we have determined that to get essentially optimal results one would need to go to high enough order that the polynomial expansion would become intractable. One can determine the optimal approximation factor by using a high order expansion and numerically optimizing or simply by randomly sampling over some “net” of the parameter space. As stated in the introduction, our observed approximation factors under these approaches are:

**Conjecture 12.** *Algorithm 9 constitutes a randomized  $\beta(k)$  approximation algorithm for strictly quadratic projectors where*

$$\beta(k) = \begin{cases} 0.498 & \text{if } k = 1 \\ 0.653 & \text{if } k = 2 \\ 0.821 & \text{if } k = 3. \end{cases} \quad (16)$$

Proving an approximation factor as large as the observed performance of our algorithm (say with techniques from algebraic geometry) is the subject of future work.

## 2.5 Analysis Overview

We present an overview of our analysis for the strictly quadratic case, which will also carry over to the general case with additional bookkeeping and bounding for the 1-local terms. Suppose we are given an instance of QLHP (Problem 6) on which we execute Algorithm 9 to produce a random solution  $\rho$ . For  $i, j \in [n]$ , the 2-qubit marginals of  $\rho$  are

$$\rho_{ij} = \frac{1}{4} (\mathbb{I} + \theta_{i1} \sigma^1 + \theta_{i2} \sigma^2 + \theta_{i3} \sigma^3) \otimes (\mathbb{I} + \theta_{j1} \sigma^1 + \theta_{j2} \sigma^2 + \theta_{j3} \sigma^3), \quad (17)$$

and the objective value of  $\rho$  is  $\sum_{e \in E} \text{Tr}[H_e \rho] = \sum_{e \in E} w_e APX_e$ , where  $APX_e := \text{Tr}[P_e \rho_{e_1 e_2}]$  is the unweighted contribution to the objective value from edge  $e$ . Let  $M^*$  and  $\rho_{e_1 e_2}^*$  for  $e \in E$  be the SDP solution obtained by Algorithm 9, and let  $SDP_e = \text{Tr}[P_e \rho_{e_1 e_2}^*]$  be the unweighted contribution



to the SDP objective value from edge  $e$ . The approximation ratio, which we seek to bound from below, is consequently:

$$\alpha = \mathbb{E} \left[ \frac{\sum_{e \in E} w_e APX_e}{\sum_{e \in E} w_e SDP_e} \right] = \frac{\sum_{e \in E} w_e \mathbb{E}[APX_e]}{\sum_{e \in E} w_e SDP_e}.$$

Observe that  $APX_e \geq 0$  and  $SDP_e \geq 0$  since  $P_e$ ,  $\rho_{e_1 e_2}$ , and  $\rho_{e_1 e_2}^*$  are all PSD. Since all the terms in the denominator are nonnegative, it follows from an elementary argument that

$$\alpha \geq \frac{\sum_{e \in E} w_e \mathbb{E}[APX_e]}{\sum_{e \in E} w_e SDP_e} \geq \min_{e \in E} \frac{\mathbb{E}[APX_e]}{SDP_e}.$$

Thus it suffices to bound the approximation ratio for the worst case occurring on a single edge.

**Bounding a worst-case edge.** We now focus our attention on a single edge  $e = 12$  on qubits 1, 2. We collect the vectors  $\mathbf{v}_{ik}$ , obtained from a Cholesky decomposition of the SDP solution  $M^*$ , into matrices  $V_i = [\mathbf{v}_{i1}, \mathbf{v}_{i2}, \mathbf{v}_{i3}] \in \mathbb{R}^{d \times 3}$ , for  $i = 1, 2$ . We define an objective matrix  $C \in \mathbb{R}^{3 \times 3}$ , containing scaled 2-local Pauli-basis coefficients of  $P_{12}$ , in the vein of Equation (3):  $C(\sigma_1^k, \sigma_2^l) := \text{Tr}[\sigma_1^k \otimes \sigma_2^l P_{12}]$ , for  $k, l \in [3]$  (note that  $C$  is not symmetric). With these definitions in hand, observe that  $V_i^T V_i = \mathbb{I}_3$ , by the SDP constraints (7) and (8), and that  $4\text{Tr}[C_{12} M^*] = \text{Tr}[V_1 C V_2^T]$ . The hyperplane rounding produces unit vectors  $\theta_i^T = [\theta_{i1}, \theta_{i2}, \theta_{i3}] = \mathbf{r}^T V_i / \|V_i^T \mathbf{r}\|$ . In terms of these variables, we have:

$$SDP_e = \text{Tr}[P_{12} \rho_{12}^*] = \frac{1}{4} (\text{rank}(P_{12}) + \text{Tr}[V_1 C V_2^T]), \text{ and}$$

$$\mathbb{E}[APX_e] = \mathbb{E}[\text{Tr}[P_{12} \rho_{12}]] = \frac{1}{4} (\text{rank}(P_{12}) + \mathbb{E}[\theta_1^T C \theta_2]) = \frac{1}{4} \left( \text{rank}(P_{12}) + \mathbb{E}_{\mathbf{r}} \left[ \frac{\mathbf{r}^T V_1 C V_2^T \mathbf{r}}{\|V_1^T \mathbf{r}\| \|V_2^T \mathbf{r}\|} \right] \right),$$

by Equation (4) and because  $\text{Tr}[P_{12}] = \text{rank}(P_{12})$ , since  $P_{12}$  is a projector. Thus, setting  $k = \text{rank}(P_{12})$ , the quantity we seek to bound is

$$\alpha \geq \min_{V_1, V_2, C} \frac{k + \mathbb{E}_{\mathbf{r}} \left[ \frac{\mathbf{r}^T V_1 C V_2^T \mathbf{r}}{\|V_1^T \mathbf{r}\| \|V_2^T \mathbf{r}\|} \right]}{k + \text{Tr}[V_1 C V_2^T]}.$$

The bulk of our analysis lies in (i) simplifying the above to reduce the number of parameters in the minimization and expectation (Section 4), and (ii) deriving analytical bounds on the expectation (Section 5).

**Simplifying the Gaussian expectation.** The first simplification follows from observing that  $V_i^T \mathbf{r} \in \mathbb{R}^3$  are multivariate Gaussians for  $i = 1, 2$  since they are linear combinations of Gaussians,  $\mathbf{r} \sim \mathcal{N}(0, \mathbb{I})$ . If we let  $\mathbf{z}^T = [z_1, z_2, z_3] = \mathbf{r}^T V_1$  and  $(\mathbf{z}')^T = [z'_1, z'_2, z'_3] = \mathbf{r}^T V_2$ , then  $[\mathbf{z}, \mathbf{z}'] \sim \mathcal{N}(0, \Sigma)$ , where

$$\Sigma = \begin{bmatrix} \mathbb{I} & V_1^T V_2 \\ V_2^T V_1 & \mathbb{I} \end{bmatrix} \in \mathbb{R}^{6 \times 6}.$$

The Gaussians  $z_i$  are mutually independent as well as the  $z'_i$ , and the covariance between  $\mathbf{z}$  and  $\mathbf{z}'$  is given by  $M = V_1^T V_2 \in \mathbb{R}^{3 \times 3}$ . Our bound now depends on a constant number of parameters, the 18 entries of  $C$  and  $M$ :

$$\alpha \geq \min_{M, C} \frac{k + \mathbb{E}_{\mathbf{z}, \mathbf{z}'} \left[ \frac{\mathbf{z}^T C \mathbf{z}'}{\|\mathbf{z}\| \|\mathbf{z}'\|} \right]}{k + \text{Tr}[C^T M]}. \quad (18)$$

For classical hyperplane rounding algorithms,  $C$  and  $M$  simply reduce to scalars, and one may resort to a numerical argument to furnish the desired bound. However, in the case of QLH above, numerical bounds exhibit poor precision or convergence due to the number of parameters. Thus we press on, and our next observation is that only the singular values of  $M$  matter for the analysis. The above arguments are detailed in Lemma 15 in Section 4.2, which also shows that we may assume:

$$C = \begin{bmatrix} p & 0 & 0 \\ 0 & q & 0 \\ 0 & 0 & r \end{bmatrix} \text{ and } M = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, \quad (19)$$

where  $a, b, c$  are the singular values of  $V_1^T V_2$ . This reduction to 6 parameters puts analysis of  $\alpha$  within reach. The special case when  $a = b = c$  turns out to be equivalent to the recently studied quantum analog of Max Cut related to the quantum Heisenberg model [GP19, AGM20]. For this case, a representation of the expectation,

$$\mathbb{E}_{\mathbf{z}, \mathbf{z}'} \left[ \frac{\mathbf{z}^T C \mathbf{z}'}{\|\mathbf{z}\| \|\mathbf{z}'\|} \right] \quad (20)$$

as a hypergeometric function follows from work of Briët, de Oliveira Filho, and Vallentin (the expectation ends up being equivalent to the one in Lemma 2.1 from [BdOFV14], when  $\mathbf{u} \cdot \mathbf{v}$  in the lemma equals  $a = b = c$ ). To the best of our knowledge, no elementary representation is known when  $a, b, c$  may be distinct. We appeal to Hermite analysis to express the expectation (20) as a polynomial that we are subsequently able to bound; this is carried out in Lemmas 17-18/Section 5.

**Introducing constraints from positivity.** The matrices  $C$  and  $M$  from (19) are related to the quadratic Pauli-basis coefficients of  $P_{12}$  and  $\rho_{12}^*$ , respectively. The other ingredient of our analysis of the bound (18) is restricting  $C$  and  $M$  based on the facts that  $P_{12} \succeq 0$  and  $\rho_{12}^* \succeq 0$ , which is undertaken in Section 4.1 and Lemma 14. The bound we obtain is

$$\alpha \geq \min_{\substack{[a,b,c] \in \mathcal{S} \\ [p,q,r] \in \mathcal{P}_k}} \frac{k + \mathbb{E}_{\mathbf{z}, \mathbf{z}'} \left[ \frac{pz_1 z'_1 + qz_2 z'_2 + rz_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right]}{k + ap + bq + cr}, \quad (21)$$

where  $\mathcal{S}$  and  $\mathcal{P}_k$  are specific polytopes ( $\mathcal{S}$  is a simplex as is  $\mathcal{P}_k$  for  $k \neq 2$ ) derived from the positivity of  $P_{12}$  and  $\rho_{12}^*$ . Finally, Lemmas 17 and 18 in Section 5 derive the bounds in the main theorems 10 and 11, respectively.

**Analysis roadmap.** First, we describe the notations which will be used throughout the paper in Section 3. Following this, in Section 4, we will describe our reduction of the expectation of an edge to a standard form (Section 4.2), as well as the tools required for the proof of these lemmas (primarily Lemma 14 and singular value decomposition). In Section 5 we will cover the main technical lemmas in this work, Lemma 17 and Lemma 18. These are simply lower bounds on a constrained minimization of a particular expectation, e.g. Equation (21). Proving these lemmas is the brunt of the technical work in this paper, and will require tools from orthogonal polynomials (Section 5.1), optimization (Section 5.2), and some elementary calculus (Section 5.3).

### 3 Notations

We will adopt several mostly standard notations throughout the paper. Note that  $[n] := \{1, 2, \dots, n\}$  takes its standard definition, as well as the double factorial:  $a!! := a(a-2)\dots 4 \cdot 2$  if  $a$  is even and

$a!! := a(a-2)(a-4)\dots\cdot 3\cdot 1$  if  $a$  is odd. By convention,  $(-1)!! = 1 = 0!!$ . The notation  $\mathbf{z} \sim \mathcal{N}(\mu, \Sigma)$  will be used to indicate that  $\mathbf{z}$  is a multivariate normal distribution with mean  $\mu$  and covariance matrix  $\Sigma$ . Consistent with the notation from [O'D14], we define  $L^2(\mathbb{R}^n, \gamma)$  as the set of multivariate functions which are square integrable under the Gaussian measure. More formally  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is an element of  $L^2(\mathbb{R}^n, \gamma)$  if:

$$\int_{\mathbb{R}^n} |f(\mathbf{x})|^2 d\gamma(\mathbf{x}) := \int_{\mathbb{R}^n} |f(\mathbf{x})|^2 \frac{e^{-\|\mathbf{x}\|^2/2}}{(2\pi)^{n/2}} d\mathbf{x} < \infty. \quad (22)$$

We define an inner product on this space as:

$$\langle f, g \rangle = \int_{\mathbb{R}^n} f(\mathbf{x})g(\mathbf{x})d\gamma(\mathbf{x}). \quad (23)$$

(Not to be confused with ‘‘Bra-ket’’ notation, meaning will always be obvious from context)

We will make use of the Cholesky decomposition [Vaz13]. Let  $M$  be a real, symmetric,  $n \times n$  PSD matrix. The Cholesky decomposition guarantees the existence of efficiently-computable vectors  $\{\mathbf{v}_i\}_{i=1}^n$ , each of which has dimension at most  $n$ , such that  $M_{ij} = \mathbf{v}_i^T \mathbf{v}_j$ . We will refer to the  $\mathbf{v}_i$  simply as the Cholesky vectors of  $M$ .

The Gamma and Hypergeometric functions will take their standard definitions:

$$\Gamma(z) := \int_0^\infty x^{z-1} e^{-x} dx, \quad \text{and} \quad {}_2F_1 \left[ \begin{matrix} a, b \\ c \end{matrix}; z \right] = \sum_{n=0}^\infty \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!}, \quad (24)$$

where the rising factorial  $(x)_n = 1$  if  $n = 0$ , or  $(x)_n = x(x+1)\dots(x+n-1)$  if  $n > 0$ .

We will need to define a convex hull. Given some set  $S = \{\mathbf{v}_1, \dots, \mathbf{v}_m\}$  of points in  $\mathbb{R}^p$ , we denote the convex hull of  $S$  as  $\text{conv}(S)$  or  $\text{conv } S$ :

$$\text{conv}(S) = \{\lambda_1 \mathbf{v}_1 + \dots + \lambda_m \mathbf{v}_m : \sum_i \lambda_i = 1, \lambda_i \geq 0 \forall i\}. \quad (25)$$

We will reserve  $\mathcal{S}$ ,  $-\mathcal{S}$  and  $\mathcal{T}$  for specific convex hulls:

$$\begin{aligned} \mathcal{S} &= \text{conv} \{[-1, -1, -1], [-1, 1, 1], [1, -1, 1], [1, 1, -1]\} \\ -\mathcal{S} &= \text{conv} \{[1, 1, 1], [1, -1, -1], [-1, 1, -1], [-1, -1, 1]\} \\ \mathcal{T} &= \text{conv} \{[2, 0, 0], [0, 2, 0], [0, 0, 2], [-2, 0, 0], [0, -2, 0], [0, 0, -2]\} \end{aligned} \quad (26)$$

We reserve  $\Sigma' : \mathbb{R}^n \rightarrow \mathbb{R}^{2n \times 2n}$  for the matrix-valued function:

$$\Sigma'(a_1, a_2, \dots, a_n) = \begin{bmatrix} & & & & a_1 & 0 & \dots & 0 \\ & & & & 0 & a_2 & & \vdots \\ & & & & \vdots & & \ddots & 0 \\ & & & & 0 & \dots & 0 & a_n \\ a_1 & 0 & \dots & 0 & & & & \\ 0 & a_2 & & \vdots & & & & \\ \vdots & & \ddots & 0 & & & & \\ 0 & \dots & 0 & a_n & & & & \end{bmatrix} \mathbb{I}_n. \quad (27)$$

The vector-valued function  $\text{diag} : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^n$  has its standard definition:  $\text{diag}(M) = [M_{11}, M_{22}, \dots, M_{nn}]$ .

## 4 Standard Form for the Expectation of an Edge

**Overview.** Recall we will use linearity of expectation to reduce the expected approximation factor for the algorithm as a whole to the expected approximation factor for the “worst-case” edge. The next step is to reduce the number of parameters we need to consider by converting a generic edge to a “standard” form. Roughly speaking, this is accomplished by applying a singular value decomposition to the moment matrix, followed by the application of some well-known facts concerning the structure of 2-qubit density matrices and projectors. In this section we will first give formal statements for the properties of 2-qubit states we will need (Section 4.1), then integrate these facts with our SVD argument to provide the main lemmas for this reduction (Section 4.2).

### 4.1 Moment Matrix Description of Two Qubit Quantum States

Moment matrices for 2-qubit states are well studied objects [HH96, KŽ01, SHH11, JPJR14, Gam16] (see [Gam16] for an extensive list of references), and the lemmas in this section are all presented in some form in these previous works. It is a well-known fact that tensor product of Pauli matrices form a complete basis for Hermitian operators. Hence, any Hermitian matrix is completely described by its coefficients in this basis, and we can associate this Hermitian matrix with some real matrix containing the coefficients. As an example let us consider a state,  $\rho$ , or a projector  $P$ , on two qubits. Let them have Pauli decompositions:

$$\rho = \frac{1}{4} \sum_{l,m=0}^3 \Gamma_{lm} \sigma^l \otimes \sigma^m \quad \text{or} \quad P = \frac{1}{4} \sum_{l,m=0}^3 \Gamma_{lm} \sigma^l \otimes \sigma^m. \quad (28)$$

We will think of  $\Gamma_{lm}$  as comprising a real matrix, which gives a complete description of the corresponding quantum state/projector.

$$\Gamma = \begin{bmatrix} 1 & \Gamma_{01} & \Gamma_{02} & \Gamma_{03} \\ \Gamma_{10} & \Gamma_{11} & \Gamma_{12} & \Gamma_{13} \\ \Gamma_{20} & \Gamma_{21} & \Gamma_{22} & \Gamma_{23} \\ \Gamma_{30} & \Gamma_{31} & \Gamma_{32} & \Gamma_{33} \end{bmatrix} = \begin{bmatrix} 1 & \mathbf{v}^T \\ \mathbf{u} & R \end{bmatrix}. \quad (29)$$

By the above we mean that the matrix  $R$  corresponds to the submatrix of  $\Gamma$  with indices in  $\{1, 2, 3\}$ ,  $\mathbf{u} = [\Gamma_{10}, \Gamma_{20}, \Gamma_{30}]^T$  and  $\mathbf{v} = [\Gamma_{01}, \Gamma_{02}, \Gamma_{03}]^T$ . We will refer to the vectors  $\mathbf{u}$  and  $\mathbf{v}$  as the 1-local parts and  $R$  as the quadratic part.

On several occasions in this work, we will be given an arbitrary  $4 \times 4$  real matrix  $\Gamma$  or a  $3 \times 3$  matrix  $R$ . We will say  $\Gamma$  (resp.  $R$ ) is a *valid 2-moment* ( resp. *valid quadratic 2-moment*) if it corresponds to a physical density matrix or projector as in Equation (28).

**Definition 13.** 1. Given some density matrix  $\rho$  or projector  $P$  as in Equation (28) we will say that  $\Gamma$  (resp.  $R$ ) as in Equation (29) is its 2-moment (resp. quadratic 2-moment).

2. Similarly if  $\Gamma$  (resp.  $R$ ) is a given  $4 \times 4$  (resp.  $3 \times 3$ ) real matrix, we say  $\Gamma$  (resp.  $R$ ) is a valid 2-moment (resp. valid quadratic 2-moment) for a density matrix or projector if it is the 2-moment (resp. quadratic 2-moment) of a (physical) density matrix or projector.

This notation will be important because moments are more restricted than an arbitrary  $3 \times 3$  or  $4 \times 4$  matrix, and these restrictions will imply simplifications which will be crucial in the analysis.

**Lemma 14.** Let  $P = 1/4 \sum_{l,m=0}^3 \Gamma_{lm} \sigma^l \otimes \sigma^m$  be some 2-qubit projector of rank  $k$ , and let  $\rho = 1/4 \sum_{l,m=0}^3 \Delta_{lm} \sigma^l \otimes \sigma^m$  be a density matrix on 2 qubits. Let  $G$  be the quadratic 2-moment for  $P$ , and let  $D$  be the quadratic 2-moment for  $\rho$  with  $D = L \Sigma N^T$  for  $\Sigma$  diagonal and  $L, N \in SO(3)$ .

1. ([HH96], equation 8) Let  $P, Q \in SO(3)$ . Then,  $G' = PGQ^T$  (resp.  $D' = PDQ^T$ ) is a valid quadratic 2-moment for state  $U_1 \otimes U_2 \rho U_1^\dagger \otimes U_2^\dagger$  (resp. projector  $U_1 \otimes U_2 P U_1^\dagger \otimes U_2^\dagger$ ) for some local unitaries  $U_1, U_2$ .
2. ([HH96], Proposition 1) The diagonal elements of  $\Sigma$  reside in  $\mathcal{S}$  (recall  $\mathcal{S}$  was defined in Equation (26)).
- 3.

$$\text{If } \begin{cases} k = 1 \\ k = 2 \\ k = 3 \end{cases} \text{ then } [\Gamma_{11}, \Gamma_{22}, \Gamma_{33}] \in \begin{cases} \mathcal{S} \\ \mathcal{T} \\ -\mathcal{S} \end{cases}, \quad (30)$$

where  $\mathcal{S}, \mathcal{T}$  and  $-\mathcal{S}$  are defined in Equation (26).

4. Further, it holds that

$$-k \leq 4\text{Tr}[P\rho] - k \leq 4 - k, \quad (31)$$

- 5.

$$-k \leq \Sigma_{11}\Gamma_{11} + \Sigma_{22}\Gamma_{22} + \Sigma_{33}\Gamma_{33} \leq 4 - k, \quad (32)$$

- 6.

$$\|[\Gamma_{10}, \Gamma_{20}, \Gamma_{30}]\|, \|[\Gamma_{01}, \Gamma_{02}, \Gamma_{03}]\| \leq \begin{cases} 1 & \text{if } k = 1 \text{ or } 3 \\ 2 & \text{if } k = 2 \end{cases}, \quad (33)$$

7. and

$$\|[\Delta_{10}, \Delta_{20}, \Delta_{30}]\|, \|[\Delta_{01}, \Delta_{02}, \Delta_{03}]\| \leq 1. \quad (34)$$

*Proof.* See Appendix, page 53. □

## 4.2 SVD Argument

**Lemma 15** (Quadratic Part of Expectation). Let  $V_1 = [\mathbf{v}_{11}, \mathbf{v}_{12}, \mathbf{v}_{13}] \in \mathbb{R}^{d \times 3}$  and  $V_2 = [\mathbf{v}_{21}, \mathbf{v}_{22}, \mathbf{v}_{23}] \in \mathbb{R}^{d \times 3}$  be real matrices with normalized columns ( $\|\mathbf{v}_{ij}\|^2 = 1$ ) such that  $V_1^T V_2$  is a valid quadratic 2-moment for some density matrix. Let  $C$  be a quadratic 2-moment for some projector of rank  $k$ , and let  $\mathbf{r} \sim \mathcal{N}(0, \mathbb{I})$  be the same size as the vectors  $\mathbf{v}_{ij}$ . Then,

$$\mathbb{E}_{\mathbf{r}} \left[ \frac{\mathbf{r}^T V_1 C V_2^T \mathbf{r}}{\|V_1^T \mathbf{r}\| \|V_2^T \mathbf{r}\|} \right] = \mathbb{E} \left[ \frac{pz_1 z'_1 + qz_2 z'_2 + rz_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right] \quad (35)$$

$$\text{and } \text{Tr}(V_1 C V_2^T) = ap + bq + cr. \quad (36)$$

with  $[z_1, z_2, z_3, z'_1, z'_2, z'_3] \sim \mathcal{N}(0, \Sigma'(a, b, c))$  for some constants  $(a, b, c, p, q, r)$ . Further, we can

assume without loss of generality that  $[a, b, c] \in \mathcal{S}$  and  $[p, q, r] \in \mathcal{P}_k$  where

$$\mathcal{P}_k = \begin{cases} \mathcal{S} & \text{if } k = 1 \\ \mathcal{T} & \text{if } k = 2 \\ -\mathcal{S} & \text{if } k = 3 \end{cases}. \quad (37)$$

*Proof.* Take a singular value decomposition of  $V_1^T V_2 = L \Sigma N^T$ . Let us assume that  $\det(L) = 1 = \det(N)$  by possibly absorbing signs into

$$\Sigma = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}. \quad (38)$$

Since  $L$  and  $N$  are special orthogonal matrices we can re-write Equation (35) (resp. Equation (36)) as:

$$\mathbb{E}_{\mathbf{r}} \left[ \frac{\mathbf{r}^T V_1 C V_2^T \mathbf{r}}{\|V_1^T \mathbf{r}\| \|V_2^T \mathbf{r}\|} \right] = \mathbb{E}_{\mathbf{r}} \left[ \frac{\mathbf{r}^T (V_1 L) (L^T C N) (N^T V_2^T) \mathbf{r}}{\|L^T V_1^T \mathbf{r}\| \|N^T V_2^T \mathbf{r}\|} \right], \quad (39)$$

$$\text{Tr} [V_1 P V_2^T] = \text{Tr} [(V_1 L) (L^T C N) (N^T V_2^T)]. \quad (40)$$

Setting

$$W_1 = V_1 L, \quad W_2 = V_2 N, \quad \text{and} \quad C' = L^T C N, \quad (41)$$

we can re-write Equation (39) (resp. Equation (40)) as:

$$= \mathbb{E}_{\mathbf{r}} \left[ \frac{\mathbf{r}^T W_1 C' W_2^T \mathbf{r}}{\|W_1^T \mathbf{r}\| \|W_2^T \mathbf{r}\|} \right] \quad (42)$$

$$= \text{Tr} [W_1 C' W_2^T] \quad (43)$$

Defining the random variables  $[z_1, z_2, z_3]^T := W_1^T \mathbf{r}$  and  $[z'_1, z'_2, z'_3]^T := W_2^T \mathbf{r}$ , we can re-write Equation (42) (resp. Equation (43)) as:

$$= \mathbb{E} \left[ \frac{[z_1, z_2, z_3] C' [z'_1, z'_2, z'_3]^T}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right], \quad (44)$$

$$= \text{Tr} [W_1 C' W_2^T]. \quad (45)$$

Since the variable  $\mathbf{r}$  is a multivariate Gaussian distribution, linear combinations of its components are also distributed according to the multivariate Gaussian. Since each of  $(z_1, z_2, z_3, z'_1, z'_2, z'_3)$  is the sum of normal random variables with zero mean, they must each have zero mean. Observe that, by construction,  $W_1^T W_2 = \Sigma$ . So, if we let  $\mathbf{w}_1^1$  be the first column of  $W_1$  and  $\mathbf{w}_1^2$  be the first column of  $W_2$  then

$$\begin{aligned} \mathbb{E}[z_1 z'_1] &= \mathbb{E}[(\mathbf{r} \cdot \mathbf{w}_1^1)(\mathbf{r} \cdot \mathbf{w}_1^2)] = \mathbb{E}[\mathbf{r}^T \mathbf{w}_1^1 (\mathbf{w}_1^2)^T \mathbf{r}] = \mathbb{E}[\text{Tr}(\mathbf{r}^T \mathbf{w}_1^1 (\mathbf{w}_1^2)^T \mathbf{r})] = \mathbb{E}[\text{Tr}(\mathbf{r} \mathbf{r}^T \mathbf{w}_1^1 (\mathbf{w}_1^2)^T)] \\ &= \text{Tr}[\mathbb{E}(\mathbf{r} \mathbf{r}^T) \mathbf{w}_1^1 (\mathbf{w}_1^2)^T] = a. \end{aligned}$$

Similar calculations show that the covariance of these random variables are equal to the overlap of the corresponding vectors. Hence,  $[z_1, z_2, z_3, z'_1, z'_2, z'_3] \sim \mathcal{N}(\mathbf{0}, \Sigma'(a, b, c))$  with  $\Sigma'$  defined as in Equation (27).

Observe that

$$\mathbb{E} \left[ \frac{z_1 C'_{1,2} z'_2}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right] = 0,$$

since it is the expectation of an odd function with respect to  $z_1$ , and similarly for the other off-diagonal elements. Let  $(p, q, r)$  be the diagonal elements of  $C'$ . Then we have shown that Equation (44) is equal to

$$\mathbb{E} \left[ \frac{pz_1 z'_1 + qz_2 z'_2 + rz_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right].$$

Additionally, it is easy to see  $\text{Tr} [W_1 C' W_2^T] = ap + bq + cr$  by taking the trace starting with the columns of  $W_1$ .

To complete the lemma, we need to establish that we can restrict  $(a, b, c, p, q, r)$  as described in the statement.  $[p, q, r]$  are the diagonal elements of a valid 2-moment for a projector, by Lemma 14, since we obtained  $C'$  by multiplying it on the left and right by special orthogonal matrices. Hence, Equation (30) establishes  $[p, q, r] \in \mathcal{P}_k$ . Since  $\Sigma$  is obtained from  $V_1^T V_2$  by multiplying on the right and left by special orthogonal matrices, Lemma 14 item 2 implies  $\text{diag}(\Sigma) \in \mathcal{S}$ .  $\square$

**Lemma 16** (Linear part of Expectation). *Let  $V = [\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3] \in \mathbb{R}^{d \times 3}$ ,  $\mathbf{v} \in \mathbb{R}^d$  with  $VV^T = \mathbb{I}$  and  $\mathbf{v}^T \mathbf{v} = 1$ . Further, let  $\mathbf{p} \in \mathbb{R}^3$ ,  $a = \|V^T \mathbf{v}\|$ ,  $\mathbf{r} \sim \mathcal{N}(\mathbf{0}, \mathbb{I}_d)$ . Let  $U \in SO(3)$  satisfy  $UV^T \mathbf{v} = [a, 0, 0]^T$ , and define  $\tilde{\mathbf{p}}_1 = (U\mathbf{p})_1$ . Then,*

$$\mathbb{E}_{\mathbf{r}} \left[ \frac{(\mathbf{r}^T V \mathbf{p})(\mathbf{v}^T \mathbf{r})}{\|\mathbf{r}^T V\| \cdot |\mathbf{r}^T \mathbf{v}|} \right] = \mathbb{E}_{X, Y, Z, S} \left[ \frac{\tilde{\mathbf{p}}_1 X S}{\sqrt{X^2 + Y^2 + Z^2} |S|} \right], \quad (46)$$

where  $[S, X, Y, Z] \sim \mathcal{N}(\mathbf{0}, \Sigma)$  with

$$\Sigma = \begin{bmatrix} 1 & a & 0 & 0 \\ a & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Additionally,  $\text{Tr}[V \mathbf{p} \mathbf{v}^T] = a \tilde{\mathbf{p}}_1$ .

*Proof.* Define  $\tilde{V} = VU^T$ . Then,

$$\mathbb{E}_{\mathbf{r}} \left[ \frac{(\mathbf{r}^T V \mathbf{p})(\mathbf{v}^T \mathbf{r})}{\|\mathbf{r}^T V\| \cdot |\mathbf{r}^T \mathbf{v}|} \right] = \mathbb{E}_{\mathbf{r}} \left[ \frac{(\mathbf{r}^T \tilde{V} \tilde{\mathbf{p}})(\mathbf{v}^T \mathbf{r})}{\|\mathbf{r}^T \tilde{V}\| \cdot |\mathbf{r}^T \mathbf{v}|} \right]. \quad (47)$$

Let  $\tilde{V} = [\mathbf{v}_X, \mathbf{v}_Y, \mathbf{v}_Z]$ , and define  $S = \mathbf{v}^T \mathbf{r}$ ,  $X = \mathbf{v}_X^T \mathbf{r}$ ,  $Y = \mathbf{v}_Y^T \mathbf{r}$  and  $Z = \mathbf{v}_Z^T \mathbf{r}$ . Equation (47) can then be written as:

$$= \mathbb{E} \left[ \frac{[X, Y, Z] \tilde{\mathbf{p}} S}{\sqrt{X^2 + Y^2 + Z^2} |S|} \right],$$

with  $\tilde{\mathbf{p}} = U\mathbf{p}$ . Equation (46) follows with the observation that

$$\mathbb{E} \left[ \frac{YS}{\sqrt{X^2 + Y^2 + Z^2} |S|} \right] = 0 = \mathbb{E} \left[ \frac{ZS}{\sqrt{X^2 + Y^2 + Z^2} |S|} \right].$$

To finish the lemma, observe  $\text{Tr}[V \mathbf{p} \mathbf{v}^T] = \mathbf{v}^T V \mathbf{p} = \mathbf{v}^T \tilde{V} \tilde{\mathbf{p}} = a \tilde{\mathbf{p}}_1$ .  $\square$

## 5 Proof of the Main Technical Lemmas

In this section we give a proof and the requisite tools for the proofs of the main technical lemmas. Recall we have some expectations from the rounding algorithm, and we are interested in finding lower bounds for these expectations over some set of the parameters. We seek proofs of the following:

**Lemma 17** (Strictly Quadratic Case). *Let  $\mathcal{S}$ ,  $\mathcal{T}$ , and  $-\mathcal{S}$  be defined according to Equation (26). Further let  $k \in \{1, 2, 3\}$  be fixed, and let*

$$\mathcal{P}_k = \begin{cases} \mathcal{S} & \text{if } k = 1 \\ \mathcal{T} & \text{if } k = 2 \\ -\mathcal{S} & \text{if } k = 3 \end{cases} \quad (48)$$

Let  $[z_1, z_2, z_3, z'_1, z'_2, z'_3] \sim \mathcal{N}(0, \Sigma'(a, b, c))$ . Then, we have the following lower bound:

$$\min_{\substack{[a,b,c] \in \mathcal{S} \\ [p,q,r] \in \mathcal{P}_k}} \frac{k + \mathbb{E} \left[ \frac{pz_1z'_1 + qz_2z'_2 + rz_3z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right]}{k + ap + bq + cr} \geq \begin{cases} 22/(15\pi) \approx 0.467 & \text{if } k = 1 \\ 1/3 + 24/(25\pi) \approx 0.639 & \text{if } k = 2 \\ 1/2 + 388/(405\pi) \approx 0.804 & \text{if } k = 3 \end{cases}. \quad (49)$$

**Lemma 18** (General Case). *Let  $k \in \{1, 2, 3\}$  be fixed. Let  $a, b, c, p, q, r, t_i, d_i, t_j, d_j$  be constants, such that  $|a|, |b|, |c|, |d_i|, |d_j| \leq 1$ , and which satisfy:*

$$\begin{aligned} -k &\leq ap + bq + rc \leq 4 - k, \\ -k &\leq ap + bq + rc + u_id_i + u_jd_j \leq 4 - k, \\ |t_id_i| &\leq l, \\ \text{and } |t_jd_j| &\leq l, \end{aligned} \quad (50)$$

where  $l$  is 1 if  $k = 1$  or 3 and  $l = 2$  if  $k = 2$ . Further, let  $[z_1, z_2, z_3, z'_1, z'_2, z'_3] \sim \mathcal{N}(0, \Sigma'(a, b, c))$ , let  $[s_1, x_1, x_2, x_3] \sim \mathcal{N}(0, \Sigma_1)$  for

$$\Sigma_1 = \begin{bmatrix} 1 & d_i & 0 & 0 \\ d_i & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

and let  $[s_2, y_1, y_2, y_3] \sim \mathcal{N}(0, \Sigma_2)$  for

$$\Sigma_2 = \begin{bmatrix} 1 & d_j & 0 & 0 \\ d_j & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$



Then,

$$\begin{aligned}
& \left( k + \mathbb{E} \left[ \frac{pz_1z'_1 + qz_2z'_2 + rz_3z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right] + \mathbb{E} \left[ \frac{t_i x_1 s_1}{\sqrt{x_1^2 + x_2^2 + x_3^2} |s_1|} \right] \right. \\
& \quad \left. + \mathbb{E} \left[ \frac{t_j y_1 s_2}{\sqrt{y_1^2 + y_2^2 + y_3^2} |s_2|} \right] \right) / \left( k + ap + bq + cr + t_i d_i + t_j d_j \right) \\
& \geq \begin{cases} 2/\pi - 1/4 \approx 0.387 & \text{if } k = 1 \\ 16/(9\pi) \approx 0.565 & \text{if } k = 2 \\ 3/8 + 11/(9\pi) \approx 0.764 & \text{if } k = 3 \end{cases}
\end{aligned} \tag{51}$$

Demonstrating these lemmas will require several tools. Roughly we will need some results in Hermite polynomial analysis (Section 5.1), some elementary considerations for optimizing rational functions over polytopes (Section 5.2), and some analytical bounds for specific functions of interest (Section 5.3). Finally we conclude with the proofs of these two technical statements, (Section 5.4 and Section 5.5). We give rough outlines of the two proofs in this section, so that the reader may have context for the following results:

**Quadratic Case** Given an expectation of the form Equation (49), the first task is to reduce the optimization of  $[p, q, r] \in \mathcal{P}_k$  to the optimization of  $[p, q, r]$  over only the extreme points of  $\mathcal{P}_k$ . This is accomplished by fixing  $[a, b, c]$  and minimizing with respect to  $[p, q, r]$ . From this perspective, we are minimizing a rational function on  $(p, q, r)$  and it will easily follow (using Section 5.2) that we may lower bound the expectation by fixing  $[p, q, r]$  to be an extreme point of  $\mathcal{P}_k$ . The next step is to lower bound the resulting expression over all choices  $[a, b, c] \in \mathcal{S}$ . Using Hermite polynomials (Section 5.1), we may evaluate the expectation in terms of a convergent series. Bounding the error obtained from a truncation of this series (Section 5.1.3), and using calculus to bound the truncation (Section 5.3 and the Appendix) yield the final result.

**General Case** Given an expectation of the form of Equation (51) with parameters restricted as in Equation (50), we can once again evaluate the expectation using an expansion in the Hermite polynomials. The resulting expression yields a convergent series for the quadratic part, as in the previous case, as well as a hypergeometric function for the linear part. Cutting off the Hermite expansion to linear order, and bounding the special function to linear order we obtain a lower bound which is once again a rational function in the parameters (subject to the assumed constraints). Then, we can “coarse grain” the optimization to get an optimization in fewer parameters as a lower bound, and apply the same bounds for optimizing functions over polytopes (Section 5.2 and Section 5.3) to complete the proof.

## 5.1 Hermite Polynomials

### 5.1.1 Definitions and Expansions

Hermite polynomials [O’D14, LaSt72] are orthogonal polynomials which provide an orthonormal basis for  $L^2(\mathbb{R}, \gamma)$  under the inner product introduced in Equation (23). One can obtain these polynomials by applying the Gram-Schmidt procedure to the set of standard monomials  $\{1, z, z^2, z^3, \dots\}$  using the inner product in Equation (23) or alternatively they can be defined using a generating

function:

$$e^{tz-t^2/2} = \sum_{j=0}^{\infty} \frac{h_j(z)t^j}{\sqrt{j!}}. \quad (52)$$

Hermite polynomials have a well-known explicit expression [LaSt72], and since we will make use of the formula in many proofs, we define the Hermite polynomials according to this expression:

**Definition 19.** *The  $n$ th Hermite polynomial is defined as:*

$$h_n(z) = \frac{\sqrt{n!}}{2^{n/2}} \sum_{l=0}^{n/2} \frac{(-1)^{n/2-l} (\sqrt{2}z)^{2l}}{(2l)!(n/2-l)!} \text{ if } n \text{ is even,}$$

$$\text{and } h_n(z) = \frac{\sqrt{n!}}{2^{n/2}} \sum_{l=0}^{(n-1)/2} \frac{(-1)^{(n-1)/2-l} (\sqrt{2}z)^{2l+1}}{(2l+1)!((n-1)/2-l)!} \text{ if } n \text{ is odd.}$$

The generalization of these polynomials to the multivariate case is simply found by taking products of single variable Hermite polynomials:

**Definition 20.** *Given some multi-index  $\mu \in [\mathbb{Z}_{\geq 0}]^n$ , we define the multivariate Hermite polynomials as  $h_\mu(\mathbf{z}) := \prod_{i=1}^n h_{\mu_i}(z_i)$ .*

As stated previously, our interest in these polynomials stems from the fact that they are a basis for the space  $L^2(\mathbb{R}^n, \gamma)$ . Hence, any function in this space can be expanded in the Hermite polynomials:

**Proposition 21.** 1. [O'D14] *Let  $f \in L^2(\mathbb{R}^n, \gamma)$ . We can define  $f(\mathbf{z}) = \sum_{\mu \in [\mathbb{Z}_{\geq 0}]^n} \hat{f}_\mu h_\mu(\mathbf{z})$  where  $\hat{f}_\mu = \langle f, h_\mu \rangle = \int f(\mathbf{z}) h_\mu(\mathbf{z}) d\gamma(\mathbf{z})$  and convergence is in  $L^2(\mathbb{R}^n, \gamma)$  (converges in norm defined by Equation (23)).*

2. [O'D14] *If  $z, z'$  are bivariate normal with  $\mathbb{E}[zz'] = \rho$  and  $|\rho| \leq 1$ , then  $\langle h_i(z), h_j(z') \rangle = \delta_{ij} \rho^i$  where  $\delta_{ij}$  is the standard discrete delta function.*

3. *Let  $f \in L^2(\mathbb{R}^3, \gamma)$  with Hermite expansion  $f(\mathbf{z}) = \sum_{i,j,k} \hat{f}_{i,j,k} h_i(z_1) h_j(z_2) h_k(z_3)$ . Let  $(z_1, z_2, z_3, z'_1, z'_2, z'_3) \sim \mathcal{N}(0, \Sigma'(a, b, c))$  with  $|a|, |b|, |c| \leq 1$ . Then,  $\mathbb{E}[f(\mathbf{z}) f(\mathbf{z}')] = \sum_{i,j,k} \hat{f}_{i,j,k}^2 a^i b^j c^k$ .*

For completeness, we supply a short proof of item 3 in the appendix on Page 48 (although it is implicit in [O'D14]).

### 5.1.2 Hermite Expansions of Interest

We wish to apply the general theory of Hermite polynomials to two specific functions of interest:

$$g(z) = \text{sign}(z) = \begin{cases} 1 & \text{if } z \geq 0 \\ -1 & \text{if } z < 0 \end{cases} \quad \text{and} \quad f(z_1, z_2, z_3) = \frac{z_1}{\sqrt{z_1^2 + z_2^2 + z_3^2}}.$$

First observe that the Hermite expansions of both of these functions are well defined, since both  $\int g(z)^2 d\gamma(z)$  and  $\int f(\mathbf{z})^2 d\gamma(\mathbf{z})$  are  $< \infty$ <sup>4</sup>. The first function has a well-known Hermite expansion:

<sup>4</sup>The latter case holds because  $\int f(z_1, z_2, z_3)^2 + f(z_2, z_1, z_3)^2 + f(z_3, z_1, z_2)^2 d\gamma(\mathbf{z}) = 1$ , so by symmetry  $\int f(\mathbf{z})^2 d\gamma(\mathbf{z}) = 1/3$ .

**Lemma 22** ([LaSt72]). Let  $g(z) = \text{sign}(z) = \begin{cases} 1 & \text{if } z \geq 0 \\ -1 & \text{if } z < 0 \end{cases}$ . Then,  $g(z) = \sum_i \hat{g}_i h_i(z)$  where

$$\hat{g}_i = \begin{cases} 0 & \text{if } i \text{ is even} \\ \frac{\sqrt{i!}(-1)^q}{2^{q-1/2}\sqrt{\pi}q!(1+2q)} & \text{if } i \text{ is odd} \end{cases},$$

with  $q = (i - 1)/2$ .

For the second function, let us define:

$$f(z_1, z_2, z_3) := \sum_{(i,j,k)} \hat{f}_{i,j,k} h_i(z_1) h_j(z_2) h_k(z_3) \quad (53)$$

We have chosen the notation  $\{\hat{f}_{i,j,k}\}$  rather than  $\{\hat{f}_{i,j,k}\}$  because  $\hat{f}_{i,j,k} = \hat{f}_{i,k,j}$ , which follows from the fact that  $f(z_1, z_2, z_3) = f(z_1, z_3, z_2)$  and orthonormality of the Hermite polynomials. Indeed,

$$\hat{f}_{i,j,k} = \int f(z_1, z_2, z_3) h_i(z_1) h_j(z_2) h_k(z_3) d\gamma(\mathbf{z}) = \int f(z_1, z_3, z_2) h_i(z_1) h_j(z_2) h_k(z_3) d\gamma(\mathbf{z}) = \hat{f}_{i,k,j}.$$

We can also see that  $\hat{f}_{i,j,k} = 0$  if  $i$  is even,  $j$  is odd or  $k$  is odd. This follows from the explicit form of the Hermite polynomials (Definition 19) as well as the fact that the integral of an anti-symmetric function along the entire real line is zero. Having stated some symmetries of the coefficients we now state their analytic values:

**Lemma 23.** Let  $f$  have Hermite expansion  $\{\hat{f}_{i,j,k}\}$  as defined in Equation (53). If  $i$  is odd,  $j$  is even, and  $k$  is even, the Hermite coefficients take values:

$$\hat{f}_{i,j,k} = 2\sqrt{\frac{2}{\pi}} \frac{\sqrt{i!j!k!}(-1)^p}{(i-1)!!j!!k!!(1+2p)(3+2p)} \quad (54)$$

for  $p = (i + j + k - 1)/2$ . Otherwise, the coefficients are zero.

*Proof.* This is a special case of Lemma 37 found in the Appendix.  $\square$

With these facts in hand, we can evaluate one of the expectations of interest to us:

**Lemma 24.** Let  $(z, z_1, z_2, z_3) \sim \mathcal{N}(0, \Sigma)$  with

$$\Sigma = \begin{bmatrix} 1 & a & 0 & 0 \\ a & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Then,

$$\mathbb{E} \left[ \frac{\text{sign}(z)z_1}{\sqrt{z_1^2 + z_2^2 + z_3^2}} \right] = \frac{4a}{3\pi} {}_2F_1 \left[ \begin{matrix} 1/2, 1/2 \\ 5/2 \end{matrix}; a^2 \right]. \quad (55)$$

*Proof.* See page 47 in the Appendix.  $\square$

This falls out of the Hermite expansions we have so far, plus some elementary combinatorial identities. For some intuition on how this fact is used in the analysis, note that the expectation in Equation (55) looks a lot like the component of one of the Bloch vectors in the rounding algorithm. Indeed, analyzing that expectation will be exactly the purpose of this lemma.

### 5.1.3 Bounds on Hermite Expansions

While exact computations like Lemma 24 would be desirable for all expansions of interest, in order to establish rigorous results we will need bounds in some cases. The first lemma we describe concerns the symmetries of an inner product of random vectors with covariance as defined in Equation (27):

**Lemma 25.** *Let  $[z_1, z_2, z_3, z'_1, z'_2, z'_3] \sim \mathcal{N}(0, \Sigma'(a, b, c))$  and let  $\{\hat{f}_{i,jk}\}$  be the Hermite expansion of  $f(\mathbf{z}) = z_1/\sqrt{z_1^2 + z_2^2 + z_3^2}$ . Let  $\mathbf{v}_1, \mathbf{v}_2$  be vectors with components defined as  $(\mathbf{v}_1)_i := z_i/\sqrt{z_1^2 + z_2^2 + z_3^2}$  and  $(\mathbf{v}_2)_i = z'_i/\sqrt{(z'_1)^2 + (z'_2)^2 + (z'_3)^2}$  for  $i \in \{1, 2, 3\}$ . Further, define the following families of symmetric polynomials:*

$$p_{i,jk}(a, b, c) = \begin{cases} a^i b^j c^k + a^i b^k c^j + b^i a^j c^k + b^i a^k c^j + c^i a^j b^k + c^i a^k b^j & \text{if } k \neq j \\ a^i b^j c^j + b^i a^j c^j + c^i a^j b^j & \text{if } k = j \end{cases}, \quad (56)$$

$$\text{and } u_{i,jk}(a, b, c) = \begin{cases} a^i b^j c^k + a^i b^k c^j & \text{if } k \neq j \\ a^i b^j c^j & \text{if } k = j \end{cases}. \quad (57)$$

Then,

$$\mathbb{E}[\mathbf{v}_1 \cdot \mathbf{v}_2] = \sum_{\substack{i,j,k \\ j \leq k}} \hat{f}_{i,jk}^2 p_{i,jk}(a, b, c), \quad (58)$$

$$\text{and } \mathbb{E}[(\mathbf{v}_1)_1 (\mathbf{v}_2)_1] = \sum_{\substack{i,j,k \\ j \leq k}} \hat{f}_{i,jk}^2 u_{i,jk}(a, b, c) \quad (59)$$

if  $|a|, |b|, |c| \leq 1$ .

*Proof.* By Proposition 21,

$$\mathbb{E}[f(z_1, z_2, z_3) f(z'_1, z'_2, z'_3)] = \sum_{i,j,k} \hat{f}_{i,jk}^2 a^i b^j c^k. \quad (60)$$

We can use the invariance  $\hat{f}_{i,jk} = \hat{f}_{i,kj}$  to write this as:

$$\mathbb{E}[f(z_1, z_2, z_3) f(z'_1, z'_2, z'_3)] = \sum_i \left[ \sum_{j < k} \hat{f}_{i,jk}^2 (a^i b^j c^k + a^i b^k c^j) + \sum_j \hat{f}_{i,jj}^2 a^i b^j c^j \right]. \quad (61)$$

This establishes Equation (59). For Equation (58), we wish to calculate:

$$\mathbb{E} \left[ \frac{z_1 z'_1 + z_2 z'_2 + z_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right] = \mathbb{E}[f(z_1, z_2, z_3) f(z'_1, z'_2, z'_3) + f(z_2, z_1, z_3) f(z'_2, z'_1, z'_3) + f(z_3, z_1, z_2) f(z'_3, z'_1, z'_2)]. \quad (62)$$

We can permute the variables in Equation (61) to obtain analogous expressions for  $\mathbb{E}[f(z_2, z_1, z_3) f(z'_2, z'_1, z'_3)]$  and  $\mathbb{E}[f(z_3, z_1, z_2) f(z'_3, z'_1, z'_2)]$ . Substituting these expressions into Equation (62) gives Equation (58).  $\square$

We will also need a remainder bound for finite Hermite expansions. We will state it for our Hermite expansion of interest,  $\{\hat{f}_{i,j,k}\}$ , although it clearly generalizes to others.

**Lemma 26.** *Let  $Q$  be a finite subset of  $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0}$  such that  $j \leq k$  for all  $(i, j, k) \in Q$ . Then,*

$$\left| \sum_{\substack{i,j,k \\ j \leq k}} \hat{f}_{i,j,k}^2 p_{i,j,k}(a, b, c) - \sum_{(i,j,k) \in Q} \hat{f}_{i,j,k}^2 p_{i,j,k}(a, b, c) \right| \leq 3 \left( 1/3 - \sum_{(i,j,k) \in Q} 2^{1-\delta_{jk}} \hat{f}_{i,j,k}^2 \right),$$

where  $\delta_{jk}$  is the standard discrete delta function and  $|a|, |b|, |c| \leq 1$ .

*Proof.* <sup>5</sup>

$$\left| \sum_{\substack{i,j,k \\ j \leq k}} \hat{f}_{i,j,k}^2 p_{i,j,k}(a, b, c) - \sum_{(i,j,k) \in Q} \hat{f}_{i,j,k}^2 p_{i,j,k}(a, b, c) \right| \leq \sum_{i,j \leq k} \hat{f}_{i,j,k}^2 \delta_{(i,j,k) \in -Q} |p_{i,j,k}(a, b, c)|, \quad (63)$$

where  $-Q = \{(j, k, l) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0} : k \leq l \text{ and } (j, k, l) \notin Q\}$  and  $\delta_{(i,j,k) \in -Q}$  is the standard discrete delta function which evaluates to 1 if the condition is met and 0 otherwise. Observe that for  $|a|, |b|, |c| \leq 1$ ,  $|p_{i,j,k}(a, b, c)| \leq 6$  when  $j \neq k$  and  $|p_{i,j,j}(a, b, c)| \leq 3$  so we can uniformly upper bound  $|p_{i,j,k}(a, b, c)| \leq 3 \cdot 2^{1-\delta_{jk}}$ . Hence, we can upper bound Equation (63) as:

$$\begin{aligned} \sum_{i,j \leq k} \hat{f}_{i,j,k}^2 \delta_{(i,j,k) \in -Q} 3 \cdot 2^{1-\delta_{jk}} &= \sum_{i,j \leq k} \hat{f}_{i,j,k}^2 3 \cdot 2^{1-\delta_{jk}} - \sum_{(i,j,k) \in Q} \hat{f}_{i,j,k}^2 3 \cdot 2^{1-\delta_{jk}} \\ &= 3 \left( \sum_{i,j < k} \frac{\hat{f}_{i,j,k}^2 + \hat{f}_{i,k,j}^2}{2} 2^{1-\delta_{jk}} + \sum_{i,j} \hat{f}_{i,j,j}^2 - \sum_{(i,j,k) \in Q} \hat{f}_{i,j,k}^2 2^{1-\delta_{jk}} \right), \end{aligned} \quad (64)$$

where we used  $\hat{f}_{i,j,k} = \hat{f}_{i,k,j}$ . We can now write Equation (64) as:

$$= 3 \left( \sum_{i,j,k} \hat{f}_{i,j,k}^2 - \sum_{(i,j,k) \in Q} \hat{f}_{i,j,k}^2 2^{1-\delta_{jk}} \right) = 3 \left( \frac{1}{3} - \sum_{(i,j,k) \in Q} \hat{f}_{i,j,k}^2 2^{1-\delta_{jk}} \right).$$

□

We can use the same proof with the upper bound  $|u_{i,j,k}(a, b, c)| \leq 2^{1-\delta_{j,k}}$  to obtain:

**Lemma 27.** *Let  $Q$  be a finite subset of  $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0}$  such that  $k \leq l$  for all  $(j, k, l) \in Q$ . Then,*

$$\left| \sum_{\substack{i,j,k \\ j \leq k}} \hat{f}_{i,j,k}^2 u_{i,j,k}(a, b, c) - \sum_{(i,j,k) \in Q} \hat{f}_{i,j,k}^2 u_{i,j,k}(a, b, c) \right| \leq 1/3 - \sum_{(i,j,k) \in Q} 2^{1-\delta_{j,k}} \hat{f}_{i,j,k}^2$$

where  $\delta_{j,k}$  is the standard discrete delta function and  $|a|, |b|, |c| \leq 1$ .

<sup>5</sup>There are some details here concerning convergence, etc. that we will not mention. For reference, observe that the series  $\sum_{i,j \leq k} \hat{f}_{i,j,k}^2 p_{i,j,k}(a, b, c)$  is absolutely convergent since  $|p_{i,j,k}(a, b, c)|$  is bounded and  $\sum_{i,j,k} \hat{f}_{i,j,k}^2 = 1/3$ , so the convergence proofs are a simple exercise in real analysis [Rud64].

## 5.2 Optimizing Rational Functions over Polytopes

The next set of important tools concerns optimizing rational functions over polytopes. These observations will be used to simplify analysis, by reducing the parameter space we need to examine for a lower bound on the approximation factor.

**Lemma 28.** *Let  $\mathcal{P}$  be some polytope, and  $(A, B, C, a, b, c, k)$  be constants such that*

$$\begin{aligned} k + [p, q, r] \cdot [A, B, C] &\geq 0 \\ \text{and } k + [p, q, r] \cdot [a, b, c] &\geq 0 \end{aligned}$$

for all  $[p, q, r] \in \mathcal{P}$ . Then,

$$\min_{[p, q, r] \in \mathcal{P}} \frac{k + [p, q, r] \cdot [A, B, C]}{k + [p, q, r] \cdot [a, b, c]} \geq \min_{[p, q, r] \in \mathcal{B}} \frac{k + [p, q, r] \cdot [A, B, C]}{k + [p, q, r] \cdot [a, b, c]}, \quad (65)$$

where  $\mathcal{B}$  is the set of extreme points for which the objective is defined.

*Proof.* By definition any  $[p, q, r]$  can be written as a convex combination of the extreme points of  $\mathcal{P}$ :  $[p, q, r] = \sum_i \lambda_i [p_i, q_i, r_i]$ . Re-write both the numerator and denominator under this decomposition:

$$\frac{\sum_i \lambda_i (k + [p_i, q_i, r_i] \cdot [A, B, C])}{\sum_i \lambda_i (k + [p_i, q_i, r_i] \cdot [a, b, c])}.$$

Apply the elementary fact that if  $\{a_1, \dots, a_n, b_1, \dots, b_n\}$  are non-negative constants, then:

$$\frac{a_1 + \dots + a_n}{b_1 + \dots + b_n} \geq \min_{i: b_i \neq 0} \frac{a_i}{b_i}.$$

The lemma follows. □

The above lemma indicates that if we have a restriction  $[p, q, r] \in \mathcal{P}$  we need only examine the extreme points for a lower bound. We can take it a step further and use the symmetries of our specific problem to demonstrate that a single extreme point is sufficient in each case, for the quadratic part of the objective. We obtain the following:

**Lemma 29.** *Fix  $k \in \{1, 2, 3\}$ . Let  $\mathcal{B}_1$  be the set of extreme points of  $\mathcal{S}$ , let  $\mathcal{B}_2$  be the set of extreme points of  $\mathcal{T}$ , and let  $\mathcal{B}_3$  be the set of extreme points of  $-\mathcal{S}$ . Let us further define*

$$\begin{aligned} A &= \mathbb{E} \left[ \frac{z_1 z'_1}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right], \\ B &= \mathbb{E} \left[ \frac{z_2 z'_2}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right], \\ \text{and } C &= \mathbb{E} \left[ \frac{z_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right], \end{aligned}$$

where  $[z_1, z_2, z_3, z'_1, z'_2, z'_3] \sim \mathcal{N}(0, \Sigma'(a, b, c))$ . Then,

$$\min_{\substack{[a, b, c] \in \mathcal{S} \\ [p, q, r] \in \mathcal{B}_k}} \frac{k + [p, q, r] \cdot [A, B, C]}{k + [p, q, r] \cdot [a, b, c]} = \min_{[a, b, c] \in \mathcal{S}} \frac{k + [p_k, q_k, r_k] \cdot [A, B, C]}{k + [p_k, q_k, r_k] \cdot [a, b, c]},$$

where

$$[p_k, q_k, r_k] = \begin{cases} [-1, -1, -1] & \text{if } k = 1 \\ [2, 0, 0] & \text{if } k = 2 \\ [1, 1, 1] & \text{if } k = 3 \end{cases}.$$

*Proof.* It is easy to see that the region  $\mathcal{S}$  is invariant under permutation of coordinates, as well as the linear transformation  $[a, b, c] \rightarrow [-a, -b, c]$ . Hence,

$$\begin{aligned} & \min_{[a,b,c] \in \mathcal{S}} \frac{k + [p, q, r] \cdot [A, B, C]}{k + [p, q, r] \cdot [a, b, c]} = \min_{[a,b,c] \in \mathcal{S}} \frac{k + [-p, -q, r] \cdot [-A, -B, C]}{k + [-p, -q, r] \cdot [-a, -b, c]} \\ &= \min_{[-a,-b,c] \in \mathcal{S}} \frac{k + [-p, -q, r] \cdot [A, B, C]}{k + [-p, -q, r] \cdot [a, b, c]} = \min_{[a,b,c] \in \mathcal{S}} \frac{k + [-p, -q, r] \cdot [A, B, C]}{k + [-p, -q, r] \cdot [a, b, c]}. \end{aligned}$$

For every  $k$ , all the points in  $\mathcal{B}_k$  are related by a permutation of coordinates followed by  $[p, q, r] \rightarrow [-p, -q, r]$ . Hence, all points in  $\mathcal{B}_k$  result in the same value when we minimize and we are free to pick an arbitrary extreme point.  $\square$

### 5.3 Technical Estimates for Functions of Interest

There are several functions we will need bounds on. The first set of functions (Lemma 30 - Lemma 32) come from truncations of Hermite expansions, and the second set (Lemma 33) is related to the (exact) expansion for the linear part of the objective. In this direction let us define the following three functions:

$$\begin{aligned} q_1(a, b, c) &= \hat{f}_{1,00}^2 p_{1,00}(a, b, c) + \hat{f}_{1,02}^2 p_{1,02}(a, b, c) + \hat{f}_{3,00}^2 p_{3,00}(a, b, c) \\ &\quad + 3(1/3 - \hat{f}_{1,00}^2 - 2\hat{f}_{1,02}^2 - \hat{f}_{3,00}^2), \\ q_2(a, b, c) &= \hat{f}_{1,00}^2 a + \hat{f}_{1,02}^2 a(b^2 + c^2) + \hat{f}_{3,00}^2 a^3 \\ &\quad - (1/3 - \hat{f}_{1,00}^2 - 2\hat{f}_{1,02}^2 - \hat{f}_{3,00}^2), \\ \text{and } q_3(a, b, c) &= \hat{f}_{1,00}^2 p_{1,00}(a, b, c) + \hat{f}_{1,02}^2 p_{1,02}(a, b, c) + \hat{f}_{3,00}^2 p_{3,00}(a, b, c) \\ &\quad - 3(1/3 - \hat{f}_{1,00}^2 - 2\hat{f}_{1,02}^2 - \hat{f}_{3,00}^2). \end{aligned}$$

Each of these functions will correspond to a different rank,  $k$ . Since  $[p, q, r]$  changes in each case, we will need different functions to lower bound the expectation, i.e. we may need to add or subtract the remainder term based on the signs of  $[p, q, r]$ . For completeness, note the following analytic values of Hermite coefficients from Equation (54):

$$\hat{f}_{1,00}^2 = \frac{8}{9\pi}, \quad \hat{f}_{1,02}^2 = \frac{4}{225\pi}, \quad \text{and} \quad \hat{f}_{3,00}^2 = \frac{4}{75\pi}.$$

With these definitions in hand, we can give the lower bounds we use for functions of interest:

**Lemma 30** (Rank 1 Lemma). *It holds that*

$$\min_{[a,b,c] \in \mathcal{S}} \frac{1 - \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a, b, c)}{1 - a - b - c} \geq \frac{1 - q_1(-1, -1, -1)}{4} = \frac{22}{15\pi}.$$

**Lemma 31** (Rank 2 Lemma). *It holds that*

$$\min_{[a,b,c] \in \mathcal{S}} \frac{1 + \sum_{i,j \leq k} \hat{f}_{i,jk}^2 u_{i,jk}(a, b, c)}{1 + a} \geq \frac{1 + q_2(1, 0, 0)}{1 + 1} = \frac{1}{3} + \frac{24}{25\pi}.$$

**Lemma 32** (Rank 3 Lemma). *It holds that*

$$\min_{[a,b,c] \in \mathcal{S}} \frac{3 + \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a, b, c)}{3 + a + b + c} \geq \frac{3 + q_3(1/3, 1/3, 1/3)}{3 + 1} = \frac{1}{2} + \frac{388}{405\pi}.$$

*Proof.* See page 48 in the Appendix. □

The final bound we need before proving the main technical lemmas follows:

**Lemma 33.** *Let  $a \in \mathbb{R}$  with  $|a| \leq 1$ , and let  $b(a) = \frac{4a}{3\pi} {}_2F_1 \left[ \begin{matrix} 1/2, 1/2 \\ 5/2 \end{matrix}; a^2 \right]$ . Then,*

$$b(a) \geq \begin{cases} (1/2)a & \text{if } a \leq 0 \\ 4/(3\pi)a & \text{if } a \geq 0 \end{cases}$$

and

$$b(a) \leq \begin{cases} 4/(3\pi)a & \text{if } a \leq 0 \\ (1/2)a & \text{if } a \geq 0 \end{cases}.$$

*Proof.* If  $a \geq 0$  it is easy to see that the first and second derivatives of  $b(a)$  are positive, so the function is increasing and convex in this region. this implies that  $b'(0)a \leq b(a) \leq (b(1) - b(0))a$  in this region. Similarly, it is easy to see that if  $a \leq 0$  the first derivative is positive and the second is negative. Hence,  $b(a)$  is increasing and concave so the rest of the lemma follows. □

#### 5.4 Proof of Lemma 17

Recall we are interested in evaluating

$$\xi := \min_{\substack{[a,b,c] \in \mathcal{S} \\ [p,q,r] \in \mathcal{P}_k}} \frac{k + \mathbb{E} \left[ \frac{pz_1 z'_1 + qz_2 z'_2 + rz_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right]}{k + ap + bq + cr}.$$

Observe that both the numerator and denominator correspond to  $Tr(\mathcal{O}\phi)$  for some PSD observable  $\mathcal{O}$  and some valid density matrix  $\phi$  in all cases. Hence, for fixed  $[a, b, c]$  the numerator and denominator are non-negative and we can apply Lemma 28:

$$\xi \geq \min_{\substack{[a,b,c] \in \mathcal{S} \\ [p,q,r] \in \mathcal{B}_k}} \frac{k + \mathbb{E} \left[ \frac{pz_1 z'_1 + qz_2 z'_2 + rz_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right]}{k + ap + bq + cr},$$

where  $\mathcal{B}_k$  is the set of extreme points of  $\mathcal{P}_k$ . Now let us apply Lemma 29, which implies that we need only consider a particular  $[p_k, q_k, r_k]$  for each case. We obtain:

$$\xi \geq \min_{[a,b,c] \in \mathcal{S}} \frac{k + \mathbb{E} \left[ \frac{p_k z_1 z'_1 + q_k z_2 z'_2 + r_k z_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right]}{k + ap_k + bq_k + cr_k},$$

for

$$[p_k, q_k, r_k] = \begin{cases} [-1, -1, -1] & \text{if } k = 1 \\ [2, 0, 0] & \text{if } k = 2 \\ [1, 1, 1] & \text{if } k = 3 \end{cases}.$$



Now we can expand in the Hermite polynomials according to Lemma 25:

$$\xi \geq \begin{cases} \min_{[a,b,c] \in \mathcal{S}} \frac{1 - \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a,b,c)}{1-a-b-c} & \text{if } k = 1 \\ \min_{[a,b,c] \in \mathcal{S}} \frac{2+2 \sum_{i,j \leq k} \hat{f}_{i,jk}^2 u_{i,jk}(a,b,c)}{2+2a} & \text{if } k = 2 \\ \min_{[a,b,c] \in \mathcal{S}} \frac{3 + \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a,b,c)}{3+a+b+c} & \text{if } k = 3 \end{cases} .$$

Finally, we can apply Lemmas 26-27 and Lemmas 30-32.

## 5.5 Proof of Lemma 18

Recall we are interested in finding a lower bound for:

$$\begin{aligned} \xi := \min & \left( k + \mathbb{E} \left[ \frac{pz_1z'_1 + qz_2z'_2 + rz_3z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right] + \mathbb{E} \left[ \frac{t_i x_1 s_1}{\sqrt{x_1^2 + x_2^2 + x_3^2} |s_1|} \right] \right. \\ & \left. + \mathbb{E} \left[ \frac{t_j y_1 s_2}{\sqrt{y_1^2 + y_2^2 + y_3^2} |s_2|} \right] \right) / \left( k + ap + bq + cr + t_i d_i + t_j d_j \right) \\ & \text{s.t. } -k \leq ap + bq + rc \leq 4 - k, \\ & -k \leq ap + bq + rc + t_i d_i + t_j d_j \leq 4 - k, \\ & |t_i d_i| \leq l, \\ & \text{and } |t_j d_j| \leq l. \end{aligned} \quad (66)$$

We can evaluate the expectations according to Lemma 24 and Lemma 25 to write the objective of Equation (66) as:

$$\begin{aligned} \xi = \min & \left( k + p \sum_{i,j \leq k} \hat{f}_{i,jk}^2 u_{i,jk}(a,b,c) + q \sum_{i,j \leq k} \hat{f}_{i,jk}^2 u_{i,jk}(b,a,c) + r \sum_{i,j \leq k} \hat{f}_{i,jk}^2 u_{i,jk}(c,a,b) \right. \\ & \left. + t_i \frac{4d_i}{3\pi} {}_2F_1 \left[ \begin{matrix} 1/2, 1/2 \\ 5/2 \end{matrix}; d_i^2 \right] + t_j \frac{4d_j}{3\pi} {}_2F_1 \left[ \begin{matrix} 1/2, 1/2 \\ 5/2 \end{matrix}; d_j^2 \right] \right) / \left( k + ap + bq + cr + t_i d_i + t_j d_j \right) \\ & \text{s.t. } \dots \end{aligned}$$

We can apply Lemma 27 with  $Q = \{(1, 0, 0)\}^6$ , and Lemma 33 to obtain the minimization:

$$\begin{aligned} \xi \geq \min & \frac{k + \frac{8}{9\pi}(ap + bq + rc) - 3\left(\frac{1}{3} - \frac{8}{9\pi}\right) + c_i t_i d_i + c_j t_j d_j}{k + ap + bq + rc + t_i d_i + t_j d_j} \\ & \text{s.t. } -k \leq ap + bq + rc \leq 4 - k, \\ & -k \leq ap + bq + rc + t_i d_i + t_j d_j \leq 4 - k, \\ & c_i, c_j \in \left\{ \frac{1}{2}, \frac{4}{3\pi} \right\}, \\ & |t_i d_i| \leq l, \\ & \text{and } |t_j d_j| \leq l. \end{aligned}$$

Let us ‘‘coarse grain’’ the optimization by setting  $v_1 = ap + bq + rc$ ,  $v_2 = t_i d_i$  and  $v_3 = t_j d_j$ . The optimization we obtain by coarse graining is a lower bound on the original optimization by the

<sup>6</sup>Note that the remainder term,  $3(1/3 - 8/(9\pi))$ , can be taken negative for a lower bound because the denominator is always non-negative by the constraints.

following: Introduce variables  $v_1$ ,  $v_2$  and  $v_3$  into the program with constraints  $v_1 = ap + bq + rc$ ,  $v_2 = t_i d_i$  and  $v_3 = t_j d_j$ . We can rewrite the objective and the original constraints so that they are written entirely in  $(v_1, v_2, v_3)$ , while maintaining an equivalent problem (we have simply renamed things). Then, disregard the constraints  $v_1 = ap + bq + rc$ ,  $v_2 = t_i d_i$  and  $v_3 = t_j d_j$ . Since we are de-constraining the problem the new objective is non-increasing, and the variables  $(a, b, c, p, q, r, t_i, t_j)$  are redundant and can be ignored. The program obtained from this is:

$$\begin{aligned} \xi \geq \min & \frac{k + \frac{8}{9\pi}v_1 - 3(\frac{1}{3} - \frac{8}{9\pi}) + c_i v_2 + c_j v_3}{k + v_1 + v_2 + v_3} \\ & s.t. -k \leq v_1 \leq 4 - k, \\ & -k \leq v_1 + v_2 + v_3 \leq 4 - k, \\ & c_i, c_j \in \left\{ \frac{1}{2}, \frac{4}{3\pi} \right\}, \\ & |v_2| \leq l, \\ & \text{and } |v_3| \leq l. \end{aligned} \tag{67}$$

We will proceed by solving the above optimization problem for any fixed choice of  $(c_i, c_j)$ . For any fixed  $(c_i, c_j)$ ,  $[v_1, v_2, v_3]$  is constrained to a polytope, so we may apply Lemma 28 (the denominator is non-negative by the constraints, and it is easy to check that the numerator is always non-negative) to lower bound the optimization problem as:

$$\min \dots s.t. [v_1, v_2, v_3] \in \mathcal{Q}_k \geq \min \dots s.t. [v_1, v_2, v_3] \in \mathcal{B}_k,$$

where  $\mathcal{Q}_k$  is the polytope defined via the constraints of Equation (67), and  $\mathcal{B}_k$  is the corresponding set of extreme points. Hence, for any fixed  $(c_i, c_j)$ , we have a lower bound on the optimization by checking extreme points of some polytope. By checking these points and checking all possible values of  $c_i, c_j$  we obtain the following minima:

$$\xi \geq \begin{cases} 2/\pi - 1/4 \approx 0.387 & \text{if } k = 1 \\ 16/(9\pi) \approx 0.565 & \text{if } k = 2 \\ 3/8 + 11/(9\pi) \approx 0.764 & \text{if } k = 3 \end{cases} .$$

## 6 Proof of Main Theorems

We will first provide a proof of Theorem 11, since techniques used in this proof will be similar to the general case.

### 6.1 Proof of Theorem 11 (strictly quadratic case)

The proof will follow the outline in Section 2.5, but here we will point to the formal components needed in each step. Suppose we are given an instance of Problem 6 and let

$$\rho = \bigotimes_{i=1}^n \left( \frac{\mathbb{I} + \theta_{iX}\sigma^1 + \theta_{iY}\sigma^2 + \theta_{iZ}\sigma^3}{2} \right) \tag{68}$$

be the (random) output of the approximation algorithm (Algorithm 9). The total expected cost is:

$$\mathbb{E} \left[ \sum_{ij} Tr[H_{ij}\rho] \right] = \sum_{ij} \mathbb{E}[Tr[H_{ij}\rho]]. \tag{69}$$

Now observe that, if we can find a worst case lower bound on

$$\frac{\mathbb{E}[\text{Tr}[H_{ij}\rho]]}{\text{rank}(H_{ij})/4 + \text{Tr}[M^*C_{ij}]}, \quad (70)$$

we would be able to uniformly bound each term, and hence be able to provide Theorem 11. The majority of the technical work presented here is aimed at precisely such a goal<sup>7</sup>. Note that we can WLOG assume that  $H_{ij}$  is a strictly quadratic projector since the positive constant of proportionality will cancel from the numerator and denominator.

Let  $D$  be a nonzero off-diagonal block of  $C_{ij}$  corresponding to qubits  $i$  and  $j$ . Formally,  $D \in \mathbb{R}^{3 \times 3}$  such that  $D_{k,l} = C_{ij}(\sigma_i^k, \sigma_j^l)$  for all  $k, l$  in  $[3]$ . Let  $M^*$  be the optimal moment matrix with a Cholesky vectors  $\{\mathbf{v}_{ik}\}$  for all  $i \in [n], k \in [3]$  and let  $\mathbf{v}_0$  be the Cholesky vector corresponding to index  $\mathbb{I}$ . By this we mean that  $\mathbf{v}_{ik}^T \mathbf{v}_{jl} = M^*(\sigma_i^k, \sigma_j^l)$  and  $\mathbf{v}_0^T \mathbf{v}_{jl} = M^*(\mathbb{I}, \sigma_j^l)$ . Define  $V_i = [\mathbf{v}_{i1}, \mathbf{v}_{i2}, \mathbf{v}_{i3}]$  and  $V_j = [\mathbf{v}_{j1}, \mathbf{v}_{j2}, \mathbf{v}_{j3}]$ . We can rewrite our expectation of interest as:

$$\mathbb{E}[\text{Tr}[H_{ij}\rho]] = \frac{\text{rank}(H_{ij})}{4} + 2\mathbb{E}\left[\frac{\mathbf{r}^T V_i D V_j^T \mathbf{r}}{\|V_i^T \mathbf{r}\| \|V_j^T \mathbf{r}\|}\right], \quad (71)$$

where the factor of 2 is used to account for the symmetry of  $C_{ij}$ . Note, as we have previously mentioned,  $\text{sign}(\mathbf{v}_0^T \mathbf{r})$  is squared, so it has no effect on the objective. Hence we can rewrite Equation (70):

$$\frac{\mathbb{E}[\text{Tr}[H_{ij}\rho]]}{\text{rank}(H_{ij})/4 + \text{Tr}[M^*C_{ij}]} = \frac{k + \mathbb{E}\left[\frac{\mathbf{r}^T V_i (8D) V_j^T \mathbf{r}}{\|V_i^T \mathbf{r}\| \|V_j^T \mathbf{r}\|}\right]}{k + \text{Tr}[V_i (8D) V_j^T]}. \quad (72)$$

Observe that  $V_i^T V_j$  must correspond to a valid quadratic 2-moment for density matrix  $\rho_{ij}$  in the relaxation, since the SDP is constrained in that way (recall the SDP is defined so that the moment matrix is consistent with *local* density matrices). Similarly,  $8D$  must correspond to a 2-moment for a strictly quadratic projector of rank  $k$  by our assumptions on the problem instance (compare with Equation (3)). Hence, we can apply Lemma 15 (note that  $C$  in the context of that Lemma is the same as  $8D$  in the context of this proof), which implies

$$\frac{\mathbb{E}[\text{Tr}[H_{ij}\rho]]}{\text{rank}(H_{ij})/4 + \text{Tr}[M^*C_{ij}]} \geq \min_{\substack{[a,b,c] \in \mathcal{S} \\ [p,q,r] \in \mathcal{P}_k}} \frac{k + \mathbb{E}\left[\frac{pz_1 z'_1 + qz_2 z'_2 + rz_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}}\right]}{k + ap + bq + cr}, \quad (73)$$

where for fixed  $[a, b, c]$ ,  $[\mathbf{z}, \mathbf{z}'] \sim \mathcal{N}(0, \Sigma'(a, b, c))$  and  $\mathcal{P}_k = \begin{cases} -\mathcal{S} & \text{if } k = 3 \\ \mathcal{T} & \text{if } k = 2 \\ \mathcal{S} & \text{if } k = 1 \end{cases}$ . Note that we have met exactly the conditions of Lemma 17, so we may apply it to obtain the theorem.

## 6.2 Proof of Theorem 10

Just as before, we will lower bound the expected approximation factor by arguing a worst case lower bound on the expected approximation factor of a single edge. Let  $H_{ij} = P_{ij} \otimes \mathbb{I}_{[n] \setminus \{i,j\}}$  be some projector of rank  $k$  and let  $C_{ij}$  be the corresponding value matrix defined by Equation (3).

<sup>7</sup>The careful reader will notice there is an edge case, where  $\text{rank}(H_{ij})/4 + \text{Tr}[M^*C_{ij}] = 0$ . However, for this case the SDP is earning 0, and since  $H_{ij} \succeq 0$ , any  $\rho$  has objective at least the SDP.

Just as in the previous proof, let  $D \in \mathbb{R}^{3 \times 3}$  be the ‘‘quadratic part’’ of  $C_{ij}$  with elements defined as  $D_{k,l} = C_{ij}(\sigma_i^k, \sigma_j^l)$  for all  $k, l \in [3]$ . Let  $\mathbf{w}_i/\mathbf{w}_j$  be the 1-local part of  $C_{ij}$  corresponding to qubit  $i/j$ . Specifically,  $\mathbf{w}_i$  and  $\mathbf{w}_j$  are vectors in  $\mathbb{R}^3$  satisfying

$$(\mathbf{w}_i)_k = C_{ij}(\sigma_i^k, \mathbb{I}), \quad (74)$$

$$\text{and } (\mathbf{w}_j)_l = C_{ij}(\mathbb{I}, \sigma_j^l). \quad (75)$$

Let  $M^*$  be the optimal moment matrix with a Cholesky vectors  $\{\mathbf{v}_{ik}\}$  for all  $i \in [n], k \in [3]$  and let  $\mathbf{v}_0$  be the Cholesky vector corresponding to index  $\mathbb{I}$ . By this we mean that  $\mathbf{v}_{ik}^T \mathbf{v}_{jl} = M(\sigma_i^k, \sigma_j^l)$  and  $\mathbf{v}_0^T \mathbf{v}_{jl} = M(\mathbb{I}, \sigma_j^l)$ . Lastly, we will define  $V_i = [\mathbf{v}_{i1}, \mathbf{v}_{i2}, \mathbf{v}_{i3}]$  and  $V_j = [\mathbf{v}_{j1}, \mathbf{v}_{j2}, \mathbf{v}_{j3}]$ , matrices composed of Cholesky vectors for a single qubit.

Just as in Theorem 11, we can write the expected approximation factor as the quotient of the expected objective from the rounding algorithm and the value that the relaxation obtains. We can write:

$$\begin{aligned} & \frac{\mathbb{E}[\text{Tr}[H_{ij}\rho]]}{k/4 + \text{Tr}[C_{ij}M]} \\ &= \left( \frac{k}{4} + 2\mathbb{E} \left[ \frac{\mathbf{r}^T V_i D V_j^T \mathbf{r}}{\|V_i^T \mathbf{r}\| \|V_j^T \mathbf{r}\|} \right] + 2\mathbb{E} \left[ \frac{(\mathbf{r}^T V_i \mathbf{w}_i)(\mathbf{v}_0^T \mathbf{r})}{\|V_i^T \mathbf{r}\| |\mathbf{v}_0^T \mathbf{r}|} \right] + 2\mathbb{E} \left[ \frac{(\mathbf{r}^T V_j \mathbf{w}_j)(\mathbf{v}_0^T \mathbf{r})}{\|V_j^T \mathbf{r}\| |\mathbf{v}_0^T \mathbf{r}|} \right] \right) \\ & \quad / \left( \frac{k}{4} + 2\text{Tr}[V_i D V_j^T] + 2\text{Tr}[V_i \mathbf{w}_i \mathbf{v}_0^T] + 2\text{Tr}[V_j \mathbf{w}_j \mathbf{v}_0^T] \right). \end{aligned} \quad (76)$$

Multiplying the numerator and denominator by 4 we can find an expected approximation factor of

$$= \frac{k + \mathbb{E} \left[ \frac{\mathbf{r}^T V_i (8D) V_j^T \mathbf{r}}{\|V_i^T \mathbf{r}\| \|V_j^T \mathbf{r}\|} \right] + \mathbb{E} \left[ \frac{\mathbf{r}^T V_i (8\mathbf{w}_i) \mathbf{v}_0^T \mathbf{r}}{\|V_i^T \mathbf{r}\| |\mathbf{v}_0^T \mathbf{r}|} \right] + \mathbb{E} \left[ \frac{\mathbf{r}^T V_j (8\mathbf{w}_j) \mathbf{v}_0^T \mathbf{r}}{\|V_j^T \mathbf{r}\| |\mathbf{v}_0^T \mathbf{r}|} \right]}{k + \text{Tr}[V_i (8D) V_j^T] + \text{Tr}[V_i (8\mathbf{w}_i) \mathbf{v}_0^T] + \text{Tr}[V_j (8\mathbf{w}_j) \mathbf{v}_0^T]}. \quad (77)$$

We can apply Lemma 15 and Lemma 16 to convert Equation (77) to a standard form:

$$= \frac{k + \mathbb{E} \left[ \frac{pz_1 z'_1 + qz_2 z'_2 + rz_3 z'_3}{\sqrt{(z_1^2 + z_2^2 + z_3^2)((z'_1)^2 + (z'_2)^2 + (z'_3)^2)}} \right] + \mathbb{E} \left[ \frac{t_i x_1 s_1}{\sqrt{x_1^2 + x_2^2 + x_3^2} |s_1|} \right] + \mathbb{E} \left[ \frac{t_j y_1 s_2}{\sqrt{y_1^2 + y_2^2 + y_3^2} |s_2|} \right]}{k + ap + bq + cr + t_i d_i + t_j d_j}, \quad (78)$$

where random variables are distributed according to:

$$\begin{aligned} [z_1, z_2, z_3, z'_1, z'_2, z'_3] &\sim \mathcal{N}(0, \Sigma'(a, b, c)), \\ [s_1, x_1, x_2, x_3] &\sim \mathcal{N}(0, \Sigma_1), \\ \text{and } [s_2, y_1, y_2, y_3] &\sim \mathcal{N}(0, \Sigma_2) \end{aligned}$$

with

$$\Sigma_1 = \begin{bmatrix} 1 & d_i & 0 & 0 \\ d_i & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad \Sigma_2 = \begin{bmatrix} 1 & d_j & 0 & 0 \\ d_j & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}. \quad (79)$$

Now let us note several constraints on the parameters. Just as in the proof of Theorem 11, Lemma 15 implies that  $[a, b, c] \in \mathcal{S}$  and  $[p, q, r] \in \mathcal{P}_k$  where  $\mathcal{P}_k$  is defined in Equation (37). Equation (32) implies

that  $-k \leq ap + bq + rc \leq 4 - k$ . Note further that  $ap + bq + rc + t_i d_i + t_j d_j = 4\text{Tr}(P\rho) - k$  for some 2 qubit projector  $P$  of rank  $k$  and some density matrix on two qubits (here we are crucially using the constraints that force the moment matrix to correspond to a valid 2 qubits marginals), hence Equation (31) implies  $-k \leq ap + bq + rc + t_i d_i + t_j d_j \leq 4 - k$ . Finally, let us note that Equations 33, 34 imply that  $|t_i d_i|, |t_j d_j| \leq 1$  if  $k = 1$  or  $3$  and  $|t_i d_i|, |t_j d_j| \leq 2$  if  $k = 2$ . Let us say that  $|t_i d_i|, |t_j d_j| \leq l$  where  $l$  is 1 if  $k = 1$  or  $3$  and  $l$  is 2 if  $k = 2$ . Hence, we have met the conditions required for Lemma 18 which completes the proof.

## 7 Conclusion

In this work we have demonstrated several new approximation algorithms for interesting cases of the 2-Local Hamiltonian problem. As is the theme in many works [BBT09, GK12, BH16], we have given evidence that the geometry of 2-Local interactions can drastically effect approximability for traceless Hamiltonians since we demonstrate the bipartite case has a constant factor approximation algorithm and the unconstrained case is known to have no constant factor algorithm [BGKT19]. In addition to this, we have given a novel approximation algorithm/analysis for the 2-Local Hamiltonian with local terms which are also projectors. This is especially interesting given the scarcity of approximation algorithms for quantum problems. Indeed, the rank 3 case, has been open for some time [GK12, HLP20]. Furthermore, we have provided new techniques for rounding to product states that we believe will have additional applications in quantum information. Our rounding algorithm is quite “natural” given the solution of the SDP, and the ability to understand the expectation through Hermite polynomial analysis seems likely to extend to other Hamiltonians/problems.

Future work includes tightening our analysis with more sophisticated methods. As stated in the introduction, we believe the strictly quadratic case is a very interesting special case of the general problem, and given the symmetry of the problem, the analysis may be amenable to techniques from algebraic geometry. In particular, we hope to replace the tedious analysis from lemmas 30-32 with a more sophisticated approach that can handle higher order Hermite expansions. Such a method would also probably allow for a better analysis of the approximation algorithm itself, potentially allowing us to prove what we believe to be the attained approximation factor for our rounding algorithm. Another natural direction is removing the assumption we have that the terms are projectors, and replacing it with generic PSD terms. Implicit in this work is a 0.387-approximation to such problems, by writing a given PSD term as a positive combination of projectors, the question is whether or not a better approximation can be obtained, perhaps with additional assumptions.

### 7.1 Acknowledgments

Sandia National Laboratories is a multimission laboratory managed and operated by National Technology and Engineering Solutions of Sandia, LLC., a wholly owned subsidiary of Honeywell International, Inc., for the U.S. Department of Energy’s National Nuclear Security Administration under contract DE-NA-0003525. This work was supported by the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, Accelerated Research in Quantum Computing and Quantum Algorithms Teams programs.

## References

- [ABE<sup>+</sup>05] Sanjeev Arora, Eli Berger, Hazan Elad, Guy Kindler, and Muli Safra. On non-approximability for quadratic programs. In *46th Annual IEEE Symposium on Founda-*

- tions of Computer Science (FOCS'05), pages 206–215. IEEE, 2005.
- [AGM20] Anurag Anshu, David Gosset, and Karen Morenz. Beyond Product State Approximations for a Quantum Analogue of Max Cut. In Steven T. Flammia, editor, *15th Conference on the Theory of Quantum Computation, Communication and Cryptography (TQC 2020)*, volume 158 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 7:1–7:15, Dagstuhl, Germany, 2020. Schloss Dagstuhl–Leibniz-Zentrum für Informatik.
- [AS48] Milton Abramowitz and Irene A Stegun. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, volume 55. US Government printing office, 1948.
- [BBT09] Nikhil Bansal, Sergey Bravyi, and Barbara M. Terhal. Classical approximation schemes for the ground-state energy of quantum and classical ising spin hamiltonians on planar graphs. *Quantum Info. Comput.*, 9(7):701–720, July 2009.
- [BdOFV10] Jop Briët, Fernando Mário de Oliveira Filho, and Frank Vallentin. The positive semidefinite grothendieck problem with rank constraint. In *International Colloquium on Automata, Languages, and Programming*, pages 31–42. Springer, 2010.
- [BdOFV14] Jop Briët, Fernando Mário de Oliveira Filho, and Frank Vallentin. Grothendieck inequalities for semidefinite programs with rank constraint. *Theory of Computing*, 10(4):77–105, 2014.
- [Bet31] Hans Bethe. Zur theorie der metalle. *Zeitschrift für Physik*, 71(3-4):205–226, 1931.
- [BG19] Anne Broadbent and Alex B Grilo. Zero-knowledge for qma from locally simulatable proofs. *arXiv preprint arXiv:1911.07782*, 2019.
- [BGG<sup>+</sup>18] Vijay Bhattiprolu, Mrinalkanti Ghosh, Venkatesan Guruswami, Euiwoong Lee, and Madhur Tulsiani. Approximating operator norms via generalized krivine rounding. *arXiv preprint arXiv:1804.03644*, 2018.
- [BGKT19] Sergey Bravyi, David Gosset, Robert König, and Kristan Temme. Approximation algorithms for quantum many-body problems. *Journal of Mathematical Physics*, 60(3):032203, 2019.
- [BH16] Fernando GSL Brandao and Aram W Harrow. Product-state approximations to quantum states. *Communications in Mathematical Physics*, 342(1):47–80, 2016.
- [BMMN13] Mark Braverman, Konstantin Makarychev, Yury Makarychev, and Assaf Naor. The grothendieck constant is strictly smaller than krivine’s bound. In *Forum of Mathematics, Pi*, volume 1. Cambridge University Press, 2013.
- [Bra11] Sergey Bravyi. Efficient algorithm for a quantum analogue of 2-sat. *Contemporary Mathematics*, 536:33–48, 2011.
- [Bri11] Jop Briët. *Grothendieck Inequalities, Nonlocal Games and Optimization*. PhD thesis, Universiteit van Amsterdam, 2011.
- [BS07] Salman Beigi and Peter W Shor. On the complexity of computing zero-error and holevo capacity of quantum channels. *arXiv preprint arXiv:0709.2090*, 2007.

- [BV04] Stephen Boyd and Lieven Vandenberghe. *Convex Optimization*. Cambridge University Press, 2004.
- [CW04] Moses Charikar and Anthony Wirth. Maximizing quadratic programs: Extending grothendieck’s inequality. In *45th Annual IEEE Symposium on Foundations of Computer Science*, pages 54–60. IEEE, 2004.
- [DLTW08] Andrew C Doherty, Yeong-Cherng Liang, Ben Toner, and Stephanie Wehner. The quantum moment problem and bounds on entangled multi-prover games. In *2008 23rd Annual IEEE Conference on Computational Complexity*, pages 199–210. IEEE, 2008.
- [DS05] Irit Dinur and Samuel Safra. On the hardness of approximating minimum vertex cover. *Annals of mathematics*, pages 439–485, 2005.
- [FL20] Shmuel Friedland and Lek-Heng Lim. Symmetric grothendieck inequality. *arXiv preprint arXiv:2003.07345*, 2020.
- [Gam16] Omar Gamel. Entangled bloch spheres: Bloch matrix and two-qubit state space. *Physical Review A*, 93(6):062320, 2016.
- [GK12] Sevag Gharibian and Julia Kempe. Approximation algorithms for qma-complete problems. *SIAM Journal on Computing*, 41(4):1028–1050, 2012.
- [GP19] Sevag Gharibian and Ojas Parekh. Almost Optimal Classical Approximation Algorithms for a Quantum Generalization of Max-Cut. In Dimitris Achlioptas and László A. Végh, editors, *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2019)*, volume 145 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 31:1–31:17, Dagstuhl, Germany, 2019. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik.
- [GW95] Michel X Goemans and David P Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *Journal of the ACM (JACM)*, 42(6):1115–1145, 1995.
- [Hås99] Johan Håstad. Clique is hard to approximate within  $1-\epsilon$ . *Acta Mathematica*, 182(1):105–142, 1999.
- [HH96] Ryszard Horodecki and Michał Horodecki. Information-theoretic aspects of inseparability of mixed states. *Physical Review A*, 54(3):1838, 1996.
- [HLP20] Sean Hallgren, Eunou Lee, and Ojas Parekh. An approximation algorithm for the max-2-local hamiltonian problem. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2020)*. Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2020.
- [HM17] Aram W Harrow and Ashley Montanaro. Extremal eigenvalues of local hamiltonians. *Quantum*, 1:6, 2017.
- [JPJR14] Sania Jevtic, Matthew Pusey, David Jennings, and Terry Rudolph. Quantum steering ellipsoids. *Physical review letters*, 113(2):020402, 2014.
- [KKR06] Julia Kempe, Alexei Kitaev, and Oded Regev. The complexity of the local hamiltonian problem. *SIAM Journal on Computing*, 35(5):1070–1097, 2006.

- [Kri77] Jean-Louis Krivine. Sur la constante de grothendieck. *CR Acad. Sci. Paris Ser. AB*, 284(8):A445–A446, 1977.
- [KSVV02] Alexei Yu Kitaev, Alexander Shen, Mikhail N Vyalyi, and Mikhail N Vyalyi. *Classical and Quantum Computation*. Number 47. American Mathematical Soc., 2002.
- [KŻ01] Marek Kuś and Karol Życzkowski. Geometry of entangled states. *Physical Review A*, 63(3):032307, 2001.
- [LaSt72] Nikolai N. Lebedev (author) and Richard Silverman (translator). *Special Functions and their Applications*. Courier Corporation, 1972.
- [Liu06] Yi-Kai Liu. Consistency of local density matrices is qma-complete. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, pages 438–449. Springer, 2006.
- [LLZ02] Michael Lewin, Dror Livnat, and Uri Zwick. Improved rounding techniques for the max 2-sat and max di-cut problems. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 67–82. Springer, 2002.
- [NC10] Michael A. Nielsen and Isaac L. Chuang. *Quantum Computation and Quantum Information: 10th Anniversary Edition*. Cambridge University Press, 2010.
- [O’D14] Ryan O’Donnell. *Analysis of Boolean Functions*. Cambridge University Press, 2014.
- [Osb12] Tobias J Osborne. Hamiltonian complexity. *Reports on Progress in Physics*, 75(2):022001, 2012.
- [PM15] Stephen Piddock and Ashley Montanaro. The complexity of antiferromagnetic interactions and 2d lattices. *arXiv preprint arXiv:1506.04014*, 2015.
- [PMS19] Joanna Piotrowska, Jonah M Miller, and Erik Schnetter. Spectral methods in the presence of discontinuities. *Journal of Computational Physics*, 390:527–547, 2019.
- [PNA10] Stefano Pironio, Miguel Navascués, and Antonio Acin. Convergent relaxations of polynomial optimization problems with noncommuting variables. *SIAM Journal on Optimization*, 20(5):2157–2180, 2010.
- [PWZ97] Marko Petkovsek, Herbert Wilf, and Doron Zeilberger.  $A=B$ . 1997. Available at <https://www.math.upenn.edu/~wilf/AeqB.html>.
- [RT12] Prasad Raghavendra and Ning Tan. Approximating csps with global cardinality constraints using sdp hierarchies. In *Proceedings of the twenty-third annual ACM-SIAM symposium on Discrete Algorithms*, pages 373–387. SIAM, 2012.
- [Rud64] Walter Rudin. *Principles of Mathematical Analysis*, volume 3. McGraw-hill New York, 1964.
- [SHH11] Christoph Spengler, Marcus Huber, and Beatrix C Hiesmayr. A geometric comparison of entanglement and quantum nonlocality in discrete systems. *Journal of Physics A: Mathematical and Theoretical*, 44(6):065304, 2011.



- [SV09] Norbert Schuch and Frank Verstraete. Computational complexity of interacting electrons and fundamental limitations of density functional theory. *Nature Physics*, 5(10):732–735, 2009.
- [Vaz13] Vijay V Vazirani. *Approximation Algorithms*. Springer Science & Business Media, 2013.
- [WB03] Pawel Wocjan and Thomas Beth. The 2-local hamiltonian problem encompasses np. *International Journal of Quantum Information*, 1(03):349–357, 2003.
- [WS11] David P Williamson and David B Shmoys. *The Design of Approximation Algorithms*. Cambridge university press, 2011.

## Appendix

The Appendix covers a range of results. First we present connections between classical constraint satisfaction problems and quantum local Hamiltonian problems. Next, we list some combinatorial identities (Appendix B) used in this work. These identities are all standard themselves, or follow from standard techniques [PWZ97]. Third, we cover some calculations relevant to the Hermite expansions used in the paper (Appendix C). In the paper we were primarily concerned with the 3-dimensional function  $f$  (Equation (53)), but here we give Hermite coefficients for the analogous  $r$ -dimensional function for arbitrary  $r \in \mathbb{Z}_{\geq 0}$ . This calculation allows us to reproduce technical tools from another work [Bri11]. Then, in Appendix D, we give our calculus-based lower bounds for the functions of interest. The following section (Appendix E) concerns some proofs for the properties of quantum states/projectors we used in the paper. Lastly, in Appendix F we give an approximation algorithm for the bipartite, traceless 2-Local Hamiltonian problem which achieves a novel approximation factor.

### A Connecting 2-Local Hamiltonians and Classical 2-CSPs

Here we motivate why 2-Local Hamiltonian is a generalization of classical 2-CSPs. We start with the Pauli matrices as defined in Equation (1) in Section 2.1. We will use the notions of a qubit and classical Boolean variable interchangeably and assume we have  $n$  such objects. Here we favor different notation than in Section 2.1 to better emphasize the connection between boolean polynomials and Hamiltonians.

We use the notation  $P_i$  to denote a Pauli matrix  $P \in \{X = \sigma^1, Y = \sigma^2, Z = \sigma^3\}$  acting on qubit  $i$ , i.e.  $P_i := \mathbb{I} \otimes \mathbb{I} \otimes \dots \otimes P \otimes \dots \otimes \mathbb{I} \in \mathbb{C}^{2^n \times 2^n}$ , where the  $P$  occurs at position  $i$ . As an example comparing notation, for  $n = 4$ ,  $X_1 Y_3 = \sigma_1^1 \otimes \sigma_3^2 \otimes \mathbb{I}_{2,4} = X \otimes \mathbb{I} \otimes Y \otimes \mathbb{I}$ . This notation will allow us to avoid explicitly using tensor-product notation, which is replaced with matrix multiplication; moreover, it will allow us to express Hermitian matrices as multilinear polynomials.

For  $n = 2$ , consider the matrix  $C = \frac{1}{2}(\mathbb{I} - Z_1 Z_2) = |01\rangle\langle 01| + |10\rangle\langle 10|$ :

$$C = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad Q = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Finding a maximum eigenvector of  $C$  corresponds to solving the classical 2-CSP  $\text{Max} \frac{1}{2}(1 - z_1 z_2)$  over  $z_1, z_2 \in \{\pm 1\}$ . This is because  $C$  is diagonal matrix with Boolean entries, so that maximum eigenvectors correspond to  $|b_1 b_2\rangle$  where the corresponding diagonal entry of  $C$  (i.e.,  $\langle b_1 b_2 | C | b_1 b_2 \rangle$ )

is 1. If we associate  $|0\rangle$  with the value 1 and  $|1\rangle$  with -1, then  $\frac{1}{2}(1 - z_1 z_2)$  is the Boolean indicator function for the diagonal of  $C$ . More generally, if we are given a *density matrix*  $\rho \in \mathbb{C}^{4 \times 4}$ , satisfying  $\rho \succeq 0$  and  $\text{Tr}[\rho] = 1$ , then  $\text{Tr}[C\rho] \in [0, 1]$  may be viewed as the expected value of  $\frac{1}{2}(1 - z_1 z_2)$  over a probability distribution specified by the diagonal of  $\rho$ . In this way, any rank  $r$  projector in  $\mathbb{C}^{4 \times 4}$  corresponds to Boolean clause on 2 variables, where  $r$  indicates the number of satisfying assignments. Such a clause may be extended to act on variables  $i$  and  $j$  among  $n$  variables by taking  $C_{ij} \otimes \mathbb{I}_{[n] \setminus \{i, j\}}$  ( $C_{ij}$  denotes that  $C$  acts on qubits  $i$  and  $j$ ). Summing over such extended clauses yields a 2-local Hamiltonian instance corresponding to a Boolean 2-CSP. Theorem 39 demonstrates how Max 2-QSAT generalizes Max 2-SAT.

Multilinear polynomials over  $\mathbb{I}$  and  $Z_i$  of degree  $k$  are diagonal  $k$ -local Hamiltonians encoding classical  $k$ -CSPs, while multilinear polynomials over  $\mathbb{I}, X_i, Y_i, Z_i$  of degree  $k$  are more general non-diagonal  $k$ -local Hamiltonians that are able to represent quantum phenomena.

We obtain a quantum generalization of Boolean 2-CSPs by considering non-diagonal matrices (in the standard basis). Let  $Q = \frac{1}{4}(\mathbb{I} - X_1 X_2 - Y_1 Y_2 - Z_1 Z_2)$  as depicted above. Here  $C$  represents Max Cut on a single edge, while  $Q$  is a quantum generalization of Max Cut [GP19]. The unique maximum eigenvector of  $Q$  is  $|\psi\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$ , with corresponding density matrix  $\rho = |\psi\rangle\langle\psi|$ , achieving an eigenvalue of  $\text{Tr}[Q\rho] = 1$ . This is a non-classical solution, which in general may require an exponential amount of information for representation. In contrast *product states*, which exhibit no quantum entanglement, may be represented as

$$\rho = \prod_i \frac{1}{2}(\mathbb{I} + \alpha_i X_i + \beta_i Y_i + \gamma_i Z_i).$$

Such  $\rho$  has  $\text{Tr}[\rho] = 1$  by construction, and  $\rho \succeq 0$  is guaranteed if  $\alpha_i^2 + \beta_i^2 + \gamma_i^2 \leq 1$  for all  $i$ . A product state maximizing  $\text{Tr}[Q\rho]$  is  $\rho = \frac{1}{4}(\mathbb{I} + Z_1)(\mathbb{I} - Z_2) = |01\rangle\langle 01|$ , achieving a value of  $\frac{1}{2}$ . Thus for  $Q$  there is a gap of  $\frac{1}{2}$  between the best product state and the best arbitrary quantum density matrix.

## B Identities

We will make use of several identities concerning the generalized Hypergeometric function. The following standard technique will be used in proving our combinatorial identities:

**Theorem 34** ([PWZ97]). 1. Let  $Q = \sum_{m=0}^p t_m$  be some finite sum. If

$$\frac{t_{m+1}}{t_m} = \frac{(m+a)(m+b)}{(m+c)(m+1)} \quad (80)$$

for  $m = \{0, \dots, p-1\}$ , then  $Q = t_0 {}_2F_1 \left[ \begin{matrix} a, b \\ c \end{matrix}; 1 \right]$ .

2. Let  $Q = \sum_{m=0}^{\infty} t_m x^m$ . If Equation (80) holds for all  $m$ , then  $Q = t_0 {}_2F_1 \left[ \begin{matrix} a, b \\ c \end{matrix}; x \right]$ .

Now, we list several identities that will be used in proving the results in the paper:

**Lemma 35.**

1.

$$\int_0^\pi \sin^a t \cos^b t dt = \begin{cases} \frac{2(a-1)!(b-1)!!}{(b+a)!!} & a \text{ is odd, } b \text{ is even} \\ \frac{\pi(a-1)!(b-1)!!}{(b+a)!!} & a \text{ is even, } b \text{ is even} \end{cases} \quad (81)$$

2.

$$\int_0^{2\pi} \sin^a t \cos^b t dt = \frac{2\pi(a-1)!(b-1)!!}{(a+b)!!} \text{ for } a \text{ and } b \text{ even.} \quad (82)$$

3. [AS48] If  $n$  is an integer:

$$\Gamma(n+1/2) = \sqrt{\pi} \frac{(2n-1)!!}{2^n}. \quad (83)$$

4. [PWZ97] (Gauss's Hypergeometric Theorem) If  $c > a + b$  (and  $a, b, c \in \mathbb{R}$ ):

$${}_2F_1 \left[ \begin{matrix} a, b \\ c \end{matrix}; 1 \right] = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}. \quad (84)$$

5. If  $r$  is odd:

$$\sum_{m=0}^p \frac{(-1)^m (2m+r-1)!!}{(2m+r)!!} \binom{p}{m} = \frac{(r-1)!!}{(2p+r)(2p+r-2)\dots(2p+1)}. \quad (85)$$

6. Generalized Vandermonde's identity:

$$\sum_{k_1+\dots+k_p=m} \binom{n_1}{k_1} \binom{n_2}{k_2} \dots \binom{n_p}{k_p} = \binom{n_1+\dots+n_p}{m}. \quad (86)$$

7.

$$\sum_{q=0}^{\infty} a^{2q+1} \frac{(2q+1)!}{2^{2q+1}(1+2q)^2(3+2q)(q!)^2} = \frac{a}{6} {}_2F_1 \left[ \begin{matrix} 1/2, 1/2 \\ 5/2 \end{matrix}; a^2 \right]. \quad (87)$$

*Proof.* Equation (81) and Equation (82) are easy to establish with integration by parts and induction, and Equation (86) follows from a simple counting argument. To prove Equation (85) we use Theorem 34, item 1. Let

$$\sum_{m=0}^p \frac{(-1)^m (2m+r-1)!!}{(2m+r)!!} \binom{p}{m} = \sum_{m=0}^p t_m. \quad (88)$$

We can calculate:

$$\frac{t_{m+1}}{t_m} = \frac{(m-p)(m+(r+1)/2)}{(m+(r+2)/2)(m+1)}. \quad (89)$$

Theorem 34 then implies that:

$$\sum_{m=0}^p t_m = t_0 {}_2F_1 \left[ \begin{matrix} -p, (r+1)/2 \\ (r+2)/2 \end{matrix}; 1 \right] = t_0 \frac{\Gamma((r+2)/2)\Gamma(p+1/2)}{\Gamma((r+2)/2+p)\Gamma(1/2)}. \quad (90)$$

where we applied Equation (84) for the last equality. Now apply Equation (83) and our assumption that  $r$  is odd to obtain:

$$\sum_{m=0}^p t_m = t_0 \frac{r!!(2p-1)!!}{(r+2p)!!} = \frac{(r-1)!!}{(2p+r)(2p+r-2)\dots(2p+1)}. \quad (91)$$

We prove Equation (87) in the same way. In the context of Theorem 34, item 2 we can take write:

$$\sum_{q=0}^{\infty} a^{2q+1} \frac{(2q+1)!}{2^{2q+1}(1+2q)^2(3+2q)(q!)^2} = a \sum_{q=0}^{\infty} t_q (a^2)^q, \quad (92)$$

and calculate:

$$\frac{t_{q+1}}{t_q} = \frac{\frac{(2(q+1)+1)!}{2^{2(q+1)+1}(1+2(q+1))^2(3+2(q+1))(q+1)!^2}}{\frac{(2q+1)!}{2^{2q+1}(1+2q)^2(3+2q)(q!)^2}} = \frac{(q+1/2)^2}{(q+5/2)(q+1)}, \quad (93)$$

where  $t_0 = a/6$ . Equation (87) follows.  $\square$

## C Calculation of Hermite Coefficients

We will make use of these identities to determine the Hermite coefficients for our function of interest. First, let us compute the moments:

**Lemma 36.** *Let  $\mathbf{z} \sim \mathcal{N}(0, \mathbb{I})$ . If  $b_1, \dots, b_r$  are even,*

$$\mathbb{E} \left[ \frac{z_1^{b_1} z_2^{b_2} \dots z_r^{b_r}}{\sqrt{z_1^2 + \dots + z_r^2}} \right] = c_r \frac{(b_1 + \dots + b_r + r - 3)!! (b_1 - 1)!! (b_2 - 1)!! \dots (b_r - 1)!!}{(b_1 + b_2 + \dots + b_r + r - 2)!!}, \quad (94)$$

where  $c_r$  is  $\sqrt{2/\pi}$  if  $r$  is odd and  $\sqrt{\pi/2}$  if  $r$  is even.

*Proof.* From the definition of the multivariate Gaussian,

$$\xi := \mathbb{E} \left[ \frac{z_1^{b_1} z_2^{b_2} \dots z_d^{b_d}}{\sqrt{z_1^2 + z_2^2 + \dots + z_d^2}} \right] = \int \frac{z_1^{b_1} z_2^{b_2} \dots z_d^{b_d}}{\sqrt{z_1^2 + z_2^2 + \dots + z_d^2}} \left( \frac{1}{(2\pi)^{d/2}} e^{-\|z\|^2/2} \right) dz.$$

Let us re-write this using generalized spherical coordinates:

$$\begin{aligned} \xi = & \frac{1}{(2\pi)^{d/2}} \int \frac{1}{r} \left( r \cos \phi_1 \right)^{b_1} \left( r \sin \phi_1 \cos \phi_2 \right)^{b_2} \left( r \sin \phi_1 \sin \phi_2 \cos \phi_3 \right)^{b_3} \dots \left( r \sin \phi_1 \right. \\ & \left. \dots \sin \phi_{d-3} \cos \phi_{d-2} \right)^{b_{d-2}} \left( r \sin \phi_1 \dots \sin \phi_{d-2} \cos \phi_{d-1} \right)^{b_{d-1}} \left( r \sin \phi_1 \dots \sin \phi_{d-2} \sin \phi_{d-1} \right)^{b_d} \\ & e^{-r^2/2} \left( r^{d-1} \sin^{d-2} \phi_1 \sin^{d-3} \phi_2 \dots \sin \phi_{d-2} \right) dr d\phi_1 \dots d\phi_{d-1}, \end{aligned}$$

where  $\phi_1, \dots, \phi_{d-2} \in [0, \pi]$  and  $\phi_{d-1} \in [0, 2\pi]$ . Now we can break it up into smaller integrals:

$$\begin{aligned} \xi = & \frac{1}{(2\pi)^{d/2}} \int_0^\infty r^{b_1+b_2+\dots+b_d+d-2} e^{-r^2/2} dr \int_0^\pi \cos^{b_1} \phi_1 (\sin \phi_1)^{b_2+\dots+b_d+d-2} d\phi_1 \\ & \int_0^\pi \cos^{b_2} \phi_2 (\sin \phi_2)^{b_3+\dots+b_d+d-3} d\phi_2 \int_0^\pi \cos^{b_3} \phi_3 (\sin \phi_3)^{b_4+\dots+b_d+d-4} d\phi_3 \\ & \dots \int_0^\pi \cos^{b_{d-3}} \phi_{d-3} (\sin \phi_{d-3})^{b_{d-2}+b_{d-1}+b_d+2} d\phi_{d-3} \int_0^\pi \cos^{b_{d-2}} \phi_{d-2} (\sin \phi_{d-2})^{b_{d-1}+b_d+1} d\phi_{d-2} \\ & \cdot \int_0^{2\pi} \cos^{b_{d-1}} \phi_{d-1} \sin^{b_d} \phi_{d-1} d\phi_{d-1}. \end{aligned}$$

Now we will apply Equation (81) and Equation (82). There are two cases here depending on if  $d$  is even or odd. If  $d$  is even:

$$\xi = \frac{1}{(2\pi)^{d/2}} \left[ 2^{(b_1+\dots+b_d+d-3)/2} \Gamma\left(\frac{b_1+\dots+b_d+d-1}{2}\right) \left[ \frac{\pi(b_1-1)!(b_2+\dots+b_d+d-3)!!}{(b_1+b_2+\dots+b_d+d-2)!!} \right] \right. \\ \left. \left[ \frac{2(b_2-1)!(b_3+\dots+b_d+d-4)!!}{(b_2+b_3+\dots+b_d+d-3)!!} \right] \left[ \frac{\pi(b_3-1)!(b_4+\dots+b_d+d-5)!!}{(b_3+\dots+b_d+d-4)!!} \right] \right. \\ \dots \left. \left[ \frac{\pi(b_{d-3}-1)!(b_{d-2}+b_{d-1}+b_d+1)!!}{(b_{d-3}+\dots+b_d+2)!!} \right] \left[ \frac{2(b_{d-2}-1)!(b_{d-1}+b_d)!!}{(b_{d-2}+b_{d-1}+b_d+1)!!} \right] \right. \\ \left. \left[ \frac{2\pi(b_{d-1}-1)!(b_d-1)!!}{(b_{d-1}+b_d)!!} \right] \right.$$

Note that the product telescopes. Since  $d$  is even, we can count a factor  $(2\pi)^{(d-2)/2+1} = (2\pi)^{d/2}$ . Continuing,

$$\xi = \left[ 2^{(b_1+\dots+b_d+d-3)/2} \Gamma\left(\frac{b_1+\dots+b_d+d-1}{2}\right) \right] \frac{(b_1-1)!(b_2-1)!\dots(b_d-1)!!}{(b_1+b_2+\dots+b_d+d-2)!!} \\ = \sqrt{\pi} 2^{(b_1+\dots+b_d+d-3)/2} \frac{(b_1+\dots+b_d+d-3)!!(b_1-1)!(b_2-1)!\dots(b_d-1)!!}{2^{(b_1+\dots+b_d+d-2)/2} (b_1+b_2+\dots+b_d+d-2)!!} \\ = \sqrt{\frac{\pi}{2}} \frac{(b_1+\dots+b_d+d-3)!!(b_1-1)!(b_2-1)!\dots(b_d-1)!!}{(b_1+b_2+\dots+b_d+d-2)!!}.$$

In the odd case, the constant factor we get is  $2(2\pi)^{(d-3)/2+1}$ , so the overall constant is  $\sqrt{2/\pi}$ . We end up getting the same thing with a different constant:

$$\xi = \sqrt{\frac{2}{\pi}} \frac{(b_1+\dots+b_d+d-3)!!(b_1-1)!(b_2-1)!\dots(b_d-1)!!}{(b_1+b_2+\dots+b_d+d-2)!!}.$$

□

Given these moments we can explicitly calculate the Hermite coefficients:

**Lemma 37.** *Let  $f(\mathbf{z}) = z_1/\sqrt{z_1^2+\dots+z_r^2}$  for  $r$  odd, and let it have Hermite expansion  $f(\mathbf{z}) = \sum_{\mu} \hat{f}_{\mu} h_{\mu}(\mathbf{z})$ . Then, if  $\mu_1$  is odd and  $\mu_i$  is even for all  $i \in \{2, \dots, r\}$ ,*

$$\hat{f}_{\mu} = \sqrt{\frac{2}{\pi}} \frac{\sqrt{\mu_1!\dots\mu_r!}(-1)^p}{(\mu_1-1)!!\mu_2!!\dots\mu_r!!} \frac{(r-1)!!}{(2p+r)(2p+r-2)\dots(2p+1)} \quad (95)$$

where  $p = (\mu_1 + \dots + \mu_r - 1)/2$ . Otherwise,  $\hat{f}_{\mu} = 0$ .

*Proof.* The fact that  $\hat{f}_{\mu} = 0$  when  $\mu_1$  is even or at least one of  $\mu_i$  is odd for  $i \in \{2, \dots, r\}$  follows from the same idea in the paper, i.e. the integral of an anti-symmetric function is zero. For the other values of  $\mu$ , use Definition 19 to obtain:

$$\hat{f}_{\mu} = \mathbb{E} \left[ \frac{z_1}{\sqrt{z_1^2+\dots+z_r^2}} \frac{\sqrt{\mu_1!}}{2^{\mu_1/2}} \sum_{l_1=0}^{(\mu_1-1)/2} \frac{(-1)^{(\mu_1-1)/2-l_1} (\sqrt{2}z_1)^{2l_1+1}}{(2l_1+1)! \left(\frac{\mu_1-1}{2} - l_1\right)!} \frac{\sqrt{\mu_2!}}{2^{\mu_2/2}} \sum_{l_2=0}^{\mu_2/2} \frac{(-1)^{\mu_2/2-l_2} (\sqrt{2}z_2)^{2l_2}}{(2l_2)! \left(\frac{\mu_2}{2} - l_2\right)!} \right. \\ \left. \dots \frac{\sqrt{\mu_r!}}{2^{\mu_r/2}} \sum_{l_r=0}^{\mu_r/2} \frac{(-1)^{\mu_r/2-l_r} (\sqrt{2}z_r)^{2l_r}}{(2l_r)! \left(\frac{\mu_r}{2} - l_r\right)!} \right]$$

$$= \frac{\sqrt{\mu_1! \dots \mu_r!}}{2^{(\mu_1 + \dots + \mu_r)/2}} \sum_{l_1, \dots, l_r} \frac{(-1)^{(\mu_1 + \dots + \mu_r - 1)/2 - (l_1 + \dots + l_r)} \sqrt{2}^{2(l_1 + \dots + l_r) + 1}}{[(2l_1 + 1)!(2l_2)! \dots (2l_r)!] \left[ \left( \frac{\mu_1 - 1}{2} - l_1 \right)! (\mu_2/2 - l_2)! \dots (\mu_r/2 - l_r)! \right]} \cdot \mathbb{E} \left[ \frac{z_1^{2l_1 + 2} z_2^{2l_2} \dots z_r^{2l_r}}{\sqrt{z_1^2 + \dots + z_r^2}} \right].$$

By Lemma 36,

$$\begin{aligned} \hat{f}_\mu &= \frac{\sqrt{\mu_1! \dots \mu_r!}}{2^{(\mu_1 + \dots + \mu_r)/2}} \sum_{l_1, \dots, l_r} \frac{(-1)^{(\mu_1 + \dots + \mu_r - 1)/2 - (l_1 + \dots + l_r)} \sqrt{2}^{2(l_1 + \dots + l_r) + 1}}{[(2l_1 + 1)!(2l_2)! \dots (2l_r)!] \left[ \left( \frac{\mu_1 - 1}{2} - l_1 \right)! (\mu_2/2 - l_2)! \dots (\mu_r/2 - l_r)! \right]} \\ &\quad \cdot c_r \frac{(2l_1 + 2 + 2l_2 + \dots + 2l_r + r - 3)!! (2l_1 + 1)!! (2l_2 - 1)!! \dots (2l_r - 1)!!}{(2l_1 + 2 + 2l_2 + \dots + 2l_r + r - 2)!!} \\ &= c_r \frac{\sqrt{\mu_1! \dots \mu_r!}}{2^{(\mu_1 + \dots + \mu_r)/2}} (-1)^{(\mu_1 + \dots + \mu_r - 1)/2} \sum_{l_i} \frac{\sqrt{2}^{2(l_1 + \dots + l_r) + 1} (2l_1 + \dots + 2l_r + r - 1)!! (-1)^{l_1 + \dots + l_r}}{(2l_1)!! (2l_2)!! \dots (2l_r)!! \left( \frac{\mu_1 - 1}{2} - l_1 \right)! (\mu_2/2 - l_2)! \dots (\mu_r/2 - l_r)!} \\ &\quad \cdot \frac{1}{(2l_1 + \dots + 2l_r + r)!!}. \end{aligned} \tag{96}$$

Now let us apply the identity:

$$(2l)!!(q - l)! = 2^l l!(q - l)! = 2^l \frac{q!}{\binom{q}{l}},$$

to rewrite Equation (96) as:

$$\begin{aligned} \hat{f}_\mu &= \frac{\sqrt{2} c_r \sqrt{\mu_1! \dots \mu_r!}}{2^{(\mu_1 + \dots + \mu_r)/2}} (-1)^{(\mu_1 + \dots + \mu_r - 1)/2} \sum_{l_i} \frac{2^{l_1 + \dots + l_r} (2l_1 + \dots + 2l_r + r - 1)!! \binom{(\mu_1 - 1)/2}{l_1} \binom{\mu_2/2}{l_2} \dots \binom{\mu_r/2}{l_r}}{2^{l_1 + \dots + l_r} \left( \frac{\mu_1 - 1}{2} \right)! (\mu_2/2)! \dots (\mu_r/2)! (2l_1 + \dots + 2l_r + r)!!} \\ &= \frac{\sqrt{2} c_r \sqrt{\mu_1! \dots \mu_r!} (-1)^{(\mu_1 + \dots + \mu_r - 1)/2}}{2^{(\mu_1 + \dots + \mu_r)/2} \left( \frac{\mu_1 - 1}{2} \right)! (\mu_2/2)! \dots (\mu_r/2)!} \sum_{m=0}^{(\mu_1 + \dots + \mu_r - 1)/2} (-1)^m \frac{(2m + r - 1)!!}{(2m + r)!!} \\ &\quad \cdot \sum_{l_1 + \dots + l_r = m} \binom{(\mu_1 - 1)/2}{l_1} \binom{\mu_2/2}{l_2} \dots \binom{\mu_r/2}{l_r}. \end{aligned}$$

Now apply the generalized Vandermonde's equality, Equation (86):

$$\hat{f}_\mu = \frac{\sqrt{2} c_r \sqrt{\mu_1! \dots \mu_r!} (-1)^{(\mu_1 + \dots + \mu_r - 1)/2}}{2^{(\mu_1 + \dots + \mu_r)/2} \left( \frac{\mu_1 - 1}{2} \right)! (\mu_2/2)! \dots (\mu_r/2)!} \sum_m (-1)^m \frac{(2m + r - 1)!!}{(2m + r)!!} \binom{(\mu_1 - 1)/2 + \mu_2/2 + \dots + \mu_r/2}{m}.$$

Let us set  $p = (\mu_1 - 1)/2 + \mu_2/2 + \dots + \mu_r/2$ . Using Equation (85),

$$\begin{aligned} \hat{f}_\mu &= \frac{\sqrt{2} c_r \sqrt{\mu_1! \dots \mu_r!} (-1)^p}{2^{(\mu_1 + \dots + \mu_r)/2} \left( \frac{\mu_1 - 1}{2} \right)! (\mu_2/2)! \dots (\mu_r/2)!} \frac{(r - 1)!!}{(2p + r)(2p + r - 2) + \dots + (2p + 1)} \\ &= \frac{\sqrt{2} c_r \sqrt{\mu_1! \dots \mu_r!} (-1)^p (r - 1)!!}{2^{(\mu_1 + \dots + \mu_r)/2} (\mu_1 - 1)!! (\mu_2)!! \dots (\mu_r)!! 2^{-p} (2p + r) \dots (2p + 1)} \\ &= \frac{c_r \sqrt{\mu_1! \dots \mu_r!} (-1)^p (r - 1)!!}{(\mu_1 - 1)!! \mu_2!! \dots \mu_r!! (2p + r) \dots (2p + 1)}. \end{aligned}$$

Note that  $c_r = \sqrt{2/\pi}$  since  $r$  is odd. □

We can do a ‘‘sanity check’’ for the calculations presented thus far by giving a proof sketch of a specialization of our result which yields a result from [Bri11]:

**Proposition 38.**

$$\sum_{i,j,k} \hat{f}_{i,j,k}^2 a^{i+j+k} = \frac{8}{3\pi} {}_2F_1 \left[ \begin{matrix} 1/2, 1/2 \\ 5/2 \end{matrix}; a^2 \right]$$

*Proof. (sketch)* It is sufficient to collect terms  $\hat{f}_{i,j,k}$  with fixed  $i + j + k$  and compare them to the Taylor series given in the statement of Lemma 5.2.1. from [Bri11]. For this we must demonstrate that:

$$\sum_{i+j+k=2l+1} \frac{i!j!k!}{[(i-1)!!j!!k!!]^2} = \frac{1}{3} \frac{(2l-1)!!(1+2l)(3+2l)}{(2l)!!},$$

where we have canceled some common factors. This can be rewritten as:

$$= \sum_{i=[0:2:2l]} \left( \frac{(i+1)!}{(i!!)^2} \sum_{j=[0:2:2l-i]} \frac{j!(2l-i-j)!}{[j!!(2l-i-j)!!]^2} \right),$$

where we have redefined  $i$  and used the (MATLAB) notation  $i = [0 : 2 : 2l]$  to indicate a sum from 0 to  $2l$  by 2. The second sum can be proven to be 1 using Theorem 34 and Gauss’s theorem (Equation (84)). The remainder of the proof follows from the identity

$$\sum_{q=0}^l \frac{(2q+1)!}{[(2q)!!]^2} = \frac{1}{3} \frac{(2l+3)!!}{(2l)!!},$$

which is simple to establish with induction. □

*Proof of Lemma 24.* We have two functions of interest here:

$$g(z) = \text{sign}(z) = \sum_i \hat{g}_i h_i(z)$$

$$\text{and } f(z_1, z_2, z_3) = \frac{z_1}{\sqrt{z_1^2 + z_2^2 + z_3^2}} = \sum_{i,j,k} \hat{f}_{i,j,k} h_i(z_1) h_j(z_2) h_k(z_3).$$

for  $\{\hat{g}_i\}$  and  $\{\hat{f}_{i,j,k}\}$  Hermite expansions given in Lemma 22 and Lemma 23. We want:

$$\mathbb{E}[g(z)f(z_1, z_2, z_3)] = \sum_{i,j,k} \hat{f}_{i,j,k} \hat{g}_i \mathbb{E}[h_i(z_1)h_i(z)] \mathbb{E}[h_j(z_2)] \mathbb{E}[h_k(z_3)] = \sum_{i \text{ odd}} \hat{f}_{i,00} \hat{g}_i a^i,$$

where in the second equality we used that fact that  $\delta_{0j} = \mathbb{E}[h_0(z_2)h_j(z_2)]$ . If we apply the identity  $(i-1)!! = [(i-1)/2]! 2^{\frac{i-1}{2}}$  and use the previously derived values of  $\hat{f}_{i,00}$  and  $\hat{g}_i$ , the sum is:

$$\begin{aligned} \sum_{i \text{ odd}} a^i \hat{g}_i \hat{f}_{i,00} &= \sum_{i \text{ odd}} a^i \left( \frac{\sqrt{i!}(-1)^q}{2^{i/2}} \frac{2}{\sqrt{\pi}q!(1+2q)} \right) \left( \frac{4}{\sqrt{\pi}} \frac{\sqrt{i!}}{2^{(i)/2}} \frac{(-1)^q}{\left(\frac{i-1}{2}\right)!(1+2q)(3+2q)} \right) \\ &= \frac{8}{\pi} \sum_{i \text{ odd}} a^i \frac{i!}{2^i(1+2q)^2(3+2q)(q!)^2}. \end{aligned}$$

where  $q = (i - 1)/2$ . We can equivalently sum over  $q$  to get a sum from 0 to infinity by 1, (i.e. not every other integer):

$$\begin{aligned} &= \frac{8}{\pi} \sum_{q=0}^{\infty} a^{2q+1} \frac{(2q+1)!}{2^{2q+1}(1+2q)^2(3+2q)(q!)^2} = \frac{8a}{\pi 6} {}_2F_1 \left[ \begin{matrix} 1/2, 1/2 \\ 5/2 \end{matrix}; a^2 \right] \\ &= \frac{4a}{3\pi} {}_2F_1 \left[ \begin{matrix} 1/2, 1/2 \\ 5/2 \end{matrix}; a^2 \right], \end{aligned}$$

where we used Equation (87). □

We also present a proof of this fact for completeness:

*Proof of Proposition 21, Item 3.*

$$\begin{aligned} &\left| \mathbb{E}[f(\mathbf{z})f(\mathbf{z}')] - \sum_{i,j,k \leq N} \hat{f}_{i,j,k}^2 a^i b^j c^k \right|^2 \tag{97} \\ &= \left| \mathbb{E} \left[ \left( f(\mathbf{z}) - \sum_{i,j,k \leq N} \hat{f}_{i,j,k} h_i(\mathbf{z}_1) h_j(\mathbf{z}_2) h_k(\mathbf{z}_3) \right) \left( f(\mathbf{z}') + \sum_{i,j,k \leq N} \hat{f}_{i,j,k} h_i(\mathbf{z}'_1) h_j(\mathbf{z}'_2) h_k(\mathbf{z}'_3) \right) \right] \right|^2 \\ &\leq \mathbb{E} \left[ \left( f(\mathbf{z}) - \sum_{i,j,k \leq N} \hat{f}_{i,j,k} h_i(\mathbf{z}_1) h_j(\mathbf{z}_2) h_k(\mathbf{z}_3) \right)^2 \right] \mathbb{E} \left[ \left( f(\mathbf{z}') + \sum_{i,j,k \leq N} \hat{f}_{i,j,k} h_i(\mathbf{z}'_1) h_j(\mathbf{z}'_2) h_k(\mathbf{z}'_3) \right)^2 \right] \end{aligned}$$

by Cauchy-Schwarz. The second term is:

$$\begin{aligned} &\mathbb{E} \left[ f(\mathbf{z}')^2 + 2 \sum_{i,j,k} \hat{f}_{i,j,k} h_i(\mathbf{z}'_1) h_j(\mathbf{z}'_2) h_k(\mathbf{z}'_3) f(\mathbf{z}') \right] + \sum_{i,j,k \leq N} \hat{f}_{i,j,k}^2 = \langle f, f \rangle + 3 \sum_{i,j,k \leq N} \hat{f}_{i,j,k}^2 \\ &\leq 4 \langle f, f \rangle. \end{aligned}$$

So to get the RHS of Equation (97) less than  $\epsilon$ , take  $N$  large enough that

$$\mathbb{E} \left[ \left( f(\mathbf{z}) - \sum_{i,j,k \leq N} \hat{f}_{i,j,k} h_i(\mathbf{z}_1) h_j(\mathbf{z}_2) h_k(\mathbf{z}_3) \right)^2 \right] \leq \frac{\epsilon}{4 \langle f, f \rangle}.$$

We know this can be done since  $\sum_{i,j,k \leq N} \hat{f}_{i,j,k} h_i(\mathbf{z}_1) h_j(\mathbf{z}_2) h_k(\mathbf{z}_3) \rightarrow f$  in this norm. □

## D Lower Bounds for the Expectation of an Edge

For the result we have described, we crucially needed several lower bounds on the expectation of a single edge. In this direction we present proofs of lemmas 30-32:



*Proof of Lemma 30. Rank 1 case* We will divide the region  $\mathcal{S}$  into two regions and use different order expansions in each region for a bound. Define:

$$\mathcal{S}_1 = \text{conv}\{-1/2, -1/2, -1\}, [-1/2, -1, -1/2], [-1, -1/2, -1/2], [-1, -1, -1]\},$$

and  $\mathcal{S}_2 = \mathcal{S} \setminus \mathcal{S}_1$ .

Let us apply Lemma 26 with  $Q = \{(1, 0, 0)\}$  to obtain

$$\frac{1 - \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a, b, c)}{1 - a - b - c} \geq \frac{1 - \hat{f}_{1,00}^2(a + b + c) - 3(1/3 - \hat{f}_{1,00}^2)}{1 - a - b - c}.$$

Define  $x = a + b + c$  and observe that the RHS is a function of  $x$ , so define

$$l(x) := \frac{1 - \hat{f}_{1,00}^2 x - 3(1/3 - \hat{f}_{1,00}^2)}{1 - x}.$$

It is easy to see that this function is increasing as a function of  $x$ , and that  $l(-2) \geq 0.47$ , so we have the lower bound on  $\mathcal{S}_2$ :

$$\min_{(a,b,c) \in \mathcal{S}_2} \frac{1 - \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a, b, c)}{1 - a - b - c} \geq 0.47. \quad (98)$$

It remains to bound the function over  $\mathcal{S}_1$ . For this we will need a higher order Hermite expansion to get a good approximation factor. Let us apply Lemma 26 with  $Q = \{(1, 0, 0), (1, 0, 2), (3, 0, 0)\}$  to obtain:

$$\frac{1 - \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a, b, c)}{1 - a - b - c} \geq \left( 1 - \hat{f}_{1,00}^2 p_{1,00}(a, b, c) - \hat{f}_{1,02}^2 p_{1,02}(a, b, c) - \hat{f}_{3,00}^2 p_{3,00}(a, b, c) \right. \\ \left. - 3(1/3 - \hat{f}_{1,00}^2 - 2\hat{f}_{1,02}^2 - \hat{f}_{3,00}^2) \right) / (1 - a - b - c) =: q(a, b, c). \quad (99)$$

We will show that  $q(a, b, c)$  is minimized at  $[-1, -1, -1]$ , demonstrating the lemma. First calculate the partial derivative with respect to  $a$ :

$$\frac{\partial q}{\partial a} = \frac{4(115 + 6a^3 - b^2(1 + 2b) + 2a(-1 + b + c)(b + c) - c^2(1 + 2c) + a^2(-9 + 10b + 10c))}{225(-1 + a + b + c)^2 \pi}.$$

We can easily check that this is never 0 since the constant term is large enough that the numerator is always strictly positive:

$$115 + 6a^3 - b^2(1 + 2b) + 2a(-1 + b + c)(b + c) - c^2(1 + 2c) + a^2(-9 + 10b + 10c) \\ \geq 115 - 6 - (1 + 2) - 2(3)(2) - (1 + 2) + (-9 - 10 - 10) = 62.$$

Note that we applied the bounds  $|a| \leq 1$ ,  $|b| \leq 1$  and  $|c| \leq 1$ . Hence there are no internal critical points.

Let us now consider a horizontal line along one of the ‘‘vertical faces’’ (line 1 in Figure 1). One such a line has the form  $[a, b, c] = [s, s, -1] + t[0, -1, 1]$  for  $s$  some fixed constant  $s \in [-1/2, -1]$  and  $t$  some parameter that varies along the line between 0 and  $s + 1$ . We can calculate the derivative of  $q$  along this line to obtain

$$\frac{dq(s, s - t, -1 + t)}{dt} = -\frac{4(-4 + 5s)(1 + s - 2t)}{225\pi(-1 + s)}.$$




Figure 1: Relevant parameterizations for proof of Lemma 30, Region  $\mathcal{S}_1$

The only potential critical points here are in the middle of the line when  $t = (s + 1)/2$  and when  $s = 4/5$ . The later point is outside of our region of interest, so we can ignore it. For the first point it is clear we must examine the point which is in the middle of the line for each value of  $s$ . For this consider the line which goes down the “middle” of a vertical face (line 2 in Figure 1). This line can be parameterized as  $[a, b, c] = [-1, -1, -1] + t[1/2, 1/4, 1/4]$  for  $t$  between 0 and 1. We can once again evaluate the derivative of  $q$  along this line:

$$\frac{dq}{dt} = \frac{560 + t(344 + t(-109 + 11t))}{450\pi(-4 + t)^2}.$$

We can again see that the constant term is large enough that the numerator is always strictly positive, so the only possible critical points are the endpoints.

It remains to check the “top face”, line 3, and line 4. Observe that we can parameterize the top face as  $[a, b, -2 - a - b]$  under some suitable restriction of  $a$  and  $b$ . The gradient under this parameterization can be written as

$$\begin{aligned} \frac{\partial q(a, b, -2 - a - b)}{\partial a} &= \frac{8(2 + 2a + b)(8 + 3b)}{675\pi}, \\ \frac{\partial q(a, b, -2 - a - b)}{\partial b} &= \frac{8(8 + 3a)(2 + a + 2b)}{675\pi}. \end{aligned}$$

The only possible critical point is  $[-2/3, -2/3, -2/3]$ .

For the line 3 parameterize as  $[a, b, c] = [-1/2, -1/2, -1] + t[0, -1/2, 1/2]$  and calculate:

$$\frac{dq(-1/2, -1/2(1 + t), -1 + t/2)}{dt} = \frac{13(-1 + 2t)}{675\pi}.$$

Only critical point is in the middle of the line.

Lastly, we need to check line 4:  $[a, b, c] = [-1, -1, -1] + t[1, 1, 0]$ . The derivative is:

$$\frac{dq(-1 + t, -1 + t, -1)}{dt} = \frac{4(35 + 52t - 37t^2 + 8t^3)}{225\pi(-2 + t)^2}.$$

Finding the roots of the numerator (which can be determined analytically by a well known formula), it is clear there are no critical points along the line, so we only need to check its endpoints.

The previous analysis as well as the symmetries of the function  $q(a, b, c)$  imply that we need only check the following points to find a minimum:

$$\begin{aligned} q(-2/3, -2/3, -2/3) &\approx 0.508, \\ q(-1, -1, -1) &\approx 0.467, \\ q(-1/2, -1/2, -1) &\approx 0.510, \\ \text{and } q(-1/2, -3/4, -3/4) &\approx 0.509. \end{aligned}$$

The lemma follows. □

*Proof of Lemma 31.* As before let us lower bound this expectation using Lemma 27 with  $Q = \{(1, 0, 0), (1, 0, 2), (3, 0, 0)\}$ .

$$\frac{1 + \sum_{i,j \leq k} \hat{f}_{i,jk}^2 u_{i,jk}(a, b, c)}{1 + a} \geq \frac{1 + \hat{f}_{1,00}^2 a + \hat{f}_{1,02}^2 a(b^2 + c^2) + \hat{f}_{3,00}^2 a^3 - (1/3 - \hat{f}_{1,00}^2 - 2\hat{f}_{1,02}^2 - \hat{f}_{3,00}^2)}{1 + a}. \quad (100)$$

Since  $b^2 + c^2 \geq 0$ , we can uniformly lower bound this expression as:

$$\frac{1 + \hat{f}_{1,00}^2 a + \hat{f}_{3,00}^2 a^3 - (1/3 - \hat{f}_{1,00}^2 - 2\hat{f}_{1,02}^2 - \hat{f}_{3,00}^2)}{1 + a} =: l(a). \quad (101)$$

It is easy to check that the derivative is always negative, so the function is minimized at  $a = 1$ . The lemma follows since  $l(1) \approx 0.639$ . □

*Proof of Lemma 32.* This proof is very similar to the proof of the rank 1 case, however the function behaves a bit differently so different regions are needed. Let

$$\begin{aligned} \mathcal{S}_1 &= \text{conv}\{[-1, -1, -1], [r, r, -1], [r, -1, r], [-1, r, r]\}, \\ \text{and } \mathcal{S}_2 &= \mathcal{S} \setminus \mathcal{S}_1, \end{aligned}$$

where  $r = 0.6$  here and throughout the proof. Just as before, we will use a low order Hermite expansion for  $\mathcal{S}_1$  and a higher order expansion along with some tedious analysis for  $\mathcal{S}_2$ . Let us first apply Lemma 26 with  $Q = \{(1, 0, 0)\}$  to obtain:

$$\frac{3 + \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a, b, c)}{3 + a + b + c} \geq \frac{3 + \hat{f}_{1,00}^2(a + b + c) - 3(1/3 - \hat{f}_{1,00}^2)}{3 + a + b + c}.$$

The RHS is a function of  $x := a + b + c$ , so we define:

$$l(x) := \frac{3 + \hat{f}_{1,00}^2 x - 3(1/3 - \hat{f}_{1,00}^2)}{3 + x}.$$

In  $\mathcal{S}_1$   $x$  ranges from  $-3$  to  $2r - 1$ . It is easy to see  $l$  is decreasing as a function of  $x$ , so we can obtain a lower bound on  $\mathcal{S}_1$  of  $l(2r - 1) \approx 0.907$ .

Now we turn to a bound on the function in  $\mathcal{S}_2$ . First apply Lemma 26 with  $Q = \{(1, 0, 0), (1, 0, 2), (3, 0, 0)\}$  to obtain:

$$\begin{aligned} \frac{3 + \sum_{i,j \leq k} \hat{f}_{i,jk}^2 p_{i,jk}(a, b, c)}{3 + a + b + c} &\geq \left( 3 + \hat{f}_{1,00}^2 p_{1,00}(a, b, c) + \hat{f}_{1,02}^2 p_{1,02}(a, b, c) + \hat{f}_{3,00}^2 p_{3,00}(a, b, c) \right. \\ &\quad \left. - 3(1/3 - \hat{f}_{1,00}^2 - 2\hat{f}_{1,02}^2 - \hat{f}_{3,00}^2) \right) / \left( 3 + a + b + c \right) =: q(a, b, c). \end{aligned}$$




Figure 2: Relevant parameterizations for proof of Lemma 32, region  $\mathcal{S}_2$

First, let us examine *inside*  $\mathcal{S}_2$ . Just as before we can take a partial derivative:

$$\frac{\partial q}{\partial a} = \left( 2(-30 + 12a^3 + 6b^2 - 4b^3 + 6c^2 - 4c^3 + a^2(54 + 20b + 20c) + 4a(b^2 + c(3 + c) + b(3 + 2c)) - 225\pi) \right) / \left( 225(3 + a + b + c)^2\pi \right),$$

and check that the constant term has large enough magnitude that the numerator is always negative:

$$\left( 2(-30 + 12a^3 + 6b^2 - 4b^3 + 6c^2 - 4c^3 + a^2(54 + 20b + 20c) + 4a(b^2 + c(3 + c) + b(3 + 2c)) - 225\pi) \right) \leq \left( 2(-30 + 12 + 6 + 4 + 6 + 4 + (54 + 20 + 20) + 4(1 + (3 + 1) + (3 + 2)) - 225\pi) \right) \approx -1100.$$

So, the numerator is always negative and we know the denominator is always positive ( $\mathcal{S}_2$  does not contain  $[-1, -1, -1]$ ) so their quotient is always negative.

Consider a vertical face (see Figure 2). We can parameterize a horizontal line (line 1) across the face as  $[a, b, c] = [s, s - t, -1 + t]$ . We can take the derivative of  $q(s, s - t, -1 + t)$  and simplify to obtain:

$$\frac{dq(s, s - t, -1 + t)}{dt} = -\frac{4(-4 + 5s)(1 + s - 2t)}{225\pi(1 + s)}.$$

Critical points occur only when  $t = (1 + s)/2$  or  $s = 4/5$ . For  $s = 4/5$ , this calculation shows that  $q$  is constant along the line, hence minima can be taken to be the endpoints of the line. The case  $t = (1 + s)/2$  corresponds to a vertical line through the center of the face (line 2 in Figure 2). We can parameterize this vertical line as  $[a, b, c] = [t + r(1 - t), \frac{r-1}{2}(1 - t), \frac{r-1}{2}(1 - t)]$ . We calculate:

$$\frac{dq(t + r(1 - t), \frac{r-1}{2}(1 - t), \frac{r-1}{2}(1 - t))}{dt} = \left( (-1 + r)(225\pi + (21 + 22r(-1 + t) - 22t)(1 + r + t - rt)^2) \right) / \left( 225\pi(1 + r + t - rt)^2 \right).$$

Just as before, the constant term in the numerator is large enough that the derivative never changes sign, so the function must be minimized at an endpoint.

Now let us consider the “top” and “bottom” faces in the Figure 2. They are both parameterized as  $[a, b, w - a - b]$  where  $w = 1$  for the top face and  $w = 2r - 1$  for the bottom face. The two components of the gradient can be factored as:

$$\begin{aligned} \frac{\partial q}{\partial a} &\propto 8(3b - 4w)(2a + b - w) \\ \text{and } \frac{\partial q}{\partial b} &\propto 8(3a - 4w)(a + 2b - w) \end{aligned}$$

On the top face, the only solution is  $\{[1/3, 1/3, 1/3]\}$ . On the bottom face, we have critical points when

$$(a, b) \in \{(w/3, w/3), (4w/3, -5w/3), (-5w/3, 4w/3), (4w/3, 4w/3)\}.$$

The only region of  $\mathcal{S}_2$  which has not yet been covered explicitly, or implicitly by symmetry is line 3 in the Figure 2. We can parameterize this line as  $(a, b, c) = (r + t, r + t, -1)$  for  $t \in [0, 1 - r]$ . We take the derivative with respect to  $t$  to obtain:

$$\frac{dq(r + t, r + t, -1)}{dt} = \frac{4(-5 + 8r + 8t)(1 + r + t)^2 - 225\pi}{225\pi(1 + r + t)^2}.$$

Once again the constants in the numerator are large enough that the derivative never changes sign. Our analysis and symmetry of the function imply that we need to only check 5 points:

$$\begin{aligned} q(1, 1, -1) &\approx 0.811, \\ q(w/3, w/3, w/3) &\approx 0.934, \\ q(4w/3, -5w/3, 4w/3) &\approx 0.934, \\ q(r, r, -1) &\approx 0.933, \\ q(1/3, 1/3, 1/3) &\approx 0.805, \\ q(r, (r - 1)/2, (r - 1)/2) &\approx 0.935, \\ \text{and } q(1, 0, 0) &\approx 0.808. \end{aligned}$$

□

## E Technical Quantum Facts

In this subsection we present the proofs for the quantum facts we used in the paper.

*Proof of Lemma 14.* For this proof we will have the Bell states take their usual definition:

$$\begin{aligned} |\Phi^+\rangle &= \frac{|00\rangle + |11\rangle}{\sqrt{2}}, & |\Phi^-\rangle &= \frac{|00\rangle - |11\rangle}{\sqrt{2}}, \\ |\Psi^+\rangle &= \frac{|01\rangle + |10\rangle}{\sqrt{2}}, & \text{and } |\Psi^-\rangle &= \frac{|01\rangle - |10\rangle}{\sqrt{2}}. \end{aligned} \tag{102}$$

If  $\text{rank}(P) = 1$  then  $P = |\phi\rangle\langle\phi|$ . Write  $|\phi\rangle$  in the Bell basis:  $|\phi\rangle = \alpha_1 |\Phi^+\rangle + \alpha_2 |\Phi^-\rangle + \alpha_3 |\Psi^+\rangle + \alpha_4 |\Psi^-\rangle$ . Then,

$$\begin{aligned} \Gamma_{11} &= \text{Tr}[\sigma^1 \otimes \sigma^1 |\phi\rangle\langle\phi|] = |\alpha_1|^2 - |\alpha_2|^2 + |\alpha_3|^2 - |\alpha_4|^2, \\ \Gamma_{22} &= \text{Tr}[\sigma^2 \otimes \sigma^2 |\phi\rangle\langle\phi|] = -|\alpha_1|^2 + |\alpha_2|^2 + |\alpha_3|^2 - |\alpha_4|^2, \\ \text{and } \Gamma_{33} &= \text{Tr}[\sigma^3 \otimes \sigma^3 |\phi\rangle\langle\phi|] = |\alpha_1|^2 + |\alpha_2|^2 - |\alpha_3|^2 - |\alpha_4|^2. \end{aligned}$$

So,  $[\Gamma_{11}, \Gamma_{22}, \Gamma_{33}] = |\alpha_1|^2[1, -1, 1] + |\alpha_2|^2[-1, 1, 1] + |\alpha_3|^2[1, 1, -1] + |\alpha_4|^2[-1, -1, -1]$ .

If  $\text{rank}(P) = 2$ ,  $P = |\phi_1\rangle\langle\phi_1| + |\phi_2\rangle\langle\phi_2|$ . Once again let us write these states in the Bell basis:

$$\begin{aligned} |\phi_1\rangle &= \alpha_1 |\Phi^+\rangle + \alpha_2 |\Phi^-\rangle + \alpha_3 |\Psi^+\rangle + \alpha_4 |\Psi^-\rangle \\ \text{and } |\phi_2\rangle &= \beta_1 |\Phi^+\rangle + \beta_2 |\Phi^-\rangle + \beta_3 |\Psi^+\rangle + \beta_4 |\Psi^-\rangle, \end{aligned}$$

where  $\sum_i \alpha_i^* \beta_i = 0$  since  $|\phi_1\rangle$  and  $|\phi_2\rangle$  must be orthogonal.  $[\Gamma_{11}, \Gamma_{22}, \Gamma_{33}]$  can be evaluated in the same way:

$$\begin{aligned} \Gamma_{11} &= \text{Tr}[\sigma^1 \otimes \sigma^1 P] = |\alpha_1|^2 - |\alpha_2|^2 + |\alpha_3|^2 - |\alpha_4|^2 + |\beta_1|^2 - |\beta_2|^2 + |\beta_3|^2 - |\beta_4|^2, \\ \Gamma_{22} &= \text{Tr}[\sigma^2 \otimes \sigma^2 P] = -|\alpha_1|^2 + |\alpha_2|^2 + |\alpha_3|^2 - |\alpha_4|^2 - |\beta_1|^2 + |\beta_2|^2 + |\beta_3|^2 - |\beta_4|^2, \\ \text{and } \Gamma_{33} &= \text{Tr}[\sigma^3 \otimes \sigma^3 P] = |\alpha_1|^2 + |\alpha_2|^2 - |\alpha_3|^2 - |\alpha_4|^2 + |\beta_1|^2 + |\beta_2|^2 - |\beta_3|^2 - |\beta_4|^2. \end{aligned}$$

So,  $[\Gamma_{11}, \Gamma_{22}, \Gamma_{33}] = (|\alpha_1|^2 + |\beta_1|^2)[1, -1, 1] + (|\alpha_2|^2 + |\beta_2|^2)[-1, 1, 1] + (|\alpha_3|^2 + |\beta_3|^2)[1, 1, -1] + (|\alpha_4|^2 + |\beta_4|^2)[-1, -1, -1]$ . Observe that the hull  $\mathcal{T}$  is defined by the following inequalities:

$$[x, y, z] \in \mathcal{T} \Leftrightarrow -2 \leq x + y + z \leq 2, \quad (103)$$

$$-2 \leq x - y + z \leq 2, \quad (104)$$

$$-2 \leq -x + y + z \leq 2, \quad (105)$$

$$-2 \leq x + y - z \leq 2. \quad (106)$$

We will prove this case by demonstrating our derived expression for  $[\Gamma_{11}, \Gamma_{22}, \Gamma_{33}]$  satisfies all these equations.

**Equation (103)**

$$\begin{aligned} \Gamma_{11} + \Gamma_{22} + \Gamma_{33} &= (|\alpha_1|^2 + |\beta_1|^2) + (|\alpha_2|^2 + |\beta_2|^2) + (|\alpha_3|^2 + |\beta_3|^2) - 3(|\alpha_4|^2 + |\beta_4|^2) \\ &= (1 - |\alpha_4|^2) + (1 - |\beta_4|^2) - 3(|\alpha_4|^2 + |\beta_4|^2) = 2(1 - 2(|\alpha_4|^2 + |\beta_4|^2)). \end{aligned}$$

The case follows since  $0 \leq |\alpha_4|^2 + |\beta_4|^2 \leq 1$ . To see this, complete the vectors  $\alpha$  and  $\beta$  to a basis for  $\mathbb{C}^4$ :  $\{\alpha, \beta, \gamma, \eta\}$ . Also let  $e_4 = [0, 0, 0, 1]$ . Then,  $\text{Tr}((|\alpha\rangle\langle\alpha| + |\beta\rangle\langle\beta| + |\gamma\rangle\langle\gamma| + |\eta\rangle\langle\eta|) |e_4\rangle\langle e_4|) = 1$ .

**Equation (104)**

$$\begin{aligned} \Gamma_{11} - \Gamma_{22} + \Gamma_{33} &= 3(|\alpha_1|^2 + |\beta_1|^2) - (|\alpha_2|^2 + |\beta_2|^2) - (|\alpha_3|^2 + |\beta_3|^2) - (|\alpha_4|^2 + |\beta_4|^2) \\ &= 3(|\alpha_1|^2 + |\beta_1|^2 - (2 - |\alpha_1|^2 - |\beta_1|^2)) = 2(2(|\alpha_1|^2 + |\beta_1|^2) - 1). \end{aligned}$$

This case follows since  $0 \leq |\alpha_1|^2 + |\beta_1|^2 \leq 1$  by the same argument.

**Equation (105)**

$$\begin{aligned} -\Gamma_{11} + \Gamma_{22} + \Gamma_{33} &= (-1)(|\alpha_1|^2 + |\beta_1|^2) + 3(|\alpha_2|^2 + |\beta_2|^2) - (|\alpha_3|^2 + |\beta_3|^2) - (|\alpha_4|^2 + |\beta_4|^2) \\ &= 2(2(|\alpha_2|^2 + |\beta_2|^2) - 1). \end{aligned}$$

This case follows since  $0 \leq |\alpha_2|^2 + |\beta_2|^2 \leq 1$  by the same argument.

**Equation (106)**

$$\begin{aligned} \Gamma_{11} + \Gamma_{22} - \Gamma_{33} &= (-1)(|\alpha_1|^2 + |\beta_1|^2) - (|\alpha_2|^2 + |\beta_2|^2) + 3(|\alpha_3|^2 + |\beta_3|^2) - (|\alpha_4|^2 + |\beta_4|^2) \\ &= 2(2(|\alpha_3|^2 + |\beta_3|^2) - 1). \end{aligned}$$

This case follows since  $0 \leq |\alpha_3|^2 + |\beta_3|^2 \leq 1$  by the same argument.

If  $\text{rank}(P) = 3$ , then  $P = \mathbb{I} - |\phi\rangle\langle\phi|$  so if the moment matrix for  $|\phi\rangle\langle\phi|$  has values  $[\Gamma_{11}, \Gamma_{22}, \Gamma_{33}]$ , then the moment matrix for  $P$  has values  $[\Gamma'_{11}, \Gamma'_{22}, \Gamma'_{33}] = -[\Gamma_{11}, \Gamma_{22}, \Gamma_{33}]$ . It is easy to verify that  $\{-[\Gamma_{11}, \Gamma_{22}, \Gamma_{33}] : [\Gamma_{11}, \Gamma_{22}, \Gamma_{33}] \in \mathcal{S}\} = -\mathcal{S}$  as defined so Equation (30) is proven. To prove Equation (31), note:

$$0 \leq \text{Tr}(P\rho) \leq 1 \Rightarrow 0 \leq 4\text{Tr}(P\rho) \leq 4 \Rightarrow -k \leq 4\text{Tr}(P\rho) - k \leq 4 - k.$$

To prove Equation (32) set  $[\Sigma_{11}, \Sigma_{22}, \Sigma_{33}] = [a, b, c]$  and  $[\Gamma_{11}, \Gamma_{22}, \Gamma_{33}] = [p, q, r]$ . We will show that if  $[a, b, c]$  is restricted to some polytope,  $\mathcal{P}_1$ , and  $[p, q, r]$  is restricted to some polytope,  $\mathcal{P}_2$ , then

$$\min_{\substack{[a,b,c] \in \mathcal{P}_1 \\ [p,q,r] \in \mathcal{P}_2}} [a, b, c] \cdot [p, q, r] \geq \min_{\substack{[a,b,c] \in \mathcal{B}_1 \\ [p,q,r] \in \mathcal{B}_2}} [a, b, c] \cdot [p, q, r], \quad (107)$$

where  $\mathcal{B}_i$  is the set of extreme points of  $\mathcal{P}_i$ . Similarly, we will show

$$\max_{\substack{[a,b,c] \in \mathcal{P}_1 \\ [p,q,r] \in \mathcal{P}_2}} [a, b, c] \cdot [p, q, r] \leq \max_{\substack{[a,b,c] \in \mathcal{B}_1 \\ [p,q,r] \in \mathcal{B}_2}} [a, b, c] \cdot [p, q, r]. \quad (108)$$

For Equation (107) fix  $[p, q, r] \in \mathcal{P}_2$  and write it as a convex combination of extreme points  $[p, q, r] = \sum_i \lambda_i [p_i, q_i, r_i]$ .

$$\min_{[a,b,c] \in \mathcal{P}_1} \sum_i \lambda_i [a, b, c] \cdot [p_i, q_i, r_i]. \quad (109)$$

Since  $\sum_i \lambda_i = 1$  we may lower bound this quantity by

$$\geq \min_{\substack{[a,b,c] \in \mathcal{P}_1 \\ [p',q',r'] \in \mathcal{B}_2}} [a, b, c] \cdot [p', q', r'].$$

Since this argument holds for all  $(p, q, r)$  we can uniformly lower bound:

$$\min_{\substack{[a,b,c] \in \mathcal{P}_1 \\ [p,q,r] \in \mathcal{P}_2}} [a, b, c] \cdot [p, q, r] \geq \min_{\substack{[a,b,c] \in \mathcal{P}_1 \\ [p',q',r'] \in \mathcal{B}_2}} [a, b, c] \cdot [p', q', r'].$$

For fixed  $[p, q, r] \in \mathcal{B}_2$  apply the same argument to  $[a, b, c]$  to get Equation (107). The argument for Equation (108) is similar. Applying Equation (107) and Equation (108) to the relevant polytopes provides Equation (32).

Equation (34) follows from [Gam16]. To prove Equation (33) if  $k = 1$  or  $3$  note that in this case the 1-local parts of the projector are proportional to the 1-local parts for a pure state, hence [Gam16] provides this case. For  $k = 2$  observe that the 1-local part of the 2-moment is the sum of 1-local parts from two pure states. Hence, by the triangle inequality the length of the 1-local part is at most 2. □

We also give the map between standard (classical) and Max 2-QSAT:

**Theorem 39.** *Max 2-QSAT generalizes Max 2-SAT.*

*Proof.* An instance of Max 2-SAT corresponds to a set of variables  $\{x_i\}_{i=1}^n$  as well as a multiset of clauses  $E = \{(y_i, y_j)\}$  where each  $y_i$  is either  $x_i$  or  $\neg x_i$ . Let  $OPT$  be the maximum number of clauses which are satisfiable with some Boolean assignment to the variables  $\{x_i\}$ . Define the following function:

$$b(y_i) = \begin{cases} 1 & \text{if } y_i = \neg x_i \\ 0 & \text{if } y_i = x_i \end{cases},$$

and consider the following mapping between clauses and Hamiltonians,

$$(y_i, y_j) \leftrightarrow \frac{3}{4} \mathbb{I}_{ij} - (-1)^{b(y_i)} \frac{\sigma_i^3 \otimes \mathbb{I}_j}{4} - (-1)^{b(y_j)} \frac{\mathbb{I}_i \otimes \sigma_j^3}{4} - (-1)^{b(y_i)+b(y_j)} \frac{\sigma_i^3 \otimes \sigma_j^3}{4}.$$

Denote the 2-Local term on the RHS as  $H_e(y_i, y_j)$ . Then, we claim

$$OPT = \lambda_{max} \left( \sum_{(y_i, y_j) \in E} H_e(y_i, y_j) \right).$$

This holds because the Hamiltonian is diagonal in the computational basis, hence its largest eigenstate can be assumed to be a computational basis state. Such a state has 0 and 1 values corresponding to *False* and *True* assignments. □

The next result we present is that our constants are nearly tight for quadratic Hamiltonians, in that our approximation factors are about as good as we can expect for approximation algorithms which yield product states.

**Theorem 40.** *There exist instances of the 2-Local Hamiltonian problem,  $\{H_e\}$  with strictly quadratic projectors where each  $H_e$  has rank  $k$  such that*

$$\max_{|\phi_1\rangle \in \mathbb{C}^2, |\phi_2\rangle \in \mathbb{C}^2} \langle \phi_1 | \otimes \langle \phi_2 | \left( \sum_e H_e \right) | \phi_1 \rangle \otimes | \phi_2 \rangle \leq \beta(k) \lambda_{max} \left( \sum_e H_e \right), \quad (110)$$

$$\text{where } \beta(k) = \begin{cases} 1/2 & \text{if } k = 1 \\ 2/3 & \text{if } k = 2 \\ 5/6 & \text{if } k = 3 \end{cases}.$$

*Proof.* Let the Bell states be defined as in Equation (102). If  $k = 1$  consider the Hamiltonian  $H = |\Psi^-\rangle \langle \Psi^-|$ . From, e.g., [GP19], we know that if we optimize  $|\phi\rangle$  over product states we can get at most 1/2 for  $\langle \phi | H | \phi \rangle$ , while it is apparent that  $\lambda_{max} = 1$  since we can take the singlet,  $|\Phi^-\rangle$ .

If  $k = 2$  consider the Hamiltonian (recall the problem formulation allows multi-edges):

$$H = \frac{1}{3} (|\Psi^-\rangle \langle \Psi^-| + |\Phi^+\rangle \langle \Phi^+|) + \frac{1}{3} (|\Psi^-\rangle \langle \Psi^-| + |\Phi^-\rangle \langle \Phi^-|) \quad (111)$$

$$+ \frac{1}{3} (|\Psi^-\rangle \langle \Psi^-| + |\Psi^+\rangle \langle \Psi^+|) \\ = \frac{1}{3} \mathbb{I} + \frac{2}{3} |\Psi^-\rangle \langle \Psi^-|. \quad (112)$$

Similarly, we can expect objective at most 1/2 for the singlet if we optimize over product states, hence  $\max_{|\phi\rangle \in PROD} \langle \phi | H | \phi \rangle = 2/3$ . Observe that we can achieve  $\lambda_{max} = 1$  by taking the singlet once again.

If  $k = 3$  consider:

$$H = \frac{1}{3} (\mathbb{I} - |\Phi^+\rangle \langle \Phi^+|) + \frac{1}{3} (\mathbb{I} - |\Phi^-\rangle \langle \Phi^-|) + \frac{1}{3} (\mathbb{I} - |\Psi^+\rangle \langle \Psi^+|) \quad (113)$$

$$= \frac{2}{3} \mathbb{I} + \frac{1}{3} |\Psi^-\rangle \langle \Psi^-|. \quad (114)$$

We get at most 5/6 for product states in the same way, while  $\lambda_{max} = 1$ . □



## F Traceless, Bipartite, and Strictly Quadratic Hamiltonians

The final result we present is an approximation algorithm for strictly quadratic (Definition 5) instances of traceless 2-Local Hamiltonian on bipartite interaction graphs. Since this result is relatively self-contained and very different from the main results of this paper, we present both the statements and the analysis here. We demonstrate that this problem can be approximated using Krivine rounding [Kri77], with the same product-state ansatz as in [BGKT19]. More generally we show how traceless instances of 2-LH can be approximated using an approximation algorithm for a related classical 2-CSP.

An instance,  $\{H_e\}$ , of QLH on  $n$  qubits (Problem 4) is *traceless* if  $\text{Tr}[\sum_e H_e] = 0$ . For a traceless instance, we may assume that each term  $H_e$  is traceless by replacing it with  $H'_e = H_e - \frac{1}{2^n} \text{Tr}[H_e] \mathbb{I}$ , which preserves the maximum eigenvalue of  $H = \sum_e H_e$ . We will also assume for our result that each  $H_e$  is strictly quadratic, so that we may assume each  $H_e = w_e \sigma_i^k \otimes \sigma_j^l \otimes \mathbb{I}_{[n] \setminus \{i,j\}}$  for  $w_e \neq 0$  and  $k, l \in [3]$ . Given such a Hamiltonian, we will create a ‘‘Pauli’’ interaction graph  $G = (V, E)$  where the set of vertices have a one-to-one correspondence with nontrivial single qubit Pauli operators. If there are  $n$  qubits there would be  $3n$  such vertices, one for each  $\sigma_i^k$  for  $i \in [n]$  and  $k \in [3]$ . We place an edge between vertices if there is a term  $H_e$  in the description of the local Hamiltonian problem which contains both Pauli matrices, i.e. vertex  $(i, k)$  is connected to  $(j, l)$  if there is an  $H_e = w_e \sigma_i^k \otimes \sigma_j^l \otimes \mathbb{I}_{[n] \setminus \{i,j\}}$ . If the corresponding graph is bipartite, then we say that the corresponding (strictly quadratic) 2-local Hamiltonian is traceless and bipartite. For ease of reference, we provide a formal definition:

**Definition 41** (Pauli Interaction Graph). *Let  $\{H_e\}$  be a set of 2-local terms on  $n$  qubits where each  $H_e = w_e \sigma_i^k \otimes \sigma_j^l \otimes \mathbb{I}_{[n] \setminus \{i,j\}}$  with  $w_e \neq 0$  and  $k, l \in [3]$ . Let  $V = \{(i, k) \mid i \in [n] \text{ and } k \in [3]\}$ . Construct a set of pairs  $E \subseteq [3n] \times [3n]$  with  $(ik, jl) \in E$  if and only if there exists  $e$  such that  $H_e = w_e \sigma_i^k \otimes \sigma_j^l \otimes \mathbb{I}_{[n] \setminus \{i,j\}}$ . We say that  $\{H_e\}$  is strictly quadratic, bipartite and traceless if  $G = (V, E)$  is bipartite.*

Our general approach is to solve a classical CSP on the Pauli interaction graph of a 2-local Hamiltonian. In particular, we seek to maximize the weight earned from the edges  $\sigma_i^k \otimes \sigma_j^l \otimes \mathbb{I}_{[n] \setminus \{i,j\}}$ , where we now assign values in  $\{\pm 1\}$  to each variable  $\sigma_i^k$ , represented by a vertex in  $V$ . Such an assignment is converted to a product state by treating the unit vector  $[\sigma_i^1, \sigma_i^2, \sigma_i^3]/\sqrt{3}$  as a Bloch vector for qubit  $i$ . We note that this approach works for any traceless instance of 2-LH, where a classical  $\alpha$ -approximation algorithm for the classical 2-CSP on the Pauli interaction graph yields a classical  $\frac{\alpha}{3}$ -approximation for the original 2-LH instance.

The strictly quadratic, bipartite, and traceless case of the 2-Local Hamiltonian problem is still *QMA*-hard, since many (strictly quadratic) families of Hamiltonians retain *QMA*-hardness even when the edges are restricted to a  $2-d$  lattice [PM15]. For this problem, the relevant relaxation is a weakening of our main SDP, Problem 7. We opt to drop many of the constraints present in Problem 7 for consistency with the result we use as a black box [Kri77, BdOFV14]. Given a local term of the form  $H_e = w_e \sigma_i^k \otimes \sigma_j^l \otimes \mathbb{I}_{[n] \setminus \{i,j\}}$ , where in the present context we allow  $w_e$  to be negative, let  $D_e \in \mathbb{R}^{3n \times 3n}$  be a symmetric matrix indexed by nontrivial single-qubit Pauli operators (i.e.  $[n] \times [3]$ ), such that:

$$D_e(\sigma_i^k, \sigma_j^l) = D_e(\sigma_j^l, \sigma_i^k) = w_e/2, \quad (115)$$

and  $D_e = 0$  otherwise.

**Problem 42.** Given a strictly quadratic instance of QLH on  $n$  qubits with terms  $\{H_e\}$ , for each  $H_e$  let  $D_e$  be defined as in Equation (115). Solve the following SDP:

$$\begin{aligned} & \max \sum_e \text{Tr}[D_e M] \\ & \text{s.t.} \quad M(\sigma_i^k, \sigma_i^k) = 1 \quad \forall (i, k) \in V, \\ & \quad \mathcal{S}(\mathbb{R}^{3n \times 3n}) \ni M \succeq 0, \end{aligned} \tag{116}$$

where  $\mathcal{S}(\cdot)$  refers to the class of symmetric matrices.

Given this relaxation, which is equivalent to the relaxation used by Goemans and Williamson [GW95], we appeal to a classical rounding scheme, used as a black box, to obtain the result. Given some graph  $G = (V, E)$  with  $|V| = n$ , let  $A$  be a symmetric matrix with rows and columns indexed by elements of  $V$ . For each  $v \in V$  we associate a  $\mathbf{w}_v \in \mathbb{R}^n$ . The main observation is the following:

**Theorem 43** ( [Kri77] and Remark 2 on p. 95 in [BdOFV14]). *If  $G$  is a bipartite graph, then there is a polynomial-time randomized algorithm which outputs variables  $\{z_v\}_{v \in V}$  where each  $z_v \in \{\pm 1\}$  and*

$$\mathbb{E} \left[ \sum_{uv \in E} A_{uv} z_u z_v \right] \geq \frac{2(\ln(1 + \sqrt{2}))}{\pi} \sum_{uv \in E} A_{uv} \mathbf{w}_u \cdot \mathbf{w}_v.$$

The above gives an approximation algorithm for the Grothendieck problem if the underlying graph is bipartite. The right hand side corresponds to the semidefinite relaxation of some NP-hard optimization problem and the left hand side corresponds to some feasible (but likely not optimal) solution to the optimization problem. We obtain the following corollary:

**Corollary 44.** *Let  $H = \sum_e H_e$  be traceless, strictly quadratic, and bipartite in the sense of Definition 41. Then, there is a randomized polynomial-time algorithm which produces a pure product state  $|\psi\rangle$  such that:*

$$\mathbb{E} \left[ \langle \psi | \sum_e H_e | \psi \rangle \right] \geq \frac{2 \ln(1 + \sqrt{2})}{3\pi} \lambda_{\max} \left( \sum_e H_e \right). \tag{117}$$

*Proof.* Construct a  $3n \times 3n$  matrix  $A$  by taking

$$A(\sigma_i^k, \sigma_j^l) = \text{Tr} \left[ (\sigma_i^k \otimes \sigma_j^l \otimes \mathbb{I}_{[n] \setminus \{i, j\}}) \sum_e H_e \right] / 2^{n+1}.$$

Observe that the matrix  $A$  is bipartite in the same sense as Theorem 43: Let  $G = (V, E)$  be the graph described previously, where we have a vertex for each single qubit Pauli matrix and vertices are connected if they interact through  $H$ . We have a partition of the vertices into two sets  $B_1$  and  $B_2$  where no edge links two vertices in the same  $B_i$ . Then,  $A(\sigma_i^k, \sigma_j^l) = 0$  if  $[(i, k) \in B_1 \text{ and } (j, l) \in B_1]$  or  $[(i, k) \in B_2 \text{ and } (j, l) \in B_2]$ .

Now let us define the following optimization problem:

$$\max_{\substack{z_{ik} \in \{\pm 1\} \\ \forall (i, k) \in V}} \sum_{(ik, jl) \in E} A(\sigma_i^k, \sigma_j^l) z_{ik} z_{jl}. \tag{118}$$

We can take its natural semidefinite relaxation [GW95] to obtain:

$$\max_{\substack{M(\sigma_i^k, \sigma_i^k) = 1 \quad \forall i, k \\ \mathcal{S}(\mathbb{R}^{3n \times 3n}) \ni M \succeq 0}} \sum_{(ik, jl) \in E} A(\sigma_i^k, \sigma_j^l) M(\sigma_i^k, \sigma_j^l),$$

or written in terms of the Cholesky vectors:

$$\max_{\substack{\mathbf{w}_{ik} \in \mathbb{R}^{3n} \\ \|\mathbf{w}_{ik}\|=1 \forall (i,k) \in V}} \sum_{(ik,jl) \in E} A(\sigma_i^k, \sigma_j^l) \mathbf{w}_{ik} \cdot \mathbf{w}_{jl}.$$

Observe that Equation (118) is exactly the relaxation for the 2-Local Hamiltonian problem, Problem 42, since  $\sum_e D_e = A$ . Theorem 43, Krivine rounding, provides a random set of variables  $z_{ik} \in \{\pm 1\}$  such that:

$$\mathbb{E} \left[ \sum_{(ik,jl) \in E} A(\sigma_i^k, \sigma_j^l) z_{ik} z_{jl} \right] \geq \frac{2 \ln(1 + \sqrt{2})}{\pi} \lambda_{max} \left( \sum_e H_e \right).$$

Dividing both sides by 3, we obtain:

$$\mathbb{E} \left[ \sum_{(ik,jl) \in E} A(\sigma_i^k, \sigma_j^l) \frac{z_{ik}}{\sqrt{3}} \frac{z_{jl}}{\sqrt{3}} \right] \geq \frac{2 \ln(1 + \sqrt{2})}{3\pi} \lambda_{max} \left( \sum_e H_e \right).$$

Note that the objective of the LHS corresponds to a valid quantum product state just as in [BGKT19]:

$$\rho = \bigotimes_{i=1}^n \left( \frac{\mathbb{I} + (z_{i1}/\sqrt{3})\sigma^1 + (z_{i2}/\sqrt{3})\sigma^2 + (z_{i3}/\sqrt{3})\sigma^3}{2} \right).$$

□