

# TIME REGULARITY FOR GENERALIZED MEHLER SEMIGROUPS

ALESSANDRA LUNARDI

**ABSTRACT.** We study continuity and Hölder continuity of  $t \mapsto P_t f$ , where  $P_t$  is a generalized Mehler semigroup in  $C_b(X)$ , the space of the continuous and bounded functions from a Banach space  $X$  to  $\mathbb{R}$ , and  $f \in C_b(X)$ . The generators  $L$  of such semigroups are realizations of a class of differential and pseudo-differential operators, both in finite and in infinite dimension. Examples of operators  $L$  to which this theory is applicable include Ornstein-Uhlenbeck operators with fractional diffusion in finite dimension, and Ornstein-Uhlenbeck operators with associated strong-Feller semigroups, in infinite dimension.

## 1. INTRODUCTION

Let  $X$  be a real Banach space, and let  $C_b(X)$  denote the space of the continuous and bounded functions from  $X$  to  $\mathbb{R}$ , endowed with the sup norm. A generalized Mehler semigroup  $P_t$  is a semigroup of bounded operators in  $C_b(X)$  that may be represented by

$$P_t f(x) = \int_X f(T_t x + y) \mu_t(dy), \quad t \geq 0, f \in C_b(X), x \in X, \quad (1.1)$$

where  $T_t$  is a strongly continuous semigroup of bounded operators on  $X$ , and  $\{\mu_t : t \geq 0\}$  is a family of Borel probability measures in  $X$  such that  $\mu_0 = \delta_0$  (the Dirac measure at  $0 \in X$ ),  $t \mapsto \mu_t$  is weakly continuous in  $[0, +\infty)$ , and

$$\mu_{t+s} = (\mu_t \circ T_s^{-1}) * \mu_s, \quad t, s > 0, \quad (1.2)$$

that is in fact an algebraic necessary and sufficient for  $P_t$  be a semigroup.

They arise as transition semigroups of (weak or mild) solutions to stochastic differential equations such as

$$dX(t) = AX(t)dt + dL_t, \quad t > 0; X(0) = x, \quad (1.3)$$

where  $A : D(A) \subset X \rightarrow X$  is the infinitesimal generator of  $T_t$ , and  $\{L_t : t \geq 0\}$  is a Lévy process in  $X$ . Detailed discussions about several features of such semigroups, and about connections with the stochastic differential equations (1.3), are in [7, 9, 13, 14, 15, 17, 18, 19, 26, 27, 33] and in the papers quoted therein; see also the review paper [1].

It is not hard to see that for every  $f \in C_b(X)$  and  $x \in X$  the function  $(t, x) \mapsto P_t f(x)$  is continuous in  $[0, +\infty) \times X$  (e.g., [7, Lemma 2.1]). But  $P_t$  is not strongly continuous in general, as well as its restriction to the space  $BUC(X)$  of the bounded and uniformly continuous functions from  $X$  to  $\mathbb{R}$ . The generator  $L$  of  $P_t$  is defined through its resolvent,

$$R(\lambda, L)f(x) = \int_0^\infty e^{-\lambda t} P_t f(x) dt, \quad \lambda > 0, f \in C_b(X), x \in X, \quad (1.4)$$

however it is not the infinitesimal generator in the standard sense.

In this paper we study continuity and Hölder continuity of the function  $[0, +\infty) \mapsto C_b(X)$ ,  $t \mapsto P_t f$ , for  $f \in C_b(X)$ .

---

2010 *Mathematics Subject Classification.* 35B65, 35R15, 47D07.

*Key words and phrases.* Hölder regularity, generalized Mehler semigroups, Ornstein-Uhlenbeck semigroups, fractional diffusion.

As a simple first result, we show that for every  $f \in C_b(X)$ ,  $t \mapsto P_t f$  is continuous if and only if  $\lim_{\lambda \rightarrow +\infty} \lambda R(\lambda, L)f = f$ , if and only if  $f \in \overline{D(L)}$ . So,  $\overline{D(L)}$  is the subspace of strong continuity of  $P_t$ . This characterization is shared by (not necessarily strongly continuous) analytic semigroups in general Banach spaces, see e.g. [21, Sect. 2.1]. But in general, generalized Mehler semigroups are not analytic in  $C_b(X)$  nor in  $BUC(X)$ . Even worse, they are not necessarily eventually norm continuous: see [29] for Ornstein-Uhlenbeck semigroups.

Then we prove that for every  $\alpha \in (0, 1)$ ,  $t \mapsto P_t f$  is  $\alpha$ -Hölder continuous if and only if  $\sup_{\lambda > 0} \|\lambda^\alpha L R(\lambda, L)f\|_\infty < +\infty$ , if and only if  $f$  is in the real interpolation space  $(C_b(X), D(L))_{\alpha, \infty}$ . This characterization is shared by strongly continuous or analytic semigroups of nonnegative type.

In fact, both characterizations are proved for a larger class of semigroups, and precisely for all bounded semigroups  $P_t$  such that  $(t, x) \mapsto P_t f(x)$  belongs to  $C([0, +\infty) \times X)$  for every  $f \in C_b(X)$ .

Going back to generalized Mehler semigroups, if  $P_t$  enjoys suitable smoothing assumptions it is possible to provide more explicit descriptions of both  $\overline{D(L)}$  and  $(C_b(X), D(L))_{\alpha, \infty}$ .

Concerning  $\overline{D(L)}$ , if  $P_t$  maps  $C_b(X)$  to  $BUC(X)$  we have

$$\overline{D(L)} = \{f \in BUC(X) : \lim_{t \rightarrow 0} \|f(T_t \cdot) - f\|_\infty = 0\}.$$

Concerning  $(C_b(X), D(L))_{\alpha, \infty}$ , we need better smoothing properties of  $P_t$ . Precisely, we assume that the following hypothesis holds.

**Hypothesis 1.1.** *Each  $\mu_t$  is Radon, and Fomin differentiable along the range of  $T_t$ . There exist  $C > 0$ ,  $\omega \in \mathbb{R}$  such that denoting by  $\beta_{t,h}$  the Fomin derivative of  $\mu_t$  along  $T_t h$ , we have*

$$\|\beta_{t,h}\|_{L^1(X, \mu_t)} \leq \frac{C e^{\omega t}}{t^\theta} \|h\|, \quad t > 0, h \in X. \quad (1.5)$$

Then it is possible to show that  $P_t$  is strong-Feller (namely, if  $f$  is Borel measurable and bounded, then  $P_t f$ , still defined by (1.1), belongs to  $C_b(X)$ ) and moreover it maps  $C_b(X)$  into the space  $C_b^k(X)$  of the  $k$  times Fréchet differentiable functions with bounded derivatives up to the order  $k$ , for  $k \in \mathbb{N}$ ,  $k < 1/\theta$ , and there is  $C_k > 0$  such that

$$\|P_t f\|_{C_b^k(X)} \leq \frac{C_k}{t^{k\theta}} \|f\|_\infty, \quad f \in C_b(X), \quad 0 < t \leq 1, \quad k < 1/\theta. \quad (1.6)$$

Hypothesis 1.1 was introduced in [25], where Schauder type theorems were proved. Estimates (1.6) are used to prove that for every  $\alpha \in (0, 1)$  we have

$$(C_b(X), D(L))_{\alpha, \infty} \subset \begin{cases} C_b^{\alpha/\theta}(X), & \alpha/\theta \notin \mathbb{N}, \\ Z_b^{\alpha/\theta}(X), & \alpha/\theta \in \mathbb{N}, \end{cases} \quad (1.7)$$

with continuous embedding. Here, for  $k \in \mathbb{N}$ ,  $\sigma \in (0, 1)$ ,  $C_b^{k+\sigma}(X)$  consists of the functions  $f \in C_b^k(X)$  having  $\sigma$ -Hölder continuous  $k$ -th order derivative, and  $Z_b^k(X)$  is the Zygmund space defined in Sect. 2. Besides estimates (1.6), the proof of (1.7) relies on results proved in [25] and on the continuous embeddings

$$(C_b(X), C_b^\beta(X))_{\alpha, \infty} \subset \begin{cases} C_b^{\alpha\beta}(X), & \alpha\beta \notin \mathbb{N}, \\ Z_b^{\alpha\beta}(X), & \alpha\beta \in \mathbb{N}, \end{cases}$$

valid for every Banach space  $X$  and for every  $\beta > 0$ ,  $\alpha \in (0, 1)$ .

From the representation formula (1.1) we see that  $P_t$  is a contraction semigroup in all spaces  $C_b^\beta(X)$ , and also in all Zygmund spaces. Therefore (1.7) yields that if  $t \mapsto P_t f(x)$  is  $\alpha$ -Hölder

continuous uniformly with respect to  $x$ , then  $x \mapsto P_t f(x)$  belongs to  $C_b^{\alpha/\theta}(X)$  (or to  $Z_b^{\alpha/\theta}(X)$ ), uniformly with respect to  $t$ .

So, we have an extension to the present general setting of the heuristic principle “time regularity implies space regularity” that holds for semigroups associated to uniformly elliptic operators with bounded regular coefficients in  $\mathbb{R}^N$ . However the converse does not hold; in other words the embeddings in (1.7) cannot be replaced by equalities, in general. Under a further assumption on the moments of the measures  $\mu_t$ , namely

$$\exists \gamma, C > 0 : \int_X \|x\|^\gamma \mu_t(dx) \leq C t^{\gamma\theta}, \quad 0 < t \leq 1, \quad (1.8)$$

we prove that for every  $\alpha \in (0, 1 \wedge \theta) \cap (0, \gamma\theta]$  we have

$$(C_b(X), D(L))_{\alpha, \infty} = C_b^{\alpha/\theta}(X) \cap Y_\alpha, \quad (1.9)$$

with equivalence of the respective norms, where

$$Y_\alpha := \left\{ f \in C_b(X) : [f]_{Y_\alpha} := \sup_{t>0} \frac{\|f(T_t \cdot) - f\|_\infty}{t^\alpha} < +\infty \right\}, \quad \|f\|_{Y_\alpha} = \|f\|_\infty + [f]_{Y_\alpha}.$$

If in addition all the measures  $\mu_t$  are centered, namely  $\mu_t(B) = \mu_t(-B)$  for every Borel set  $B$ , and  $\theta < 1$ , the above equivalence holds also for all  $\alpha \in (\theta, 2\theta) \cap (0, \gamma\theta]$ .

Both in finite and in infinite dimension, popular examples of generalized Mehler semigroups are the Ornstein-Uhlenbeck semigroups (§5.1), in which case the measures  $\mu_t$  are Gaussian. We recall that if  $X = \mathbb{R}^N$ , given any matrices  $A, Q = Q^* \geq 0$ , and setting  $Q_t = \int_0^t e^{sA} Q e^{sA^*} ds$ , the relevant Ornstein-Uhlenbeck semigroup  $P_t$  is defined by (1.1) where  $T_t = e^{tA}$ , and  $\mu_t$  is the Gaussian measure with mean 0 and covariance  $Q_t$ . The generator  $L$  is a realization of the operator  $\mathcal{L}u(x) = \text{Tr}(QD^2u(x))/2 + \langle Ax, \nabla u(x) \rangle$ . If in addition  $Q > 0$ , estimates (1.6) holds with  $\theta = 1/2$  and (1.8) holds for any  $\gamma > 0$ . The above characterization of  $(C_b(\mathbb{R}^N), D(L))_{\alpha, \infty}$  was already proved in [11]; in the subsequent paper [2] it was re-discovered that  $t \mapsto P_t f$  is  $\alpha$ -Hölder continuous with values in  $C_b(\mathbb{R}^N)$  for  $\alpha < 1/2$  iff  $f \in C_b^{2\alpha}(\mathbb{R}^N) \cap Y_\alpha$ . A characterization of  $(C_b(\mathbb{R}^N), D(L))_{\alpha, \infty}$  is available also in the case that  $\text{Det } Q = 0$  but  $\mathcal{L}$  is hypoelliptic ([23]).

Still for Ornstein-Uhlenbeck semigroups, in infinite dimension sufficient conditions for Hypothesis 1.1 to hold are well known in the case that  $X$  is a separable Hilbert space; for any  $\theta \geq 1/2$  there are examples such that (1.5) is satisfied (e.g., [8, 12, 2]). Instead, it is not clear whether (1.8) holds for some  $\gamma > 0$ . Therefore, we can prove only the embeddings (1.7).

Nontrivial new examples of generators of generalized Mehler semigroups satisfying both Hypothesis 1.1 and (1.8) are Ornstein-Uhlenbeck operators with fractional diffusion in finite dimension (§5.2), such as

$$(\mathcal{L}u)(x) = \frac{1}{2}(\text{Tr}^s(QD^2u))(x) - \langle Ax, \nabla u(x) \rangle, \quad x \in \mathbb{R}^N,$$

with  $s \in (0, 1)$  and  $Q, A$  matrices such that  $Q = Q^* > 0$ . Here  $\text{Tr}^s(QD^2)$  is the pseudo-differential operator with symbol  $-\langle Q\xi, \xi \rangle^s$ . In this case (1.5) holds with  $\theta = 1/(2s)$ , the measures  $\mu_t$  are absolutely continuous with respect to the Lebesgue measure, (1.8) holds for every  $\gamma < 2s$ , and we obtain

$$(C_b(X), D(L))_{\alpha, \infty} = C_b^{2s\alpha}(\mathbb{R}^N) \cap Y_\alpha, \quad \alpha \in (0, 1), \alpha \neq 1/(2s).$$

## 2. INTERPOLATION IN SPACES OF CONTINUOUS AND BOUNDED FUNCTIONS

**2.1. Function spaces.** Let  $X, Y$  be Banach spaces.

By  $\mathcal{B}_b(X; Y)$  and  $C_b(X; Y)$  we denote the space of all bounded Borel measurable (resp. bounded continuous) functions  $F : X \mapsto Y$ , endowed with the sup norm  $\|F\|_\infty := \sup_{x \in X} \|F(x)\|_Y$ . If  $X = \mathbb{R}$  we set  $\mathcal{B}_b(X; \mathbb{R}) := \mathcal{B}_b(X)$  and  $C_b(X; \mathbb{R}) := C_b(X)$ .

For  $\alpha \in (0, 1)$  we denote by  $C_b^\alpha(X; Y)$  the space of the bounded and  $\alpha$ -Hölder continuous functions  $F : X \mapsto Y$ , endowed with the Hölder norm  $\|F\|_{C_b^\alpha(X; Y)} = \|F\|_\infty + [F]_{C^\alpha(X; Y)}$ , where  $[F]_{C^\alpha(X; Y)} := \sup_{x \in X, h \in X \setminus \{0\}} \|F(x+h) - F(x)\|_Y \|h\|_X^{-\alpha}$ .

For  $\alpha = 1$  we will not consider the Lipschitz condition, but a weaker one. We set

$$Z_b^1(X, Y) := \left\{ F \in C_b(X; Y) : [F]_{Z^1(X, Y)} := \sup_{x, h \in X, h \neq 0} \frac{\|F(x+2h) - 2F(x+h) + F(x)\|_Y}{\|h\|_X} < +\infty \right\}.$$

The space  $Z^1(X; Y)$  is called *Zygmund space*, and it is endowed with the norm

$$\|F\|_{Z_b^1(X, Y)} := \sup_{x \in X} \|F(x)\|_Y + [F]_{Z^1(X, Y)}.$$

For every  $n \in \mathbb{N}$  we denote by  $C_b^k(X)$  the space of the bounded and  $n$  times continuously Fréchet differentiable functions  $f : X \mapsto \mathbb{R}$  with bounded Fréchet derivatives up to the order  $k$ . Its norm is

$$\|f\|_{C_b^k(X)} := \|f\|_\infty + \sum_{j=1}^k \sup_{x \in X} \|D^j f(x)\|_{\mathcal{L}^j(X)}, \quad (2.1)$$

where  $\mathcal{L}^j(X)$  is the space of the  $k$ -linear continuous functions from  $X^j$  to  $\mathbb{R}$ , endowed with the norm  $\|T\|_{\mathcal{L}^j(X)} := \sup \left\{ \frac{|T(h_1, \dots, h_j)|}{\|h_1\| \cdots \|h_j\|} : h_i \in X \setminus \{0\} \right\}$ .

For  $\sigma \in (0, 1)$  and  $k \in \mathbb{N}$  we set

$$C_b^{\sigma+k}(X) := \{f \in C_b^k(X) : D^k f \in C^\sigma(X, \mathcal{L}^k(X))\}, \quad (2.2)$$

$$\|f\|_{C_b^{\sigma+k}(X)} := \|f\|_{C_b^k(X)} + [D^k f]_{C^\sigma(X, \mathcal{L}^k(X))},$$

and for  $k \in \mathbb{N}$ ,  $k \geq 2$ , the higher order Zygmund spaces are defined by

$$Z_b^k(X) := \{f \in C_b^{k-1}(X) : D^{k-1} f \in Z^1(X, \mathcal{L}^{k-1}(X))\}, \quad (2.3)$$

$$\|f\|_{Z_b^k(X)} := \|f\|_{C_b^{k-1}(X)} + [D^{k-1} f]_{Z^1(X, \mathcal{L}^{k-1}(X))}.$$

**2.2. Interpolation.** We recall that if  $E, F$  are Banach spaces and  $F$  is continuously embedded in  $E$ , the real interpolation space  $(E, F)_{\alpha, \infty}$  is defined by

$$(E, F)_{\alpha, \infty} = \{f \in E : \|f\|_{(E, F)_{\alpha, \infty}} := \sup_{\xi > 0} \xi^{-\alpha} K(\xi, f, E, F) < +\infty\}$$

where

$$K(\xi, f, E, F) := \inf\{\|a\|_E + \xi\|b\|_F : f = a + b, a \in E, b \in F\}, \quad \xi > 0, f \in E.$$

There are several equivalent characterizations of real interpolation spaces. In the following we shall use the next one (e.g., [34, Thm. 1.5.3]).

**Proposition 2.1.**  $(E, F)_{\alpha, \infty}$  consists of the elements  $f \in E$  such that  $f = \int_0^\infty \frac{u(t)}{t} dt$ , where  $u \in C((0, +\infty); F)$  is such that  $t \mapsto t^{1-\alpha} u(t) \in L^\infty((0, +\infty); F)$  and  $t \mapsto t^{-\alpha} u(t) \in L^\infty((0, +\infty); E)$ . The norm

$$f \mapsto \inf \left\{ \text{ess sup}_{t>0} \|t^{1-\alpha} u(t)\|_F + \text{ess sup}_{t>0} \|t^{-\alpha} u(t)\|_E : f = \int_0^\infty \frac{u(t)}{t} dt \right\}$$

is equivalent to the norm of  $(E, F)_{\alpha, \infty}$ .

Moreover, given  $\theta \in (0, 1)$  and three Banach spaces  $G \subset F \subset E$ , with continuous embeddings, we say that  $F$  belongs to the class  $J_\theta$  between  $E$  and  $G$  if there exists  $C > 0$  such that

$$\|x\|_F \leq C\|x\|_G^\theta \|x\|_E^{1-\theta}, \quad x \in G.$$

In this case the Reiteration Theorem (e.g., [34, Thm. 1.10.2], [22, Thm. 1.23(ii)]) yields, for every  $\alpha \in (0, 1)$ ,  $\alpha \neq \theta$ ,

$$(E, G)_{\alpha, \infty} \subset \begin{cases} (E, F)_{\alpha/\theta, \infty}, & \text{if } \alpha < \theta, \\ (F, G)_{(\alpha-\theta)/(1-\theta), \infty}, & \text{if } \alpha > \theta, \end{cases} \quad (2.4)$$

with continuous embeddings.

From now on we take  $E = C_b(X)$ , and  $F = C_b^\beta(X)$ , with  $\beta > 0$ . Next statements are far from surprising. However, the known proofs in the case  $X = \mathbb{R}^N$  do not seem to be immediately extendable to the infinite dimensional case, and therefore we provide independent proofs.

**Lemma 2.2.** *For every  $k \in \mathbb{N}$ ,  $k \geq 2$ , there exists  $C_k > 0$  such that*

$$\|D^h f\|_{L^\infty(X, \mathcal{L}^h(X))} \leq C_k \|f\|_\infty^{1-h/k} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))}, \quad h = 1, \dots, k-1, \quad f \in C_b^k(X). \quad (2.5)$$

Similarly, for every noninteger  $\beta > 1$ ,  $\beta = k + \sigma$  with  $k \in \mathbb{N}$  and  $\sigma \in (0, 1)$  there exists  $C_\beta > 0$  such that

$$\|D^h f\|_{L^\infty(X, \mathcal{L}^h(X))} \leq C_\beta \|f\|_\infty^{1-h/\beta} [D^k f]_{C^\sigma(X, \mathcal{L}^k(X))}, \quad h = 1, \dots, k, \quad f \in C_b^\beta(X). \quad (2.6)$$

*Proof.* Let us prove a basic estimate, for  $f \in C_b^{1+\sigma}(X)$  with  $\sigma \in (0, 1]$ . For every  $x \in X$ ,  $h \in X$  with  $\|h\| = 1$  and  $t > 0$ , we have

$$\begin{aligned} |Df(x)(h)| &\leq \left| Df(x)(h) - \frac{f(x+th) - f(x)}{t} \right| + \left| \frac{f(x+th) - f(x)}{t} \right| \\ &\leq \frac{1}{t} \int_0^t |Df(x)(h) - Df(x+\tau h)(h)| d\tau + \frac{2}{t} \|f\|_\infty \\ &\leq [[Df]]_\sigma \frac{t^\sigma}{1+\sigma} + \frac{2}{t} \|f\|_\infty, \end{aligned}$$

where  $[[Df]]_\sigma := [Df]_{C^\sigma(X, X^*)}$  if  $\sigma \in (0, 1)$ ,  $[[Df]]_\sigma := \|D^2 f\|_{L^\infty(X, \mathcal{L}^2(X))}$  if  $\sigma = 1$ . In both cases, for every  $x \in X$ ,

$$\|Df(x)\|_{X^*} \leq [[Df]]_\sigma \frac{t^\sigma}{1+\sigma} + \frac{2}{t} \|f\|_\infty, \quad t > 0.$$

Taking the minimum of the right hand side over  $t > 0$  (or else, choosing  $t = (\|f\|_\infty / [[Df]]_\sigma)^{1/(1+\sigma)}$ ), we see that there exists  $C$  such that

$$\|Df(x)\|_{X^*} \leq C ([[Df]]_\sigma^{1/(\sigma+1)} (\|f\|_\infty)^{1-1/(\sigma+1)}), \quad x \in X, \quad (2.7)$$

which proves (2.5) for  $k = 2$  and (2.6) for  $\beta \in (1, 2)$ .

Estimate (2.7) is readily extended as follows: for every  $k \in \mathbb{N}$ , and  $\sigma \in (0, 1)$  there exists  $K_{k, \sigma} > 0$  such that

$$\|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))} \leq K_{k, \sigma} (\|D^{k-1} f\|_{L^\infty(X, \mathcal{L}^{k-1}(X))}^{1-1/(\sigma+1)} [D^k f]_{C^\sigma(X, \mathcal{L}^k(X))})^{1/(\sigma+1)}, \quad f \in C_b^{k+\sigma}(X), \quad (2.8)$$

while for  $\sigma = 1$  there exists  $K_k > 0$  such that

$$\|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))} \leq K_k (\|D^{k-1} f\|_{L^\infty(X, \mathcal{L}^{k-1}(X))}^{1/2} \|D^{k+1} f\|_{L^\infty(X, \mathcal{L}^{k+1}(X))})^{1/2}, \quad f \in C_b^{k+1}(X), \quad (2.9)$$

(for  $k \geq 2$  it is sufficient to replace  $f$  by  $D^{k-1} f$  and argue as above).

Now we are ready to prove (2.5), by recurrence. We just proved that (2.5) holds for  $k = 2$ . Assume that (2.5) holds for some  $k \geq 2$ . Using first (2.9) and then the recurrence assumption, for every  $f \in C_b^{k+1}(X)$  we have

$$\begin{aligned} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))} &\leq K_k \|D^{k-1} f\|_{L^\infty(X, \mathcal{L}^{k-1}(X))}^{1/2} \|D^{k+1} f\|_{L^\infty(X, \mathcal{L}^{k+1}(X))}^{1/2} \\ &\leq K_k (C_k \|f\|_\infty^{1/k} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))}^{(k-1)/k})^{1/2} \|D^{k+1} f\|_{L^\infty(X, \mathcal{L}^{k+1}(X))}^{1/2} \\ &= K_k C_k^{1/2} \|f\|_\infty^{1/(2k)} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))}^{(k-1)/(2k)} \|D^{k+1} f\|_{L^\infty(X, \mathcal{L}^{k+1}(X))}^{1/2} \end{aligned}$$

so that

$$\|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))} \leq (K_k C_k^{1/2})^{2k/(k+1)} \|f\|_\infty^{1/(k+1)} \|D^{k+1} f\|_{L^\infty(X, \mathcal{L}^{k+1}(X))}^{k/(k+1)}. \quad (2.10)$$

For  $h < k$  we use the recurrence assumption and then (2.10), estimating

$$\begin{aligned} \|D^h f\|_{L^\infty(X, \mathcal{L}^h(X))} &\leq C_k \|f\|_\infty^{1-h/k} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))}^{h/k} \\ &\leq C_k (K_k C_k^{1/2})^{2k/(k+1)} \|f\|_\infty^{1-h/k} (\|f\|_\infty^{1/(k+1)} \|D^{k+1} f\|_{L^\infty(X, \mathcal{L}^{k+1}(X))}^{k/(k+1)})^{h/k} \\ &= C_k^{1+h/(k+1)} K_k^{2h/(k+1)} \|f\|_\infty^{1-h/(k+1)} \|D^{k+1} f\|_{L^\infty(X, \mathcal{L}^{k+1}(X))}^{h/(k+1)}. \end{aligned}$$

Such estimates and (2.10) yield that (2.5) holds for  $k + 1$ .

Now we prove that (2.6) holds. Let  $\beta = k + \sigma$  with  $k \in \mathbb{N}$ ,  $k \geq 2$ , and  $\sigma \in (0, 1)$ , and let  $f \in C_b^\beta(X)$ . Estimates (2.8) and (2.5) with  $h = k - 1$  yield

$$\begin{aligned} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))} &\leq K_{k,\sigma} (\|D^{k-1} f\|_{L^\infty(X, \mathcal{L}^{k-1}(X))}^{1-1/(\sigma+1)} [D^k f]_{C^\sigma(X, \mathcal{L}^k(X))})^{1/(\sigma+1)} \\ &\leq K_{k,\sigma} (C_k \|f\|_\infty^{1/k} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))}^{1-1/k})^{1-1/(\sigma+1)} [D^k f]_{C^\sigma(X, \mathcal{L}^k(X))}^{1/(\sigma+1)} \end{aligned}$$

so that

$$\|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))} \leq (K_{k,\sigma} C_k^{1-1/(\sigma+1)})^{k(\sigma+1)/\beta} \|f\|_\infty^{\sigma/\beta} [D^k f]_{C^\sigma(X, \mathcal{L}^k(X))}^{k/\beta}$$

which gives (2.6) with  $h = k$ . For  $h < k$  we use (2.5) and the above estimate, to get

$$\begin{aligned} \|D^h f\|_{L^\infty(X, \mathcal{L}^h(X))} &\leq C_k \|f\|_\infty^{1-h/k} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))}^{h/k} \\ &\leq C_k (K_{k,\sigma} C_k^{1-1/(\sigma+1)})^{k(\sigma+1)/\beta} \|f\|_\infty^{1-h/\beta} [D^k f]_{C^\sigma(X, \mathcal{L}^k(X))}^{h/\beta}, \end{aligned}$$

and (2.6) is proved.  $\square$

**Corollary 2.3.** *For every  $k \in \mathbb{N}$  the mapping  $f \mapsto \|f\|_\infty + \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))}$  is a norm in  $C_b^k(X)$ , equivalent to the norm (2.1), and for every non integer  $\beta > 0$ ,  $\beta = k + \sigma$  with  $k \in \mathbb{N}$ ,  $\sigma \in (0, 1)$ , the mapping  $f \mapsto \|f\|_\infty + [D^k f]_{C^\sigma(X, \mathcal{L}^k(X))}$  is a norm in  $C_b^\beta(X)$ , equivalent to the norm (2.2). Moreover, for every  $\beta > 1$  and for every  $h \in \mathbb{N}$ ,  $h < \beta$ , the space  $C_b^h(X)$  belongs to the class  $J_{h/\beta}$  between  $C_b(X)$  and  $C_b^\beta(X)$ .*

**Proposition 2.4.** *For every  $\beta > 0$  and  $\gamma \in (0, 1)$  we have*

$$(C_b(X), C_b^\beta(X))_{\gamma, \infty} \subset \begin{cases} C_b^{\gamma\beta}(X), & \gamma\beta \notin \mathbb{N}, \\ Z_b^{\gamma\beta}(X), & \gamma\beta \in \mathbb{N}, \end{cases} \quad (2.11)$$

with continuous embeddings.

*Proof.* By Proposition 2.1, each  $f \in (C_b(X), C_b^\beta(X))_{\gamma, \infty}$  may be represented as  $f = \int_0^\infty \frac{u(t)}{t} dt$ , where  $u \in C((0, +\infty); C_b^\beta(X))$ ,  $t \mapsto t^{1-\gamma}u(t) \in L^\infty((0, +\infty); C_b^\beta(X))$ , and  $t \mapsto t^{-\gamma}u(t) \in L^\infty((0, +\infty); C_b(X))$ . By the norm equivalence of Proposition 2.1 there is  $C = C(\gamma) > 0$  such that

$$(i) \|u(t)\|_{C_b(X)} \leq Ct^\gamma \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}, \quad (ii) \|u(t)\|_{C_b^\beta(X)} \leq Ct^{\gamma-1} \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}, \quad t > 0, \quad (2.12)$$

and by (2.12) and Lemma 4.3 there is  $C = C(\beta, \gamma)$  such that for every  $k \in \mathbb{N}$ ,  $k < \beta$  we have

$$\sup_{x \in X} \|D^k u(t)(x)\|_{\mathcal{L}^k(X)} \leq Ct^{\gamma-k/\beta} \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}, \quad t > 0. \quad (2.13)$$

Having such estimates at our disposal, the proof is now similar to the ones of Theorems 3.8(i) and 3.9(i) of [25].

First we consider the case that  $\gamma\beta$  is not integer. Let  $n = [\gamma\beta]$  be the integral part of  $\gamma\beta$ .

If  $n = 0$ , namely  $\gamma\beta < 1$ , for every  $x, y \in X$  and  $\lambda > 0$  we have

$$|f(x) - f(y)| \leq \int_0^\lambda \frac{|u(t)(x) - u(t)(y)|}{t} dt + \int_\lambda^\infty \frac{|u(t)(x) - u(t)(y)|}{t} dt$$

where by (2.12)(i) the first integral is bounded by

$$2 \int_0^\lambda \frac{\|u(t)\|_\infty}{t} dt \leq 2 \int_0^\lambda t^{\gamma-1} dt C \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} = \frac{2\lambda^\gamma C}{\gamma} \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}.$$

If  $\beta \leq 1$ , by (2.12)(ii) the second integral is bounded by

$$\begin{aligned} \int_\lambda^\infty \frac{\|u(t)\|_{C_b^\beta(X)}}{t} dt \|x - y\|^\beta &\leq \int_\lambda^\infty t^{\gamma-2} dt C \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \|x - y\|^\beta \\ &= \frac{\lambda^{\gamma-1} C}{1-\gamma} \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \|x - y\|^\beta; \end{aligned}$$

while if  $\beta > 1$ , by (2.13) with  $k = 1$  the second integral is bounded by

$$\begin{aligned} \int_\lambda^\infty \frac{\|u(t)\|_{C_b^1(X)}}{t} dt \|x - y\| &\leq \int_\lambda^\infty t^{\gamma-1/\beta-1} dt C \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \|x - y\| \\ &= \frac{\lambda^{\gamma-1/\beta} C}{1/\beta - \gamma} \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \|x - y\|. \end{aligned}$$

In both cases, taking  $\lambda = \|x - y\|^\beta$  we get

$$|f(x) - f(y)| \leq K \|x - y\|^{\gamma\beta},$$

with  $K > 0$  independent of  $f, x, y$ . Therefore  $f \in C_b^{\gamma\beta}(X)$  and  $[f]_{C^{\gamma\beta}(X)} \leq K \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}$ .

Let now  $n \in \mathbb{N}$ . In this case, first we show that  $f \in C_b^n(X)$ , splitting  $f = \int_0^\lambda \frac{u(t)}{t} dt + \int_\lambda^\infty \frac{u(t)}{t} dt$  for every  $\lambda > 0$ . The first integral has values in  $C_b^n(X)$  since  $\gamma > n\beta$ , so that  $t \mapsto u(t)/t \in L^1((0, \lambda); C_b^n(X))$  by (2.13). By (2.12)(ii), the second integral has values in  $C_b^\beta(X)$  and therefore in  $C_b^n(X)$ . In both cases,  $D^n$  commutes with the integral.



Now we prove that  $D^n f \in C_b^{\gamma\beta-n}(X, \mathcal{L}^n(X))$ . As in the case  $n = 0$ , for every  $x, y \in X$  and  $\lambda > 0$  we split and estimate as follows,

$$\begin{aligned} \|(D^n f(x) - D^n f(y))\|_{\mathcal{L}^n(X)} &\leq \int_0^\lambda \frac{\|D^n u(t)(x) - D^n u(t)(y)\|_{\mathcal{L}^n(X)}}{t} dt \\ &\quad + \int_\lambda^\infty \frac{\|D^n u(t)(x) - D^n u(t)(y)\|_{\mathcal{L}^n(X)}}{t} dt. \end{aligned}$$

In the first integral we estimate  $\|D^n u(t)(x) - D^n u(t)(y)\|_{\mathcal{L}^n(X)} \leq 2\|D^n u(t)\|_{L^\infty(X, \mathcal{L}^n(X))}$  and we use (2.13) with  $k = n$ ; in the second integral we estimate  $\|D^n u(t)(x) - D^n u(t)(y)\|_{\mathcal{L}^n(X)} \leq [D^n u(t)]_{C^{\beta-n}(X, \mathcal{L}^n(X))} \|x - y\|^{\beta-n}$  and we use (2.12)(ii) if  $\beta - n < 1$ , we estimate  $\|D^n u(t)(x) - D^n u(t)(y)\|_{\mathcal{L}^n(X)} \leq 2\|D^{n+1} u(t)\|_{L^\infty(X, \mathcal{L}^{n+1}(X))} \|x - y\|$  and we use (2.13) with  $k = n + 1$  if  $\beta - n \geq 1$ , getting respectively

$$\begin{aligned} &\|(D^n f(x) - D^n f(y))\|_{\mathcal{L}^n(X)} \\ &\leq \int_0^\lambda 2t^{\gamma-n/\beta-1} dt C \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} + \int_\lambda^\infty \|x - y\|^{\beta-n} t^{\gamma-2} dt C \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \\ &= C \left( \frac{2\lambda^{\gamma-n/\beta}}{\gamma - n/\beta} + \frac{\|x - y\|^{\beta-n}}{(1-\gamma)\lambda^{1-\gamma}} \right) \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \end{aligned}$$

if  $\beta - n < 1$ , and

$$\begin{aligned} &\|(D^n f(x) - D^n f(y))\|_{\mathcal{L}^n(X)} \\ &\leq \int_0^\lambda 2Ct^{\gamma-n/\beta-1} dt \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} + \int_\lambda^\infty C \|x - y\| t^{\gamma-(n+1)/\beta-1} dt \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \\ &= C \left( \frac{2\lambda^{\gamma-n/\beta}}{\gamma - n/\beta} + \frac{\|x - y\|^{\beta-n}}{((n+1)/\beta - \gamma)\lambda^{(n+1)/\beta-\gamma}} \right) \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \end{aligned}$$

if  $\beta - n \geq 1$  (recall that  $n = \lceil \gamma\beta \rceil$ , so that  $(n+1)/\beta - \gamma > 0$ ). In both cases, taking again  $\lambda = \|x - y\|^\beta$ , we get  $\|(D^n f(x) - D^n f(y))\|_{\mathcal{L}^n(X)} \leq K \|x - y\|^{\beta\gamma-n} \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}$  for some  $K > 0$  independent of  $f, x, y$ . So,  $f \in C_b^{\beta\gamma}(X)$ , and  $(C_b(X), C_b^\beta(X))_{\gamma, \infty}$  is continuously embedded in  $C_b^{\beta\gamma}(X)$  by such estimate and Corollary 2.3.

If  $\gamma\beta = n \in \mathbb{N}$  the proof is similar. For  $n = 1$ , for every  $x, y \in X$  and  $\lambda > 0$ , we estimate

$$\begin{aligned} &|f(x) + f(y) - 2f((x+y)/2)| \leq \\ &\leq \int_0^\lambda \frac{|u(t)(x) + u(t)(y) - 2u((x+y)/2)|}{t} dt + \int_\lambda^\infty \frac{|u(t)(x) + u(t)(y) - 2u((x+y)/2)|}{t} dt. \end{aligned}$$

In the first integral we use estimate (2.12)(i); in the second integral we estimate

$$|u(t)(x) + u(t)(y) - 2u((x+y)/2)| \leq [Du(t)]_{C^{\beta-1}(X, X^*)} \|x - y\|^\beta \leq Ct^{\gamma-1} \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \|x - y\|^\beta$$

by (2.12)(ii) if  $\beta \in (1, 2)$ ; and

$$|u(t)(x) + u(t)(y) - 2u((x+y)/2)| \leq \|D^2 u\|_{L^\infty(X, \mathcal{L}^2(X))} \|x - y\|^2 \leq Ct^{\gamma-2/\beta-1} \|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \|x - y\|^2$$



if  $\beta \geq 2$ , by (2.13) with  $k = 2$ . In the first case we get

$$\begin{aligned} |f(x) + f(y) - 2f((x+y)/2)| &\leq \left(4 \int_0^\lambda t^{\gamma-1} dt + \|x-y\|^\beta \int_\lambda^\infty t^{\gamma-2} dt\right) C\|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \\ &= \left(\frac{4\lambda^\gamma}{\gamma} + \frac{\|x-y\|^\beta}{(1-\gamma)\lambda^{1-\gamma}}\right) C\|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}, \end{aligned}$$

while in the second case we get

$$\begin{aligned} |f(x) + f(y) - 2f((x+y)/2)| &\leq \left(4 \int_0^\lambda t^{\gamma-1} dt + \|x-y\|^2 \int_\lambda^\infty t^{\gamma-2/\beta-1} dt\right) C\|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}} \\ &= \left(\frac{4\lambda^\gamma}{\gamma} + \frac{\|x-y\|^2}{(2/\beta-\gamma)\lambda^{2/\beta-\gamma}}\right) C\|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}. \end{aligned}$$

In both cases choosing  $\lambda = \|x-y\|^\beta$  and recalling that  $\gamma\beta = 1$  we get  $|f(x) + f(y) - 2f((x+y)/2)| \leq K\|f\|_{(C_b(X), C_b^\beta(X))_{\gamma, \infty}}\|x-y\|$  for some  $K$  independent of  $f, x, y$ , so that  $f \in Z_b^1(X)$  and  $(C_b(X), C_b^\beta(X))_{1/\beta, \infty}$  is continuously embedded in  $Z_b^1(X)$ .

If  $n > 1$  the procedure is similar, replacing  $f$  by  $D^{n-1}f$  and distinguishing the cases  $\beta \in (n, n+1)$  and  $\beta \geq n+1$ . The details are left to the reader.  $\square$

It would be desirable to have equivalences instead of embeddings in (2.11), which is true in the case  $X = \mathbb{R}^N$ . In the infinite dimensional case the only characterization of this kind that we are aware of is  $(BUC(X), BUC^1(X))_{\gamma, \infty} = C_b^\gamma(X)$  for every  $\gamma \in (0, 1)$ , proved in [10] when  $X$  is a separable Hilbert space.

### 3. GOOD SEMIGROUPS IN $C_b(X)$

Although the aim of this paper is the study of generalized Mehler semigroups, we start with a few properties satisfied by a larger class of semigroups. Indeed in this section we consider semigroups of bounded operators  $P_t$  in  $C_b(X)$ , such that

$$\|P_t\|_{\mathcal{L}(C_b(X))} \leq Me^{\beta t}, \quad t > 0,$$

for some  $M > 0$ ,  $\beta \in \mathbb{R}$ , and such that the function  $(t, x) \mapsto P_t f(x)$  is continuous in  $[0, +\infty) \times X$ .

Then there exists a closed operator  $L : D(L) \subset C_b(X) \mapsto C_b(X)$ , such that the resolvent set of  $L$  contains  $(\beta, +\infty)$ , and

$$R(\lambda, L)f(x) = \int_0^\infty e^{-\lambda t} P_t f(x) dt, \quad \lambda > \beta, f \in C_b(X), x \in X. \quad (3.1)$$

The (easy) proof of this statement may be found e.g. in [20, §2(c)]. From (3.1) we get immediately

$$\|R(\lambda, L)\|_{\mathcal{L}(C_b(X))} \leq \frac{M}{\lambda - \beta}, \quad \lambda > \beta.$$

Some properties of strongly continuous semigroups are shared by such semigroups. The first one is in the next lemma.

**Lemma 3.1.** *For every  $s > 0$  and  $f \in D(L)$ ,  $P_s f \in D(L)$ , we have  $P_s Lf = LP_s f$ , and*

$$P_t f(x) - f(x) = \int_0^t LP_s f(x) ds = \int_0^t P_s Lf(x) ds, \quad t > 0, x \in X. \quad (3.2)$$

Consequently, for every  $x \in X$  the function  $t \mapsto P_t f(x)$  is differentiable at any  $t \geq 0$  with derivative  $P_t Lf(x)$ , and moreover

$$\|P_t f - f\|_\infty \leq t M \max\{e^\beta, 1\} \|Lf\|_\infty, \quad 0 < t \leq 1. \quad (3.3)$$

*Proof.* By formula (3.1),  $P_s$  commutes with  $R(\lambda, L)$  for every  $\lambda > \beta$ , because it commutes with the Riemann sums  $\frac{1}{nT} \sum_{k=1}^n e^{-kT/n} P_{kT/n} f$ , for every  $T > 0$  and  $n \in \mathbb{N}$ . Therefore it commutes with  $L$  on  $D(L)$ . Fixed any  $f \in D(L)$  and  $\lambda > \beta$  set  $g = \lambda f - Lf$  so that  $f = R(\lambda, L)g$ . For every  $t > 0$  and  $x \in X$  we have

$$\begin{aligned} P_t f(x) - f(x) &= \int_0^\infty e^{-\lambda s} P_{t+s} g(x) ds - \int_0^\infty e^{-\lambda s} P_s g(x) ds \\ &= \int_t^\infty e^{-\lambda(\sigma-t)} P_\sigma g(x) d\sigma - \int_0^\infty e^{-\lambda s} P_s g(x) ds \\ &= (e^{\lambda t} - 1) \int_0^\infty e^{-\lambda \sigma} P_\sigma g(x) d\sigma - \int_0^t e^{-\lambda \sigma} P_\sigma g(x) d\sigma \\ &= (e^{\lambda t} - 1) f(x) - \int_0^t e^{-\lambda \sigma} P_\sigma (\lambda f - Lf)(x) d\sigma. \end{aligned}$$

Although the proof of this formula makes sense only for  $\lambda > \beta$ , the formula holds for every  $\lambda \in \mathbb{C}$  because the right-hand side is a holomorphic function of  $\lambda$ . Letting  $\lambda \rightarrow 0$ , we get  $P_t f(x) - f(x) = \int_0^t (P_\sigma Lf)(x) d\sigma$  and (3.2) follows.

Since  $s \mapsto P_s Lf(x)$  is continuous in  $[0, +\infty)$ , by (3.2)  $t \mapsto P_t f(x)$  is differentiable at any  $t \geq 0$  with derivative  $P_t Lf(x)$ . Estimate (3.3) follows immediately from (3.2).  $\square$

**Corollary 3.2.** *For every  $f \in C_b(X)$  we have*

$$\lim_{t \rightarrow 0} \|P_t f - f\|_\infty = 0 \iff \lim_{\lambda \rightarrow \infty} \|\lambda R(\lambda, L)f - f\|_\infty = 0 \iff f \in \overline{D(L)}.$$

*Proof.* For every  $f \in D(L)$  we have  $\lim_{t \rightarrow 0} \|P_t f - f\|_\infty = \lim_{\lambda \rightarrow \infty} \|\lambda R(\lambda, L)f - f\|_\infty = 0$ , respectively by (3.3) and by the equality  $\lambda R(\lambda, L)f = R(\lambda, L)Lf + f$ . Since there is  $C > 0$  such that  $\|P_t - I\|_{\mathcal{L}(C_b(X))} \leq C$  for every  $t \in (0, 1)$ ,  $\|\lambda R(\lambda, L) - I\|_{\mathcal{L}(C_b(X))} \leq C$  for every  $\lambda > \beta + 1$ , one gets as well  $\lim_{t \rightarrow 0} \|P_t f - f\|_\infty = \lim_{\lambda \rightarrow \infty} \|\lambda R(\lambda, L)f - f\|_\infty = 0$  for every  $f \in \overline{D(L)}$ .

Conversely, if  $\lim_{\lambda \rightarrow \infty} \|\lambda R(\lambda, L)f - f\|_\infty = 0$  then  $f \in \overline{D(L)}$  since  $\lambda R(\lambda, L)f \in D(L)$  for every  $\lambda > 0$ . If  $\lim_{t \rightarrow 0} \|P_t f - f\|_\infty = 0$ , for every  $\varepsilon > 0$  let  $\delta > 0$  be such that  $\|P_t f - f\|_\infty \leq \varepsilon$  for every  $t \in [0, \delta]$ . For each  $\lambda > \beta$  and  $x \in X$  we have

$$\lambda R(\lambda, L)f(x) - f(x) = \left( \int_0^\delta + \int_\delta^\infty \right) \lambda e^{-\lambda t} (P_t f(x) - f(x)) dt,$$

so that  $\|\lambda R(\lambda, L)f - f\|_\infty \leq \delta \varepsilon + \sup_{t \geq \delta} \lambda e^{-\lambda t} (M e^{\beta t} + 1) \|f\|_\infty$ , and therefore  $\lim_{\lambda \rightarrow \infty} \|\lambda R(\lambda, L)f - f\|_\infty = 0$ , which implies  $f \in \overline{D(L)}$ .  $\square$

Now we consider Hölder continuity. It is well known that for every strongly continuous semigroup  $P_t$  in a Banach space  $E$  and for every  $f \in E$ ,  $\alpha \in (0, 1)$ , the function  $t \mapsto P_t f$  belongs to  $C^\alpha([0, T]; E)$  for some/all  $T \in (0, +\infty)$  if and only if  $f \in (E, D(L))_{\alpha, \infty}$ , where  $L$  is the infinitesimal generator of  $P_t$ . This characterization goes back to the pioneering paper by Lions and Peetre [20] and may be found in (almost) any book about interpolation theory. See e.g. [34, Sect. 1.13]. It still holds for analytic semigroups not necessarily strongly continuous at 0 ([21, Sect. 2.2]) and for some non analytic semigroups in spaces of continuous and bounded functions ([22, Ch. 5]). In fact in the proof of the next Proposition 3.3 we use a result from [22].

**Proposition 3.3.** *If  $\beta \leq 0$ , for every  $f \in C_b(X)$  and  $\alpha \in (0, 1)$  the following conditions are equivalent:*

- (i)  $[f]_\alpha^{(1)} := \sup_{t \in (0, 1)} t^{-\alpha} \|P_t f - f\|_\infty < +\infty$ ,
- (ii)  $[f]_\alpha^{(2)} := \sup_{t > 0} t^{-\alpha} \|P_t f - f\|_\infty < +\infty$ ,
- (iii)  $[f]_\alpha^{(3)} := \sup_{\lambda > 0} \|\lambda^\alpha LR(\lambda, L)f\|_\infty < +\infty$ ,
- (iv)  $f \in (C_b(X), D(L))_{\alpha, \infty}$ ,

and the norms  $f \mapsto \|f\|_\infty + [f]_\alpha^{(1)}$ ,  $f \mapsto \|f\|_\infty + [f]_\alpha^{(2)}$ ,  $f \mapsto \|f\|_\infty + [f]_\alpha^{(3)}$  are equivalent to the norm of  $(C_b(X), D(L))_{\alpha, \infty}$ .

If  $\beta > 0$ , for every  $f \in C_b(X)$  and  $\alpha \in (0, 1)$  condition (i) is equivalent to (iv), and the norm  $f \mapsto \|f\|_\infty + [f]_\alpha^{(1)}$  is equivalent to the norm of  $(C_b(X), D(L))_{\alpha, \infty}$ .

*Proof.* Let  $\beta \leq 0$ . Since  $\|P_t\|_{\mathcal{L}(C_b(X))} \leq M$  for every  $t$ , the equivalence between (i) and (ii) is obvious, as well as the inequalities  $[f]_\alpha^{(1)} \leq [f]_\alpha^{(2)} \leq \max\{[f]_\alpha^{(1)}, (M+1)\|f\|_\infty\}$  for every  $f \in C_b(X)$  and  $\alpha \in (0, 1)$ .

Let us prove that (ii) implies (iii). For every  $\lambda > 0$  we have  $LR(\lambda, L)f = \lambda R(\lambda, L)f - f$ , so that for every  $x \in X$

$$|\lambda^\alpha LR(\lambda, L)f(x)| = \left| \int_0^{+\infty} \lambda^{\alpha+1} t^\alpha e^{-\lambda t} \frac{P_t f(x) - f(x)}{t^\alpha} dt \right|$$

Therefore,

$$\|\lambda^\alpha LR(\lambda, L)f\|_\infty \leq \Gamma(\alpha + 1)[f]_\alpha^{(1)}$$

so that (iii) holds, and  $[f]_\alpha^{(3)} \leq \Gamma(\alpha + 1)[f]_\alpha^{(1)}$ .

The equivalence between (iii) and (iv), as well as the relevant norms, follows from [22, Prop. 3.1], where no assumption on the density of  $D(L)$  was made.

If (iv) holds, for every  $t > 0$  and for every decomposition  $f = a + b$ , with  $a \in C_b(X)$ ,  $b \in D(L)$ , we have  $\|P_t b - b\|_\infty \leq t \|Lb\|_\infty$  by (3.3), and therefore

$$\frac{\|P_t f - f\|_\infty}{t^\alpha} \leq \frac{\|P_t a - a\|_\infty}{t^\alpha} + \frac{\|P_t b - b\|_\infty}{t^\alpha} \leq \frac{2\|a\|_\infty}{t^\alpha} + \frac{t\|Lb\|_\infty}{t^\alpha}.$$

Taking the infimum over all the decompositions we get

$$\frac{\|P_t f - f\|_\infty}{t^\alpha} \leq 2t^{-\alpha} K(t, f, C_b(X), D(L)), \quad t > 0,$$

so that (ii) holds, and  $[f]_\alpha^{(2)} \leq 2\|f\|_{(C_b(X), D(L))_{\alpha, \infty}}$ .

If  $\beta > 0$  we consider the semigroup  $S_t : e^{-\beta t} P_t$ , whose generator is  $L_\beta := L - \beta I : D(L) \mapsto C_b(X)$ , and  $R(\lambda, L_\beta) = R(\lambda + \beta, L)$  for  $\lambda > 0$ . So, conditions (i) to (iv) are equivalent for  $S_t$ . Since the graph norm of  $L_\beta$  is equivalent to the graph norm of  $L$ , we still obtain (i)  $\iff$  (iv), with equivalence of the norms  $f \mapsto \|f\|_\infty + [f]_\alpha^{(1)}$ ,  $f \mapsto \|f\|_{(C_b(X), D(L))_{\alpha, \infty}}$ .  $\square$

Using the semigroup law for  $P_t$  yields that for every  $T > 0$  the function  $t \mapsto P_t f =: u(t)$  belongs to  $C^\alpha([0, T]; C_b(X))$  if and only if  $f \in (C_b(X), D(L))_{\alpha, \infty}$ . If  $\beta \leq 0$ , in such a case  $u \in C^\alpha([0, +\infty); C_b(X))$ . We may go on, characterizing the initial functions  $f$  such that  $t \mapsto P_t f$  is more regular.

**Corollary 3.4.** *For every  $f \in C_b(X)$  and  $\alpha \in (0, 1)$ , the function  $t \mapsto P_t f$  belongs to  $C^1([0, +\infty); C_b(X))$  if and only if  $f \in D(L)$  and  $Lf \in \overline{D(L)}$ ; it belongs to  $C^{1+\alpha}([0, T]; C_b(X))$  for some/every  $T > 0$  if and only if  $f \in D(L)$  and  $Lf \in (C_b(X), D(L))_{\alpha, \infty}$ .*

*Proof.* Set  $u(t) := P_t f$ . If  $f \in D(L)$  and  $Lf \in \overline{D(L)}$  then  $s \mapsto P_s Lf$  belongs to  $C([0, +\infty); C_b(X))$ , so that  $u$  belongs to  $C^1([0, +\infty); C_b(X))$  by (3.2) and  $u'(t) = P_t Lf = Lu(t)$ . If in addition  $Lf \in (C_b(X), D(L))_{\alpha, \infty}$ , then  $u' \in C^\alpha([0, T]; C_b(X))$  for every  $T > 0$  and therefore  $u \in C^{1+\alpha}([0, T]; C_b(X))$  for every  $T > 0$ .

Conversely, assume that  $u$  is continuously differentiable. Then  $u'(t) = \lim_{h \rightarrow 0^+} P_t((P_h f - f)/h) = P_t u'(0)$ . Since  $u'$  is continuous,  $u'(0) \in \overline{D(L)}$  by Corollary 3.2, and moreover integrating by parts in (1.4) we get, for every  $\lambda > \beta$ ,

$$\lambda R(\lambda, L)f = \int_0^\infty e^{-\lambda t} u'(t) dt + f = R(\lambda, L)u'(0) + f$$

which yields  $f = \overline{R(\lambda, L)}(\lambda f - u'(0)) \in D(L)$  and  $R(\lambda, L)Lf = \lambda R(\lambda, L)f - f = R(\lambda, L)u'(0)$  so that  $Lf = u'(0) \in \overline{D(L)}$ . If in addition  $u'(t) = P_t Lf$  belongs to  $C^\alpha([0, T]; C_b(X))$  for some  $T > 0$ , then  $Lf \in (C_b(X), D(L))_{\alpha, \infty}$  by Proposition 3.3 and therefore  $u \in C^{1+\alpha}([0, T]; C_b(X))$  for every  $T > 0$ .  $\square$

#### 4. GENERALIZED MEHLER SEMIGROUPS

Throughout this section  $P_t$  is the generalized Mehler semigroup defined by (1.1), where  $T_t$  is any strongly continuous semigroup of bounded operators on  $X$ , and  $\{\mu_t : t \geq 0\}$  is a family of Borel probability measures in  $X$  such that  $\mu_0 = \delta_0$ ,  $t \mapsto \mu_t$  is weakly continuous in  $[0, +\infty)$  and (1.2) holds.

Notice that

$$\|P_t\|_{\mathcal{L}(C_b(X))} = 1, \quad t > 0, \quad \|R(\lambda, L)\|_{\mathcal{L}(C_b(X))} = \frac{1}{\lambda}, \quad \lambda > 0. \quad (4.1)$$

The inequalities  $\leq$  are an immediate consequence of (1.1), the equalities follow taking  $f \equiv 1$ . Another immediate consequence of the representation formula (1.1) is that  $P_t$  preserves  $C_b^k(X)$  for every  $k \in \mathbb{N}$ . In particular,

$$DP_t f(x) = T_t^* \int_X Df(T_t x + y) \mu_t(dy), \quad f \in C_b^1(X), \quad x \in X,$$

and therefore there are  $M > 0$ ,  $\omega \in \mathbb{R}$  such that

$$\|DP_t f\|_{L^\infty(X, X^*)} \leq M e^{\omega t} \|Df\|_{L^\infty(X, X^*)}, \quad t > 0, \quad f \in C_b^1(X). \quad (4.2)$$

Similarly, for every  $k \in \mathbb{N}$ ,

$$\|D^k P_t f\|_{L^\infty(X, \mathcal{L}^k(X))} \leq M e^{k\omega t} \|D^k f\|_{L^\infty(X, \mathcal{L}^k(X))}, \quad t > 0, \quad f \in C_b^k(X). \quad (4.3)$$

It is not hard to check that for every  $f \in C_b(X)$  the function  $(t, x) \mapsto P_t f(x)$  is continuous in  $[0, +\infty) \times X$ . See e.g. [7, Lemma 2.1]. Therefore the results of Section 3 hold for  $P_t$ .

Notice that the semigroup  $S_t$  defined by  $S_t f(x) := f(T_t x)$  is itself a generalized Mehler semigroup, with  $\mu_t = \delta_0$  for every  $t$ . We denote by  $L_0$  its generator, and by  $Y_0$  the subspace of strong continuity of  $S_t$ , namely

$$Y_0 := \{f \in C_b(X) : \lim_{t \rightarrow 0} \|f(T_t \cdot) - f\|_\infty = 0\}.$$

For  $\alpha \in (0, 1)$  we denote by  $Y_\alpha$  the subspace of  $\alpha$ -Hölder continuity of  $S_t$ , namely

$$Y_\alpha := \left\{ f \in C_b(X) : [f]_{Y_\alpha} := \sup_{t > 0} \frac{\|f(T_t \cdot) - f\|_\infty}{t^\alpha} < +\infty \right\},$$

endowed with the norm

$$\|f\|_{Y_\alpha} := \|f\|_\infty + [f]_{Y_\alpha}.$$

By Proposition 3.3 it coincides with the interpolation space  $(C_b(X), D(L_0))_{\alpha, \infty}$ , and the respective norms are equivalent.

Notice that if  $g \in R(\lambda, L)(C_b^1(X))$  for some  $\lambda > 0$  and  $x \in D(A)$ , integrating by parts in (1.4) we get  $L_0g(x) = g'(x)(Ax)$ , so that  $L_0$  is a realization of a drift operator.

Going back to the general case, Corollary 3.2 gives an abstract characterization of the subspace of strong continuity of  $P_t$ . Under some additional assumptions we prove an explicit characterization of it.

**Proposition 4.1.** *Assume that  $P_t$  maps  $C_b(X)$  into  $BUC(X)$ , for every  $t > 0$ . Then*

$$\overline{D(L)} = BUC(X) \cap Y_0.$$

*Proof.* Let  $f \in BUC(X)$ . Fixed any  $\varepsilon > 0$  let  $r$  be such that  $|f(z+y) - f(y)| \leq \varepsilon$  for every  $z \in X$  and  $y \in B(0, r)$ . Then for every  $t > 0$  and  $x \in X$  we have

$$\begin{aligned} |P_t f(x) - f(T_t x)| &\leq \int_X |f(T_t x + y) - f(T_t x)| \mu_t(dy) \\ &\leq \left( \int_{B(0, r)} + \int_{X \setminus B(0, r)} \right) |f(T_t x + y) - f(T_t x)| \mu_t(dy) \\ &\leq \varepsilon + 2\|f\|_\infty \mu_t(X \setminus B(0, r)). \end{aligned}$$

Since  $\mu_t$  weakly converges to  $\delta_0$ ,  $\lim_{t \rightarrow 0} \mu_t(X \setminus B(0, r)) = 0$ . Therefore, for  $t$  small enough we have  $|P_t f(x) - f(T_t x)| \leq 2\varepsilon$ , so that

$$\lim_{t \rightarrow 0} \|P_t f - f(T_t \cdot)\|_\infty = 0, \quad f \in BUC(X). \quad (4.4)$$

Now, let  $f \in \overline{D(L)}$ . Then  $\|P_t f - f\|_\infty \rightarrow 0$  as  $t \rightarrow 0$  by Corollary 3.2, and since  $P_t f \in BUC(X)$  for  $t > 0$ , also  $f \in BUC(X)$ . Moreover,  $\|f(T_t \cdot) - f\|_\infty \leq \|f(T_t \cdot) - P_t f\|_\infty + \|P_t f - f\|_\infty$  vanishes as  $t \rightarrow 0$  by (4.4) and Corollary 3.2.

Conversely, if  $f \in BUC(X) \cap Y_0$  then  $\lim_{t \rightarrow 0} \|P_t f - f\|_\infty \rightarrow 0$  by (4.4).  $\square$

As for strong continuity, also for Hölder continuity we give an explicit characterization under further assumptions. Preliminarily we recall some definitions and properties of Borel measures in Banach spaces, taken from [6].

A Borel probability measure  $\mu$  in  $X$  is called *Fomin differentiable* along  $v \in X$  if for every Borel set  $B$  the incremental ratio  $(\mu(B + tv) - \mu(B))/t$  has finite limit as  $t \rightarrow 0$ . Such a limit is a signed measure, called  $d_v \mu(B)$ ; setting  $\mu_v(B) := \mu(B + v)$  for every Borel set  $B$  we have  $\lim_{t \rightarrow 0} \left\| \frac{\mu_{tv} - \mu}{t} - d_v \mu \right\| = 0$ , where  $\|\cdot\|$  denotes the total variation norm.

Moreover,  $d_v \mu$  is absolutely continuous with respect to  $\mu$ . The density  $\beta_v^\mu \in L^1(X, \mu)$  is called *Fomin derivative* of  $\mu$  along  $v$ , and it satisfies

$$\int_X \frac{\partial f}{\partial v} \mu(dx) = - \int_X \beta_v^\mu f \mu(dx), \quad f \in C_b^1(X). \quad (4.5)$$

Conversely, if (4.5) holds for some  $\beta_v^\mu \in L^1(X, \mu)$  and for every  $f \in C_b^1(X)$ , then  $\mu$  is Fomin differentiable along  $v$  ([6, Thm. 3.6.8]).

Under Hypothesis 1.1 it was proved in [25] that  $P_t f$  has Gateaux derivatives of any order, for every  $t > 0$  and for every  $f \in C_b(X)$ . To be more precise, in [25] it was assumed that  $X$  is separable; however this condition was used only to guarantee that every Borel measure in  $X$  is Radon, in order to apply the results on differentiable measures contained in [6, §3.3]. Here we prove better smoothing properties, showing that  $P_t$  is strong-Feller (namely, that it maps  $B_b(X)$  into  $C_b(X)$ ), and that all the Gateaux derivatives of  $P_t f$  are in fact Fréchet derivatives.

**Lemma 4.2.** *Let Hypothesis 1.1 hold. For each  $f \in B_b(X)$ ,  $P_t f \in C_b^k(X)$  for every  $t > 0$  and  $k \in \mathbb{N}$ . Moreover for every  $k \in \mathbb{N}$ , the function  $(t, x) \mapsto P_t f(x)$  belongs to  $C((0, +\infty) \times X)$  if  $f \in B_b(X)$ , and to  $C([0, +\infty) \times X)$  if  $f \in C_b(X)$ .*

*Proof.* Fix  $f \in B_b(X)$ . For every  $x, h \in X$  we have

$$P_t f(x+h) = \int_X f(T_t x + T_t h + y) \mu_t(dy) = \int_X f(T_t x + z) (\mu_t)_{T_t h}(dz),$$

so that

$$|P_t f(x+h) - P_t f(x)| = \left| \int_X f(T_t x + z) ((\mu_t)_{T_t h}(dz) - \mu_t(dz)) \right| \leq \|f\|_\infty \|(\mu_t)_{T_t h} - \mu_t\|.$$

By [6, Thm. 3.3.7(i)] we have  $\|(\mu_t)_{T_t h} - \mu_t\| \leq \|\beta_{t,h}\|_{L^1(X, \mu_t)}$ , so that

$$|P_t f(x+h) - P_t f(x)| \leq \frac{C e^{\omega t}}{t^\theta} \|h\|, \quad t > 0, h \in X.$$

Therefore,  $P_t f$  is Lipschitz continuous for  $t > 0$ . Let us prove that in fact it belongs to  $C_b^k(X)$  for every  $k \in \mathbb{N}$ .

In [25] we remarked that if  $g \in BUC(X)$ , then  $P_t g$  is Fréchet differentiable at every  $x \in X$ , since the Gateaux derivative is continuous with values in  $X^*$ . Now we extend this remark to derivatives of arbitrary order  $n$ , through the representation formula for the  $n$ -th order Gateaux derivative  $D_G^n P_t g$  of [25, Prop. 3.3(ii)],

$$\begin{aligned} D_G^n P_t g(x)(h_1, \dots, h_n) &= (-1)^n \int_X \cdots \int_X g(T_t x + T_{\frac{n-1}{n}t} y_1 + \cdots + T_{\frac{1}{n}t} y_{n-1} + y_n) \cdot \\ &\quad \cdot \beta_{t/n, h_n}(y_n) \cdots \beta_{t/n, h_1}(y_1) \mu_{t/n}(dy_n) \cdots \mu_{t/n}(dy_1). \end{aligned}$$

So, for  $x, x_0 \in X$  we have

$$\begin{aligned} &|(D_G^n P_t g(x) - D_G^n P_t g(x_0))(h_1, \dots, h_n)| \leq \\ &\int_X \cdots \int_X |g(T_t x + T_{\frac{n-1}{n}t} y_1 + \cdots + T_{\frac{1}{n}t} y_{n-1} + y_n) - g(T_t x_0 + T_{\frac{n-1}{n}t} y_1 + \cdots + T_{\frac{1}{n}t} y_{n-1} + y_n)| \cdot \\ &\quad \cdot |\beta_{t/n, h_n}(y_n)| \cdots |\beta_{t/n, h_1}(y_1)| \mu_{t/n}(dy_n) \cdots \mu_{t/n}(dy_1) \\ &\leq (C e^{\omega t/n} n^\theta t^{-\theta})^n \|h_1\| \cdots \|h_n\| \sup_{z \in X} |g(T_t x + z) - g(T_t x_0 + z)| \end{aligned}$$

By the uniform continuity of  $g$ ,  $D_G^n P_t g$  is continuous at  $x_0$  with values in  $\mathcal{L}^n(X)$ , and therefore it is a Fréchet derivative.

Now, let  $f \in B_b(X)$ . For  $t > 0$ , the function  $g = P_{t/2} f$  belongs to  $BUC(X)$ , by the first part of the proof, and therefore  $P_t f = P_{t/2} g$  has Fréchet derivatives of any order.

The continuity of the function  $[0, +\infty) \times X \mapsto \mathbb{R}$ ,  $(t, x) \mapsto P_t f(x)$  if  $f \in C_b(X)$  was proved in [7, Lemma 2.1]. If  $f \in B_b(X)$ ,  $P_{t_0} f \in C_b(X)$  for every  $t_0 > 0$ , and therefore  $(t, x) \mapsto P_t f(x) = (P_{t-t_0} P_{t_0} f)(x)$  is continuous in  $[t_0, +\infty) \times X$ . Since  $t_0$  is arbitrary,  $(t, x) \mapsto P_t f(x) \in C((0, +\infty) \times X)$ .  $\square$

By Lemma 4.2, the pointwise estimates on the Gateaux derivatives of  $P_t f$  in [25] are in fact estimates on the Fréchet derivatives. They amount to

$$\|D^n P_t f(x)\|_{\mathcal{L}^n(X)} \leq K_n \max\{1, t^{-n\theta}\} \|f\|_\infty, \quad t > 0, x \in X, f \in B_b(X), \quad (4.6)$$

for some  $K_n > 0$  ([25, Prop. 3.3(ii)]). By Lemma 2.1(i) of [25], with  $Y = \mathcal{L}^n(X)$ , we obtain estimates for Hölder seminorms, namely for every  $n \in \mathbb{N}$ ,  $\alpha \in (0, 1)$  there exists  $K_{n,\alpha} > 0$  such that

$$[D^n P_t f]_{C^\alpha(X, \mathcal{L}^n(X))} \leq K_{n,\alpha} \max\{1, t^{-(n+\alpha)\theta}\} \|f\|_\infty, \quad t > 0, x \in X, f \in B_b(X). \quad (4.7)$$

Moreover, Lemma 4.2 and Theorem 3.8(i) of [25] yield

$$D(L) \subset \begin{cases} C_b^{1/\theta}(X), & 1/\theta \notin \mathbb{N}, \\ Z_b^{1/\theta}(X), & 1/\theta \in \mathbb{N}, \end{cases} \quad (4.8)$$

with continuous embedding.

**Proposition 4.3.** *If Hypothesis 1.1 holds, we have*

$$(C_b(X), D(L))_{\alpha, \infty} \subset \begin{cases} C_b^{\alpha/\theta}(X), & \alpha/\theta \notin \mathbb{N}, \\ Z_b^{\alpha/\theta}(X), & \alpha/\theta \in \mathbb{N}, \end{cases}$$

with continuous embedding.

*Proof.* If  $1/\theta \notin \mathbb{N}$ , the statement is a direct consequence of the embedding (4.8) and of Proposition 2.4, recalling that if  $F_1 \subset F_2 \subset E$  with continuous embeddings then  $(E, F_1)_{\alpha, \infty} \subset (E, F_2)_{\alpha, \infty}$  with continuous embedding, for every  $\alpha \in (0, 1)$ .

If  $1/\theta \in \mathbb{N}$ , we show that for every non integer  $\beta \in (0, 1/\theta)$  the space  $C_b^\beta(X)$  belongs to the class  $J_{\theta\beta}$  between  $C_b(X)$  and  $D(L)$ .

Let  $\beta = k + \sigma$ , with  $k \in \mathbb{N} \cup \{0\}$  and  $\sigma \in (0, 1)$ .

For every  $\varphi \in D(L) \setminus \{0\}$ , set  $f = \lambda\varphi - (L - I)\varphi$ , so that  $\varphi(x) = \int_0^\infty e^{-(\lambda+1)t} P_t f(x) dt$  for every  $x \in X$ . By proposition 3.7 of [25] we have  $D^j \varphi(x)(h_1, \dots, h_j) = \int_0^\infty e^{-(\lambda+1)t} (D^h P_t f)(x)(h_1, \dots, h_j) dt$  for every  $j \leq k$ ,  $(h_1, \dots, h_j) \in X^j$  and  $x \in X$ . By estimates (4.6), for every  $j \leq k$  we have

$$\begin{aligned} |D^j \varphi(x)(h_1, \dots, h_j)| &\leq \left( \int_0^1 \frac{e^{-(\lambda+1)t}}{t^{\theta j}} dt + \int_1^\infty e^{-(\lambda+1)t} dt \right) K_j \|f\|_\infty \prod_{l=1}^j \|h_l\| \\ &\leq \left( \frac{\Gamma(1 - \theta j)}{(\lambda + 1)^{(1 - \theta j)}} + \frac{1}{\lambda + 1} \right) K_j \|f\|_\infty \prod_{l=1}^j \|h_l\| \end{aligned}$$

so that there exists  $C_1 > 0$  such that  $\|D^j \varphi\|_{L^\infty(X, \mathcal{L}^j(X))} \leq C_1 \lambda^{\theta\beta-1} \|f\|_\infty$  for every  $j \leq k$ . Similarly, using (4.7) with  $n = k$  and  $\alpha = \sigma$ , we get  $[D^k \varphi]_{C^\sigma(X, \mathcal{L}^k(X))} \leq C_2 \lambda^{\theta\beta-1} \|f\|_\infty$  for some  $C_2 > 0$ . Summing up, there exists  $C > 0$  such that

$$\|\varphi\|_{C_b^\beta(X)} \leq C \lambda^{\theta\beta-1} \|f\|_\infty \leq C(\lambda^{\theta\beta} \|\varphi\|_\infty + \lambda^{\theta\beta-1} \|(L - I)\varphi\|_\infty), \quad \lambda > 0.$$

Choosing  $\lambda = \|(L - I)\varphi\|_\infty / \|\varphi\|_\infty$  we get

$$\|\varphi\|_{C_b^\beta(X)} \leq 2C \|(L - I)\varphi\|_\infty^{\theta\beta} \|\varphi\|_\infty^{1-\theta\beta} \leq 2C \|\varphi\|_{D(L)}^{\theta\beta} \|\varphi\|_\infty^{1-\theta\beta},$$

so that  $C_b^\beta(X)$  belongs to the class  $J_{\theta\beta}$  between  $C_b(X)$  and  $D(L)$ .

Now we fix  $\beta \in (\alpha/\theta, 1/\theta)$ . By the Reiteration Theorem,  $(C_b(X), D(L))_{\alpha, \infty}$  is continuously embedded in  $(C_b(X), C_b^\beta(X))_{\alpha/(\theta\beta), \infty}$ . In its turn,  $(C_b(X), C_b^\beta(X))_{\alpha/(\theta\beta), \infty}$  is continuously embedded in  $C_b^{\alpha/\theta}(X)$  or in  $Z_b^{\alpha/\theta}(X)$  by Proposition 2.4, and the statement follows.  $\square$

Under further (optimal) estimates on the moments of  $\mu_t$  the spaces  $(C_b(X), D(L))_{\alpha, \infty}$  may be characterized.



**Proposition 4.4.** *Let Hypothesis 1.1 hold, and assume in addition that there exist  $\gamma, C > 0$  such that*

$$\int_X \|x\|^\gamma \mu_t(dx) \leq Ct^{\gamma\theta}, \quad 0 < t \leq 1. \quad (4.9)$$

Then for every  $\alpha \in (0, 1 \wedge \theta) \cap (0, \gamma\theta]$  we have

$$(C_b(X), D(L))_{\alpha, \infty} = C_b^{\alpha/\theta}(X) \cap Y_\alpha, \quad (4.10)$$

and the respective norms are equivalent.

If  $\theta < 1$  and moreover the measures  $\mu_t$  are centered for every  $t > 0$ , (4.10) holds for every  $\alpha \in (0, 1 \wedge 2\theta) \cap (0, \gamma\theta]$ ,  $\alpha \neq \theta$ , still with equivalence of the respective norms.

*Proof.* Proposition 4.3 yields the embeddings  $\subset$ . Concerning the embeddings  $\supset$ , since  $\alpha/\theta \leq \gamma$ , assumption (4.9) and the Hölder inequality yield

$$\int_X \|y\|^{\alpha/\theta} \mu_t(dy) \leq Ct^\alpha, \quad 0 < t \leq 1. \quad (4.11)$$

Fix  $f \in C_b^{\alpha/\theta}(X) \cap Y_\alpha$ . If  $\alpha < \theta$ , for every  $t \in (0, 1]$  and  $x \in X$  using (4.11) we get

$$|P_t f(x) - f(T_t x)| \leq \int_X |f(T_t x + y) - f(T_t x)| \mu_t(dy) = \int_X [f]_{C^{\alpha/\theta}} \|y\|^{\alpha/\theta} \mu_t(dy) \leq C \|f\|_{C_b^{\alpha/\theta}(X)} t^\alpha.$$

Since  $f \in Y_\alpha$ ,

$$|P_t f(x) - f(x)| \leq |P_t f(x) - f(T_t x)| + |f(T_t x) - f(x)| \leq C \|f\|_{C_b^{\alpha/\theta}(X)} t^\alpha + [f]_{Y_\alpha} t^\alpha, \quad 0 < t \leq 1, x \in X, \quad (4.12)$$

and the first statement follows from Proposition 3.3.

Let now  $\theta < 1$  and assume that the measures  $\mu_t$  are centered for every  $t > 0$ . Then,  $\int_X \varphi(y) \mu_t(dy) = 0$  for every  $\varphi \in X^*$  and  $t > 0$ . Take  $\alpha \in (0, 1) \cap (\theta, 2\theta) \cap (0, \gamma\theta]$ . For every  $f \in C_b^{\alpha/\theta}(X) \cap Y_\alpha$  and  $t \in (0, 1]$ ,  $x \in X$ , we have

$$|P_t f(x) - f(T_t x)| = \left| \int_X f(T_t x + y) - f(T_t x) - Df(T_t x)(y) \mu_t(dy) \right|,$$

so that, using the estimate

$$|f(z + y) - f(z) - Df(z)(y)| = \left| \int_0^1 (Df(z + \sigma y)(y) - Df(z)(y)) d\sigma \right| \leq [Df]_{C^{\alpha/\theta-1}(X, X^*)} \|y\|^{\alpha/\theta},$$

that holds for every  $z, y \in X$ , and (4.11), we get

$$|P_t f(x) - f(T_t x)| \leq [Df]_{C^{\alpha/\theta-1}(X, X^*)} \int_X \|y\|^{\alpha/\theta} \mu_t(dy) \leq Ct^\alpha \|f\|_{C_b^{\alpha/\theta}(X)},$$

and, as before, (4.12) holds, so that  $f \in (C_b(X), D(L))_{\alpha, \infty}$  by Proposition 3.3.  $\square$

## 5. EXAMPLES

**5.1. Ornstein-Uhlenbeck operators in infinite dimension.** Here  $X$  is an infinite dimensional separable Banach space and the measures  $\mu_t$  are Gaussian and centered. We refer to the book [5] for the general theory of Gaussian measures in Banach spaces.

Ornstein-Uhlenbeck semigroups are defined as follows. We fix a strongly continuous semigroup  $T_t$  in  $X$  generated by a linear operator  $A : D(A) \subset X \mapsto X$ , and an operator  $Q \in \mathcal{L}(X^*, X)$  which is non-negative (namely,  $f(Qf) \geq 0$  for every  $f \in X^*$ ) and symmetric (namely,  $f(Qg) = g(Qf)$  for every  $f, g \in X^*$ ). We assume that the operators  $Q_t$  defined by

$$Q_t := \int_0^t T_s Q T_s^* ds, \quad t > 0,$$

are Gaussian covariances, and for all  $t > 0$  we denote by  $\mu_t$  the Gaussian measure with mean 0 and covariance  $Q_t$ . In this case the measures  $\mu_t$  satisfy (1.2), and taking  $\mu_0 := \delta_0$  the mapping  $t \mapsto \mu_t$  is weakly continuous in  $[0, +\infty)$ . The semigroup  $P_t$  defined by (1.1) is called *Ornstein-Uhlenbeck semigroup*.

As well known, if  $X$  is a Hilbert space any bounded self-adjoint, non-negative operator is the covariance of a Gaussian measure if and only if its trace is finite. If  $X$  is just a Banach space, establishing whether a given non-negative symmetric operator in  $\mathcal{L}(X^*, X)$  is a Gaussian covariance is not as simple; necessary and sufficient conditions are in [35].

The theory of Ornstein-Uhlenbeck operators and semigroups in infinite dimensional spaces is very rich; we refer to the book [12] for the basic theory in Hilbert spaces, to the survey paper [16] for the theory in Banach spaces, and to the more recent survey paper [20] for an up-to-date account.

Hypothesis 1.1 is satisfied provided  $T_t(X)$  is contained in the Cameron-Martin space  $H_t$  of  $\mu_t$  for every  $t > 0$ , and there exist  $C, \theta > 0, \omega \in \mathbb{R}$  such that

$$\|T_t x\|_{H_t} \leq \frac{C e^{\omega t}}{t^\theta} \|x\|, \quad t > 0, x \in X. \quad (5.1)$$

See [25, §5.1] for more details. Propositions 4.1 and 4.3 yield

**Proposition 5.1.** *Assume that (5.1) holds. Then*

$$\overline{D(L)} = BUC(X) \cap Y_0.$$

Moreover, for every  $\alpha \in (0, 1)$ ,

$$(C_b(X), D(L))_{\alpha, \infty} \subset \begin{cases} C_b^{\alpha/\theta}(X), & \alpha/\theta \notin \mathbb{N}, \\ Z_b^{\alpha/\theta}(X), & \alpha/\theta \in \mathbb{N}, \end{cases}$$

with continuous embedding.

Therefore, for any  $f \in C_b(X)$  and  $\alpha \in (0, 1)$ ,  $\alpha/\theta \notin \mathbb{N}$  we have

$$\sup_{t>0, x \in X} \frac{|P_t f(x) - f(x)|}{t^\alpha} < \infty \implies f \in C_b^{\alpha/\theta}(X),$$

while if  $\alpha/\theta \in \mathbb{N}$ ,

$$\sup_{t>0, x \in X} \frac{|P_t f(x) - f(x)|}{t^\alpha} < \infty \implies f \in Z_b^{\alpha/\theta}(X).$$

We recall that if  $X$  is a Hilbert space the Cameron-Martin space of  $\mu_t$  is just the range of  $Q_t^{1/2}$ , therefore (5.1) holds if and only if  $T_t(X) \subset Q_t^{1/2}(X)$  for every  $t$  and  $\sup_{0 < t \leq 1} t^\theta \|Q_t^{-1/2} T_t\|_{\mathcal{L}(X)} < +\infty$ , where  $Q_t^{-1/2}$  is the pseudo-inverse of  $Q_t$ .

Basic examples such that (5.1) holds, with any  $\theta \geq 1/2$ , were given in [12, Ex. 6.2.11]. A general family of Ornstein-Uhlenbeck semigroups such that (5.1) holds with  $\theta = 1/2$  was considered in [2], where it was proved that if  $t \mapsto P_t f \in C^\alpha([0, +\infty); C_b(X))$  with  $\alpha < 1/2$ , then  $f \in C_b^{2\alpha}(X)$ . The norm  $\|f\|_\infty + [f]_\alpha^{(2)}$  was called “semigroup norm” there.

However, in such examples condition (4.9) is not satisfied if  $X$  is infinite dimensional, for any  $\gamma > 0$ , so that Proposition 4.4 is not applicable.

Instead, if  $X = \mathbb{R}^N$  and  $Q = Q^* > 0$ , Hypothesis 1.1 holds with  $\theta = 1/2$  and (4.9) holds for any  $\gamma > 0$ . The equalities  $\overline{D(L)} = BUC(\mathbb{R}^N) \cap Y_0$  and  $(C_b(\mathbb{R}^N), D(L))_{\alpha, \infty} = C_b^{2\alpha}(\mathbb{R}^N) \cap Y_\alpha$  for  $\alpha \in (0, 1) \setminus \{1/2\}$ ,  $(C_b(\mathbb{R}^N), D(L))_{1/2, \infty} = Z_b^1(\mathbb{R}^N) \cap Y_{1/2}$  were proved in [11], so that applying Proposition 4.4 does not give any new information.

**5.2. Ornstein-Uhlenbeck operators with fractional diffusion in finite dimension.** Here we take  $X = \mathbb{R}^N$  and we fix a symmetric positive definite matrix  $Q$ , a matrix  $A$ , and any  $s \in (0, 1)$ . The corresponding Ornstein-Uhlenbeck operator with fractional diffusion is given by

$$(\mathcal{L}u)(x) = \frac{1}{2}(\text{Tr}^s(QD^2u))(x) + \langle Ax, \nabla u(x) \rangle, \quad x \in \mathbb{R}^N. \quad (5.2)$$

where  $\text{Tr}^s(QD^2)$  is the pseudo-differential operator with symbol  $-\langle Q\xi, \xi \rangle^s$ . The associated semi-group is given by (e.g., [3, 30])

$$P_t f(x) = \int_{\mathbb{R}^N} f(e^{tA}x + y)g_t(y)dy, \quad t > 0, f \in C_b(\mathbb{R}^N), x \in \mathbb{R}^N,$$

where

$$g_t(y) = \frac{1}{(2\pi)^N} \int_{\mathbb{R}^N} e^{-\frac{1}{2} \int_0^t \|Q^{1/2} e^{\sigma A^*} \xi\|^2 d\sigma} e^{-i\langle \xi, y \rangle} d\xi, \quad t > 0, y \in \mathbb{R}^N. \quad (5.3)$$

So,  $P_t$  is written in the form (1.1), with  $T_t = e^{-tA}$ ,  $\mu_t(dy) := g_t(y)dy$ , for  $t > 0$ . For  $t = 0$  we set  $P_0 = I$  and  $\mu_0 = \delta_0$ .

It has been checked in [25, Sect. 4.2] that  $P_t$  is a generalized Mehler semigroup, and moreover the function  $g_t$  belongs to  $W^{1,1}(\mathbb{R}^N)$  for every  $t > 0$  and

$$\sup_{0 < t \leq 1} t^{1/(2s)} \left\| \frac{\partial g_t}{\partial x_k} \right\|_{L^1(\mathbb{R}^N)} < +\infty, \quad k = 1, \dots, N, \quad (5.4)$$

so that Hypothesis 1.1 is satisfied with  $\theta = 1/(2s)$ . Moreover, since  $g_t(y) = g_t(-y)$  for every  $y \in \mathbb{R}^N$  and  $t > 0$ , then each  $\mu_t$  is centered. Corollary 3.2 and Proposition 4.1 yield

**Proposition 5.2.** *For every  $f \in C_b(\mathbb{R}^N)$ , we have*

$$\lim_{t \rightarrow 0} \|P_t f - f\|_\infty = 0 \iff f \in \overline{D(L)} \iff f \in BUC(\mathbb{R}^N) \text{ and } \lim_{t \rightarrow 0} \|f(e^{tA} \cdot) - f\|_\infty = 0.$$

To study Hölder continuity of  $t \mapsto P_t f$  through Proposition 4.4 it remains to check that (4.9) holds, for some  $\gamma > 0$ .

**Lemma 5.3.** *For every  $\gamma < 2s$  there is  $C = C(\gamma)$  such that*

$$\int_{\mathbb{R}^N} \|y\|^\gamma g_t(y) dy \leq C t^{\gamma/(2s)}, \quad 0 < t \leq 1. \quad (5.5)$$

*Proof.* By the change of variables  $Q^{1/2}\xi = \eta$  in (5.3) we obtain

$$g_t(y) = \frac{1}{(2\pi)^N (\text{Det } Q)^{1/2}} \int_{\mathbb{R}^N} e^{-\frac{1}{2} \int_0^t \|e^{\sigma B^*} \eta\|^2 d\sigma} e^{-i\langle \xi, Q^{-1/2}y \rangle} d\xi,$$

with  $B = Q^{1/2}AQ^{-1/2}$ , and the proof of (5.5) is reduced to the case  $Q = I$ , just using the inverse change of variables  $Q^{-1/2}y = x$  in the left-hand side of (5.5).

So, without loss of generality we may assume  $Q = I$ , in which case  $P_t$  is the transition semigroup of a stochastic differential equation,

$$dX_t = AX_t + dL_t, \quad X_0 = x, \quad (5.6)$$

where  $\{L_t : t \geq 0\}$  is a  $2s$ -stable standard Lévy process, whose laws  $\nu_t$  have Fourier transforms  $\widehat{\nu}_t(h) = e^{-t|h|^{2s}}$  for every  $t > 0$ . See [30] for this explicit example, and [4, 32] for the general theory of Lévy processes. So, we have

$$P_t f(x) = \mathbb{E}(f(X_t)), \quad t > 0, \quad (5.7)$$

$X_t(x)$  being the (unique) mild solution to (5.6), and

$$\int_{\mathbb{R}^N} |y|^\gamma g_t(y) dy = P_t(|\cdot|^\gamma)(0) = \mathbb{E}(|X_t(0)|^\gamma), \quad t > 0.$$

Estimates for the above integrals are classical if  $A = 0$ , in which case  $X_t = x + L_t$  and  $\int_{\mathbb{R}^N} |y|^\gamma g_t(y) dy = \mathbb{E}(|L_t|^\gamma) \leq ct^{\gamma/(2s)}$  for  $\gamma < 2s$ , while  $\int_{\mathbb{R}^N} |y|^\gamma g_t(y) dy = +\infty$  for  $\gamma \geq 2s$  (e.g., [32, Ex. 25.10]).

In the general case, by (5.6) (with  $x = 0$ ) we obtain  $X_t - L_t = \int_0^t AX_\tau d\tau$ , so that  $|X_t| \leq |L_t| + \|A\| \int_0^t |X_\tau| d\tau$ , and by the Gronwall Lemma there exists  $C > 0$  such that

$$|X_t| \leq C \sup_{0 < \tau \leq t} |L_\tau|, \quad 0 < t \leq 1,$$

and therefore

$$\mathbb{E}(|X_t|^\gamma) \leq C \mathbb{E}[(\sup_{0 \leq \tau \leq t} |L_\tau|)^\gamma], \quad 0 < t \leq 1. \quad (5.8)$$

To estimate the right hand side, we notice that for every  $t \in (0, 1]$  we have

$$\mathbb{E}[\sup_{0 \leq \tau \leq t} |L_\tau|^\gamma] = \mathbb{E}[\sup_{0 \leq \tau \leq 1} |L_{t\tau}|^\gamma] = t^{\gamma/(2s)} \mathbb{E}[(\sup_{0 \leq \tau \leq 1} t^{-1/(2s)} |L_{t\tau}|)^\gamma].$$

It is known (e.g. [4, Ch. 8]) that the process  $\{L_r^* := \sup_{0 \leq \tau \leq r} |L_\tau| : r \geq 0\}$  enjoys the scaling property of index  $2s$ , namely for every  $k > 0$  the rescaled process  $\{k^{-1/(2s)} L_{kr}^* : r \geq 0\}$  has the same finite dimensional distributions of  $\{L_r^* : r \geq 0\}$ ; in particular for every  $r > 0$  and  $k > 0$  the random variables  $k^{-1/(2s)} L_{kr}^*$  and  $L_r^*$  have the same law. Taking  $k = t$ ,  $r = 1$ , we obtain

$$\mathbb{E}[(\sup_{0 \leq \tau \leq 1} t^{-1/(2s)} |L_{t\tau}|)^\gamma] = \mathbb{E}[(\sup_{0 \leq \tau \leq 1} |L_\tau|)^\gamma]$$

and the latter is finite, due to [32, Ex. 25.10, Thm. 25.18]. Replacing in (5.8), we get  $\mathbb{E}(|X_t|^\gamma) \leq Ct^{\gamma/(2s)}$ , and (5.5) follows.  $\square$

Applying the results of Sections 3 and 4 we obtain the following proposition. We recall that  $Y_\alpha = \{f \in C_b(\mathbb{R}^N) : \sup_{t>0, x \in \mathbb{R}^N} t^{-\alpha} |f(e^{tA}x) - f(x)| < +\infty\}$ .

**Proposition 5.4.** (i) For every  $s \in (0, 1)$  we have

$$\overline{D(L)} = \{f \in BUC(\mathbb{R}^N) : \lim_{t \rightarrow 0} \|f(e^{tA} \cdot) - f\|_\infty = 0\},$$

so that, given any  $f \in C_b(\mathbb{R}^N)$ ,  $P_t f$  converges uniformly to  $f$  as  $t \rightarrow 0$  if and only if  $f \in BUC(\mathbb{R}^N)$  and  $\lim_{t \rightarrow 0} \|f(e^{tA} \cdot) - f\|_\infty = 0$ .

(ii) For every  $s \in (0, 1)$  and  $\alpha \in (0, 1) \setminus \{1/(2s)\}$  we have

$$(C_b(\mathbb{R}^N), D(L))_{\alpha, \infty} = C_b^{2s\alpha}(\mathbb{R}^N) \cap Y_\alpha$$

so that, given any  $f \in C_b(\mathbb{R}^N)$ , we have  $\sup_{t>0} t^{-\alpha} \|P_t f - f\|_\infty < +\infty$  if and only if  $C_b^{2s\alpha}(\mathbb{R}^N) \cap Y_\alpha$ .

*Proof.* Statement (i) is a consequence of Corollary 3.2 and Proposition 4.1. Statement (ii), with  $\alpha \neq 1/(2s)$ , follows from Propositions 3.3 and 4.4, since the assumptions of Proposition 4.4 are satisfied with  $\theta = 1/(2s)$  by the above discussion and Lemma 5.3.  $\square$

**Remark 5.5.** The results in this section hold also for  $A = 0$ . In this case if we take  $Q = 2^s I$  the operator  $-L$  is just the realization of the fractional Laplacian  $(-\Delta)^s$  in  $C_b(\mathbb{R}^N)$ . However the characterization of the interpolation spaces  $D_L(\alpha, \infty)$  may be obtained for free from the general theory of powers of operators and the characterization for the Laplacian (the case of general  $Q > 0$  may be easily reduced to this one).

Indeed, for every Banach space  $X$  and for every linear operator  $T : D(T) \subset X \mapsto X$  such that  $\sup_{\lambda>0} \lambda \|(\lambda I + T)^{-1}\|_{\mathcal{L}(X)} < +\infty$ , we have the continuous embeddings  $(X, D(T))_{s, 1} \subset D(-T)^s \subset (X, D(T))_{s, \infty}$  for every  $s \in (0, 1)$ . See [34, Sect. 1.14.1], [22, Prop. 4.7]; in both books it is assumed that  $0 \in \rho(T)$ , but an inspection to the proof shows that this assumption is not essential. By the

Reiteration Theorem we get  $(X, D(T)^s)_{\alpha, \infty} = (X, D(T))_{\alpha s, \infty}$  for every  $\alpha \in (0, 1)$ , with equivalence of the respective norms. In our case,  $T = -\Delta$ , we have (e.g., [21, Thm. 3.1.12])

$$(X, D(T))_{\theta, \infty} = \begin{cases} C_b^{2\theta}(\mathbb{R}^N), & \theta \neq 1/2, \\ Z_b^1(\mathbb{R}^N), & \theta = 1/2. \end{cases}$$

As a consequence, for every  $\alpha \in (0, 1)$  we get

$$(X, D((-\Delta)^s)_{\alpha, \infty} = \begin{cases} C_b^{2\alpha s}(\mathbb{R}^N), & \alpha \neq 1/(2s), \\ Z_b^1(\mathbb{R}^N), & \alpha = 1/(2s). \end{cases}$$

## 6. ACKNOWLEDGEMENTS

The Author is a member of GNAMPA-INdAM. Thanks are due to E. Priola for suggestions about §5.2.

## REFERENCES

- [1] D. APPLEBAUM, *Infinite dimensional Ornstein-Uhlenbeck processes driven by Lévy processes*, Prob. Surveys **12** (2015), 33-54.
- [2] S.A. ATHREYA, R.F. BASS, E.A. PERKINS, *Hölder norm estimates for elliptic operators on finite and infinite dimensional spaces*, Trans. Amer. Math. Soc. **357** (2005), 5001-5029.
- [3] P. ALPHONSE, J. BERNIER, *Smoothing properties of fractional Ornstein-Uhlenbeck semigroups and null controllability*, Bull. Sci. Math. **165** (2020), 102914.
- [4] J. BERTOIN, *Lévy Processes*, Cambridge Univ. Press, 1996.
- [5] V.I. BOGACHEV, *Gaussian Measures*, American Mathematical Society, 1998.
- [6] V.I. BOGACHEV, *Differentiable Measures and the Malliavin Calculus*, American Mathematical Society, 2010.
- [7] V.I. BOGACHEV, M. RÖCKNER, B. SCHMULAND, *Generalized Mehler semigroups and applications*, Prob. Th. Rel. Fields **105** (1996), 193-225.
- [8] S. CERRAI, *Second Order PDEs in Finite and Infinite Dimension. A probabilistic approach*, Lecture Notes in Mathematics, 1762. Springer-Verlag, Berlin, 2001.
- [9] A. CHOJNOWSKA-MICHALIK, *On processes of Ornstein-Uhlenbeck type in Hilbert space*, Stochastics **21** (1987), no. 3, 251-286.
- [10] G. DA PRATO, P. CANNARSA, *Infinite dimensional elliptic equations with Hölder continuous coefficients*, Adv. Diff. Equations **1** (1996), 425-452.
- [11] G. DA PRATO, A. LUNARDI, *On the Ornstein-Uhlenbeck operator in spaces of continuous functions*, J. Funct. Anal. **131** (1995), 94-114.
- [12] G. DA PRATO, J. ZABCZYK, *Second Order Partial Differential Equations in Hilbert spaces*, Cambridge Univ. Press, Cambridge, 2002.
- [13] D. A. DAWSON, Z. LI, *Skew convolution semigroups and affine Markov processes*, Ann. Probab. **34** (2006), no. 3, 1103-1142.
- [14] D. A. DAWSON, Z. LI, B. SCHMULAND, W. SUN, *Generalized Mehler semigroups and catalytic branching processes with immigration*, Potential Anal. **21** (2004), no. 1, 75-97.
- [15] M. FUHRMAN, M. RÖCKNER, *Generalized Mehler semigroups: the non-Gaussian case*, Potential Anal. **12** (2000), 1-47.
- [16] B. GOLDYS, J. VAN NEERVEN, *Transition semigroups of Banach space valued Ornstein-Uhlenbeck processes*, Acta Appl. Math. **76** (2003), 283-330. Revised version ArXiv 0606785v1.
- [17] P. LESCOT, M. RÖCKNER, *Generators of Mehler type semigroups as pseudo-differential operators*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **5** (2002), 297-315.
- [18] P. LESCOT, M. RÖCKNER, *Perturbations of generalized Mehler semigroups and applications to stochastic heat equations with Levy noise and singular drift*, Potential Anal. **20** (2004), 317-344.
- [19] Z. LI, *Measure-valued branching Markov processes*, Probability and its Applications (New York). Springer, Heidelberg, 2011.
- [20] J.L. LIONS, J. PEETRE: *Sur une classe d'espaces d'interpolation*, Publ. I.H.E.S. **19** (1964), 5-68.
- [21] A. LUNARDI, *Analytic semigroups and optimal regularity in parabolic equations*, Birkhäuser, Basel, 1995. Second Edition 2013.

- [22] A. LUNARDI, *Interpolation Theory*. Third edition, Lecture Notes. Scuola Normale Superiore di Pisa (New Series), 16. Edizioni della Normale, Pisa, 2018.
- [23] A. LUNARDI, *Schauder estimates for a class of degenerate elliptic and parabolic operators with unbounded coefficients in  $\mathbb{R}^n$* , Ann. Sc. Norm. Sup. Pisa, IV **24** (1997), 133-164.
- [24] A. LUNARDI, D. PALLARA, *Ornstein-Uhlenbeck semigroups in infinite dimension*, Phil. Trans. A **378** (2020), 20190620.
- [25] A. LUNARDI, M. RÖCKNER, *Schauder theorems for a class of (pseudo-)differential operators on finite and infinite dimensional state spaces*, arXiv:1907.06237. To appear in J. London Math. Soc.
- [26] S.-X. OUYANG, M. RÖCKNER, *Time inhomogeneous generalized Mehler semigroups and skew convolution equations*, Forum Math. **28** (2016), no. 2, 339-376.
- [27] S.-X. OUYANG, M. RÖCKNER, F.-Y. WANG, *Harnack inequalities and applications for Ornstein-Uhlenbeck semigroups with jump*, Potential Anal. **36** (2012), no. 2, 301-315.
- [28] E. PRIOLA, *Pathwise uniqueness for singular SDEs driven by stable processes*, Osaka J. Math. **49** (2012), 421-447.
- [29] E. PRIOLA, J. M. A. M. VAN NEERVEN, *Norm discontinuity and spectral properties of Ornstein-Uhlenbeck semigroups*, J. Evol. Equ. **5** (2005), 557-576.
- [30] E. PRIOLA, J. ZABCZYK, *Densities for Ornstein-Uhlenbeck processes with jumps*, Bull. Lond. Math. Soc. **41** (2009), 41-50.
- [31] M. RÖCKNER, F.-Y. WANG, *Harnack and functional inequalities for generalized Mehler semigroups*, J. Funct. Anal. **203** (2003), 237-261.
- [32] K. SATO, *Lévy processes and infinitely divisible distributions*, Cambridge Studies in Advanced Mathematics, 68. Cambridge University Press, Cambridge, 2013.
- [33] B. SCHMULAND, W. SUN, *On the equation  $\mu_{t+s} = \mu_s * T_s \mu_t$* , Statist. Probab. Lett. **52** (2001), 183-188.
- [34] H. TRIEBEL *Interpolation Theory, Function Spaces, Differential Operators*, North-Holland, Amsterdam, 1978.
- [35] J. VAN NEERVEN, L. WEIS, *Stochastic integration of functions with values in a Banach space*, Studia Math. **166** (2005), 132-170.

DIPARTIMENTO DI SCIENZE MATEMATICHE, FISICHE E INFORMATICHE, UNIVERSITÀ DI PARMA, PARCO AREA DELLE SCIENZE, 53/A, 43124 PARMA, ITALY

*Email address:* `alessandra.lunardi@unipr.it`