

FUNCTION SPACES OVER PRODUCTS WITH ORDINALS

RAUSHAN BUZYAKOVA

ABSTRACT. We are concerned with the question of when a homeomorphism between $C_p(X \times \tau)$ and $C_p(Y \times \tau)$ implies the existence of a homeomorphism between $C_p(X \times \tau')$ and $C_p(Y \times \tau')$, where τ and τ' are spaces of ordinals with some additional desired properties.

1. INTRODUCTION

We will be concerned with the following general problem.

Problem. *Let $C_p(X \times Z)$ be homeomorphic to $C_p(Y \times Z)$. Suppose that Z' is a subspace or a superspace of Z . What additional conditions on X, Y, Z, Z' guarantee that $C_p(X \times Z')$ is homeomorphic to $C_p(Y \times Z')$? That X is homeomorphic to Y ?*

Note that for any X and Y we can find Z such that $X \times Z$ is homeomorphic to $Y \times Z$ (in particular, $Z = X^\omega \times Y^\omega$ works). Therefore, it is natural to require that candidates for Z in the above problems have rigid structures. There are many interesting results around this problem. Since most of Z 's in this study will be ordered, we would like to mention O. Okunev's result (see, for example, [1, the argument of Theorem 0.6.2]) that if $C_p(X \times \mathbb{R})$ and $C_p(Y \times \mathbb{R})$ are homeomorphic, then so are $C_p(X)^\omega$ and $C_p(Y)^\omega$. In this work, we will primarily target cases when Z is a space of ordinals. Our major restriction on the topology of a space will be a network size. We will show, in particular, that under certain restrictions on a network size of X and Y , a homeomorphism between $C_p(X \times \tau)$ and $C_p(Y \times \tau)$, where τ is regular and uncountable, implies the existence of a homeomorphism between $C_p(X \times \lambda)$ and $C_p(Y \times \lambda)$ for some $\lambda < \tau$. This statement and other observations lead to some natural generalizations as well as questions for possible further exploration.

Recall that a network of X is a family of sets with the same properties as a basis except that elements of the family need not be open. By $A(\tau)$ we denote the Alexandroff single-point compactification of a discrete space of size τ with ∞

2020 *Mathematics Subject Classification.* 54C35, 54B10, 54E99.

Key words and phrases. function spaces in the topology of point-wise convergence, space of ordinals, product spaces, homeomorphism.

being its only non-isolated point. All spaces under consideration are Tychonov. In notation and terminology of general topological nature we follow [3]. To distinguish ordered pairs from open intervals, we will denote the former by $\langle a, b \rangle$ and the latter by (a, b) . We also refer the reader to [1] for general facts and notations involving function spaces endowed with the topology of point-wise convergence.

2. STUDY

In our arguments we will use the following folklore fact for which we outline a proof for convenience.

Proposition 2.1. *(Folklore) Let an ordinal τ have cofinality greater than the density of an infinite space X and let $f \in C_p(X \times \tau) \cup C_p(X \times (\tau + 1))$. Then there exists $\lambda < \tau$ such that $f|_{\{x\} \times [\lambda, \tau]}$ is constant for every $x \in X$.*

Proof. To see why the statement is true let $D = \{d_\alpha : \alpha < d(X)\}$ be dense in X , where $d(X)$ is the density of X . For each $\alpha < d(X)$, fix $\tau_\alpha < \tau$ such that f is constant on $\{d_\alpha\} \times [\tau_\alpha, \tau)$. Such a τ_α exists due to countable compactness of τ . Since $d(X) < cf(\tau)$, we conclude that $\lambda = \sup\{\tau_\alpha : \alpha < d(X)\} < \tau$. By continuity of f and density of D , λ is as desired. \square

Theorem 2.2. *Let X and Y have networks of cardinality less than τ , where τ is regular and uncountable. Let $C_p(X \times \tau)$ be homeomorphic to $C_p(Y \times \tau)$. Then $C_p(X \times \lambda)$ is homeomorphic to $C_p(Y \times \lambda)$ for some non-zero $\lambda < \tau$.*

Proof. Fix a homeomorphism $H : C_p(X \times \tau) \rightarrow C_p(Y \times \tau)$. Next for $\alpha < \tau$, define $F(X, \alpha)$ as follows: $f \in F(X, \alpha)$ if and only if f is constant on $\{x\} \times [\alpha, \tau)$ for every $x \in X$. Similarly, we define $F(Y, \alpha)$.

Claim 1. $F(X, \mu)$ is homeomorphic to $C_p(X \times (\mu + 1))$.

To prove the claim, for each $f \in C_p(X \times (\mu + 1))$, let $f_\mu \in F(X, \mu)$ be the function such that $f(x, \alpha) = f_\mu(x, \alpha)$ for each $\langle x, \alpha \rangle \in X \times (\mu + 1)$. Clearly, the map $M : C_p(X \times (\mu + 1)) \rightarrow F(X, \mu)$ defined by $M(f) = f_\mu$ is a homeomorphism.

Claim 2. $F(X, \alpha)$ is closed in $C_p(X \times \tau)$ for any $\alpha < \tau$

To prove the claim fix $f \in C_p(X \times \tau) \setminus F(X, \alpha)$. By the description of $F(X, \alpha)$, there exist $x \in X$ and $\lambda \in (\alpha, \tau)$ such that $f(x, \lambda) \neq f(x, \alpha)$. Put $\epsilon = |f(x, \lambda) - f(x, \alpha)|$. Then the following open set contains f and misses $F(X, \alpha)$:

$$\begin{aligned} & \{g : g(x, \lambda) \in (f(x, \lambda) - \epsilon/3, f(x, \lambda) + \epsilon/3)\} \cap \\ & \cap \{g : g(x, \alpha) \in (f(x, \alpha) - \epsilon/3, f(x, \alpha) + \epsilon/3)\} \end{aligned}$$

The claim is proved.

Claim 3. Let $\alpha < \tau$. Then $H(F(X, \alpha)) \subset F(Y, \alpha')$ for some $\alpha' < \tau$.

To prove the claim, recall that X has netweight less than τ and τ is regular. Therefore, $X \times (\alpha + 1)$ has netweight less than τ . Therefore, $C_p(X \times (\alpha + 1))$ has density less than τ . By Claim 1, $F(X, \alpha)$ has density less than τ too. By Proposition 2.1, any subset of $C_p(Y \times \tau)$ of density less than τ is a subset of $F(Y, \alpha')$ for some $\alpha' < \tau$.

By Claim 3, we can create a sequence $\alpha_1 < \alpha_2 < \dots$ with the following two properties:

P1: $H(F(X, \alpha_n)) \subset F(Y, \alpha_{n+1})$ if n is odd.

P2: $H^{-1}(F(Y, \alpha_n)) \subset F(X, \alpha_{n+1})$ if n is even.

Put $\mu = \sup\{\alpha_n\}_n$. By Claim 1, it remains to show that $H(F(X, \mu)) = F(Y, \mu)$. Since $\{\alpha_n\}_n$ is an increasing sequence we conclude that $H(\cup_n F(\alpha_n, X)) = \cup_n F(\alpha_n, Y)$. Since H is a homeomorphism, $H(\overline{\cup_n F(\alpha_n, X)}) = \overline{\cup_n F(\alpha_n, Y)}$. It remains to show that $\overline{\cup_n F(\alpha_n, X)} = F(\mu, X)$ and $\overline{\cup_n F(\alpha_n, Y)} = F(\mu, Y)$. We will prove the former equality. Since $F(\alpha_n, X)$ is a subset of $F(\mu, X)$ and the latter is closed by Claim 2, we conclude that $\overline{\cup_n F(\alpha_n, X)} \subset F(\mu, X)$. To show the reverse inclusion, fix $f \in F(\mu, X)$ and a standard open neighborhood U of f . We may assume that $U = U_1 \cap U_2$, where $U_1 = U_1(p_1, \dots, p_n, B_1, \dots, B_n)$ and $U_2 = U_2(q_1, \dots, q_k, O_1, \dots, O_k)$ for some $p_1, \dots, p_n \in X \times \mu$ and $q_1, \dots, q_k \in (X \times \tau) \setminus (X \times \mu)$. To show that U meets $\overline{\cup_n F(\alpha_n, X)}$ let N be such that $p_1, \dots, p_n \in X \times \alpha_N$. Define g by letting $g(x, \beta) = f(x, \mu)$ for every $\beta > \alpha_N$ and $g(x, \beta) = f(x, \beta)$ for every $\beta \leq \alpha_N$. Clearly, $g \in U \cap F(\alpha_n, X)$. \square

The following is a by-product of the argument of Theorem 2.2.

Theorem 2.3. Let X and Y have networks of cardinality less than τ , where τ is regular and uncountable. Let $C_p(X \times (\tau+1))$ be homeomorphic to $C_p(Y \times (\tau+1))$. Then $C_p(X \times \lambda)$ is homeomorphic to $C_p(Y \times \lambda)$ for some non-zero $\lambda < \tau$.

One may wonder if a homeomorphism between $C_p(X \times \kappa)$ and $C_p(Y \times \kappa)$ for some ordinal κ implies the existence of a homeomorphism between $C_p(X \times \lambda)$ and $C_p(Y \times \lambda)$ for some $0 < \lambda < \kappa$. It is, however, not the case. Note that the function spaces over any discrete space X is homeomorphic to $\mathbb{R}^{|X|}$. Therefore, $C_p(2 \times \omega)$ and $C_p(3 \times \omega)$ are homeomorphic but $C_p(2 \times n)$ and $C_p(3 \times n)$ are not homeomorphic for any positive integer n .

For future reference let us isolate the case of $\tau = \omega_1$ of Theorem 2.2 into a corollary.

Corollary 2.4. *Let X and Y have countable networks. If $C_p(X \times \omega_1)$ is homeomorphic to $C_p(Y \times \omega_1)$, then $C_p(X \times \lambda)$ is homeomorphic to $C_p(Y \times \lambda)$ for some non-zero countable ordinal λ .*

The above corollary prompts the following question.

Question 2.5. *Let $C_p(X \times (\omega + 1))$ and $C_p(Y \times (\omega + 1))$ be homeomorphic. Is it true that $C_p(X \times \mathbb{Q})$ and $C_p(Y \times \mathbb{Q})$ are homeomorphic? What additional conditions on X and Y guarantee an affirmative answer?*

An argument similar to that in Theorem 2.2 leads to our next statement. Since the argument of Theorem 2.2 needs only notation changes for our next result, we will provide the strategy without repeating the proof details of Theorem 2.2.

Theorem 2.6. *Let X and Y have countable networks and let $C_p(X \times A(\tau))$ be homeomorphic to $C_p(Y \times A(\tau))$ for some infinite cardinal τ . Then $C_p(X \times A(\omega))$ is homeomorphic to $C_p(Y \times A(\omega))$.*

Proof. We may assume that τ is uncountable. Fix a homeomorphism $\phi : C_p(X \times A(\tau)) \rightarrow C_p(Y \times A(\tau))$. Given $A \subset \tau$, define $F(A, X) \subset C_p(X \times A(\tau))$ as follows:

Definition of $F(A, X)$: $f \in C_p(X \times A(\tau))$ is in $F(A, X)$ if and only if $f(x, \beta) = f(x, \infty)$ for every $\beta \notin A$ and every $x \in X$.

Claim 1. *For every $f \in C_p(X \times A(\tau))$ there exists a countable $A \subset \tau$ such that $f \in F(A, X)$.*

To prove the claim, assume the contrary. Then there exists a subset $\{\langle x_\alpha, \lambda_\alpha \rangle : \alpha < \omega_1\} \subset X \times A(\tau)$ such that $f(x_\alpha, \lambda_\alpha) \neq f(x_\alpha, \infty)$ for each $\alpha < \omega_1$, and $\lambda_\alpha \neq \lambda_\beta$ for distinct $\alpha, \beta < \omega_1$. Without loss of generality we may assume that there exist $p, q \in \mathbb{Q}$ such that $f(x_\alpha, \lambda_\alpha) < p < q < f(x_\alpha, \infty)$ for all $\alpha < \omega_1$. Since X is Lindelöf, there exists $\langle x, \infty \rangle$ a complete accumulation point of $\{\langle x_\alpha, \infty \rangle : \alpha < \omega_1\}$, and therefore, $f(x, \infty) \geq q$. By the product topology, $\langle x, \infty \rangle$ is also a complete accumulation point for $\{\langle x_\alpha, \lambda \rangle : \alpha < \omega_1\}$, and therefore, $f(x, \infty) \leq p$. We have arrived at a contradiction with $p < q$, which proves the claim.

The next two claims (as Claim 1) are proved using the same arguments as the corresponding claims of Theorem 2.2.

Claim 2. *Let K be an infinite subset of τ of cardinality κ . Then $F(X, K)$ is homeomorphic to $C_p(X \times A(\kappa))$.*

Claim 3. *$F(X, A)$ is closed in $C_p(X \times A(\tau))$ for any $A \subset \tau$.*

By Claim 2, it remains to find a countable $A \subset \tau$ such that $\phi(F(X, A)) = F(Y, A)$. Inductively, we will define A_n that will be building blocks for our desired A .

Step 0: $A_0 = \omega$.

Assumption: A_k is defined for all $k = 0, \dots, n - 1$ and $A_m \subset A_k$ if $m < k$.

Step n: Our definition of A_n will depend on whether n is odd or even.

Assume that n is odd. Since X has a countable network, by Claim 2, $F(A_{n-1}, X)$ is separable. By Claims 1 and 3, there exists a countable $A_n \subset \tau$ that contains A_{n-1} such that $\phi(F(A_{n-1}, X)) \subset F(A_n, Y)$.

Assume that n is even. Similarly, we choose a countable $A_n \subset \tau$ such that A_n contains A_{n-1} and $\phi^{-1}(F(A_{n-1}, Y)) \subset F(A_n, X)$.

Our induction construction is complete. Put $A = \cup_n A_n$. By Claim 2, it remains to show that $\phi(F(A, X)) = F(A, Y)$. The argument is identical to that in Theorem 2.2. \square

A more general statement is as follows.

Theorem 2.7. *Let τ be an infinite cardinal. Let X and Y have networks of cardinality $\tau < \kappa$. If $C_p(X \times A(\kappa))$ is homeomorphic to $C_p(Y \times A(\kappa))$, then $C_p(X \times A(\tau))$ is homeomorphic to $C_p(Y \times A(\tau))$.*

It would be interesting to consider scenarios in a direction somewhat opposite to Theorems 2.2 - 2.7, namely, in a direction aligned with the following question.

Question 2.8. *Assume that $C_p(X \times \alpha)$ is homeomorphic to $C_p(Y \times \alpha)$ for every $\alpha < \omega_1$. Is it true that $C_p(X \times \omega_1)$ is homeomorphic to $C_p(Y \times \omega_1)$? What if ω_1 is replaced by an arbitrary fixed ordinal?*

Note that there are many examples of pairs X and Y that satisfy the assumption of Question 2.8. For example, it was shown by van Mill [6] that any two non-locally compact subspaces of rational numbers have homeomorphic C_p 's. This result was later generalized to any infinite non-discrete subspaces of rationals by Dobrowolski, Marciszewski, and Mogilski [2]. In connection with these nice facts and the statement of Theorem 2.4, the following special case of Question 2.8 might be of interest:

Question 2.9. *Let X and Y be infinite non-discrete subspaces of rationals. Is it true that $C_p(X \times \omega_1)$ is homeomorphic to $C_p(Y \times \omega_1)$? What if ω_1 is replaced by $\omega_1 + 1$ or any other larger ordinal?*

Next, we would like to remark on a positive outcome in the scenario "from smaller to larger" ordinal. A particular case of Okunev's main result in [5] states that if $C_p(X)$ and $C_p(Y)$ are homeomorphic for pseudocompact X and Y , then $C_p(\beta X)$ and $C_p(\beta Y)$ are homeomorphic too. It is also a classical fact that if X is compact, then $X \times \omega_1$ is pseudocompact and $\beta(X \times \omega_1) = X \times (\omega_1 + 1)$. These facts imply that if X and Y are compact and $C_p(X \times \omega_1)$ is homeomorphic to $C_p(Y \times \omega_1)$, then $C_p(X \times (\omega_1 + 1))$ is homeomorphic to $C_p(Y \times (\omega_1 + 1))$. Clearly, the statement holds if we replace ω_1 with any ordinal of uncountable cofinality. Let us summarize this discussion as follows.

Remark 2.10. *(a Corollary to Okunev's Theorem in [5]) Let X and Y be compact spaces and let an ordinal τ have uncountable cofinality. If $C_p(X \times \tau)$ is homeomorphic to $C_p(Y \times \tau)$, then $C_p(X \times (\tau + 1))$ is homeomorphic to $C_p(Y \times (\tau + 1))$.*

A few natural questions arise from the above discussion.

Question 2.11. *Let X and Y be compact spaces. Suppose that $C_p(X \times (\omega_1 + 1))$ is homeomorphic to $C_p(Y \times (\omega_1 + 1))$. Is it true that $C_p(X \times \omega_1)$ is homeomorphic to $C_p(Y \times \omega_1)$?*

Question 2.12. *Are there spaces X and Y (not necessarily compact) such that $C_p(X \times (\omega_1 + 1))$ is homeomorphic to $C_p(Y \times (\omega_1 + 1))$ but $C_p(X \times \omega_1)$ is not homeomorphic to $C_p(Y \times \omega_1)$? Or, such that $C_p(X \times \omega_1)$ is homeomorphic to $C_p(Y \times \omega_1)$ but $C_p(X \times (\omega_1 + 1))$ is not homeomorphic to $C_p(Y \times (\omega_1 + 1))$?*

We would like to finish with two general questions related to our study that are also particular cases of the general problem that motivated this work.

Question 2.13. *Assume that $C_p(X \times \tau)$ and $C_p(Y \times \tau)$ are homeomorphic for some ordinal $\tau > 1$. What additional conditions on X, Y, τ guarantee that $C_p(X)$ and $C_p(Y)$ homeomorphic? That X and Y are homeomorphic?*

Question 2.14. *Is there an example of X and Y with homeomorphic C_p 's such that $C_p(X \times \tau)$ and $C_p(Y \times \tau)$ are not homeomorphic for some τ ?*

In connection with this question, a notable pair to test is the set of reals and a closed segment. It is due to Gulko and Khmyleva [4] that $C_p(\mathbb{R})$ and $C_p([0, 1])$ are homeomorphic. It might be interesting to know if there is an ordinal λ such that $C_p(\mathbb{R} \times \lambda)$ and $C_p([0, 1] \times \lambda)$ are not homeomorphic. For an overview of non-homeomorphic spaces with homeomorphic C_p 's, which may address Question 2.14, we refer the reader to [7].

REFERENCES

- [1] A. Arhangel'skii, *Topological Function Spaces*, Math. Appl., vol. 78, Kluwer Academic Publishers, Dordrecht, 1992
- [2] T. Dobrowolski, W. Marcsiszewski, and J. Mogilski, *On topological classification of functions spaces $C_p(X)$ of low Borel complexity*, Trans. Amer. Math. Soc. 678(1991), 307-324
- [3] R. Engelking, *General Topology*, PWN, Warszawa, 1977
- [4] S. P. Gulko and T. E. Khmyleva, *Compactness is not preserved by t -equivalence*, Mat. Zametki, 39 (1986), 895-903, in Russian
- [5] O. G. Okunev, *Spaces of functions in the topology of pointwise convergence: the Hewitt extension and tau-continuous functions*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. 1985, no. 4, 78-80, in Russian
- [6] J. van Mill, *Topological Equivalence of Certain Function Spaces*, Composito. Math., 63(1987), 195-201
- [7] J. van Mill, *The Infinite- Dimensional Topology of Function Spaces*, Elsevier, 2001
Email address: Raushan.Buzyakova@yahoo.com