

First-Order Quantum Correction in Coherent State Expectation Value of Loop-Quantum-Gravity Hamiltonian: Overview and Results

Cong Zhang¹ Shicong Song² Muxin Han^{2,3}

¹*Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland*

²*Department of Physics, Florida Atlantic University, 777 Glades Road, Boca Raton, FL 33431-0991, USA*

³*Institut für Quantengravitation, Universität Erlangen-Nürnberg, Staudtstr. 7/B2, 91058 Erlangen, Germany*

E-mail: [czhang\(AT\)fuw.edu.pl](mailto:czhang@fuw.edu.pl), [ssong2019\(AT\)fau.edu](mailto:ssong2019@fau.edu), [hanm\(AT\)fau.edu](mailto:hanm@fau.edu)

ABSTRACT: Given the Loop-Quantum-Gravity (LQG) non-graph-changing Hamiltonian $\widehat{H}[N]$, the coherent state expectation value $\langle \widehat{H}[N] \rangle$ admits a semiclassical expansion in ℓ_p^2 . In this paper, we compute explicitly the expansion of $\langle \widehat{H}[N] \rangle$ on the cubic graph to the linear order in ℓ_p^2 , when the coherent state is peaked at the homogeneous and isotropic data of cosmology. In our computation, a powerful algorithm is developed to overcome the complexity in computing $\langle \widehat{H}[N] \rangle$. In particular, some key innovations in our algorithm substantially reduce the computational complexity in the Lorentzian part of $\langle \widehat{H}[N] \rangle$. Moreover, the algorithm developed in the present work makes it possible to compute the expectation value of arbitrary monomial of holonomies and fluxes on one edge up to arbitrary order of ℓ_p^2 .

Contents

1	Introduction	1
2	Preliminaries	4
2.1	Quantization and Hamiltonian	4
2.2	Coherent states	6
3	Expectation values of operators on one edge	7
3.1	the first step	8
3.2	the second step	8
3.3	the third step	9
4	Power counting	9
5	Cosmological expectation value	11
5.1	Symmetries of the expectation value	12
5.1.1	Symmetries of the Euclidean part	12
5.1.2	Symmetries of the Lorentzian part	13
5.2	Introducing the algorithm: A case study	17
6	Quantum correction in the expectation value	19
7	Conclusion and outlook	21

1 Introduction

Loop Quantum Gravity (LQG) is an approach toward the background independent and nonperturbative quantum gravity in four and higher dimensions [1–4]. The research on the quantum dynamics of LQG is active and leads to important recent progresses. In particular, there have been tremendous progresses in both canonical and covariant LQG on the semiclassical limit and the consistency with classical gravity e.g. [5–12]. However, there has been less progress on quantum corrections in the full theory of LQG dynamics (see e.g. [13, 14] for some results in the covariant approach). It is important that as a candidate of quantum gravity theory, LQG should provide predictions on quantum corrections to the classical theory of gravity.

The present paper focuses on the canonical LQG. In canonical LQG, the dynamics is encoded in the Hamiltonian constraint operator $\widehat{H}[N] = \widehat{H}_E[N] + (1 + \beta^2)\widehat{H}_L[N]$, where N is the lapse function, $\widehat{H}_E[N]$ and $\widehat{H}_L[N]$ are the Euclidean part and Lorentzian part of $\widehat{H}[N]$, and β is referred to as the Barbero-Immirzi parameter [15]. There has been long-term confusion that the quantum dynamics of LQG might not be computable analytically due to the non-polynomial Hamiltonian constraint operator $\widehat{H}[N]$ [16]. This confusion has been partially resolved by [6], where the authors schematically show that the coherent state expectation value of the Hamiltonian/master constraint

are computable order-by-order by the semiclassical expansion in \hbar . The scheme proposed in [6] is applicable to a wide class of non-polynomial operators used in studying LQG dynamics. Although this scheme was proposed as early as in 2006 when [6] firstly appeared, the expectation value $\langle \widehat{H[N]} \rangle$ has only been computed at the 0-th order (in \hbar) which is the classical limit, while the $O(\hbar)$ quantum correction was not studied in the literature, due to the complexity of the operator, especially the complexity of the Lorentzian part $\widehat{H_L[N]}$ in $\widehat{H[N]}$.

The main purpose of our work is to fill this gap by computing explicitly the $O(\hbar)$ quantum correction in the coherent state expectation value $\langle \widehat{H[N]} \rangle$. In this paper, a powerful algorithm is developed to overcome the complexity of $\widehat{H[N]}$, so that we can compute the quantum correction in $\langle \widehat{H[N]} \rangle$, where $\widehat{H[N]}$ is the non-graph-changing Hamiltonian on a cubic lattice γ . Applying the algorithm, we explicitly expand $\langle \widehat{H[N]} \rangle$ to linear order in \hbar , when the coherent state is peaked at the homogeneous and isotropic data of cosmology. Namely we explicitly compute H_0 and H_1 in

$$\langle \widehat{H[1]} \rangle = H_0 + \ell_p^2 H_1 + O(\ell_p^4), \quad \ell_p^2 = \hbar \kappa \quad (1.1)$$

where $\kappa = 8\pi G_{\text{Newton}}$ and the lapse function $N = 1$. The 0-th order H_0 reproduces the μ_0 -scheme cosmological effective Hamiltonian [17–20]. Our new result, H_1 , gives the first order quantum correction. The explicit expression of H_1 is given in Section 6. Note that in our work, the coherent state for $\langle \widehat{H[N]} \rangle$ is not $\text{SU}(2)$ gauge invariant (see below for the motivation).

This work closely relates to the reduced phase space formulation of LQG (see e.g. [21, 22]). In this formulation, gravity is coupled to some matter fields known as clock fields. These matter fields serve as material reference frames used to transform gravity variables to gauge invariant Dirac observables. This procedure resolves the Diffeomorphism and Hamiltonian constraints at the classical level and results in the reduced phase space \mathcal{P}_{red} of Dirac observables. The dynamics of the gravity-clock system is described by the material-time evolution generated by the physical Hamiltonian \mathbf{H} on \mathcal{P}_{red} . As an interesting model, we choose the clock fields to be the Gaussian dust [22, 23]. Then the resulting \mathcal{P}_{red} is identified to the pure-gravity unconstrained phase space. This identification defines the pure-gravity Hamiltonian constraint with unit lapse $H[1]$ on \mathcal{P}_{red} , and we find $\mathbf{H} = H[1]$ for gravity coupled to Gaussian dust. In this model, the LQG quantization of \mathcal{P}_{red} is the same as quantizing the pure-gravity unconstrained phase space, and leads to the physical Hilbert space \mathcal{H} identical to the kinematical Hilbert space in the usual LQG. \mathcal{H} is free of constraint because it is from the quantization of \mathcal{P}_{red} . The physical Hamiltonian operator is obtained by $\widehat{\mathbf{H}} = \frac{1}{2}(\widehat{H[1]} + \widehat{H[1]}^\dagger)$ with the LQG quantization $\widehat{H[N]}$ [24, 25]. Therefore from the reduced phase space LQG perspective, our work computes the expectation value $\langle \widehat{\mathbf{H}} \rangle$ at the coherent state peaked at cosmological data on the graph γ , and gives $\langle \widehat{\mathbf{H}} \rangle$ equals the real part of (1.1).

Recent research works have been focus on building models of LQG on a single graph γ [18, 26–32]. In particular, the quantum dynamics of the reduced phase space LQG is formulated on the cubic lattice γ as a path integral [7, 19]

$$A_{[g],[g']} = \int dh[dg] \nu[g] e^{S[g,h]/\ell_p^2}, \quad (1.2)$$

which is the canonical-LQG analog of the spinfoam formulation. $A_{[g],[g']}$ is the transition amplitude of $\widehat{\mathbf{H}}$ between the initial and final $\text{SU}(2)$ gauge invariant coherent states $|[g]\rangle, |[g']\rangle$. The integration variables contains trajectories of $g \in \mathcal{P}_{red,\gamma}$ and $\text{SU}(2)$ gauge transformations h on γ . $\nu[g]$ is a measure factor. As a feature of the path integral formula (1.2), the $\text{SU}(2)$ gauge invariant amplitude $A_{[g],[g']}$ is expressed as the integral in terms of $\text{SU}(2)$ gauge non-invariant variables g and h . The

action $S[g, h]$ is linear to the expectation value $\langle \widehat{\mathbf{H}} \rangle$ at $SU(2)$ gauge non-invariant coherent states when trajectories of g are continuous in time. In contrast to usual path integrals of quantum field theories, $S[g, h]$ contains not only the 1-loop quantum correction but also the $O(\hbar)$ correction from $\langle \widehat{\mathbf{H}} \rangle$. Our work precisely computes this $O(\hbar)$ correction from $\langle \widehat{\mathbf{H}} \rangle$ in $S[g, h]$ in the case of cosmological dynamics.

The semiclassical limit $\hbar \rightarrow 0$ of $A_{[g],[g']}$ gives the equation of motion studied in general in [7] and applied in cosmology in [19, 31]. The cosmological dynamics as $\hbar \rightarrow 0$ gives the μ_0 -scheme effective cosmological dynamics which reduces to the classical FRLW cosmology at low energy density. As the next step, it is important to discover the $O(\hbar)$ correction to the effective cosmological dynamics. The effective dynamics with $O(\hbar)$ correction can be obtained by the *quantum effective action* Γ [33] from the path integral (1.2). Perturbatively, the $O(\hbar)$ in Γ for cosmology contains 3 contributions: (1) $O(\hbar)$ in $S[g, h]$ which is computed in this work, (2) $\log \nu[g]$ where $\nu[g]$ has been given explicitly in [19], and (3) $\frac{1}{2} \log \det(\mathfrak{H})$ where the “1-loop determinant” $\det(\mathfrak{H})$ is the determinant of the Hessian matrix \mathfrak{H} of $S[g, h]$. The g - g matrix elements in \mathfrak{H} is computed in [31]. The study of $\frac{1}{2} \log \det(\mathfrak{H})$ is postponed to the future work. Therefore from the perspective of the quantum correction in the effective cosmological dynamics, our work computes an important part in the $O(\hbar)$ of the quantum effective action Γ .

After introducing several motivations in the above, let us summarize the key steps in our computation: First of all, an important complication in $\widehat{H[N]}$ is the volume operator $\widehat{V}_v = \sqrt{|\widehat{Q}_v|}$ which contains the square-root and absolute-value so that $\widehat{H[N]}$ is non-polynomial. In the case of the coherent state expectation value $\langle \widehat{H[N]} \rangle$, this issue is overcome by substituting \widehat{V}_v with the semiclassical expansion [6]

$$\widehat{V}_{GT}^{(v)} = \langle \widehat{Q}_v \rangle^{2q} \left[1 + \sum_{n=1}^{2k+1} (-1)^{n+1} \frac{q(1-q) \cdots (n-1-q)}{n!} \left(\frac{\widehat{Q}_v^2}{\langle \widehat{Q}_v \rangle^2} - 1 \right)^n \right] + O(\hbar^{k+1}) \quad (1.3)$$

where \widehat{Q}_v is a polynomial of flux operators and $q = 1/4$. Inserting in $\widehat{V}_{GT}^{(v)}$ truncated at a finite k reduces $\langle \widehat{H[N]} \rangle$ to the expectation value of a polynomial operator. Note that the current work uses the volume operator introduced in [34], which is the same as that studied in [6]. However, this volume operator is not the unique choice to define $\widehat{H[N]}$. For instance, one can use the volume operators given in [35, 36]. Even though it is still open that whether these operators can be replaced by some operator polynomials as far as the expectation value is concerned, it is expected that these operators should give the same leading order result in \hbar if they possess desirable semiclassical properties.

The resulting polynomial sums a huge number of terms ($\sim 10^{19}$), each of which is a monomial of holonomies and fluxes. Computing expectation values of all individual terms would give a large computational complexity. The complexity mainly comes from the Lorentzian part of $\widehat{H[N]}$. Several key methods in our work are used to reduce the number of terms needed for computation:

- The expectation value of every monomial term in $\widehat{H[N]}$ can be factorized into expectation values of holonomy-flux monomials on individual edges. We only need to compute expectation values of various types of monomials on a single edge. We further reduce the number of types by the commutation relations, and derive several general formulae for the expectation values of resulting types (see Section 3).
- We develop a power-counting argument to find the power of \hbar as the leading order behavior of each expectation value of the monomial operator (see Section 4). Since we focus on expanding

$\langle \widehat{H}[N] \rangle$ to the linear order in \hbar , a large number of monomials only contributing to higher order can be neglected.

- When we focus on the expectation values at coherent states peaked at homogeneous and isotropic data. There are a large amount of symmetries which identify different terms and significantly reduce the number of terms needed for computation (see Section 5).

Our method exponentially reduces the computational complexity and is particularly useful in computing the expectation value of Lorentzian part $\widehat{H}_L[N]$ in $\widehat{H}[N]$. In Section 5.2, we demonstrate the reduction in a typical example which is an operator contains 3^{3m-1} (m can be large) monomials. By applying our method, only 5 monomials are finally needed for computing the expectation values.

The purpose of this paper is to give an overview our computation and present the results. The detailed derivations are presented in [37]. Our computation is carried out in Mathematica on the High Performance Computation server with two 48-Core Processors (AMD EPYC 7642). The Mathematica codes are available at [38].

The resulting $O(\hbar)$ quantum correction in $\langle \widehat{H}[N] \rangle$ is summarized in Section 6. In order to demonstrate the physical application of our results and effects from the $O(\hbar)$ correction, we adopt the proposal in [18]: We view $\Re\langle \widehat{H}[1] \rangle$ in (1.1) as the effective Hamiltonian function on the 2-dimensional phase space \mathcal{P}_{cos} of homogeneous and isotropic cosmology. $\Re\langle \widehat{H}[1] \rangle$ generates the Hamiltonian time evolution on \mathcal{P}_{cos} . We plot the time evolution of the homogeneous spatial volume, and compare with the evolution generated by $\langle \widehat{H}[1] \rangle$ at $\hbar \rightarrow 0$. The comparison demonstrates the effect from the $O(\hbar)$ correction in $\langle \widehat{H}[1] \rangle$ (see Section 6 for details). We emphasize that here the proposal that we adopt for the cosmological evolution is not as rigorous as the path integral formula (1.2), and we have argued above that the $O(\hbar)$ correction in $\langle \widehat{H}[1] \rangle$ is only a partial contribution to the quantum effective action Γ which ultimately determines the quantum effect in the dynamics. The cosmological dynamics studied in Section 6 is only for the purpose of displaying the effect of the $O(\hbar)$ correction in $\langle \widehat{H}[1] \rangle$, and is not yet a rigorous prediction from the principle of LQG.

The paper is organized as follows. Section 2 reviews briefly the theory of LQG on a cubic lattice, including the Hamiltonian and the coherent state. In Section 3, we describe the procedure to compute the expectation value of operators at a single edge. In Section 4, we develop a power-counting argument to reduce the computational complexity. Section 5 discusses $\langle \widehat{H}[1] \rangle$ at coherent states peaked at homogeneous and isotropic data, and discuss the symmetries which further reduce the computational complexity. Section 6 summarizes the results of the quantum correction in $\langle \widehat{H}[1] \rangle$. In Sec. 7, we conclude and discuss a few future perspectives.

Note added in proof: During finalizing this paper, we became aware that the $O(\hbar)$ correction of the Euclidean part \widehat{H}_E in $\langle \widehat{H}[1] \rangle$ was derived independently in [32]¹.

2 Preliminaries

2.1 Quantization and Hamiltonian

Classically general relativity can be reformulated with the Ashtekar-Barbero variables (A_a^i, E_i^a) consisting of an $SU(2)$ connection A_a^i and its canonically conjugate densitized triad field E_i^a defined on the spatial manifold Σ [15]. We choose the coordinate denoted by (x, y, z) on Σ . Let $\gamma \subset \Sigma$ be

¹The result of $\langle \widehat{H}_E \rangle$ in [32] is the same as ours (see (6.1)) up to an overall constant and $(\eta, \xi) \rightarrow (-\eta, -\xi)$.

a finite cubic lattice whose edges are parallel to the axes of the coordinates. The sets of edges and vertices in γ are denoted by $E(\gamma)$ and $V(\gamma)$. Taking advantage of γ , we define holonomies along the edges of γ ,

$$h_e(A) = \mathcal{P} \exp \int_e A = 1 + \sum_{n=1}^{\infty} \int_0^1 dt_n \int_0^{t_n} dt_{n-1} \cdots \int_0^{t_2} dt_1 A(t_1) \cdots A(t_n), \quad \forall e \in E(\gamma), \quad (2.1)$$

and gauge covariant fluxes on the 2-faces S_e in the dual lattices γ^* ,

$$p_{\mathfrak{s}}^i(e) := -\frac{2}{\beta a^2} \text{tr} \left[\tau^i \int_{S_e} dx^a dx^b \varepsilon_{abc} h(\rho_e^{\mathfrak{s}}(\sigma)) E^c(\sigma) h(\rho_e^{\mathfrak{s}}(\sigma)^{-1}) \right], \quad (2.2)$$

where $S_e \in \gamma^*$ is the 2-face, $\rho^{\mathfrak{s}}(\sigma) : [0, 1] \rightarrow \Sigma$ is a path connecting the source point $\mathfrak{s}_e \in e$ to $\sigma \in S_e$ such that $\rho_e^{\mathfrak{s}}(\sigma) : [0, 1/2] \rightarrow e$ and $\rho_e^{\mathfrak{s}}(\sigma) : [1/2, 1] \rightarrow S_e$. The parameter a , identified with a length scale of the system, is chosen to be (cosmological constant) $^{-1/2}$ in our paper. Alternatively, one can choose the target point $\mathfrak{t}_e \in e$ rather than \mathfrak{s}_e to define

$$p_{\mathfrak{t}}^i(e) := \frac{2}{\beta a^2} \text{tr} \left[\tau^i \int_{S_e} dx^a dx^b \varepsilon_{abc} h(\rho_e^{\mathfrak{t}}(\sigma)) E^c(\sigma) h(\rho_e^{\mathfrak{t}}(\sigma)^{-1}) \right]. \quad (2.3)$$

where $\rho^{\mathfrak{t}}(\sigma) : [0, 1] \rightarrow \Sigma$ is a path connecting the target $\mathfrak{t}_e \in e$ to $\sigma \in S_e$ such that $\rho_e^{\mathfrak{t}}(\sigma) : [0, 1/2] \rightarrow e$ and $\rho_e^{\mathfrak{t}}(\sigma) : [1/2, 1] \rightarrow S_e$.

The quantization of this classical lattice theory gives us LQG based on the graph γ . The Hilbert space \mathcal{H}_γ consists of the square integrable functions of the holonomies. Given two functions $\psi_i : \{h_e\}_{e \in E(\gamma)} \rightarrow \mathbb{C}$, the inner produce is

$$\langle \psi_1 | \psi_2 \rangle = \int_{\text{SU}(2)^{|E(\gamma)|}} d\mu_{\text{H}} \overline{\psi_1(\{h_e\}_{e \in E(\gamma)})} \psi_2(\{h_e\}_{e \in E(\gamma)}) \quad (2.4)$$

where $|E(\gamma)|$ denote the number of elements (i.e. cardinality) of $E(\gamma)$ and μ_{H} is the Haar measure. \mathcal{H}_γ is the kinematical Hilbert space in the operator-constraint formalism of the canonical LQG. However \mathcal{H}_γ modulo gauge transformations is the physical Hilbert space in the reduced phase space LQG, where $h_e(A)$, $p_{\mathfrak{s}}^i(e)$ and $p_{\mathfrak{t}}^i(e)$ are Dirac observables from the deparametrization by coupling to clock fields [39].

On \mathcal{H}_γ , the holonomy becomes the multiplication operator and, $p_{\mathfrak{s}}^i(e)$ and $p_{\mathfrak{t}}^i(e)$ are quantized as the right- and left-invariant vector field, namely

$$\begin{aligned} (\hat{p}_{\mathfrak{s}}^i(e)\psi)(h_{e'}, \dots, h_e, \dots, h_{e''}) &= it \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} \psi(h_{e'}, \dots, e^{\epsilon\tau^i} h_e, \dots, h_{e''}) \\ (\hat{p}_{\mathfrak{t}}^i(e)\psi)(h_{e'}, \dots, h_e, \dots, h_{e''}) &= -it \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} \psi(h_{e'}, \dots, h_e e^{\epsilon\tau^i}, \dots, h_{e''}) \end{aligned} \quad (2.5)$$

where $t = \kappa\hbar/a^2 =: \ell_{\text{p}}^2/a^2$ and $\tau^j = (-i/2)\sigma^j$ with σ^j the Pauli matrix. The commutators between the basic operators are

$$\begin{aligned} [h(e), h(e')] &= 0 = [\hat{p}_{\mathfrak{s}}^i(e), \hat{p}_{\mathfrak{t}}^j(e')] \\ [\hat{p}_{\mathfrak{s}}^i(e), \hat{p}_{\mathfrak{s}}^j(e')] &= -it\delta_{ee'}\epsilon_{ijk}\hat{p}_{\mathfrak{s}}^k(e), \\ [\hat{p}_{\mathfrak{t}}^i(e), \hat{p}_{\mathfrak{t}}^j(e')] &= -it\delta_{ee'}\epsilon_{ijk}\hat{p}_{\mathfrak{t}}^k(e), \\ [\hat{p}_{\mathfrak{s}}^i(e), h(e')] &= it\delta_{ee'}\tau^i h(e), \\ [\hat{p}_{\mathfrak{t}}^i(e), h(e')] &= -it\delta_{ee'}h(e)\tau^i. \end{aligned} \quad (2.6)$$

It is useful to introduce the flux operators with respect to the spherical basis. We define

$$\hat{p}_v^{\pm 1}(e) := \mp \frac{1}{\sqrt{2}} (\hat{p}_v^x(e) \pm i\hat{p}_v^y(e)), \quad \hat{p}_v^0(e) = \hat{p}_v^z(e) \quad (2.7)$$

with $v = \mathfrak{s}, \mathfrak{t}$. In the following context, $\alpha, \beta, \dots = 0, \pm 1$ is used to denote the indices in the spherical basis, and $i, j, k \dots = 1, 2, 3$, the indices in the Cartesian basis.

In the operator-constraint formalism, the dynamics of LQG is encoded in the Hamiltonian constraint, which can be written as

$$\widehat{H}[N] = \widehat{H}_E[N] + (1 + \beta^2)\widehat{H}_L[N] \quad (2.8)$$

where $\widehat{H}_E[N]$ is called the Euclidean part and $\widehat{H}_L[N]$ is the Lorentzian part. N is the smeared function. $\widehat{H}[N]$ is constructed by using the Thiemann's trick [24, 25]. The operator corresponding to the Euclidean part is

$$\widehat{H}_E[N] = \frac{1}{2i\beta a^2 t} \sum_{v \in V(\gamma)} N(v) \sum_{e_I, e_J, e_K \text{ at } v} \epsilon^{IJK} \text{tr}(h_{\alpha_{IJ}} h_{e_K} [\hat{V}_v, h_{e_K}^{-1}]) \quad (2.9)$$

where e_I, e_J and e_K are oriented to be outgoing from v , $\epsilon^{IJK} = \text{sgn}[\det(e_I \wedge e_J \wedge e_K)]$, α_{IJ} is the minimal loop around a plaquette consisting of e_I and e_J , where it goes out via e_I and comes back through e_J , taking v as its end point. The volume operator \hat{V}_v at v reads

$$\hat{V}_v = \sqrt{|\hat{Q}_v|} \quad (2.10)$$

where the operator \hat{Q}_v in terms of the flux operators with respect to the spherical basis (2.7) is

$$\hat{Q}_v = -i(\beta a^2)^3 \epsilon_{\alpha\beta\gamma} \frac{\hat{p}_\mathfrak{s}^\alpha(e_x^+) - \hat{p}_\mathfrak{t}^\alpha(e_x^-)}{2} \frac{\hat{p}_\mathfrak{s}^\beta(e_y^+) - \hat{p}_\mathfrak{s}^\beta(e_y^-)}{2} \frac{\hat{p}_\mathfrak{s}^\gamma(e_z^+) - \hat{p}_\mathfrak{s}^\gamma(e_z^-)}{2} \quad (2.11)$$

with $\epsilon_{\alpha\beta\gamma}$ defined by $\epsilon_{-1,0,1} = 1$. With the same notion, the Lorentzian part reads

$$\widehat{H}_L[N] = \frac{-2}{i\beta^7 a^{10} t^5} \sum_v N(v) \sum_{e_I, e_J, e_K \text{ at } v} \epsilon^{IJK} \text{tr}([h_{e_I}, [\hat{V}, \hat{H}_E]] h_{e_I}^{-1} [h_{e_J}, [\hat{V}, \hat{H}_E]] h_{e_J}^{-1} [h_{e_K}, \hat{V}_v] h_{e_K}^{-1}). \quad (2.12)$$

2.2 Coherent states

Choosing a canonical orientation for each edge $e \in E(\gamma)$, the classical phase space based on the graph γ is

$$\Gamma_\gamma \cong [\text{SL}(2, \mathbb{C})]^{|E(\gamma)|}. \quad (2.13)$$

The complexifier coherent state $\Psi_{\mathbf{g}}$ is [40]

$$\Psi_{\mathbf{g}} = \bigotimes_{e \in E(\gamma)} \psi_{g_e}^t, \quad \psi_{g_e}^t(h_e) = \sum_j d_j e^{-\frac{\xi}{2} j(j+1)} \chi_j(g_e h_e^{-1}) \quad (2.14)$$

where $\psi_{g_e}^t$ is the $\text{SU}(2)$ coherent state at the edge e . The character $\chi_j(g_e h_e^{-1})$ is the trace of the j -representation of $g_e h_e^{-1}$. The property $\chi_j(g_e h_e^{-1}) = \chi_j(g_e^{-1} h_e)$ leads to the useful relation

$$\psi_{g_e}(h_e) = \psi_{g_{e^{-1}}}(h_{e^{-1}}).$$

Given $g = e^{ip_k \tau^k} u \in \text{SL}(2, \mathbb{C})$, one can find $n^s \in \text{SU}(2)$ such that $g = n^s e^{i\eta \tau_3} (n^s)^{-1} u$. Due to $(n^s)^{-1} u \in \text{SU}(2)$, one can decompose it as $(n^s)^{-1} u = e^{-\xi \tau_3} (n^t)^{-1}$. Finally, we have

$$g = n^s e^{i(\eta + i\xi) \tau_3} (n^t)^{-1}. \quad (2.15)$$

In this decomposition, one has $n^s \tau_3 (n^s)^{-1} = \vec{p} \cdot \vec{\tau} / \eta$, which implies $|\eta| = \sqrt{\vec{p} \cdot \vec{p}}$. In the current we choose the convention with $\eta = -\sqrt{\vec{p} \cdot \vec{p}}$. Moreover, this decomposition associates to $n^v \in \text{SU}(2)$ ($v = \mathfrak{s}, \mathfrak{t}$) a vector \vec{n}^v :

$$n^v \tau_3 (n^v)^{-1} = \vec{n}^v \cdot \vec{\tau}, \quad v = \mathfrak{s}, \mathfrak{t}. \quad (2.16)$$

Indeed, one has $\vec{n}^s = \vec{p} / \eta$ and $\vec{n}^t = (u^{-1} \triangleright \vec{p}) / \eta$ with $u^{-1} \triangleright \vec{p}$ denoting the vector of \vec{p} rotated by u^{-1} . It is shown in [41] and is revisited shortly that

$$\langle \psi_{g_e}^t | \vec{\hat{p}}_s(e) | \psi_{g_e}^t \rangle = -\eta_e \vec{n}_e^s + O(t), \quad \langle \psi_{g_e}^t | \vec{\hat{p}}_t(e) | \psi_{g_e}^t \rangle = \eta_e \vec{n}_e^t + O(t) \quad (2.17)$$

which indicates that $\eta_e \vec{n}_e^t$ is the classical limit of the flux operator at e .

The following properties of the ψ_g^t [40–42] are useful in our analysis. Firstly, the inner product of these states read

$$\langle \psi_{g_1}^t | \psi_{g_2}^t \rangle = \psi_{g_1^\dagger g_2}^{2t} (1) = \frac{2\sqrt{\pi} e^{t/4}}{t^{3/2}} \frac{\zeta e^{\frac{\zeta^2}{t}}}{\sinh(\zeta)} + O(t^\infty) \quad (2.18)$$

where $\text{tr}(g_1^\dagger g_2) = 2 \cosh(\zeta)$ and $\Im(\zeta) \in [0, \pi]$ with $\Im(\zeta)$ the imaginary part of ζ ². Consequently, the norm of the coherent state is

$$\langle 1 \rangle_g := \langle \psi_g^t | \psi_g^t \rangle = \frac{2\sqrt{\pi} e^{t/4}}{t^{3/2}} \frac{p e^{\frac{p^2}{t}}}{\sinh(p)} + O(t^\infty), \quad (2.19)$$

where $p = \sqrt{\vec{p} \cdot \vec{p}}$. Secondly, ψ_g^t satisfy the completeness condition

$$\int d\nu_t(g) |\psi_g^t\rangle \langle \psi_g^t| = \mathbb{I}, \quad (2.20)$$

where the measure $d\nu_t(g)$ is

$$d\nu_t(g) = \frac{2\sqrt{2} e^{-t/4}}{(2\pi t)^{3/2}} \frac{\sinh(p)}{p} e^{-\frac{p^2}{t}} d\mu_H(u) d^3 p = \frac{2}{\langle 1 \rangle_g \pi t^3} d\mu_H(u) d^3 p. \quad (2.21)$$

Let us complete this section with some discussions on the volume operator contained in the Hamiltonian operator $\widehat{H}[N]$. Because of the square root in the definition of the volume operator, matrix elements of these operators are difficult to compute analytically. However, as far as the coherent state expectation value is concerned, the volume operators \hat{V}_v in $\widehat{H}[N]$ can be replaced by Giesel-Thiemann's volume $\hat{V}_{GT}^{(v)}$ as aforementioned in (1.3) with $q = 1/4$. Substituting \hat{V}_v with $\hat{V}_{GT}^{(v)}$ truncated at finite k transforms $\widehat{H}[N]$ to be a polynomial of holonomies and fluxes. Up to higher order in t , it is now manageable to compute the expectation value of $\widehat{H}[N]$, through computing the expectation value of a polynomial of holonomies and fluxes.

3 Expectation values of operators on one edge

Because we only consider gauge *variant* coherent state, then the gauge *variant* coherent state can be written simply as the tensor product of SU(2) coherent state on each edge. As a consequence, computing the expectation value of $\widehat{H}[N]$ with respect to a coherent state can be reduced to computing expectation values of operator monomials on individual edges. In this section, let us firstly focus on the expectation value of operators on one edge.

²Here we used the following result shown in [41]. For any complex number $z = R + iI$, there exist real numbers $s \in \mathbb{R}$ and $\phi \in [0, \pi]$ such that $\cosh(s + i\phi) = z$. s and ϕ are uniquely determined except in the case $I = 0$ and $|R| > 1$ in which case the s is determined up to its sign.

Given a monomial of holonomies and fluxes on an edge e , its expectation value with respect to the coherent state $\psi_{g_e}^t$ labelled by $g_e = n_e^s e^{i z_e \tau_3} (n_e^t)^{-1}$ relates to its expectation value with respect to $\psi_{e^{i z_e \tau_3}}^t \equiv \psi_{z_e}^t$, by a gauge transformation generated by n_e^s and n_e^t [26]:

$$\begin{aligned} & \langle \psi_{g_e}^t | P(\{\hat{p}_s^{\alpha_i}(e)\}, \{\hat{p}_t^{\alpha_j}(e)\}, \{D_{a_k b_k}^{\iota_k}(h_e)\}) | \psi_{g_e}^t \rangle \\ &= \langle \psi_{z_e}^t | P(\{\hat{p}_s^{\beta_i}(e) D_{\beta_i \alpha_i}^1((n_e^s)^{-1})\}, \{\hat{p}_t^{\beta_j}(e) D_{\beta_j \alpha_j}^1((n_e^t)^{-1})\}, \{D_{a_k c_k}^{\iota_k}(n_e^s) D_{c_k d_k}^{\iota_k}(h_e) D_{d_k b_k}^{\iota_k}((n_e^t)^{-1})\}) | \psi_{z_e}^t \rangle, \end{aligned} \quad (3.1)$$

where $P(x, y, z)$ represents any monomial of x , y and z . In the following context, we denote

$$\langle \psi_{z_e}^t | \hat{F}_e | \psi_{z_e}^t \rangle =: \langle \hat{F}_e \rangle_{z_e}. \quad (3.2)$$

According to (3.1), in order to compute $\langle \psi_{g_e}^t | \hat{F}_e | \psi_{g_e}^t \rangle$, it is sufficient to compute expectation values with respect to $\psi_{z_e}^t$. The algorithm to compute the expectation value with respect to $\psi_{z_e}^t$ is divided into three steps.

3.1 the first step

Let \hat{O} be a monomial of flux and holonomy operators. To compute the expectation value of \hat{O} , we firstly move all the holonomies operators to the right, re-ordering the operators, by implementing the basic commutation relations (2.6). To this end, we need

$$\hat{O} = \hat{O}_2 \cdots \hat{O}_m \hat{O}_1 + \sum_{k=1}^{m-1} \sum_{\mathcal{I}_k} \left(\prod_{l \in \mathcal{I} - \mathcal{I}_k} \hat{O}_l \right) [[\cdots [[\hat{O}_1, \hat{O}_{i_1}], \hat{O}_{i_2}] \cdots], \hat{O}_{i_k}]. \quad (3.3)$$

with $\hat{O} = \hat{O}_1 \hat{O}_2 \cdots \hat{O}_m$, $\mathcal{I} := \{2, \dots, m\}$ and $\mathcal{I}_k := \{i_1, i_2, \dots, i_k\} \subset \mathcal{I}$ containing k elements with $i_1 < i_2 < \cdots < i_k$. In Eq. (3.3), the terms at k carry k -fold commutator. Due to the factor t in the right hand side of the commutation relations (2.6), the k -fold commutator produces a factor t^k in the final result, which implies that the contributions of these terms to the expectation value of \hat{O} are at least at t^k -order.

3.2 the second step

Once all of the holonomies are moved to the right, they can be merged into a single Wigner-D matrix by using the angular momentum recoupling theory

$$D_{m_1 n_1}^{j_1}(h) D_{m_2 n_2}^{j_2}(h) = \sum_{J=j_1-j_2}^{j_1+j_2} d_J (-1)^{M-N} \begin{pmatrix} j_1 & j_2 & J \\ m_1 & m_2 & -M \end{pmatrix} \begin{pmatrix} j_1 & j_2 & J \\ n_1 & n_2 & -N \end{pmatrix} D_{MN}^J(h). \quad (3.4)$$

Thus it is sufficient to consider the operators of the form $\hat{O}_1 = \hat{p}_s^{\alpha_1}(e) \cdots \hat{p}_s^{\alpha_m}(e) \hat{p}_t^{\beta_1}(e) \cdots \hat{p}_t^{\beta_n}(e) D_{ab}^{\iota}(h_e)$. To calculate the expectation value of \hat{O}_1 , we transform $\hat{p}_t^{\alpha}(e)$ to $\hat{p}_s^{\alpha}(e)$ with the identity [37]

$$\begin{aligned} & \langle p_s^{\alpha_1}(e) \cdots p_s^{\alpha_m}(e) p_t^{\beta_1}(e) \cdots p_t^{\beta_n}(e) D_{ab}^{\iota}(h_e) \rangle_{z_e} \\ &= (-1)^n e^{-(\beta_1 + \cdots + \beta_n) \bar{z}_e} \langle p_s^{\beta_n}(e) \cdots p_s^{\beta_1}(e) p_s^{\alpha_1}(e) \cdots p_s^{\alpha_m}(e) D_{ab}^{\iota}(h_e) \rangle_{z_e} \end{aligned} \quad (3.5)$$

where the factor $(-1)^n$ in the right hand side is because of the minus sign in the definition of $p_t^{\alpha}(e)$. In summary, the second step is to apply Eqs. (3.4) and (3.5) to simplify the concerning operator into a linear combination of operators taking the form

$$\hat{F}_{\iota ab}^{\alpha_1 \cdots \alpha_m} = \hat{p}_s^{\alpha_1}(e) \cdots \hat{p}_s^{\alpha_m}(e) D_{ab}^{\iota}(h_e).$$

The the last step is to calculate the expectation value of $\hat{F}_{\iota ab}^{\alpha_1 \cdots \alpha_m}$.

3.3 the third step

The strategy to compute the expectation value of the operator $\hat{F}_{iab}^{\alpha_1 \dots \alpha_m}$ is to write the action of the operator on the coherent state ψ_{z_e} in terms of Wigner $3jm$ -symbols [37, 43],

$$\langle \hat{F}_{iab}^{\alpha_1 \dots \alpha_m} \rangle_{z_e} = t^m e^{bz_e} \sum_{j, j'} e^{-\frac{t}{2}(j(j+1)+j'(j'+1))} F_\iota(j, j', \frac{\partial_\eta}{2}) \frac{\sinh((2j'+1)\eta)}{\sinh(\eta)} \quad (3.6)$$

with \hat{F}_ι expressed by Wigner $2j$ - and $3j$ - symbols and j , for each fixed half integer j' , taking values of $|j' - \iota|, |j' - \iota| + 1, \dots, |j' + \iota|$. The summation over j' will be computed with the Poisson summation formula. To do this, we extend the sum from $j' \geq 0$ to $j' \leq 0$ by applying the symmetry

$$F_\iota(j' + \Delta, j', k') = -F_\iota(j' - \Delta, j', k') \Big|_{j' \rightarrow -j'-1}, \quad (3.7)$$

which is proved in details in [37]. Then, summing over j' with $j = j' + \Delta$ and $j = j' - \Delta$ can be combined and rewritten as summing over all positive and negative half-integers j' , so that the Poisson summation formula can be applied. One can refer to [37] for the explicit expression of F_ι .

After applying the Poisson summation formula, Eq. (3.6) will simplified to take the form:

$$\langle \hat{F}_{iab}^{\alpha_1 \dots \alpha_m} \rangle_{z_e} = \sum_{\substack{0 \leq d \leq \iota \\ d + \iota \in \mathbb{Z}}} \int dx W_d(x) e^{-\frac{1}{4}t(x^2 - 2dx)} F_\iota\left(\frac{x-1}{2} - d, \frac{x-1}{2}, \frac{\partial_\eta}{2}\right) \frac{\sinh(x\eta)}{\sinh(\eta)} + O(t^{-\infty}) \quad (3.8)$$

with some function W_d . Then, according to the explicit result in [37], for at least $\iota \leq 20$, $F_\iota(\frac{x-1}{2} - d, \frac{x-1}{2}, \frac{\partial_\eta}{2}) \frac{\sinh(x\eta)}{\sinh(\eta)}$, as a function of x , is a summation of polynomials of x and the functions $\frac{\sinh(\eta(x-n))}{x-n}$ with n being some integers. Thus, the integral in Eq. (3.8) is a linear combination of integrals $I_1 = \int_{-\infty}^{\infty} dx e^{-ax^2+bx} \frac{\sinh(x\eta)}{x}$ and $I_2 = \int_{-\infty}^{\infty} dx e^{-ax^2+bx} x^n e^{\pm\eta x}$ where $a > 0, b \in \mathbb{R}$. The integrals I_1 and I_2 are easily to be computed. The only remaining problem is how to realize the concrete linear combination form of I_1 and I_2 , which can be quite technical and illustrated by the derivation of $\langle \hat{F}_{iab}^{\alpha_1 \dots \alpha_m} \rangle_{z_e}$ for $\iota = 1/2$ and $\iota = 1$ in [37].

We would like to compare our algorithm presented here with the known results in [28]. At first, our algorithm generalizes the known results in literature [28], in the sense that our formula gives the results for arbitrary lists $\{\alpha_i\}_{i=1}^m$ of flux indices and triples (ι, a, b) with at least $\iota \leq 20$. Moreover, with our formula, one can get the expectation values to arbitrary order of t . However, in [28], the authors give only the results for the special cases where the list $\{\alpha_i\}_{i=1}^m$ contains at most either a single -1 , or a single 1 , or a pair of $(-1, 1)$. They are all the cases such that the expectation values have non-vanishing $O(t^0)$ or $O(t)$ -term. Other cases are also interesting when we study the higher-order correction, even though the higher-order correction is beyond the present work. Our formula reduce to these known results at the special cases. The current work only use these special cases, but our codes [38] are designed based on the generalization algorithm, since the generalized formulae have the potential for computing higher-order correction.

In summary, we have outlined a general procedure to compute the expectation value of any monomial of holonomies and fluxes. According to this mechanism, we develop an algorithm in Mathematica to obtain analytical result [38].

4 Power counting

Applying $V_{GT}^{(v)}$ in (1.3) and truncating to a finite k , $\widehat{H[N]}$ is of the form $\sum_{\vec{\alpha}} \mathcal{T}^{\alpha_1 \dots \alpha_m} \hat{O}_{\alpha_1 \dots \alpha_m}$ with $\mathcal{T}^{\alpha_1 \dots \alpha_m}$ being some numerical factors and $\hat{O}_{\alpha_1 \dots \alpha_m}$ being some polynomial operators of holonomies

and fluxes. We consider gauge *variant* coherent state, namely, we do not introduce intertwiners and group averaging procedure. Then, the gauge *variant* coherent state can be written simply as the tensor product of SU(2) coherent state on each edge, so that the expectation can be factored into single edge expectation values. Thus the algorithm introduced in Sec. 3 can be applied to calculate the expectation value of $\hat{O}_{\alpha_1 \dots \alpha_m}$. To do this, in principle, we would need to compute the expectation values of $\hat{O}_{\alpha_1 \dots \alpha_m}$ for all indices $\alpha_1 \dots \alpha_m$. The computational complexity comes from the huge number of terms in the sum over $\vec{\alpha}$. Since we only focus on expanding $\langle \widehat{H[N]} \rangle$ to $O(t)$, the complexity can be reduced by certain power-counting argument: We count the least power of t contains in each $\langle \hat{O}_{\alpha_1 \dots \alpha_m} \rangle$ before explicit computation, then we omit those terms only contribute to higher order than $O(t)$ in $\langle \widehat{H[N]} \rangle$. It turns out that a large degree of complexity can be reduced in this manner.

To illustrate how we do power counting before explicit computation, let us introduce some conventions at first. In this section, we denote $\Psi_{\vec{g}}$ defined in (2.14) by $|\Psi_{\vec{g}}\rangle$ with $\vec{g} = \{g_e\}_{e \in E(\gamma)}$, namely

$$|\Psi_{\vec{g}}\rangle = \bigotimes_{e \in E(\gamma)} |\psi_{g_e}^t\rangle. \quad (4.1)$$

Similarly, $|\Psi_{\vec{g}^{(i)}}\rangle$ denotes the coherent state in which $|\psi_{g_e}^t\rangle$ is at edge e . Let \hat{O} take the form of

$$\hat{O} = \hat{O}_1 \hat{O}_2 \dots \hat{O}_k, \quad (4.2)$$

with \hat{O}_i being polynomial of fluxes and holonomies. Besides, we assume that the matrix elements \hat{O}_i with respect to coherent states $\psi_{g_e^{(1)}}^t$ and $\psi_{g_e^{(2)}}^t$ on e take the form

$$\langle \Psi_{\vec{g}^{(1)}} | \hat{O}_i | \Psi_{\vec{g}^{(2)}} \rangle = \langle \Psi_{\vec{g}^{(1)}} | \Psi_{\vec{g}^{(2)}} \rangle \left(E_0^{(i)}(\vec{g}^{(1)}, \vec{g}^{(2)}) + t E_1^{(i)}(\vec{g}^{(1)}, \vec{g}^{(2)}) + O(t^\infty) \right). \quad (4.3)$$

Note that this assumption is always satisfied by the operators considered in the current work. Indeed, according to the results in [41], if \hat{O}_i is just a flux or holonomy on some edge e , its matrix elements with respect to coherent states $\psi_{g_e^{(1)}}^t$ and $\psi_{g_e^{(2)}}^t$ on e take the form

$$\langle \psi_{g_e^{(1)}}^t | \hat{O}_i | \psi_{g_e^{(2)}}^t \rangle = \langle \psi_{g_e^{(1)}}^t | \psi_{g_e^{(2)}}^t \rangle \left(E_0(g_e^{(1)}, g_e^{(2)}) + t E_1(g_e^{(1)}, g_e^{(2)}) + O(t^\infty) \right). \quad (4.4)$$

Let N_0 be the number of operators $\hat{O}_m \in \{\hat{O}_i\}_{i=1}^k$ such that

$$\frac{\langle \Psi_{\vec{w}} | \hat{O}_m | \Psi_{\vec{w}} \rangle}{\langle \Psi_{\vec{w}} | \Psi_{\vec{w}} \rangle} = O(t), \quad (4.5)$$

where the $O(t^0)$ term vanishes on the RHS. Using (4.3), it turns out that the expectation value of \hat{O} with respect to the coherent state $|\Psi_{\vec{w}}\rangle$ satisfies [37]

$$\frac{\langle \Psi_{\vec{w}} | \hat{O} | \Psi_{\vec{w}} \rangle}{\langle \Psi_{\vec{w}} | \Psi_{\vec{w}} \rangle} = O(t^n), \quad \text{with } n \geq \lfloor \frac{N_0 + 1}{2} \rfloor \quad (4.6)$$

where $\lfloor x \rfloor$ is the largest integer no larger than x .

In the practical computation of $\langle \widehat{H[N]} \rangle$ with the application of $\hat{V}_{GT}^{(v)}$, these N_0 operators \hat{O}_m satisfying (4.5) can be the operator $\frac{\hat{Q}}{\langle \hat{Q} \rangle} - 1$. Then we use Eq. (4.6) to count the power of t for the term with $\left(\frac{\hat{Q}^2}{\langle \hat{Q} \rangle^2} - 1 \right)^k$ in $\langle \widehat{H[N]} \rangle$. Note that Eq. (4.6) needs the matrix elements of \hat{O}_i . We have to factorize $\frac{\hat{Q}^2}{\langle \hat{Q} \rangle^2} - 1 = \left(\frac{\hat{Q}}{\langle \hat{Q} \rangle} + 1 \right) \left(\frac{\hat{Q}}{\langle \hat{Q} \rangle} - 1 \right)$, because every matrix element of \hat{Q} is a polynomial of matrix elements of flux operators, while that of \hat{Q}^2 is not.

Moreover, these N_0 operators satisfying (4.5) can also be $\hat{p}_s^{\pm 1}(e)$, $\hat{p}_t^{\pm 1}(e)$ and $D_{ab}^k(h_e)$ with $a \neq b$ whose expectation values with respect to $\Psi_{\vec{\omega}}$ vanish. Then this theorem can be used to study the leading order of monomial of holonomies and fluxes. Let us fix $\hat{p}^\beta(e)$ to be either $\hat{p}_s^\beta(e)$ or $\hat{p}_t^\beta(e)$ and, use \mathcal{M} to denote the monomial of holonomies $\{D_{a_j b_j}^{\frac{1}{2}}(h_e)\}_{j=1}^k$ and fluxes $\{\hat{p}^{\beta_i}(e)\}_{i=1}^m$. Let N_\pm be the number of $\hat{p}^{\pm 1}(e) \in \{\hat{p}^{\beta_i}(e)\}_{i=1}^m$ and M_+ (respectively M_-) be the number of $D_{-\frac{1}{2} \frac{1}{2}}^{\frac{1}{2}}(h_e) \in \{D_{a_j b_j}^{\frac{1}{2}}(h_e)\}_{j=1}^k$ (respectively $D_{\frac{1}{2} -\frac{1}{2}}^{\frac{1}{2}}(h_e) \in \{D_{a_j b_j}^{\frac{1}{2}}(h_e)\}_{j=1}^k$). According to our analysis above, the expectation value of \mathcal{M} with respect to the coherent state $|\psi_{z_e}\rangle$ with $z_e \in \mathbb{C}$ is non-vanishing provided that

$$\sum_{i=1}^m \beta_i + \sum_{j=1}^k (b_j - a_j) = 0. \quad (4.7)$$

Thus it has

$$N_+ + M_+ = N_- + M_-. \quad (4.8)$$

Therefore, this theorem tells us that the leading order the expectation value of $\langle \mathcal{M} \rangle_{z_e}$ is of order $O(t^{N_+ + M_+})$ or higher.

We have more discussions on this case. Since the matrix elements of $\hat{p}^\alpha(e)$ and $D_{ab}^{\frac{1}{2}}(h_e)$ are computable, the results on the leading order of \mathcal{M} can be calculated more exactly by applying the generalized stationary phase approximation analysis. A straightforward calculation tells us

$$\langle \mathcal{M} \rangle_{z_e} \cong (\langle \hat{p}_s^0(e) \rangle_{z_e})^{N_{0,s}} (\langle \hat{p}_t^0(e) \rangle_{z_e})^{N_{0,t}} (\langle D_{\frac{1}{2} \frac{1}{2}}^{\frac{1}{2}}(h_e) \rangle_{z_e})^{M_{0,+}} (\langle D_{-\frac{1}{2} -\frac{1}{2}}^{\frac{1}{2}}(h_e) \rangle_{z_e})^{M_{0,-}} \langle \mathcal{M}' \rangle_{z_e} \quad (4.9)$$

where \cong means the leading-order terms, i.e., the $O(t^{M_+ + N_+})$ terms, of the left and right hand sides are equal to each other, \mathcal{M}' is the operator resulting from \mathcal{M} by deleting all factors $\hat{p}^0(e)$ and $D_{aa}^{\frac{1}{2}}(h_e)$, $N_{0,s}$ and $N_{0,t}$ are the number of $\hat{p}_s^0(e)$ and $\hat{p}_t^0(e)$ in \mathcal{M} , and $M_{0,+}$ and $M_{0,-}$ the number of $D_{\frac{1}{2} \frac{1}{2}}^{\frac{1}{2}}(h_e)$ and $D_{-\frac{1}{2} -\frac{1}{2}}^{\frac{1}{2}}(h_e)$ in \mathcal{M} .

5 Cosmological expectation value

We apply our computation of expectation values to coherent states labelled by homogeneous and isotropic data. The symmetry group of the homogeneous and isotropic cosmology is $\mathbb{T} \rtimes F$ with F being the isotropy subgroup, \mathbb{T} being the translation subgroup. Let S_γ be the subgroup of $\mathbb{T} \rtimes F$ preserving γ . A classical state \mathbf{g} is called symmetric with respect to S_γ if $s^* \mathbf{g} := \mathbf{g} \circ s$ are the same as \mathbf{g} up to a gauge transformation for all $s \in S_\gamma$. By this definition, a classically symmetric state \mathbf{g} takes the form

$$\mathbf{g} : e \mapsto g_e = n_e e^{iz\tau_3} n_e^{-1} \quad (5.1)$$

where $n_e \in \text{SU}(2)$ satisfies

$$n_e \tau_3 n_e^{-1} = \vec{n}_e \cdot \vec{\tau} \quad (5.2)$$

\vec{n}_e is the unit vector pointing to direction of edge e . Then for each $s = (T, f) \in \mathbb{T} \rtimes F$, it has

$$\mathbf{g} \circ s = \text{Ad}_f \circ \mathbf{g} \quad (5.3)$$

where $\text{Ad}_f \circ \mathbf{g}(e) = f \mathbf{g}(e) f^{-1}$ for all $e \in E(\gamma)$ with $f \in \text{SU}(2)$.

5.1 Symmetries of the expectation value

Let \hat{F}_e be an operator of polynomial of fluxes and holonomies on e . For $s = (T, f) \in \mathbb{T} \rtimes F$, Eq. (5.3) leads to that

$$\langle \psi_{g_{s(e)}} | \hat{F}_{s(e)} | \psi_{g_{s(e)}} \rangle = \langle \psi_{fg_e f^{-1}} | \hat{F}_e | \psi_{fg_e f^{-1}} \rangle = f \triangleright \langle \psi_{g_e} | \hat{F}_e | \psi_{g_e} \rangle \quad (5.4)$$

where $f \triangleright \langle \psi_{g_e} | \hat{F}_e | \psi_{g_e} \rangle$ means the gauge transformation to the expectation value $\langle \psi_{g_e} | \hat{F}_e | \psi_{g_e} \rangle$ and Eq. (3.1) was used to derive the last equality.

To expand the expectation value of \hat{H}_E and \hat{H}_L to order $O(t)$, we first replace the operator \hat{V}_v by $\hat{V}_{GT}^{(v)}$ in (1.3). Then the Euclidean part $\widehat{H}_E[\mathcal{N}]$ can be rewritten in terms of (there is no summation over I, J, K here)

$$\hat{H}_E^{(n)}(v; e_I, e_J, e_K) = \frac{1}{i\beta a^2 t} \epsilon_{IJK} \text{tr}(h_{\alpha_{IJ}} [h_{e_K}, \hat{Q}_v^{2n}] h_{e_K}^{-1}), \quad (5.5)$$

and the Lorentzian part, in terms of

$$\begin{aligned} & \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_I, e_J, e_K) \\ &= \frac{-2}{i\beta^7 a^{10} t^5} \epsilon^{IJK} \text{tr}([h_{e_I}, [\hat{Q}_{v_1}^{2k_1}, \hat{H}_E^{(k_2)}(v_2)]] h_{e_I}^{-1} [h_{e_J}, [\hat{Q}_{v_3}^{2k_3}, \hat{H}_E^{(k_4)}(v_4)]] h_{e_J}^{-1} [h_{e_K}, \hat{Q}_v^{2k_5}] h_{e_K}^{-1}) \end{aligned} \quad (5.6)$$

where $\vec{k} = (k_1, k_2, k_3, k_4, k_5)$. Let us define

$$\hat{H}_E^{(n)}(v) = \sum_{e_I, e_J, e_K} \hat{H}_E^{(n)}(v; e_I, e_J, e_K) \quad (5.7)$$

and

$$\hat{H}_L^{(\vec{k})}(v) = \sum_{v_1, v_2, v_3, v_4, e_I, e_J, e_K} \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_I, e_J, e_K) \quad (5.8)$$

The Euclidean and Lorentzian parts with the replacement \hat{V}_v by $\hat{V}_{GT}^{(v)}$, truncated at a finite k of (1.3), are given by linear combinations of $\hat{H}_E^{(n)}(v)$ and $\hat{H}_L^{(\vec{k})}(v)$ for various n and \vec{k} respectively.

5.1.1 Symmetries of the Euclidean part

By Eq. (5.4) and the fact that $\hat{H}_E^{(n)}(v; e_I, e_J, e_K)$ is gauge invariant, we obtain that

$$\langle \hat{H}_E^{(n)}(v; e_I, e_J, e_K) \rangle = \langle \hat{H}_E^{(n)}(s(v); s(e_I), s(e_J), s(e_K)) \rangle \quad (5.9)$$

where $\langle \cdot \rangle$ denotes the expectation value with respect to the cosmological coherent state defined by (5.1) and $s = (T, f)$ as defined in Eq. (5.4) is a symmetry of the graph. According to this relation, Eq. (5.7) can be simplified as

$$\hat{H}_E^{(n)}(v) = 24(\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+) + \hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^-)) \quad (5.10)$$

where the prefactor 24 is because there are 48 terms in the RHS of (5.7).

Moreover, $[h_{e_z^\pm}, \hat{Q}_v^{2n}] h_{e_z^\pm}^{-1}$ appearing in $\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^\pm)$ helps us to obtain the relation between the operators $\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^\pm)$. Indeed,

$$[h_{e_z^\pm}, \hat{Q}_v^{2n}] h_{e_z^\pm}^{-1} = \sum_{l=1}^{2n} \sum_{\mathcal{P}_l} \left(\mp i t \frac{(\beta a^2)^3}{8} \right)^l \hat{Q}_v^{p_1} \epsilon_{\alpha_1 \beta_1 \gamma_1} \hat{X}^{\alpha_1} \hat{Y}^{\beta_1} \tau^{\gamma_1} \hat{Q}_v^{p_2} \epsilon_{\alpha_2 \beta_2 \gamma_2} \hat{X}^{\alpha_2} \hat{Y}^{\beta_2} \tau^{\gamma_2} \dots \hat{Q}_v^{p_l} \epsilon_{\alpha_l \beta_l \gamma_l} \hat{X}^{\alpha_l} \hat{Y}^{\beta_l} \tau^{\gamma_l} \hat{Q}_v^{p_{l+1}} \quad (5.11)$$

where e_z^\pm are oriented such that $\mathfrak{s}(e_z^+) = v = \mathfrak{s}(e_z^-)$, $\hat{X}^\alpha = \hat{p}_\mathfrak{s}^\alpha(e') - \hat{p}_\mathfrak{t}^\alpha(e')$ and $\hat{Y}^\alpha = \hat{p}_\mathfrak{s}^\alpha(e'') - \hat{p}_\mathfrak{t}^\alpha(e'')$, and $\mathcal{P} = \{p_1, p_2, \dots, p_{l+1}\}$ with $p_i \in \mathbb{Z}$, $p_i \geq 0$ and $\sum_{i=1}^{l+1} p_i = 2n - l$.³ Substituting the last equation, we have that

$$\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+) + \hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^-) = 2\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+) \quad (5.12)$$

where $\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+)$ is the operator of $\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+)$ with the replacement

$$\begin{aligned} [h_{e_z^+}, \hat{Q}_v^{2n}] h_{e_z^+}^{-1} \rightarrow & \sum_{l \text{ is odd}} \sum_{\mathcal{P}_l} \left(-it \frac{(\beta a^2)^3}{8} \right)^l \hat{Q}_v^{p_1} \epsilon_{\alpha_1 \beta_1 \gamma_1} \hat{X}^{\alpha_1} \hat{Y}^{\beta_1} \tau^{\gamma_1} \hat{Q}_v^{p_2} \epsilon_{\alpha_2 \beta_2 \gamma_2} \hat{X}^{\alpha_2} \hat{Y}^{\beta_2} \tau^{\gamma_2} \dots \hat{Q}_v^{p_l} \times \\ & \epsilon_{\alpha_l \beta_l \gamma_l} \hat{X}^{\alpha_l} \hat{Y}^{\beta_l} \tau^{\gamma_l} \hat{Q}_v^{p_{l+1}}. \end{aligned} \quad (5.13)$$

According to Eq. (5.10), $\hat{H}_E^{(n)}(v) = 48\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+)$. Thus when calculating the expectation value of the Euclidean part, we only consider the operator $\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+)$ instead of $\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^\pm)$.

Moreover, Eq. (5.11) tells us that the Euclidean Hamiltonian takes the form

$$\hat{H}_E^{(n)}(v; e_I, e_J, e_K) = \epsilon_{IJK} \text{tr}(h_{\alpha_I J} \tau^\alpha) \hat{O}_\alpha,$$

with \hat{O}_α a polynomial of fluxes. Because of $\text{tr}(h\tau^\alpha) = -\text{tr}(h^{-1}\tau^\alpha)$, we obtain that

$$\hat{H}_E^{(n)}(v; e_I, e_J, e_K) = \hat{H}_E^{(n)}(v; e_J, e_I, e_K). \quad (5.14)$$

In summary, originally there are 48 terms for each $\hat{H}_E^{(n)}(v)$ in Eq. (5.7). Once we take into account the symmetries discussed in this section, we have $\hat{H}_E^{(n)}(v) = 48\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+)$. Thus we only need to compute the expectation value of the single term $\hat{H}_E^{(n)}(v; e_x^+, e_y^+, e_z^+)$.

5.1.2 Symmetries of the Lorentzian part

Given a list of vertices and edges $(v; v_1, v_2, v_3, v_4; e_I, e_J, e_K)$ where e_I, e_J and e_K are outgoing from v , (e_I, e_J, e_K) is either left-handed or right-handed. Thus there exists a rotation f , leaving v invariant, such that $(v; f(v_1), f(v_2), f(v_3), f(v_4); f(e_I), f(e_J), f(e_K))$ is either $(v; f(v_1), f(v_2), f(v_3), f(v_4); e_x^+, e_y^+, e_z^+)$ or $(v; f(v_1), f(v_2), f(v_3), f(v_4); e_x^+, e_y^+, e_z^-)$. Therefore, (5.8) can be simplified as

$$\hat{H}_L^{(\vec{k})}(v) = 24 \sum_{v_1, v_2, v_3, v_4} \left(\hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^+) + \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^-) \right) \quad (5.15)$$

Moreover, because the term $[h_{e_z^+}, \hat{Q}_v^{2n}] h_{e_z^+}^{-1}$ also appears in $\hat{H}_L^{(\vec{k})}$, Eq. (5.11) can be applied again which simplifies Eq. (5.15) further to

$$\hat{H}_L^{(\vec{k})}(v) = 48 \sum_{v_1, v_2, v_3, v_4} \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^+) \quad (5.16)$$

where the $\hat{H}_L^{(\vec{k})}$ operator is obtained from $\hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^+)$ by the replacement of Eq. (5.13). As a result, we only need to compute the expectation value of the operator $\hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^+)$ for various vertices v_1, v_2, v_3 and v_4 .

³A general equation can be obtained analogously if the holonomy $h_{e_z^\pm}$ is replaced by a holonomy along other edges. In the following context, Eq. (5.11) will be usually referred as this general equation.

The above discussion helps us to simplify our computation for the Lorentzian part. However, much more symmetries are needed so that the computation time is acceptable. For this purpose, let us firstly look at the term $[h_e[\hat{Q}_{v_1}^m, h_e^{-1}], \hat{Q}_{v_2}]$ which comes from the commutator between the volume operator and the Euclidean Hamiltonian. We obtain the following proposition which can be proven by using Eq. (5.11).

Proposition 1. *Given an edge e with the source $s(e)$ and the target $t(e)$, $[h_e[\hat{Q}_{s(e)}^m, h_e^{-1}], \hat{Q}_v] = 0$ for all $v \neq s(e)$.*

With this proposition, let us consider the the commutator $[\hat{Q}_{v_1}^{2k}, \hat{H}_E^{(n)}(v_2; e_I, e_J, e_K)]$ which defines the operator \hat{K} .

$$\begin{aligned} & [\hat{Q}_{v_1}^{2k}, \hat{H}_E^{(n)}(v_2; e_I, e_J, e_K)] \\ &= \frac{1}{i\beta a^2 t} \epsilon_{IJK} \left(\text{tr}([\hat{Q}_{v_1}^{2k}, h_{\alpha_{IJ}}][h_{e_K}, \hat{Q}_{v_2}^{2n}]h_{e_K}^{-1}) + \text{tr}(h_{\alpha_{IJ}}[\hat{Q}_{v_1}^{2k}, [h_{e_K}, \hat{Q}_{v_2}^{2n}]h_{e_K}^{-1}]) \right) \end{aligned} \quad (5.17)$$

The commutators on the RHS of Eq. (5.17) can be expanded in t with Eq. (3.3) and simplified by the basic commutation relation (2.6), which can finally simplify $\hat{H}_L^{(\vec{k})}(v)$ of Eq. (5.16) to be in terms of

$$\frac{C}{t^2} \text{tr}(h_{e_x^+} F_1 h_{e_x^+}^{-1} h_{e_y^+} F_2 h_{e_y^+}^{-1} G_1) \quad (5.18)$$

with C being some constant of order t^0 or higher, F_i being some monomials of holonomies and $(p_s^\alpha(e^+) \pm p_t^\alpha(e^-))$ and G_1 being a monomial of $(p_s^\alpha(e^+) - p_t^\alpha(e^-))$.

We apply the results in Sec. 4 to reduce the computational complexity. To use these results, we need to use the basic computation relations (2.6) to simplify the Hamiltonian operator, so that the operators after the simplification are written in terms of $C\hat{P}$ where C is some constant of order t^0 or higher and \hat{P} is some monomial of holonomies and fluxes.

In order to achieve so, we need to permute $h_{e_x^+}$ and \hat{F}_1 , as well as $h_{e_y^+}$ and \hat{F}_2 , in Eq. (5.18) by Eq. (3.3). Take permuting $h_{e_x^+}$ and \hat{F}_1 as an example: Implementing the result of Eq. (3.3), we substitute \hat{O}_1 by $h_{e_x^+}$, and \hat{O}_{i_k} by $\hat{p}_s^\alpha(e_x^+)$ and/or $\hat{p}_t^\alpha(e_x^+)$. One of many such substitutions inevitably generates some special terms whose commutators only contain $\hat{p}_s^\alpha(e_x^+)$. Computing these commutators with (2.6) leads to results that are proportional to $h_{e_x^+}$. After substituting the permuted results into Eq. (5.18), the $h_{e_x^+}$ eventually cancels with $h_{e_x^+}^{-1}$. (similar to Eq. (5.11)). One can apply the same mechanism for permuting $h_{e_y^+}$ and \hat{F}_2 .

Let us collect these special terms from permuting $h_{e_x^+}$ and \hat{F}_1 and permuting $h_{e_y^+}$ and \hat{F}_2 . We denote the partial sum of these special terms in $\hat{H}_L^{(k)}$ by $\text{alt}\hat{H}_L^{(\vec{k})}$. Because of the cancellation between holonomies and their inverses discussed above, $\text{alt}\hat{H}_L^{(\vec{k})}$ does not depend on $h_{e_x^+}$ and $h_{e_y^+}$.

It turns out that $\text{alt}\hat{H}_L^{(\vec{k})}$ has more symmetries which are discussed shortly below. These special terms can be equivalently selected by considering only the non-commutativity between $h_{e_x^+}$ and $p_s^\alpha(e_x^+)$ while ignoring the non-commutativity between $h_{e_x^+}$ and $p_t^\alpha(e_x^+)$. Hence we have

$$\text{the special terms of } h_{e_x^+} \hat{F}_1 h_{e_x^+}^{-1} = h_{s_x^+} \hat{F}_1 h_{s_x^+}^{-1} \quad (5.19)$$

where s_x^+ is the segment within e_x^+ and does not contain the target $t(e_x^+)$. Note that the length of the segment does not cause any ambiguity because of the cancellation between the holonomies and

their inverses discussed above. Indeed, Eq. (5.19) leads to

$$\begin{aligned} & \text{alt } \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_I, e_J, e_K) \\ &= \frac{-2}{i\beta^7 a^{10} t^5} \epsilon^{IJK} \text{tr} \left([h_{s_I}, [\hat{Q}_{v_1}^{2k_1}, \hat{H}_E^{(k_2)}(v_2)]] h_{s_I}^{-1} [h_{s_J}, [\hat{Q}_{v_3}^{2k_3}, \hat{H}_E^{(k_4)}(v_4)]] h_{s_J}^{-1} [h_{s_K}, \hat{Q}_v^{2k_5}] h_{s_K}^{-1} \right). \end{aligned} \quad (5.20)$$

Interestingly, the RHS may be understood as from an alternative definition of the Lorentzian part,

$$\text{alt } \widehat{H}_L[N] = \frac{-2}{i\beta^7 a^{10} t^5} \sum_v N(v) \sum_{s_I, s_J, s_K \text{ at } v} \epsilon^{IJK} \text{tr} \left([h_{s_I}, [\hat{V}, \hat{H}_E]] h_{s_I}^{-1} [h_{s_J}, [\hat{V}, \hat{H}_E]] h_{s_J}^{-1} [h_{s_K}, \hat{V}_v] h_{s_K}^{-1} \right). \quad (5.21)$$

where all edges e_I, e_J, e_K are replaced by the segments s_I, s_J, s_K with $s_I \subset e_I$. $\text{alt } \widehat{H}_L[N]$ may be obtained by a different regularization/quantization,

$$\{K, e^a A_a(x)\} \rightarrow \frac{-1}{2\kappa(i\hbar\beta)^2} [h_{s_e}, [\hat{V}, \hat{H}_E]] h_{s_e}^{-1},$$

with e^a the vector tangent to the edge e and $s_e \subset e$ the segment. More details on this aspect is discussed in [37].

We collect terms in $\hat{H}_L^{(k)}$ that are other than the special terms discussed above, and denote their sum by $\text{extra } \hat{H}_L^{(\vec{k})}$, namely

$$\begin{aligned} & \text{extra } \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_I, e_J, e_K) \\ &= \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_I, e_J, e_K) - \text{alt } \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_I, e_J, e_K). \end{aligned} \quad (5.22)$$

The operators $\text{alt } \hat{H}_L^{(\vec{k})}$ and $\text{extra } \hat{H}_L^{(\vec{k})}$ are treated separately in our algorithm.

For $\text{alt } \hat{H}_L^{(\vec{k})}$, the above simplification procedures lead to

$$\frac{C}{t^2} h_{s_x^\pm} F_1 h_{s_x^\pm}^{-1} h_{s_y^\pm} F_2 h_{s_y^\pm}^{-1} G_1, \quad (5.23)$$

instead of Eq. (5.18). Since $[h_{s_I}, \hat{p}_t^\alpha(e_I)] = 0$, these terms can be simplified with

$$h_{s^\pm} \prod_{i=1}^m (\sigma_i^+ p_s^{\alpha_i}(e^+) + \sigma_i^- p_t^{\alpha_i}(e^-)) h_{s^\pm}^{-1} = \sum_{\mathcal{I}} (it)^{|\mathcal{I}|} \prod_{i \notin \mathcal{I}} (\sigma_i^+ p_s^{\alpha_i}(e^+) + \sigma_i^- p_t^{\alpha_i}(e^-)) \prod_{j \in \mathcal{I}} \sigma_j^\pm \tau^{\alpha_j} \quad (5.24)$$

where \mathcal{I} is a subsets of $\{1, 2, \dots, m\}$ with length $|\mathcal{I}|$, s^+ and s^- are segments of e^+ and $(e^-)^{-1}$ respectively, $\sigma_i^+ = 1$ and $\sigma_i^- = \pm 1$. Because of the summation over e_I, e_J and e_K in Eq. (5.8) which lets us to compute

$$h_{s^+} \prod_{i=1}^m (\sigma_i^+ p_s^{\alpha_i}(e^+) + \sigma_i^- p_t^{\alpha_i}(e^-)) h_{s^+}^{-1} - h_{s^-} \prod_{i=1}^m (\sigma_i^+ p_s^{\alpha_i}(e^+) + \sigma_i^- p_t^{\alpha_i}(e^-)) h_{s^-}^{-1}, \quad (5.25)$$

we can use the replacement

$$h_{s^+} \prod_{i=1}^m (\sigma_i^+ p_s^{\alpha_i}(e^+) + \sigma_i^- p_t^{\alpha_i}(e^-)) h_{s^+}^{-1} \rightarrow \sum_{\mathcal{J}} (it)^{|\mathcal{J}|} \prod_{i \notin \mathcal{J}} (\sigma_i^+ p_s^{\alpha_i}(e^+) + \sigma_i^- p_t^{\alpha_i}(e^-)) \prod_{j \in \mathcal{J}} \tau^{\alpha_j} \quad (5.26)$$

with $\mathcal{J} \subset \{1, 2, \dots, m\}$ such that

$$\prod_{j \in \mathcal{J}} \sigma_j^- = -1. \quad (5.27)$$

Substituting Eq. (5.26) into (5.23), we cancel the prefactor $1/t^2$ to simplify (5.23) to be in terms of $C\hat{P}$ with some constant C of order t^0 or higher, so that the results in Sec. 4 can be applied.

Moreover, $\text{alt } \hat{H}_L^{(\vec{k})}$ brings more symmetries, which is described as below: Consider a π -rotation which transforms either e_x^+ to e_x^- or e_y^+ to e_y^- . Denote this rotation by \mathfrak{s} . Let us rewrite F_i in Eq. (5.23) as $F_i(v, e)$ to indicate the dependence of F_i on vertices and edges. Then Eq. (5.24) tells us

$$\begin{aligned} & h_{\mathfrak{s}(s_x^+)} F_1(\mathfrak{s}(v), \mathfrak{s}(e)) h_{\mathfrak{s}(s_x^+)}^{-1} h_{\mathfrak{s}(s_y^+)} F_2(\mathfrak{s}(v), \mathfrak{s}(e)) h_{\mathfrak{s}(s_y^+)}^{-1} G_1(\mathfrak{s}(v), \mathfrak{s}(e)) \\ &= h_{s_x^+} F_1(\mathfrak{s}(v), \mathfrak{s}(e)) h_{s_x^+}^{-1} h_{s_y^+} F_2(\mathfrak{s}(v), \mathfrak{s}(e)) h_{s_y^+}^{-1} G_1(\mathfrak{s}(v), \mathfrak{s}(e)). \end{aligned} \quad (5.28)$$

Furthermore, recalling Eq. (5.4), we obtain that

$$\begin{aligned} & \langle h_{s_x^+} F_1(v, e) h_{s_x^+}^{-1} h_{s_y^+} F_2(v, e) h_{s_y^+}^{-1} G_1(v, e) \rangle \\ &= \langle h_{s_x^+} F_1(\mathfrak{s}(v), \mathfrak{s}(e)) h_{s_x^+}^{-1} h_{s_y^+} F_2(\mathfrak{s}(v), \mathfrak{s}(e)) h_{s_y^+}^{-1} G_2(\mathfrak{s}(v), \mathfrak{s}(e)) \rangle \end{aligned} \quad (5.29)$$

which reduces the number of contributing vertices and edges in the computation of $\text{alt } \hat{H}_L^{(\vec{k})}$.

For the operator $\text{extr } \hat{H}_L^{(\vec{k})}$, Eq. (5.24) can no longer be used and, thus, there does not exist the symmetry implied by (5.29) yet. We propose the following strategy to reduce the complexity of the computation. Let \mathfrak{r} be the rotation about the axis $(1/\sqrt{2}, 1/\sqrt{2}, 0)$ for π radians, which exchanges the x - and y -axes, and flips the z -axis. We have

$$\hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^+) = \hat{H}_L^{(\vec{k})}(v; \mathfrak{r}(v_1), \mathfrak{r}(v_2), \mathfrak{r}(v_3), \mathfrak{r}(v_4); e_y^+, e_x^+, e_z^+) \quad (5.30)$$

where $\hat{H}_L^{(\vec{k})}$ is defined in (5.16) and $\mathfrak{r}(v) = v$ is assumed without loss of generality. Let us consider the operator

$$\begin{aligned} & \hat{F}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^+) \\ &:= \hat{H}_L^{(\vec{k})}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^+) + \hat{H}_L^{(\vec{k})}(v; \mathfrak{r}(v_3), \mathfrak{r}(v_4), \mathfrak{r}(v_1), \mathfrak{r}(v_2); e_x^+, e_y^+, e_z^+) \end{aligned} \quad (5.31)$$

By (5.30), \hat{F} , up to an overall factor, is

$$\begin{aligned} & \hat{F}(v; v_1, v_2, v_3, v_4; e_x^+, e_y^+, e_z^+) \\ &= -\text{tr}([h_{e_y^+}, [\hat{Q}_{\mathfrak{r}(v_1)}^{2k_1}, \hat{H}_E^{(k_2)}(\mathfrak{r}(v_2))]] h_{e_x^+}^{-1} [h_{e_x^+}, [\hat{Q}_{v_3}^{2k_3}, \hat{H}_E^{(k_4)}(\mathfrak{r}(v_4))]] h_{e_x^+}^{-1} [h_{e_z^+}, \hat{Q}_v^{2k_5}] h_{e_z^+}^{-1}) \\ &+ \text{tr}([h_{e_x^+}, [\hat{Q}_{\mathfrak{r}(v_3)}^{2k_1}, \hat{H}_E^{(k_2)}(\mathfrak{r}(v_4))]] h_{e_x^+}^{-1} [h_{e_y^+}, [\hat{Q}_{\mathfrak{r}(v_1)}^{2k_3}, \hat{H}_E^{(k_4)}(\mathfrak{r}(v_2))]] h_{e_y^+}^{-1} [h_{e_z^+}, \hat{Q}_v^{2k_5}] h_{e_z^+}^{-1}) \end{aligned} \quad (5.32)$$

For clarity, let us denote

$$\begin{aligned} \hat{X} &\equiv [h_{e_x^+}, [\hat{Q}_{v_3}^{2k_3}, \hat{H}_E^{(k_4)}(\mathfrak{r}(v_4))]] h_{e_x^+}^{-1} \\ \hat{Y} &\equiv [h_{e_y^+}, [\hat{Q}_{\mathfrak{r}(v_1)}^{2k_1}, \hat{H}_E^{(k_2)}(\mathfrak{r}(v_2))]] h_{e_y^+}^{-1} \\ \hat{Z} &\equiv [h_{e_z^+}, \hat{Q}_v^{2k_5}] h_{e_z^+}^{-1}. \end{aligned} \quad (5.33)$$

Because of the holonomies in them, they are all operator-valued matrices whose entries are denoted as \hat{X}_{ab} , \hat{Y}_{ab} and \hat{Z}_{ab} respectively. With this notion, we have

$$\hat{F} = \hat{X}_{ab} \hat{Y}_{bc} \hat{Z}_{ca} - \hat{Y}_{ab} \hat{X}_{bc} \hat{Z}_{ca} = \hat{Y}_{bc} \hat{X}_{ab} \hat{Z}_{ca} - \hat{Y}_{ab} \hat{X}_{bc} \hat{Z}_{ca} + [\hat{X}_{ab}, \hat{Y}_{bc}] \hat{Z}_{ca}. \quad (5.34)$$

For the first two terms, because of Eq. (3.1), the expectation value of their subtraction can be simplified as

$$\langle \hat{Y}_{bc} \hat{X}_{ab} \hat{Z}_{ca} - \hat{Y}_{ab} \hat{X}_{bc} \hat{Z}_{ca} \rangle_{\{g_e\}} = (C_{\bar{a}\bar{b}\bar{c}\bar{d}\bar{e}\bar{f}} - C'_{\bar{a}\bar{b}\bar{c}\bar{d}\bar{e}\bar{f}}) \langle \hat{Y}'_{\bar{a}\bar{b}} \hat{X}'_{\bar{c}\bar{d}} \hat{Z}'_{\bar{e}\bar{f}} \rangle_{\{z_e\}} \quad (5.35)$$

where $C_{\bar{a}\bar{b}\bar{c}\bar{d}\bar{e}\bar{f}}$ and $C'_{\bar{a}\bar{b}\bar{c}\bar{d}\bar{e}\bar{f}}$ are two sets of constant coefficients. $(C_{\bar{a}\bar{b}\bar{c}\bar{d}\bar{e}\bar{f}} - C'_{\bar{a}\bar{b}\bar{c}\bar{d}\bar{e}\bar{f}})$ can be computed explicitly. \hat{X}' , \hat{Y}' and \hat{Z}' are the operators gauge transformed from X, Y, Z according to Eq. (3.1). For the last term in (5.34), since $[\hat{X}_{ab}, \hat{Y}_{bc}]$ itself is $O(t)$, we only need to compute the leading order expectation value of this term. Thus the commutator $[\hat{X}_{ab}, \hat{Y}_{bc}]$ can be easily computed with the results in Sec. 4.

Let us summarize the effects of symmetries. Let N_1 be the original number of terms in $\text{alt } \hat{H}_L^{(k)}$. We firstly use the symmetries implied by Eqs. (5.12) and (5.14) to reduce the number to $N_1/4^2$, where the power 2 is because there are two \hat{H}_E in $\text{alt } \hat{H}_L^{(k)}$. Eq. (5.16) further reduces this number to $N_1/(4^2 \times 48)$. Finally the symmetry (5.29) reduces the number to

$$\frac{N_1}{4^2 \times 48 \times 4} = \frac{N_1}{3072}.$$

Let N_2 be the original number of terms in $\text{extr } \hat{H}_L^{(k)}$. We firstly use the symmetries implied by Eqs. (5.12) and (5.14) to reduce the number to $N_2/4^2$. Then Eq. (5.16) reduces the number to $N_2/(4^2 \times 48)$. Eq. (5.34) further reduce this number to about ⁴

$$\frac{N_2}{4^2 \times 48 \times 2} = \frac{N_2}{1536}.$$

5.2 Introducing the algorithm: A case study

The procedure to compute the expectation value of the Hamiltonian operator can be illustrated by the following simple example. The complete implementation is presented in [37]. Let \hat{O} be the operator

$$\hat{O} = \frac{1}{t} [D_{ab}^l(h_{e_z^+}), \hat{Q}_v^m] D_{bc}^l(h_{e_z^+}^{-1}) \quad (5.36)$$

where $v = s(e_z^+)$. Then, in the case when we are only interested in $O(t)$ expansion, the expectation value of \hat{O} can be simplified by Eq. (5.11),

$$\begin{aligned} \langle \hat{O} \rangle_g &= \left(i \frac{(\beta a^2)^3}{8} \right) \sum_k \left(\prod_{i=1}^m \mathcal{T}_{\alpha_i \beta_i \gamma_i} \right) D_{ac}^l(\tau^{\gamma_k}) \left\langle \prod_{i=1}^m \hat{X}^{\alpha_i} \right\rangle_z \left\langle \prod_{i=1}^m \hat{Y}^{\beta_i} \right\rangle_z \left\langle \prod_{i \neq k} \hat{Z}^{\gamma_i} \right\rangle_z \\ &+ \left(i \frac{(\beta a^2)^3}{8} \right)^2 t \sum_{k < l} \left(\prod_{i=1}^m \mathcal{T}_{\alpha_i \beta_i \gamma_i} \right) D_{ac}^l(\tau^{\gamma_k} \tau^{\gamma_l}) \left\langle \prod_{i=1}^m \hat{X}^{\alpha_i} \right\rangle_z \left\langle \prod_{i=1}^m \hat{Y}^{\beta_i} \right\rangle_z \left\langle \prod_{i \neq k, l} \hat{Z}^{\gamma_i} \right\rangle_z \\ &=: O_0 + tO_1 \end{aligned} \quad (5.37)$$

where $\mathcal{T}_{\alpha\beta\gamma}$ is the contraction of $\epsilon_{\alpha'\beta'\gamma'}$ with the gauge transformation matrices according to Eq. (3.1).

Now let us show how to apply the results in Sec. 4 to reduce the computation complexity. Take O_0 as an example. The result of O_0 can be divided into two parts: the expectation-value parts

$$\left\langle \prod_{i=1}^m \hat{X}^{\alpha_i} \right\rangle_z \left\langle \prod_{i=1}^m \hat{Y}^{\beta_i} \right\rangle_z \left\langle \prod_{i \neq k} \hat{Z}^{\gamma_i} \right\rangle_z$$

and the factor parts

$$\left(\prod_{i=1}^m \mathcal{T}_{\alpha_i \beta_i \gamma_i} \right) D_{ac}^l(\tau^{\gamma_k}).$$

⁴The word ‘‘about’’ is because there exist the cases with $\hat{X} = \hat{Y}$ in Eq. (5.34).

For the expectation-value parts, without the results given in Sec. 4, we would have to consider 3^{3m-1} cases for various values of the indices α_i , β_i and γ_i . However, once the results in Sec. 4 are applied, since we only expand the expectation value to the next-to-leading order, we only have the following options

- (1) all indices are 0;
- (2) these indices contain a single +1 or a single -1;
- (3) these indices contain a pair of (+1, -1) or a pair (-1, +1).

Moreover, the results in Sec. 4 also tell us that for the last two options, positions of the ± 1 among these indices do not influence the expectation value, as far as the leading and next-to-leading orders are concerned. Therefore, we finally only need to compute $1 + 2 + 2 = 5$ cases. Comparing with the original number of cases 3^{3m-1} , we conclude that the computational complexity is reduced exponentially by the algorithm.

For the factor part, we only need to combine γ_k of $D_{ac}^l(\tau^{\gamma_k})$ with the indices in the expectation-value part. Then we consider all possibilities of permutations and all values of γ_k so that all possible values of the factor part can be computed. The result of O_0 can be obtained by summing over all possibilities.

Finally, let us complete this section with discussing the values of n and \vec{k} in $\hat{H}_E^{(n)}$ and $\hat{H}_L^{(\vec{k})}$. When we replace \hat{V}_v in \widehat{H}_E by $\hat{V}_{GT}^{(v)}$, a term $(\hat{Q}^2/\langle\hat{Q}\rangle^2 - 1)^k$ in $\hat{V}_{GT}^{(v)}$ has the following contribution to \widehat{H}_E

$$\begin{aligned} & \frac{1}{t} \text{tr}(h_{\alpha_{IJ}} h_{s_K} [(\hat{Q}^2/\langle\hat{Q}\rangle^2 - 1)^k, h_{s_K}^{-1}]) \\ &= \sum_{m=0}^{k-1} \text{tr}(h_{\alpha_{IJ}} h_{s_K} (\hat{Q}^2/\langle\hat{Q}\rangle^2 - 1)^m \frac{[\hat{Q}^2/\langle\hat{Q}\rangle^2, h_{s_K}^{-1}]}{t} (\hat{Q}^2/\langle\hat{Q}\rangle^2 - 1)^{k-1-m}) \end{aligned} \quad (5.38)$$

where the unimportant overall coefficient is neglected. Because $\hat{Q}^2/\langle\hat{Q}\rangle^2 - 1 = (\hat{Q}/\langle\hat{Q}\rangle + 1)(\hat{Q}/\langle\hat{Q}\rangle - 1)$ and the expectation value of $\hat{Q}/\langle\hat{Q}\rangle - 1$ is order $O(t)$, Eq. (4.6) tells us that the leading order of the operator in Eq. (5.38) is of $O(t^{\lfloor k/2 \rfloor})$. Thus as far as expanding the expectation value to $O(t)$ is considered, it is sufficient to choose $k \leq 3$ which means that the value of n in $\hat{H}_E^{(n)}$ is smaller than or equal to 3. A very similar discussion for $\hat{H}_L^{(\vec{k})}$ gives us that \vec{k} satisfies

$$\frac{|k_1 + k_2 - 3| + (k_1 + k_2 - 3)}{2} + \frac{|k_3 + k_4 - 3| + (k_3 + k_4 - 3)}{2} + k_5 \leq 3. \quad (5.39)$$

6 Quantum correction in the expectation value

We summarize the resulting expectation value of the Hamiltonian with unit lapse $\widehat{H}[1] = \widehat{H}_E + (1 + \beta^2) \widehat{H}_L$ at coherent states with cosmological data ($\eta < 0$ in our convention)

$$\begin{aligned} \langle \widehat{H}_E \rangle = & 6a\sqrt{-\beta\eta} \sin^2(\xi) - \frac{3}{4}at\sqrt{-\frac{\beta}{\eta^3}} \sin^2\left(\frac{\xi}{2}\right) \cos\left(\frac{\xi}{2}\right) \left\{ \cos\left(\frac{\xi}{2}\right) \left[8\eta^2 + 8\eta(4 \cosh(\eta) - 3) \operatorname{csch}(\eta) - 9 \right] \right. \\ & \left. - 12i\eta \sin\left(\frac{\xi}{2}\right) \right\} + O(t^2), \end{aligned} \quad (6.1)$$

$$\begin{aligned} \langle \widehat{H}_L \rangle = & -\frac{6a\sqrt{-\beta\eta} \sin^2(\xi) \cos^2(\xi)}{\beta^2} - \frac{3at}{262144(-\beta\eta)^{3/2}} \left\{ 2(3 - 220\eta^2) \cos(6\xi) \right. \\ & + 4i\eta(4838 \sin(\xi) - 6284 \sin(2\xi) + 4685 \sin(3\xi) - 5222 \sin(4\xi) - 105 \sin(5\xi)) \\ & + 2(-3611 + 8\eta(492\eta + 11i)) \cos(\xi) - 2(-789 + 4\eta(305\eta + 18i)) \cos(2\xi) \\ & + (4413 - 4\eta(928\eta + 49i)) \cos(3\xi) + 8(-1978 + \eta(4192\eta - 7i)) \cos(4\xi) \\ & + (-7 + 4(-272\eta + 5i)\eta) \cos(5\xi) - 4\eta \coth(\eta) \left[536 \cos(\xi) + 1731 \cos(2\xi) \right. \\ & \left. + 1524 \cos(3\xi) - 40548 \cos(4\xi) + 116 \cos(5\xi) + 117 \cos(6\xi) + 37292 \right] \\ & \left. + 8\eta \operatorname{csch}(\eta) \left[130 \cos(\xi) + 918 \cos(2\xi) + 801 \cos(3\xi) - 18618 \cos(4\xi) \right. \right. \\ & \left. \left. + 125 \cos(5\xi) + 58 \cos(6\xi) + 16362 \right] + 8(1436 + \eta(-4056\eta + 25i)) \right\} + O(t^2). \end{aligned} \quad (6.2)$$

By (5.22), $\langle \widehat{H}_L \rangle = \langle^{\text{extr}} \widehat{H}_L \rangle + \langle^{\text{alt}} \widehat{H}_L \rangle$ where $\langle^{\text{alt}} \widehat{H}_L \rangle$ is the expectation value of (5.21),

$$\begin{aligned} \langle^{\text{alt}} \widehat{H}_L \rangle = & -\frac{3a\sqrt{-\beta\eta} \sin^2(2\xi)}{8\beta^2} - \frac{3at}{32768(-\beta\eta)^{3/2}} \left\{ 4(104\eta^2 - 79) \cos(\xi) + (68 - 160\eta^2) \cos(2\xi) \right. \\ & + 4(55 - 104\eta^2) \cos(3\xi) + (944\eta^2 - 501) \cos(4\xi) - 848\eta^2 \\ & - 2\eta \left[\coth(\eta)(-262 \cos(\xi) - 6(46 \cos(2\xi) + 65 \cos(3\xi) - 380 \cos(4\xi) + 366)) \right. \\ & \left. - \operatorname{csch}(\eta)(-292 \cos(\xi) + 268 \cos(2\xi) + 420 \cos(3\xi) - 2035 \cos(4\xi) + 1799) \right. \\ & \left. \left. - 64i(6 \sin(\xi) - 10 \sin(2\xi) + 6 \sin(3\xi) - 7 \sin(4\xi)) \right] + 241 \right\} + O(t^2), \end{aligned} \quad (6.3)$$

$$\begin{aligned}
\langle^{\text{extr}}\widehat{H}_L\rangle = & -\frac{9a\sqrt{-\beta\eta}\sin^2(2\xi)}{8\beta^2} + \frac{3at}{262144(-\beta\eta)^{3/2}} \left\{ -2(580\eta^2 + 72i\eta - 517)\cos(2\xi) \right. \\
& + 2(3 - 220\eta^2)\cos(6\xi) + 4i\eta\left[3302\sin(\xi) - 3724\sin(2\xi) + 3149\sin(3\xi) \right. \\
& \left. \left. - 35(98\sin(4\xi) + 3\sin(5\xi))\right] + 2(-2347 + 8\eta(284\eta + 11i))\cos(\xi) \right. \\
& + (2653 - 4\eta(96\eta + 49i))\cos(3\xi) + 56(-211 + \eta(464\eta - i))\cos(4\xi) \\
& + (-7 + 4(-272\eta + 5i)\eta)\cos(5\xi) - 4\eta\coth(\eta)\left[1584\cos(\xi) + 627\cos(2\xi) \right. \\
& \left. - 36\cos(3\xi) - 31428\cos(4\xi) + 116\cos(5\xi) + 117\cos(6\xi) + 28508\right] \\
& + 8\eta\text{csch}(\eta)\left[714\cos(\xi) + 382\cos(2\xi) - 39\cos(3\xi) \right. \\
& \left. - 14548\cos(4\xi) + 125\cos(5\xi) + 58\cos(6\xi) + 12764\right] + 8\eta(-3208\eta + 25i) + 9560 \left. \right\} + O(t^2). \tag{6.4}
\end{aligned}$$

Note that the above results are written based on the graph with only a single vertex. Moreover, the results are complex because the operators \widehat{H}_E and \widehat{H}_L are not symmetry operators. Usually, we define the symmetrized operator as $\frac{1}{2}(\widehat{H}_E + \widehat{H}_E^\dagger)$ and $\frac{1}{2}(\widehat{H}_L + \widehat{H}_L^\dagger)$. Then the expectation values $\langle\frac{1}{2}(\widehat{H}_E + \widehat{H}_E^\dagger)\rangle$, and $\langle\frac{1}{2}(\widehat{H}_L + \widehat{H}_L^\dagger)\rangle$ are just the real parts of $\langle\widehat{H}_E\rangle$ and $\langle\widehat{H}_L\rangle$ respectively. Furthermore, when we express $\langle\widehat{H}[1]\rangle = H_0 + tH_1 + O(t^2)$, we notice that the $O(t)$ contribution H_1 contains η in the denominator, thus is divergent if $\eta \rightarrow 0$. This feature descends from the fact that $\widehat{V}_{GT}^{(v)}$ is divergent if $\langle\widehat{Q}_v\rangle \rightarrow 0$ because $\langle\widehat{Q}_v\rangle \sim |\eta|^3$. Thus our expansion of $\langle\widehat{H}[1]\rangle$ becomes invalid if η is too small. More precisely, when we express $\langle\widehat{H}[1]\rangle = \sqrt{|\eta|} [f_0 + (t/\eta^2)f_1 + O(t^2)]$, we have that f_0 is independent of η , and f_1 is regular at $\eta \rightarrow 0$, so we conclude that our expansion is valid when $\eta^2 \gg t$. Our expansion is valid for large $|\eta|$, because f_1 is regular at $|\eta| \rightarrow \infty$.

Even though there have been literature (e.g. [18, 26, 28]) considering the same issue and telling the leading order of the expectation value, our work gives the $O(\hbar)$ correction of the both Euclidean part and Lorentzian part. It should be noted that the $O(t)$ correction presented in the work is just the correction to the Hamiltonian operator. To get the total quantum correction of the whole mode, i.e. the total quantum correction of the effective action, one also needs to consider the 1-loop correction as in the usual quantum field theory.

Finally, we would like to compare our results with the results in loop quantum cosmology. In loop quantum cosmology, there are two types of dynamics, i.e., the μ_0 -scheme dynamics and the $\bar{\mu}$ -scheme dynamics [44, 45]. Since the μ_0 -scheme dynamics possesses some physically undesirable features, saying, that the bounce could occur in some classical regime. People prefer the $\bar{\mu}$ -scheme dynamics solving those problems in the μ_0 -scheme model. However, according to our results, our work can only recover the μ_0 scheme of loop quantum cosmology, and further adds some incomplete $O(t)$ corrections for cosmology. It is still open whether one can recover the $\bar{\mu}$ -scheme dynamics in the current framework or not once all quantum corrections are considered. Beyond the current framework, another attempt to recover the $\bar{\mu}$ -scheme cosmology is to considering the graph-changing Hamiltonian operator. For instance, in [46], by considering a model where the lattice changes in time, one gets results qualitatively similar to the $\bar{\mu}$ -scheme model.

7 Conclusion and outlook

In this paper, we develop an algorithm to overcome the complexity in the LQG Hamiltonian operator $\widehat{H[N]}$, and we compute explicitly the $O(\hbar)$ quantum correction in the expectation value $\langle \widehat{H[N]} \rangle$ at the coherent state peaked at the homogeneous and isotropic data of cosmology. With our algorithm and results, there are several perspectives which should be addressed in the future:

An important future task is to complete the computation of the quantum effective action Γ mentioned in Section 1. After the completion of the present work, the only missing ingredient in the $O(\hbar)$ terms in Γ is the “1-loop determinant” $\det(\mathfrak{H})$. Therefore a research to be carried out immediately is to compute $\det(\mathfrak{H})$ at the homogeneous and isotropic background. Once we obtain the complete $O(\hbar)$ contribution to Γ , the variation of Γ should give the quantum corrected effective equations which demonstrate the quantum correction to cosmology implied by LQG.

The next step of generalizing our computation is to study the expectation values $\langle \widehat{H[N]} \rangle$ at coherent states peaked at cosmological perturbations. The semiclassical limit of the expectation value and the cosmological perturbation theory from the path integral (1.2) are studied in [31]. Thus it is interesting to compute the $O(\ell_p)$ correction to the cosmological perturbation theory.

The computation of the quantum correction in $\langle \widehat{H[N]} \rangle$ should be extended to couplings of standard matter fields. The contributions from matter fields to $\widehat{H[N]}$ have been studied in the literature [47, 48]. The matter parts in $\widehat{H[N]}$ is much simpler than the Lorentzian part in $\widehat{H[N]}$, so computing their expectation values should not be hard. Studying the matter contributions and their quantum corrections is a project currently undergoing [49].

Acknowledgements

M.H. acknowledges Andrea Dapor, Klaus Liegener, and Hongguang Liu for various discussions motivating this work. M.H. receives support from the National Science Foundation through grant PHY-1912278. C. Z. acknowledges the support by the Polish Narodowe Centrum Nauki, Grant No. 2018/30/Q/ST2/00811

References

- [1] T. Thiemann, *Modern canonical quantum general relativity*, Cambridge University Press (2007).
- [2] C. Rovelli and F. Vidotto, *Covariant Loop Quantum Gravity: An Elementary Introduction to Quantum Gravity and Spinfoam Theory*, Cambridge Monographs on Mathematical Physics, Cambridge University Press (2014).
- [3] M. Han, Y. Ma and W. Huang, *Fundamental structure of loop quantum gravity*, *International Journal of Modern Physics D* **16** (2007) 1397.
- [4] A. Ashtekar and J. Lewandowski, *Background independent quantum gravity: a status report*, *Classical and Quantum Gravity* **21** (2004) R53.
- [5] K. Giesel and T. Thiemann, *Algebraic Quantum Gravity (AQG). II. Semiclassical Analysis*, *Class. Quant. Grav.* **24** (2007) 2499 [[gr-qc/0607100](#)].
- [6] K. Giesel and T. Thiemann, *Algebraic quantum gravity (AQG). III. Semiclassical perturbation theory*, *Class. Quant. Grav.* **24** (2007) 2565 [[gr-qc/0607101](#)].
- [7] M. Han and H. Liu, *Semiclassical limit of new path integral formulation from reduced phase space loop quantum gravity*, *Phys. Rev. D* **102** (2020) 024083 [[2005.00988](#)].

- [8] F. Conrady and L. Freidel, *On the semiclassical limit of 4d spin foam models*, *Phys. Rev. D* **78** (2008) 104023 [[0809.2280](#)].
- [9] J.W. Barrett, R. Dowdall, W.J. Fairbairn, F. Hellmann and R. Pereira, *Lorentzian spin foam amplitudes: Graphical calculus and asymptotics*, *Class.Quant.Grav.* **27** (2010) 165009 [[0907.2440](#)].
- [10] M. Han and M. Zhang, *Asymptotics of spinfoam amplitude on simplicial manifold: Lorentzian theory*, *Class.Quant.Grav.* **30** (2013) 165012 [[1109.0499](#)].
- [11] E. Bianchi, L. Modesto, C. Rovelli and S. Speziale, *Graviton propagator in loop quantum gravity*, *Class. Quant. Grav.* **23** (2006) 6989 [[gr-qc/0604044](#)].
- [12] M. Han, Z. Huang and A. Zipfel, *Emergent four-dimensional linearized gravity from a spin foam model*, *Phys. Rev. D* **100** (2019) 024060 [[1812.02110](#)].
- [13] M. Han, Z. Huang, H. Liu and D. Qu, *Numerical computations of next-to-leading order corrections in spinfoam large- j asymptotics*, *Phys. Rev. D* **102** (2020) 124010 [[2007.01998](#)].
- [14] P. Donà, M. Fanizza, G. Sarno and S. Speziale, *Numerical study of the Lorentzian Engle-Pereira-Rovelli-Livine spin foam amplitude*, *Phys. Rev. D* **100** (2019) 106003 [[1903.12624](#)].
- [15] J.F. Barbero G., *Real ashtekar variables for lorentzian signature space-times*, *Phys. Rev. D* **51** (1995) 5507.
- [16] H. Nicolai, K. Peeters and M. Zamaklar, *Loop quantum gravity: An Outside view*, *Class. Quant. Grav.* **22** (2005) R193 [[hep-th/0501114](#)].
- [17] J. Yang, Y. Ding and Y. Ma, *Alternative quantization of the hamiltonian in loop quantum cosmology*, *Physics Letters B* **682** (2009) 1.
- [18] A. Dapor and K. Liegener, *Cosmological Effective Hamiltonian from full Loop Quantum Gravity Dynamics*, *Phys. Lett. B* **785** (2018) 506 [[1706.09833](#)].
- [19] M. Han and H. Liu, *Effective dynamics from coherent state path integral of full loop quantum gravity*, *Physical Review D* **101** (2020) 046003.
- [20] W. Kamiński and K. Liegener, *Symmetry restriction and its application to gravity*, [2009.06311](#).
- [21] K. Giesel and T. Thiemann, *Algebraic quantum gravity (aqg): Iv. reduced phase space quantization of loop quantum gravity*, *Classical and Quantum Gravity* **27** (2010) 175009.
- [22] K. Giesel and T. Thiemann, *Scalar material reference systems and loop quantum gravity*, *Classical and Quantum Gravity* **32** (2015) 135015.
- [23] K.V. Kuchar and C.G. Torre, *Gaussian reference fluid and interpretation of quantum geometrodynamics*, *Phys. Rev.* **D43** (1991) 419.
- [24] T. Thiemann, *Quantum spin dynamics (qsd)*, *Classical and Quantum Gravity* **15** (1998) 839.
- [25] K. Giesel and T. Thiemann, *Algebraic Quantum Gravity (AQG). I. Conceptual Setup*, *Class. Quant. Grav.* **24** (2007) 2465 [[gr-qc/0607099](#)].
- [26] A. Dapor and K. Liegener, *Cosmological coherent state expectation values in loop quantum gravity I. Isotropic kinematics*, *Class. Quant. Grav.* **35** (2018) 135011 [[1710.04015](#)].
- [27] C. Zhang, J. Lewandowski, H. Li and Y. Ma, *Bouncing evolution in a model of loop quantum gravity*, *Physical Review D* **99** (2019) 124012.
- [28] K. Liegener and E.-A. Zwicknagel, *Expectation values of coherent states for $SU(2)$ Lattice Gauge Theories*, *JHEP* **02** (2020) 024 [[2001.00032](#)].

- [29] M. Han and H. Liu, *Improved $\bar{\mu}$ -scheme effective dynamics of full loop quantum gravity*, *Phys. Rev. D* **102** (2020) 064061 [[1912.08668](#)].
- [30] A. Dapor and K. Liegener, *Modifications to Gravitational Wave Equation from Canonical Quantum Gravity*, *Eur. Phys. J. C* **80** (2020) 741 [[2002.00834](#)].
- [31] M. Han, H. Li and H. Liu, *Manifestly gauge-invariant cosmological perturbation theory from full loop quantum gravity*, *Phys. Rev. D* **102** (2020) 124002 [[2005.00883](#)].
- [32] K. Liegener and L. Rudnicki, *Algorithmic approach to Cosmological Coherent State Expectation Values in LQG*, 12, 2020 [[2012.07813](#)].
- [33] M. Peskin and D. Schroeder, *An Introduction To Quantum Field Theory*, Frontiers in Physics, Avalon Publishing (1995).
- [34] A. Ashtekar and J. Lewandowski, *Quantum theory of geometry ii: Volume operators*, *Advances in Theoretical and Mathematical Physics* **1** (1997) 388.
- [35] C. Rovelli and L. Smolin, *Discreteness of area and volume in quantum gravity*, *Nuclear Physics B* **442** (1995) 593.
- [36] J. Yang and Y. Ma, *New volume and inverse volume operators for loop quantum gravity*, *Phys. Rev. D* **94** (2016) 044003.
- [37] C. Zhang, S. Song and M. Han, *First-order quantum correction in coherent state expectation value of loop-quantum-gravity hamiltonian*, *Physical Review D* **105** (2022) 064008.
- [38] https://github.com/czhangUW/LQG_expectation_values.
- [39] K. Giesel and T. Thiemann, *Algebraic quantum gravity (aqg): Iv. reduced phase space quantization of loop quantum gravity*, *Classical and Quantum Gravity* **27** (2010) 175009.
- [40] T. Thiemann, *Gauge field theory coherent states (gcs): I. general properties*, *Classical and Quantum Gravity* **18** (2001) 2025.
- [41] T. Thiemann and O. Winkler, *Gauge field theory coherent states (gcs): Iii. ehrenfest theorems*, *Classical and Quantum Gravity* **18** (2001) 4629.
- [42] T. Thiemann and O. Winkler, *Gauge field theory coherent states (gcs): Ii. peakedness properties*, *Classical and Quantum Gravity* **18** (2001) 2561.
- [43] J. Yang and Y. Ma, *Graphical calculus of volume, inverse volume and hamiltonian operators in loop quantum gravity*, *The European Physical Journal C* **77** (2017) 235.
- [44] A. Ashtekar, T. Pawłowski and P. Singh, *Quantum nature of the big bang: an analytical and numerical investigation*, *Physical Review D* **73** (2006) 124038.
- [45] A. Ashtekar, T. Pawłowski and P. Singh, *Quantum nature of the big bang: Improved dynamics*, *Phys. Rev. D* **74** (2006) 084003.
- [46] M. Han and H. Liu, *Loop quantum gravity on dynamical lattice and improved cosmological effective dynamics with inflaton*, *Physical Review D* **104** (2021) 024011.
- [47] T. Thiemann, *QSD 5: Quantum gravity as the natural regulator of matter quantum field theories*, *Class. Quant. Grav.* **15** (1998) 1281 [[gr-qc/9705019](#)].
- [48] H. Sahlmann and T. Thiemann, *Towards the QFT on curved space-time limit of QGR. 1. A General scheme*, *Class. Quant. Grav.* **23** (2006) 867 [[gr-qc/0207030](#)].
- [49] M. Han, C.-H. Hsiao, S. Song and C. Zhang, *Matter Couplings in Path Integral Formulation of Reduced Phase Space Loop Quantum Gravity*, [in preparation](#).