

AN EXPLICIT MINORANT FOR THE AMENABILITY CONSTANT OF THE FOURIER ALGEBRA

Y. CHOI

24th December 2021

Abstract

For a locally compact group G , let $\text{AM}(A(G))$ denote the amenability constant of the Fourier algebra of G . We show that $\text{AM}(A(G)) \geq 3/2$ for every non-abelian G , extending a result of Johnson (JLMS, 1994) who obtained this for finite non-abelian groups. This lower bound, which is known to be best possible, improves on results by previous authors and answers a question raised by Runde (PAMS, 2006).

To do this we study a minorant for $\text{AM}(A(G))$, related to the anti-diagonal in $G \times G$, which was implicitly used in Runde's work but hitherto not studied in depth. Our main novelty is an explicit formula for this minorant when G is a countable virtually abelian group, in terms of the Plancherel measure for G . As further applications, we characterize those non-abelian groups where the minorant attains its minimal value, and present some examples to support the conjecture that the minorant always coincides with the amenability constant.

MSC 2020: 46H20 (primary); 43A30, 46J10 (secondary)

1 Introduction

1.1 Background context

The Fourier algebra of a locally compact group G , denoted by $A(G)$, is a Banach algebra of functions on G whose norm encodes the group structure of G . By a celebrated theorem of Walter [Wal72], $A(G)$ is a complete invariant for G in the following sense: given two locally compact groups G_1 and G_2 the Banach algebras $A(G_1)$ and $A(G_2)$ are isometrically algebra-isomorphic if and only if G_1 and G_2 are topologically group-isomorphic. Thus $A(G)$ is a finer invariant than the commutative C^* -algebra $C_0(G)$, which only remembers the underlying topological space of G (and in particular cannot distinguish between discrete groups of the same cardinality).

It is therefore natural to study various invariants of Banach algebras, specialized to the setting of Fourier algebras, to see what information they remember about the group one starts with. Typically, algebraically motivated invariants of Banach algebras are *isomorphic* rather than *isometric* invariants, and hence cannot distinguish between groups whose Fourier algebras are isomorphic as topological algebras. For instance, if G_1 and G_2 are finite groups with the same cardinality n , then $A(G_1)$ and $A(G_2)$ are both (non-isometrically) isomorphic as unital Banach algebras to $(\mathbb{C}^n, \|\cdot\|_\infty)$. In particular, since the latter Banach algebra is known to be amenable, so are $A(G_1)$ and $A(G_2)$, and the notion of amenability cannot distinguish between $A(G_1)$ and $A(G_2)$ even though G_1 and G_2 may have very different group structure.

However, if A is an amenable Banach algebra one may consider its amenability constant $\text{AM}(A)$. This is an isometric invariant, and so has the potential to distinguish between two Fourier algebras that happen to be non-isometrically isomorphic; moreover, it is a numerical invariant, and hence lends itself to estimates and inequalities once one exploits its functorial properties. Even for finite groups, it displays interesting behaviour, as demonstrated by Johnson in his study [Joh94] of the (non-)amenability properties of Fourier algebras of *compact* groups.

Specifically, when G is a finite group, [Joh94, Theorem 4.1] provides an explicit and unexpectedly simple formula for $\text{AM}(A(G))$:

$$\text{AM}(A(G)) = \frac{1}{|G|} \sum_{\pi \in \widehat{G}} (d_\pi)^3 \quad (1.1)$$

where \widehat{G} denotes the set of (equivalence classes of) irreducible representations of G , and d_π denotes the dimension of the representation π . As we will see in Theorem 1.4, the right-hand side of (1.1) has independent meaning, which seems to have been overlooked in [Joh94] and the work of subsequent authors.

The formula (1.1) has several applications. For instance, it shows that $\text{AM}(A(G)) = 1$ for any *finite abelian* G . With the aid of some basic group theory it also implies the following “gap theorem” for the possible values of $\text{AM}(A(G))$ in the class of finite groups.

THEOREM 1.1 ([Joh94, Proposition 4.3]). *If G is a finite non-abelian group then $\text{AM}(A(G)) \geq 3/2$. Equality can be achieved (for instance, take G to be the dihedral group of order 8 or the quaternion group).*

REMARK 1.2. Combining Johnson’s formula with some basic properties of character tables, one can obtain a complete characterization of those finite non-abelian groups G where $\text{AM}(A(G))$ attains its minimum. Surprisingly this seems not to have been done before. Details of this character-theoretic argument will appear in future work, as part of a more systematic investigation of $\text{AM}(A(G))$ for finite groups.

Another consequence of (1.1), already observed by Johnson [Joh94, Corollary 4.2], is the following formula:

$$\text{AM}(A(G_1 \times G_2)) = \text{AM}(A(G_1)) \text{AM}(A(G_2)) \quad \text{for any finite groups } G_1 \text{ and } G_2. \quad (1.2)$$

This formula was used in conjunction with Theorem 1.1 to establish the non-amenability of the Fourier algebras of a large class of profinite groups (see [Joh94, Theorem 4.5]). Note that Equation (1.2) is not obvious from the general definition of the amenability constant of a Banach algebra! It is true that for arbitrary Banach algebras A and B one has $\text{AM}(A \widehat{\otimes} B) \leq \text{AM}(A) \text{AM}(B)$, where $\widehat{\otimes}$ denotes the projective tensor product; and one can show that equality holds if A and B are commutative, semisimple and finite-dimensional. However, this would only tell us about $A(G_1) \widehat{\otimes} A(G_2)$, which in general is not isometrically isomorphic to $A(G_1 \times G_2)$.

REMARK 1.3. There are difficulties in extending [Joh94, Theorem 4.1] to the non-compact setting. Johnson’s proof of the explicit formula (1.1) is indirect, since it proceeds by analyzing an auxiliary Banach algebra $A_\gamma(G)$ and showing that a particular idempotent in $A_\gamma(G)$ is mapped isometrically to the unique (virtual) diagonal for $A(G)$. Although the definition of $A_\gamma(G)$ can be extended to locally compact virtually abelian G , the relationship with $A(G)$ is less clear in non-compact cases. Moreover, when G is infinite, one no longer has a unique virtual diagonal for $A(G)$.

1.2 A minorant for the amenability constant

The study of (non-)amenability in [Joh94] relied heavily on the matrix-valued Fourier transform available for compact groups. Using a different perspective, Forrest and Runde proved that $A(G)$ is amenable if and only if G is *virtually abelian* [FR05, Theorem 2.3]. A key step in their proof is the following result: if $A(G)$ is amenable, then the indicator function of the set $\text{adiag}(G) := \{(x, x^{-1}) : x \in G\}$ belongs to the Fourier–Stieltjes algebra of $G_d \times G_d$. Here G_d denotes G equipped with the discrete topology.

In fact, as observed by Runde in a follow-up paper [Run06], the argument in [FR05] yields a more precise statement: $\text{AM}(A(G))$ is bounded below by the Fourier–Stieltjes norm of $1_{\text{adiag}(G)}$. This quantity, which we shall denote by $\text{AD}(G_d)$, is the “minorant” alluded to in the title; it can be defined without reference to amenability, and depends only on the underlying group of G rather than its topological group structure. As an invariant of G , it also enjoys slightly better functorial properties than $\text{AM}(A(\cdot))$. For instance, given two (discrete) virtually abelian groups Γ_1 and Γ_2 , one has

$$\text{AD}(\Gamma_1 \times \Gamma_2) = \text{AD}(\Gamma_1) \text{AD}(\Gamma_2)$$

(see Proposition 2.11 below), while for amenability constants of Fourier algebras this is only known in particular cases.

Besides its intrinsic interest, AD is a natural tool if one seeks to obtain *lower bounds* on $\text{AM}(A(G))$ for particular G , which would be hard to ascertain directly from the definition of the amenability constant. This was already noted in [Run06, Section 3], where after some partial results the following question is raised.

Runde’s question, paraphrased (see also [Jus19, Remark 4.9]).

- (Strong form.) Does the lower bound in Theorem 1.1 remain true for all locally compact non-abelian groups, not just the finite ones?
- (Weak form.) Can we find some $C > 1$ such that $\text{AM}(A(G)) \geq C$ for every locally compact non-abelian group G ?

Progress on these questions has been hampered by the lack of a formula such as (1.1) for infinite groups (see Remark 1.3). Nevertheless, the weak form of Runde’s question was answered positively by Forrest and Runde, by using results on norms of idempotent Schur multipliers to obtain lower bounds on $\text{AD}(G_d)$ (although they do not use this notation). In particular, it is observed in the remarks following [FR11, Theorem 3.4] that one can take $C = 2/\sqrt{3} \approx 1.1547$. An improved value of $C = 9/7 \approx 1.2857$ can be extracted¹ from results of Mudge and Pham [MP16]; strictly speaking, this is not stated explicitly in their paper, but it follows from the proof of [MP16, Theorem 2.2] and observations on the preceding page.

Despite this progress, the original stronger version of Runde’s question has remained unanswered until now. Naively, since $\text{AM}(A(G)) \geq \text{AM}(A(H))$ whenever H is a closed subgroup of G , one might hope to prove that $\text{AM}(A(G)) \geq 3/2$ by locating non-abelian finite subgroups of G and applying Theorem 1.1. However, as we shall see in Section 2.1, there are virtually abelian non-abelian groups in which every finite subgroup is abelian, and such groups cannot be handled with this strategy.

¹The author is grateful to N. Juselius for sharing a copy of [Jus19], which provides a useful synopsis of these developments. The results of Forrest–Runde and Mudge–Pham actually prove something sharper, concerning lower bounds on cb-multiplier norms of functions rather than their Fourier–Stieltjes norms, but we shall not discuss this extra refinement here.

1.3 Main results of this paper

In this paper we develop new machinery which yields, as a by-product, a positive answer to the strong form of Runde’s question. By [FR05, Theorem 2.3], it is enough to restrict our attention to virtually abelian groups, where more tools are available. Rather than using the perspective of operator theory, and trying to find improved estimates of Schur multiplier norms, we use the theory of the operator-valued Fourier transform. (In a sense this is more in the spirit of Johnson’s original paper [Joh94], although as indicated in Remark 1.3 we have to go beyond the techniques of [Joh94] once we leave the realm of compact groups.)

If one wishes to check how tight $\text{AD}(G_d)$ is as a lower bound for $\text{AM}(A(G))$, then a natural place to start is with finite groups where Johnson’s formula (1.1) is available. For finite G , the Fourier–Stieltjes algebra of $G_d \times G_d$ is just the Fourier algebra of $G \times G$, and one can calculate norms in this algebra in terms of the matrix-valued Fourier transform for $G \times G$. Working through the details, it turns out that the Fourier coefficients of $1_{\text{adiag}(G)}$ have a particular simple form, yielding a surprising result.

THEOREM 1.4 (Johnson’s formula, revisited). *Let G be a finite group. Then*

$$\text{AD}(G) = \frac{1}{|G|} \sum_{\pi \in \widehat{G}} (d_\pi)^3 = \text{AM}(A(G)). \quad (1.3)$$

Our result shows that [Joh94, Theorem 4.1] should not merely be regarded as a numerical coincidence, where $\text{AM}(A(G))$ happens to be equal to some “random number” defined in terms of \widehat{G} , but rather that an inequality relating two *a priori* different invariants for arbitrary G is actually an equality for all finite groups.

When Γ is infinite and virtually abelian, one still has a version of the Plancherel formula and hence a version of the Fourier inversion theorem. By using the explicit formula for the Fourier coefficients of $1_{\text{adiag}(G)}$ in the finite case, and making a suitable *Ansatz*, we obtain an explicit formula for $\text{AD}(\Gamma)$, at least when Γ is countable.

THEOREM 1.5 (The main formula). *Let Γ be countable and virtually abelian, with Plancherel measure ν normalized so that $\nu(\widehat{\Gamma}) = 1$. For each $n \in \mathbb{N}$, let $\Omega_n = \{\pi \in \widehat{\Gamma} : d_\pi = n\}$. Then*

$$\text{AD}(\Gamma) = \int_{\widehat{\Gamma}} (d_\pi)^2 d\nu(\pi) = \sum_{n \in \mathbb{N}} n^2 \nu(\Omega_n). \quad (1.4)$$

Although the new part of Theorem 1.4 is a special case of Theorem 1.5, it seems more natural to state the results separately and prove the special case first, since this approach provides crucial motivation for the key *Ansatz* in the proof of the general result. Moreover, the proof of the general result is technically more involved, relying on a non-abelian version of Bochner’s theorem that was developed in work of Arsac [Ars76, Section 3G].

Combining the formula (1.4) with an upper bound on $\nu(\Omega_1)$, and exploiting a “monotonicity principle” for AD , we are able to improve the existing lower bounds on $\text{AM}(A(G))$ for locally compact non-abelian groups, and show that the “strong form” of Runde’s question has a positive answer.

THEOREM 1.6 (Sharp lower bound). *If Γ is a (discrete) non-abelian group then $\text{AD}(\Gamma) \geq 3/2$. Consequently $\text{AM}(A(G)) \geq 3/2$ for every locally compact group non-abelian group G .*

With more work, we can go further and determine exactly which non-abelian groups attain the lower bound on $\text{AD}(\Gamma)$ in Theorem 1.6. Note that the next result does not require any countability assumptions.

THEOREM 1.7 (Non-abelian groups where AD attains its minimum). *Let Γ be a (discrete) group. Then $\text{AD}(\Gamma) = 3/2$ if and only if $|\Gamma : Z(\Gamma)| = 4$.*

REMARK 1.8. If $|\Gamma : Z(\Gamma)| = 4$ then basic group theory implies that $\Gamma/Z(\Gamma)$ is non-cyclic and hence is isomorphic to $C_2 \times C_2$ (see Lemma 4.5 below). Thus the groups described by Theorem 1.7 are exactly the non-trivial central extensions of $C_2 \times C_2$. In principle they can be classified by standard methods from group cohomology, although we have not found an explicit list in the literature.

REMARK 1.9. For a finite group G , we may combine Theorems 1.4 and 1.7 to obtain the following result, which has not appeared before in the literature: $\text{AM}(\text{A}(G)) = 3/2$ if and only if $|G : Z(G)| = 4$. However, this approach would be slightly misleading; as mentioned in Remark 1.2, the “correct” proof is much easier than that of Theorem 1.7, and proceeds by combining basic character theory with Johnson’s formula (1.3).

In view of Theorem 1.4, it is natural to ask if the inequality $\text{AD}(G_d) \leq \text{AM}(\text{A}(G))$ is always an equality. While we are unable to resolve this question at present, we can show that equality holds for certain infinite families of semidirect product groups (see also Remark 1.12).

To put these examples in context, we return briefly to the results of [Run06]. If G is a locally compact virtually abelian group, then $\text{maxdeg}(G) := \sup\{d_\pi : \pi \in \widehat{G}\}$ is finite. Runde showed that for such G we have a chain of inequalities

$$\text{AD}(G_d) \leq \text{AM}(\text{A}(G)) \leq \text{maxdeg}(G). \quad (1.5)$$

(The first inequality is proved in [Run06, Lemma 3.1] and the second inequality is proved in [Run06, Theorem 2.7].) We can now state our final main result.

THEOREM 1.10 (Sufficient conditions for equality to hold in (1.5)). *Let N be a torsion-free locally compact abelian group, and let p be a prime. Let α be a non-trivial action of the cyclic group C_p on N , and let $G = N \rtimes_\alpha C_p$. Then*

$$\text{AD}(G_d) = \text{AM}(\text{A}(G)) = \text{maxdeg}(G) = p. \quad (1.6)$$

These are the first recorded examples of groups not of the form (abelian) \times (finite), where the amenability constant of the Fourier algebra is finite and can be calculated explicitly.

REMARK 1.11.

- (i) The groups covered by Theorem 1.10 include examples which are discrete, (non-connected) Lie, or compact: let V be either \mathbb{Z} , \mathbb{R} or the 2-adic integers respectively, and then take $N = V^p$ equipped with the natural permutation action of C_p .
- (ii) If G is a finite non-abelian group, then it follows immediately from Johnson’s formula (1.1) that $\text{AM}(\text{A}(G)) < \text{maxdeg}(G)$. Hence the phenomenon in Theorem 1.10 is special to the setting of infinite G .

REMARK 1.12 (Crystal groups). The proof of Theorem 1.10 uses the fact that a suitably chosen countable subgroup $\Gamma \leq G$ has the following property: the set $\{\pi \in \widehat{\Gamma} : d_\pi < \text{maxdeg}(\Gamma)\}$ has Plancherel measure zero. In response to an earlier version of this paper, K. F. Taylor has pointed out (personal communication) that the same property holds for any crystal group, i.e. a discrete subgroup $\Gamma < \text{Isom}(\mathbb{R}^n)$ such that the orbit space \mathbb{R}^n/Γ is compact. Using this fact, it follows easily from Theorem 1.5 that for a crystal group Γ we also have $\text{AD}(\Gamma) = \text{AM}(\text{A}(\Gamma)) = \text{maxdeg}(\Gamma)$. Moreover, $\text{maxdeg}(\Gamma)$ is equal to the index in Γ of its translation subgroup (equivalently, equal to the order of the point group of Γ).

After some preliminaries in Section 2, the proofs of Theorems 1.4, 1.5, 1.6, 1.7 and 1.10 will be given in the following sections. We finish the paper with some questions and suggestions for future work.

2 Preliminaries

2.1 Group-theoretic background

A group G is said to be *virtually abelian* (VA for short) if it has an abelian subgroup of finite index. For convenience, we shall also use the abbreviation VANA, which stands for “virtually abelian non-abelian”.

Note that this property does not depend on any topological group structure that G may or may not possess. Note also that subgroups of VA groups are VA: if H has finite index in G and $\Gamma \leq G$ then $|\Gamma : H \cap \Gamma| \leq |G : H|$.

Clearly every finite group is virtually abelian, as is every abelian group. However the class of VA groups displays diverse behaviour, much more than the “trivial” examples of the form (abelian) \times (finite) might suggest. We present two examples, which are close to abelian in the sense that they have abelian subgroups of index 2, and in which every non-trivial finite subgroup has order 2 (so in particular, is abelian).

EXAMPLE 2.1 (Generalized dihedral groups). Let N be any torsion-free LCA group and let $G = N \rtimes C_2$, where C_2 acts on N by $x \mapsto -x$ (using additive notation). Regard N as a normal subgroup of G , via the canonical embedding $N \rightarrow N \rtimes C_2$. Then every non-identity element of N has infinite order in G , while every element of $G \setminus N$ is an involution; moreover, $xy \in N$ for all $x, y \in G \setminus N$. Therefore the only finite subgroups of G are the trivial subgroup $\{e\}$ and subgroups of the form $\{e, t\}$ for some $t \in G \setminus N$.

The following example seems to be well known to specialists in group theory, and can be found in [Fol16, Section 6.8]. It may be viewed as a discrete analogue of the “reduced Heisenberg group”.

EXAMPLE 2.2. Consider the group

$$\mathbb{H} = \left\{ \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} : a, b, c \in \mathbb{Z} \right\} \leq \mathrm{SL}_3(\mathbb{Z})$$

whose elements will be denoted by triples (a, b, c) for sake of conciseness. Define \mathbb{H}_2 to be the quotient of \mathbb{H} by the central subgroup $\{(0, 0, c) : c \in 2\mathbb{Z}\}$. We denote elements of \mathbb{H}_2 by triples $(a, b, [c])$ where $a, b \in \mathbb{Z}$ and $c \in \mathbb{Z}/2\mathbb{Z}$. There is a surjective homomorphism $q : \mathbb{H}_2 \rightarrow \mathbb{Z}^2$, given by $q(a, b, [c]) = (a, b)$, and since \mathbb{Z}^2 is torsion-free, the only finite-order elements in \mathbb{H}_2 are those in $\ker(q)$. Thus the only finite subgroups of \mathbb{H}_2 are the trivial subgroup and the 2-element subgroup $\ker(q)$. On the other hand, one can show by direct calculation that

$$N := \{(a, b, [c]) : a \in \mathbb{Z}, b \in 2\mathbb{Z}, [c] \in \mathbb{Z}/2\mathbb{Z}\}$$

is an abelian subgroup of \mathbb{H}_2 , and clearly $|\mathbb{H}_2 : N| = 2$.

In fact, for both of these examples, one can calculate the exact value of the anti-diagonal constant using the new results described in the introduction. The group G from Example 2.1 falls into the class of groups covered by Theorem 1.10, and therefore satisfies $\mathrm{AD}(G) = 2$.

The group \mathbb{H}_2 in Example 2.2 turns out to have centre $\{(a, b, [c]) : a \in 2\mathbb{Z}, b \in 2\mathbb{Z}, [c] \in \mathbb{Z}/2\mathbb{Z}\}$, and quotienting out by the centre gives a copy of $C_2 \times C_2$; so \mathbb{H}_2 falls into the class of groups covered by Theorem 1.7, and hence² $\text{AD}(\mathbb{H}_2) = 3/2$.

2.2 Fourier and Fourier–Stieltjes algebras

For the definition of the Fourier algebra $A(G)$ we refer the reader to [Eym64] or the recent text [KL18]. We shall not discuss amenability of Banach algebras in this paper, as we are only concerned with the minorant $\text{AD}(G_d)$ rather than $\text{AM}(A(G))$ itself. However, to study $\text{AD}(G_d)$ we must first discuss norms of elements inside Fourier–Stieltjes algebras of discrete groups. Since there are various equivalent definitions of the Fourier–Stieltjes algebra and equivalent definitions of its canonical norm, we review some terminology and notation.

An important convention throughout this paper is that the word “representation” *always means* “unitary representation”; this short-hand is merely to avoid unnecessary repetition.

DEFINITION 2.3. Let Γ be a discrete group. For a representation $\pi : \Gamma \rightarrow \mathcal{U}(H)$, a *coefficient function* of π is a function $\Gamma \rightarrow \mathbb{C}$ of the form $x \mapsto \langle \pi(x)\xi, \eta \rangle$ where $\xi, \eta \in H$. The *Fourier–Stieltjes algebra* of Γ is

$$B(\Gamma) := \bigcup_{\pi} \{\text{coefficient functions of } \pi\} \subseteq \ell^\infty(\Gamma).$$

where the union is taken over all representations of Γ . Clearly $B(\Gamma)$ is closed under scalar multiplication; by taking direct sums and tensor products of representations, we see that it is also closed under pointwise addition and multiplication.

Let Γ be a discrete group and let $b \in c_{00}(\Gamma)$. If $\theta : \Gamma \rightarrow \mathcal{U}(H)$ is a representation, we write $\theta(b)$ for the (finite) sum $\sum_{x \in \Gamma} b(x)\theta(x) \in \mathcal{B}(H)$. We then define $\|b\|_{C^*(\Gamma)} := \sup_{\theta} \|\theta(b)\|$ where the supremum is taken over all representations of Γ .

DEFINITION 2.4. For a discrete group Γ and $f \in B(\Gamma)$, we define

$$\|f\|_B := \sup \left\{ \sum_{x \in \Gamma} f(x)b(x) : b \in c_{00}(\Gamma), \|b\|_{C^*(\Gamma)} \leq 1 \right\} \quad (2.1)$$

Sometimes we shall write this as $\|f\|_{B(\Gamma)}$ to emphasize which group Γ is being considered.

The following result is standard but included here for convenient reference.

LEMMA 2.5. *Let $f \in B(\Gamma)$. If f can be expressed as a coefficient function $\langle \theta(\cdot)\xi, \eta \rangle$ for some representation $\theta : \Gamma \rightarrow \mathcal{U}(H)$ and vectors $\xi, \eta \in H$, then $\|f\|_B \leq \|\xi\|\|\eta\|$. Moreover, one can always find some choice of H , θ , ξ and η for which $f = \langle \theta(\cdot)\xi, \eta \rangle$ and $\|f\|_B = \|\xi\|\|\eta\|$.*

The first part of the lemma follows from Definition 2.4 and the Cauchy–Schwarz inequality. The second part follows by considering a suitable GNS representation and using polar decomposition; see e.g. [KL18, Lemma 2.1.9].

LEMMA 2.6. (a) *Let Λ be a discrete group and let $f \in B(\Lambda)$. If Λ_0 is a subgroup of Λ , then $f|_{\Lambda_0} \in B(\Lambda_0)$ and $\|f|_{\Lambda_0}\|_B \leq \|f\|_B$.*

²One can also give a more direct calculation of $\text{AD}(\mathbb{H}_2)$ using the main formula in Theorem 1.5, see Example 3.10 below.

(b) Let Λ_1 and Λ_2 be discrete groups, and let $f_i \in \mathcal{B}(\Lambda_i)$ for $i = 1, 2$. Define the function $f_1 \otimes f_2 : \Lambda_1 \times \Lambda_2 \rightarrow \mathbb{C}$ by $(f_1 \otimes f_2)(x, y) = f_1(x)f_2(y)$. Then $f_1 \otimes f_2 \in \mathcal{B}(\Lambda_1 \times \Lambda_2)$ and

$$\|f_1 \otimes f_2\|_{\mathcal{B}} = \|f_1\|_{\mathcal{B}}\|f_2\|_{\mathcal{B}}.$$

Proof. Part (a) is a straightforward consequence of Lemma 2.5, which we leave to the reader. The proof of part (b) requires more work. By Lemma 2.5, for $i = 1, 2$ we can write $f_i = \langle \theta_i(\cdot)\xi_i, \eta_i \rangle$ and $\|f_i\|_{\mathcal{B}} = \|\xi_i\|\|\eta_i\|$, for some representations $\theta_i : \Lambda_i \rightarrow \mathcal{U}(H_i)$ and vectors $\xi_i, \eta_i \in H_i$. Put $H = H_1 \otimes H_2$. Then $\theta_1 \otimes \theta_2 : \Lambda_1 \times \Lambda_2 \rightarrow \mathcal{U}(H)$ is a representation and

$$(f_1 \otimes f_2)(x, y) = \langle (\theta_1 \otimes \theta_2)(x, y)(\xi_1 \otimes \xi_2), \eta_1 \otimes \eta_2 \rangle.$$

Hence $f_1 \otimes f_2 \in \mathcal{B}(\Lambda_1 \times \Lambda_2)$ and

$$\|f_1 \otimes f_2\|_{\mathcal{B}} \leq \|\xi_1 \otimes \xi_2\|_H \|\eta_1 \otimes \eta_2\|_H \leq \|\xi_1\|\|\xi_2\|\|\eta_1\|\|\eta_2\| = \|f_1\|_{\mathcal{B}}\|f_2\|_{\mathcal{B}}.$$

For the converse inequality, observe that for any $b_1 \in c_{00}(\Lambda_1)$ and $b_2 \in c_{00}(\Lambda_2)$,

$$\begin{aligned} \|b_1 \otimes b_2\|_{C^*(\Lambda_1 \times \Lambda_2)} \|f_1 \otimes f_2\|_{\mathcal{B}} &\geq \left| \sum_{(x,y) \in \Lambda_1 \times \Lambda_2} (f_1 \otimes f_2)(x, y)(b_1 \otimes b_2)(x, y) \right| \\ &= \left| \sum_{x \in \Lambda_1} f_1(x)b_1(x) \sum_{y \in \Lambda_2} f_2(y)b_2(y) \right|, \end{aligned}$$

Therefore, if we can prove that

$$\|b\|_{C^*(\Lambda_1 \times \Lambda_2)} \leq \|b_1\|_{C^*(\Lambda_1)} \|b_2\|_{C^*(\Lambda_2)} \quad (*)$$

it will follow that $\|f_1 \otimes f_2\|_{\mathcal{B}} \geq \|f_1\|_{\mathcal{B}}\|f_2\|_{\mathcal{B}}$ as required.

Let $\theta : \Lambda_1 \times \Lambda_2 \rightarrow \mathcal{U}(H)$ be a representation and let $\theta_1(x) = \theta(x, e_2)$, $\theta_2(y) = \theta(e_1, y)$, where e_1 and e_2 denote the identity elements of Λ_1 and Λ_2 respectively. Then

$$\theta(b) = \sum_{(x,y) \in \Lambda_1 \times \Lambda_2} b_1(x)b_2(y)\theta(x, y) = \sum_{(x,y) \in \Lambda_1 \times \Lambda_2} b_1(x)b_2(y)\theta_1(x)\theta_2(y) = \theta_1(b_1)\theta_2(b_2)$$

so that $\|\theta(b)\| \leq \|b_1\|_{C^*(\Lambda_1)} \|b_2\|_{C^*(\Lambda_2)}$. Taking the supremum over all θ yields (*). \square

The following lemma is a special case of more general results, usually proved by considering extension problems for positive-definite functions. Given our earlier approach to defining $\|\cdot\|_{\mathcal{B}}$, we shall state a weaker result and provide a more direct proof.

LEMMA 2.7. *Let Λ be a discrete group and let $\Delta \leq \Lambda$ be a subgroup. Given $b \in c_{00}(\Delta)$, regard b as an element of $c_{00}(\Lambda)$ by setting $b(x) = 0$ for all $x \in \Lambda \setminus \Delta$. Then $\|b\|_{C^*(\Delta)} \leq \|b\|_{C^*(\Lambda)}$.*

Proof. Let $\pi : \Delta \rightarrow \mathcal{U}(H_\pi)$ be a representation, and form the induced representation $\Pi = \text{Ind}_\Delta^\Lambda \pi$. If we fix a transversal $T \subset \Lambda$ for the left coset space Λ/Δ , then Π can be realized on the Hilbert space

$$\mathcal{H}(\Lambda; \pi) := \left\{ F \in (H_\pi)^\Lambda : \sum_{x \in T} \|F(x)\|^2 < \infty, F(xt) = \pi(t^{-1})F(x) \forall x \in \Lambda \forall t \in \Delta \right\}$$

with $\Pi(x)F(y) := F(x^{-1}y)$. (See [KT13, Definition 2.1].)

We claim that $\Pi|_{\Delta}$ has a subrepresentation equivalent to π . Explicitly, defining $\phi : H_{\pi} \rightarrow \mathcal{H}(\Lambda; \pi)$ by

$$\phi(f)(x) = \begin{cases} \pi(x^{-1})f & \text{if } x \in \Delta, \\ 0 & \text{if } x \notin \Delta, \end{cases}$$

one can verify that ϕ is an isometry which intertwines π with Π . Hence $\pi(b) \in \mathcal{B}(H_{\pi})$ is unitarily equivalent to a compression of $\Pi(b) \in \mathcal{B}(\mathcal{H}(\Lambda; \pi))$, and so $\|\pi(b)\| \leq \|\Pi(b)\| \leq \|b\|_{C^*(\Lambda)}$. Taking the supremum over all π completes the proof. \square

LEMMA 2.8. *Let Λ be a discrete group and let $f \in B(\Lambda)$. Then there exists a countable subgroup $\Lambda_0 \leq \Lambda$ such that $\|f|_{\Lambda_0}\|_B = \|f\|_B$.*

This result is surely not new, but since we did not find an explicit reference in the literature, a proof is included here for sake of completeness.

Proof. Choose a sequence (b_n) in $c_{00}(\Lambda)$ such that $\|b_n\|_{C^*(\Lambda)} \leq 1$ and

$$\|f\|_B - \frac{1}{n} \leq \left| \sum_{x \in \Lambda} f(x)b_n(x) \right| \leq \|f\|_B.$$

For each n let $X_n \subset \Lambda$ be a finite subset such that $\text{supp}(b_n) \subseteq X_n$, and let Λ_0 be the subgroup of Λ generated by $\bigcup_{n \geq 1} X_n$, which is clearly countable. Each b_n may be viewed as an element of $c_{00}(\Lambda_0)$, and by Lemma 2.7 $\|b_n\|_{C^*(\Lambda_0)} \leq \|b_n\|_{C^*(\Lambda)} \leq 1$. Thus

$$\left| \sum_{x \in \Lambda} f(x)b_n(x) \right| = \left| \sum_{x \in \Lambda_0} f(x)b_n(x) \right| \leq \|f|_{\Lambda_0}\|_B.$$

Hence $\|f\|_B - \frac{1}{n} \leq \|f|_{\Lambda_0}\|_B \leq \|f\|_B$, and the result follows. \square

2.3 The anti-diagonal constant of a discrete group

We recall some definitions and notation that were already used in the introduction. Given a (locally compact) group G , the anti-diagonal of $G \times G$ is the set $\text{adiag}(G) = \{(x, x^{-1}) : x \in G\}$.

DEFINITION 2.9. Given a discrete group Γ we define the anti-diagonal constant of Γ to be

$$\text{AD}(\Gamma) = \|1_{\text{adiag}(\Gamma)}\|_{B(\Gamma \times \Gamma)}$$

with the convention that $\text{AD}(\Gamma) = \infty$ if $1_{\text{adiag}(\Gamma)} \notin B(\Gamma \times \Gamma)$.

As remarked in the introduction, Forrest and Runde showed that $\text{AD}(\Gamma) < \infty$ if and only if Γ is virtually abelian. We also note the following consequence of [Run06, Proposition 3.4]; we shall not need it for proving the main formula (Theorem 1.5) or the sharp lower bound (Theorem 1.6), but it will be convenient to use it in Section 4 when proving Theorem 1.7.

LEMMA 2.10 (Runde). *Let Γ be a group. Then $\text{AD}(\Gamma) = 1$ if and only if Γ is abelian.*

Both statements in the next proposition follow immediately from the corresponding parts of Lemma 2.6. We sometimes refer to the first statement as monotonicity of AD.

PROPOSITION 2.11. *If Γ is a discrete group and $\Gamma_0 \leq \Gamma$ then $\text{AD}(\Gamma_0) \leq \text{AD}(\Gamma)$. If Γ_1 and Γ_2 are discrete groups then $\text{AD}(\Gamma_1 \times \Gamma_2) = \text{AD}(\Gamma_1) \text{AD}(\Gamma_2)$*

For most of the paper, monotonicity of AD will be our main tool, since it allows us to get lower bounds on $\text{AD}(\Gamma)$ by choosing an appropriate countable subgroup of Γ . The next result is less essential, but has independent interest, and it will be convenient to make use of it in Section 4.

PROPOSITION 2.12 (Countable saturation for AD). *Every VA group Γ has a countable subgroup Γ_0 such that $\text{AD}(\Gamma) = \text{AD}(\Gamma_0)$.*

Proof. By Lemma 2.8 applied to $\text{B}(\Gamma \times \Gamma)$, there is a countable $\Lambda_0 \leq \Gamma \times \Gamma$ such that $\|\mathbf{1}_{\text{adiag}(\Gamma)}\|_{\text{B}} = \|\mathbf{1}_{\text{adiag}(\Gamma)}|_{\Lambda_0}\|_{\text{B}}$. Let Γ_0 be the subgroup of Γ generated by the subsets $\{x : (x, y) \in \Lambda_0\}$ and $\{y : (x, y) \in \Lambda_0\}$. Then Γ_0 is countable and $\Lambda_0 \leq \Gamma_0 \times \Gamma_0$, so that (by two applications of Lemma 2.6(a))

$$\|\mathbf{1}_{\text{adiag}(\Gamma)}\|_{\text{B}} \geq \|\mathbf{1}_{\text{adiag}(\Gamma_0)}\|_{\text{B}} \geq \|\mathbf{1}_{\text{adiag}(\Gamma)}|_{\Lambda_0}\|_{\text{B}}.$$

Hence $\|\mathbf{1}_{\text{adiag}(\Gamma)}\|_{\text{B}} = \|\mathbf{1}_{\text{adiag}(\Gamma_0)}\|_{\text{B}}$, as required. \square

2.4 Irreducible representations and the unitary dual

Since we only consider unitary representations in this paper, the notions of algebraic irreducibility and topological irreducibility coincide. We can therefore speak unambiguously of a representation being irreducible, and will abbreviate the phrase “irreducible representation” to “irrep”. Given a locally compact group G , we write \widehat{G} for the unitary dual of G : that is, the set of unitary equivalence classes of continuous irreps of G .

To simplify formulas and statements of results, for most of this paper we shall follow a standard abuse of notation, and identify \widehat{G} with a set of representative irreps of G : that is, we tacitly fix a function $z \mapsto \pi_z$ with the property that $[\pi_z] = z$. With this convention, already seen several times in Section 1, we refer to particular irreps as being elements of \widehat{G} .

In both the proof and the applications of the main formula (Equation (1.4)), we will freely use results and ideas from the theory of the non-commutative Fourier transform and Fourier inversion. We give an abridged review of the necessary notation and results from the literature, specialized to the setting of countable VA groups. For an overview of more general results, see [Fol16, Section 7.5].

Let Γ be a countable VA group. As remarked in the introduction, $\max\text{deg}(\Gamma) < \infty$. Moreover, since Γ is Type I and (2nd-)countable, the (Borel) σ -algebra on $\widehat{\Gamma}$ generated by the Fell topology is standard in the sense of measure theory.

Given $f \in c_{00}(\Gamma)$, note that when π is a finite-dimensional representation of Γ the scalar $\text{Tr}(\pi(f)\pi(f)^*)$ depends only on the unitary equivalence class of π . Hence each such f yields a well-defined function $\widehat{\Gamma} \rightarrow \mathbb{C}$, given by $[\pi] \mapsto \text{Tr}(\pi(f)\pi(f)^*)$, and one can show this function is continuous with respect to the Fell topology. Furthermore, there is a non-negative (Radon) measure ν on $\widehat{\Gamma}$, the Plancherel measure, such that

$$\|f\|_2^2 = \int_{\widehat{\Gamma}} \text{Tr}(\pi(f)\pi(f)^*) d\nu(\pi) \quad \text{for every } f \in c_{00}(\Gamma). \quad (2.2)$$

Taking $f = \delta_e$ we have $1 = \int_{\widehat{\Gamma}} d_\pi d\nu(\pi)$. (Note that we are tacitly adopting the normalization where Γ carries counting measure.) Moreover, by combining (2.2) with a polarization argument, we deduce that

$$\langle f, g \rangle_{\ell^2(\Gamma)} = \int_{\widehat{\Gamma}} \text{Tr}(\pi(f)\pi(g)^*) d\nu(\pi) \quad \text{for every } f, g \in c_{00}(\Gamma). \quad (2.3)$$

Note that in Equations (2.2) and (2.3) we are following the convention mentioned earlier, treating $\widehat{\Gamma}$ as a set of irreps by means of some tacit selection map. This practice will be more convenient when working with the main formula for $\text{AD}(\Gamma)$. However, in order to obtain this formula (i.e. in order to prove Theorem 1.5) we need to address measurability issues for certain operator fields defined on $\widehat{\Gamma}$ and $(\Gamma \times \Gamma)^\wedge$, and here it seems more appropriate to make explicit this selection of an irrep from each equivalence class.

We therefore finish this section with a précis of the required facts concerning measurable fields of Hilbert spaces, operators and representations. For all relevant definitions concerning these objects, the reader may consult the overview in [Fol16, Section 7.4], or the foundational presentation in [Dix81, Part II, Ch. 1–3]. The following terminology, however, is non-standard and made purely for convenience of later reference.

DEFINITION 2.13. Let G be a locally compact group and equip \widehat{G} with the σ -algebra generated by the Fell topology. A measurable selection for \widehat{G} is a pair $(\mathcal{H}_\bullet, \beta_\bullet)$, where:

- for each $z \in \widehat{G}$, β_z is a (continuous) representation of G acting on a Hilbert space \mathcal{H}_z , such that β_z belongs to the equivalence class z ;
- $z \mapsto \mathcal{H}_z$ is a measurable field of Hilbert spaces;
- for each $g \in G$, $z \mapsto \beta_z(g)$ is a measurable operator field with respect to \mathcal{H} .

It is not immediately obvious that we can always find measurable selections in the sense of the previous definition, even for countable VA groups. The following result will be sufficient for our needs: it is the specialization to the countable VA setting of results, due originally to Mackey, that hold for second-countable Type I groups.

LEMMA 2.14 (c.f. [Fol16, Lemma 7.31]). *Let Δ be a countable VA group, and equip $\widehat{\Delta}$ with the σ -algebra generated by the Fell topology. For each $z \in \widehat{\Delta}$, let d_z be the degree of any irrep belonging to the equivalence class z , and equip \mathbb{C}^{d_z} with its standard inner product. Then $(\mathbb{C}^{d_z})_{z \in \widehat{\Delta}}$ is a measurable field of Hilbert spaces on $\widehat{\Delta}$, and there is a measurable field of representations on $\widehat{\Delta}$, denoted by $z \mapsto \beta_z$, such that β_z acts on \mathbb{C}^{d_z} and $\pi_z \in z$ for each $z \in \widehat{\Delta}$. In particular, we have a measurable selection for Δ .*

(The field $(\mathbb{C}^{d_z})_{z \in \widehat{\Delta}}$ is often referred to as the canonical field of Hilbert spaces over $\widehat{\Delta}$.)

3 Fourier coefficients of the anti-diagonal

In this section we will prove Theorems 1.4, 1.5 and 1.6.

For a compact group K and $h \in A(K)$, the Fourier norm of h can be calculated from its matrix-valued Fourier coefficients. We only need the case when K is a finite group; with our earlier convention that $\sigma(h) = \sum_{x \in K} h(x)\sigma(x)$, we obtain

$$\|h\|_{A(K)} = \sum_{\sigma \in \widehat{K}} \frac{d_\sigma}{|K|} \|\sigma(h)\|_{(1)}. \quad (3.1)$$

In particular, for finite G we can calculate $\text{AD}(G)$ by working out the Fourier coefficients of $1_{\text{adiag}(G)}$, which turn out to have an explicit and tractable form if one exploits Schur orthogonality relations for G itself.

The calculation of the Fourier coefficients works just as well as for an infinite compact G , provided that one regards $1_{\text{adiag}(G)}$ as a singular measure supported on the anti-diagonal.

Given a Radon measure γ on a compact group K , we define its matrix-valued Fourier coefficients $(\pi(\gamma))_{\pi \in \widehat{K}}$ as integrals in the weak sense, i.e.

$$\langle \pi(\gamma)\xi, \eta \rangle := \int_K \langle \pi(x)\xi, \eta \rangle d\gamma(x) \quad (\xi, \eta \in H_\pi)$$

For K finite, we may view γ as a function on K , and then this definition of $\pi(\gamma)$ agrees with the earlier notation used for discrete groups.

Given a compact group G equipped with a choice of Haar measure μ , we introduce a Radon measure $\mu^{(\text{ad})}$ on $G \times G$, which is defined to be the pushforward of μ along the embedding $\text{adiag} : G \rightarrow G \times G$. Explicitly,

$$\int_{G \times G} f(x, y) d\mu^{(\text{ad})}(x, y) := \int_G f(x, x^{-1}) d\mu(x) \quad (f \in C(G \times G)). \quad (3.2)$$

Note that if G is finite and μ is counting measure, then the Radon–Nikodym derivative of $\mu^{(\text{ad})}$ with respect to counting measure on $G \times G$ is $1_{\text{adiag}(G)}$.

REMARK 3.1. If G is finite, but we chose to equip G and $G \times G$ with their uniform measures, as is more common when discussing Fourier analysis on compact groups, then the Radon–Nikodym derivative of $\mu^{(\text{ad})}$ with respect to uniform measure on $G \times G$ would be $|G|1_{\text{adiag}(G)}$.

DEFINITION 3.2. Let H be a Hilbert space. The flip map on $H \otimes_2 H$ is defined to be the unique bounded operator $\mathsf{X}_H : H \otimes_2 H \rightarrow H \otimes_2 H$ which satisfies $\mathsf{X}_H(\xi \otimes \xi') = \xi' \otimes \xi$ for every $\xi, \xi' \in H$.

PROPOSITION 3.3 (Fourier coefficients of the anti-diagonal). *Let G be compact and let $\mu, \mu^{(\text{ad})}$ be as above. Let $\pi, \sigma \in \widehat{G}$. Then*

$$(\pi \otimes \sigma)(\mu^{(\text{ad})}) = \begin{cases} 0 & \text{if } \pi \not\sim \sigma, \\ d_\pi^{-1} \mu(G) \mathsf{X}_\pi & \text{if } \pi = \sigma, \end{cases} \quad (3.3)$$

where by mild abuse of notation X_π denotes the flip map on H_π .

Proof. By linearity and continuity it suffices to check that the identities in (3.3) hold when both sides are evaluated on elementary tensors in $H_\pi \otimes H_\sigma$. Let $\xi, \eta \in H_\pi$ and $\xi', \eta' \in H_\sigma$. Then

$$\begin{aligned} \langle (\pi \otimes \sigma)(\mu^{(\text{ad})})(\xi \otimes \xi'), \eta \otimes \eta' \rangle &= \int_{G \times G} \langle \pi(x)\xi, \eta \rangle \langle \sigma(y)\xi', \eta' \rangle d\mu^{(\text{ad})}(x, y) \\ &= \int_G \langle \pi(x)\xi, \eta \rangle \langle \sigma(x^{-1})\xi', \eta' \rangle d\mu(x) \\ &= \int_G \langle \pi(x)\xi, \eta \rangle \overline{\langle \sigma(x)\eta', \xi' \rangle} d\mu(x) \end{aligned}$$

By the Schur orthogonality relations for G , the right-hand side vanishes if $\pi \not\sim \sigma$, while for $\pi = \sigma$ we have

$$\int_G \langle \pi(x)\xi, \eta \rangle \overline{\langle \pi(x)\eta', \xi' \rangle} d\mu(x) = \frac{\mu(G)}{d_\pi} \langle \xi', \eta \rangle \langle \xi, \eta' \rangle = \frac{\mu(G)}{d_\pi} \langle \mathsf{X}_\pi(\xi \otimes \xi'), \eta \otimes \eta' \rangle$$

as claimed. \square

REMARK 3.4. When $\dim(H) < \infty$ there is an explicit formula for X_H with respect to a choice of o.n. basis for H and the ensuing o.n. basis for $H \otimes_2 H$ (in fact, this was how the formulas in Proposition 3.3 were originally obtained). Let (E_{ij}) be the standard matrix units for $M_n(\mathbb{C})$ and consider $\sum_{i,j=1}^n E_{ij} \otimes E_{ji}$ viewed as an operator on $\mathbb{C}^n \otimes \mathbb{C}^n$. A direct calculation shows that this operator implements the flip map, and hence it equals $\mathsf{X}_{\mathbb{C}^n}$.

The proof of Theorem 1.4. It suffices to prove the first equality in (1.3), since the second one is Johnson’s formula (1.1). Let G be a finite group. Then $B(G \times G) = A(G \times G)$, so equipping $G \times G$ with counting measure and identifying $\mu^{(\text{ad})}$ with $1_{\text{adiag}(G)}$, we have

$$\begin{aligned} \text{AD}(G) &= \|1_{\text{adiag}(G)}\|_{A(G \times G)} = \sum_{\pi \otimes \rho \in (G \times G)^\wedge} \frac{d_{\pi \otimes \rho}}{|G \times G|} \|(\pi \otimes \rho)(1_{\text{adiag}(G)})\|_{(1)} \\ &= \sum_{\pi \in \widehat{G}} \frac{(d_\pi)^2 |G|}{|G|^2 d_\pi} \|\mathbf{X}_\pi\|_{(1)} \quad (\text{by Proposition 3.3}) \end{aligned}$$

To calculate the trace norm of \mathbf{X}_π , note that \mathbf{X}_π is a Hermitian involution on $H_\pi \otimes_2 H_\pi$, and so with respect to a suitable o.n. basis of $H_\pi \otimes_2 H_\pi$ it is represented by a diagonal matrix with ± 1 entries. Hence $\|\mathbf{X}_\pi\|_{(1)} = \dim(H_\pi \otimes_2 H_\pi) = (d_\pi)^2$, and the desired identity for $\text{AD}(G)$ follows. \square

While Proposition 3.3 does not apply to infinite discrete groups, it does suggest how we might proceed for groups which have a well-behaved notion of Fourier transform and Fourier inversion. However, since $1_{\text{adiag}(\Gamma)}$ does not belong to $c_{00}(\Gamma \times \Gamma)$, we cannot calculate its Fourier–Stieltjes norm by a direct application of the Plancherel theorem for $\Gamma \times \Gamma$. What is required is an isometric identification of $B(\Gamma \times \Gamma)$ with some space of “operator-valued measures” on $(\Gamma \times \Gamma)^\wedge$, in analogy with Bochner’s theorem which isometrically identifies the Fourier–Stieltjes algebra of a LCA group with the space of finite Radon measures on the dual group.

Such a result can be found in the PhD thesis of Arsac, specifically [Ars76, Section 3G]. The following is a special case of [Ars76, (3.55)].

THEOREM 3.5 (Arsac). *Let G be a second-countable Type I group, and fix a measurable selection $(\mathcal{H}_\bullet, \beta_\bullet)$ for \widehat{G} . If μ is a σ -finite Borel measure on \widehat{G} , and $T = (T_z)_{z \in \widehat{G}}$ is a μ -measurable operator field acting on \mathcal{H} , such that*

$$\|T\|_{1,\mu} := \int_{\widehat{G}} \|T_z\|_{(1)} d\mu(z) < \infty,$$

then the function $\Psi_\mu(T) : G \rightarrow \mathbb{C}$ defined by

$$\Psi_\mu(T)(s) := \int_{\widehat{G}} \text{Tr}(T_z \beta_z(s)) d\mu(z)$$

belongs to $B(G)$ and satisfies $\|\Psi(T)\|_B = \|T\|_{1,\mu}$.

We will apply this theorem with $G = \Gamma \times \Gamma$ for Γ countable and virtually abelian: the measure μ will be the push-forward of Plancherel measure on $\widehat{\Gamma}$ along the canonical map $\text{diag} : \widehat{\Gamma} \rightarrow (\Gamma \times \Gamma)^\wedge$, $[\pi] \mapsto [\pi \otimes \pi]$; and informally, the operator field T will be supported on $\text{diag}(\widehat{\Gamma})$, and its value on $[\pi \otimes \pi]$ will be the tensor flip map on an appropriate Hilbert space.

REMARK 3.6. Defining the push-forward presumes that diag is measurable with respect to the Borel σ -algebras on $\widehat{\Gamma}$ and $(\Gamma \times \Gamma)^\wedge$. Clearly this would follow from continuity with respect to the Fell topologies on both sides; while this seems to be a well-known fact, we are not aware of an explicit reference in the literature, and so a proof is included in the appendix. In fact, one can prove continuity of $\text{diag} : \widehat{G} \rightarrow (G \times G)^\wedge$ for a general locally compact group G , without any Type I assumptions (Theorem A.1); note that injectivity also holds in this more general setting, see e.g. [Fol16, Corollary 7.14].

We will also need the following property of the flip map, which we state as a separate lemma since it has independent interest.

LEMMA 3.7. *Let B and C be trace-class operators on a Hilbert space H . Then*

$$\mathrm{Tr}(\mathbf{X}_H \cdot (B \otimes C)) = \mathrm{Tr}(BC).$$

Proof. By linearity and continuity it suffices to prove this when B and C are rank-one operators on H , say $B(\xi) = \langle \xi, \beta_2 \rangle \beta_1$ and $C(\eta) = \langle \eta, \gamma_2 \rangle \gamma_1$ for some $\beta_1, \beta_2, \gamma_1, \gamma_2 \in H$. Then $BC(\eta) = \langle \eta, \gamma_2 \rangle \langle \gamma_1, \beta_2 \rangle \beta_1$, and so

$$\mathrm{Tr}(\mathbf{X}_H \cdot (B \otimes C)) = \langle \mathbf{X}_H(\beta_1 \otimes \gamma_1), \beta_2 \otimes \gamma_2 \rangle = \langle \gamma_1, \beta_2 \rangle \langle \beta_1, \gamma_2 \rangle = \mathrm{Tr}(BC)$$

as required. \square

(Note that if $\dim H < \infty$ we could give a simpler proof using the formula in Remark 3.4.)

The proof of the main formula (Theorem 1.5). Let $\Delta = \Gamma \times \Gamma$, and let $(\mathcal{H}_\bullet, \beta_\bullet)$ be a measurable selection for $\widehat{\Delta}$; such a selection exists by Lemma 2.14. Recall that $\mathrm{diag} : \widehat{\Gamma} \rightarrow \widehat{\Delta}$ denotes the function $[\pi] \rightarrow [\pi \otimes \pi]$. Informally, the idea is to construct an explicit operator field T over $\widehat{\Delta}$, and a Borel measure $\nu^{(d)}$ on $\widehat{\Delta}$, such that in the notation of Theorem 3.5 with $G = \widehat{\Delta}$ we obtain $\Psi_{\nu^{(d)}}(T) = 1_{\mathrm{ad}\mathrm{diag}(\Gamma)}$.

The measure $\nu^{(d)}$ is the easier part. As observed earlier in Remark 3.6, since $\mathrm{diag} : \widehat{\Gamma} \rightarrow \widehat{\Delta}$ is measurable we can define $\nu^{(d)}$ to be the pushforward measure $\mathrm{diag}_*(\nu)$; this is a finite Borel measure on $\widehat{\Delta}$ satisfying

$$\int_{\widehat{\Delta}} h(z) d\nu^{(d)}(z) = \int_{\widehat{\Gamma}} h(\mathrm{diag}(y)) d\nu(y) \quad \text{for every bounded measurable function } h.$$

To define the operator field T , we need to modify both \mathcal{H}_\bullet and β_\bullet on the subset $\mathrm{diag}(\widehat{\Gamma})$. Using Lemma 2.14, fix a measurable selection for $\widehat{\Gamma}$, say $(\mathcal{F}_\bullet, \alpha_\bullet)$. Then $\mathcal{F}^{(2)} := (\mathcal{F}_y \otimes_2 \mathcal{F}_y)_{y \in \widehat{\Gamma}}$ is also a measurable field of Hilbert spaces over $\widehat{\Gamma}$, and $(\alpha_y \otimes \alpha_y)_{y \in \widehat{\Gamma}}$ is a measurable field (over $\widehat{\Gamma}$) of representations of Δ acting on $\mathcal{F}^{(2)}$. Moreover, if we write \mathbf{X}_y for the tensor flip on $\mathcal{F}_y \otimes_2 \mathcal{F}_y$, then $(\mathbf{X}_y)_{y \in \widehat{\Gamma}}$ is a measurable operator field (over $\widehat{\Gamma}$), acting on $\mathcal{F}^{(2)}$.

Now we make the following definitions:

- $\widetilde{\mathcal{H}}$ is the field of Hilbert spaces defined by $\widetilde{\mathcal{H}}_z = \mathcal{F}_y \otimes \mathcal{F}_y$ if $z = \mathrm{diag}(y)$ and \mathcal{H}_z otherwise;
- $(\widetilde{\beta}_z)_{z \in \widehat{\Delta}}$ is the field of representations of Δ , acting on $\widetilde{\mathcal{H}}$, defined by $\widetilde{\beta}_z = \alpha_y \otimes \alpha_y$ if $z = \mathrm{diag}(y)$ and $\widetilde{\beta}_z = \beta_z$ otherwise;
- $T = (T_z)_{z \in \widehat{\Delta}}$ is the operator field defined by $T_z = \mathbf{X}_y$ if $z = \mathrm{diag}(y)$ and $T_z = 0$ otherwise.

We claim that these are all measurable fields, in the appropriate senses, over $\widehat{\Delta}$. For since Γ and Δ are 2nd-countable Type I groups, $\widehat{\Gamma}$ and $\widehat{\Delta}$ are standard measurable spaces; hence, by a result of Mackey (see e.g. [Dix77, B21]), every measurable injection $j : \widehat{\Gamma} \rightarrow \widehat{\Delta}$ has measurable image and its co-restriction $j : \widehat{\Gamma} \rightarrow j(\widehat{\Gamma})$ has measurable inverse. Applying this to $\mathrm{diag} : \widehat{\Gamma} \rightarrow \widehat{\Delta}$, we see that the fields $\widetilde{\mathcal{H}}$, $(\widetilde{\beta}_z)_{z \in \widehat{\Delta}}$ and T are obtained by gluing together measurable fields on $\mathrm{diag}(\widehat{\Gamma})$ with measurable fields on the complement $\widehat{\Delta} \setminus \mathrm{diag}(\widehat{\Gamma})$, and hence by routine calculations are themselves measurable, as claimed.

Thus $(\tilde{\mathcal{H}}_\bullet, \tilde{\beta}_\bullet)$ is also a measurable selection for $\widehat{\Delta}$, and T is a measurable operator field over $\widehat{\Delta}$ acting on $\tilde{\mathcal{H}}$. As seen earlier, for each $y \in \widehat{\Gamma}$ the trace norm of \mathbf{X}_y equals $(d_y)^2$, where d_y denotes the degree of any irrep in the equivalence class y . Since $\nu^{(d)}$ is a finite measure and $\max \deg(\Gamma) < \infty$, this shows that T is integrable in the sense required by Theorem 3.5 (we shall give a more precise calculation later). Now, given $s, t \in \Gamma$,

$$\begin{aligned}
\Psi_{\nu^{(d)}}(T)(s, t) &= \int_{\widehat{\Delta}} \text{Tr}(T_z \tilde{\beta}_z(s, t)) d\nu^{(d)}(z) && \text{(definition of } \Psi(T)) \\
&= \int_{\widehat{\Gamma}} \text{Tr}(T_{\text{diag}(y)} \tilde{\beta}_{\text{diag}(y)}) d\nu(\pi) && \text{(definition of } \nu^{(d)}) \\
&= \int_{\widehat{\Gamma}} \text{Tr}(\mathbf{X}_y \cdot (\alpha_y(s) \otimes \alpha_y(t))) d\nu(y) && \text{(definition of } T \text{ and } \tilde{\beta}_\bullet) \\
&= \int_{\widehat{\Gamma}} \text{Tr}(\alpha_y(s) \alpha_y(t)) d\nu(y) && \text{(Lemma 3.7)} \\
&= \langle \delta_s, \delta_{t^{-1}} \rangle && \text{(Equation (2.3)).}
\end{aligned}$$

Thus $\Psi(T) = 1_{\text{diag}(\Gamma)}$, and so by Theorem 3.5, $\text{AD}(\Gamma) = \|T\|_{1, \nu^{(d)}}$, which in turn is equal to

$$\int_{\widehat{\Delta}} \|T_z\|_{(1)} d\nu^{(d)}(z) = \int_{\widehat{\Gamma}} \|\mathbf{X}_y\|_{(1)} d\nu(y) = \int_{\widehat{\Gamma}} (d_y)^2 d\nu(y)$$

as required. \square

REMARK 3.8.

- (i) The modification procedure in the previous proof is somewhat cumbersome, and was only necessary because we chose to take Theorem 3.5 as a black box; informally, both the proof and statement of that theorem ignore the extra structure present when G has the form $\Gamma \times \Gamma$. One can give longer but more natural proofs of Theorem 1.5, by working through the arguments in [Ars76, Section 3G] and building in this extra structure.
- (ii) An alternative approach would be to start by showing that when Γ is countable and virtually abelian, the canonical injection $\widehat{\Gamma} \times \widehat{\Gamma} \rightarrow (\Gamma \times \Gamma)^\wedge$ is an isomorphism of measurable spaces. This seems to be folklore but, as with Remark 3.6, we are not aware of an explicit reference. Note that if we had followed this approach, the definition of $\nu^{(d)}$ and T becomes simpler, but we would still need to verify that the tensor square of a measurable selection for $\widehat{\Gamma}$ gives a measurable selection for $(\Gamma \times \Gamma)^\wedge$, in order to apply Theorem 3.5.

It is natural to ask if Theorem 1.5 can be extended to cover uncountable VA groups. The arguments in [Ars76, Section 3G] work by studying the coefficient space of the representation $\int_{\widehat{G}}^\oplus \sigma d\mu(\sigma)$, and decomposing this representation as a sum of direct integrals over constant fields of Hilbert spaces. Since our groups are virtually abelian, the fibres of these constant fields would be finite-dimensional Hilbert spaces with a uniform bound on the dimension, and it seems plausible that this would compensate for the fact that \widehat{G} could be non-standard as a measure space. There is no problem in defining the pushforward measure $\nu^{(d)} = \text{diag}_*(\nu)$: the Plancherel theorem can be extended to cover certain unimodular Type I groups which are not second-countable, in particular all discrete VA groups.

However, the proofs in [Ars76] rely on the disintegration results listed in [Dix77] (which in turn rely on foundational results for fields of von Neumann algebras, see e.g. [Dix81]). So one would need to work through the proofs of those results and check that the special cases

needed here can indeed be extended to non-standard measure spaces. All this would be very laborious, with limited rewards since Theorem 1.5 is most useful for countable groups anyway. We have therefore chosen to avoid such technicalities in this paper.

Let us now see how the main formula can be applied to prove the sharp lower bound (Theorem 1.6).

PROPOSITION 3.9. *Let Γ be a countable VANA group and let $\Omega_1 = \{\pi \in \widehat{\Gamma} : d_\pi = 1\}$. Let ν be Plancherel measure on $\widehat{\Gamma}$, normalized so that $\int_{\widehat{\Gamma}} d_\pi d\nu(\pi) = 1$. Then $\nu(\Omega_1) \leq 1/2$.*

Proof. Pick x and y in Γ with $xy \neq yx$, and note that whenever $\pi \in \Omega_1$, $\pi(xy) = \pi(yx)$ is a complex number of modulus 1. The Plancherel formula (Equation 2.3) then gives us

$$0 = \langle \delta_{xy}, \delta_{yx} \rangle_{\ell^2(\Gamma)} = \int_{\widehat{\Gamma}} \text{Tr}(\pi(xy)\pi(yx)^*) d\nu(\pi) = \nu(\Omega_1) + \int_{\widehat{\Gamma} \setminus \Omega_1} \text{Tr}(\pi(xy)\pi(yx)^*) d\nu(\pi)$$

and so

$$\begin{aligned} \nu(\Omega_1) &\leq \left| \int_{\widehat{\Gamma} \setminus \Omega_1} \text{Tr}(\pi(xy)\pi(yx)^*) d\nu(\pi) \right| \leq \int_{\widehat{\Gamma} \setminus \Omega_1} d_\pi d\nu(\pi) \\ &= \int_{\widehat{\Gamma}} d_\pi d\nu(\pi) - \int_{\Omega_1} d_\pi d\nu(\pi) \\ &= 1 - \nu(\Omega_1). \end{aligned}$$

Rearranging gives $2\nu(\Omega_1) \leq 1$ as required. \square

The proof of Theorem 1.6. The second statement follows from the first statement and the fact that $\text{AM}(A(G)) \geq \text{AD}(G_d)$ for any locally compact group. Hence it suffices to prove that $\text{AD}(\Gamma) \geq 3/2$ whenever Γ is non-abelian.

Since $\text{AD}(\Gamma) = +\infty$ if Γ is not virtually abelian, we may assume without loss of generality that Γ is VANA. Pick $x, y \in \Gamma$ which do not commute, and let Γ_0 be the subgroup of Γ generated by x and y . Then Γ_0 is countable and VANA. For each $n \in \mathbb{N}$, let $\Omega_n = \{\pi \in \widehat{\Gamma}_0 : d_\pi = n\}$. Then by Theorem 1.5

$$\text{AD}(\Gamma_0) - \nu(\Omega_1) = \sum_{n \geq 2} n^2 \nu(\Omega_n) \geq \sum_{n \geq 2} 2n \nu(\Omega_n) = 2(1 - \nu(\Omega_1)). \quad (3.4)$$

Rearranging and using Proposition 3.9 yields $\text{AD}(\Gamma_0) \geq 2 - 1/2 = 3/2$. By monotonicity of AD (Proposition 2.11) we deduce that $\text{AD}(\Gamma) \geq \text{AD}(\Gamma_0) \geq 3/2$, as required, \square

In the next section we will characterize those non-abelian groups for which our lower bound on AD is sharp. To motivate some of the techniques, we finish this section by returning to an earlier example and calculating its anti-diagonal constant.

EXAMPLE 3.10. Let \mathbb{H}_2 be the group from Example 2.2. Recall that \mathbb{H}_2 is non-abelian yet has no finite non-abelian subgroups; in particular it is not of the form (finite) \times (abelian). Despite this, we shall see that it attains the lower bound in Theorem 1.6.

The irreps of \mathbb{H}_2 can be found by a standard application of the Mackey machine, with details in e.g. [Fol16, Section 6.8]. We only require the following facts:

- every such representation is either 1-dimensional or 2-dimensional;
- if π is a 2-dimensional irrep of \mathbb{H}_2 , then $\pi(0, 0, [1]) = -I_2$.

Let ν be Plancherel measure on $\widehat{\mathbb{H}_2}$. The first property implies that $1 = \nu(\Omega_1) + 2\nu(\Omega_2)$. If we put $x = (1, 0, [0])$ and $y = (0, 1, [0])$, direct calculation shows that $[x, y] = (0, 0, [1])$. Arguing as in the proof of Proposition 3.9, we obtain

$$0 = \langle \delta_{xy}, \delta_{yx} \rangle = \int_{\pi \in \widehat{\mathbb{H}_2}} d_\pi \operatorname{Tr}(\pi([x, y])) d\nu(\pi) = \nu(\Omega_1) - 2\nu(\Omega_2).$$

Therefore $\nu(\Omega_1) = 1/2$ and $\nu(\Omega_2) = 1/4$, which yields $\operatorname{AD}(\mathbb{H}_2) = \nu(\Omega_1) + 4\nu(\Omega_2) = 3/2$.

4 Characterizing groups with anti-diagonal constant $3/2$

In this section we show that the groups Γ satisfying $\operatorname{AD}(\Gamma) = 3/2$ are exactly those satisfying $|\Gamma : Z(\Gamma)| = 4$ (Theorem 1.7). Note that we do not impose any countability restriction on Γ . However, since our “main formula” for the anti-diagonal constant is only established for countable VA groups, our approach inevitably passes through the countable setting.

Although the natural approach is to start with the condition $\operatorname{AD}(\Gamma) = 3/2$ and see what can be deduced, it is convenient to set up some general tools that can be used repeatedly.

DEFINITION 4.1. For a group Γ let $\operatorname{comm}(\Gamma) = \{[x, y] : x, y \in \Gamma\}$.

Note that $e \in \operatorname{comm}(\Gamma)$, and $|\operatorname{comm}(\Gamma)| = 1$ if and only if Γ is abelian. Thus the groups Γ for which $|\operatorname{comm}(\Gamma)| = 2$ are, informally, the non-abelian ones where there are as few commutators as possible. This has strong structural consequences; in particular, it follows from Lemma 4.2(i) below that such groups are nilpotent of class 2.

LEMMA 4.2. *Let Γ be a group satisfying $|\operatorname{comm}(\Gamma)| = 2$, and let z be the non-identity element of $\operatorname{comm}(\Gamma)$.*

- (i) $z^2 = e$ and $z = Z(\Gamma)$.
- (ii) *Let $\pi \in \widehat{\Gamma}$ with $d_\pi > 1$. Then $\pi(z) = -I_\pi$. Hence, if x and y are noncommuting elements of Γ , the operators $\pi(x)$ and $\pi(y)$ anti-commute, that is $\pi(x)\pi(y) + \pi(y)\pi(x) = 0$.*

Proof. If φ is an automorphism of Γ then it maps $\operatorname{comm}(\Gamma) \setminus \{e\}$ to itself. Thus z is fixed by every automorphism of Γ , in particular by all inner automorphisms, so $z \in Z(\Gamma)$. Since $([a, b])^{-1} = [b, a]$ for every $a, b \in \Gamma$, we have $z^{-1} \in \operatorname{comm}(\Gamma) \setminus \{e\} = \{z\}$. That is, $z^{-1} = z$. This proves part (i).

Let $\pi \in \widehat{\Gamma}$ with $d_\pi > 1$. Since $z \in Z(\Gamma)$ and π is irreducible, $\pi(z) \in \mathbb{C}I_\pi$ by Schur’s lemma. Moreover, $\pi(z) = \pm I_\pi$ since $z^2 = e$. Suppose $\pi(z) = I_\pi$: then since $\operatorname{comm}(\Gamma) = \{e, z\}$ we have $[\pi(a), \pi(b)] = \pi([a, b]) = I_\pi$ for all $a, b \in \Gamma$. But this implies that $\pi(\Gamma)$ is abelian which contradicts irreducibility of π . Therefore $\pi(z) = -I_\pi$, and since $\pi(z) = \pi(x)\pi(y)\pi(x)^{-1}\pi(y)^{-1}$, rearranging gives $\pi(x)\pi(y) = -\pi(y)\pi(x)$. This proves part (ii). \square

The relevance of the condition $|\operatorname{comm}(\Gamma)| = 2$ to the study of $\operatorname{AD}(\Gamma)$ is not immediately apparent, but is partially explained by the next two results. In both results we restrict attention to countable VA groups, to make use of the Plancherel identity (2.3).

PROPOSITION 4.3. *Let Λ be a countable VA group with Plancherel measure ν . Let $\Omega_1 = \{\pi \in \widehat{\Lambda} : d_\pi = 1\}$. Then $|\operatorname{comm}(\Lambda)| = 2$ if and only if $\nu(\Omega_1) = 1/2$.*

Proof. We may assume that Λ is non-abelian (since this follows from either of the assumptions $|\text{comm}(\Lambda)| = 2$ or $\nu(\Omega_1) = 1/2$).

Note that $\text{Re Tr}(V) \leq d_\pi$ for any unitary $V \in \mathcal{U}(H_\pi)$, with equality holding if and only if $V = I_\pi$. Hence, as already observed in the proof of Proposition 3.9:

$$\begin{aligned} \text{for all } z \in \text{comm}(\Lambda) \setminus \{e\}, \quad \nu(\Omega_1) &= \int_{\widehat{\Lambda} \setminus \Omega_1} -\text{Tr } \pi(z) d\nu(\pi) \\ &= \int_{\widehat{\Lambda} \setminus \Omega_1} \text{Re Tr}(-\pi(z)) d\nu(\pi) \\ &\leq \int_{\widehat{\Lambda} \setminus \Omega_1} d_\pi d\nu(\pi) = 1 - \nu(\Omega_1), \end{aligned}$$

and equality holds if and only if $\pi(z) = -I_\pi$ for ν -a.e. $\pi \in \widehat{\Lambda} \setminus \Omega_1$.

Note that $\nu(\Omega_1) = 1/2$ if and only if $\nu(\Omega_1) = 1 - \nu(\Omega_1)$. By the previous remarks, the following statements are therefore equivalent:

- (i) $\nu(\Omega_1) = 1/2$;
- (ii) $\pi(z) = -I_\pi$ for every $z \in \text{comm}(\Lambda) \setminus \{e\}$ and ν -a.e. $\pi \in \widehat{\Lambda} \setminus \Omega_1$.

If $|\text{comm}(\Lambda)| = 2$, then (ii) holds by Lemma 4.2(ii). On the other hand, suppose that (ii) holds. Let $z, z' \in \text{comm } \Lambda \setminus \{e\}$. By assumption $\pi(z) = \pi(z')$ for ν -a.e. $\pi \in \widehat{\Lambda} \setminus \Omega_1$; and for every $\pi \in \Omega_1$ we have $\pi(z) = \pi(z')$. By injectivity of the non-abelian Fourier transform (or, alternatively, by applying the Plancherel identity (2.3) to $\langle \delta_z, \delta_{z'} \rangle$) it follows that $z = z'$. Thus $|\text{comm}(\Lambda)| = 2$. \square

LEMMA 4.4. *Let Λ be a countable VA group and let $\Omega_n = \{\pi \in \widehat{\Lambda} : d_\pi = n\}$. Let ν denote Plancherel measure on $\widehat{\Lambda}$. The following are equivalent:*

- (i) $\text{AD}(\Lambda) = 3/2$;
- (ii) $\nu(\Omega_1) = 1/2$ and $\nu(\Omega_n) = 0$ for all $n \geq 3$.

Proof. We already observed in the proof of Theorem 1.6 — specifically, in the inequality (3.4) — that $\text{AD}(\Lambda) \geq 2 - \nu(\Omega_1)$, with equality if and only if $n^2\nu(\Omega_n) = 2n\nu(\Omega_n)$ for all $n \geq 2$. In turn, this is equivalent to requiring $\nu(\Omega_n) = 0$ for all $n \geq 3$.

Now, Proposition 3.9 implies that $2 - \nu(\Omega_1) \geq 3/2$ with equality if and only if $\nu(\Omega_1) = 1/2$. Thus (i) and (ii) are equivalent. \square

Let us sketch where we are headed. Lemma 4.4 and Proposition 4.3 indicate that, at least for countable groups, the assumption $\text{AD}(\Lambda) = 3/2$ places strong restrictions on both the set of commutators in Λ and the set of possible degrees of irreps of Λ . These restrictions turn out to imply that for any non-central $x \in \Lambda$, the centralizer of x has index 2 in Λ but also contains $Z(\Lambda)$ as a subgroup of index 2; this will immediately imply that $|\Lambda : Z(\Lambda)| = 4$. In the converse direction, if Γ is any group (not necessarily countable), assuming that $|\Gamma : Z(\Gamma)| = 4$ turns out to have strong structural consequences for Γ , which can be used to control both $\text{comm}(\Gamma)$ and the degrees of irreps of Γ .

For technical reasons, let us switch attention from the assumption $\text{AD}(\Gamma) = 3/2$ and its consequences to the assumption $|\Gamma : Z(\Gamma)| = 4$ and its consequences. The following basic group-theoretic lemma can be found as a standard exercise in various texts.

LEMMA 4.5. *Let Δ be a non-abelian group. Then $\Delta/Z(\Delta)$ is not cyclic. In particular $|\Delta:Z(\Delta)| \geq 4$.*

Outline of a proof. The first part follows from a stronger statement: if Δ is a group and N is a normal subgroup contained in $Z(\Delta)$ such that Δ/N is cyclic, then Δ is abelian. To prove this, lift the generator of Δ/N to some $r \in \Delta$, and note that $\Delta = \bigcup_{n \geq 1} r^n N \subseteq \bigcup_{n \geq 1} r^n Z(\Delta)$. For the second part, it suffices to note that all groups of order ≤ 3 are cyclic. \square

The next result characterizes the groups in which the centre has index 4, as those non-abelian groups where the set of commutators and the set of degrees of irreps are as small as possible. We have not found it stated in the literature, so we provide a detailed proof, since it may have interest independent of the application to anti-diagonal constants. Note that we do not require any countability assumptions.

THEOREM 4.6. *Let Γ be a group. The following conditions are equivalent.*

- (i) $|\Gamma:Z(\Gamma)| = 4$.
- (ii) $|\text{comm}(\Gamma)| = 2$ and $\text{maxdeg}(\Gamma) = 2$.
- (iii) $|\text{comm}(\Gamma)| = 2$ and $\{\pi \in \widehat{\Gamma}: d_\pi \leq 2\}$ separates the points of Γ .

REMARK 4.7. The equivalence of parts (ii) and (iii) is a special case of general results for the maxdeg function, which follow in turn from results on polynomial identities for noncommutative algebras. However, this does not seem to significantly shorten the proof of the equivalence (iii) \implies (i), which is what we need in the application to $\text{AD}(\Gamma)$.

Proof of Theorem 4.6(i) \implies (ii). Note that the commutator map $\Gamma \times \Gamma \rightarrow \Gamma$ factors through $(\Gamma/Z(\Gamma)) \times (\Gamma/Z(\Gamma))$. More precisely, let $q: \Gamma \rightarrow \Gamma/Z(\Gamma)$ be the quotient homomorphism, and define an equivalence relation \sim on Γ by $x \sim y \iff q(x) = q(y)$. Given $x \sim x'$ and $y \sim y'$, a direct calculation shows that $[x, y] = [x', y']$. Therefore, if T is a transversal for $Z(\Gamma)$ in Γ , we have $\text{comm}(\Gamma) = \{[x, y]: x, y \in T\}$.

Since $\Gamma/Z(\Gamma)$ is non-cyclic (Lemma 4.5) and has order 4, it must be isomorphic to the Klein-four group, and hence has the the following properties:

- each non-identity element has order 2;
- any two elements commute with each other;
- multiplying two distinct non-identity elements gives the third non-identity element.

Choose a transversal $\{e, a, b, c\}$ for $Z(\Gamma)$ in Γ . Then the properties listed above imply:

$$a \sim a^{-1}, b \sim b^{-1}, c \sim c^{-1} \quad \text{and} \quad ab \sim c \sim ba, bc \sim a \sim cb, ca \sim b \sim ac.$$

Since $a \sim a^{-1}$ and $b \sim ac$, we have

$$[a, b] = [a^{-1}, ac] = a^{-1}(ac)a(ac)^{-1} = [c, a];$$

and since $a \sim cb \sim cb^{-1}$, we also have

$$[a, b] = [cb^{-1}, b] = (cb^{-1})b(bc^{-1})b^{-1} = [c, b].$$

Thus $[a, b] = [c, a] = [c, b]$. By symmetry, these identities remain valid under any permutation of the symbols a, b, c , so we also obtain $[b, c] = [a, b] = [a, c]$ and $[c, a] = [b, c] = [b, a]$. We

have shown that $[a, b], [b, c], [c, a], [b, a], [c, b], [a, c]$ are all equal; denoting their common value by z_0 , it follows from our previous remarks that

$$\text{comm}(\Gamma) = \{[x, y] : x, y \in T\} = \{e, z_0\}.$$

Thus $|\text{comm}(\Gamma)| \leq 2$, and equality must hold since Γ is non-abelian.

Now let $\pi \in \widehat{\Gamma}$ with $d_\pi \geq 2$; we must show that H_π is 2-dimensional. Pick x and y in Γ which do not commute. Their images in $\Gamma/Z(\Gamma)$ are distinct, and so they generate all of $\Gamma/Z(\Gamma)$, by the properties of the Klein-four group mentioned above. Hence Γ is generated by $\{x, y\} \cup Z(\Gamma)$. Moreover, $\pi(Z(\Gamma)) \subseteq \mathbb{C}I_\pi$ (by Schur's lemma), and so $\pi(\Gamma)$ is generated by $\{\pi(x), \pi(y)\} \cup \mathbb{C}I_\pi$. Therefore, if we can produce a 2-dimensional subspace V which is $\pi(x)$ -invariant and $\pi(y)$ -invariant, this will force $V = H_\pi$ by irreducibility.

Let v_1 be an eigenvector of $\pi(y)$, with eigenvalue λ , and let $v_2 = \pi(x)v_1$. Since $\Gamma/Z(\Gamma)$ is commutative, $[x, y]$ is central in Γ , and so by Schur's lemma $\pi([x, y]) = \mu I_\pi$ for some³ $\mu \in \mathbb{C}$. Thus $\pi(x)\pi(y) = \mu\pi(y)\pi(x)$, which implies $\pi(y)v_2 = \lambda\mu v_1$. Also, since every element of $\Gamma/Z(\Gamma)$ squares to the identity, $x^2 \in Z(\Gamma)$ and once again Schur's lemma yields $\pi(x)^2 \in \mathbb{C}I_\pi$. Hence $\pi(x)v_2 = \pi(x^2)v_1 \in \mathbb{C}v_1$. Thus $V := \text{lin}\{v_1, v_2\}$ is an invariant subspace for $\pi(x)$ and $\pi(y)$, as required. \square

Proof of Theorem 4.6(ii) \implies (iii). This follows immediately from the fact that $\widehat{\Gamma}$ separates the points of Γ (note that this does not require any form of the Plancherel theorem). \square

To complete the circle of implications in Theorem 4.6, we need a closer look at $\pi(\Gamma)$ when $\pi \in \widehat{\Gamma}$ with $d_\pi = 2$. For clarity, we isolate some steps in a preliminary lemma.

LEMMA 4.8. *Let Γ be a group with $|\text{comm}(\Gamma)| = 2$ and let $\pi \in \widehat{\Gamma}$ with $d_\pi = 2$. Let $x \in \Gamma \setminus Z(\Gamma)$.*

- (i) *If $c \in Z_\Gamma(x) \setminus Z(\Gamma)$, then $\pi(c)$ is a scalar multiple of $\pi(x)$.*
- (ii) *If $a, b \in \Gamma \setminus Z_\Gamma(x)$, then $\pi(a^{-1}b)$ commutes with $\pi(x)$.*

Proof. We start by diagonalizing the matrix $\pi(x)$. Since $\pi(x)$ is unitary, its eigenvalues lie on the unit circle. On the other hand, pick $y \in \Gamma$ with $xy \neq yx$. Then by Lemma 4.2(ii), $\pi(y^{-1})\pi(x)\pi(y) = -\pi(x)$; taking traces on both sides and rearranging yields $\text{Tr } \pi(x) = 0$. Therefore $\pi(x)$ has distinct eigenvalues λ and $-\lambda$ for some $\lambda \in \mathbb{T}$, and by taking corresponding eigenvectors v_+ and v_- we obtain a basis for H_π . If $c \in Z_\Gamma(x) \setminus Z(\Gamma)$, then since $\pi(c)$ commutes with $\pi(x)$ it must be diagonal with respect to the basis $\{v_+, v_-\}$. But since c is non-central, repeating the argument used for $\pi(x)$ shows that the eigenvalues of $\pi(c)$ lie on the unit circle and sum to zero. Thus $\pi(c)v_\pm = \pm\mu v_\pm$ for some $\mu \in \mathbb{T}$, so that $\pi(c) = \lambda^{-1}\mu\pi(x)$, and we have proved part (i).

For part (ii), it is convenient to introduce the matrix $J = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, so that $\pi(x)$ is represented by the matrix λJ with respect to the basis $\{v_+, v_-\}$. Now let $a, b \in \Gamma$ with $ax \neq xa$ and $bx \neq xb$. By Lemma 4.2(ii), the operators $\pi(a)$ and $\pi(x)$ *anti-commute*. A direct calculation shows that an element of $M_2(\mathbb{C})$ which anti-commutes with J must have zero entries on the diagonal, and so $\pi(a)$ swaps the two subspaces $\mathbb{C}v_+$ and $\mathbb{C}v_-$. The same is true for $\pi(b)$, and so $\pi(a^{-1}b) = \pi(a)^{-1}\pi(b)$ preserves each of these subspaces, i.e. it is diagonal with respect to the basis $\{v_+, v_-\}$. Hence $\pi(a^{-1}b)$ commutes with $\pi(x)$, and part (ii) is proved. \square

³In fact, it follows from the proof of Lemma 4.2 that $\mu = -1$, but this is more than we need.

Proof of Theorem 4.6(iii) \implies (i). Pick a non-central $x \in \Gamma$. We will show that $|Z_\Gamma(x) : Z(\Gamma)| = 2$ and $|\Gamma : Z_\Gamma(x)| = 2$, which together imply that $|\Gamma : Z(\Gamma)| = 4$.

For convenience let $\Omega_n = \{\pi \in \widehat{\Gamma} : d_\pi = n\}$. We know that $Z_\Gamma(x) \supseteq Z(\Gamma) \cup xZ(\Gamma)$. To prove the converse inclusion, let $c \in Z_\Gamma(x) \setminus Z(\Gamma)$. It suffices to prove that $x^{-1}c \in Z(\Gamma)$, and since $\Omega_1 \cup \Omega_2$ separates the points of Γ , it suffices to prove that $\pi(x^{-1}c) \in \mathbb{C}I_\pi$ for all $\pi \in \Omega_1 \cup \Omega_2$. This holds trivially if $\pi \in \Omega_1$, and for $\pi \in \Omega_2$ it follows from Lemma 4.8(i).

Similarly, to prove that $|\Gamma : Z_\Gamma(x)| = 2$, let a and b be elements of $\Gamma \setminus Z_\Gamma(x)$. It suffices to prove that $\pi(a^{-1}bx) = \pi(xa^{-1}b)$ for all $\pi \in \Omega_1 \cup \Omega_2$. This holds trivially if $\pi \in \Omega_1$, and for $\pi \in \Omega_2$ it follows from Lemma 4.8(ii). \square

We can now give a proof of Theorem 1.7 for countable groups.

- Let Λ be a countable group satisfying $\text{AD}(\Lambda) = 3/2$. Since $\text{AD}(\Lambda) < \infty$, Λ is virtually abelian. Let ν be Plancherel measure for $\widehat{\Lambda}$. By Lemma 4.4 $\nu(\Omega_1) = 1/2$ and $\nu(\{\pi \in \widehat{\Lambda} : d_\pi \geq 3\}) = 0$. Hence, by Proposition 4.3 $|\text{comm}(\Lambda)| = 2$. Moreover, if $x, x' \in \Lambda$ satisfy $\pi(x) = \pi(x')$ for all $\pi \in \widehat{\Lambda}$ with $d_\pi \leq 2$, then by injectivity of the non-abelian Fourier transform (or, alternatively, by applying the Plancherel identity (2.3) to $\langle \delta_x, \delta_{x'} \rangle$) we must have $x = x'$. By (iii) \implies (ii) in Theorem 4.6, $|\Lambda : Z(\Lambda)| = 4$.
- Conversely, suppose Λ is a countable group with $|\Lambda : Z(\Lambda)| = 4$. By (i) \implies (ii) in Theorem 4.6, $|\text{comm}(\Lambda)| = 2$ and $\max\text{deg}(\Lambda) = 2$. In particular, $\{\pi : d_\pi \geq 3\}$ is empty and so has measure zero, while by Proposition 4.3 $\nu(\Omega_1) = 1/2$. Applying Lemma 4.4 in the other direction we conclude that $\text{AD}(\Lambda) = 3/2$.

To obtain the general case we need another group-theoretic lemma.

LEMMA 4.9 (Control by finitely generated subgroups). *Let Γ be a group and let $m \in \mathbb{N}$. If $|\Lambda : Z(\Lambda)| \leq m$ for every finitely generated subgroup $\Lambda \leq \Gamma$, then $|\Gamma : Z(\Gamma)| \leq m$.*

Proof. We prove the contrapositive statement. Suppose $|\Gamma : Z(\Gamma)| \geq m+1$, and pick a_1, \dots, a_{m+1} which belong to different cosets of $Z(\Gamma)$. Then for each $(i, j) \in \mathbb{N}^2$ with $1 \leq i < j \leq m+1$, we have $a_i^{-1}a_j \notin Z(\Gamma)$, so there exists $t_{ij} \in \Gamma$ such that $a_i^{-1}a_j$ does not commute with t_{ij} . Let Λ be the subgroup of Γ generated by $\{a_1, \dots, a_{m+1}\} \cup \{t_{ij} : 1 \leq i < j \leq m+1\}$. Then Λ is a finitely generated subgroup of Γ , and $|\Lambda : Z(\Lambda)| \geq m+1$ since a_1, \dots, a_{m+1} belong to different cosets of $Z(\Lambda)$. \square

Proof of Theorem 1.7 in the general case. Suppose Γ is a group with $\text{AD}(\Gamma) = 3/2$. Since Γ is non-abelian, $|\Gamma : Z(\Gamma)| \geq 4$ by Lemma 4.5. It suffices to show that $|\Gamma : Z(\Gamma)| \leq 4$.

Let $\Lambda \leq \Gamma$ be a finitely generated subgroup. If Λ is abelian $|\Lambda : Z(\Lambda)| = 1$. If Λ is non-abelian, then by our sharp lower bound (Theorem 1.6) and monotonicity of AD (Proposition 2.11), $3/2 \leq \text{AD}(\Lambda) \leq \text{AD}(\Gamma) = 3/2$. So $\text{AD}(\Lambda) = 3/2$, and since Λ is countable, the previous reasoning tells us that $|\Lambda : Z(\Lambda)| = 4$. Thus $|\Lambda : Z(\Lambda)| \leq 4$ for every finitely generated subgroup $\Lambda \leq \Gamma$. By Lemma 4.9 $|\Gamma : Z(\Gamma)| \leq 4$, as required.

Conversely, suppose Γ is a group satisfying $|\Gamma : Z(\Gamma)| = 4$. By countable saturation for AD (Proposition 2.12) there is a countable subgroup $\Lambda \leq \Gamma$ such that $\text{AD}(\Lambda) = \text{AD}(\Gamma)$. Recall (Lemma 2.10) that a group has antidiagonal constant 1 if and only if it is abelian. Therefore, since Γ is non-abelian so is Λ , and hence by Lemma 4.5 $|\Lambda : Z(\Lambda)| \geq 4$. On the other hand, since $Z(\Lambda) \supseteq \Lambda \cap Z(\Gamma)$,

$$|\Lambda : Z(\Lambda)| \leq |\Lambda : \Lambda \cap Z(\Gamma)| \leq |\Gamma : Z(\Gamma)| = 4,$$

and so $|\Lambda : Z(\Lambda)| = 4$. Since Λ is countable, the previous reasoning tells us that $\text{AD}(\Lambda) = 3/2$, and so $\text{AD}(\Gamma) = 3/2$ as required. \square

5 Examples where $\text{AD}(G)$ is as large as possible

In this section we prove Theorem 1.10. We shall build up to this result in stages. Throughout this section, $[\Gamma, \Gamma]$ denotes the subgroup of Γ generated by the set of commutators (the commutator subgroup or derived subgroup of Γ).

LEMMA 5.1. *Let Γ be a countable VA group, equipped with counting measure, and let ν be the corresponding Plancherel measure on $\widehat{\Gamma}$. Let $\Omega_1 = \{\pi \in \widehat{\Gamma} : d_\pi = 1\}$. If $[\Gamma, \Gamma]$ is infinite then $\nu(\Omega_1) = 0$.*

Proof. We prove the contrapositive statement. Assume Ω_1 has strictly positive measure, and for each $x \in \Gamma$ put $h(x) = \int_{\Omega_1} \pi(x) d\nu(\pi)$. Note that $h(x) = \nu(\Omega_1) > 0$ for all $x \in [\Gamma, \Gamma]$. We claim that $h \in \ell^2(\Gamma)$. Assuming for the moment that this holds, we deduce that

$$0 < |[\Gamma, \Gamma]|^{1/2} \nu(\Omega_1) \leq \|h\|_2 < \infty$$

and conclude that $|[\Gamma, \Gamma]| < \infty$ as required.

The claim follows from non-abelian Fourier inversion for Γ , but we can give a proof that only requires⁴ Equation (2.2). Let $f \in c_{00}(\Gamma)$, and note that by interchanging finite summation with integration we have

$$\sum_{x \in \Gamma} f(x)h(x) = \sum_{x \in \Gamma} f(x) \int_{\Omega_1} \pi(x) d\nu(\pi) = \int_{\Omega_1} \pi(f) d\nu(\pi).$$

Therefore, applying Cauchy–Schwarz and Equation (2.2) yields

$$\left| \sum_{x \in \Gamma} f(x)h(x) \right| \leq \nu(\Omega_1)^{1/2} \left(\int_{\Omega_1} |\pi(f)|^2 d\nu(\pi) \right)^{1/2} \leq \nu(\Omega_1)^{1/2} \|f\|_2.$$

By standard duality arguments in $\ell^2(\Gamma)$ we deduce that $\|h\|_2 \leq \nu(\Omega_1)^{1/2} < \infty$, as claimed. This completes the proof. \square

REMARK 5.2. The proof of Lemma 5.1 can be generalized to any 2nd-countable unimodular Type I group G , and yields the following result: if the commutator subgroup in G has non-compact closure, then the set of one-dimensional representations is a Plancherel-null subset of \widehat{G} . We leave the details to the reader, since we have no applications at present for this more general result.

PROPOSITION 5.3. *Let Γ be a countable, VA group. Suppose that the following two conditions hold:*

- (a) $[\Gamma, \Gamma]$ is infinite;
- (b) there exists some $d \geq 2$ such that $d_\pi \in \{1, d\}$ for all $\pi \in \widehat{\Gamma}$.

Then $d = \text{AD}(\Gamma) = \text{AM}(\text{A}(\Gamma)) = \text{maxdeg}(\Gamma)$.

(Example 3.10 shows that the assumption (a) is necessary.)

⁴That is, we only need the fact that the non-abelian Fourier transform is an isometry from $\ell^2(\Gamma)$ to a suitable space of operator fields, rather than the subtler issue of characterizing the range of this isometry

Proof. Recall (see (1.5) above) that by results in [Run06] we have $d = \max \deg(\Gamma) \geq \text{AM}(A(\Gamma)) \geq \text{AD}(\Gamma)$. It therefore suffices to prove that $\text{AD}(\Gamma) = d$.

Let ν be the Plancherel measure on $\widehat{\Gamma}$ corresponding to counting measure on Γ . By Assumption (b),

$$1 = \int_{\widehat{\Gamma}} d_\pi d\nu(\pi) = \nu(\Omega_1) + d\nu(\Omega_d),$$

while by Theorem 1.5,

$$\text{AD}(\Gamma) = \int_{\widehat{\Gamma}} d_\pi^2 d\nu(\pi) = \nu(\Omega_1) + d^2\nu(\Omega_d).$$

Hence $\text{AD}(\Gamma) = \nu(\Omega_1) + d(1 - \nu(\Omega_1))$. But, by Assumption (a) and Lemma 5.1, $\nu(\Omega_1) = 0$, and the result follows. \square

LEMMA 5.4. *Let N be a torsion-free LCA group, let $m \geq 2$, and let $\alpha : C_m \rightarrow \text{Aut}(N)$. Let $G = N \rtimes_\alpha C_m$ be the corresponding semidirect product. Then $d_\pi \mid m$ for all $\pi \in \widehat{G}$.*

Proof. This follows from (a simple case of) the Mackey machine, as described in e.g. [KT13]. Given $\chi \in \widehat{N}$, let D_χ denote the stabilizer of χ with respect to the adjoint action of C_p on \widehat{N} , and let $G_\chi = N \rtimes_\alpha D_\chi \subset G$. Then each $\pi \in \widehat{G}$ is unitarily equivalent to one obtained by inducing some one-dimensional representation $G_\chi \rightarrow \mathbb{T}$. (See [KT13, Theorem 4.40] for a more general result.) In particular, $d_\pi = |G : G_\chi| = |C_m : D_\chi|$, which divides $|C_m| = m$ by Lagrange's theorem. \square

The proof of Theorem 1.10. Recall the setup: N is a torsion-free LCA group, p is a prime, and $G = N \rtimes_\alpha C_p$ for some non-trivial action α . By Lemma 5.4 we have $\max \deg(G) \leq p$. (In fact, equality holds, but we will obtain this for free later on.)

By assumption there exists $x_0 \in N$ whose α -orbit contains at least one other point of N . Let N_{00} be the subgroup of N (not necessarily closed!) that is generated by this orbit; then N_{00} is countable, torsion-free abelian, and α -invariant. Let $\Gamma = N_{00} \rtimes_\alpha C_p$: this is a subgroup of G_d , and so by monotonicity of AD,

$$\text{AD}(\Gamma) \leq \text{AD}(G_d) \leq \text{AM}(A(G)) \leq p. \quad (*)$$

Now observe the following:

- (a) Let x_1 be any point in the C_p -orbit of x_0 with $x_1 \neq x_0$. Since x_0 and x_1 are conjugate in Γ , $x_0^{-1}x_1$ is a non-trivial commutator in Γ ; also $x_0^{-1}x_1 \in N_{00}$, which is torsion-free. Thus $[\Gamma, \Gamma]$ contains a free abelian subgroup, and in particular is infinite.
- (b) Applying Lemma 5.4 with N replaced by N_{00} , and recalling that p is prime, we see that $d_\pi \in \{1, p\}$ for every $\pi \in \widehat{\Gamma}$.

Applying Proposition 5.3 we conclude that $\text{AD}(\Gamma) = p$, and so all the inequalities in (*) must be equalities. \square

6 Concluding remarks and further questions

The most obvious and important question is:

QUESTION 6.1. Do we always have $\text{AM}(A(G)) = \text{AD}(G_d)$?

A positive answer, combined with Equation (1.4) and monotonicity, would provide a powerful tool for estimating or calculating amenability constants of Fourier algebras.

To see why the answer to Question 6.1 is not obvious, we give a sketch of how the inequality $\text{AM}(A(G)) \geq \text{AD}(G_d)$ is proved in [FR05, Run06]. By definition, $\text{AM}(A(G))$ is the infimum of norms of virtual diagonals for $A(G)$. We will not define virtual diagonals here: they are elements $\mathbf{M} \in (A(G) \widehat{\otimes} A(G))^{**}$ satisfying certain algebraic properties, where $\widehat{\otimes}$ denotes the projective tensor product of Banach spaces. There is a linear contraction $J : A(G) \widehat{\otimes} A(G) \rightarrow A(G \times G)$ satisfying $J(f \otimes g)(x, y) = f(x)g(y^{-1})$, and one can show this extends to a linear contraction $\tilde{J} : (A(G) \widehat{\otimes} A(G))^{**} \rightarrow B(G_d \times G_d)$. Moreover, the algebraic properties of \mathbf{M} imply that $\tilde{J}(\mathbf{M}) = 1_{\text{adiag}(G)}$. Hence

$$\text{AD}(G_d) = \|1_{\text{adiag}(G)}\|_B \leq \inf_{\mathbf{M}} \|\mathbf{M}\| = \text{AM}(A(G)).$$

The point is that for non-abelian G , J is never an isometry (although it is bijective if G is virtually abelian), and \tilde{J} is not a quotient map of Banach spaces. Hence even if one knows that the set $\tilde{J}^{-1}(\{1_{\text{adiag}(G)}\})$ contains a virtual diagonal \mathbf{M} , there is no immediate reason why one should be able to find such an \mathbf{M} whose norm equals that of $1_{\text{adiag}(G)}$.

On the other hand, if G is finite and non-abelian, then even though $J : A(G) \widehat{\otimes} A(G) \rightarrow A(G \times G)$ need not be isometric we know from Theorem 1.4 that we do have $\text{AM}(A(G)) = \text{AD}(G)$. So if there is a group G for which $\text{AM}(A(G)) > \text{AD}(G_d)$, there must be some deeper obstruction behind this which is not visible for finite groups; Theorem 1.10 shows that we can have groups that look very different from “finite times abelian” where $\text{AM}(A(G)) = \text{AD}(G_d)$.

An additional motivation for Question 6.1 is that a positive answer, combined with Theorem 1.7, would give a complete characterization of those locally compact non-abelian G where $\text{AM}(A(G))$ attains its minimal value. We phrase this as a separate question, which might be easier to attack directly.

QUESTION 6.2. Which non-abelian G attain the minimal value for $\text{AM}(A(G))$?

If $\text{AM}(A(G)) = 3/2$ then, since G is non-abelian, our sharp lower bound (Theorem 1.6) implies that

$$3/2 \leq \text{AD}(G_d) \leq \text{AM}(A(G)) = 3/2$$

and so by Theorem 1.7 $|G : Z(G)| = 4$. Thus Question 6.2 is really asking if the converse holds, i.e. does $|G : Z(G)| = 4$ imply $\text{AM}(A(G)) = 3/2$?

QUESTION 6.3. Can we bound $\text{maxdeg}(G)$ from above by some explicit function of $\text{AD}(G_d)$?

This was done for finite groups in [LLW96, Theorem 4.9] but to the author’s knowledge it remains an open problem for infinite VANA groups.

QUESTION 6.4. Can we classify those Γ such that $3/2 < \text{AD}(\Gamma) < 2$?

Note that if Λ is countable and $|\llbracket \Lambda, \Lambda \rrbracket|$ is infinite, then by Lemma 5.1 and the main formula (Equation (1.4)) we have $\text{AD}(\Lambda) \geq 2$. Hence the condition that $\text{AD}(\Gamma) < 2$ forces $[\Gamma, \Gamma]$ to be finite (if it were infinite then there would be a countable $\Lambda \leq \Gamma$ witnessing this). There is a substantial literature on groups whose commutator subgroups are finite (so-called FD groups) and one might be able to use known structure theory to attack Question 6.4.

On the other hand, the answer to Question 6.4 does not seem to be known even for finite groups. While investigating Question 6.4 for finite groups, all examples we found satisfying $\text{AD}(\Gamma) \leq 2$ also satisfy $\text{maxdeg}(\Gamma) \leq 2$.

QUESTION 6.5. Let Γ be a finite group with $\text{AD}(\Gamma) \leq 2$. Must we also have $\text{maxdeg}(\Gamma) \leq 2$?

The author thanks G. Robinson (personal communication) for some preliminary discussions. These indicate that if the answer to Question 6.5 is negative, then a minimal counterexample must have rather strange or restrictive properties. On the other hand, so far this falls short of compelling evidence for a positive answer.

Acknowledgements

The challenge of finding optimal lower bounds on $\text{AM}(\text{A}(G))$ for infinite non-abelian groups originally arose during conversations in 2016 with M. Alaghmandan, who the author thanks for some stimulating discussions. He also thanks M. Ghandehari for several valuable discussions about the overall project, and is indebted to her for the crucial suggestion to make use of results in [Ars76, Section 3G]. Thanks also go to: Y. Cornulier for pointing out via *MathOverflow* that an example which the author had laboriously constructed by hand was in fact isomorphic to the group in Example 2.2; N. Juselius for sharing a copy of [Jus19]; and K. F. Taylor for sharing some private notes on the Plancherel theorem for crystal groups, which led to the additional examples mentioned in Remark 1.12.

Key progress towards the main formula (Theorem 1.5) was made, and some special cases of Theorem 1.10 were obtained, through discussions with Ghandehari during a visit to the University of Delaware in March 2020. The author thanks his hosts for their hospitality, and acknowledges financial support from the Visitor Fund of the Department of Mathematics and Statistics at Lancaster University.

Important revisions to an earlier version of this paper were worked out while the author was visiting the Laboratoire de mathématiques de Besançon, Université de Franche-Comté, in November 2021. The author thanks his hosts for a welcoming and pleasant working environment, and acknowledges financial support from the Alliance Hubert Curien programme.

A Continuity of the diagonal embedding $\widehat{G} \rightarrow (G \times G)^\wedge$

In this appendix we provide a proof of the following result, which may well be folklore.

THEOREM A.1. *Let G be a locally compact group, and define $\text{diag} : \widehat{G} \rightarrow (G \times G)^\wedge$ by $\text{diag}([\pi]) = [\pi \otimes \pi]$. Then diag is continuous with respect to the Fell topologies on \widehat{G} and $(G \times G)^\wedge$.*

Here, we are not claiming that $(G \times G)^\wedge = \widehat{G} \times \widehat{G}$ as sets; this can fail in general, even for countable G , although it will hold if G is Type I.

REMARK A.2. The function diag in Theorem A.1 is well-defined, because:

- if $\pi : G \rightarrow \mathcal{U}(H_\pi)$ is an irrep, then $\pi \otimes \pi : G \times G \rightarrow \mathcal{U}(H_\pi \otimes_2 H_\pi)$ is also an irrep;
- if π and σ are unitarily equivalent representations of G , then $\pi \otimes \pi$ and $\sigma \otimes \sigma$ are unitarily equivalent representations of $G \times G$.

There are various equivalent definitions of the Fell topology on the dual of a locally compact group, usually in terms of a closure operator rather than a (sub-)base for the topology. In this context, note that given two topological spaces X and Y and a function $f : X \rightarrow Y$, f is continuous if and only if $\overline{f^{-1}(T)} \subseteq f^{-1}(\overline{T})$ for all $T \subseteq Y$ (this is a straightforward exercise).

The following result is originally due to Fell. A convenient source is Section 18.1 of Dixmier's book [Dix77]. In the form stated below, it is (part of) [Dix77, Proposition 18.1.5]: see [Dix77, Paragraphs 13.4.6 and 18.1.1] for some of the terminology, and [Dix77, Paragraph 18.1.3] for a roadmap of the C^* -algebraic results needed in the proof.

PROPOSITION A.3. *Let $S \subseteq \widehat{G}$ and let σ be an irrep of G . Then $[\sigma] \in \overline{S}$ if and only if there exist*

- a non-zero $\eta \in H_\sigma$,
- a net $(\pi_i)_{i \in I}$ where each π_i is an irrep of G and $[\pi_i] \in S$,
- for each $i \in I$, some $\xi_i \in H_{\pi_i}$

such that the net of coefficient functions $\langle \pi_i(\cdot)\xi_i, \xi_i \rangle$ converges to the coefficient function $\langle \sigma(\cdot)\eta, \eta \rangle$ uniformly on compact subsets of G .

Proof of Theorem A.1. Let $T \subseteq (G \times G)^\wedge$ and let $S = \text{diag}^{-1}(T) = \{[\pi] \in \widehat{G} : [\pi \otimes \pi] \in T\}$. Given $\sigma \in \overline{S}$, let η , $(\pi_i)_{i \in I}$ and $(\xi_i)_{i \in I}$ have the properties described in Proposition A.3. Then:

- (i) $\eta \otimes \eta$ is a non-zero vector in $H_{\sigma \otimes \sigma} \equiv H_\sigma \otimes_2 H_\sigma$;
- (ii) we may form the net $(\pi_i \otimes \pi_i)_{i \in I}$, noting that $[\pi_i \otimes \pi_i] \in T$ since $[\pi_i] \in S$;
- (iii) for each $i \in I$, we have $\xi_i \otimes \xi_i \in H_{\pi_i \otimes \pi_i} \equiv H_{\pi_i} \otimes_2 H_{\pi_i}$.

To ease notation, let

$$\begin{aligned} f_i(x, y) &:= \langle (\pi_i \otimes \pi_i)(x, y)(\xi_i \otimes \xi_i), \xi_i \otimes \xi_i \rangle = \langle \pi_i(x)\xi_i, \xi_i \rangle \langle \pi_i(y)\xi_i, \xi_i \rangle \\ h(x, y) &:= \langle (\sigma \otimes \sigma)(x, y)(\eta \otimes \eta), \eta \otimes \eta \rangle = \langle \sigma(x)\eta, \eta \rangle \langle \sigma(y)\eta, \eta \rangle \end{aligned}$$

Since $\langle \pi_i(\cdot)\xi_i, \xi_i \rangle$ converges to $\langle \sigma(\cdot)\eta, \eta \rangle$ uniformly on compact subsets of G , f_i converges to h uniformly on compact subsets of $G \times G$. By the way we have defined f_i and h , and properties (i)–(iii) above, it follows from Proposition A.3 that $[\sigma \otimes \sigma] \in \overline{T}$. This completes the proof. \square

References

- [Ars76] Gilbert Arsac. Sur l'espace de Banach engendr e par les coefficients d'une repr esentation unitaire. *Publ. D ep. Math. (Lyon)*, 13(2):1–101, 1976.
- [Dix77] Jacques Dixmier. *C*-algebras*, volume 15 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1977. Translated from the French by Francis Jollett.
- [Dix81] Jacques Dixmier. *Von Neumann algebras*, volume 27 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1981. Translated from the second French edition by F. Jollett.
- [Eym64] Pierre Eymard. L'alg ebre de Fourier d'un groupe localement compact. *Bull. Soc. Math. France*, 92:181–236, 1964.
- [Fol16] Gerald B. Folland. *A course in abstract harmonic analysis*. Textbooks in Mathematics. CRC Press, Boca Raton, FL, second edition, 2016.

- [FR05] Brian E. Forrest and Volker Runde. Amenability and weak amenability of the Fourier algebra. *Math. Z.*, 250(4):731–744, 2005.
- [FR11] Brian E. Forrest and Volker Runde. Norm one idempotent cb -multipliers with applications to the Fourier algebra in the cb -multiplier norm. *Canad. Math. Bull.*, 54(4):654–662, 2011.
- [Joh94] Barry E. Johnson. Non-amenability of the Fourier algebra of a compact group. *J. London Math. Soc. (2)*, 50(2):361–374, 1994.
- [Jus19] Nicholas A. Juselius. Bounds on the amenability constants of Fourier algebras in the completely bounded multiplier norm. Master’s thesis, University of Alberta, 2019.
- [KL18] Eberhard Kaniuth and Anthony To-Ming Lau. *Fourier and Fourier-Stieltjes algebras on locally compact groups*, volume 231 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2018.
- [KT13] Eberhard Kaniuth and Keith F. Taylor. *Induced representations of locally compact groups*, volume 197 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2013.
- [LLW96] Anthony To-Ming Lau, Richard J. Loy, and George A. Willis. Amenability of Banach and C^* -algebras on locally compact groups. *Studia Math.*, 119(2):161–178, 1996.
- [MP16] Jayden Mudge and Hung Le Pham. Idempotents with small norms. *J. Funct. Anal.*, 270(12):4597–4603, 2016.
- [Run06] Volker Runde. The amenability constant of the Fourier algebra. *Proc. Amer. Math. Soc.*, 134(5):1473–1481, 2006.
- [Wal72] Martin E. Walter. W^* -algebras and nonabelian harmonic analysis. *J. Functional Analysis*, 11:17–38, 1972.

YEMON CHOI, DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LANCASTER LA1 4YF, UNITED KINGDOM.

y.choi1@lancaster.ac.uk