

Periodic orbit evaluation of a spectral statistic of quantum graphs without the semiclassical limit

Jon Harrison*

Baylor University, Department of Mathematics, Sid Richardson Building, 1410 S. 4th Street, Waco, TX 76706, USA

Tori Hudgins†

University of Dallas, Department of Mathematics, 1845 E Northgate Dr, Irving, TX 75062, USA

(Dated: April 21, 2022)

We evaluate the variance of coefficients of the characteristic polynomial of the quantum evolution operator for chaotic 4-regular quantum graphs (networks) via periodic orbits without taking the semiclassical limit. The variance of the n -th coefficient is precisely determined by the number of primitive pseudo orbits (sets of distinct primitive periodic orbits) with n bonds that fall in the following classes: those with no self-intersections, and those where all the self-intersections consist of two sections of the pseudo orbit crossing at a single vertex (2-encounters of length zero).

The analysis of chaotic quantum systems via their periodic orbits originates with Gutzwiller’s trace formula [1] which has been applied to analyze spectral statistics like the form factor in the semiclassical limit. This starts with Berry [2] who computes a diagonal approximation using [3]. Subsequently, Sieber and Richter [4, 5] evaluated a first-order correction by considering figure eight orbits with a self-intersection. This was extended to higher orders in [6, 7]. Quantum graphs are used to study quantum systems with complex topology in fields from Anderson localization to carbon nanostructures, see e.g. [8]. Kottos and Smilansky [9, 10] introduced quantum graphs to study quantum mechanics in a system that is classically chaotic. They demonstrated that, in the semiclassical limit, spectral statistics can be traced to topological properties of the periodic orbits. Spectra of quantum graphs are investigated experimentally in microwave networks [11, 12]. We show that, for a large class of chaotic quantum graphs, one can precisely compute a spectral statistic from the periodic orbit structure without the semiclassical limit.

A graph is a network of V vertices connected by B bonds. We consider 4-regular graphs where every vertex is connected by a bond to four neighboring vertices. We take the graph to be directed, so every vertex has two incoming and two outgoing bonds. (This is always possible as 4-regular graphs admit an Euler tour.) Figure 1 shows a 4-regular directed graph. A bond $b = (u, v)$ is an ordered pair of vertices; $o(b) = u$ and $t(b) = v$, the origin and terminus of b , respectively. In a quantum graph the bonds are assigned a positive length which we collect in a diagonal matrix $L = \text{diag}\{L_1, \dots, L_B\}$. At each vertex v there is an associated unitary vertex scattering matrix $\sigma^{(v)}$ which describes scattering from incoming to outgoing bonds. The elements of these matrices are collected in a $B \times B$ bond scattering matrix,

$$S_{b',b} = \delta_{o(b'),v} \delta_{t(b),v} \sigma_{b',b}^{(v)}. \quad (1)$$

The spectrum of a quantum graph corresponds to roots

of a secular equation,

$$\det(Se^{ikL} - \mathbf{I}) = 0, \quad (2)$$

where k is the spectral parameter [9, 10]. The matrix $U(k) = Se^{ikL}$ is the quantum evolution operator.




FIG. 1. Binary de Bruijn graph with 2^3 vertices.

Given a Hamiltonian, the vertex scattering matrices are determined by the matching conditions at the vertices. If $k \neq 0$ is a root of (2) then k^2 is an eigenvalue with the same multiplicity. We adopt a complimentary approach to quantize graphs where we simply specify unitary scattering matrices at the vertices [13]. This defines a unitary stochastic matrix ensemble $U(k)$. The vertex scattering matrix we use to quantize 4-regular graphs is the discrete Fourier transform (DFT) matrix,

$$\sigma^{(v)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}. \quad (3)$$

The DFT matrix is frequently used to quantize graphs [14]. It is democratic, in the sense that the transmission probability $|\sigma_{b',b}^{(v)}|^2$ is the same for every pair b, b' .

The characteristic polynomial of a quantum graph is,

$$\det(U(k) - \zeta \mathbf{I}) = \sum_{n=0}^B a_n \zeta^{B-n}. \quad (4)$$

The left hand side of (2) is the characteristic polynomial at $\zeta = 1$. The unitarity of $U(k)$ produces a Riemann-Siegel lookalike formula for the coefficients $a_n = a_B \bar{a}_{B-n}$ [10]. The n -th coefficient can be written as a sum over

sets of distinct primitive periodic orbits (primitive pseudo orbits) [15]. Averaging over k , the spectral parameter, $\langle a_n \rangle = 0$ for $0 < n < B$. So the first non-trivial moment is the variance,

$$\langle |a_n|^2 \rangle = \sum_{\bar{\gamma}, \bar{\gamma}' \in \mathcal{P}^n} (-1)^{m_{\bar{\gamma}} + m_{\bar{\gamma}'}} A_{\bar{\gamma}} \bar{A}_{\bar{\gamma}'} \delta_{L_{\bar{\gamma}}, L_{\bar{\gamma}'}} , \quad (5)$$

where \mathcal{P}^n is the set of primitive pseudo orbits with n bonds. A primitive pseudo orbit $\bar{\gamma}$ consists of $m_{\bar{\gamma}}$ distinct primitive periodic orbits, so $\bar{\gamma} = \{\gamma_1, \gamma_2, \dots, \gamma_{m_{\bar{\gamma}}}\}$ where γ_i is a periodic orbit that is not a repetition of a shorter orbit and $\gamma_i \neq \gamma_j$ for $i \neq j$. Associated to an orbit γ_i is a stability amplitude A_{γ_i} , the product of the scattering amplitudes of the vertex scattering matrices associated with γ_i . Then $A_{\bar{\gamma}}$ is the product of the stability amplitudes of the orbits in $\bar{\gamma}$. The length of a periodic orbit is the sum of the lengths its bonds, and $L_{\bar{\gamma}}$ is the sum of the lengths of the orbits in $\bar{\gamma}$. Notice that pairs of primitive pseudo orbits $\bar{\gamma}, \bar{\gamma}'$ that contribute to the variance must have the same length $L_{\bar{\gamma}} = L_{\bar{\gamma}'}$. We take the set of bond lengths to be incommensurate (not rationally related), so if two pseudo orbits have the same length they visit each bond the same number of times. Let,

$$C_{\bar{\gamma}} = \sum_{\bar{\gamma}' \in \mathcal{P}_{\bar{\gamma}}} (-1)^{m_{\bar{\gamma}} + m_{\bar{\gamma}'}} A_{\bar{\gamma}} \bar{A}_{\bar{\gamma}'}, \quad (6)$$

where $\mathcal{P}_{\bar{\gamma}}$ is the set of primitive pseudo orbits of length $L_{\bar{\gamma}}$. The variance of the coefficients (5) is then,

$$\langle |a_n|^2 \rangle = \sum_{\bar{\gamma} \in \mathcal{P}^n} C_{\bar{\gamma}}. \quad (7)$$

We now state the main result,

$$\langle |a_n|^2 \rangle = \frac{1}{2^n} \left(|\mathcal{P}_0^n| + \sum_{N=1}^n 2^N |\hat{\mathcal{P}}_N^n| \right), \quad (8)$$

where \mathcal{P}_0^n is the set of primitive pseudo orbits of length n with no self-intersections and $\hat{\mathcal{P}}_N^n$ is the set of primitive pseudo orbits of length n with N self-intersections, all of which are 2-encounters of length zero. We can use n as the last term in the sum, as an orbit of length n cannot have more than n self-intersections.

The first term in (8) is straightforward to explain. From (3) we see that, for a pseudo orbit of n bonds, $|A_{\bar{\gamma}}|^2 = 1/2^n$. If $\bar{\gamma}$ has no self-intersections it is not possible to reorder the bonds of $\bar{\gamma}$ to produce a different pseudo orbit with the same length. Hence $\mathcal{P}_{\bar{\gamma}} = \{\bar{\gamma}\}$ and $C_{\bar{\gamma}} = 1/2^n$. However, if $\bar{\gamma}$ has one or more self-intersections it is possible to rearrange sections of $\bar{\gamma}$ to produce other pseudo orbits of the same length. We will see that typically, if $\bar{\gamma}$ has self-intersections, $C_{\bar{\gamma}} = 0$.

A self-intersection is a section of a pseudo orbit that is repeated in the pseudo orbit. The repeated section

can be one that appears in more than one of the orbits in the pseudo orbit or is repeated multiple times in a single periodic orbit or some combination. If a section is repeated three or more times there will also be preceding and subsequent sections that are repeated fewer times, as there are only two incoming and two outgoing bonds at each vertex. We call the maximally repeated section the encounter. An l -encounter is an encounter that is repeated l times in the pseudo orbit.

Figure 2 shows a subgraph from which we can construct pseudo orbits with self-intersections. In the figure, the encounter is the sequence of adjacent vertices $\text{enc} = (v_0, \dots, v_r)$ where r is the length of the encounter. Sequences of adjacent vertices joining the end of an encounter to the start of an encounter are called links. A pseudo orbit can be defined by specifying the sequences of links in each orbit. Consider $\bar{\gamma} = (\gamma_1, \dots, \gamma_m)$ where there are no self-intersections in $\gamma_2, \dots, \gamma_m$ and $\gamma_1 = (12)$, by which we mean the orbit follows link 1, then the encounter, followed by link 2, and the encounter a second time,

$$\gamma_1 = (f_1 \dots, s_1, \text{enc}, f_2, f_2' \dots, s_2', s_2, \text{enc}, f_1), \quad (9)$$

then $\bar{\gamma}$ has a 2-encounter. Similarly, we could define a primitive pseudo orbit with a 3-encounter by taking $\gamma_1 = (123)$. In this case there are bonds preceding and following the encounter that are repeated twice, (s_2, v_0) and (v_r, f_2) . If $v_0 = v_r$ there are no bonds in the encounter, and we have an encounter of length zero. While the previous examples had the encounter repeated in a single orbit we could define a pseudo orbit with a 2-encounter by taking $\gamma_1 = (1)$ and $\gamma_2 = (2)$ with the remaining orbits not intersecting any other orbit. Finally, we also point out that pseudo orbits with an l -encounter with $l > 2$ can repeat links. So $\gamma_1 = (1232)$ defines an orbit with a 4-encounter which uses link 2 twice.




FIG. 2. Subgraph on which pseudo orbits with self-intersections can be constructed.

The goal is to show that, for most primitive pseudo orbits with self-intersections, $C_{\bar{\gamma}}$ is zero. While this requires technical steps for certain classes of pseudo orbits, in each case the result relies on a parity argument. The parity argument is straightforward when $\bar{\gamma}$ has a single l -encounter of positive length and no repeated links. The

set of partner orbits of the same length $\mathcal{P}_{\bar{\gamma}}$ is labeled by the symmetric group S_l . So, for example, if $\bar{\gamma}$ has a single 3-encounter, $\gamma_1 = (123)$, there are six primitive partner pseudo orbits $\bar{\gamma}'$ with the same length that correspond to replacing γ_1 with an element of

$$S_3 = \{(1)(2)(3), (12)(3), (13)(2), (1)(23), (123), (132)\}. \quad (10)$$

As the encounter has positive length, $A_{\bar{\gamma}'} = A_{\bar{\gamma}}$. Then,

$$C_{\bar{\gamma}} = \pm 2^{-n} \sum_{\sigma \in S_l} (-1)^{|\sigma|}, \quad (11)$$

where $|\sigma|$ is the number of cycles in σ . It is well known that there are an equal number of permutations with odd and even parity; hence $C_{\bar{\gamma}} = 0$.

If there are repeated links in $\bar{\gamma}$, the primitive partner pseudo orbits $\bar{\gamma}'$ are not labeled by elements of S_l , but by Lyndon tuples. A Lyndon word, over an alphabet of l letters, is a word that is strictly earlier in lexicographic order than its rotations. Hence, Lyndon words, in the links of a single l -encounter, uniquely label primitive periodic orbits. Primitive pseudo orbits then correspond to sets of distinct Lyndon words, Lyndon tuples. A multiset $M = [1^{m_1}, 2^{m_2}, \dots, l^{m_l}]$ over the alphabet of l letters records the number of times each of the l links is used in $\bar{\gamma}$. Then the set of primitive partner pseudo orbits is labeled by the set $\mathcal{L}(M)$ of all Lyndon tuples over M . For example,

$$\mathcal{L}([1^2, 2^2]) = \{(1122), (122)(1), (2)(112), (2)(12)(1)\}. \quad (12)$$

Setting $|M| = \sum_{j=1}^l m_j$, the index of a Lyndon tuple with k words is $|M| - k$ and the tuple is odd or even depending on whether the index is odd or even. There are equal numbers of odd and even Lyndon tuples [16] from which we obtain $C_{\bar{\gamma}} = 0$ for pseudo orbits with a single encounter of positive length.

If the encounter has zero length, $A_{\bar{\gamma}} \bar{A}_{\bar{\gamma}'} = \pm 2^{-n}$, where the sign depends on tracking the number of times that the negative scattering amplitude in (3) appears in $\bar{\gamma}'$ relative to $\bar{\gamma}$. The equality of the number of even and odd Lyndon tuples in $\mathcal{L}(M)$ is established by obtaining a bijection from even to odd Lyndon tuples. By describing a bijection for $l > 2$ that does not change the number of times the negative scattering coefficient is used, the parity argument extends to show $C_{\bar{\gamma}} = 0$ for l -encounters of length zero.

If $\bar{\gamma}$ has multiple encounters we will assume there is an l -encounter with positive length or $l > 2$, the first encounter. We construct a subset of primitive partner pseudo orbits by taking $\bar{\gamma}$ and rearranging the links at the first encounter. Summing over this subset produces zero. Then taking a primitive partner pseudo orbit that was not included in the sum, we repeat the procedure, generating partners from it by rearranging at the first encounter. Summing over this new subset again produces

zero. Iterating, the procedure terminates as \mathcal{P}^n is finite. Hence, $C_{\bar{\gamma}} = 0$ for pseudo orbits with any l -encounter of positive length or with $l > 2$.

The remaining primitive pseudo orbits are those where all the self-intersections are 2-encounters of length zero. Let N be the number of such 2-encounters and j the number of those encounters at which links in the partner orbit $\bar{\gamma}'$ are reordered relative to $\bar{\gamma}$. Each encounter at which $\bar{\gamma}'$ is reordered changes the number of periodic orbits in the pseudo orbit by ± 1 . Hence $(-1)^{m_{\bar{\gamma}'}} = (-1)^{m_{\bar{\gamma}}+j}$. Also at every encounter where $\bar{\gamma}'$ is reordered there is an additional change in the sign of the stability amplitude, so $A_{\bar{\gamma}} \bar{A}_{\bar{\gamma}'} = (-1)^j 2^{-n}$. Consequently,

$$C_{\bar{\gamma}} = 2^{-n} \sum_{j=0}^N (-1)^{2j} \binom{N}{j} = 2^{N-n}. \quad (13)$$

We have shown $C_{\bar{\gamma}}$ can be evaluated for all primitive pseudo orbits and substituting these results in (7) produces (8). While we have presented the central idea of the parity argument, it is described in full in [17].

Families of 4-regular directed graphs called binary graphs were studied by Tanner [13, 18, 19]. Binary graphs have $V = p2^r$ where p is odd, with the bonds defined by the adjacency matrix,

$$[A]_{ij} = \begin{cases} \delta_{2i,j} + \delta_{2i+1,j}, & 0 \leq i < V/2, \\ \delta_{2i-V,j} + \delta_{2i+1-V,j}, & V/2 \leq i < V, \end{cases} \quad (14)$$

where $[A]_{ij} = 1$ if i is connected to j and is zero otherwise. Binary de Bruijn graphs are the family of binary graphs with $p = 1$. Figure 1 shows the binary de Bruijn graph with $V = 2^3$.

To demonstrate the result, tables I and II compare the variance of the first half of the coefficients of the characteristic polynomial evaluated using (8) with the numerically computed variance. (The variance of the remaining coefficients is determined by the symmetry about $n = B/2$.) The averages over k were computed for spectral intervals containing 50 million eigenvalues. The sets of primitive pseudo orbits used to compute the variance from (8) are in [17]. In each case we see agreement to at least four decimal places.

We have shown that there is a quantum chaotic system for which a spectral statistic can be computed dynamically without the semiclassical limit. The result is surprising as only particular sets of pseudo orbits contribute. However, in the semiclassical limit, the asymptotic variance can be obtained from a diagonal argument which weights all the orbits equally [19, 20]. The semiclassical limit is the limit of increasing spectral density, which for quantum graphs is the limit of a sequence of graphs with an increasing number of bonds [8, 9, 14]. For families of binary graphs with p fixed and $B = p2^{r+1}$ the semiclassical limit is $r \rightarrow \infty$. To take the semiclassical limit in (8) we must determine the proportion of orbits where all

TABLE I. Variance of coefficients of the characteristic polynomial of a binary de Bruijn quantum graph with 2^3 vertices.

n	$ \mathcal{P}_0^n $	$ \widehat{\mathcal{P}}_1^n $	$ \widehat{\mathcal{P}}_2^n $	$\langle a_n ^2 \rangle$	Numerics	Error
0	1	0	0	1	1.000000	0.000000
1	2	0	0	1	0.999991	0.000009
2	2	0	0	1/2	0.499999	0.000001
3	4	0	0	1/2	0.499999	0.000001
4	8	0	0	1/2	0.499999	0.000001
5	8	8	0	3/4	0.749998	0.000002
6	8	20	0	3/4	0.749986	0.000014
7	16	16	8	5/8	0.624989	0.000011
8	16	16	24	9/16	0.562501	-0.000001

TABLE II. Variance of coefficients of the characteristic polynomial of a binary quantum graph with $3 \cdot 2$ vertices.

n	$ \mathcal{P}_0^n $	$ \widehat{\mathcal{P}}_1^n $	$\langle a_n ^2 \rangle$	Numerics	Error
0	1	0	1	1.000000	0.000000
1	2	0	1	1.000000	0.000000
2	3	0	3/4	0.750001	-0.000001
3	6	0	3/4	0.750003	-0.000003
4	10	4	7/8	0.874999	0.000001
5	8	4	1/2	0.499998	0.000002
6	8	8	3/8	0.374999	0.000001

the self-intersections are 2-encounters of length zero on large graphs, fixing the ratio n/B .

Following a pseudo orbit with a 2-encounter, when you reach the initial encounter vertex for the second time, there are two outgoing bonds, one of which has already appeared in the orbit. The probability to scatter in each direction is $1/2$ and so, for long pseudo orbits on a large graph, half of primitive pseudo orbits with a 2-encounter will have encounter length zero. In order to have a 3-encounter the pseudo orbit must already have passed through the encounter twice and then hit an encounter vertex again. As the graph is mixing, the probability to end on any bond after a large number of steps is $1/B$. Hence, for long orbits on large graphs, the proportion of 3-encounters is vanishingly small compared to the number of 2-encounters. Let \mathcal{P}_N^n denote the set of primitive pseudo orbits of length n with N encounters. Then $|\widehat{\mathcal{P}}_N^n| \approx 2^{-N} |\mathcal{P}_N^n|$ and consequently from (8),

$$\langle |a_n|^2 \rangle \approx 2^{-n} \sum_{N=0}^n |\mathcal{P}_N^n| = 2^{-n} |\mathcal{P}^n|. \quad (15)$$

For binary de Bruijn graphs $|\mathcal{P}^n| = 2^{n-1}$ [20] and hence $\langle |a_n|^2 \rangle \approx 1/2$, which is the result predicted from the diagonal contribution [19, 20]. Figure 3 shows the convergence of the variance to the asymptotic result for the family of binary de Bruijn graphs.

JH would like to thank Mark Pollicott and the University of Warwick for their hospitality during his sabbatical.




FIG. 3. Variance of coefficients of the characteristic polynomial of binary de Bruijn quantum graphs showing convergence to the asymptotic value $1/2$.

JH was supported by the Baylor University research leave program. This work was supported by a grant from the Simons Foundation (354583 to Jonathan Harrison).

* jon_harrison@baylor.edu

† vhudgins@udallas.edu

- [1] M. G. Gutzwiller, *Chaos in Classical and Quantum Mechanics*, edited by F. John, Interdisciplinary Applied Mathematics Vol. 1 (Springer-Verlag, 1990).
- [2] M. V. Berry, Proc. R. Soc. A **400**, 229 (1985).
- [3] J. Hannay and A. M. Ozorio de Almeida, J. Phys. A: Math. Gen. **17**, 3429 (1984).
- [4] M. Sieber, J. Phys. A: Math. Gen. **35**, L613 (2002).
- [5] K. Richter and M. Sieber, Phys. Rev. Lett. **89**, 206801 (2002).
- [6] S. Müller, S. Heusler, P. Braun, F. Haake, and A. Altland, Phys. Rev. Lett. **93**, 014103 (2004).
- [7] S. Müller, S. Heusler, P. Braun, F. Haake, and A. Altland, Phys. Rev. E **72**, 046207 (2005).
- [8] G. Berkolaiko and P. Kuchment, *Introduction to Quantum Graphs*, Mathematical Surveys and Monographs, Vol. 186 (AMS, 2013).
- [9] T. Kottos and U. Smilansky, Phys. Rev. Lett. **79**, 4794 (1997).
- [10] T. Kottos and U. Smilansky, Ann. Phys. **274**, 76 (1999).
- [11] O. Hul, S. Bauch, P. Pakoński, N. Savytsky, K. Życzkowski, and L. Sirko, Phys. Rev. E **69**, 056205 (2004).
- [12] A. Rehemanjiang, M. Allgaier, C. H. Joyner, S. Müller, M. Sieber, U. Kuhl, and H.-J. Stöckmann, Phys. Rev. Lett. **117**, 064101 (2016).
- [13] G. Tanner, J. Phys. A: Math. Gen. **34**, 8485 (2001).
- [14] S. Gnutzmann and U. Smilansky, Adv. Phys. **55**, 527 (2006).
- [15] R. Band, J. M. Harrison, and C. H. Joyner, J. Phys. A: Math. Theor. **45**, 325204 (2012).
- [16] H. T. Faal, Int. J. Found. Comput. S. **30**, 683 (2019).
- [17] J. M. Harrison and T. Hudgins, arXiv:2011.05213.
- [18] G. Tanner, J. Phys. A: Math. Gen. **33**, 3567 (2000).
- [19] G. Tanner, J. Phys. A: Math. Gen. **35**, 5985 (2002).
- [20] R. Band, J. M. Harrison, and S. Sepanski, J. Math. Anal. Appl. **470**, 135 (2019).