

THE SIZE, MULTIPARTITE RAMSEY NUMBERS FOR nK_2 VERSUS PATH-PATH AND CYCLE

YASER ROWSHAN¹ AND MOSTAFA GHOLAMI¹

ABSTRACT. For given graphs G_1, G_2, \dots, G_n and any integer j , the size of the multipartite Ramsey number $m_j(G_1, G_2, \dots, G_n)$ is the smallest positive integer t such that any n -coloring of the edges of $K_{j \times t}$ contains a monochromatic copy of G_i in color i for some i , $1 \leq i \leq n$, where $K_{j \times t}$ denotes the complete multipartite graph having j classes with t vertices per each class. In this paper we compute the size of the multipartite Ramsey number $m_j(K_{1,2}, P_4, nK_2)$ for any $j, n \geq 2$ and $m_j(nK_2, C_7)$, for any $j \leq 4$ and $n \geq 2$.

1. INTRODUCTION

In this paper, we are only concerned with undirected, simple and finite graphs. We follow [1] for terminology and notations not defined here. For a given graph G , we denote its vertex set, edge set, maximum degree and minimum degree by $V(G)$, $E(G)$, $\Delta(G)$ and $\delta(G)$, respectively. For a vertex $v \in V(G)$, we use $\deg_G(v)$ and $N_G(v)$ to denote the degree and neighbours of v in G , respectively. The neighbourhood of a vertex $v \in V(G)$ are denoted by $N_G(v) = \{u \in V(G) \mid uv \in E(G)\}$ and $N_{X_j}(v) = \{u \in V(X_j) \mid uv \in E(G)\}$.

As usual, a cycle and a path on n vertices are denoted by C_n and P_n , respectively. Also by a stripe mK_2 we mean a graph on $2m$ vertices and m independent edges. As usual the complete multipartite graph with the partite set (X_1, X_2, \dots, X_j) , $|X_i| = s$ for $i = 1, 2, \dots, j$, denoted by $K_{j \times s}$. We use $[X_i, X_j]$ to denote the set of edges between partite sets X_i and X_j . The complement of a graph G , denoted by \overline{G} , is a graph with the same vertices as G and contains those edges which are not in G . Let $T \subseteq V(G)$ be any subset of vertices of G . Then the induced subgraph $G[T]$ is the graph whose vertex set is T and whose edge set consists of all of the edges in $E(G)$ that have both endpoints in T .

For given graphs G_1, G_2, \dots, G_n and integer j , the size of the multipartite Ramsey number $m_j(G_1, G_2, \dots, G_n)$ is the smallest integer t such that any n -coloring of the edges of $K_{j \times t}$ contains a monochromatic copy of G_i in color i for some i , $1 \leq i \leq n$, where $K_{j \times t}$ denotes the complete multipartite graph having j classes with t vertices per each class. G is n -colorable to (G_1, G_2, \dots, G_n) if there exist a t -edges decomposition of G say (H_1, H_2, \dots, H_n) where $G_i \not\subseteq H_i$ for each $i = 1, 2, \dots, n$.

The existence of such a positive integer is guaranteed by a result in [4]. The size of the multipartite Ramsey numbers of small paths versus certain classes of graphs have been studied in [5, 6, 9]. The size of the multipartite Ramsey numbers of stars versus certain classes of graphs have been studied in [7, 8]. In [2, 3], Burger, Stipp, Vuuren, and Grobler, investigated the multipartite ramsey numbers $m_j(G_1, G_2)$, where G_1 and G_2 are complete balanced multipartite graph, which can be naturally extended to several colors. Recently the numbers $m_j(G_1, G_2)$ have been investigated for special classes: stripes versus cycles, stars versus cycles, see [5] and its references.

In [7], Lusiani et al. determined the size multipartite Ramsey numbers of $m_j(K_{1,m}, H)$, for $j = 2, 3$

2010 *Mathematics Subject Classification.* MSC 05D10, MSC 05C55.

Key words and phrases. Ramsey numbers, Multipartite Ramsey numbers, Stripes, Paths, Cycle.

where H is a path or a cycle on n vertices, and $K_{1,m}$ is a star of order $m+1$. In this paper we compute the size of the multipartite Ramsey number $m_j(K_{1,2}, P_4, nK_2)$ for $n, j \geq 2$ and $m_j(nK_2, C_7)$, for $j \leq 4$ and $n \geq 2$ which are new results of multipartite Ramsey numbers. In particular, as the first target of this work we prove the following theorem:

Theorem 1.1. $m_j(K_{1,2}, P_4, nK_2) = \lfloor \frac{2n}{j} \rfloor + 1$ where $j, n \geq 2$.

In [5], Jayawardene, et al. determined the size multipartite Ramsey numbers $m_j(nK_2, C_m)$ where $j \geq 2$ and $m \in \{3, 4, 5, 6\}$. The second goal of this work generalizes these results, as stated below.

Theorem 1.2. Let $j \in \{2, 3, 4\}$ and $n \geq 2$. Then

$$m_j(nK_2, C_7) = \begin{cases} \infty & j = 2, n \geq 2, \\ 2 & (j, n) = (4, 2), \\ 3 & (j, n) = (3, 2), (4, 3), \\ n & j = 3, n \geq 3, \\ \lceil \frac{n+1}{2} \rceil & j = 4, n \geq 4. \end{cases}$$

2. PROOF OF THEOREM 1.1

In order to simplify the comprehension, let us split the proof of Theorem 1.1 into small parts. We begin with a simple but very useful general lower bound in the following theorem:

Theorem 2.1. $m_j(K_{1,2}, P_4, nK_2) \geq \lfloor \frac{2n}{j} \rfloor + 1$ where $j, n \geq 2$.

Proof. Consider $G = K_{j \times t}$ where $t = \lfloor \frac{2n}{j} \rfloor$ with partition sets X_i , $X_i = \{x_1^i, x_2^i, \dots, x_t^i\}$ for $i \in \{1, 2, \dots, j\}$. Consider $x_1^1 \in X_1$, decompose the edges of $K_{j \times t}$ into graphs G_1, G_2 , and G_3 , where G_1 is a null graph and $G_2 = \overline{G_3}$, where G_3 is $G[X_1 \setminus \{x_1^1\}, X_2, \dots, X_j]$. In fact G_2 is isomorphic to $K_{1, (j-1)t}$ and:

$$E(G_2) = \{x_1^1 x_i^r \mid r = 2, 3, \dots, j \text{ and } i = 1, 2, \dots, t\}$$

Clearly $E(G_t) \cap E(G_{t'}) = \emptyset$, $E(G) = E(G_1) \cup E(G_2) \cup E(G_3)$, $K_{1,2} \not\subseteq G_1$ and $P_4 \not\subseteq G_2$. Since $|V(K_{j \times t})| = j \times \lfloor \frac{2n}{j} \rfloor \leq 2n$, we have $|V(G_3)| \leq 2n - 1$, that is, $nK_2 \not\subseteq G_3$, which means that $m_3(K_{1,2}, P_4, nK_2) \geq \lfloor \frac{2n}{j} \rfloor + 1$ and the proof is complete. ■

Observation 2.1.1. Let $G = K_{2,3}$ (or $K_4 - e$). For any subgraph of G , say H , either H has a subgraph isomorphism to $K_{1,2}$ or \overline{H} has a subgraph isomorphism to P_4 .

Proof. Let $H \subseteq G = K_{2,3}$, for $G = K_4 - e$ the proof is same. Without loss of generality (W.l.g) let $X = \{x_1, x_2\}$ and $Y = \{y_1, y_2, y_3\}$ be a partition set of $V(G)$ and P be a maximum path in H . If $|P| \geq 3$, then H has a subgraph isomorphic to $K_{1,2}$, so let $|P| \leq 2$. If $|P| = 1$, then $\overline{H} (= G)$ has a subgraph isomorphic to P_4 . Hence we may assume that $|P| = 2$, w.l.g let $P = x_1 y_1$. Since $|P| = 2$, $x_1 y_2, x_1 y_3$ and $x_2 y_1$ are in $E(\overline{H})$ and there is at least one edge of $\{x_2 y_2, x_2 y_3\}$ in \overline{H} , in any case, $P_4 \subseteq \overline{H}$ and the proof is complete. ■

We determined the exact value of the multipartite Ramsey number of $m_2(K_{1,2}, P_4, nK_2)$ for $n \geq 2$ in the following lemma:

Lemma 2.2. $m_2(K_{1,2}, P_4, nK_2) = n + 1$ for $n \geq 2$.

Proof. Let $X = \{x_1, x_2, \dots, x_{n+1}\}$ and $Y = \{y_1, y_2, \dots, y_{n+1}\}$ be a partition set of $G = K_{n+1, n+1}$. Consider a 3-edge coloring G^r, G^b and G^g of G . By Theorem 2.1 the lower bound holds. Now let M be the maximum matching in G^g . If $|M| \geq n$, then the lemma holds, so let $|M| \leq n-1$. If $|M| \leq n-2$, then we have $K_{3,3} \subseteq \overline{G^g}$ and by Observation 2.1.1 the lemma holds, so let $|M| = n-1$. W.l.g we may assume that $M = \{x_1y_1, x_2y_2, \dots, x_{n-1}y_{n-1}\}$. By considering the edges between $\{x_n, x_{n+1}\}$ and $Y \setminus \{y_n, y_{n+1}\}$ and the edges between $\{y_n, y_{n+1}\}$ and $X \setminus \{x_n, x_{n+1}\}$, we have $K_{3,2} \subseteq G^r \cup G^b$. Hence by Observation 2.1.1 the lemma holds. ■

In the next two lemmas, we consider $m_3(K_{1,2}, P_4, nK_2)$ for certain values of n . In particular, we prove that $m_3(K_{1,2}, P_4, nK_2) = n$ for $n = 2, 3$ in Lemma 2.3 and $m_3(K_{1,2}, P_4, 4K_2) = 3$ in Lemma 2.4.

Lemma 2.3. $m_3(K_{1,2}, P_4, nK_2) = n$ for $n = 2, 3$.

Proof. Let $X_i = \{x_1^i, x_2^i, \dots, x_n^i\}$ for $i \in \{1, 2, 3\}$ be a partition set of $G = K_{3 \times n}$. Consider a 3-edge coloring G^r, G^b and G^g of G . By Theorem 2.1 the lower bound holds. Now let M be the maximum matching in G^g and consider the following cases:

Case 1: $n = 2$. If $|M| \geq 2$ then $nK_2 \subseteq G^g$ and the proof is complete. So let $|E(M)| \leq 1$. W.l.g we may assume that $x_1^1x_1^2 \in E(M)$, hence we have $K_4 - e \cong G[x_2^1, x_2^2, X_3] \subseteq G^r \cup G^b$ and by Observation 2.1.1 the proof is complete.

Case 2: $n = 3$. In this case, if $|E(M)| \leq 1$ or $|E(M)| \geq 3$, then the proof is same as case 1. So let $|E(M)| = 2$ and w.l.g we may assume that $E(M) = \{e_1, e_2\}$. Considering any e_1 and e_2 in $E(G)$. In any case we have $G^r \cup G^b$ has a subgraph isomorphic to $K_{3,2}$, hence by Observation 2.1.1 the lemma holds. Therefore we have $m_3(K_{1,2}, P_4, 3K_2) = 3$. Now by cases 1 and 2, the proof is complete. ■

Lemma 2.4. $m_3(K_{1,2}, P_4, 4K_2) = 3$.

Proof. Let $X_i = \{x_1^i, x_2^i, x_3^i\}$ for $i \in \{1, 2, 3\}$ be a partition set of $G = K_{3 \times 3}$. By Theorem 2.1 the lower bound holds. Consider a 3-edge coloring (G^r, G^b, G^g) of G where $4K_2 \not\subseteq G^g$. Let M be a maximum matching in G^g , if $|M| \leq 2$ then the proof is same as Lemma 2.3. Hence we may assume that $|M| = 3$ and w.l.g let $E(M) = \{e_1, e_2, e_3\}$. By Observation 2.1.1 there is at least one edge between X_1 and X_2 in G^g , say $e_1 = x_1^1x_1^2$, similarly there is at least one edge between X_3 and $\{x_2^1, x_3^1\}$ in G^g , say $e_2 = x_2^1x_3^1$, otherwise $K_{3,2} \subseteq G^r \cup G^b$ and the proof is complete. Now by Observation 2.1.1 there is at least one edge between $\{x_3^1, x_2^3, x_3^3\}$ and $\{x_2^2, x_3^2\}$ in G^g , let e_3 be this edge. If $x_3^1 \notin V(e_3)$ (say $e_3 = x_2^2x_3^2$), then $K_3 \subseteq G^r \cup G^b[x_3^1, x_3^2, x_3^3]$. Now consider the vertex x_1^1 and x_1^2 , since $|M| = 3$ and $e_1 = x_1^1x_1^2$, it is easy to check that $x_1^1x_3^3, x_1^2x_3^3 \in E(G^g)$ and $x_1^1x_3^2, x_1^2x_3^2 \in E(\overline{G^g})$, otherwise $K_4 - e \subseteq \overline{G^g}$ and the proof is complete. Similarly we have $x_2^1x_3^3, x_3^1x_3^3 \in E(G^g)$ and $x_2^1x_3^2, x_3^1x_3^2 \in E(\overline{G^g})$. Now by considering the edges of $G[X_1, x_1^2, x_2^3, x_3^3]$ it is easy to check that $K_4 - e \subseteq G^r \cup G^b$ and the lemma holds. Hence we have $x_3^1 \in V(e_3)$ (say $e_3 = x_3^1x_2^2$), in this case we have $K_{2,2} \cong G[x_2^2, x_2^3, x_3^2, x_3^3] \subseteq G^r \cup G^b$, otherwise if there exist at least one edge between $\{x_2^3, x_3^3\}$ and $\{x_2^2, x_3^2\}$ in G^g , say e , set $e = e_3$ and the proof is same. Hence by considering the vertex x_1^1 and x_1^2 , since $|M| = 3$ and $e_1 = x_1^1x_1^2$, it is easy to check that $K_{3,2} \subseteq G^r \cup G^b$ and by Observation 2.1.1 the proof is complete. ■

Theorem 2.5. $m_3(K_{1,2}, P_4, nK_2) \leq \lfloor \frac{2n}{3} \rfloor + 1$ for each $n \geq 2$.

Proof. Let $X_i = \{x_1^i, x_2^i, \dots, x_t^i\}$ for $i \in \{1, 2, 3\}$ be a partition sets of $G = K_{3 \times t}$ where $t = \lfloor \frac{2n}{3} \rfloor + 1$. We will prove this theorem by induction. For the base step of the induction, since $\lfloor \frac{2 \times 2}{3} \rfloor + 1 = 2$, $\lfloor \frac{2 \times 3}{3} \rfloor + 1 = 3$ and $\lfloor \frac{2 \times 4}{3} \rfloor + 1 = 3$, theorem holds by Lemmas 2.3 and 2.4. Suppose that $n \geq 5$ and

$m_3(K_{1,2}, P_4, n'K_2) \leq \lfloor \frac{2n'}{3} \rfloor + 1$ for each $n' < n$. We will show that $m_3(K_{1,2}, P_4, nK_2) \leq \lfloor \frac{2n}{3} \rfloor + 1$. By contradiction, we may assume that $m_3(K_{1,2}, P_4, nK_2) > \lfloor \frac{2n}{3} \rfloor + 1$, that is, $K_{3 \times (\lfloor \frac{2n}{3} \rfloor + 1)}$ is 3-colorable to $(K_{1,2}, P_4, nK_2)$. Consider a 3-edge coloring (G^r, G^b, G^g) of G , such that $K_{1,2} \not\subseteq G^r$, $P_4 \not\subseteq G^b$ and $nK_2 \not\subseteq G^g$. By the induction hypothesis and Theorem 2.1 we have $m_3(K_{1,2}, P_4, (n-1)K_2) = \lfloor \frac{2(n-1)}{3} \rfloor + 1 \leq \lfloor \frac{2n}{3} \rfloor + 1$. Therefore since $K_{1,2} \not\subseteq G^r$ and $P_4 \not\subseteq G^b$ we have $(n-1)K_2 \subseteq G^g$. Now we have the following cases:

Case 1: $\lfloor \frac{2n}{3} \rfloor = \lfloor \frac{2(n-1)}{3} \rfloor + 1$.

Since $\lfloor \frac{2n}{3} \rfloor = \lfloor \frac{2(n-1)}{3} \rfloor + 1$ we have a copy of $H = K_{3 \times (\lfloor \frac{2(n-1)}{3} \rfloor + 1)}$ in G . In other words, for each $i \in \{1, 2, 3\}$, there is a vertex, say $x \in X_i$, such that $x \in V(G) \setminus V(H)$. W.l.g we may assume that $A = \{x_1^1, x_1^2, x_1^3\}$ be this vertices. Since $H \subseteq G$, we have $K_{1,2} \not\subseteq G^r[V(H)]$ and $P_4 \not\subseteq G^b[V(H)]$. Hence by the induction hypothesis, we have $M = (n-1)K_2 \subseteq G^g[V(H)] \subseteq G^g$. We consider that the three vertices do not belong to $V(H)$, i.e., A . Since $nK_2 \not\subseteq G^g$, we have $G[A] \subseteq G^r \cup G^b$. Now we consider the following Claim:

Claim 2.6. $n \in B \cup D$ where $B = \{3k \mid k = 1, 2, \dots\}$ and $D = \{3k + 2 \mid k = 1, 2, \dots\}$.

Proof. By contradiction we may assume that $n \notin B \cup D$. In other words, let $n = 3k + 1$, then we have

$$\begin{aligned} 2k &= \lfloor \frac{6k}{3} \rfloor = \lfloor \frac{6k}{3} + \frac{2}{3} \rfloor = \lfloor \frac{6k+2}{3} \rfloor = \lfloor \frac{2(3k+1)}{3} \rfloor \\ &= \lfloor \frac{2n}{3} \rfloor = \lfloor \frac{2(n-1)}{3} \rfloor + 1 = \lfloor \frac{2(3k)}{3} \rfloor + 1 = 2k + 1, \end{aligned}$$

which is a contradiction implying that $n \in B \cup D$.

Claim 2.7. *There is at least one vertex in $V(H) \setminus V(M)$.*

Proof. Let $M = (n-1)K_2 \subseteq G^g$, Then $|V(M)| = 2(n-1) = 2n - 2$. Since $\lfloor \frac{2n}{3} \rfloor = \lfloor \frac{2(n-1)}{3} \rfloor + 1$, by Claim 1, if $n \in B$ we have $n = 3k$ for $k \geq 2$. Now we have

$$\lfloor \frac{2(n-1)}{3} \rfloor + 1 = \lfloor \frac{2(3k-1)}{3} \rfloor + 1 = \lfloor \frac{2(3k)}{3} - \frac{2}{3} \rfloor + 1 = 2k - 1 + 1 = 2k.$$

Hence we have $|V(H)| = 3 \times (2k) = 6k = 2n$ and thus $|V(H) \setminus V(M)| = 2$. If $n \in D$ then we have

$$\lfloor \frac{2(n-1)}{3} \rfloor + 1 = \lfloor \frac{2(3k+1)}{3} \rfloor + 1 = \lfloor \frac{2(3k)}{3} + \frac{2}{3} \rfloor + 1 = 2k + 1.$$

Hence $|V(H)| = 3 \times (2k + 1) = 6k + 3 = 2n - 1$. Therefore $|V(H) \setminus V(M)| = 1$.

Now by Claim 2.7, let $x \in V(H) \setminus V(M)$. Since $nK_2 \not\subseteq G^g$ we have $K_4 - e \cong G[A \cup \{x\}] \subseteq G^r \cup G^b$. Hence by Observation 2.1.1 again a contradiction.

Case 2: $\lfloor \frac{2n}{3} \rfloor = \lfloor \frac{2(n-1)}{3} \rfloor$.

In this case, by Claim 2.6 we have $n = 3k + 1$. Since $K_{1,2} \not\subseteq G^r$ and $P_4 \not\subseteq G^b$, by the induction hypothesis we have $M = (n-1)K_2 \subseteq G^g$. Now we have the following claim:

Claim 2.8. $|V(G) \setminus V(M)| = 3$.

Proof. Let $M = (n-1)K_2 \subseteq G^g$. Since $|V(X_j)| = \lfloor \frac{2n}{3} \rfloor + 1$ and $n = 3k + 1$, we have $\lfloor \frac{2n}{3} \rfloor + 1 = \lfloor \frac{2(3k+1)}{3} \rfloor + 1 = \lfloor \frac{6k}{3} + \frac{2}{3} \rfloor + 1 = 2k + 1$ and, therefore, $|V(G)| = 3 \times (2k + 1) = 6k + 3 = 2(3k + 1) + 1 = 2n + 1$, that is, $|V(G) \setminus V(M)| = (2n + 1) - (2n - 2) = 3$.

By Claim 2.8 we have $|V(G) \setminus V(M)| = 3$. W.l.g we may assume that $A' = \{x, y, z\}$ is this vertices, since $nK_2 \not\subseteq G^g$ we have $G[A'] \subseteq G^r \cup G^b$. We consider the three vertices belonging to A' , now we have the following subcases:

Subcase 2-1: $A' \subseteq X_j$ for only one $j \in \{1, 2, 3\}$. W.l.g we may assume that $A' \subseteq X_1$ and $E(M) = \{e_i \mid i = 1, 2, \dots, (n-1)\}$. Since $k \geq 2$ and $3k + 1 = n \geq 7$ we have $|X_j| \geq 5$ and $|E(M) \cap E(G[X_2, X_3])| \geq 3$, otherwise, $K_{3,3} \subseteq G^r \cup G^b$ and by Observation 2.1.1; a contradiction. W.l.g we may assume that $\{x_i^2 x_i^3 \mid i = 1, 2, 3\} \subseteq (E(M) \cap E(G^g[X_2, X_3]))$. Consider $G' = G[A', x_1^2, x_2^2, x_3^2, x_1^3, x_2^3, x_3^3] \cong K_{3 \times 3}$. Since $nK_2 \not\subseteq G^g$, if M' is a maximum matching in G'^g , then $|M'| \leq 3$, otherwise we have $nK_2 = M \setminus \{e_1, e_2, e_3\} \cup M' \subseteq G^g$; a contradiction again. Since $m_3(K_{1,2}, P_4, 4K_2) = 3$ and $|M'| \leq 3$, we have $K_{1,2} \subseteq G'^r \subseteq G^r$ or $P_4 \subseteq G'^b \subseteq G^b$; also a contradiction.

Subcase 2-2: $|A' \cap X_j| = 1$ for each $j \in \{1, 2, 3\}$. W.l.g we may assume that $x \in X_1, y \in X_2$ and $z \in X_3$. Hence $G[A'] \cong K_3 \subseteq G^r \cup G^b$. Since $|X_j| \geq 5$ we have $|E(M) \cap E(G^g[X_i, X_j])| \geq 2$ for each $i, j \in \{1, 2, 3\}$. W.l.g we may assume that $x'y' \in E(M) \cap E(G^g[X_1 \setminus \{x\}, X_2 \setminus \{y\}])$, $x' \in X_1$ and $y' \in X_2$. If $x'y$ and $x'z \in E(G^r \cup G^b)$ then we have $K_4 - e \subseteq G^r \cup G^b$ and by Observation 2.1.1; a contradiction. So let $x'y$ or $x'z \in E(G^g)$. If $x'y \in E(G^g)$, then, since $nK_2 \not\subseteq G^g$, we have $y'x, y'z \in E(G^r \cup G^b)$, that is, $K_4 - e \subseteq G^r \cup G^b$; a contradiction again. So let $x'z \in E(G^g)$ and $x'y \in E(G^r \cup G^b)$. Since $nK_2 \not\subseteq G^g$, we have $y'x \in E(G^r \cup G^b)$. If $|E(G^r) \cap E(G[A'])| \neq 0$ then we have $P_4 \subseteq G^b$. So let $xy, yz, zx \in E(G^b)$ and $xy', yx' \in E(G^r)$. Since $|E(M) \cap E(G^g[X_i, X_j])| \geq 2$ there is at least one edge, say $y''z'' \in E(M) \cap E(G^g[X_2 \setminus \{y\}, X_3 \setminus \{z\}])$. W.l.g we may assume that $y'' \in X_2$ and $z'' \in X_3$. Since $K_{1,2} \not\subseteq G^r$ and $P_4 \not\subseteq G^b$ we have $y''x, z''y \in E(G^g)$. Hence we have $nK_2 = M \setminus \{y''z''\} \cup \{y''x, z''y\}$; a contradiction.

Subcase 2-3: $|A' \cap X_j| = 2$ for only one $j \in \{1, 2, 3\}$. W.l.g we may assume that $x, y \in X_1$ and $z \in X_2$. Hence we have $G'[A'] \cong P_3 \subseteq G^r \cup G^b$. Since $k \geq 2$ we have $|X_j| \geq 5$, that is, $|E(M) \cap E(G^g[X_2, X_3])| \geq 3$. W.l.g we may assume that $vu, v'u' \in E(M) \cap E(G^g[X_2, X_3])$ where $v, v' \in X_2$ and $u, u' \in X_3$. Now we have the following claim:

Claim 2.9. $|N_{G^g}(x) \cap \{v, v'\}| = |N_{G^g}(y) \cap \{v, v'\}| = 0$.

Proof. By contradiction, w.l.g we may assume that $xv \in E(G^g)$. Since $nK_2 \not\subseteq G^g$, we have $yu, zu \in E(G^r \cup G^b)$. Consider $A'' = \{y, z, u\}$ and $M' = M \setminus \{vu\} \cup \{xv\}$. Hence $M' = (n-1)K_2 \subseteq G^g$ and $|A'' \cap X_j| \neq 0$ for each $j \in \{1, 2, 3\}$; a contradiction to subcase 2-2.

Now by Claim 2.9 we have $K_{2,3} = G[A' \cup \{v, v'\}] \subseteq G^r \cup G^b$. In this case by Observation 2.1.1 we have $K_{1,2} \subseteq G^r$ or $P_4 \subseteq G^b$; a contradiction again.

Therefore by Cases 1 and 2 we have $m_3(K_{1,2}, P_4, nK_2) \leq \lfloor \frac{2n}{3} \rfloor + 1$ for $n \geq 2$. ■

Now by Theorems 2.1 and 2.5 we have the following theorem:

Theorem 2.10. $m_3(K_{1,2}, P_4, nK_2) = \lfloor \frac{2n}{3} \rfloor + 1$ for $n \geq 2$.

In the next two theorems, we consider $m_j(K_{1,2}, P_4, nK_2)$ for each values of $n \geq 2$ and $j \geq 4$. In particular, we prove that $m_j(K_{1,2}, P_4, nK_2) = \lfloor \frac{2n}{j} \rfloor + 1$ for $n \geq 2$ and $j \geq 4$. We start with the following theorem:

Theorem 2.11. Let $j \geq 4$ and $n \geq 2$. Given that $m_j(K_{1,2}, P_4, (n-1)K_2) = \lfloor \frac{2(n-1)}{j} \rfloor + 1$, it follows that $m_j(K_{1,2}, P_4, nK_2) \leq \lfloor \frac{2n}{j} \rfloor + 1$.

Proof. Let $j \geq 4$ and $n \geq 2$. For $i \in \{1, 2, \dots, j\}$ let $X_i = \{x_1^i, x_2^i, \dots, x_j^i\}$ be partition set of $G = K_{j \times t}$ where $t = \lfloor \frac{2n}{j} \rfloor + 1$. Assume that $m_j(K_{1,2}, P_4, (n-1)K_2) = \lfloor \frac{2(n-1)}{j} \rfloor + 1$ is true. To prove $m_j(K_{1,2}, P_4, nK_2) \leq \lfloor \frac{2n}{j} \rfloor + 1$. Consider a 3-edge coloring (G^r, G^b, G^g) of G . Suppose that $nK_2 \not\subseteq G^g$, we prove that $K_{1,2} \subseteq G^r$ or $P_4 \subseteq G^b$. Let M^* be the maximum matching in G^g . Hence by the assumption, $|M^*| \leq n-1$, that is $|V(K_{j \times t}) \cap V(M^*)| \leq 2(n-1)$. Now, we have the following claim:

Claim 2.12. $|V(K_{j \times t}) \setminus V(M^*)| \geq 3$.

Proof. Consider the following cases:

Case 1: Let $2n = jk$ ($2n \equiv 0 \pmod{j}$). In this case, we have

$$|V(G)| = j \times t = j \times (\lfloor \frac{2n}{j} \rfloor + 1) = j \times \lfloor \frac{2n}{j} \rfloor + j = jk + j = j(k+1).$$

Hence

$$|V(G) \setminus V(M^*)| \geq j(k+1) - 2(n-1) = jk + j - 2n + 2 = j + 2 \geq 6 \quad (j \geq 4).$$

Case 2: Let $2n = jk + r$ ($2n \equiv r \pmod{j}$ where $r \in \{1, 2, \dots, j-1\}$). In this case, we have $|V(G)| = j \times (\lfloor \frac{2n}{j} \rfloor + 1) = j \times (\lfloor \frac{jk+r}{j} \rfloor + 1) = j \times (\lfloor \frac{jk}{j} \rfloor + \lfloor \frac{r}{j} \rfloor + 1) = j \times \lfloor \frac{jk}{j} \rfloor + j = jk + j$. Hence we have

$$|V(G) \setminus V(M^*)| \geq j(k+1) - 2(n-1) = jk + j - 2n + 2 = jk + j - jk - r + 2 = j - r + 2 \geq 3.$$

By Claim 2.12, G contains three vertices, say x, y and z in $V(K_{j \times t}) \setminus V(M^*)$. Consider the vertex set $\{x, y, z\}$ and let $\{x, y, z\} \subseteq A = V(G) \setminus V(M^*)$. Now we have the following cases:

Case 1: Let $x \in X_1, y \in X_2$ and $z \in X_3$ where X_i for $i = 1, 2, 3$ are distinct partition sets of $G = K_{j \times t}$. Note that all vertices of A are adjacent to each other in $\overline{G^g}$. Since $t \geq 2$ we have $|X_i| \geq 2$. Consider the partition X_j for $j \geq 4$. Since $|X_j| \geq 2$, if $|A \cap X_j| \geq 1$ for at least one $j \geq 4$, then we have $K_4 \subseteq G^r \cup G^b$ and the proof is complete by Observation 2.1.1. Now let $|A \cap X_j| = 0$ for each $j \geq 4$. Hence for $x_1^4 \in X_4$ there exists a vertex, say u such that $x_1^4 u \in E(M^*)$. Consider $N_{G^g}(x_1^4) \cap \{x, y, z\}$. If $|N_{G^g}(x_1^4) \cap \{x, y, z\}| \leq 1$, then we have $K_4 - e \subseteq G^r \cup G^b$ and, by Observation 2.1.1, the proof is complete. Therefore let $|N_{G^g}(x_1^4) \cap \{x, y, z\}| \geq 2$. W.l.g we may assume that $\{x, y\} \subseteq N_{G^g}(x_1^4) \cap \{x, y, z\}$. In this case, we have $|N_{G^g}(u) \cap \{x, y, z\}| = 0$. On the contrary, let $xu \in E(G^g)$ and set $M' = M^* \setminus \{x_1^4 u\} \cup \{x_1^4 y, ux\}$. Clearly M' is a matching where $|M'| > |M^*|$, which contradicts the maximality of M^* . Hence we have $|N_{G^g}(u) \cap \{x, y, z\}| = 0$. Therefore we have $K_4 - e \subseteq G^r \cup G^b[x, y, z, u]$ and, by Observation 2.1.1, the proof is complete.

Case 2: Let $x, y \in X_i$ and $z \in X_{i'}$ where $X_i, X_{i'}$ are distinct partition sets of G . W.l.g let $i = 1$ and $i' = 2$. Consider the partition X_j ($j \neq 1, 2$). Since $|X_j| \geq 2$, if $|A \cap X_j| \geq 1$, then we have $K_4 - e \subseteq G^r \cup G^b$ and, by Observation 2.1.1, the proof is complete. So let $|A \cap X_j| = 0$ for each $j \geq 3$. Now, we have the following claim.

Claim 2.13. Let $e = v_1 v_2 \in E(M^*)$, and w.l.g let $|N_{G^g}(v_1) \cap \{x, y, z\}| \geq |N_{G^g}(v_2) \cap \{x, y, z\}|$. If $|N_{G^g}(v_1) \cap \{x, y, z\}| \geq 2$, then $|N_{G^g}(v_2) \cap \{x, y, z\}| = 0$. If $|N_{G^g}(v_1) \cap \{x, y, z\}| = |N_{G^g}(v_2) \cap \{x, y, z\}| = 1$, then v_1, v_2 has the same neighbour in $\{x, y, z\}$.

Proof. Let $|N_{G^g}(v_1) \cap \{x, y, z\}| \geq 2$. W.l.g we may assume that $\{w, w'\} \subseteq N_{G^g}(v_1) \cap \{x, y, z\}$. By contradiction let $|N_{G^g}(v_2) \cap \{x, y, z\}| \neq 0$, w.l.g let $w'' \in N_{G^g}(v_2) \cap \{x, y, z\}$. In this case, we set $M' = (M^* \setminus \{v_1 v_2\}) \cup \{v_1 w, v_2 w''\}$. Clearly M' is a matching with $|M'| > |M^*|$, which contradicts the maximality of M^* . So let $|N_{G^g}(v_i) \cap \{x, y, z\}| = 1$ for $i = 1, 2$, if v_i has a different neighbour then the proof is same.

Claim 2.14. *There is at least one edge, say $e = u_i u_j \in E(M^*)$, such that $u_i, u_j \notin X_1, X_2$.*

Proof. If $|X_j| \geq 3$ then there is at least one edge, say $e = u_i u_j \in E(M^*)$, such that $u_i, u_j \notin X_1, X_2$. Otherwise, we have $K_{3,2} \subseteq G^r \cup G^b[X_j, X_{j'}]$ where $j, j' \geq 3$, hence by Observation 2.1.1; a contradiction. So let $|X_j| = 2$. In this case, if $j \geq 5$ then the proof is same. Now let $j = 4$. We have $|M^*| \leq 2$, that is, $n \leq 3$. Hence there is at least one vertex, say $w \in (X_3 \cup X_4) \cap A$; a contradiction to $|A \cap X_j| = 0$.

By Claim 2.14 there is at least one edge, say $e = u_i u_j \in E(M^*)$, such that $u_i, u_j \notin X_1, X_2$. W.l.g let $e = u_1 u_2 \in E(M^*)$ such that $u_i \notin X_1, X_2$, also w.l.g assume that $|N_{G^g}(u_1) \cap \{x, y, z\}| \geq |N_{G^g}(u_2) \cap \{x, y, z\}|$. If $|N_{G^g}(u_1) \cap \{x, y, z\}| \geq 2$, then by Claim 2.14 we have $|N_{G^g}(u_2) \cap \{x, y, z\}| = 0$. Hence we have $K_4 - e \subseteq G^r \cup G^b$. So let $|N_{G^g}(u_1) \cap \{x, y, z\}| = |N_{G^g}(u_2) \cap \{x, y, z\}| = 1$, in this case, by Claim 2.14, we have $N_{G^g}(u_1) \cap \{x, y, z\} = N_{G^g}(u_2) \cap \{x, y, z\}$, and if x or y is this vertex, then $K_4 - e \subseteq G^r \cup G^b$, otherwise $K_{3,2} \subseteq G^r \cup G^b$. In any case, by Observation 2.1.1, the proof is complete.

Case 3: Let $x, y, z \in X_i$ where X_i is a partition set of $G = K_{j \times t}$, say $i = 1$. If there exists a vertex, say $w \in X_j \cap A$, where $j \neq 1$, then the proof is same as case 2. Hence let $|A \cap X_j| = 0$. Since $|X_j| \geq 3$, there exists an edge, say $e = vu \in E(M^*)$, such that $v, u \notin X_1$. Consider the neighbours of vertices v and u in X_1 . W.l.g let $|N_{G^g}(v) \cap \{x, y, z\}| \geq |N_{G^g}(u) \cap \{x, y, z\}|$. If $|N_{G^g}(v) \cap \{x, y, z\}| = 0$ then we have $K_{3,2} \subseteq G^r \cup G^b$, so let $|N_{G^g}(v) \cap \{x, y, z\}| \geq 1$. In this case, by Claim 2.14, we have $|N_{G^g}(u) \cap \{x, y, z\}| \leq 1$. Hence w.l.g we may assume that yu and zu be in $E(G^r \cup G^b)$ and $x \in N_{G^g}(v)$. Now set $M^{**} = (M^* \setminus \{vu\}) \cup \{vx\}$ and $A' = (A \setminus \{x\}) \cup \{u\}$, the proof is same as case 2 and the proof is complete.

According to the cases 1, 2 and 3 we have $m_j(K_{1,2}, P_4, nK_2) \leq \lfloor \frac{2n}{j} \rfloor + 1$. ■

The results of Theorems 2.1, 2.10 and 2.11 and Lemmas 2.2, 2.3 and 2.4 concludes the proof of Theorem 1.1.

3. PROOF OF THEOREM 1.2

In this section, we investigate the size multipartite Ramsey number $m_j(nK_2, C_7)$ for $j \leq 4$ and $n \geq 2$. In order to simplify the comprehension, let us split the proof of Theorem 1.2 into small parts. For $j = 2$, since the bipartite graph has no odd cycle, we have $m_2(nK_2, C_7) = \infty$. For other cases we start with the following proposition:

Proposition 3.1. $m_3(nK_2, C_7) = 3$ where $n = 2, 3$.

Proof. Clearly $m_3(nK_2, C_7) \geq 3$. Consider $K_{3 \times 3}$ with the partition set $X_i = \{x_1^i, x_2^i, x_3^i\}$ for $i = 1, 2, 3$. Let G be a subgraph of $K_{3 \times 3}$. For $n = 2$ if $2K_2 \subseteq G$, then the proof is complete, so let $2K_2 \not\subseteq G$. In this case we have $K_{3,2,2} \subseteq \overline{G}$, hence $C_7 \subseteq \overline{G}$, that is, $m_3(2K_2, C_7) = 3$. For $n = 3$ by contradiction, we may assume that $m_3(3K_2, C_7) > 3$, that is, $K_{3 \times 3}$ is 2-colorable to $(3K_2, C_7)$, say $3K_2 \not\subseteq G$ and $C_7 \not\subseteq \overline{G}$. Since $m_3(3K_2, C_6) = 3$ [5], and $3K_2 \not\subseteq G$ we have $C_6 \subseteq \overline{G}$. Let $A = V(C_6)$ and $Y_i = A \cap X_i$ for $i = 1, 2, 3$. If there exists $i \in \{1, 2, 3\}$ such that $|Y_i| = 0$, say $i = 1$, then we have $A = X_2 \cup X_3$ and $C_6 \subseteq \overline{G}[X_2, X_3]$. Let $C_6 = w_1 w_2 \dots w_6 w_1$. Since $C_7 \not\subseteq \overline{G}$, for each $x_i \in X_1$ in \overline{G} , x_i cannot be adjacent to w_i and w_{i+1} for $i = 1, 2, \dots, 6$. Hence we have $|N_G(x_i) \cap V(C_6)| \geq 3$ for each $x_i \in X_1$. One can easily check that in any case, we have $3K_2 \subseteq G$; a contradiction, hence let $|Y_i| \geq 1$ for each $i = 1, 2, 3$. Set $B = (|Y_1|, |Y_2|, |Y_3|)$. Now we have the following cases:

Case 1: $B = (3, 2, 1)$. let $A = X_1 \cup \{x_1^2, x_2^2, x_3^2\}$. In this case, we have $C_6 \cong x_1^1 x_1^2 x_2^2 x_2^3 x_3^3 x_3^1 x_1^1$. Consider the vertex set $A' = V(K_{3 \times 3}) \setminus A = \{x_3^2, x_2^3, x_3^3\}$. Since $C_7 \not\subseteq \overline{G}$, we have $|N_{\overline{G}}(x_2^3) \cap \{x_1^1, x_2^2\}| \leq 1$. Hence $|N_G(x_2^3) \cap \{x_1^1, x_2^2\}| \geq 1$. W.l.g let $x_2^3 x_1^1 \in E(G)$. By similarity, we have

$|N_G(x_3^3) \cap \{x_2^1, x_2^2\}| \geq 1$ and $|N_G(x_3^2) \cap \{x_3^1, x_3^3\}| \geq 1$, see Figure 1. In any case, we have $3K_2 \subseteq G$; a contradiction again.




FIGURE 1. $B = (3, 2, 1)$

Case 2: $B = (2, 2, 2)$. W.l.g let $Y_i = \{x_1^i, x_2^i\}$ for $i = 1, 2, 3$. In this case, we have $C_6 \cong w_1 w_2 w_3 w_4 w_5 w_6 w_1$. W.l.g let $w_1 = x_1^1, w_2 = x_2^1$. Since $|Y_3| = 2$ and $w_4 w_5 \in E(C_6)$ we have $|\{w_3, w_6\} \cap Y_3| \geq 1$. If $|\{w_3, w_6\} \cap Y_3| = 2$, then considering Figure 2(a), the proof is same as case 1. So let $|\{w_3, w_6\} \cap Y_3| = 1$. W.l.g let $w_3 = x_1^3, x_2^3 = w_5, x_2^1 = w_4, x_2^2 = w_6$. In this case, consider Figure 2(b) and the proof is same as case 1. Hence, in any case, we have $3K_2 \subseteq G$; again a contradiction.




FIGURE 2. (a) $|\{w_3, w_6\} \cap Y_3| = 2$, (b) $|\{w_3, w_6\} \cap Y_3| = 1$

By cases 1 and 2 we have $3K_2 \subseteq G$. Thus the proof is complete and the proposition holds. \blacksquare

We determined the exact value of the multipartite Ramsey number $m_3(nK_2, C_7)$ for $n \geq 3$ in the following theorem:

Theorem 3.2. *For each $n \geq 3$ we have $m_3(nK_2, C_7) = n$.*

Proof. First we show that $m_3(nK_2, C_7) \geq n$. Consider the coloring given by $K_{3 \times (n-1)} = G^r \cup G^b$ where $G^r \cong K_{n-1, n-1}$ and $G^b \cong K_{n-1, 2(n-1)}$. Since $|V(G^r)| = 2(n-1)$ and G^b is bipartite, we have $nK_2 \not\subseteq G^r$ and $C_7 \not\subseteq G^b$, that is, $m_3(nK_2, C_7) \geq n$. For the upper bound, consider $K_{3 \times n}$ with partite sets $X_i = \{x_1^i, x_2^i, \dots, x_n^i\}$ for $i = 1, 2, 3$. We will prove this by induction. For $n = 3$, by Proposition 3.1, the theorem holds. Suppose that $m_3(nK_2, C_7) \leq n$ for each $n \geq 4$. We will show that $m_3((n+1)K_2, C_7) \leq n+1$ as follows: By contradiction we may assume that $m_3((n+1)K_2, C_7) > n+1$, that is, $K_{3 \times (n+1)}$ is 2-colorable to $((n+1)K_2, C_7)$, say $(n+1)K_2 \not\subseteq G$ and $C_7 \not\subseteq \overline{G}$. Let $X'_i = X_i \setminus \{x_1^i\}$. Hence by the induction hypothesis we have $m_3(nK_2, C_7) \leq n$. Therefore since $|X'_i| = n$ and $C_7 \not\subseteq \overline{G}[X'_1, X'_2, X'_3]$ we have $M = nK_2 \subseteq G[X'_1, X'_2, X'_3]$. If there

exists i and j such that $x_1^i x_1^j \in E(G)$, then we have $(n+1)K_2 \subseteq G$; a contradiction. Hence we have $x_1^i x_1^j \in E(\overline{G})$ for $i, j \in \{1, 2, 3\}$. Let $A = V(K_{3 \times n}) \setminus V(M)$. Hence we have $|A| = 3n - 2n = n$. Since $(n+1)K_2 \not\subseteq G$ we have $G[A, x_1^1, x_1^2, x_1^3] \subseteq \overline{G}$. Since $|A| = n \geq 4$ one can easily check that, in any case, we have $H \subseteq \overline{G}$, where $H \in \{K_{5,1,1}, K_{4,2,1}, K_{3,3,1}, K_{3,2,2}\}$. If $H \in \{K_{3,3,1}, K_{3,2,2}\}$, one can easily check that we have $C_7 \subseteq H \subseteq \overline{G}$; a contradiction again. So let $H \in \{K_{5,1,1}, K_{4,2,1}\}$ and consider the following cases:

Case 1: $A \subseteq X_i$ for only one i , that is, $H = K_{5,1,1}$. W.l.g let $A \subseteq X_1$ and $\{x_2^1, x_3^1, \dots, x_5^1\} \subseteq A$. Then we have $K_{n+1,1,1} \subseteq \overline{G}$ and $M \subseteq G[X_2, X_3]$. Since $n \geq 4$, we have $|M| \geq 4$, that is, there exists at least two edges, say $e_1 = x_1 y_1$ and $e_2 = x_2 y_2$ in $E(M)$, where $\{x_1, x_2, y_1, y_2\} \subseteq X_2 \cup X_3$. W.l.g let $|N_G(x_i) \cap A| \geq |N_G(y_i) \cap A|$ for $i = 1, 2$. One can easily check that $|N_G(y_i) \cap A| \leq 1$, otherwise we have $(n+1)K_2 \subseteq G$; a contradiction. Since $|N_G(y_i) \cap A| \leq 1$ and $|A| \geq 5$, we have $|N_{\overline{G}}(y_i) \cap A| \geq 4$. Hence we have $|N_{\overline{G}}(y_1) \cap N_{\overline{G}}(y_2) \cap A| \geq 3$. W.l.g we may assume that $\{x_1^1, x_2^1, x_3^1\} \subseteq N_{\overline{G}}(y_1) \cap N_{\overline{G}}(y_2) \cap A$. In this case, we have $C_7 \subseteq \overline{G}[x_1^1, x_2^1, x_3^1, x_1^2, x_1^3, y_1, y_2] \subseteq \overline{G}$; a contradiction again.

Case 2: $H = K_{4,2,1}$. W.l.g let $|A \cap X_1| = n-1$ and $|A \cap X_2| = 2$. Let $\{x_2^1, x_3^1, \dots, x_4^1\} \subseteq A \cap X_1$ and $x_2^2 \in A \cap X_2$, that is, we have $K_{4,2,1} \subseteq K_{n,2,1} = G[A, x_1^1, x_2^1, x_3^1] \subseteq \overline{G}$ and $M \subseteq K_{1,n-1,n}$. That is, there exists at least one edge, say $e = xy$, where $x \in X_2$ and $y \in X_3$. W.l.g let $|N_G(x) \cap A| \geq |N_G(y) \cap A|$. One can easily check that $|N_G(y) \cap A| \leq 1$. Hence we have $|N_{\overline{G}}(y) \cap A| \geq 3$ and the proof is same as case 1.

By cases 1 and 2 we have the assumption that $m_3((n+1)K_2, C_7) > n+1$ does not hold. Now we have $m_3(nK_2, C_7) = n$ for each $n \geq 3$. This completes the induction step and the proof. \blacksquare

Lemma 3.3. For $j \geq 3$ and $n \geq j$ we have $m_j(nK_2, C_7) \geq \lceil \frac{2n+2}{j} \rceil$.

Proof. To show that $m_j(nK_2, C_7) \geq \lceil \frac{2n+2}{j} \rceil$, assume that $\lceil \frac{2n+2}{j} \rceil \geq 1$. Consider the coloring given by $K_{j \times t_0} = G^r \cup G^b$ where $t_0 = \lceil \frac{2n+2}{j} \rceil - 1$ such that $G^r \cong K_{(j-1) \times t_0}$ and $G^b \cong K_{t_0, (j-1)t_0}$. Since G^b is bipartite, we have $C_7 \not\subseteq G^b$, and

$$\begin{aligned} |V(G^r)| &= (j-1) \times t_0 = (j-1) \left(\lceil \frac{2n+2}{j} \rceil - 1 \right) = (j-1) \left(\lceil \frac{2n+2}{j} \rceil \right) - (j-1) \\ &\leq (j-1) \left(\frac{2n+2}{j} + 1 \right) - (j-1) = j \times \left(\frac{2n+2}{j} \right) - \frac{2n+2}{j}. \end{aligned}$$

Since $n \geq j$ we have $|V(G^r)| < 2n$. Hence we have $nK_2 \not\subseteq G^r$. Since $K_{j \times t_0} = G^r \cup G^b$ we have $m_j(nK_2, C_7) \geq \lceil \frac{2n+2}{j} \rceil$ for $n \geq j \geq 3$. \blacksquare

Lemma 3.4. $m_4(4K_2, C_7) = 3$.

Proof. By Lemma 3.3 we have $m_4(4K_2, C_7) \geq 3$. For the upper bound consider the coloring given by $K_{4 \times 3} = G^r \cup G^b$ such that $C_7 \not\subseteq G^b$. Since $m_3(3K_2, C_7) = 3$, we have $3K_2 \subseteq G^r[X_1, X_2, X_3] \subseteq G^r$. Let $M = 3K_2$, hence we have $|V(X_1 \cup X_2 \cup X_3) \setminus V(M)| = 3$. W.l.g let $A = \{w_1, w_2, w_3\}$ be this vertices. If $E(G^r) \cap E(G[X_4, A]) \neq \emptyset$ then we have $4K_2 \subseteq G^r$. So let $K_{3,3} \subseteq G[X_4, A] \subseteq G^b$. Consider the edge $e = v_1 v_2 \in E(M)$, it is easy to show that $|N_{G^b}(v_i) \cap X_4| \geq 2$ for some $i \in \{1, 2\}$, otherwise we have $4K_2 \subseteq G^r$. In any case, one can easily check that $C_7 \subseteq G^b$; a contradiction. Thus giving $m_4(4K_2, C_7) = 3$. \blacksquare

Lemma 3.5. For $n \geq 4$ we have $m_4(nK_2, C_7) = \lceil \frac{n+1}{2} \rceil$.

Proof. By Lemma 3.3 we have $m_4(nK_2, C_7) \geq \lceil \frac{n+1}{2} \rceil$. To prove $m_4(nK_2, C_7) \leq \lceil \frac{n+1}{2} \rceil$, consider $K_{4 \times t}$ with partite set $X_i = \{x_1^i, x_2^i, \dots, x_t^i\}$ for $i = 1, 2, 3, 4$, where $t = \lceil \frac{n+1}{2} \rceil$. We will prove this by induction. For $n = 4$ by Lemma 3.4, the lemma holds. Now we consider the following cases:

Case 1: $n = 2k$, where $k \geq 3$. Suppose that $m_4(n'K_2, C_7) \leq \lceil \frac{n'+1}{2} \rceil$ for each $n' < n$. We will show that $m_4(nK_2, C_7) \leq \lceil \frac{n+1}{2} \rceil$ as follows: By contradiction, we may assume that $m_4(nK_2, C_7) > \lceil \frac{n+1}{2} \rceil$, that is, $K_{4 \times t}$ is 2-colorable to (nK_2, C_7) , say $nK_2 \not\subseteq G$ and $C_7 \not\subseteq \overline{G}$. Let $X'_i = X_i \setminus \{x_1^i\}$ for $i = 1, 2, 3, 4$. Hence by the induction hypothesis, we have $m_4((n-1)K_2, C_7) \leq \lceil \frac{n}{2} \rceil = k$. Therefore, since $|X'_i| = k = \frac{n}{2}$ and $C_7 \not\subseteq \overline{G}$ we have $M = (n-1)K_2 \subseteq G[X'_1, X'_2, X'_3, X'_4]$. If there exists $i, j \in \{1, 2, 3, 4\}$, where $x_1^i x_1^j \in E(G)$, then $nK_2 \subseteq G$; a contradiction. Now we have $K_4 \cong \overline{G}[x_1^1, x_1^2, x_1^3, x_1^4] \subseteq \overline{G^c}$. Since $nK_2 \not\subseteq G$ and $\lceil \frac{n+1}{2} \rceil = \lceil \frac{2k+1}{2} \rceil = k+1$, we have $|V(K_{4 \times k}) \setminus V(M)| = 2n - 2(n-1) = 2$, that is, there exists two vertices, say w_1 and w_2 in $V(K_{4 \times k}) \setminus V(M)$. Since $nK_2 \not\subseteq G$ we have $G[S] \subseteq \overline{G}$, where $S = \{x_1^i \mid i = 1, 2, 3, 4\} \cup \{w_1, w_2\}$. Hence we have the following claim:

Claim 3.6. *Let $e = v_1 v_2 \in E(M)$ and w.l.g we may assume that $|N_G(v_1) \cap S| \geq |N_G(v_2) \cap S|$. If $|N_G(v_1) \cap S| \geq 2$ then $|N_G(v_2) \cap S| = 0$. If $|N_G(v_1) \cap S| = 1$ then $|N_G(v_2) \cap S| \leq 1$. If $|N_G(v_i) \cap S| = 1$ then v_1 and v_2 have the same neighbour in S .*

Proof. By contradiction. We may assume that $\{w, w'\} \subseteq N_G(v_1) \cap S$ and $w'' \in N_G(v_2) \cap S$, in this case, we set $M' = (M \setminus \{v_1 v_2\}) \cup \{v_1 w, v_2 w''\}$. Clearly, M' is a matching which $|M'| > |M| = n-1$, with contradicts the $nK_2 \not\subseteq G$. If $|N_G(v_i) \cap S| = 1$ and v_i has a different neighbour then the proof is same.

Since $n \geq 4$ and $|M| \geq 3$. If $\{w_1, w_2\} \subseteq X_i$, say X_1 , then there is at least one edge, say $e = vu \in E(M)$ such that $v, u \notin X_1$. Otherwise, we have $C_7 \subseteq K_{3 \times 3} \subseteq \overline{G}[X_2, X_3, X_4]$; again a contradiction. W.l.g let $|N_G(v) \cap S| \geq |N_G(u) \cap S|$. Now by Claim 3.6 we have $|N_G(u) \cap S| \leq 1$. One can easily check that in any case, we have $C_7 \subseteq \overline{G}[S \cup \{u\}]$; again a contradiction. So w.l.g let $w_1 \in X_1$ and $w_2 \in X_2$. In this case, since $|N_G(u) \cap S| \leq 1$, we have $C_7 \subseteq \overline{G}[S \cup \{u\}]$; a contradiction again.

Case 2: $n = 2k + 1$ where $k \geq 2$, $|X_i| = k + 1$. Suppose that $m_4((n-2)K_2, C_7) \leq \lceil \frac{n-2+1}{2} \rceil$ for $n \geq 2$. We show that $m_4(nK_2, C_7) \leq \lceil \frac{n+1}{2} \rceil$ as follows: By contradiction, we may assume that $m_4(nK_2, C_7) > \lceil \frac{n+1}{2} \rceil$, that is, $K_{4 \times t}$ is 2-colorable to (nK_2, C_7) , say $nK_2 \not\subseteq G$ and $C_7 \not\subseteq \overline{G}$. Let $X'_i = X_i \setminus \{x_1^i\}$. By the induction hypothesis, we have $m_4((n-2)K_2, C_7) \leq \lceil \frac{n-1}{2} \rceil = \lceil \frac{2k}{2} \rceil = k$. Therefore, since $|X'_i| = k$ and $C_7 \not\subseteq \overline{G}$ we have $M = (n-2)K_2 \subseteq G[X'_1, X'_2, X'_3, X'_4]$ and thus we have the following claim:

Claim 3.7. *There exist two edges, say $e_1 = uv$ and $e_2 = u'v'$ in $E(M) = E((n-2)K_2)$, such that v, v', u and u' are in different partite.*

Proof. W.l.g assume that $v \in X'_1$ and $u \in X'_2$. By contradiction, assume that $|E(M) \cap E(G[X'_3, X'_4])| = 0$, that is, $G[X'_3, X'_4] \subseteq \overline{G}$. Since $|V(M)| = 2(n-2)$ and $|X'_i| = k$, we have $|V(M) \cap X'_i| \geq k-2$. Since $k \geq 3$, $|V(M) \cap X'_j| \geq 1$ ($j = 3, 4$). W.l.g let $e'_j = x_j y_j \in E(M)$ where $x_j \in V(M) \cap X'_j$. And w.l.g we may assume that $y_3 \in V(M) \cap X'_1$. Hence we have $y_4 \in V(M) \cap X'_1$. In other words, take $e_1 = x_3 y_3$ and $e_2 = x_4 y_4$ and the proof is complete. Hence we have $|E(M) \cap E(G[X'_2, X'_j])| = 0$ for $j = 3, 4$, in other words, if there exists $e'' \in E(M) \cap E(G[X'_2, X'_j])$, then set $e_1 = e'_1$ and $e_2 = e''$ and the proof is complete. Therefore, for each $e \in E(M)$ we have $v(e) \cap X'_1 \neq \emptyset$ which means that $|M| \leq |X'_1| = k$; a contradiction to $|M|$.

Now by Claim 3.7 there exist two edges, say $e_1 = uv$ and $e_2 = u'v'$ in $E(M) = E((n-2)K_2)$, such that v, v', u and u' are in different partite. W.l.g let $e_1 = x_1x_2$ and $e_2 = x_3x_4$ be these edges and let $x_i \in X'_i$ for $i = 1, 2, 3, 4$. Set $X''_i = X_i \setminus \{x_i\}$, hence we have $|X''_i| = k$. Since $C_7 \not\subseteq \overline{G}$ we have $C_7 \not\subseteq \overline{G}[X''_1, X''_2, X''_3, X''_4]$. Therefore, by the induction hypothesis, we have $(n-2)K_2 \subseteq G[X''_1, X''_2, X''_3, X''_4]$. Let $M = (n-2)K_2 \subseteq G[X''_1, X''_2, X''_3, X''_4]$, set $M^* = M \cup \{e_1, e_2\}$ hence $|M^*| = n$, that is, $nK_2 \subseteq G$; again a contradiction. Hence the assumption that $m_4(nK_2, C_7) > \lceil \frac{n+1}{2} \rceil$ does not hold and we have $m_4(nK_2, C_7) \leq \lceil \frac{n+1}{2} \rceil$. This completes the induction step and the proof is complete. By cases 1, 2 we have $m_4(nK_2, C_7) = \lceil \frac{n+1}{2} \rceil$ for $n \geq 4$. ■

The results of Proposition 3.1, Lemmas 3.4 and 3.5 and Theorem 3.2 concludes the proof of Theorem 1.2.

REFERENCES

- [1] Bondy, J.A., Murty, U.S.R.: Graph theory with applications. American Elsevier Publishing Co., Inc., New York (1976)
- [2] Burger, A.P., Grobler, P.J.P., Stipp, E.H., van Vuuren, J.H.: Diagonal Ramsey numbers in multipartite graphs. Util. Math. **66**, 137–163 (2004)
- [3] Burger, A.P., Vuuren, J.H.V.: Ramsey numbers in complete balanced multipartite graphs. II. Size numbers. Discrete Math. **283**(1-3), 45–49 (2004). DOI 10.1016/j.disc.2004.02.003. URL <https://doi.org.creativapplus.uaslp.mx/10.1016/j.disc.2004.02.003>
- [4] Erdős, P., Rado, R.: A partition calculus in set theory. Bull. Amer. Math. Soc. **62**, 427–489 (1956). DOI 10.1090/S0002-9904-1956-10036-0. URL <https://doi.org.creativapplus.uaslp.mx/10.1090/S0002-9904-1956-10036-0>
- [5] Jayawardene, C., Baskoro, E.T., Samarasekara, L., Sy, S.: Size multipartite Ramsey numbers for stripes versus small cycles. Electron. J. Graph Theory Appl. (EJGTA) **4**(2), 157–170 (2016). DOI 10.5614/ejgta.2016.4.2.4. URL <https://doi.org.creativapplus.uaslp.mx/10.5614/ejgta.2016.4.2.4>
- [6] L. uczak, T., Polcyn, J.: The multipartite Ramsey number for the 3-path of length three. Discrete Math. **341**(5), 1270–1274 (2018). DOI 10.1016/j.disc.2018.01.015. URL <https://doi.org.creativapplus.uaslp.mx/10.1016/j.disc.2018.01.015>
- [7] Lusiani, A., Baskoro, E.T., Saputro, S.W.: On size multipartite Ramsey numbers for stars versus paths and cycles. Electron. J. Graph Theory Appl. (EJGTA) **5**(1), 43–50 (2017). DOI 10.5614/ejgta.2017.5.1.5. URL <https://doi.org.creativapplus.uaslp.mx/10.5614/ejgta.2017.5.1.5>
- [8] Perondi, P.H., Carmelo, E.L.M.: Set and size multipartite Ramsey numbers for stars. Discrete Appl. Math. **250**, 368–372 (2018). DOI 10.1016/j.dam.2018.05.016. URL <https://doi.org.creativapplus.uaslp.mx/10.1016/j.dam.2018.05.016>
- [9] Sy, S.: On the size multipartite Ramsey numbers for small path versus cocktail party graphs. Far East J. Appl. Math. **55**(1), 53–60 (2011)

¹DEPARTMENT OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDIES IN BASIC SCIENCES (IASBS), ZANJAN 66731-45137, IRAN

Email address: y.rowshan@iasbs.ac.ir

Email address: gholami.m@iasbs.ac.ir