

# Potentials versus Geometry\*

T. Curtright<sup>§</sup> and S. Subedi<sup>∞</sup>

Department of Physics, University of Miami, Coral Gables, Florida 33124

<sup>§</sup>curtright@miami.edu    <sup>∞</sup>sushil.subedi04@gmail.com

October 13, 2021

## Abstract

We discuss some equivalence relations between the non-relativistic quantum mechanics for particles subjected to potentials and for particles moving freely on background geometries. In particular, we illustrate how selected geometries can be used to regularize singular potentials.

Given a solution to Einstein's theory of gravity acting as a background spacetime geometry, it is well-known that particle motion on this fixed geometry can be described by an effective potential [1]. But given a potential, it is perhaps less widely recognized that an equivalent particle dynamics can be described by an effective geometry. This equivalence is discussed here in the context of non-relativistic quantum mechanics. That is to say, only the non-relativistic form of Schrödinger's equation is used in our discussion, and only spatial dimensions are allowed non-trivial geometry, while time is taken to be common to all frames and universal to all observers.

Potential problems can be related to geometrical models in ways that often make the physics easier to extract and understand from the perspective of one or the other side of the relationship. For a generic potential the corresponding geometry can be singular, especially if the potential is singular (e.g. inverse powers of  $r$ ). On the other hand, non-singular geometries can give regularized forms of singular potentials that specify the essential physical features of those potentials without mathematical ambiguities or pathologies.

Consider a non-relativistic particle of mass  $\mu$  with energy  $E = \hbar^2 \varepsilon / (2\mu)$ , subject to a potential  $V(\vec{r}) = \hbar^2 U(\vec{r}) / (2\mu)$ , and whose wave function is a solution of the Schrödinger equation in flat Euclidean space.

$$-\nabla^2 \Psi + U \Psi = \varepsilon \Psi \quad (1)$$

Equivalence between this potential problem and a suitably chosen non-flat geometry can be obtained by relating the solutions of (1) with those for *free* particle motion on a specified manifold, written as

$$-\frac{1}{\sqrt{g}} \partial_\mu (\sqrt{g} g^{\mu\nu} \partial_\nu \Psi) = \varepsilon \Psi \quad (2)$$

In particular, given a rotationally invariant potential,  $U(r)$ , for a particle with specified angular momentum there is an equivalent radial Schrödinger equation that describes a particle moving freely on a curved space whose geometry is nontrivial, and vice versa. The main result can be expressed in general as an integro-differential modification of the Riccati equation [2].

In the case of two spatial dimensions (2D) with Euclidean signature, for a particle with angular momentum  $m$  so that the particle's wave function factorizes as  $\Psi(r, \theta) = \Psi_m(r) \exp(im\theta)$ , the main result is

$$\frac{dW}{dr} + W^2 + \frac{m^2}{R_0^2} \exp\left(-4 \int_{r_0}^r W(\tau) d\tau\right) = U(r) + \frac{m^2 - 1/4}{r^2} \quad (3)$$

Upon solving (3), the function  $W$  encodes the geometry in terms of the invariant distance on the curved 2D space as

$$(ds)^2 = (dr)^2 + R^2(r) (d\theta)^2 \quad \text{with} \quad R(r) = R_0 \exp\left(2 \int_{r_0}^r W(\tau) d\tau\right) \quad \text{i.e.} \quad W(r) = \frac{d}{dr} \ln \sqrt{R(r)} \quad (4)$$

---

\*Not quite the same as *Ford v Ferrari*. Or is it?

and the invariant Laplacian on the 2-manifold is

$$\frac{1}{\sqrt{g}}\partial_\mu(\sqrt{g}g^{\mu\nu}\partial_\nu) = \frac{1}{R(r)}\partial_r(R(r)\partial_r) + \frac{1}{R(r)^2}\partial_\theta^2 = \partial_r^2 + 2W(r)\partial_r + \frac{1}{R(r)^2}\partial_\theta^2 \quad (5)$$

Alternatively, given the geometry of the surface with a specified  $R(r)$ , the corresponding effective potential in flat space for angular momentum  $m$  is given by

$$U(r) = \frac{1}{2}\frac{R''(r)}{R(r)} - \frac{1}{4}\left(\frac{R'(r)}{R(r)}\right)^2 + \left(\frac{m}{R(r)}\right)^2 - \frac{m^2 - \frac{1}{4}}{r^2} \quad (6)$$

As should be expected, for  $R(r) \neq r$  this effective radial potential will depend on the angular momentum.

To establish the relation between the potential problem in 2D flat space and the 2D curved space system without a potential, it is only necessary to eliminate the first derivative term in (5) by writing  $\Psi(r, \theta) = \exp\left(-\int_{r_0}^r W(\tau) d\tau\right) \psi(r, \theta)$  to obtain

$$\left(\partial_r^2 + 2W(r)\partial_r + \frac{1}{R(r)^2}\partial_\theta^2\right)\Psi(r, \theta) = \exp\left(-\int_{r_0}^r W(\tau) d\tau\right)\left(\partial_r^2 + \frac{1}{R(r)^2}\partial_\theta^2 - W' - W^2\right)\psi(r, \theta) \quad (7)$$

and compare this to the flat space system with a potential  $U$ , as obtained by writing  $\Psi(r, \theta) = \psi(r, \theta)/\sqrt{r}$ , namely,

$$\left(\partial_r^2 + \frac{1}{r}\partial_r + \frac{1}{r^2}\partial_\theta^2 - U(r)\right)\Psi(r, \theta) = \frac{1}{\sqrt{r}}\left(\partial_r^2 + \frac{1}{r^2}\partial_\theta^2 - U(r) + \frac{1}{4r^2}\right)\psi(r, \theta) \quad (8)$$

Separating variables as  $\psi(r, \theta) = \psi_m(r) \exp(im\theta)$  for both (7) and (8) leads to the same second-order radial equation provided

$$-\frac{m^2}{R(r)^2} - W' - W^2 = -\frac{(m^2 - 1/4)}{r^2} - U(r) \quad (9)$$

and hence (3) with  $R(r) = R_0 \exp\left(2\int_{r_0}^r W(\tau) d\tau\right)$ , or alternatively (6).

In the case of  $N$  spatial dimensions (ND) with Euclidean signature [3], on a manifold with isotropic metric of the same form as (4), a particle with angular momentum  $\ell$  has a wave function that factorizes as  $\Psi = \Psi_\ell(r) Y_{lm_1 m_2 \dots m_{N-2}}(\Omega)$ , with all the angular dependence in the hyperspherical harmonics  $Y_{lm_1 m_2 \dots m_{N-2}}(\Omega)$ . The invariant Laplacian acting on  $\Psi$  is

$$\frac{1}{\sqrt{g}}\partial_\mu(\sqrt{g}g^{\mu\nu}\partial_\nu)\Psi = \frac{1}{R(r)^{N-1}}\partial_r\left(R(r)^{N-1}\partial_r\right)\Psi - \frac{1}{R(r)^2}L^2\Psi \quad (10)$$

where all the  $N-1$  angular derivatives are contained in  $L_{jk} \equiv -i(x_j\partial_k - x_k\partial_j)$  with  $L^2 \equiv \sum_{1 \leq j < k \leq N} L_{jk}L_{jk}$ . The  $Y_{lm_1 m_2 \dots m_{N-2}}$  are eigenfunctions of  $L^2$  and form a complete set on the hypersphere,  $S_{N-1}$ , analogous to the familiar spherical harmonics  $Y_{lm}$  on  $S_2$ . Acting on  $Y_{lm_1 m_2 \dots m_{N-2}}$  the  $L^2$  eigenvalues are given by [4]

$$L^2 Y_{lm_1 m_2 \dots m_{N-2}} = l(l + N - 2) Y_{lm_1 m_2 \dots m_{N-2}} \quad (11)$$

for  $l = 0, 1, 2, \dots$ , generalizing the well-known  $N = 3$  case. The eigenvalue equation (2) then reduces to the radial equation

$$\frac{-1}{R(r)^{N-1}}\partial_r\left(R(r)^{N-1}\partial_r\Psi_l(r)\right) + \frac{l(l + N - 2)}{R(r)^2}\Psi_l(r) = \varepsilon\Psi_l(r) \quad \text{for } 0 \leq r \leq \infty \quad (12)$$

Comparison to a potential system in ND flat space, (1), again leads to the main result in the form of a modified Riccati equation.

$$\frac{dW}{dr} + W^2 + \frac{\ell(\ell + N - 2)}{R_0^2} \exp\left(\frac{4}{1 - N} \int_{r_0}^r W(\tau) d\tau\right) = U(r) + \frac{(\ell + \frac{1}{2}(N - 1))(\ell + \frac{1}{2}(N - 3))}{r^2} \quad (13)$$

Upon solving for the function  $W$ , the effective geometry is encoded in the invariant distance on the curved ND space as

$$(ds)^2 = (dr)^2 + R^2(r) (d\Omega)^2 \quad \text{with} \quad R(r) = R_0 \exp\left(\frac{2}{N-1} \int_{r_0}^r W(\tau) d\tau\right) \quad \text{i.e.} \quad W(r) = \frac{N-1}{2} \frac{d}{dr} \ln R(r) \quad (14)$$

Alternatively, given the geometry of the surface with a specified  $R(r)$ , the corresponding potential in flat space for angular momentum  $\ell$  is given by

$$U(r) = \frac{(N-1)}{2} \frac{d^2}{dr^2} \ln R(r) + \left(\frac{N-1}{2}\right)^2 \left(\frac{d}{dr} \ln R(r)\right)^2 + \frac{\ell(\ell+N-2)}{R^2(r)} - \frac{(\ell + \frac{1}{2}(N-1))(\ell + \frac{1}{2}(N-3))}{r^2} \quad (15)$$

As in the 2D case, for  $R(r) \neq r$  this effective radial potential will depend on the angular momentum.

For  $\ell = 0$  “s-wave” solutions, (13) is an *unmodified* Riccati equation, a first-order differential equation whose solutions are well-studied [2]. For a  $1/r$  potential in ND, *plus a constant*, the  $\ell = 0$  case alone provides an illustration of potential  $\implies$  effective geometry. Let

$$U(r) = \frac{\kappa}{r} + \frac{\kappa^2}{(N-1)^2} \quad (16)$$

so that (13) for  $\ell = 0$  becomes

$$\frac{dW}{dr} + W^2 = \frac{\kappa}{r} + \frac{\kappa^2}{(N-1)^2} + \frac{(N-1)(N-3)}{4r^2} \quad (17)$$

A particularly simple, rational solution is immediately seen to be

$$W(r) = \frac{N-1}{2r} + \frac{\kappa}{N-1} \quad (18)$$

with general solutions obtained by quadrature [2].

The 3D case is exceptional in that there is no  $1/r^2$  term on the RHS of the Riccati equation. This physically interesting case readily admits another solution for a pure Coulomb potential, sans constant, namely,<sup>1</sup>

$$\begin{aligned} \frac{dW(r)}{dr} + W^2(r) &= \frac{\kappa}{r} \quad \text{for} \quad N=3 \quad \text{with} \\ W(r) &= \sqrt{\frac{\kappa I_0(2\sqrt{\kappa r})}{r I_1(2\sqrt{\kappa r})}} = \frac{1}{r} + \frac{\kappa}{2} - \frac{\kappa^2 r}{12} + O(r^2) \end{aligned} \quad (19)$$

which is well-approximated by (18) for small  $r$ . But as before, this is a solution only for  $\ell = 0$ .

The effective geometry corresponding to the solution (18) is given by

$$R(r) = K r e^{2\kappa r/(N-1)^2}, \quad K = \frac{R_0}{r_0} e^{-2\kappa r_0/(N-1)^2} \quad (20)$$

The invariant distance for this particular ND manifold is then

$$(ds)^2 = (dr)^2 + K^2 r^2 e^{4\kappa r/(N-1)^2} (d\Omega)^2 \quad (21)$$

In terms of more conventional isotropic coordinates, such that  $(ds)^2 = \left(\frac{dr(\rho)}{d\rho}\right)^2 (d\rho)^2 + \rho^2 (d\Omega)^2$ , let

$$\rho^2 = K^2 r^2 e^{4\kappa r/(N-1)^2} \quad (22)$$

---

<sup>1</sup>NB This is the  $1/r$  potential expressed as it would be in supersymmetric quantum mechanics.

and solve for  $r(\rho)$  in terms of the Lambert W function. For a repulsive potential  $\kappa \geq 0$ , and thus

$$r(\rho) = \frac{(N-1)^2}{2\kappa} \text{LambertW}\left(\frac{2\kappa\rho}{K(N-1)^2}\right), \quad \frac{dr(\rho)}{d\rho} = \frac{(N-1)^2}{2\kappa\rho} \left( \frac{\text{LambertW}\left(\frac{2\kappa\rho}{K(N-1)^2}\right)}{1 + \text{LambertW}\left(\frac{2\kappa\rho}{K(N-1)^2}\right)} \right) \quad (23)$$

The resulting geometry is described by

$$(ds)^2 = \left( \frac{(N-1)^2}{2\kappa\rho} \frac{\text{LambertW}\left(\frac{2\kappa\rho}{K(N-1)^2}\right)}{1 + \text{LambertW}\left(\frac{2\kappa\rho}{K(N-1)^2}\right)} \right)^2 (d\rho)^2 + \rho^2 (d\Omega)^2 \quad (24)$$

A canonical embedding of this  $N$ -manifold into an  $N+1$  Lorentz space is given by

$$(ds)^2 = (d\rho)^2 + \rho^2 (d\Omega)^2 - c^2 (dt(\rho))^2 \quad (25)$$

with

$$c \frac{dt(\rho)}{d\rho} = \frac{\sqrt{\left(1 + \text{LambertW}\left(\frac{2\kappa\rho}{K(N-1)^2}\right)\right)^2 - \left(\frac{(N-1)^2}{2\kappa\rho} \text{LambertW}\left(\frac{2\kappa\rho}{K(N-1)^2}\right)\right)^2}}{1 + \text{LambertW}\left(\frac{2\kappa\rho}{K(N-1)^2}\right)} \quad (26)$$

For  $\kappa \geq 0$  this is a real embedding all the way down to  $\rho = 0$  if  $K \geq 1$ . For  $K < 1$  the embedding is real only for  $\rho_{\min} \leq \rho \leq \infty$  where  $\rho_{\min}$  is given by the positive real root of the radicand in (26).

Graphical representations of the embedded surface and generalizations to situations where  $\ell \neq 0$  are left as an exercise for the reader, as is the  $\kappa < 0$  situation. But it should already be evident from the  $\ell = 0$  case that a repulsive  $1/r$  model is more easily understood as a potential problem than it is from the geometrical side of the relationship. Rather than pursue the quantum mechanics on the resulting ND manifold for this example, we consider next a more troublesome singular potential which can be regularized by mapping onto a well-known geometry.

Consider a smooth, spatial ‘‘bridge’’ manifold [5] (i.e. a ‘‘wormhole’’ [6]) given by

$$(ds)^2 = (dw)^2 + R^2(w) (d\Omega)^2, \quad -\infty \leq w \leq +\infty \quad (27)$$

where  $R(w) > 0$  has an absolute minimum at  $w = 0$ , and behaves asymptotically as  $R^2(w) \underset{w \rightarrow \pm\infty}{\sim} w^2 + O(1/w^2)$ . For example, the  $w$ -form of a static Ellis metric [7] in  $N$  spatial dimensions is simply

$$R^2(w) = R_0^2 + w^2, \quad -\infty \leq w \leq +\infty \quad (28)$$

The metric (27) leads to the invariant Laplacian

$$\frac{1}{\sqrt{g}} \partial_\mu (\sqrt{g} g^{\mu\nu} \partial_\nu \Psi) = \frac{1}{R(w)^{N-1}} \partial_w \left( R(w)^{N-1} \partial_w \right) - \frac{1}{R(w)^2} L^2 \quad (29)$$

where again all the  $N-1$  angular derivatives are contained in  $L^2$ .

The non-relativistic Schrödinger energy eigenvalue problem for a particle moving freely on this manifold is again solved by separating variables.

$$\frac{1}{\sqrt{g}} \partial_\mu (\sqrt{g} g^{\mu\nu} \partial_\nu \Psi) + \varepsilon \Psi = 0, \quad \Psi = \Psi_l(w) Y_{lm_1 m_2 \dots m_{N-2}}(\Omega) \quad (30)$$

The effective radial equation is now

$$\frac{1}{R(w)^{N-1}} \partial_w \left( R(w)^{N-1} \partial_w \Psi_l(w) \right) + \left( \varepsilon - \frac{l(l+N-2)}{R(w)^2} \right) \Psi_l(w) = 0 \quad (31)$$

Rather than compare this to rotationally invariant potential scattering in ND flat space, as above, instead compare this eigenvalue equation to one-dimensional potential scattering on the line  $-\infty \leq w \leq +\infty$ , as given by

$$\partial_w^2 \Psi_l(w) + (\varepsilon - U(w)) \Psi_l(w) = 0 \quad (32)$$

To establish the sought-for relation, again just eliminate the first derivative term in (31) by writing  $\Psi_l(w) = (R(w))^{(1-N)/2} \psi_l(w)$  to obtain

$$\begin{aligned} & \left( \partial_w^2 + (N-1) \frac{R'}{R} \partial_w + \left( \varepsilon - \frac{l(l+N-2)}{R(w)^2} \right) \right) \Psi_l(w) \\ &= R^p(w) \left( \partial_w^2 + \left( \varepsilon - \frac{1}{2}(N-1) \frac{R''}{R} - \frac{1}{4}(N-1)(N-3) \left( \frac{R'}{R} \right)^2 - \frac{l(l+N-2)}{R^2} \right) \right) \psi_l(w) \end{aligned} \quad (33)$$

where primes indicate derivatives with respect to  $w$ .

The second-order equation (33) is the same as that for potential scattering on the line (32) if

$$U(w) = \frac{1}{2}(N-1) \frac{R''}{R} + \frac{1}{4}(N-1)(N-3) \left( \frac{R'}{R} \right)^2 + \frac{l(l+N-2)}{R^2} \quad (34)$$

For the Ellis metric (28),  $R' = w/R$ ,  $R'' = R_0^2/R^3$ , so

$$U(w) = (N+2l-3)(N+2l-1) \frac{1}{4R(w)^2} - (N-1)(N-5) \frac{R_0^2}{4R(w)^4} \quad (35)$$

In particular, for 3D,

$$U(w) = \frac{l(l+1)}{R_0^2 + w^2} + \frac{R_0^2}{(R_0^2 + w^2)^2} \quad \text{for } N=3 \quad (36)$$

This  $U(w)$  may be thought of as a regularized form of  $1/r^4$ , a singular potential that has been well-studied in 3D [8, 9].

The angular momentum *independent*  $1/R^4$  term in the potential (35) is repulsive for  $N \leq 4$ , absent for  $N = 5$ , and attractive for  $N \geq 6$ . (It so happens to be a regularized form of an attractive electrostatic potential for a point particle for  $N = 6$  [3, 10].) In any case, non-relativistic scattering for the regular potential  $U(w)$  has both elastic and inelastic components for all  $N \geq 2$ , where elastic scattering is interpreted as both incident and emergent probability flux on the “upper” branch of the wormhole manifold, and inelastic scattering is to be understood as emergent flux on the “lower” branch of the manifold, without any incident flux on that lower branch. For incident flux on the wormhole’s upper branch, with  $R_0 > 0$ , there is always some probability flow through the bridge joining the two branches so that particles are effectively absorbed by the wormhole, as viewed by an observer located on the upper branch. For any  $N$ , the time independent partial wave scattering amplitudes can be determined exactly for  $U(w)$  in terms of spheroidal wave functions [11]. An Argand diagram of the phase shifts clearly reveals the inelastic behavior. This is discussed in considerable detail in [12].

The singular  $1/r^4$  potential that arises from  $U(w)$  in the limit as  $R_0 \rightarrow 0$  is somewhat problematic, as explained in [9] and references cited therein. At issue is whether the Hamiltonian for such a potential admits a self-adjoint extension for  $0 \leq r \leq \infty$ , with only real eigenvalues and unitary time evolution. For a repulsive  $1/r^4$  this can be arranged with an appropriate choice of boundary conditions at  $r = 0$ . An attractive  $1/r^4$  is another story, however, with considerable ambiguity in the boundary conditions at the origin. As stated in [9]:

“Thus the basic feature of an attractive singular potential is seen to lie in the fact that physical processes are not uniquely determined. This gives rise to the possibility of imposing unusual or unconventional boundary conditions in physical problems as a means of representing particular physical processes. An example of a process of this type is provided by particle absorption or capture.”

“Since the deficiency indices are infinite, nonself-adjoint extensions are possible which would correspond to boundary conditions which describe inelastic scattering.”

“Thus, in the singular case, the long-range part of the force between particles does not alone suffice to determine their behavior; some cutoff mechanism apparently must be provided.”

The regularization defined by the wormhole geometry avoids all these ambiguities. The Hamiltonian for a non-relativistic system governed by  $U(w)$  is manifestly self-adjoint for  $R_0 > 0$  when defined on a rigged Hilbert space with the usual boundary conditions as  $w \rightarrow \pm\infty$ . But there is a price to be paid for this mathematical convenience, namely, inelastic scattering, even when the potential is repulsive. This feature for  $U(w)$  is discussed further in [12].

In conclusion, the paper has exhibited various relationships between non-relativistic quantum systems involving a potential, in flat space, and systems without a potential but defined on curved manifolds. The main general results are encoded in (13), (14), (15), and (34) for  $N$ -dimensional spatial manifolds. More specific examples have been presented involving  $1/r$  potentials and regularized  $1/R^4$  potentials on wormhole manifolds. Perhaps our discussion should be viewed as a contemporary reconsideration of Riemann’s ideas about dynamics in terms of geometry [13]. Or perhaps not. The reader can decide.

## References

- [1] C.W. Misner, K.S. Thorne, and J.A. Wheeler, *Gravitation*, Princeton University Press (2017), Chapter 25.
- [2] E.L. Ince, *Ordinary Differential Equations*, Dover, New York (1956).
- [3] G. Green, “Mathematical investigations concerning the laws of the equilibrium of fluids analogous to the electric fluid, with other similar researches” *Trans. Cambridge Phil. Soc.* 5 (1835) 1–63.
- [4] A. Sommerfeld, *Partial Differential Equations in Physics*, Academic Press (1964), Appendix IV.
- [5] A. Einstein and N. Rosen, “The Particle Problem in the General Theory of Relativity” *Phys. Rev.* 48 (1935) 73-77.
- [6] M.S. Morris and K.S. Thorne, “Wormholes in spacetime and their use for interstellar travel: A tool for teaching general relativity” *Am. J. Phys.* 56 (1988) 395-412.
- [7] H.G. Ellis, “Ether flow through a drainhole: A particle model in general relativity” *J. Math. Phys.* 14 (1973) 104–118.
- [8] K.M. Case, “Singular Potentials” *Phys. Rev.* 80 (1950) 797-806.
- [9] W.M. Frank, D.J. Land, and R.M. Spector, “Singular Potentials” *Rev. Mod. Phys.* 43 (1971) 36-98.
- [10] T.L. Curtright, et al., “Charged line segments and ellipsoidal equipotentials” *Eur. J. Phys.* 37 (2016) 035201.
- [11] H. Volkmer, Chapter 30: Spheroidal Wave Functions, NIST Digital Library of Mathematical Functions, Release 1.1.0 of 2020-12-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [12] T.L. Curtright and S. Subedi, “Holistic Scattering Theory” *unpublished, in preparation*.
- [13] Riemann’s pre-relativity thoughts on gravity as curved space, and related ideas by several others, are discussed here.