

FOUR-DIMENSIONAL QUADRATIC FORMS OVER $\mathbb{C}((t))(X)$

PARUL GUPTA

ABSTRACT. For quadratic forms in 4 variables defined over the rational function field in one variable over $\mathbb{C}((t))$, the validity of the local-global principle for isotropy with respect to different sets of discrete valuations is examined.

CLASSIFICATION (MSC 2010): 11E04, 12E30, 12J10

KEYWORDS: isotropy, local-global-principle, rational function field, valuation, completion

1. INTRODUCTION

Let E be a field of characteristic different from 2 and let $E(X)$ denote the rational function field in one variable over E .

For $E = \mathbb{C}((t))$, the field of Laurent series in one variable over the complex numbers, the quadratic form

$$Y_1^2 + tY_2^2 + tY_3^2 + X(Y_1^2 + Y_2^2 + tY_4^2)$$

in the variables Y_1, Y_2, Y_3, Y_4 over $E(X)$ has no non-trivial zero, but it has a non-trivial zero over the completion of $E(X)$ with respect to any non-trivial valuation on $E(X)$ that is trivial on E . This is in contrast to the situation when E is a finite field, by the Hasse-Minkowski Theorem (See [6, Chapter VI, Theorem 66.1]). Note that, in both cases, the field E has a unique extension of each degree in a fixed algebraic closure.

By a \mathbb{Z} -valuation, we mean a valuation with value group \mathbb{Z} . A quadratic form is *isotropic* if it has a non-trivial zero, otherwise it is *anisotropic*. In all generality, an anisotropic quadratic form over $E(X)$ of dimension at most 3 remains anisotropic over the completion of $E(X)$ with respect to some \mathbb{Z} -valuation on $E(X)$ that is trivial on E ; this follows for example from Milnor's Exact Sequence [4, Theorem IX.3.1]. The case of 4-dimensional quadratic forms is the first case over $E(X)$ where the validity of such a local-global principle for isotropy depends on the base field E .

When E is a nondyadic local field, using a result of Lichtenbaum [5], one obtains that a 4-dimensional anisotropic quadratic form over $E(X)$ remains anisotropic

Date: December 13, 2021.

This work was supported by the *Fonds Wetenschappelijk Onderzoek – Vlaanderen (FWO)* in the *FWO Odysseus Programme* (project ‘Explicit Methods in Quadratic Form Theory’), the *Bijzonder Onderzoeksfonds (BOF)*, *University of Antwerp* (project BOF-DOCPRO-4, 2865), and the *Science and Engeneering Research Board (SERB)*, India (Grant CRG/2019/000271).

over the completion of $E(X)$ with respect to some \mathbb{Z} -valuation on $E(X)$ that is trivial on E (see [1, Remark 3.8]). This resembles the case where E is a finite field.

In contrast to the situations where E is a finite field or a local field, for $E = \mathbb{C}((t))$ the example of the quadratic form above shows that the local-global principle for isotropy of 4-dimensional quadratic forms over $E(X)$ fails with respect to \mathbb{Z} -valuations that are trivial on E . However, anisotropy of this quadratic form can be detected over the larger field $\mathbb{C}(X)((t))$, by using Springer's Theorem (see [4, Proposition VI.1.9]).

Consider the more general situation where the field E is complete with respect to a non-dyadic \mathbb{Z} -valuation v . In this case, a local-global principle for isotropy was obtained in [1] using a geometric setup. Let \mathcal{O}_v denote the valuation ring of v . By a *model for $E(X)$ over \mathcal{O}_v* we mean a two-dimensional integral normal projective flat \mathcal{O}_v -scheme \mathcal{X} whose function field is isomorphic to $E(X)$. Codimension-one points on a model of $E(X)$ over \mathcal{O}_v correspond to certain \mathbb{Z} -valuations on $E(X)$. For a model \mathcal{X} of $E(X)$ over \mathcal{O}_v let $\Omega_{\mathcal{X}}$ denote the set of \mathbb{Z} -valuations given by codimension-one points of \mathcal{X} . Consider the set $\Omega = \bigcup_{\mathcal{X}} \Omega_{\mathcal{X}}$ where the union is taken over all models \mathcal{X} of $E(X)$ over \mathcal{O}_v . It follows from [1, Theorem 3.1 and Remark 3.2] that an anisotropic quadratic form over $E(X)$ remains anisotropic over the completion of $E(X)$ with respect to some \mathbb{Z} -valuation in Ω . One may ask whether this remains true if one replaces Ω by $\Omega_{\mathcal{X}}$ for some well-chosen model \mathcal{X} of $E(X)$ over \mathcal{O}_v .

The aim of this note is to show that this is not the case: if the residue field of v is separably closed then, for any model \mathcal{X} of $E(X)$ over \mathcal{O}_v , there exists an anisotropic 4-dimensional quadratic form over $E(X)$ which is isotropic over the completion of $E(X)$ with respect to any $w \in \Omega_{\mathcal{X}}$ (Corollary 2). Let $\pi \in \mathcal{O}_v$ be a uniformiser of v . For any model \mathcal{X} of $E(X)$ over \mathcal{O}_v , the set $\{w(\pi) \mid w \in \Omega_{\mathcal{X}}\}$ is finite and hence it has an upper bound. However, for any positive integer r , the quadratic form

$$\varphi_r = (X^r - \pi)Y_1^2 + (X^{r+1} + \pi)Y_2^2 + \pi XY_3^2 + X(X^r + \pi)Y_4^2$$

is anisotropic over $E(X)$, but it is isotropic over the completion of $E(X)$ with respect to any \mathbb{Z} -valuation w on $E(X)$ with $w(\pi) < r$ (Theorem). The construction of φ_r is inspired by the example in [1, Remark 3.6] of an anisotropic 6-dimensional quadratic form over $\mathbb{Q}_p(X)$ where p is an odd prime.

2. RESULTS

We assume some familiarity with basic quadratic form theory over fields, for which we refer to [4]. We first fix some notation and recall some results.

By a *quadratic form* or simply a *form* we mean a regular quadratic form. Let E always be a field of characteristic different from 2 and let E^\times denote its multiplicative group. For $a_1, \dots, a_n \in E^\times$ the diagonal form $a_1X_1^2 + \dots + a_nX_n^2$ is denoted by $\langle a_1, \dots, a_n \rangle$.

Let v be a \mathbb{Z} -valuation on E . We denote the corresponding valuation ring, its maximal ideal and its residue field respectively by \mathcal{O}_v , \mathfrak{m}_v and κ_v . For an element $a \in \mathcal{O}_v$, let \bar{a} denote the image $a + \mathfrak{m}_v$ of a under the residue map $\mathcal{O}_v \rightarrow \kappa_v$. The completion of E with respect to v is denoted by E_v . We say that v is *henselian* if it extends uniquely to every finite field extension of E . Complete discretely valued fields are henselian (see [2, Theorem 1.3.1 and Theorem 4.1.3]). We recall a consequence of Hensel's Lemma:

Lemma. *Let v be a henselian \mathbb{Z} -valuation on E such that $v(2) = 0$. Then*

- (a) *The form $\langle u_1, u_2 \rangle$ over E is isotropic if and only if $\overline{u_1 u_2} \in -\kappa_v^{\times 2}$.*
- (b) *If κ_v is separably closed, then every 3-dimensional form over E is isotropic.*

Proof: Since $\overline{u_1 u_2} \in -\kappa_v^{\times 2}$ the polynomial equation $t^2 + \overline{u_1 u_2}$ has a solution in κ_v and since $v(2) = 0$ it follows by Hensel's Lemma [2, Theorem 4.1.3(4)] that $u_1 u_2 \in -E^2$, whereby the quadratic form $\langle u_1, u_2 \rangle$ over E is isotropic. Since κ_v is separably closed with $v(2) = 0$, we have that $\bar{u} \in -\kappa_v^{\times 2}$ for all $u \in \mathcal{O}_v^\times$. Since every 3-dimensional quadratic form over E contains a 2-dimensional form isometric to $\lambda \langle 1, u \rangle$ for some $u \in \mathcal{O}_v^\times$ and $\lambda \in E^\times$; (b) follows from (a). \square

The set of all \mathbb{Z} -valuations on $E(X)$ is denoted by $\Omega_{E(X)}$. For $r \in \mathbb{N}$, we define

$$\Omega_r = \{w \in \Omega_{E(X)} \mid w(E^\times) = i\mathbb{Z} \text{ for some } 0 \leq i \leq r\}.$$

With this notation, Ω_0 is the set of all E -trivial \mathbb{Z} -valuations on $E(X)$. We recall that any monic irreducible polynomial $p \in E[X]$ determines a unique \mathbb{Z} -valuation v_p on $E(X)$ which is trivial on E and such that $v_p(p) = 1$. There is further a unique \mathbb{Z} -valuation v_∞ on $E(X)$ such that $v_\infty(f) = -\deg(f)$ for any $f \in E[X] \setminus \{0\}$. Moreover, every \mathbb{Z} -valuation w on $E(X)$ trivial on E is either equal to v_∞ or to v_p for some monic irreducible polynomial $p \in E[X]$ (see [2, Theorem 2.1.4]), and in either of the two cases the residue field is a finite field extension of E .

Theorem. *Let v be a henselian \mathbb{Z} -valuation on E such that $v(2) = 0$. Assume that κ_v is separably closed. Let $\pi \in E^\times$ be such that $v(\pi) = 1$ and let $r \in \mathbb{N}$. Then the quadratic form*

$$\varphi_r = \langle X^r - \pi, X^{r+1} + \pi, \pi X, X(X^r + \pi) \rangle$$

is isotropic over $E(X)_w$ for every \mathbb{Z} -valuation $w \in \Omega_{r-1}$ but anisotropic over $E(X)_w$ for some $w \in \Omega_r$.

Proof: Set $F = E(X)$. We first show that φ_r is isotropic over F_w for all $w \in \Omega_{r-1}$. Consider $w \in \Omega_{r-1}$.

Case 1: $w(\pi) = 0 = w(X)$. Then κ_w is a finite extension of E . Since v is henselian, there is a unique extension v' of v to κ_w , and v' again henselian. Furthermore, it follows by [2, Theorem 3.3.4] that $v'(\kappa_w^\times)$ is isomorphic to \mathbb{Z} and $\kappa_{v'}$ is separably closed. It follows by part (b) of the Lemma that every 3-dimensional quadratic form over κ_w is isotropic. We have that $w = v_p$ for some monic irreducible

polynomial $p \in E[X]$ such that $p \neq X$. Note that, in this case at least three diagonal coefficients of φ_r are units in \mathcal{O}_w . It follows by Springer's Theorem [4, Proposition VI.1.9] that φ_r is isotropic over F_w .

Case 2: $0 \leq w(\pi) < r$ and $1 \leq w(X)$. Let $u = (X^r \pi^{-1} - 1)(X^{(r+1)} \pi^{-1} + 1)$. Then $w(u) = 0$ and $\bar{u} = -1 \in -\kappa_w^{\times 2}$. It follows by part (a) of the Lemma that the form $\pi^{-1} \langle X^r - \pi, X^{r+1} + \pi \rangle$ is isotropic over F_w . Thus φ_r is isotropic over F_w .

Case 3: $w(X) < 0 \leq w(\pi) < r$. Note that κ_w is either a finite extension of E or a rational function field over a finite extension of κ_v ; since $-1 \in \kappa_v^{\times 2}$, we get in either case that $-1 \in \kappa_w^{\times 2}$. Consider $u = (1 + \pi X^{-(r+1)})(1 + \pi X^{-r})$. We have that $w(u) = 0$ and $\bar{u} = 1 \in \kappa_w^{\times 2} = -\kappa_w^{\times 2}$. It follows by part (a) of the Lemma that the form $X^{-(r+1)} \langle X^{r+1} + \pi, X(X^r + \pi) \rangle$ is isotropic over F_w . Thus φ_r is isotropic over F_w .

We have thus shown that φ_r is isotropic over F_w for every $w \in \Omega_{r-1}$. Now we show that φ_r is anisotropic over F_w for some $w \in \Omega_F$.

Let $E' = E(s)$, where $s = \sqrt[r]{\pi}$. Then v extends uniquely to a valuation on E' which we again denote by v . Note that $s^r = \pi$ in E' and hence $v(\pi) = rv(s)$. Then $v' = rv$ is a \mathbb{Z} -valuation on E' .

Let $L = E'(X)$ and let $Y = \frac{X}{s}$. Note that $L = E'(Y)$. By [2, Corollary 2.2.2], there exists a unique extension of v' to L such that $v(Y) = 0$ and \bar{Y} is transcendental of $\kappa_{v'}$; we further have that $\kappa_w = \kappa_{v'}(\bar{Y})$ and $w(L^\times) = v'(E'^\times) = \mathbb{Z}$. Since $w(Y) = 0$, we have that $w(X) = w(s) = 1$. We get that

$$\varphi_r = \langle s^r(Y^r - 1), s^r(sY + 1), s^{r+1}Y, s^{r+1}Y(Y^r + 1) \rangle$$

Consider the forms $\varphi_1 = \langle Y^r - 1, sY + 1 \rangle$ and $\varphi_2 = \langle Y, Y(Y^r + 1) \rangle$.

Since $\bar{Y}^r - 1, \bar{Y}^r + 1 \notin -\kappa_w^{\times 2}$, it follows by Springer's Theorem [4, Proposition VI.1.9] that the quadratic form $s^{-r}\varphi_r$ is anisotropic over L_w . Hence φ_r is anisotropic over L_w . We obtain that φ_r is anisotropic over $F_{w|_F}$. Note that, $w(\pi) = w(s^r) = rw(s) = r$, thus $w \in \Omega_r$. \square

We now provide a different perspective to the above theorem. For a subset $\Omega \subseteq \Omega_{E(X)}$, we say that Ω has the *finite support property* if for every $f \in E(X)^\times$ the set $\{w \in \Omega \mid w(f) \neq 0\}$ is finite. It is well-known that Ω_0 has the finite support property. When E carries a discrete valuation the set $\Omega_{E(X)}$ does not have the finite support property. However, for any model \mathcal{X} of $E(X)$ over \mathcal{O}_v , the set $\Omega_{\mathcal{X}}$ contains Ω_0 and has the finite support property. We show the following:

Corollary 1. *Let v be a henselian \mathbb{Z} -valuation on E with $v(2) = 0$. Assume that κ_v is separably closed. Let $\Omega \subseteq \Omega_{E(X)}$ be a subset with the finite support property. Then there exists an anisotropic 4-dimensional quadratic form over $E(X)$ which is isotropic over $E(X)_w$ for every $w \in \Omega$.*

Proof: Let $\pi \in E^\times$ be such that $v(\pi) = 1$. Since Ω has the finite support property, the set $\{w \in \Omega \mid w(\pi) \neq 0\}$ is finite. Set $r = 1 + \max\{w(\pi) \mid w \in \Omega\}$. Clearly

$\Omega \subseteq \Omega_{r-1}$. Then the form φ_r in the Theorem is isotropic over $E(X)_w$ for every $w \in \Omega$, but anisotropic over $E(X)$. \square

Corollary 2. *Let v be a henselian \mathbb{Z} -valuation on E with $v(2) = 0$. Assume that κ_v is separably closed. Let \mathcal{X} be a regular model of $E(X)$ over \mathcal{O}_v . Then there exists an anisotropic 4-dimensional quadratic form over $E(X)$ which is isotropic over $E(X)_w$ for every $w \in \Omega_{\mathcal{X}}$.*

Proof: By [3, Chapter II, Lemma 6.1], for every element $f \in E(X)^\times$ the set $\{w \in \Omega_{\mathcal{X}} \mid w(f) \neq 0\}$ is finite, hence the statement follows by Corollary 1. \square

Acknowledgments. The author wishes to thank David Leep, Suresh Venapally, Karim Johannes Becher, Gonzalo Manzano Flores and Marco Zaninelli for many inspiring discussions and comments related to this article. This article is based on the author's PhD-thesis, prepared under the supervision of Karim Johannes Becher (*Universiteit Antwerpen*) and Arno Fehm (*Technische Universität Dresden*) in the framework of a joint PhD at *Universiteit Antwerpen* and *Universität Konstanz*.

REFERENCES

- [1] J.-L. Colliot-Thélène, R. Parimala, and V. Suresh, *Patching and local-global principles for homogeneous spaces over function fields of p -adic curves*, Comment. Math. Helv. (2012), no. 87, 1011–1033.
- [2] A.J. Engler and A. Prestel, *Valued fields*, Springer-Verlag, 2005.
- [3] R. Hartshorne, *Algebraic geometry*, vol. 52, Springer-Verlag, New York-Berlin, 1977.
- [4] T.Y. Lam, *Introduction to quadratic forms over fields*, American Mathematical Society, 2005.
- [5] S. Lichtenbaum, *Duality theorems for curves over p -adic fields*, Inventiones Mathematicae (1969), no. 7, 120–136.
- [6] O.T. O'Meara, *Introduction to quadratic forms*, Springer-Verlag Berlin Heidelberg GmbH, 1973.

UNIVERSITEIT ANTWERPEN, DEPARTEMENT WISKUNDE, MIDDELHEIMLAAN 1, 2020 ANTWERPEN, BELGIUM.

IISER PUNE, DR. HOMI BHABHA ROAD, PASHAN, PUNE 411 008, INDIA.
Email address: parul.gupta@iiserpune.ac.in