

Introduction of frame in tensor product of n -Hilbert spaces

Prasenjit Ghosh^{1*} and Tapas Kumar Samanta²

ABSTRACT. We study the concept of frame in tensor product of n -Hilbert spaces as tensor product of n -Hilbert spaces is again a n -Hilbert space. We generalize some of the known results about bases to frames in this new Hilbert space. A relationship between frame and bounded linear operator in tensor product of n -Hilbert spaces is studied. Finally, the dual frame in tensor product of n -Hilbert spaces is discussed.

1. INTRODUCTION

There are several techniques to rebuild signals by using a family of elementary signals. One of these techniques was given by D. Gabor in 1946 [10]. Thereafter frame in Hilbert space has been developed by Duffin and Schaeffer [7]. It is very useful to study nonharmonic Fourier series, i.e., sequences of the type $\{e^{i\lambda_n x}\}_{n \in \mathbb{Z}}$, where $\{\lambda_n\}_{n \in \mathbb{Z}}$ is a family of real or complex numbers. Daubechies et al, connected frames with Gabor system and Wavelet in 1986 [4]. To translate informations, frames are more flexible tools than bases and which makes them very useful in signal processing [8], coding and communications [20], system modeling [6], filter bank theory [1] etc. In recent times, many generalizations of frames have been appeared. Some of them are g -frame [21], fusion frame [2] and g -fusion frame [19] etc. P. Ghosh and T. K. Samanta studied the stability of dual g -fusion frames and generalized atomic systems for operators in Hilbert spaces [11, 13].

In 1970, Diminnie et al, introduced the concept of 2-inner product space [5]. A generalization of 2-inner product space for $n \geq 2$ was

2020 *Mathematics Subject Classification.* 42C15, 46C07, 46C50.

Key words and phrases. Frame, Dual frame, Tensor product of Hilbert spaces, n -normed space, n -Hilbert space.

Received: dd mmmm yyyy, Accepted: dd mmmm yyyy.

* Corresponding author.

developed by A. Misiak in 1989 [16]. The basic concepts of tensor product of Hilbert spaces were presented by S. Rabinson [17].

In this paper, we study frame in the tensor product of n -Hilbert spaces and establish some of its properties. We note the result that in tensor product of n -Hilbert spaces, an image of a frame under a bounded linear operator is a frame if and only if the operator is invertible. Finally, dual of a frame in tensor product of n -Hilbert spaces is described.

Throughout this paper, X denotes separable Hilbert space associated with the inner product $\langle \cdot, \cdot \rangle_1$ and $l^2(\mathbb{N})$ denote the space of square summable scalar-valued sequences with index set of natural numbers \mathbb{N} .

2. PRELIMINARIES

Definition 2.1. [3] A sequence $\{p_i\}_{i=1}^{\infty} \subseteq X$ is said to be a frame for X if there exist positive constants A, B such that

$$A \|p\|_1^2 \leq \sum_{i=1}^{\infty} |\langle p, p_i \rangle_1|^2 \leq B \|p\|_1^2; \quad \forall p \in X.$$

The constants A and B are called frame bounds.

Definition 2.2. [3] Let $\{p_i\}_{i=1}^{\infty}$ be a frame for X . The synthesis operator, $T : l^2(\mathbb{N}) \rightarrow X$, defined by $T\{c_i\} = \sum_{i=1}^{\infty} c_i p_i$ is bounded operator and its adjoint, called the analysis operator, is given by $T^*p = \{\langle p, p_i \rangle_1\}_{i=1}^{\infty}$. The frame operator $S : X \rightarrow X$, is given by

$$Sp = TT^*p = \sum_{i=1}^{\infty} \langle p, p_i \rangle_1 p_i, \quad \text{for all } p \in X.$$

Definition 2.3. [3] A frame $\{q_i\}_{i=1}^{\infty}$ is said to be a dual of a frame $\{p_i\}_{i=1}^{\infty}$ for X if $p = \sum_{i=1}^{\infty} \langle p, q_i \rangle_1 p_i$, for all $p \in X$.

The tensor product of Hilbert spaces are introduced by several ways and it is a certain linear space of operators which was represented by Folland in [9], Kadison and Ringrose in [15].

Definition 2.4. [18] Let $(Y, \langle \cdot, \cdot \rangle_2)$ be a Hilbert space. Then the tensor product of Hilbert spaces X and Y is denoted by $X \otimes Y$ and it is defined to be an inner product space with respect to the inner product:

$$\langle p \otimes q, p' \otimes q' \rangle = \langle p, p' \rangle_1 \langle q, q' \rangle_2, \quad \text{for all } p, p' \in X; q, q' \in Y.$$

The norm on $X \otimes Y$ is given by

$$\|p \otimes q\| = \|p\|_1 \|q\|_2, \quad \text{for all } p \in X \text{ and } q \in Y.$$

The space $X \otimes Y$ is complete with respect to the above inner product. Therefore the space $X \otimes Y$ is a Hilbert space.

Tensor product of operators $Q \in \mathcal{B}(X)$ and $T \in \mathcal{B}(Y)$, is denoted by $Q \otimes T$ and defined as $(Q \otimes T)A = QAT^*$, for all $A \in X \otimes Y$. It can be easily verified that $Q \otimes T \in \mathcal{B}(X \otimes Y)$ [9].

Theorem 2.5. [9] Suppose $Q, Q' \in \mathcal{B}(X)$ and $T, T' \in \mathcal{B}(Y)$. Then

- (i) $Q \otimes T \in \mathcal{B}(X \otimes Y)$ and $\|Q \otimes T\| = \|Q\| \|T\|$.
- (ii) $(Q \otimes T)(f \otimes g) = Qf \otimes Tg$ for all $f \in X, g \in Y$.
- (iii) $(Q \otimes T)(Q' \otimes T') = (QQ') \otimes (TT')$.
- (iv) $Q \otimes T$ is invertible if and only if Q and T are invertible, in which case $(Q \otimes T)^{-1} = (Q^{-1} \otimes T^{-1})$.
- (v) $(Q \otimes T)^* = (Q^* \otimes T^*)$.

Definition 2.6. [14] A real valued function $\|\cdot, \dots, \cdot\| : H^n \rightarrow \mathbb{R}$ satisfying the following properties:

- (i) $\|x_1, x_2, \dots, x_n\| = 0$ if and only if x_1, \dots, x_n are linearly dependent,
- (ii) $\|x_1, x_2, \dots, x_n\|$ is invariant under permutations of x_1, \dots, x_n ,
- (iii) $\|\alpha x_1, x_2, \dots, x_n\| = |\alpha| \|x_1, x_2, \dots, x_n\|$, $\alpha \in \mathbb{K}$,
- (iv) $\|x + y, x_2, \dots, x_n\| \leq \|x, x_2, \dots, x_n\| + \|y, x_2, \dots, x_n\|$,

for all $x_1, x_2, \dots, x_n, x, y \in H$, is called n -norm on H . A linear space H , together with a n -norm $\|\cdot, \dots, \cdot\|$, is called a linear n -normed space.

Definition 2.7. [16] Let $n \in \mathbb{N}$ and H be a linear space of dimension greater than or equal to n over the field \mathbb{K} , where \mathbb{K} is the real or complex numbers field. A function $\langle \cdot, \cdot | \cdot, \dots, \cdot \rangle : H^{n+1} \rightarrow \mathbb{K}$ satisfying the following five properties:

- (i) $\langle x_1, x_1 | x_2, \dots, x_n \rangle \geq 0$ and $\langle x_1, x_1 | x_2, \dots, x_n \rangle = 0$

if and only if x_1, x_2, \dots, x_n are linearly dependent,

- (ii) $\langle x, y | x_2, \dots, x_n \rangle = \langle x, y | x_{i_2}, \dots, x_{i_n} \rangle$ for every permutations (i_2, \dots, i_n) of $(2, \dots, n)$,

- (iii) $\langle x, y | x_2, \dots, x_n \rangle = \overline{\langle y, x | x_2, \dots, x_n \rangle}$,

- (iv) $\langle \alpha x, y | x_2, \dots, x_n \rangle = \alpha \langle x, y | x_2, \dots, x_n \rangle$, $\alpha \in \mathbb{K}$

- (v) $\langle x + y, z | x_2, \dots, x_n \rangle = \langle x, z | x_2, \dots, x_n \rangle + \langle y, z | x_2, \dots, x_n \rangle$,
- for all $x, y, x_1, x_2, \dots, x_n \in H$, is called an n -inner product on X , and the pair $(X, \langle \cdot, \cdot | \cdot, \dots, \cdot \rangle)$ is called an n -inner product space.

Theorem 2.8. [16] Let H be an n -inner product space. Then

$$|\langle x, y | x_2, \dots, x_n \rangle| \leq \|x, x_2, \dots, x_n\| \|y, x_2, \dots, x_n\|,$$

for all $x, y, x_2, \dots, x_n \in H$, where

$$\|x_1, x_2, \dots, x_n\| = \sqrt{\langle x_1, x_1 | x_2, \dots, x_n \rangle},$$

is called Cauchy-Schwarz inequality.

Theorem 2.9. [16] Let H be an n -inner product space. Then

$$\|x_1, x_2, \dots, x_n\| = \sqrt{\langle x_1, x_1 | x_2, \dots, x_n \rangle}$$

defines a n -norm for which

$$\begin{aligned} & \langle x, y | x_2, \dots, x_n \rangle \\ &= \frac{1}{4} \left(\|x + y, x_2, \dots, x_n\|^2 - \|x - y, x_2, \dots, x_n\|^2 \right), \end{aligned}$$

and

$$\begin{aligned} & \|x + y, x_2, \dots, x_n\|^2 + \|x - y, x_2, \dots, x_n\|^2 \\ &= 2 \left(\|x, x_2, \dots, x_n\|^2 + \|y, x_2, \dots, x_n\|^2 \right) \end{aligned}$$

hold for all $x, y, x_1, x_2, \dots, x_n \in H$.

Definition 2.10. [14] A sequence $\{x_k\}$ in linear n -normed space H is said to be convergent to $x \in H$ if

$$\lim_{k \rightarrow \infty} \|x_k - x, e_2, \dots, e_n\| = 0$$

for every $e_2, \dots, e_n \in H$ and it is called a Cauchy sequence if

$$\lim_{l, k \rightarrow \infty} \|x_l - x_k, e_2, \dots, e_n\| = 0$$

for every $e_2, \dots, e_n \in H$. The space H is said to be complete if every Cauchy sequence in this space is convergent in H . A n -inner product space is called n -Hilbert space if it is complete with respect to its induce norm.

3. FRAME IN TENSOR PRODUCT OF n -HILBERT SPACES

Definition 3.1. Let H be a n -Hilbert space and $a_2, \dots, a_n \in H$. A sequence $\{p_i\}_{i=1}^{\infty}$ in H is said to be a frame associated to (a_2, \dots, a_n) for H if there exist constants $0 < A \leq B < \infty$ such that

$$(3.1) \quad A \|p, a_2, \dots, a_n\|^2 \leq \sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle|^2 \leq B \|p, a_2, \dots, a_n\|^2$$

for all $p \in H$. The constants A and B are called frame bounds. If $\{p_i\}_{i=1}^{\infty}$ satisfies the right hand inequality of (3.1), it is called a Bessel sequence associated to (a_2, \dots, a_n) in H with bound B .

By Cauchy-Schwarz inequality and the condition (i) of the definition 2.7, we may assume that every p_i and a_2, \dots, a_n are linearly independent.

Consider $F = \{a_2, a_3, \dots, a_n\}$, where a_2, a_3, \dots, a_n are fixed elements in H and L_F denote the linear subspace of H spanned by

F . The quotient space H/L_F is a normed linear space with respect to the norm, $\|p + L_F\|_F = \|p, a_2, \dots, a_n\|$ for every $p \in H$. Let M_F be the algebraic complement of L_F , then $H = L_F \oplus M_F$. Define

$$\langle p, q \rangle_F = \langle p, q | a_2, \dots, a_n \rangle \text{ on } H.$$

Then $\langle \cdot, \cdot \rangle_F$ is a semi-inner product on H and this induces an inner product on the quotient space H/L_F which is given by

$$\langle p + L_F, q + L_F \rangle_F = \langle p, q \rangle_F = \langle p, q | a_2, \dots, a_n \rangle; \forall p, q \in H.$$

By identifying H/L_F with M_F in an obvious way, we obtain an inner product on M_F . Now, for every $p \in M_F$, we define $\|p\|_F = \sqrt{\langle p, p \rangle_F}$ and it can be easily verify that $(M_F, \|\cdot\|_F)$ is a norm space. Consider H_F as the completion of the inner product space M_F .

Theorem 3.2. *Let H be a n -Hilbert space. Then $\{p_i\}_{i=1}^\infty \subseteq H$ is a frame associated to (a_2, \dots, a_n) with bounds A, B if and only if it is a frame for the Hilbert space H_F with bounds A, B .*

Proof. Let us consider that $\{p_i\}_{i=1}^\infty \subseteq H$ be a frame associated to (a_2, \dots, a_n) for H with bounds A, B . Then the inequality (3.1) can be written as

$$A \|p\|_F^2 \leq \sum_{i=1}^{\infty} |\langle p, p_i \rangle_F|^2 \leq B \|p\|_F^2; \forall p \in M_F.$$

This shows that $\{p_i\}_{i=1}^\infty$ is a frame for M_F . By the Lemma 5.1.2 of [3], the sequence $\{p_i\}_{i=1}^\infty$ is also a frame for H_F with the same bounds.

Converse part is obvious. \square

Definition 3.3. Let $\{p_i\}_{i=1}^\infty$ be a frame associated to (a_2, \dots, a_n) for H . Then the bounded linear operator $T_F : l^2(\mathbb{N}) \rightarrow H_F$, defined by $T_F \{c_i\} = \sum_{i=1}^{\infty} c_i p_i$, is called pre-frame operator and its adjoint operator described by

$$T_F^* : H_F \rightarrow l^2(\mathbb{N}), T_F^* p = \{\langle p, p_i | a_2, \dots, a_n \rangle\}_{i=1}^\infty$$

is called the analysis operator. The operator $S_F : H_F \rightarrow H_F$ given by

$$S_F p = T_F T_F^* p = \sum_{i=1}^{\infty} \langle p, p_i | a_2, \dots, a_n \rangle p_i, \text{ for all } p \in H_F,$$

is called the frame operator.

Theorem 3.4. *Let $\{p_i\}_{i=1}^\infty$ be a frame associated to (a_2, \dots, a_n) for H with bounds A, B . Then the corresponding frame operator S_F is bounded, invertible, self-adjoint and positive.*

Proof. For each $p \in H_F$, we have

$$\begin{aligned}
& \|S_F p\|_F^2 = \|S_F p, a_2, \dots, a_n\|^2 \\
& = \sup \left\{ |\langle S_F p, q | a_2, \dots, a_n \rangle|^2 : \|q, a_2, \dots, a_n\| = 1 \right\} \\
& = \sup_{\|q, a_2, \dots, a_n\|=1} \left| \left\langle \sum_{i=1}^{\infty} \langle p, p_i | a_2, \dots, a_n \rangle p_i, q | a_2, \dots, a_n \right\rangle \right|^2 \\
& \leq \sup_{\|q, a_2, \dots, a_n\|=1} \sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle|^2 \sum_{i=1}^{\infty} |\langle q, p_i | a_2, \dots, a_n \rangle|^2 \\
& \text{[using Cauchy-Schwarz inequality]} \\
& \leq B^2 \|p, a_2, \dots, a_n\|^2 = B^2 \|p\|_F^2 \\
& \text{[since } \{p_i\}_{i=1}^{\infty} \text{ is a frame associated to } (a_2, \dots, a_n) \text{]}
\end{aligned}$$

This shows that S_F is bounded. Since $S_F = T_F T_F^*$, it is easy to verify that S_F is self-adjoint. The inequality (3.1), can be written as

$$\begin{aligned}
A \langle p, p | a_2, \dots, a_n \rangle & \leq \langle S_F p, p | a_2, \dots, a_n \rangle \leq B \langle p, p | a_2, \dots, a_n \rangle \\
\text{and this gives } A I_F & \leq S_F \leq B I_F. \text{ Thus, } S_F \text{ is positive and consequently it is invertible. } \square
\end{aligned}$$

Remark 3.5. In Theorem 3.4, it is proved that $A I_F \leq S_F \leq B I_F$. Since S_F^{-1} commutes with both S_F and I_F , multiplying in the inequality, $A I_F \leq S_F \leq B I_F$ by S_F^{-1} , we get $B^{-1} I_F \leq S_F^{-1} \leq A^{-1} I_F$.

For more details on frame in n -Hilbert space one can go through the paper [12].

Let H and K be two n -Hilbert spaces associated with the n -inner products $\langle \cdot, \cdot | \cdot, \dots, \cdot \rangle_1$ and $\langle \cdot, \cdot | \cdot, \dots, \cdot \rangle_2$, respectively. The tensor product of H and K is denoted by $H \otimes K$ and it is defined to be an n -inner product space associated with the n -inner product given by

$$\begin{aligned}
& \langle f_1 \otimes g_1, f_2 \otimes g_2 | f_3 \otimes g_3, \dots, f_n \otimes g_n \rangle \\
(3.2) \quad & = \langle f_1, f_2 | f_3, \dots, f_n \rangle_1 \langle g_1, g_2 | g_3, \dots, g_n \rangle_2,
\end{aligned}$$

for all $f_1, f_2, f_3, \dots, f_n \in H$ and $g_1, g_2, g_3, \dots, g_n \in K$. The n -norm on $H \otimes K$ is defined by

$$\begin{aligned}
& \|f_1 \otimes g_1, f_2 \otimes g_2, \dots, f_n \otimes g_n\| \\
(3.3) \quad & = \|f_1, f_2, \dots, f_n\|_1 \|g_1, g_2, \dots, g_n\|_2,
\end{aligned}$$

for all $f_1, f_2, \dots, f_n \in H$ and $g_1, g_2, \dots, g_n \in K$, where the n -norms $\|\cdot, \dots, \cdot\|_1$ and $\|\cdot, \dots, \cdot\|_2$ are generated by $\langle \cdot, \cdot | \cdot, \dots, \cdot \rangle_1$

and $\langle \cdot, \cdot | \cdot, \dots, \cdot \rangle_2$, respectively. The space $H \otimes K$ is complete with respect to the above n -inner product. Therefore the space $H \otimes K$ is an n -Hilbert space.

Consider $G = \{b_2, b_3, \dots, b_n\}$, where b_2, b_3, \dots, b_n are fixed elements in K and L_G denote the linear subspace of K spanned by G . Now, we can define the Hilbert space K_G with respect to the inner product is given by

$$\langle p + L_G, q + L_G \rangle_G = \langle p, q \rangle_G = \langle p, q | b_2, \dots, b_n \rangle_2; \forall p, q \in K.$$

Remark 3.6. According to the definition 2.4, $H_F \otimes K_G$ is the Hilbert space with respect to the inner product:

$$\langle p \otimes q, p' \otimes q' \rangle = \langle p, p' \rangle_F \langle q, q' \rangle_G,$$

for all $p, p' \in H_F$ and $q, q' \in K_G$.

Remark 3.7. From the definition of ordinary frames for separable Hilbert spaces, the sequence of vectors $\{p_i \otimes q_j\}_{i,j=1}^\infty$ in $H \otimes K$ can be consider as a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$ if there exist constants $0 < A \leq B < \infty$ such that

$$(3.4) \quad \begin{aligned} & A \|p \otimes q, a_2 \otimes b_2, \dots, a_n \otimes b_n\|^2 \\ & \leq \sum_{i,j=1}^{\infty} |\langle p \otimes q, p_i \otimes q_j | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle|^2 \\ & \leq B \|p \otimes q, a_2 \otimes b_2, \dots, a_n \otimes b_n\|^2; \forall p \otimes q \in H \otimes K, \end{aligned}$$

where $\{p_i\}_{i=1}^\infty$ and $\{q_j\}_{j=1}^\infty$ be the sequences of vectors in H and K , respectively and $a_2 \otimes b_2, a_3 \otimes b_3, \dots, a_n \otimes b_n$ be fixed elements in $H \otimes K$. The constants A, B are called the frame bounds. If $A = B$ then it is called a tight frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$. If the sequence $\{p_i \otimes q_j\}_{i,j=1}^\infty$ satisfies the right hand inequality of (3.4), is called a Bessel sequence associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ in $H \otimes K$.

Theorem 3.8. Let $\{p_i\}_{i=1}^\infty$ and $\{q_j\}_{j=1}^\infty$ be the sequences of vectors in n -Hilbert spaces H and K . The sequence $\{p_i \otimes q_j\}_{i,j=1}^\infty \subseteq H \otimes K$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$ if and only if $\{p_i\}_{i=1}^\infty$ is a frame associated to (a_2, \dots, a_n) for H and $\{q_j\}_{j=1}^\infty$ is a frame associated to (b_2, \dots, b_n) for K .

Proof. Suppose that the sequence $\{p_i \otimes q_j\}_{i,j=1}^\infty$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. Then, for each $p \otimes q \in H \otimes K - \{\theta \otimes \theta\}$, there exist constants $A, B > 0$ such that

$$A \|p \otimes q, a_2 \otimes b_2, \dots, a_n \otimes b_n\|^2$$

$$\begin{aligned} &\leq \sum_{i,j=1}^{\infty} |\langle p \otimes q, p_i \otimes q_j | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle|^2 \\ &\leq B \|p \otimes q, a_2 \otimes b_2, \dots, a_n \otimes b_n\|^2. \end{aligned}$$

Using the n -norm and n -inner product on $H \otimes K$, we get

$$\begin{aligned} A \|p, a_2, \dots, a_n\|_1^2 \|q, b_2, \dots, b_n\|_2^2 &\leq \left(\sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle_1|^2 \right) \times \\ &\left(\sum_{j=1}^{\infty} |\langle q, q_j | b_2, \dots, b_n \rangle_2|^2 \right) \leq B \|p, a_2, \dots, a_n\|_1^2 \|q, b_2, \dots, b_n\|_2^2. \end{aligned}$$

Since $p \otimes q \in H \otimes K$ is non-zero element i.e., $p \in H$ and $q \in K$ are non-zero elements. Here, we may assume that every p_i and a_2, \dots, a_n are linearly independent and every q_j and b_2, \dots, b_n are linearly independent. Hence

$$\sum_{j=1}^{\infty} |\langle q, q_j | b_2, \dots, b_n \rangle_2|^2, \sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle_1|^2$$

are non-zero. Therefore, by the above inequality, we get

$$\begin{aligned} \frac{A \|q, b_2, \dots, b_n\|_2^2 \|p, a_2, \dots, a_n\|_1^2}{\sum_{j=1}^{\infty} |\langle q, q_j | b_2, \dots, b_n \rangle_2|^2} &\leq \sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle_1|^2 \\ &\leq \frac{B \|q, b_2, \dots, b_n\|_2^2}{\sum_{j=1}^{\infty} |\langle q, q_j | b_2, \dots, b_n \rangle_2|^2} \|p, a_2, \dots, a_n\|_1^2. \end{aligned}$$

This implies that

$$A_1 \|p, a_2, \dots, a_n\|_1^2 \leq \sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle_1|^2 \leq B_1 \|p, a_2, \dots, a_n\|_1^2,$$

$$\text{for all } p \in H, \text{ where } A_1 = \inf_{q \in K} \left\{ \frac{A \|q, b_2, \dots, b_n\|_2^2}{\sum_{j=1}^{\infty} |\langle q, q_j | b_2, \dots, b_n \rangle_2|^2} \right\}$$

$$\text{and } B_1 = \sup_{q \in K} \left\{ \frac{B \|q, b_2, \dots, b_n\|_2^2}{\sum_{j=1}^{\infty} |\langle q, q_j | b_2, \dots, b_n \rangle_2|^2} \right\}.$$

This shows that $\{p_i\}_{i=1}^{\infty}$ is a frame associated to (a_2, \dots, a_n) for

H . Similarly, it can be shown that $\{q_j\}_{j=1}^{\infty}$ is a frame associated to (b_2, \dots, b_n) for K .

Conversely, suppose that $\{p_i\}_{i=1}^{\infty}$ is a frame associated to (a_2, \dots, a_n) for H with bounds A, B and $\{q_j\}_{j=1}^{\infty}$ is a frame associated to (b_2, \dots, b_n) for K with bounds C, D . Then, for all $p \in H$ and $q \in K$, we have

$$A \|p, a_2, \dots, a_n\|_1^2 \leq \sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle_1|^2 \leq B \|p, a_2, \dots, a_n\|_1^2,$$

$$C \|q, b_2, \dots, b_n\|_2^2 \leq \sum_{j=1}^{\infty} |\langle q, q_j | b_2, \dots, b_n \rangle_2|^2 \leq D \|q, b_2, \dots, b_n\|_2^2.$$

Multiplying the above two inequalities and using (3.2) and (3.3), we get

$$\begin{aligned} AC \|p \otimes q, a_2 \otimes b_2, \dots, a_n \otimes b_n\|^2 \\ \leq \sum_{i,j=1}^{\infty} |\langle p \otimes q, p_i \otimes q_j | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle|^2 \\ \leq BD \|p \otimes q, a_2 \otimes b_2, \dots, a_n \otimes b_n\|^2. \end{aligned}$$

Hence, $\{p_i \otimes q_j\}_{i,j=1}^{\infty}$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. \square

Theorem 3.9. *The sequence $\{p_i \otimes q_j\}_{i,j=1}^{\infty} \subseteq H \otimes K$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$ if and only if $\{p_i \otimes q_j\}_{i,j=1}^{\infty}$ is a frame for $H_F \otimes K_G$.*

Proof. Suppose that the sequence $\{p_i \otimes q_j\}_{i,j=1}^{\infty}$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. By Theorem 3.8, $\{p_i\}_{i=1}^{\infty}$ is a frame associated to (a_2, \dots, a_n) for H and $\{q_j\}_{j=1}^{\infty}$ is a frame associated to (b_2, \dots, b_n) for K . Now, applying the Theorem 3.2, $\{p_i\}_{i=1}^{\infty}$ and $\{q_j\}_{j=1}^{\infty}$ are frames for H_F and K_G , respectively. Hence, $\{p_i \otimes q_j\}_{i,j=1}^{\infty}$ is a frame for $H_F \otimes K_G$.

The proof of the converse part is obvious. \square

Remark 3.10. Let $\{p_i \otimes q_j\}_{i,j=1}^{\infty}$ be a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. According to the definition 3.3, the frame operator $S_{F \otimes G} : H_F \otimes K_G \rightarrow H_F \otimes K_G$ is described by

$$S_{F \otimes G}(p \otimes q) = \sum_{i,j=1}^{\infty} \langle p \otimes q, p_i \otimes q_j | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle (p_i \otimes q_j)$$

for all $p \otimes q \in H_F \otimes K_G$.

Theorem 3.11. *If S_F , S_G and $S_{F \otimes G}$ are the corresponding frame operator for $\{p_i\}_{i=1}^\infty$, $\{q_j\}_{j=1}^\infty$ and $\{p_i \otimes q_j\}_{i,j=1}^\infty$, respectively, then $S_{F \otimes G} = S_F \otimes S_G$ and $S_{F \otimes G}^{-1} = S_F^{-1} \otimes S_G^{-1}$.*

Proof. Since $S_{F \otimes G}$ is the frame operator for $\{p_i \otimes q_j\}_{i,j=1}^\infty$, we have

$$\begin{aligned} S_{F \otimes G}(p \otimes q) &= \sum_{i,j=1}^{\infty} \langle p \otimes q, p_i \otimes q_j | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle (p_i \otimes q_j) \\ &= \sum_{i,j=1}^{\infty} \langle p, p_i | a_2, \dots, a_n \rangle_1 \langle q, q_j | b_2, \dots, b_n \rangle_2 (p_i \otimes q_j) \\ &= \left(\sum_{i=1}^{\infty} \langle p, p_i | a_2, \dots, a_n \rangle_1 p_i \right) \otimes \left(\sum_{j=1}^{\infty} \langle q, q_j | b_2, \dots, b_n \rangle_2 q_j \right) \\ &= S_F(p) \otimes S_G(q) = (S_F \otimes S_G)(p \otimes q) \quad \forall p \otimes q \in H_F \otimes K_G. \end{aligned}$$

Thus $S_{F \otimes G} = S_F \otimes S_G$. Since S_F and S_G are invertible, by Theorem 2.5 (iv), $S_{F \otimes G}^{-1} = (S_F \otimes S_G)^{-1} = S_F^{-1} \otimes S_G^{-1}$. \square

Theorem 3.12. *Let $\{p_i\}_{i=1}^\infty$ be a frame associated to (a_2, \dots, a_n) for H with bounds A, B and $\{q_j\}_{j=1}^\infty$ be a frame associated to (b_2, \dots, b_n) for K with bounds C, D having their corresponding frame operators S_F and S_G , respectively. Then $\Lambda = \{S_{F \otimes G}^{-1}(p_i \otimes q_j)\}_{i,j=1}^\infty$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$ with the corresponding frame operator $S_{F \otimes G}^{-1}$.*

Proof. For each $p \otimes q \in H_F \otimes K_G$, we have

$$\begin{aligned} & \sum_{i,j=1}^{\infty} \left| \langle p \otimes q, S_{F \otimes G}^{-1}(p_i \otimes q_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle \right|^2 \\ &= \sum_{i,j=1}^{\infty} \left| \langle p \otimes q, (S_F^{-1} \otimes S_G^{-1})(p_i \otimes q_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle \right|^2 \\ &= \sum_{i,j=1}^{\infty} \left| \langle p \otimes q, S_F^{-1}(p_i) \otimes S_G^{-1}(q_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle \right|^2 \\ &= \sum_{i=1}^{\infty} \left| \langle p, S_F^{-1} p_i | a_2, \dots, a_n \rangle_1 \right|^2 \sum_{j=1}^{\infty} \left| \langle q, S_G^{-1} q_j | b_2, \dots, b_n \rangle_2 \right|^2 \\ (3.5) \quad &= \sum_{i=1}^{\infty} \left| \langle S_F^{-1} p, p_i | a_2, \dots, a_n \rangle_1 \right|^2 \sum_{j=1}^{\infty} \left| \langle S_G^{-1} q, q_j | b_2, \dots, b_n \rangle_2 \right|^2 \end{aligned}$$

$$\begin{aligned}
 &\leq B \left\| S_F^{-1}(p), a_2, \dots, a_n \right\|_1^2 D \left\| S_G^{-1}(q), b_2, \dots, b_n \right\|_2^2 \\
 &[\text{since } \{p_i\}_{i=1}^\infty \text{ is a frame associated to } (a_2, \dots, a_n), \text{ and} \\
 &\{q_j\}_{j=1}^\infty \text{ is a frame associated to } (b_2, \dots, b_n)] \\
 &\leq B D \left\| S_F^{-1} \right\|^2 \left\| S_G^{-1} \right\|^2 \left\| p, a_2, \dots, a_n \right\|_1^2 \left\| q, b_2, \dots, b_n \right\|_2^2 \\
 &[\text{since } S_F^{-1}, S_G^{-1} \text{ are bounded operators on } H_F \text{ and } K_G, \text{ respectively}] \\
 &\leq \frac{B D}{A^2 C^2} \left\| p \otimes q, a_2 \otimes b_2 \dots, a_n \otimes b_n \right\|^2 [\text{using (3.3)}] \\
 &[\text{since } B^{-1} I_F \leq S_F^{-1} \leq A^{-1} I_F \text{ and } D^{-1} I_G \leq S_G^{-1} \leq C^{-1} I_G, \\
 &\text{where } I_G \text{ denote the identity operator on } K_G].
 \end{aligned}$$

Now, for $p \in H_F$, we have

$$\begin{aligned}
 \left\| p, a_2, \dots, a_n \right\|_1 &\leq \left\| S_F \right\| \left\| S_F^{-1}(p), a_2, \dots, a_n \right\|_1 \\
 (3.6) \qquad \qquad \qquad &\leq B \left\| S_F^{-1}(p), a_2, \dots, a_n \right\|_1
 \end{aligned}$$

and similarly for $q \in K_G$, we get

$$(3.7) \qquad \left\| q, b_2, \dots, b_n \right\|_2 \leq D \left\| S_G^{-1}(q), b_2, \dots, b_n \right\|_2.$$

On the other hand, from (3.5),

$$\begin{aligned}
 &\sum_{i,j=1}^\infty \left| \left\langle p \otimes q, S_{F \otimes G}^{-1}(p_i \otimes q_j) \mid a_2 \otimes b_2, \dots, a_n \otimes b_n \right\rangle \right|^2 \\
 &= \sum_{i=1}^\infty \left| \left\langle S_F^{-1} p, p_i \mid a_2, \dots, a_n \right\rangle_1 \right|^2 \sum_{j=1}^\infty \left| \left\langle S_G^{-1} q, q_j \mid b_2, \dots, b_n \right\rangle_2 \right|^2 \\
 &\geq A \left\| S_F^{-1}(p), a_2, \dots, a_n \right\|_1^2 C \left\| S_G^{-1}(q), b_2, \dots, b_n \right\|_2^2 \\
 &\geq \frac{A C}{B^2 D^2} \left\| p, a_2, \dots, a_n \right\|_1^2 \left\| q, b_2, \dots, b_n \right\|_2^2 [\text{by (3.6) and (3.7)}] \\
 &= \frac{A C}{B^2 D^2} \left\| p \otimes q, a_2 \otimes b_2 \dots, a_n \otimes b_n \right\|^2.
 \end{aligned}$$

Hence, Λ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$ with bounds $\frac{A C}{B^2 D^2}$ and $\frac{B D}{A^2 C^2}$.

Furthermore, for each $p \otimes q \in H_F \otimes K_G$, we have

$$\begin{aligned}
 &\sum_{i,j=1}^\infty \left\langle p \otimes q, S_{F \otimes G}^{-1}(p_i \otimes q_j) \mid a_2 \otimes b_2, \dots, a_n \otimes b_n \right\rangle S_{F \otimes G}^{-1}(p_i \otimes q_j) \\
 &= \sum_{i,j=1}^\infty \left\langle p \otimes q, S_F^{-1} p_i \otimes S_G^{-1} q_j \mid a_2 \otimes b_2, \dots, a_n \otimes b_n \right\rangle S_F^{-1} p_i \otimes S_G^{-1} q_j
 \end{aligned}$$

$$\begin{aligned}
&= \left(\sum_{i=1}^{\infty} \langle S_F^{-1} p, p_i | a_2, \dots, a_n \rangle_1 S_F^{-1} p_i \right) \otimes \\
&\quad \left(\sum_{j=1}^{\infty} \langle S_G^{-1} q, q_j | b_2, \dots, b_n \rangle_2 S_G^{-1} q_j \right) \\
&= S_F^{-1} S_F (S_F^{-1} p) \otimes S_G^{-1} S_G (S_G^{-1} q) \\
&= S_F^{-1} p \otimes S_G^{-1} q = S_{F \otimes G}^{-1} (p \otimes q).
\end{aligned}$$

Hence, the corresponding frame operator for Λ is $S_{F \otimes G}^{-1}$. \square

Theorem 3.13. *Let $\{p_i\}_{i=1}^{\infty}$ be a frame associated to (a_2, \dots, a_n) for H with bounds A, B and $\{q_j\}_{j=1}^{\infty}$ be a frame associated to (b_2, \dots, b_n) for K with bounds C, D having their corresponding frame operators S_F and S_G , respectively. Then $\{\Delta_{ij} = (U_1 \otimes U_2)(p_i \otimes q_j)\}_{i,j=1}^{\infty}$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$ if and only if $U_1 \otimes U_2$ is an invertible operator on $H_F \otimes K_G$.*

Proof. First we suppose that $U_1 \otimes U_2$ is an invertible on $H_F \otimes K_G$. Then by Theorem 2.5, U_1 and U_2 are invertible on H_F and K_G , respectively. For each $p \in H_F$ and $q \in K_G$, we have

$$(3.8) \quad \|p, a_2, \dots, a_n\|_1 \leq \|U_1^{-1}\| \|U_1^*(p), a_2, \dots, a_n\|_1, \text{ and}$$

$$(3.9) \quad \|q, b_2, \dots, b_n\|_2 \leq \|U_2^{-1}\| \|U_2^*(q), b_2, \dots, b_n\|_2.$$

Now, for each $p \otimes q \in H_F \otimes K_G$, we have

$$\begin{aligned}
&\sum_{i,j=1}^{\infty} |\langle p \otimes q, (U_1 \otimes U_2)(p_i \otimes q_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle|^2 \\
&= \sum_{i,j=1}^{\infty} |\langle p \otimes q, U_1(p_i) \otimes U_2(q_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle|^2 \\
(3.10) \quad &= \sum_{i=1}^{\infty} |\langle U_1^* p, p_i | a_2, \dots, a_n \rangle_1|^2 \sum_{j=1}^{\infty} |\langle U_2^* q, q_j | b_2, \dots, b_n \rangle_2|^2 \\
&\leq B D \|U_1^*(p), a_2, \dots, a_n\|_1^2 \|U_2^*(q), b_2, \dots, b_n\|_2^2 \\
&[\text{since } \{p_i\}_{i=1}^{\infty} \text{ is a frame associated to } (a_2, \dots, a_n), \text{ and} \\
&\{q_j\}_{j=1}^{\infty} \text{ is a frame associated to } (b_2, \dots, b_n)] \\
&\leq B D \|U_1^*\|^2 \|U_2^*\|^2 \|p, a_2, \dots, a_n\|_1^2 \|q, b_2, \dots, b_n\|_2^2 \\
&= B D \|U_1 \otimes U_2\|^2 \|p \otimes q, a_2 \otimes b_2, \dots, a_n \otimes b_n\|^2.
\end{aligned}$$

On the other hand, from (3.10)

$$\begin{aligned}
 & \sum_{i,j=1}^{\infty} |\langle p \otimes q, (U_1 \otimes U_2) (p_i \otimes q_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle|^2 \\
 & \geq AC \|U_1^*(p), a_2, \dots, a_n\|_1^2 \|U_2^*(q), b_2, \dots, b_n\|_2^2 \\
 & \geq \frac{AC \|p, a_2, \dots, a_n\|_1^2 \|q, b_2, \dots, b_n\|_2^2}{\|U_1^{-1}\|^2 \|U_2^{-1}\|^2} \text{ [by (3.8) and (3.9)]} \\
 & = \frac{AC}{\|(U_1 \otimes U_2)^{-1}\|^2} \|p \otimes q, a_2 \otimes b_2, \dots, a_n \otimes b_n\|^2.
 \end{aligned}$$

Therefore, $\{\Delta_{ij}\}_{i,j=1}^{\infty}$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$.

Conversely, suppose that $\{\Delta_{ij}\}_{i,j=1}^{\infty}$ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. Now, for each $p \otimes q \in H_F \otimes K_G$,

$$\begin{aligned}
 & \sum_{i,j=1}^{\infty} \langle p \otimes q, \Delta_{ij} | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle \Delta_{ij} \\
 & = \sum_{i,j=1}^{\infty} \langle p \otimes q, U_1 p_i \otimes U_2 q_j | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle (U_1 p_i \otimes U_2 q_j) \\
 & = \sum_{i=1}^{\infty} \langle U_1^* p, p_i | a_2, \dots, a_n \rangle_1 U_1 p_i \otimes \sum_{j=1}^{\infty} \langle U_2^* q, q_j | b_2, \dots, b_n \rangle_2 U_2 q_j \\
 & = U_1 S_F U_1^*(p) \otimes U_2 S_G U_2^*(q) \\
 & = (U_1 \otimes U_2) (S_F \otimes S_G) (U_1^* \otimes U_2^*) (p \otimes q) \\
 & = (U_1 \otimes U_2) S_{F \otimes G} (U_1 \otimes U_2)^* (p \otimes q).
 \end{aligned}$$

Hence, the frame operator for $\{\Delta_{ij}\}_{i,j=1}^{\infty}$ is $(U_1 \otimes U_2) S_{F \otimes G} (U_1 \otimes U_2)^*$ and therefore it is invertible. Also, we know that $S_{F \otimes G}$ is invertible and hence $U_1 \otimes U_2$ is invertible on $H_F \otimes K_G$. \square

4. DUAL FRAME IN TENSOR PRODUCT OF n -HILBERT SPACES

In this section, dual frame in n -Hilbert spaces and their tensor product are discussed.

Definition 4.1. Let $\{p_i\}_{i=1}^{\infty}$ be a frame associated to (a_2, \dots, a_n) for H . Then a frame $\{q_i\}_{i=1}^{\infty}$ associated to (a_2, \dots, a_n) satisfying

$$p = \sum_{i=1}^{\infty} \langle p, q_i | a_2, \dots, a_n \rangle_1 p_i \quad \forall p \in H$$

is called a dual frame or alternative dual frame associated to (a_2, \dots, a_n) of $\{p_i\}_{i=1}^\infty$.

Theorem 4.2. *Let $\{p_i\}_{i=1}^\infty$ and $\{q_i\}_{i=1}^\infty$ be two Bessel sequences associated to (a_2, \dots, a_n) in H . Then the following are equivalent:*

$$(i) \quad p = \sum_{i=1}^{\infty} \langle p, q_i | a_2, \dots, a_n \rangle_1 p_i; \quad \forall p \in H_F.$$

$$(ii) \quad p = \sum_{i=1}^{\infty} \langle p, p_i | a_2, \dots, a_n \rangle_1 q_i; \quad \forall p \in H_F.$$

Proof. (i) \Rightarrow (ii) Let T_F and T_G be the pre-frame operators of $\{p_i\}_{i=1}^\infty$ and $\{q_i\}_{i=1}^\infty$, respectively. Composing T_F with the adjoint of T_G , for all $p \in H_F$, we get

$$T_F T_G^* : H_F \rightarrow H_F, \quad T_F T_G^*(p) = \sum_{i=1}^{\infty} \langle p, q_i | a_2, \dots, a_n \rangle_1 p_i.$$

Now, in terms of pre-frame operators (i) can be written as $T_F T_G^* = I_F$ and this equivalent to $T_G T_F^* = I_F$. Therefore, for each $p \in H_F$,

$$p = \sum_{i=1}^{\infty} \langle p, p_i | a_2, \dots, a_n \rangle_1 q_i.$$

Similarly, (ii) \Rightarrow (i) follows. \square

Remark 4.3. Suppose that the equivalent conditions of Theorem 4.2 are satisfied. Then using Cauchy-Schwartz inequality, for every $p \in H_F$, we have

$$\begin{aligned} \|p, a_2, \dots, a_n\|_1^2 &= \langle p, p | a_2, \dots, a_n \rangle_1 \\ &= \left\langle \sum_{i=1}^{\infty} \langle p, p_i | a_2, \dots, a_n \rangle_1 q_i, p | a_2, \dots, a_n \right\rangle_1 \\ &= \sum_{i=1}^{\infty} \langle p, p_i | a_2, \dots, a_n \rangle_1 \langle q_i, p | a_2, \dots, a_n \rangle_1 \\ &\leq \left(\sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle_1|^2 \right)^{1/2} \left(\sum_{i=1}^{\infty} |\langle p, q_i | a_2, \dots, a_n \rangle_1|^2 \right)^{1/2} \\ &\leq \left(\sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle_1|^2 \right)^{1/2} B^{1/2} \|p, a_2, \dots, a_n\|_1 \end{aligned}$$

[since $\{q_i\}_{i=1}^\infty$ is a Bessel sequences associated to (a_2, \dots, a_n)]

$$\Rightarrow \frac{1}{B} \|p, a_2, \dots, a_n\|_1^2 \leq \sum_{i=1}^{\infty} |\langle p, p_i | a_2, \dots, a_n \rangle_1|^2.$$

This shows that $\{p_i\}_{i=1}^\infty$ is a frame associated to (a_2, \dots, a_n) for H . Similarly, it can be shown that $\{q_i\}_{i=1}^\infty$ is also a frame associated to (a_2, \dots, a_n) for H .

We now present the concept of a dual frame in $H \otimes K$.

Remark 4.4. Let $\{p_i \otimes q_j\}_{i,j=1}^\infty$ be a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. Then according to the definition 4.1, a frame $\{e_i \otimes h_j\}_{i,j=1}^\infty$ associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$ satisfying

$$(4.1) \quad p \otimes q = \sum_{i,j=1}^{\infty} \langle p \otimes q, e_i \otimes h_j | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle (p_i \otimes q_j),$$

for all $p \otimes q \in H \otimes K$, can be consider as a dual frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ of $\{p_i \otimes q_j\}_{i,j=1}^\infty$.

Remark 4.5. According to the Theorem 3.2, $\{p_i \otimes q_j\}_{i,j=1}^\infty$ and $\{e_i \otimes h_j\}_{i,j=1}^\infty$ are pair of dual frames associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$ if and only if $\{p_i \otimes q_j\}_{i,j=1}^\infty$ and $\{e_i \otimes h_j\}_{i,j=1}^\infty$ are pair of dual frames for $H_F \otimes K_G$.

Theorem 4.6. Let $\{p_i\}_{i=1}^\infty, \{e_i\}_{i=1}^\infty$ be a pair of dual frames associated to (a_2, \dots, a_n) for H and $\{q_j\}_{j=1}^\infty, \{h_j\}_{j=1}^\infty$ be a pair of dual frames associated to (b_2, \dots, b_n) for K . Then $\{e_i \otimes h_j\}_{i,j=1}^\infty$ is a dual frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ of $\{p_i \otimes q_j\}_{i,j=1}^\infty$.

Proof. By Theorem 3.8, $\{p_i \otimes q_j\}_{i,j=1}^\infty, \{e_i \otimes h_j\}_{i,j=1}^\infty$ are frames associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. Since $\{e_i\}_{i=1}^\infty$ and $\{h_j\}_{j=1}^\infty$ are dual frames associated to (a_2, \dots, a_n) and (b_2, \dots, b_n) of $\{p_i\}_{i=1}^\infty$ and $\{q_j\}_{j=1}^\infty$, respectively, for all $p \in H, q \in K$,

$$p = \sum_{i=1}^{\infty} \langle p, e_i | a_2, \dots, a_n \rangle_1 p_i, \quad \text{and} \quad q = \sum_{j=1}^{\infty} \langle q, h_j | b_2, \dots, b_n \rangle_2 q_j.$$

Then, for all $p \otimes q \in H \otimes K$, we have

$$\begin{aligned} p \otimes q &= \left(\sum_{i=1}^{\infty} \langle p, e_i | a_2, \dots, a_n \rangle_1 p_i \right) \otimes \left(\sum_{j=1}^{\infty} \langle q, h_j | b_2, \dots, b_n \rangle_2 q_j \right) \\ &= \sum_{i,j=1}^{\infty} \langle p, e_i | a_2, \dots, a_n \rangle_1 \langle q, h_j | b_2, \dots, b_n \rangle_2 (p_i \otimes q_j) \end{aligned}$$

$$= \sum_{i,j=1}^{\infty} \langle p \otimes q, e_i \otimes h_j | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle (p_i \otimes q_j).$$

This completes the proof. \square

Theorem 4.7. *Let $\{p_i\}_{i=1}^{\infty}, \{e_i\}_{i=1}^{\infty}$ be a pair of dual frames associated to (a_2, \dots, a_n) for H and $\{q_j\}_{j=1}^{\infty}, \{h_j\}_{j=1}^{\infty}$ be a pair of dual frames associated to (b_2, \dots, b_n) for K . Suppose $U \in \mathcal{B}(H_F)$ and $V \in \mathcal{B}(K_G)$ are unitary operators. Then $\Lambda = \{(U \otimes V)(p_i \otimes q_j)\}_{i,j=1}^{\infty}$ and $\Gamma = \{(U \otimes V)(e_i \otimes h_j)\}_{i,j=1}^{\infty}$ also form a pair of dual frames associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$.*

Proof. By Theorem 4.6, $\{p_i \otimes q_j\}_{i,j=1}^{\infty}$ and $\{e_i \otimes h_j\}_{i,j=1}^{\infty}$ form a pair of dual frames associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. Now, for each $p \otimes q \in H_F \otimes K_G$, we have

$$\begin{aligned} & \sum_{i,j=1}^{\infty} |\langle p \otimes q, (U \otimes V)(e_i \otimes h_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle|^2 \\ &= \sum_{i,j=1}^{\infty} |\langle p \otimes q, (U e_i \otimes V h_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle|^2 \\ &= \sum_{i=1}^{\infty} |\langle U^* p, e_i | a_2, \dots, a_n \rangle_1|^2 \otimes \sum_{j=1}^{\infty} |\langle V^* q, h_j | b_2, \dots, b_n \rangle_2 q_j|^2. \end{aligned}$$

Since $\{e_i\}_{i=1}^{\infty}$ is a frame associated to (a_2, \dots, a_n) for H and $\{h_j\}_{j=1}^{\infty}$ is a frame associated to (b_2, \dots, b_n) for K , the above calculation shows that Γ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. Similarly, it can be shown that Λ is a frame associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. Furthermore, for each $p \otimes q \in H_F \otimes K_G$, we have

$$\begin{aligned} & \sum_{i,j=1}^{\infty} \langle p \otimes q, (U \otimes V)(e_i \otimes h_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle (U \otimes V)(p_i \otimes q_j) \\ &= \sum_{i,j=1}^{\infty} \langle p \otimes q, (U e_i \otimes V h_j) | a_2 \otimes b_2, \dots, a_n \otimes b_n \rangle (U p_i \otimes V q_j) \\ &= U \sum_{i=1}^{\infty} \langle U^* p, e_i | a_2, \dots, a_n \rangle_1 p_i \otimes V \sum_{j=1}^{\infty} \langle V^* q, h_j | b_2, \dots, b_n \rangle_2 q_j \\ &= U U^*(p) \otimes V V^*(q) = p \otimes q \text{ [since } U, V \text{ are unitary operators]}. \end{aligned}$$

[Since $\{p_i\}_{i=1}^{\infty}, \{e_i\}_{i=1}^{\infty}$ are dual frames associated to (a_2, \dots, a_n) , and $\{q_j\}_{j=1}^{\infty}, \{h_j\}_{j=1}^{\infty}$ are dual frames associated to (b_2, \dots, b_n)].

Hence, according to the remark 4.5, Λ and Γ form a pair of dual frames associated to $(a_2 \otimes b_2, \dots, a_n \otimes b_n)$ for $H \otimes K$. \square

Acknowledgment. The authors would like to thank the editor and the referees for their helpful suggestions and comments to improve this paper.

REFERENCES

1. H. Bolcskei, F. Hlawatsch and H. G. Feichtinger, *Frame-theoretic analysis of oversampled filter banks*, IEEE Trans. Signal Process., 46 (1998), pp. 3256-3268.
2. P. G. Casazza and G. Kutyniok, *Frames of subspaces*, Contemp. Math., 345 (2004), pp. 87-114.
3. O. Christensen, *An introduction to frames and Riesz bases*, (Applied and Numerical Harmonic Analysis), Birkhauser, 2008.
4. I. Daubechies, A. Grossmann and Y. Meyer, *Painless nonorthogonal expansions*, J. Math. Phys., 27 (1986), pp. 1271-1283.
5. C. Diminnie, S. Gahler and A. White, *2-inner product spaces*, Demonstratio Mathematica, 6 (1973), pp. 525-536.
6. N. E. Dudev Ward and J. R. Partington, *A construction of rational wavelets and frames in Hardy-Sobolev space with applications to system modeling*, SIAM J. Control Optim., 36 (1998), pp. 654-679.
7. R. J. Duffin and A. C. Schaeffer, *A class of nonharmonic Fourier series*, Trans. Amer. Math. Soc., 72 (1952), pp. 341-366.
8. P. J. S. G. Ferreira, *Mathematics for multimedia signal processing II: Discrete finite frames and signal reconstruction*, In: Byrnes, J. S. (ed.) Signal processing for multimedia, IOS Press, Amsterdam (1999), pp. 35-54.
9. G. B. Folland, *A Course in Abstract Harmonic Analysis*, CRC Press, Boca Raton (1995).
10. D. Gabor, *Theory of communications*, J. Inst. Elec. Engrg., 93 (1946), pp. 429-457.
11. P. Ghosh and T. K. Samanta, *Stability of dual g -fusion frame in Hilbert spaces*, Methods Funct. Anal. Topol., 26 (2020), pp. 227-240.
12. P. Ghosh and T. K. Samanta, *Construction of frame relative to n -Hilbert space*, J. Linear. Topological. Algebra., 10 (2021), pp. 117-130.
13. P. Ghosh and T. K. Samanta, *Generalized atomic subspaces for operators in Hilbert spaces*, Mathematica Bohemica, Accepted.
14. H. Gunawan and M. Mashadi, *On n -normed spaces*, Int. J. Math. Math. Sci., 27 (2001), pp. 631-639.

15. R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras*. Vol. I, Graduate Studies in Mathematics, vol. 15. American Mathematical Society, Providence (1997) (Elementary theory, Reprint of the 1983 original).
16. A. Misiak, *n-inner product spaces*, Math. Nachr., 140 (1989), pp. 299-319.
17. S. Rabinson, *Hilbert space and tensor products*, Lecture notes September 8, 1997.
18. G. Upendar Reddy, N. Gopal Reddy and B. Krishna Reddy, *Frame operator and Hilbert-Schmidt Operator in Tensor Product of Hilbert Spaces*, J. Dyn. Syst. Geom. Theor., 7 (2009), pp. 61-70.
19. V. Sadri, Gh. Rahimlou, R. Ahmadi and Ramazan Zarghami Farfar, *Generalized Fusion Frames in Hilbert Spaces*, Inf. Dim. Anal. Quan. Prob. (IDA-QP), Vol. 23, No. 02, 2050015 (2020).
20. T. Strohmer and R. Jr. Heath, *Grassmanian frames with applications to coding and communications*, Appl. Comput. Harmon. Anal., 14 (2003), pp. 257-275.
21. W. Sun, *G-frames and G-Riesz bases*, J. Math. Anal. Appl., 322 (2006), pp. 437-452.

¹ DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF CALCUTTA, 35, BALYGUNGE CIRCULAR ROAD, KOLKATA, 700019, WEST BENGAL, INDIA.

Email address: prasenjitpuremath@gmail.com

² DEPARTMENT OF MATHEMATICS, ULUBERIA COLLEGE, ULUBERIA, HOWRAH, 711315, WEST BENGAL, INDIA.

Email address: mumpu_tapas5@yahoo.co.in