

Han's conjecture for bounded extensions

Claude Cibils, Marcelo Lanzilotta, Eduardo N. Marcos,
and Andrea Solotar *

Abstract

Let $B \subset A$ be a left or right bounded extension of finite dimensional algebras. We use the Jacobi-Zariski long nearly exact sequence to show that B satisfies Han's conjecture if and only if A does, regardless if the extension splits or not. We provide conditions ensuring that an extension by arrows and relations is left or right bounded. Finally we give a structure result for extensions of an algebra given by a quiver and admissible relations, and examples of non split left or right bounded extensions.

2020 MSC: 18G25, 16E40, 16E30, 18G15

Keywords: Hochschild, homology, relative, Han, quiver

1 Introduction

Let k be a field. We recall that for noetherian rings the left and right global dimension coincide, see [4]. Han's conjecture (see [13]) states that for a finite dimensional algebra A , if $H_*(A, A) = 0$ in large enough degrees, then A has finite global dimension. In [10] we have included a brief account of the main advances towards proving this conjecture, see [9, 5, 7, 8, 6, 21, 20, 11].

An extension $B \subset A$ of k -algebras is called left (resp. right) bounded if the B -bimodule A/B is B -tensor nilpotent, its projective dimension is finite and it is left (resp. right) B -projective. A main result of this paper is that for a left or right bounded extension $B \subset A$, the k -algebra B satisfies Han's conjecture if and only if A does. Notice that we do not assume the extension to be split.

Actually the purpose of this paper is twofold. Firstly to provide results which may be used to associate to a given algebra A an easier algebra B for which the property of satisfying Han's conjecture is equivalent to the same property for A . In [11, 10] a theory is developed for split extensions. Here we obtain results that allow us to deal with non split extensions. Even if there are algebras to which we cannot apply our methods, our guiding idea is that, starting from a given finite dimensional algebra, we can reduce the study of Han's conjecture about it to the same problem for an algebra where the computation of the Hochschild homology is more tractable. The second aim of the paper is to provide examples of this procedure. We develop

*This work has been supported by the projects UBACYT 20020170100613BA, PIP-CONICET 11220150100483CO, USP-COFECUB. The third mentioned author was supported by the thematic project of FAPESP 2014/09310-5, a research grant from CNPq 302003/2018-5 and acknowledges support from the "Brazilian-French Network in Mathematics". The fourth mentioned author is a research member of CONICET (Argentina), Senior Associate at ICTP and visiting Professor at Guangdong Technion-Israel Institute of Technology.

a theory of extensions by arrows and relations and we give conditions ensuring that the extension is left or right bounded. We also show that all extensions of an algebra given by a quiver and admissible relations are obtained through our construction in Section 5, Theorem 5.8.

We describe in what follows the structure of the paper in more detail.

Let $B \subset A$ be an extension of k -algebras. G. Hochschild defined Hochschild (resp. relative) homology with coefficients in an A -bimodule X in [14] (resp. [15]). These are vector spaces denoted $H_*(A, X)$ (resp. $H_*(A|B, X)$), for $* \geq 0$. See also the books [19, 22, 23] for a general account of Hochschild theory.

The proof of the above mentioned main result of this paper relies on the normalised relative bar resolution and the Jacobi-Zariski long nearly exact sequence obtained in [12]. We give a brief account of these results in Section 2.

It is worth noting that in Sections 3 and 4 we consider extensions of algebras which may be infinite dimensional. We first show that if an extension $B \subset A$ is left or right bounded then the Hochschild homologies of A and B are isomorphic in large enough degrees. Then we prove in Section 4 that for a left or right bounded extension, finite global dimension is preserved. Note that for the latter, the normalised relative bar resolution is also a crucial tool. We end Section 4 by proving the main quoted result about Han's conjecture for left or right bounded extensions of finite dimensional algebras.

In Section 5 we construct explicit extensions of an algebra $B = kQ/I$ where Q is a quiver and I is an admissible ideal of the path algebra kQ . A structure result is proved, which shows that this construction provides every B -finitely generated algebra extension of B . In detail, we first add new arrows to Q , to obtain a tensor algebra extension T of B . Then we add new relations through an ideal $J \subset T$ verifying $B \cap J = 0$ to obtain an extension $B \subset A = T/J$. We focus on extensions $B \subset A$ of this sort to give conditions ensuring that the B -bimodule A/B is tensor nilpotent and/or projective on one side over B . We use relative paths, they are alternate concatenations of new arrows and non-zero paths of B . If every relative cycle contains a J -interrupter (see Definition 5.13), then A/B is B -tensor nilpotent. If the projection of J to the positive part of T is generated as a B -bimodule by relative paths ending in the same new arrow, then A/B is left projective as a B -module.

Finally in Section 6 we consider examples of extensions by arrows and relations. The above criteria enables to prove that they are left or right bounded, and we show that they are not split.

We thank Konstantyn Iusenko and John MacQuarrie for their very pertinent comments on an earlier version of this article. We thank the referee for an attentive reading and useful suggestions.

2 Normalised relative bar resolution and the Jacobi-Zariski sequence

In this section we recall some results from [12] that will be useful for the sequel. We first recall the normalised relative bar resolution.

Proposition 2.1 [12, Proposition 2.3] *Let $B \subset A$ be an extension of k -algebras and let σ be a k -linear section of the canonical projection $\pi : A \rightarrow A/B$. The*

following is a B -relative resolution of A :

$$\cdots \xrightarrow{d} A \otimes_B (A/B)^{\otimes_B m} \otimes_B A \xrightarrow{d} \cdots \xrightarrow{d} A \otimes_B A/B \otimes_B A \xrightarrow{d} A \otimes_B A \xrightarrow{d} A \rightarrow 0$$

where the last d is the product of A and

$$\begin{aligned} d(a_0 \otimes \alpha_1 \otimes \cdots \otimes \alpha_{n-1} \otimes a_n) &= a_0 \sigma(\alpha_1) \otimes \alpha_2 \otimes \cdots \otimes \alpha_{n-1} \otimes a_n + \\ &\quad \sum_{i=1}^{n-2} (-1)^i a_0 \otimes \alpha_1 \otimes \cdots \otimes \pi(\sigma(\alpha_i) \sigma(\alpha_{i+1})) \otimes \cdots \otimes \alpha_{n-1} \otimes a_n + \\ &\quad (-1)^{n-1} a_0 \otimes \alpha_1 \otimes \cdots \otimes \sigma(\alpha_{n-1}) a_n. \end{aligned}$$

Moreover the differential d does not depend on the choice of the section σ .

The normalised relative bar resolution provides a chain complex for computing the relative Hochschild homology as follows.

Theorem 2.2 [12, Corollary 2.4] *Let $B \subset A$ be an extension of k -algebras and let X be an A -bimodule. Let σ be a k -linear section of $\pi : A \rightarrow A/B$. The homology of the following chain complex is the relative Hochschild homology $H_*(A|B, X)$*

$$\cdots \xrightarrow{b} X \otimes_{B^e} (A/B)^{\otimes_B m} \xrightarrow{b} \cdots \xrightarrow{b} X \otimes_{B^e} A/B \xrightarrow{b} X_B \rightarrow 0 \quad (2.1)$$

where

$$\begin{aligned} b(x \otimes \alpha_1 \otimes \cdots \otimes \alpha_n) &= x \sigma(\alpha_1) \otimes \alpha_2 \otimes \cdots \otimes \alpha_n + \\ &\quad \sum_{i=1}^{n-1} (-1)^i x \otimes \alpha_1 \otimes \cdots \otimes \pi(\sigma(\alpha_i) \sigma(\alpha_{i+1})) \otimes \cdots \otimes \alpha_n + \\ &\quad (-1)^n \sigma(\alpha_n) x \otimes \alpha_1 \otimes \cdots \otimes \alpha_{n-1}. \end{aligned}$$

The coboundary b does not depend on the choice of the section σ .

Definition 2.3 *Let B be a k -algebra. A B -bimodule M is tensor nilpotent if there exists n such that $M^{\otimes_B n} = 0$.*

As usual, we write “ $* \gg 0$ ” for “large enough $*$ ”, that is “all $*$ larger than some positive integer”.

Corollary 2.4 *Let $B \subset A$ be an extension of k -algebras with A/B tensor nilpotent. Let X be an A -bimodule. We have*

$$H_*(A|B, X) = 0 \text{ for } * \gg 0.$$

Proof.

The complex of Theorem 2.2 is zero in large enough degrees. \diamond

Next we recall the definition of a long sequence of vector spaces which is exact twice in three, as introduced in [12].

Definition 2.5 *A long nearly exact sequence is a complex of vector spaces*

$$\cdots \xrightarrow{\delta} U_m \xrightarrow{I} V_m \xrightarrow{K} W_m \xrightarrow{\delta} U_{m-1} \xrightarrow{I} V_{m-1} \rightarrow \cdots \xrightarrow{\delta} U_n \xrightarrow{I} V_n \xrightarrow{K} W_n$$

ending at a fixed $n \geq 0$ which is exact at U_* and W_* .

Its gap is the homology at V_* , namely the vector space $(\text{Ker}K/\text{Im}I)_*$.

A. Kaygun has first obtained the Jacobi-Zariski long exact sequence in the non-commutative setting for a k -algebra extension $B \subset A$ where A/B is flat as a B -bimodule, see [17, 18]. Note that for commutative algebras the Jacobi-Zariski long exact sequence has been established for André-Quillen homology in 1974, see for instance [3, p. 61] or [16]. The reason for giving the name “Jacobi-Zariski” to this sequence is explained in [3, p. 102].

We state now the Jacobi-Zariski long nearly exact sequence obtained in [12].

Theorem 2.6 [12, Theorem 4.4 and Theorem 5.1] *Let $B \subset A$ be an extension of k -algebras. Let X be an A -bimodule. There is a long nearly exact sequence*

$$\begin{aligned} \dots \xrightarrow{\delta} H_m(B, X) \xrightarrow{I} H_m(A, X) \xrightarrow{K} H_m(A|B, X) \xrightarrow{\delta} H_{m-1}(B, X) \xrightarrow{I} \dots \\ \xrightarrow{\delta} H_1(B, X) \xrightarrow{I} H_1(A, X) \xrightarrow{K} H_1(A|B, X) \end{aligned}$$

which is exact at $H_*(B, X)$ and $H_*(A|B, X)$.

Moreover, there is a spectral sequence converging to its gap $(\text{Ker}K/\text{Im}I)_*$ for $* \geq 2$. If $\text{Tor}_*^B(A/B, (A/B)^{\otimes_B n}) = 0$ for $* > 0$ and for all n , its terms at page 1 are

$$E_{p,q}^1 = \text{Tor}_{p+q}^{B^e}(X, (A/B)^{\otimes_B p}) \quad \text{for } p, q > 0$$

and 0 elsewhere.

Definition 2.7 *An extension $B \subset A$ of k -algebras is left (resp. right) bounded if*

- A/B is tensor nilpotent,
- A/B is of finite projective dimension as a B^e -module,
- A/B is a left (resp. right) projective B -module.

Remark 2.8 *The hypothesis of Theorem 2.6 for describing the first page of the spectral sequence is fulfilled if A/B is a left or right bounded extension.*

Theorem 2.9 [12, Theorem 6.5]

Let $B \subset A$ be a left or right bounded extension of k -algebras and let X be an A -bimodule.

There is a Jacobi-Zariski long exact sequence ending at some n :

$$\begin{aligned} \dots \xrightarrow{\delta} H_m(B, X) \xrightarrow{I} H_m(A, X) \xrightarrow{K} H_m(A|B, X) \xrightarrow{\delta} H_{m-1}(B, X) \xrightarrow{I} \dots \\ \xrightarrow{\delta} H_n(B, X) \xrightarrow{I} H_n(A, X) \xrightarrow{K} H_n(A|B, X). \end{aligned}$$

3 Hochschild homology of bounded extensions

In this section we prove that for any left or right bounded extension $B \subset A$ the Hochschild homology remains unchanged in large enough degrees.

Theorem 3.1 *Let $B \subset A$ be an extension of k -algebras, where the B -bimodule A/B is of finite projective dimension and tensor nilpotent. There is a linear injection*

$$H_*(B, B) \hookrightarrow H_*(A, A) \quad \text{for } * \gg 0.$$

If moreover the extension is left or right bounded then

$$H_*(B, B) \simeq H_*(A, A) \quad \text{for } * \gg 0.$$

Proof. Firstly we assert that if A/B is of finite projective dimension as B -bimodule, then $H_*(B, B)$ and $H_*(B, A)$ are isomorphic for $* \gg 0$. Indeed, the exact sequence of B -bimodules

$$0 \rightarrow B \rightarrow A \rightarrow A/B \rightarrow 0. \quad (3.1)$$

induces a long exact sequence in Hochschild homology

$$\begin{aligned} \cdots \rightarrow H_m(B, B) \rightarrow H_m(B, A) \rightarrow H_m(B, A/B) \rightarrow H_{m-1}(B, B) \rightarrow \cdots \\ \rightarrow H_0(B, B) \rightarrow H_0(B, A) \rightarrow H_0(B, A/B) \rightarrow 0. \end{aligned}$$

By definition for any B -bimodule M

$$H_*(B, M) = \mathrm{Tor}_*^{B^e}(B, M).$$

Since $M = A/B$ is of finite projective dimension as a B^e -module, we infer that $H_*(B, A/B) = 0$ for $* \gg 0$. The assertion follows.

Next we consider the Jacobi-Zariski long nearly exact sequence of Theorem 2.6 for an A -bimodule X . It is exact at $H_*(B, X)$, that is $\mathrm{Ker}I = \mathrm{Im}\delta$. Since A/B is B -tensor nilpotent, by Corollary 2.4 we have that $H_*(A|B, X) = 0$ for $* \gg 0$, thus for large enough degrees we get $\delta = 0$. Hence the maps $I : H_*(B, X) \rightarrow H_*(A, X)$ are injective for $* \gg 0$. For $X = A$, we infer from the previous assertion that

$$H_*(B, B) \hookrightarrow H_*(A, A) \text{ for } * \gg 0.$$

Note that the gap of the Jacobi-Zariski long nearly exact sequence is

$$H_*(A, A)/H_*(B, B) \text{ for } * \gg 0.$$

If the extension is left or right bounded then by Theorem 2.9 the Jacobi-Zariski sequence is exact in degrees larger than some n . Hence the gap is zero in large enough degrees and $H_*(A, A) \simeq H_*(B, B)$ for $* \gg 0$. \diamond

4 Bounded extensions, finite global dimension and Han's conjecture

4.1 Finite global dimension

We consider extensions $B \subset A$ with the aim of studying the relation between the finiteness of the global dimension B and A . For completeness we first recall the following.

Lemma 4.1 *Let $B \subset A$ be an extension of k -algebras, with A/B of finite projective dimension s as a left B -module. Any left projective A -module is of projective dimension at most s as a left B -module.*

Proof. The exact sequence (3.1) shows that A has projective dimension at most s as a left B -module. Then the result holds with the same bound for the projective dimension for any left projective A -module by standard arguments. \diamond

We recall that for noetherian rings the left and right global dimensions coincide, see [4].

Theorem 4.2 *Let $B \subset A$ be an extension of k -algebras where A has left finite global dimension a . If A/B is of finite projective dimension r as a B -bimodule and projective as a B -right module, then B has left global dimension at most $r + a$.*

Proof. Let Y be a left B -module. The exact sequence of B -bimodules (3.1) is right split, hence the sequence of left B -modules

$$0 \rightarrow B \otimes_B Y \rightarrow A \otimes_B Y \rightarrow (A/B) \otimes_B Y \rightarrow 0 \quad (4.1)$$

is exact. Of course $B \otimes_B Y$ is canonically isomorphic to Y . In order to prove that Y is of bounded projective dimension as a left B -module, we will prove that this is the case for $A \otimes_B Y$ and $(A/B) \otimes_B Y$.

Let $P_* \rightarrow A \otimes_B Y \rightarrow 0$ be a projective resolution of length a of the A -module $A \otimes_B Y$. It is an exact sequence of left B -modules by restricting the left actions to B .

By Lemma 4.1 each P_i is of projective dimension at most r as a left B -module. We infer that $A \otimes_B Y$ is of projective dimension at most $r + a$ as a left B -module.

For $(A/B) \otimes_B Y$, let $Q_* \rightarrow (A/B) \otimes_B Y \rightarrow 0$ be a finite projective resolution of the B -bimodule A/B of length r . Each Q_i is projective as a B -bimodule, then it is projective as a right B -module. Moreover A/B is projective as a right B -module. Hence $Q_* \otimes_B Y \rightarrow (A/B) \otimes_B Y \rightarrow 0$ is exact, and it is a finite projective resolution of the left B -module $(A/B) \otimes_B Y$ of length at most r . As a consequence, the projective dimension of the left B -module Y is at most $r + a + 1$. \diamond

Theorem 4.3 *Let $B \subset A$ be an extension of k -algebras where B has left finite global dimension b . Suppose that the B -bimodule A/B is tensor nilpotent and let n be the smallest positive integer such that $(A/B)^{\otimes_B n} = 0$. Suppose that A/B is projective as a B -right module. Then A has left global dimension at most $n + b - 1$.*

Proof. First we will consider A -modules of the form $A \otimes_B Y$ where Y is a left B -module, to show that they are of projective dimension at most b . Let $P_* \rightarrow Y$ be a projective resolution of Y as a B -module of length at most b . Since the exact sequence (3.1) is right split, observe that A is projective as a right B -module. Hence $A \otimes_B P_* \rightarrow A \otimes_B Y \rightarrow 0$ is exact. Moreover $A \otimes_B P_i$ is projective as a left A -module. Indeed this holds if P_i is a free left B -module, thus it holds for projectives by standard arguments. So the projective dimension of the A -module $(A \otimes_B Y)$ is at most b .

Let now X be an arbitrary A -module. We assert that there is an exact sequence of A -modules of the form

$$0 \rightarrow A \otimes_B Y_{n-1} \rightarrow \cdots \rightarrow A \otimes_B Y_0 \rightarrow X \rightarrow 0$$

where the Y_i 's are left B -modules. Note that the assertion shows that X is of projective dimension at most $n - 1 + b$.

To prove the assertion we use that the normalised bar resolution of Theorem 2.1 is zero in degrees $\geq n$. We observe that the contracting homotopy of this resolution is given by morphisms of right A -modules. Then the following sequence is exact:

$$0 \rightarrow A \otimes_B (A/B)^{\otimes_B n-1} \otimes_B A \otimes_A X \xrightarrow{d} \cdots \xrightarrow{d} A \otimes_B (A/B) \otimes_B A \otimes_A X \xrightarrow{d} A \otimes_B A \otimes_A X \xrightarrow{d} A \otimes_A X \rightarrow 0,$$

as well as the isomorphic sequence

$$0 \rightarrow A \otimes_B (A/B)^{\otimes_B n-1} \otimes_B X \xrightarrow{d} \cdots \xrightarrow{d} A \otimes_B (A/B) \otimes_B X \xrightarrow{d} A \otimes_B X \xrightarrow{d} X \rightarrow 0.$$

The modules $A \otimes_B ((A/B)^{\otimes_B i} \otimes_B X)$ are of the form $A \otimes_B Y$. By the above result they are of projective dimension at most b as left A -modules. Hence X is of projective dimension at most $n - 1 + b$. \diamond

Corollary 4.4 *Let $B \subset A$ be a right bounded extension of k -algebras. The algebra B has left finite global dimension if and only if A has left finite global dimension.*

Remark 4.5 *Analogously, if an extension is left bounded then right finiteness of the global dimension is preserved.*

4.2 Han's conjecture for bounded extensions

Next we use the results that we have obtained to prove the following. Recall that if a k -algebra is finite dimensional then the left and right global dimension are equal.

Theorem 4.6 *Let $B \subset A$ be a left or right bounded extension of finite dimensional k -algebras. The algebra B satisfies Han's conjecture if and only if A satisfies Han's conjecture.*

Proof. Assume that B satisfies Han's conjecture. Suppose $H_*(A, A) = 0$ for $* \gg 0$. Since the extension verifies that the B -bimodule A/B is of finite projective dimension and is tensor nilpotent, the first part of Theorem 3.1 gives that $H_*(B, B) = 0$ for $* \gg 0$, and then B has finite global dimension. Moreover the B -bimodule A/B is of finite projective dimension and also that A/B is a B -left (or right) projective module. Hence by Theorem 4.3 If B has finite global dimension then A also have it.

Conversely, assume that A satisfies Han's conjecture and $H_*(B, B) = 0$ for $* \gg 0$. The extension is left or right bounded, hence the second part of Theorem 3.1 gives that $H_*(A, A) = 0$ for $* \gg 0$ and therefore the algebra A has finite global dimension. The B -bimodule A/B is of finite projective dimension and A/B is projective as a left or right projective module. Theorem 4.2 ensures that B has finite global dimension. \diamond

5 Extensions by arrows and relations

In this section we define extensions of algebras given by arrows and relations, in order to provide conditions which ensure that the extensions are left or right bounded. We also give a structure result, showing that any finitely generated extension is obtained as an extension by arrows and relations.

Let Q be a finite quiver, that is a finite set of vertices Q_0 , a finite set of arrows Q_1 and two maps $s, t : Q_1 \rightarrow Q_0$ called source and target respectively. A path of length $n > 0$ is a sequence of concatenated arrows $b_n \dots b_1$, that is verifying $t(b_i) = s(b_{i+1})$ for $i = 1, \dots, n-1$. The paths of length zero are the vertices. Let Q_n be the set of paths of length n and consider the path algebra $kQ = \bigoplus_{i \geq 0} kQ_i$.

Let $\langle Q_i \rangle$ be the ideal generated by Q_i . An ideal I of kQ is *admissible* if for some $n \geq 2$ we have $\langle Q_n \rangle \subset I \subset \langle Q_2 \rangle$. Let $B = kQ/I$. It is well known that if k is algebraically closed, any finite dimensional k -algebra is Morita equivalent to an algebra of this type.

Definition 5.1 Let Q be a quiver. A set F of new arrows for Q is a finite set with two maps $s, t : F \rightarrow Q_0$. The quiver Q_F is given by $(Q_F)_0 = Q_0$ and $(Q_F)_1 = Q_1 \cup F$ with the evident maps $s, t : (Q_F)_1 \rightarrow Q_0$.

Let $B = kQ/I$ as above, let F be a set of new arrows for Q , and let $\langle I \rangle_{kQ_F}$ be the ideal of kQ_F generated by I . Let $T = kQ_F/\langle I \rangle_{kQ_F}$. Since $\langle I \rangle_{kQ_F} \cap kQ = I$ we have $B \subset T$. It is useful to observe that T is a tensor algebra over B of a projective B -bimodule as follows. Let

$$N = \bigoplus_{a \in F} Bt(a) \otimes s(a)B.$$

We have

$$T = T_B(N) = B \oplus N \oplus (N \otimes_B N) \oplus \dots$$

Definition 5.2 An extension by arrows and relations $B \subset A$ is given by

- $B = kQ/I$ as above,
- F a set of new arrows for Q ,
- J an ideal of $T = kQ_F/\langle I \rangle_{kQ_F}$ verifying $J \cap B = 0$.

The extended algebra is $A = T/J$.

Example 5.3 Let $B = kQ/I$ where Q is the full arrows quiver

$$\begin{array}{ccc} 2 & \xleftarrow{b} & 3 \\ & \downarrow a & \uparrow c \\ 5 & \xrightarrow{\alpha} & 1 \xrightarrow{d} 4 \\ & \xleftarrow{\beta} & \end{array}$$

and $I = \langle \beta\alpha \rangle$. Let $F = \{a\}$ be the dashed new arrow from 2 to 1, and let $J = \langle abcd - \alpha\beta \rangle$. Then $A = kQ_F/\langle \beta\alpha, abcd - \alpha\beta \rangle$.


Example 5.4 Let Q be a quiver and I be an admissible ideal of kQ . The algebra $\Lambda = kQ/I$ is an extension by arrows and relations of its maximal semisimple subalgebra $E = kQ_0$, where $F = Q_1$ and $J = I$.

In this case the extension $E \subset \Lambda$ is split since $\Lambda = E \oplus r$ where r is the Jacobson radical of Λ . Moreover it is well known and easy to prove that the E -bimodule $\Lambda/E = r$ is E -tensor nilpotent if and only if Q has no oriented cycles.


The next example is a relation extension algebra, see [1, 2]. The family of relation extension algebras is related with cluster theory. The example below consists in an algebra B of global dimension 2, a B -bimodule $M = \text{Ext}^2(DB, B)$ with trivial multiplication structure. The algebra A is the trivial extension $B \oplus M$.

Example 5.5 See [1, Example 2.7] and [2, Example 5.3].

Let $B = kQ/I$ where Q is the quiver



and $I = \langle ba \rangle$. Let $F = \{d\}$ be a new arrow d from 3 to 1. The quiver Q_F is



Let $J = \langle ad, db, dcd \rangle \subset kQ_F/\langle ba \rangle_{kQ_F}$. Then $A = kQ_F/\langle ba, ad, db, dcd \rangle$.

Let $B \subset A$ be an extension of k -algebras. From now on, we call A a B -algebra. The following definition is standard:

Definition 5.6 Let A be a B -algebra. A subset G of A is a B -generating set for A if every element of A is a sum of products of the form

$$b_n g_n \dots b_2 g_2 b_1 g_1 b_0 \quad (5.1)$$

where $g_i \in G$ and $b_i \in B$ for all i . The extension is called B -finitely generated if there exists a finite B -generating set for A .

Note that if G is a generating set for the B -bimodule A , then G is a B -generating set of the algebra A . If A/B is finitely generated as a B -bimodule (for instance if A is finite dimensional), then the extension is B -finitely generated.

Next we show that any extension of B which is finitely generated over B is an extension by arrows and relations.

Let B be a k -algebra and N a B -bimodule. The tensor algebra $T_B(N)$ has the following universal property. A morphism of k -algebras $\varphi : T_B(N) \rightarrow X$ is uniquely determined by a morphism of k -algebras $\varphi_B : B \rightarrow X$ (which endows X with a B -bimodule structure), and a morphism of B -bimodules $\varphi_N : N \rightarrow X$.

Let B be a k -algebra. Let BhB the free rank one B -bimodule $B \otimes B$ whose generator $1 \otimes 1$ is denoted h . Let H be a set, the free B -bimodule with basis H is denoted BHB .

The proof of the following result is clear.

Proposition 5.7 Let A be a B -algebra, G a subset of A and G' a copy of G . Consider the morphism of algebras $\varphi : T_B(BG'B) \rightarrow A$ determined by $\varphi_B : B \hookrightarrow A$ and by the B -bimodule map $\varphi_{BG'B} : BG'B \rightarrow A$ given by $g' \mapsto g$ for all $g \in G$. We have that G is a B -generating set of A if and only if φ is surjective.

Let $B = kQ/I$ where Q is a quiver and I an admissible ideal. Let A be a B -algebra. The vertices Q_0 are a complete set of orthogonal idempotents A . The corresponding Peirce decomposition of A is $A = \bigoplus_{x,y \in Q_0} yAx$. A subset $F \subset A$ is called a *Peirce subset* if $F = \bigcup_{x,y \in Q_0} F \cap yAx$.

Theorem 5.8 Let $B = kQ/I$ where Q is a quiver and I is an admissible ideal. Every finitely generated B -algebra A is an extension by arrows and relations.

Proof. Let G be a finite B -generating set of the extension $B \subset A$. We first observe that G provides a Peirce set $F = \{ygx \neq 0 \mid g \in G \text{ and } x, y \in Q_0\}$ which B -generates A . Note that $1 \otimes 1 = \sum_{x,y \in Q_0} y \otimes x$, therefore

$$B \otimes B = \bigoplus_{x,y \in Q_0} By \otimes xB.$$

Recall that G' is a set in one-to-one correspondence with G . Hence for $g \in G$, we have

$$Bg'B = \bigoplus_{x,y \in Q_0} Byg'xB$$

where $Byg'xB$ is the B -direct summand of $Bg'B$ which corresponds to $By \otimes xB$ in $B \otimes B$.

Let F' be a copy of F . If $a \in F$ belongs to yAx , we set $t(a') = y$ and $s(a') = x$. This way F' is a set of new arrows for Q . Let

$$N = \bigoplus_{a' \in F'} Bt(a')a's(a')B = \bigoplus_{a' \in F'} Bt(a') \otimes s(a')B.$$

Clearly N is a direct summand of the B -bimodule $BG'B$ with complement

$$\bigoplus_{\substack{g \in G \\ ygx=0}} Byg'xB.$$

Let $\varphi : T_B(BG'B) \rightarrow A$ be the surjective algebra morphism given by Proposition 5.7. Note that if $ygx = 0$ then by construction of φ we have

$$\varphi(yg'x) = y\varphi(g')x = ygx = 0.$$

Hence $Byg'xB \subset \text{Ker}\varphi$. Therefore $\psi = \varphi|_{T_B(N)}$ has the same image than φ , then ψ is surjective.

For $J = \text{Ker}\psi$ we infer an isomorphism of algebras $T_B(N)/J \xrightarrow{\bar{\psi}} A$. The composition $B \rightarrow T_B(N) \xrightarrow{\psi} A$ is the inclusion $B \subset A$, hence $J \cap B = 0$.

Recall that $B = kQ/I$, and that $Q_{F'}$ is the union of the quiver Q with the new arrows F' . As observed before, there is a canonical isomorphism of algebras

$$T_B(N) = \frac{kQ_{F'}}{\langle I \rangle_{kQ_{F'}}}.$$

We still denote by J the ideal in $\frac{kQ_{F'}}{\langle I \rangle_{kQ_{F'}}}$ corresponding to $\text{Ker}\psi$. We have proven that the algebras $\frac{kQ_{F'}}{\langle I \rangle_{kQ_{F'}}}/J$ and A are isomorphic. Moreover the isomorphism is the identity on B . \diamond

Let $B \subset A$ be an extension by arrows and relations. In the sequel we focus on giving a sufficient condition for A/B to be B -tensor nilpotent. First we introduce the following definitions.

Definition 5.9 Let $B \subset A$ be an extension by arrows and relations. A relative path of F -length $n \geq 0$ is a concatenated sequence

$$\beta_n a_n \dots \beta_2 a_2 \beta_1 a_1 \beta_0$$

where the β_i 's are paths of Q not in I (that is non-zero paths of B) and the a_i 's are in F .

Remark 5.10

- A relative path of F -length 0 is a path β of Q which is non-zero in B .
- The set of relative paths span T over k .
- The definition of a relative path only takes into account kQ/I and F .

Definition 5.11 A relative cycle of F -length $n \geq 1$ is a concatenated sequence

$$\beta_n a_n \dots \beta_2 a_2 \beta_1 a_1$$

where the β_i 's are paths of Q not in I , the a_i 's are in F , and $t(\beta_n) = s(a_1)$.

Remark 5.12 By abuse of language, we also call a relative cycle of F -length $n \geq 1$ a concatenated sequence

$$a_1 \beta_n a_n \dots \beta_2 a_2 \beta_1 a_1$$

where the β_i 's are paths of Q not in I and the a_i 's are in F .

In Example 5.3 (resp. 5.5), the sequence $a(bcd)a$ (resp. dcd) is a relative cycle of length 1.

In what follows, for $\gamma \in T$, we denote $\bar{\gamma}$ its class in $A = T/J$.

Definition 5.13 Let $a_1 \beta_n a_n \dots \beta_2 a_2 \beta_1 a_1$ be a relative cycle. A J -interrupter of the relative cycle is an arrow a_i such that in case $2 \leq i \leq n$ we have

$$\overline{\beta_i a_i \beta_{i-1}} \in B,$$

while for $i = 1$ we have

$$\overline{\beta_1 a_1 \beta_n} \in B.$$

In Example 5.3 the arrow a is a J -interrupter of the relative cycle $a(bcd)a$. Indeed

$$\overline{(bcd)a(bcd)} = \overline{bcd\alpha\beta} \in B.$$

On the other hand, in Example 5.5 the arrow d is not a J -interrupter of the relative cycle dcd since $\overline{cdc} \notin B$. Moreover A/B is not tensor nilpotent over B . Indeed $cd \otimes cd = cdc \otimes d$ represents a non zero element of $(A/B)^{\otimes_B 2}$, hence $cd \otimes cd \otimes \dots \otimes cd$ represents a non zero element of $(A/B)^{\otimes_B n}$ for all n .

Lemma 5.14 Let $B \subset A$ be an extension by arrows and relations. If there is no relative cycles then A/B is B -tensor nilpotent.

Before providing the proof, we underline that an oriented cycle in Q_F is not necessarily a relative cycle, as in the following example.

Example 5.15 Let $B = kQ/I$ where Q is the full arrows quiver

$$\begin{array}{ccc} 2 & \xrightarrow{b} & 3 \\ \uparrow a & & \downarrow c \\ 1 & \xleftarrow{d} & 4 \end{array}$$

and $I = \langle dcb \rangle$. Let $F = \{a\}$ be the dashed new arrow from 1 to 2. The quiver Q_F has oriented cycles but there are no relative cycles.

Proof of Lemma 5.14. Let $B = kQ/I$, where Q is a quiver and I is an admissible ideal.

We observe first that A/B is spanned over k by $\{\overline{\gamma}\}$, that is by the classes modulo J and B of relative paths γ of positive F -length.

Hence the set of m -tensors over B

$$\Gamma = \overline{\gamma_m} \otimes_B \cdots \otimes_B \overline{\gamma_1}$$

spans $(A/B)^{\otimes_B m}$ over k , where $\gamma_m, \dots, \gamma_1$ are relative paths of positive F -length. Note that since $kQ_0 \subset B$, if the relative paths are not concatenated then $\Gamma = 0$.

Recall that I is admissible, this implies that the paths β not in I are of bounded length. Hence there is a finite number n of relative paths of the form βa .

Let $\Gamma = \overline{\gamma_{n+1}} \otimes_B \cdots \otimes_B \overline{\gamma_1}$ be a k -generator of $(A/B)^{\otimes_B n+1}$, where $\gamma_{n+1}, \dots, \gamma_1$ are concatenated relative paths of positive F -length.

We consider the concatenated sequence $\gamma_{n+1} \dots \gamma_1$ and we reduce it as follows. For each occurrence

$$\cdots a'' \beta'' \otimes_B \beta' a' \cdots$$

we make the product $\beta'' \beta'$ in the sequence.

Suppose that some path $\beta'' \beta'$ belongs to I . Since

$$\cdots a'' \beta'' \otimes_B \beta' a' \cdots = \cdots a'' \beta'' \beta' \otimes_B a' \cdots \quad (5.2)$$

we infer that the sequence containing $\beta'' \beta'$ is in $\langle I \rangle_{kQ_F}$. Then the corresponding tensorand is zero, and $\Gamma = 0$.

On the contrary, in case none of the $\beta'' \beta'$ is zero in B , we infer a relative path of F -length at least $n + 1$. Then there is a repetition of some relative path of the form βa and we obtain a relative cycle

$$a \beta_m a_m \dots \beta_2 a_2 \beta a$$

for some $m \leq n$. Therefore, this case does not occur. \diamond

Theorem 5.16 *Let $B \subset A$ be an extension by arrows and relations, where $B = kQ/I$ and $A = \frac{kQ_F}{\langle I \rangle_{kQ_F}}/J$. If each relative cycle has a J -interrupter then A/B is B -tensor nilpotent.*

Proof. If there are no relative cycles then Lemma 5.14 shows that A/B is B -tensor nilpotent.

Next we assume that each relative cycle contains a J -interrupter. As before, let $\Gamma = \overline{\gamma_{n+1}} \otimes_B \cdots \otimes_B \overline{\gamma_1}$ be a k -generator of $(A/B)^{\otimes_B n+1}$, where $\gamma_{n+1}, \dots, \gamma_1$ are concatenated relative paths of positive F -length. By reducing the concatenated sequence $\gamma_{n+1} \dots \gamma_1$ as before we can assume that it is a relative path. Indeed, otherwise there is a product $\beta'' \beta'$ arising from some $\cdots a'' \beta'' \otimes_B \beta' a' \cdots$ which is in I , and by (5.2) we have $\Gamma = 0$.

Since n is the number of relative paths of the form βa , there is a repetition of some relative path βa , that is we have a relative cycle

$$a \beta_m a_m \dots \beta_2 a_2 \beta a. \quad (5.3)$$

Let c be a J -interrupter of it, with $\overline{\beta'c\beta} \in B$. If in the expression of Γ we have that β' and β belong to different tensorands, for instance $\cdots\beta'c \otimes_B \beta \cdots$, using the equality

$$\cdots\beta'c \otimes_B \beta \cdots = \cdots\beta'c\beta \otimes_B \cdots$$

we obtain an expression of Γ which contains, in one of its tensorands, the relative path $\beta'c\beta$. Modulo J , we replace $\beta'c\beta$ by a linear combination of paths of Q which are non-zero in B . Hence Γ is a linear combination of n -tensors, each of the summands having one less new arrow. If a summand has a tensorand which is in B (i.e. it has no more new arrows), then this summand is zero. For the others, we restart the process. This way we end with $\Gamma = 0$. \diamond

Next we will give conditions ensuring that A/B is projective as a left B -module. Of course reversing the arguments in the following proof gives conditions ensuring that A/B is right B -projective.

Lemma 5.17 *Let C be a k -algebra and let X and Y be respectively a left and a right C -module. Let Y' be a right C -subbimodule of Y . The C -bimodules $X \otimes \frac{Y}{Y'}$ and $\frac{X \otimes Y}{X \otimes Y'}$ are isomorphic.*

Proof. Consider the exact sequence $0 \rightarrow Y' \rightarrow Y \rightarrow \frac{Y}{Y'} \rightarrow 0$ and apply the exact functor $X \otimes -$ \diamond

Lemma 5.18 *Let $B \subset A$ be an extension by arrows and relations, where A is finite dimensional over k . There exists $n > 0$ such that any relative path of F -length $m \geq n$ is J -equivalent to a linear combination of paths of F -length smaller than n .*

Proof. The classes of relative paths k -span A , and we extract a finite basis from it. Then $n - 1$ is the maximal F -length of the relative paths in the basis. \diamond

Definition 5.19 *For a finite dimensional extension by arrows and relations $B \subset A$, the length index of the extension is the smallest n as above.*

Recall that for an extension by arrows and relations $B \subset A$, the algebra T is the tensor algebra

$$T_B(N) = B \oplus \bigoplus_{i>0} N^{\otimes_B i}$$

where $N = \bigoplus_{a \in F} Bt(a) \otimes s(a)B$. Let $T^{>0} = \bigoplus_{i>0} N^{\otimes_B i}$ be its positive ideal, and let $T^{]0,n[}$ be the B -bimodule $\bigoplus_{0 < i < n} N^{\otimes_B i}$.

Theorem 5.20 *Let $B \subset A$ be an extension by arrows and relations of length index n . Suppose that the projection $J_{]0,n[}$ of J on $T^{]0,n[}$ is generated as a B -bimodule by relations of the form $r = a \sum \lambda_\gamma \gamma$ where $a \in F$, the γ 's are relative paths concatenated with a and $\lambda_\gamma \in k$. Then A/B is left B -projective.*

Before proving the theorem, we observe that the hypothesis is verified in Example 5.3. Indeed, the ideal J is generated by $abcd - \alpha\beta$. Note that $abcd$ is a relative path of F -length two, and $\alpha\beta$ is a relative path of F -length one. In A , we have

that $abcd a = \alpha\beta a$. The relative paths of F -length ≥ 3 are zero in A . Therefore the length index of the extension is 2. Moreover $J_{]0,2[}$ is generated by $abcd$ as a B -bimodule.

Proof. Let $J_{>0}$ be the projection of J on $T^{>0}$. Since $B \cap J = 0$, as B -bimodules we have that

$$\frac{A}{B} = \frac{T^{>0}}{J_{>0}}.$$

Moreover, since the length index is n

$$\frac{T^{>0}}{J_{>0}} = \frac{T^{]0,n[}}{J_{]0,n[}}.$$

The relative path $a\beta$ corresponds to $t(a) \otimes s(a)\beta = t(a) \otimes \beta \in T$. The B -bimodule in T generated by $a\beta$ is k -spanned by $\beta''t(a) \otimes \beta\beta'$ where β'' and β' are paths of Q which are non-zero in B .

We observe that for $i \geq 1$ we have

$$N^{\otimes_B i} = \bigoplus_{a \in F} Bt(a) \otimes s(a)B \otimes_B N^{\otimes_B i-1} = \bigoplus_{a \in F} Bt(a) \otimes s(a)N^{\otimes_B i-1}.$$

Therefore

$$T^{]0,n[} = \bigoplus_{a \in F} Bt(a) \otimes s(a)T^{<n-1}.$$

For an arrow a , we consider the r 's of type a , that is of the form $r = a \sum \lambda_\gamma \gamma$. Let $r' = \sum \lambda_\gamma \gamma \in T$ and let Y_a be the right submodule of $s(a)T^{<n-1}$ generated by them. Clearly

$$J_{>0} = \bigoplus_{a \in F} Bt(a) \otimes Y_a.$$

This way

$$\frac{A}{B} = \bigoplus_{a \in F} \frac{Bt(a) \otimes s(a)T^{<n-1}}{Bt(a) \otimes Y_a}.$$

By Lemma 5.17, we obtain finally

$$\frac{A}{B} = \bigoplus_{a \in F} Bt(a) \otimes \frac{s(a)T^{<n-1}}{Y_a}.$$


Note that $Bt(a) \otimes \frac{s(a)T^{<n-1}}{Y_a}$ is isomorphic as left B -module to the direct sum of $\dim_k \left(\frac{s(a)T^{<n-1}}{Y_a} \right)$ copies of $Bt(a)$. Moreover $Bt(a)$ is the left projective B -module corresponding to the idempotent $t(a)$. Hence A/B is a left projective B -module. \diamond

6 Examples

We will use the tools that we have developed in order to consider examples of extensions by arrows and relations. We will study if the extension is left or right bounded and if it is split or not.

We recall that an extension $B \subset A$ is split if there is an ideal M of A such that $A = B \oplus M$. This is of course equivalent to the fact that the algebra inclusion $B \hookrightarrow A$ splits.

Example 6.1 We recall the algebra extension provided in Example 5.3, where $B = kQ/I$ with Q the full arrows quiver



and $I = \langle \beta\alpha \rangle$. Let $F = \{a\}$ be the dashed new arrow from 2 to 1, and let $J = \langle abcd - \alpha\beta \rangle$. Then $A = kQ_F/\langle \beta\alpha, abcd - \alpha\beta \rangle$.

As mentioned, a is a J -interrupter of the relative cycle $abcd$, and the same holds for longer relative cycles such as $abcdabcd$. So A/B is tensor nilpotent by Theorem 5.16. Let e_1 be the idempotent corresponding to the vertex $1 = t(a)$. The left B -module A/B is projective by Theorem 5.20. More precisely, as noticed before, the length index of the extension is 2. A basis of

$$\frac{s(a)T^{<1}}{Y_a} = \frac{e_2B}{bcdB}$$

is $\{e_2, b, bc\}$. Hence by Theorem 5.20 we have that $A/B \simeq Be_1 \oplus Be_1 \oplus Be_1$ as left B -modules.

The algebra B has finite global dimension, then B^e also have it and the B -bimodules are of finite projective dimension, in particular A/B is of finite projective dimension as a B -bimodule. Hence this extension by arrows and relations is left bounded.

We claim it is not split. In fact we will prove that there is not even a right B -submodule of A such that $A = B \oplus X$. Let X be a right B -submodule of A . Next we show that $B \cap X \neq 0$.

Let \mathcal{B} be a basis of paths for B with $Q_0 \subset \mathcal{B}$. The vector space A/B is k -spanned by relative paths of F -positive length, including a . Hence we can complete \mathcal{B} to a basis \mathcal{A} of A , with relative paths of positive F -length, including a . Let D be the k -vector space with basis $\mathcal{A} \setminus \{a\}$. We claim that there exists $x \in X$ such that in the basis \mathcal{A} , the coefficient x_a of a is not 0. Indeed, otherwise $X \subset D$ and since $B \subset D$ we would have $B \oplus X \subset D$. This cannot hold since $D \subsetneq A$. Then $x = x_a a + y$ with $y \in D$ and $x_a \neq 0$.

Since X is a right B -module, we have that $xbcd = x_a abcd + ybcd \in X$. We assert that $ybcd = \lambda bcd$ for $\lambda \in k$. Indeed, y is a k -linear combination of the basis $\mathcal{A} \setminus \{a\}$ of D . The only possible path of $\mathcal{A} \setminus \{a\}$ which concatenates with bcd on its left is e_2 , and this proves the assertion.

Then $xbcd = x_a abcd + \lambda bcd = x_a \beta\alpha + \lambda bcd$. Note that $\beta\alpha$ and bcd are linearly independent in B . Hence we obtain $0 \neq xbcd \in X \cap B$.

Example 6.2 Let Q be the full arrows quiver

$$\begin{array}{ccccc}
 & & 2 & \xrightarrow{d} & 3 \\
 & & \uparrow a & & \uparrow c \\
 5 & \xrightarrow{\mu} & 1 & \xrightarrow{b} & 4
 \end{array}$$

and let $B = kQ$. Let a be the dashed new arrow from 1 to 2, let $J = \langle da - cb \rangle$ and let A be the corresponding extension by arrows and relations.

We assert that the extension is right bounded and not split. There are no relative cycles, so A/B is tensor nilpotent. Moreover the conditions are satisfied to ensure that A/B is a right projective B -module by the dual version of Theorem 5.20. The algebra B is hereditary, hence B^e has finite global dimension and A/B is of finite projective dimension as a B -bimodule. As in the previous example the extension $B \subset A$ is not split.

References

- [1] Assem, I.; Brüstle, T.; Schiffler, R. Cluster-tilted algebras as trivial extensions. *Bull. Lond. Math. Soc.* 40 (2008), 151–162.
- [2] Assem, I.; Gatica, M. A.; Schiffler, R.; Taillefer, R. Hochschild cohomology of relation extension algebras. *J. Pure and Appl. Algebra* 220 (2016), 2471–2499.
- [3] André, M. *Homologie des algèbres commutatives*. Die Grundlehren der mathematischen Wissenschaften, 206. Springer-Verlag, Berlin-New York, 1974.
- [4] Auslander, M. On the dimension of modules and algebras. III. Global dimension. *Nagoya Math J.* 9 (1955), 67–77.
- [5] Avramov, L. L.; Vigué-Poirrier M. Hochschild homology criteria for smoothness. *Internat. Math. Res. Notices* 1 (1992), 17–25.
- [6] Bergh, P. A.; Erdmann, K. Homology and cohomology of quantum complete intersections. *Algebra Number Theory* 2 (2008), 501–522.
- [7] Bergh, P. A.; Madsen, D. Hochschild homology and global dimension. *Bull. Lond. Math. Soc.* 41 (2009), 473–482.
- [8] Bergh, P. A.; Madsen, D. Hochschild homology and trivial extension algebras. *Proc. Amer. Math. Soc.* 145 (2017), 1475–1480.
- [9] Buenos Aires Cyclic Homology Group. Cyclic homology of algebras with one generator. (J.A. Guccione, J. J. Guccione, M. J. Redondo, A. Solotar and O. Villamayor participated in this research) *K-Theory* 5 (1991), 51–69.
- [10] Cibils, C.; Lanzilotta, M.; Marcos, E.N.; Solotar, A. Split bounded extension algebras and Han’s conjecture. *Pacific J. Math.* 307 (2020), 63–77. <https://doi.org/10.2140/pjm.2020.307.63>

- [11] Cibils, C.; Redondo, M. J.; Solotar, A. Han's conjecture and Hochschild homology for null-square projective algebras. *Indiana Univ. Math. J.* 70 (2021), 639–668
<https://doi.org/10.1512/iumj.2021.70.8402>
- [12] Cibils, C.; Lanzilotta, M.; Marcos, E.N.; Solotar, A. Jacobi-Zariski long nearly exact sequences for associative algebras. *Bull. London Math. Soc.*
<https://doi.org/10.1112/blms.12516>
- [13] Han, Y. Hochschild (co)homology dimension. *J. London Math. Soc.* 73 (2006), 657–668.
- [14] Hochschild, G. On the cohomology groups of an associative algebra, *Ann. Math.* 46 (1945), 58–67.
- [15] Hochschild, G. Relative homological algebra, *Trans. Amer. Math. Soc.* 82 (1956), 246–269.
- [16] Iyengar, S. André-Quillen homology of commutative algebras. *Interactions between homotopy theory and algebra*, 203–234. *Contemporary Mathematics*, 436. American Mathematical Society, Providence, RI, 2007.
- [17] Kaygun, A. Jacobi-Zariski exact sequence for Hochschild homology and cyclic (co)homology, *Homology Homotopy Appl.* 14 (2012), 65–78.
- [18] Kaygun, A. Erratum to “Jacobi-Zariski exact sequence for Hochschild homology and cyclic (co)homology” *Homology Homotopy Appl.* 21 (2019), 301–303.
- [19] Loday, J.-L. *Cyclic homology*. Appendix E by María O. Ronco. Second edition. Chapter 13 by the author in collaboration with Teimuraz Pirashvili. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, 301. Springer-Verlag, Berlin, 1998.
- [20] Solotar, A.; Suárez-Alvarez, M.; Vivas, Q. Hochschild homology and cohomology of generalized Weyl algebras: the quantum case. *Ann. Inst. Fourier (Grenoble)* 63 (2013), 923–956.
- [21] Solotar, A.; Vigué-Poirrier M. Two classes of algebras with infinite Hochschild homology, *Proc. Amer. Math. Soc.* 138 (2010), 861–869.
- [22] Weibel, C. A. *An introduction to homological algebra*. *Cambridge Studies in Advanced Mathematics*, 38. Cambridge University Press, Cambridge, 1994.
- [23] Witherspoon, S. *Hochschild cohomology for algebras*. *Graduate Studies in Mathematics*, 204. American Mathematical Society, Providence, RI, 2019.

C.C.:

Institut Montpellierain Alexander Grothendieck, CNRS, Univ. Montpellier, France.
Claude.Cibils@umontpellier.fr

M.L.:

Instituto de Matemática y Estadística “Rafael Laguardia”, Facultad de Ingeniería, Universidad de la República, Uruguay.
marclan@fing.edu.uy

E.N.M.:

Departamento de Matemática, IME-USP, Universidade de São Paulo, Brazil.
enmarcos@ime.usp.br

A.S.:
IMAS-CONICET y Departamento de Matemática, Facultad de Ciencias Exactas y Naturales,
Universidad de Buenos Aires, Argentina.
asolotar@dm.uba.ar