

# Quotient maps of 2-uniform tilings of the plane on the torus

DIPENDU MAITY<sup>1</sup>, DEBASHIS BHOWMIK<sup>2</sup>, MARBARISHA M. KHARKONGOR<sup>1</sup>

<sup>1</sup>*Department of Sciences and Mathematics, Indian Institute of Information Technology Guwahati, Bongora, Assam-781 015, India.*

*{dipendu, marbarisha.kharkongor}@iiitg.ac.in/{dipendumaity, marbarisha.kharkongor}@gmail.com.*

<sup>2</sup>*Department of Mathematics, Indian Institute of Technology Patna, Patna 801 106, India. debashisiitg@gmail.com.*

May 5, 2021

## Abstract

A vertex-transitive map  $X$  is a map on a closed surface on which the automorphism group  $\text{Aut}(X)$  acts transitively on the set of vertices. If the face-cycles at all the vertices in a map are of two types, then the map is said to be a 2-semiequivelar map. A 2-uniform tiling is an edge-to-edge tiling of regular polygons having 2 distinct transitivity classes of vertices. Clearly, a 2-uniform map is 2-semiequivelar. The converse of this is not true in general. We know that there are infinitely many tilings of 2-semiequivelar type. In this article, we study 2-semiequivelar maps on the torus that are the quotient of the plane's 2-uniform lattices. We have found bounds of the number of orbits of vertices of the quotient 2-semiequivelar maps on the torus under the automorphism groups (symmetric groups).

*MSC 2010:* 52C20, 52B70, 51M20, 57M60.

*Keywords:* Polyhedral map on torus; 2-uniform maps; 2-semiequivelar maps; Symmetric group.

## 1 Introduction

A map is a connected 2-dimensional cell complex on a surface. Equivalently, it is a cellular embedding of a connected graph on a surface. In this article, a map will mean a polyhedral map on a surface, that is, non-empty intersection of any two faces is either a vertex or an edge. For a map  $\mathcal{K}$ , let  $V(\mathcal{K})$  be the vertex set of  $\mathcal{K}$  and  $u \in V(\mathcal{K})$ . The faces containing  $u$  form a cycle (called the *face-cycle* at  $u$ )  $C_u$  in the dual graph of  $\mathcal{K}$ . That is,  $C_u$  is of the form  $(F_{1,1}-\cdots-F_{1,n_1})-\cdots-(F_{k,1}-\cdots-F_{k,n_k})-F_{1,1}$ , where  $F_{i,\ell}$  is a  $p_i$ -gon for  $1 \leq \ell \leq n_i$ ,  $1 \leq i \leq k$ ,  $p_r \neq p_{r+1}$  for  $1 \leq r \leq k-1$  and  $p_k \neq p_1$ . Then, the vertex  $u$  is said to be of type  $[p_1^{n_1}, \dots, p_k^{n_k}]$  (addition in the suffix is modulo  $k$ ). A map  $\mathcal{K}$  is said to be *2-semiequivelar* of type  $[W; Z]$  if (1)  $u$  and  $v$  are of same type  $W$ , or (2)  $u$  is of type  $W$  and  $v$  is of type  $Z$  for all  $u, v \in V(\mathcal{K})$ . If  $W = Z$  then the map is called *semiequivelar* of type  $[W]$ . A *2-uniform tiling* is an edge-to-edge tiling of regular polygons having 2 distinct transitivity classes of vertices. A 2-uniform tiling will have vertices that we could label  $X$ , and others that we could label  $Y$ . Each  $X$  vertex can be mapped onto every other  $X$  vertex, but cannot be mapped to any  $Y$  vertex. Clearly, an 2-uniform map is 2-semiequivelar. Let  $F_1-\cdots-F_m-F_1$  be the face-cycle of a vertex  $u$  in a map  $\mathcal{K}$ . Then  $F_i \cap F_j$  is either  $u$  or an edge through  $u$ . Thus the face  $F_i$  must be of the form  $u_{i+1}-u-u_i-P_i-u_{i+1}$ , where  $P_i = \emptyset$  or a path &  $P_i \cap P_j = \emptyset$  for  $i \neq j$ . Here addition in the suffix is modulo  $m$ . So,  $u_1-P_1-u_2-\cdots-u_m-P_m-u_1$  is a cycle and said to be the *link-cycle* or *link* of  $u$ , denoted by  $lk(u)$  or  $lk_{\mathcal{K}}(u)$ . A face in a map of the form  $u_1-u_2-\cdots-u_n-u_1$  is also denoted by  $[u_1, \dots, u_n]$ .

A *semi-regular* tiling of the plane  $\mathbb{R}^2$  is a tiling of  $\mathbb{R}^2$  by regular polygons such that all the vertices of the tiling are of the same type. A *semi-regular* tiling of  $\mathbb{R}^2$  is also known as *Archimedean*, or *homogeneous*, or *uniform* tiling. In [11], Grünbaum and Shephard showed that there are exactly eleven types of Archimedean tilings on the plane. These types are  $[3^6]$ ,  $[3^4, 6^1]$ ,  $[3^3, 4^2]$ ,  $[3^2, 4^1, 3^1, 4^1]$ ,  $[3^1, 6^1, 3^1, 6^1]$ ,  $[3^1, 4^1, 6^1, 4^1]$ ,  $[3^1, 12^2]$ ,  $[4^4]$ ,  $[4^1, 6^1, 12^1]$ ,  $[4^1, 8^2]$ ,  $[6^3]$ . Clearly, a *semi-regular* tiling on  $\mathbb{R}^2$  gives a semiequivelar map on  $\mathbb{R}^2$ . But, there are semiequivelar maps on the plane which are not (not isomorphic to) an Archimedean tiling. In fact, there exists  $[p^q]$  equivelar maps on  $\mathbb{R}^2$  whenever  $1/p + 1/q < 1/2$  (e.g.,  $[3]$ ,  $[10]$ ). Thus, we have

**Proposition 1.1.** *There are infinitely many types of equivelar maps on the plane  $\mathbb{R}^2$ .*

We know from [7] and [8] that infinitely many equivelar maps exist on both the torus and the Klein bottle. Thus, infinitely many semiequivelar maps exist on both the torus and the Klein bottle. Only eleven types of semiequivelar maps on the torus and ten types of semiequivelar maps on the Klein bottle are known in the literature. All these are quotients of Archimedean tilings of the plane [1, 16, 17]. Since there are infinitely many equivelar maps on the plane, it is natural to ask whether there are more types of semiequivelar maps on the torus or the Klein bottle. Here we know

**Proposition 1.2.** *[5, 6] Let  $X$  be a semiequivelar map on a surface  $M$ . (a) If  $M$  is the torus then the type of  $X$  is  $[3^6]$ ,  $[6^3]$ ,  $[4^4]$ ,  $[3^4, 6^1]$ ,  $[3^3, 4^2]$ ,  $[3^2, 4^1, 3^1, 4^1]$ ,  $[3^1, 6^1, 3^1, 6^1]$ ,  $[3^1, 4^1, 6^1, 4^1]$ ,  $[3^1, 12^2]$ ,  $[4^1, 8^2]$  or  $[4^1, 6^1, 12^1]$ . (b) If  $M$  is the Klein bottle then the type of  $X$  is  $[3^6]$ ,  $[6^3]$ ,  $[4^4]$ ,  $[3^3, 4^2]$ ,  $[3^2, 4^1, 3^1, 4^1]$ ,  $[3^1, 6^1, 3^1, 6^1]$ ,  $[3^1, 4^1, 6^1, 4^1]$ ,  $[3^1, 12^2]$ ,  $[4^1, 8^2]$  or  $[4^1, 6^1, 12^1]$ .*

Proposition 1.2 and the known examples imply that there are precisely eleven types of semiequivelar maps on the torus and ten types of semiequivelar maps on the Klein bottle.

In [2], Brehm and Kühnel presented a formula to determine the number of distinct vertex-transitive equivelar maps of types  $[3^6]$  and  $[4^4]$  on the torus. It was shown in [8] that every equivelar map of type  $[3^6]$  on the torus is vertex-transitive. By similar arguments, one can easily show that an equivelar map of type  $[4^4]$  on the torus is vertex-transitive (also see [2, Proposition 6]). Thus, we have

**Proposition 1.3.** *Let  $X$  be an equivelar map on the torus. If the type of  $X$  is  $[3^6]$ ,  $[4^4]$ ,  $[6^3]$  or  $[3^3, 4^2]$  then  $X$  is vertex-transitive.*

There are 14 types  $[3^6; 3^4, 6^1]$ ,  $[3^6; 3^3; 4^2]$ ,  $[3^6; 3^2, 4^1, 3^1, 4^1]$ ,  $[3^6; 3^2, 4^1, 12^1]$ ,  $[3^4, 6^1; 3^2, 6^2]$ ,  $[3^3, 4^2; 3^2, 4^1, 3^1, 4^1]$ ,  $[3^3, 4^2; 3^1, 4^1, 6^1, 4^1]$ ,  $[3^3, 4^2; 4^4]$ ,  $[3^2, 4^1, 3^1, 4^1; 3^1, 4^1, 6^1, 4^1]$ ,  $[3^2, 6^2; 3^1, 6^1, 3^1, 6^1]$ ,  $[3^1, 4^1, 3^1, 12^1; 3^1, 12^2]$ ,  $[3^1, 4^2, 6^1; 3^1, 4^1, 6^1, 4^1]$ ,  $[3^1, 4^2, 6^1; 3^1, 6^1, 3^1, 6^1]$ ,  $[3^1, 4^1, 6^1, 4^1; 4^1, 6^1, 12^1]$  of 20 2-uniform tilings on the plane (see in Section 2). Since plane is the universal cover of torus, so, these types of maps also exist on the torus. It is natural to ask that the number of orbits of vertices under the automorphism group of the 2-semiequivelar maps on the torus. Here, we give the bound of the number of orbits of vertices under the automorphism group of the maps of the 2-semiequivelar maps on the torus that are the quotient of the plane's 2-uniform lattice. In particular,

**Theorem 1.4.** (1) *Let  $X_1$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_1[3^6; 3^4, 6^1]$ . Let the vertices of  $X_1$  form  $m_1$   $\text{Aut}(X_1)$ -orbits. Then,  $m_1 \leq 6$ .*

(2) *Let  $X_2$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_2[3^6; 3^4, 6^1]$ . Let the vertices of  $X_2$  form  $m_2$   $\text{Aut}(X_2)$ -orbits. Then,  $m_2 \leq 4$ .*

(3) Let  $X_3$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_3[3^6; 3^3, 4^2]$ . Let the vertices of  $X_3$  form  $m_3$   $\text{Aut}(X_3)$ -orbits. Then,  $m_3 = 2$ .

(4) Let  $X_4$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_4[3^6; 3^3, 4^2]$ . Let the vertices of  $X_4$  form  $m_4$   $\text{Aut}(X_4)$ -orbits. Then,  $m_4 = 2$ .

(5) Let  $X_5$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_5[3^6; 3^2, 4^1, 3^1, 4^1]$ . Let the vertices of  $X_5$  form  $m_5$   $\text{Aut}(X_5)$ -orbits. Then,  $m_5 \leq 4$ .

(6) Let  $X_6$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_6[3^6; 3^2, 4^1, 12^1]$ . Let the vertices of  $X_6$  form  $m_6$   $\text{Aut}(X_6)$ -orbits. Then,  $m_6 \leq 7$ .

(7) Let  $X_7$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_7[3^6; 3^2, 6^2]$ . Let the vertices of  $X_7$  form  $m_7$   $\text{Aut}(X_7)$ -orbits. Then,  $m_7 \leq 4$ .

(8) Let  $X_8$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_8[3^4, 6^1; 3^2, 6^2]$ . Let the vertices of  $X_8$  form  $m_8$   $\text{Aut}(X_8)$ -orbits. Then,  $m_8 \leq 3$ .

(9) Let  $X_9$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_9[3^3, 4^2; 3^2, 4^1, 3^1, 4^1]$ . Let the vertices of  $X_9$  form  $m_9$   $\text{Aut}(X_9)$ -orbits. Then,  $m_9 \leq 6$ .

(10) Let  $X_{10}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{10}[3^3, 4^2; 3^2, 4^1, 3^1, 4^1]$ . Let the vertices of  $X_{10}$  form  $m_{10}$   $\text{Aut}(X_{10})$ -orbits. Then,  $m_{10} \leq 3$ .

(11) Let  $X_{11}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{11}[3^3, 4^2; 3^1, 4^1, 6^1, 4^1]$ . Let the vertices of  $X_{11}$  form  $m_{11}$   $\text{Aut}(X_{11})$ -orbits. Then,  $m_{11} \leq 6$ .

(12) Let  $X_{12}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{12}[3^3, 4^2; 4^4]$ . Let the vertices of  $X_{12}$  form  $m_{12}$   $\text{Aut}(X_{12})$ -orbits. Then,  $m_{12} = 2$ .

(13) Let  $X_{13}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{13}[3^3, 4^2; 4^4]$ . Let the vertices of  $X_{13}$  form  $m_{13}$   $\text{Aut}(X_{13})$ -orbits. Then,  $m_{13} = 2$ .

(14) Let  $X_{14}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{14}[3^2, 4^1, 3^1, 4^1; 3^1, 4^1, 6^1, 4^1]$ . Let the vertices of  $X_{14}$  form  $m_{14}$   $\text{Aut}(X_{14})$ -orbits. Then,  $m_{14} \leq 6$ .

(15) Let  $X_{15}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{15}[3^2, 6^2; 3^1, 6^1, 3^1, 6^1]$ . Let the vertices of  $X_{15}$  form  $m_{15}$   $\text{Aut}(X_{15})$ -orbits. Then,  $m_{15} = 2$ .

(16) Let  $X_{16}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{16}[3^1, 4^1, 3^1, 12^1; 3^1, 12^2]$ . Let the vertices of  $X_{16}$  form  $m_{16}$   $\text{Aut}(X_{16})$ -orbits. Then,  $m_{16} \leq 3$ .







(17) Let  $X_{17}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{17}[3^1, 4^2, 6^1; 3^1, 4^1, 6^1, 4^1]$ . Let the vertices of  $X_{17}$  form  $m_{17}$   $\text{Aut}(X_{17})$ -orbits. Then,  $m_{17} \leq 9$ .


(18) Let  $X_{18}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{18}[3^1, 4^2, 6^1; 3^1, 6^1, 3^1, 6^1]$ . Let the vertices of  $X_{18}$  form  $m_{18}$   $\text{Aut}(X_{18})$ -orbits. Then,  $m_{18} \leq 3$ .

(19) Let  $X_{19}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{19}[3^1, 4^2, 6^1; 3^1, 6^1, 3^1, 6^1]$ . Let the vertices of  $X_{19}$  form  $m_{19}$   $\text{Aut}(X_{19})$ -orbits. Then,  $m_{19} \leq 3$ .


(20) Let  $X_{20}$  be a semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{20}[3^1, 4^1, 6^1, 4^1; 4^1, 6^1, 12^1]$ . Let the vertices of  $X_{20}$  form  $m_{20}$   $\text{Aut}(X_{20})$ -orbits. Then,  $m_{20} \leq 9$ .

## 2 2-uniform tilings of the plane







$K_7[3^6; 3^2, 6^2]$




$K_8[3^4, 6^1; 3^2, 6^2]$




$K_9[3^3, 4^2; 3^3, 4^1, 3^1, 4^1]$









$K_{10}[3^3, 4^2; 3^2, 4^1, 3^1, 4^1]$





$K_{11}[3^3, 4^2; 3^1, 4^1, 6^1, 4^1]$



$K_{12}[3^3, 4^2; 4^4]$





### 3 Proof of Theorems 1.4

Let  $K_i = K_i[A]$  ( $K_i[A]$  given in Section 2), where

$$\begin{aligned}
 A \in \{ & [3^6; 3^4, 6^1], [3^6; 3^3; 4^2], [3^6; 3^2, 4^1, 3^1, 4^1], [3^6; 3^2, 4^1, 12^1], \\
 & [3^4, 6^1; 3^2, 6^2], [3^3, 4^2; 3^2, 4^1, 3^1, 4^1], [3^3, 4^2; 3^1, 4^1, 6^1, 4^1], [3^3, 4^2; 4^4], \\
 & [3^2, 4^1, 3^1, 4^1; 3^1, 4^1, 6^1, 4^1], [3^2, 6^2; 3^1, 6^1, 3^1, 6^1], [3^1, 4^1, 3^1, 12^1; 3^1, 12^2], \\
 & [3^1, 4^2, 6^1; 3^1, 4^1, 6^1, 4^1], [3^1, 4^2, 6^1; 3^1, 6^1, 3^1, 6^1], [3^1, 4^1, 6^1, 4^1; 4^1, 6^1, 12^1] \}.
 \end{aligned}$$

Grünbaum and G. C. Shephard [11, 12] and Krötenheerdt [9] have discussed the existence and uniqueness of the 2-uniform tilings  $K_i$ ,  $i = 1, 2, \dots, 20$  of the plane. The proof of this fact may be carried out along lines analogous to those explained in connection with the 11 uniform tilings (see [5, 6] for the uniqueness of 11 uniform tilings of the plane). Thus, we have the following.

**Proposition 3.1.** *The maps  $K_i$  ( $1 \leq i \leq 20$ ) (in Section 2) are unique up to isomorphism.*

**Remark 3.2.** *Let  $S = \{0, 1, 2, \dots, m-1\}$ . Let  $S+n := \{0+n, 1+n, 2+n, \dots, m+n-1\}$ . Define  $S_m = \{S+mk : k \in \mathbb{Z} \text{ \& } 2 \mid (k-1)\}$ . Let  $K_m$  be a map on the plane defined as follows. Let the vertex set of  $K_m$  be  $\{(i, j) \in \mathbb{Z} \times \mathbb{Z}\}$  on  $\mathbb{R}^2$ . We say  $(i, j) \sim (i_1, j_1)$  if there is an edge between  $(i, j)$  and  $(i_1, j_1)$ . The edges of  $K_m$  :*

- (1)  $(i, j) \sim (i_1, j_1)$  if  $|i - i_1| = 1$  and  $j = j_1$ , or  $i = i_1$  and  $|j - j_1| = 1$ ,
- (2)  $(i, j) \sim (i+1, j+1)$ ,  $(t, j) \sim (t+1, j-1)$  if  $i \in S_m$ ,  $j$  is even and  $t \in \mathbb{Z} \setminus S_m$ .

*Thus, we get an object  $K_m$  for each  $m \geq 2$ . Clearly,  $K_x \not\cong K_y$  for  $x \neq y$  since the number of continuous quadrilaterals are different. Therefore, there are infinitely many 2-semiequivelar tilings of the plane.*

*Proof of Theorem 1.4.* Let  $X_1$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_1$ . Let the vertices of  $X_1$  form  $m_1$   $\text{Aut}(X_1)$ -orbits. Let  $K_1$  be as in Section 2. Let  $V_1 = V(K_1)$  be the vertex set of  $K_1$ . Let  $H_1$  be the group of all the translations of  $K_1$ . So,  $H_1 \leq \text{Aut}(K_1)$ .

Since  $X_1$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_1$  (as by Proposition 3.1  $K_1$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_1 : K_1 \rightarrow X_1$  where  $X_1 = K_1/\Gamma_1$  for some fixed element (vertex, edge or face)

free subgroup  $\Gamma_1 \leq \text{Aut}(K_1)$ . Hence  $\Gamma_1$  consists of translations and glide reflections. Since  $X_1 = K_1/\Gamma_1$  is orientable,  $\Gamma_1$  does not contain any glide reflection. Thus  $\Gamma_1 \leq H_1$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $a_0$  and  $a_{-2}$  of  $K_1$ . Let  $\alpha_1 := b_3 - a_0$ ,  $\beta_1 := e_0 - a_0$  and  $\gamma_1 := l_4 - a_0$  in  $K_1$ . Then

$$H_1 := \langle x \mapsto x + \alpha_1, x \mapsto x + \beta_1, x \mapsto x + \gamma_1 \rangle.$$

Under the action of  $H_1$ , vertices of  $K_1$  form twelve orbits. Consider the subgroup  $G_1$  of  $\text{Aut}(K_1)$  generated by  $H_1$  and the map (the half rotation)  $x \mapsto -x$ . So,

$$G_1 = \{ \alpha : x \mapsto \varepsilon x + m\alpha_1 + n\beta_1 + r\gamma_1 : \varepsilon = \pm 1, m, n, r \in \mathbb{Z} \} \cong H_1 \rtimes \mathbb{Z}_2.$$

Clearly, under the action of  $G_1$ , vertices of  $K_1$  form six orbits. The orbits are

$$O_1 := \langle b_{-1} \rangle, O_2 := \langle b_0 \rangle, O_3 := \langle b_1 \rangle, O_4 := \langle c_1 \rangle, O_5 := \langle d_2 \rangle, O_6 := \langle c_0 \rangle.$$

**Claim 1.** If  $K \leq H_1$  then  $K \trianglelefteq G_1$ .

Let  $g \in G_1$  and  $k \in K$ . Then  $g(x) = \varepsilon x + ma + nb + rc$  and  $k(x) = x + pa + qb + lc$  for some  $m, n, r, p, q, \ell \in \mathbb{Z}$  and  $\varepsilon \in \{1, -1\}$ . Therefore,

$$\begin{aligned} (g \circ k \circ g^{-1})(x) &= (g \circ k)(\varepsilon(x - ma - nb - rc)) \\ &= g(\varepsilon(x - ma - nb - rc) + pa + qb + lc) \\ &= x - ma - nb - rc + \varepsilon(pa + qb + lc) + ma + nb + rc \\ &= x + \varepsilon(pa + qb + lc) \\ &= k^\varepsilon(x). \end{aligned}$$

Thus,  $g \circ k \circ g^{-1} = k^\varepsilon \in K$ . This completes the claim.

By Claim 1,  $\Gamma_1$  is a normal subgroup of  $G_1$ . Therefore,  $G_1/\Gamma_1$  acts on  $X_1 = K_1/\Gamma_1$ . Since

$$O_1 := \langle b_{-1} \rangle, O_2 := \langle b_0 \rangle, O_3 := \langle b_1 \rangle, O_4 := \langle c_1 \rangle, O_5 := \langle d_2 \rangle, O_6 := \langle c_0 \rangle.$$

are the  $G_1$ -orbits, it follows that  $\eta_1(O_j)$  for  $j = 1, 2, \dots, 6$  are the  $(G_1/\Gamma_1)$ -orbits. Since the vertex set of  $K_1$  is  $\sqcup_{j=1}^6 \eta_j(O_j)$  and  $G_1/\Gamma_1 \leq \text{Aut}(X_1)$ , it follows that the number of  $\text{Aut}(X_1)$ -orbits of vertices is  $\leq 6$ . This completes the part (1).

Let  $X_2$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_2$ . Let the vertices of  $X_2$  form  $m_2$   $\text{Aut}(X_2)$ -orbits. Let  $K_2$  be as in Section 2. Let  $V_2 = V(K_2)$  be the vertex set of  $K_2$ . Let  $H_2$  be the group of all the translations of  $K_2$ . So,  $H_2 \leq \text{Aut}(K_2)$ .

Since  $X_2$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_2$  (as by Proposition 3.1  $K_2$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_2 : K_2 \rightarrow X_2$  where  $X_2 = K_2/\Gamma_2$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_2 \leq \text{Aut}(K_2)$ . Hence  $\Gamma_2$  consists of translations and glide reflections. Since  $X_2 = K_2/\Gamma_2$  is orientable,  $\Gamma_2$  does not contain any glide reflection. Thus  $\Gamma_2 \leq H_2$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $a_0$  and  $a_{-2}$  of  $K_2$  (see Section 2). Let  $\alpha_2 := a_3 - a_0$ ,  $\beta_2 := d_1 - a_0$  and  $\gamma_2 := d_{-2} - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_2 := \langle x \mapsto x + \alpha_2, x \mapsto x + \beta_2, x \mapsto x + \gamma_2 \rangle$  and

$$G_2 = \{ \alpha : x \mapsto \varepsilon x + m_1\alpha_2 + n_1\beta_2 + r_1\gamma_2 : \varepsilon = \pm 1, m_1, n_1, r_1 \in \mathbb{Z} \} \cong H_2 \rtimes \mathbb{Z}_2.$$



By the same arguments as above and in Claim 1,  $\Gamma_2 \trianglelefteq G_2$  and the number of  $G_2/\Gamma_2$ -orbits of vertices of  $X_2$  is four. Therefore,  $G_2/\Gamma_2$  acts on  $X_2 = K_2/\Gamma_2$ . Since

$$O_1 = \langle a_0 \rangle, O_2 = \langle b_0 \rangle, O_3 = \langle h_0 \rangle, O_4 = \langle h_{-1} \rangle$$

are the  $G_2$ -orbits, it follows that  $\eta_2(O_j)$  for  $j = 1, 2, 3, 4$  are the  $(G_2/\Gamma_2)$ -orbits. Since the vertex set of  $Y$  is  $\sqcup_{j=1}^4 \eta_j(O_j)$  and  $G_2/\Gamma_2 \leq \text{Aut}(X_2)$ , it follows that the number of  $\text{Aut}(X_2)$ -orbits of vertices is  $\leq 4$ . This completes the part (2).

Let  $X_3$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_3$ . Let the vertices of  $X_3$  form  $m_3$   $\text{Aut}(X_3)$ -orbits. Let  $K_3$  be as in Section 2. Let  $V_3 = V(K_3)$  be the vertex set of  $K_3$ . Let  $H_3$  be the group of all the translations of  $K_3$ . So,  $H_3 \leq \text{Aut}(K_3)$ .

Since  $X_3$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_3$  (as by Proposition 3.1  $K_3$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_3 : K_3 \rightarrow X_3$  where  $X_3 = K_3/\Gamma_3$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_3 \leq \text{Aut}(K_3)$ . Hence  $\Gamma_3$  consists of translations and glide reflections. Since  $X_3 = K_3/\Gamma_3$  is orientable,  $\Gamma_3$  does not contain any glide reflection. Thus  $\Gamma_3 \leq H_3$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $a_0$  and  $f_{-1}$  of  $K_3$  (see Section 2). Let  $\alpha_3 := a_1 - a_0$  and  $\beta_3 := d_0 - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_3 := \langle x \mapsto x + \alpha_3, x \mapsto x + \beta_3 \rangle$  and

$$G_3 = \{ \alpha : x \mapsto \varepsilon x + m_3 \alpha_3 + n_3 \beta_3 : \varepsilon = \pm 1, m_3, n_3 \in \mathbb{Z} \} \cong H_3 \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_3 \trianglelefteq G_3$  and the number of  $G_3/\Gamma_3$ -orbits of vertices of  $X_3$  is two. Therefore,  $G_3/\Gamma_3$  acts on  $X_3 = K_3/\Gamma_3$ . Since  $O_1 = \langle a_0 \rangle, O_2 = \langle b_0 \rangle$  are the  $G_3$ -orbits, it follows that  $\eta_3(O_1), \eta_3(O_2)$  are the  $(G_3/\Gamma_3)$ -orbits. Since the vertex set of  $X_3$  is  $\eta_3(O_1) \sqcup \eta_3(O_2)$  and  $G_3/\Gamma_3 \leq \text{Aut}(X_3)$ , it follows that the number of  $\text{Aut}(X_3)$ -orbits of vertices is  $\leq 2$  and hence,  $m_3 = 2$ . This completes the part (3).

Let  $X_4$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_4$ . Let the vertices of  $X_4$  form  $m_4$   $\text{Aut}(X_4)$ -orbits. Let  $K_4$  be as in Section 2. Let  $V_4 = V(K_4)$  be the vertex set of  $K_4$ . Let  $H_4$  be the group of all the translations of  $K_4$ . So,  $H_4 \leq \text{Aut}(K_4)$ .

Since  $X_4$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_4$  (as by Proposition 3.1  $K_4$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_4 : K_4 \rightarrow X_4$  where  $X_4 = K_4/\Gamma_4$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_4 \leq \text{Aut}(K_4)$ . Hence  $\Gamma_4$  consists of translations and glide reflections. Since  $X_4 = K_4/\Gamma_4$  is orientable,  $\Gamma_4$  does not contain any glide reflection. Thus  $\Gamma_4 \leq H_4$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $a_0$  and  $e_{-1}$  of  $K_4$  (see Section 2). Let  $\alpha_4 := a_1 - a_0$  and  $\beta_4 := d_0 - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_4 := \langle x \mapsto x + \alpha_4, x \mapsto x + \beta_4 \rangle$  and

$$G_4 = \{ \alpha : x \mapsto \varepsilon x + m_4 \alpha_4 + n_4 \beta_4 : \varepsilon = \pm 1, m_4, n_4 \in \mathbb{Z} \} \cong H_4 \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_4 \trianglelefteq G_4$  and the number of  $G_4/\Gamma_4$ -orbits of vertices of  $X_4$  is two. Therefore,  $G_4/\Gamma_4$  acts on  $X_4 = K_4/\Gamma_4$ . Since  $O_1 = \langle a_0 \rangle, O_2 = \langle b_0 \rangle$  are the  $G_4$ -orbits, it follows that  $\eta_4(O_1), \eta_4(O_2)$  are the  $(G_4/\Gamma_4)$ -orbits. Since the vertex set of  $X_4$  is  $\eta_4(O_1) \sqcup \eta_4(O_2)$  and  $G_4/\Gamma_4 \leq \text{Aut}(X_4)$ , it follows that the number of  $\text{Aut}(X_4)$ -orbits of vertices is  $\leq 2$  and hence,  $m_4 = 2$ . This completes the part (4).

Let  $X_5$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_5$ . Let the vertices of  $X_5$  form  $m_5$   $\text{Aut}(X_5)$ -orbits. Let  $K_5$  be as in Section 2. Let

$V_5 = V(K_5)$  be the vertex set of  $K_5$ . Let  $H_5$  be the group of all the translations of  $K_5$ . So,  $H_5 \leq \text{Aut}(K_5)$ .

Since  $X_5$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_5$  (as by Proposition 3.1  $K_5$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_5 : K_5 \rightarrow X_5$  where  $X_5 = K_5/\Gamma_5$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_5 \leq \text{Aut}(K_5)$ . Hence  $\Gamma_5$  consists of translations and glide reflections. Since  $X_5 = K_5/\Gamma_5$  is orientable,  $\Gamma_5$  does not contain any glide reflection. Thus  $\Gamma_5 \leq H_5$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $b_0$  and  $b_3$  of  $K_5$  (see Section 2). Let  $\alpha_5 := a_1 - a_0$ ,  $\beta_5 := a_7 - a_0$  and  $\gamma_5 := a_6 - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_5 := \langle x \mapsto x + \alpha_5, x \mapsto x + \beta_5, x \mapsto x + \gamma_5 \rangle$  and

$$G_5 = \{ \alpha : x \mapsto \varepsilon x + m_5 \alpha_5 + n_5 \beta_5 + r_5 \gamma_5 : \varepsilon = \pm 1, m_5, n_5, r_5 \in \mathbb{Z} \} \cong H_5 \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_5 \trianglelefteq G_5$  and the number of  $G_5/\Gamma_5$ -orbits of vertices of  $X_5$  is four. Therefore,  $G_5/\Gamma_5$  acts on  $X_5 = K_5/\Gamma_5$ . Since  $O_1 = \langle a_0 \rangle, O_2 = \langle b_0 \rangle, O_3 = \langle b_1 \rangle, O_4 = \langle b_2 \rangle$  are the  $G_5$ -orbits, it follows that  $\eta_5(O_1), \eta_5(O_2), \eta_5(O_3), \eta_5(O_4)$  are the  $(G_5/\Gamma_5)$ -orbits. Since the vertex set of  $X_5$  is  $\eta_5(O_1) \sqcup \eta_5(O_2) \sqcup \eta_5(O_3) \sqcup \eta_5(O_4)$  and  $G_5/\Gamma_5 \leq \text{Aut}(X_5)$ , it follows that the number of  $\text{Aut}(X_5)$ -orbits of vertices is  $\leq 4$ . This completes the part (5).

Let  $X_6$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_6$ . Let the vertices of  $X_6$  form  $m_6$   $\text{Aut}(X_6)$ -orbits. Let  $K_6$  be as in Section 2. Let  $V_6 = V(K_6)$  be the vertex set of  $K_6$ . Let  $H_6$  be the group of all the translations of  $K_6$ . So,  $H_6 \leq \text{Aut}(K_6)$ .

Since  $X_6$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_6$  (as by Proposition 3.1  $K_6$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_6 : K_6 \rightarrow X_6$  where  $X_6 = K_6/\Gamma_6$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_6 \leq \text{Aut}(K_6)$ . Hence  $\Gamma_6$  consists of translations and glide reflections. Since  $X_6 = K_6/\Gamma_6$  is orientable,  $\Gamma_6$  does not contain any glide reflection. Thus  $\Gamma_6 \leq H_6$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $a_0$  and  $a_6$  of  $K_6$  (see Section 2). Let  $\alpha_6 := a_3 - a_0$ ,  $\beta_6 := a_3 - a_{10}$  and  $\gamma_6 := a_8 - a_{40} \in \mathbb{R}^2$ . Similarly as above, define  $H_6 := \langle x \mapsto x + \alpha_6, x \mapsto x + \beta_6, x \mapsto x + \gamma_6 \rangle$  and

$$G_6 = \{ \alpha : x \mapsto \varepsilon x + m_6 \alpha_6 + n_6 \beta_6 + r_6 \gamma_6 : \varepsilon = \pm 1, m_6, n_6, r_6 \in \mathbb{Z} \} \cong H_6 \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_6 \trianglelefteq G_6$  and the number of  $G_6/\Gamma_6$ -orbits of vertices of  $X_6$  is seven. Therefore,  $G_6/\Gamma_6$  acts on  $X_6 = K_6/\Gamma_6$ . Since  $O_j = \langle a_j \rangle, j = 0, 1, \dots, 5$   $O_6 = \langle b_1 \rangle$  are the  $G_6$ -orbits, it follows that  $\eta_6(O_j), j = 0, 1, \dots, 6$  are the  $(G_6/\Gamma_6)$ -orbits. Since the vertex set of  $X_6$  is  $\sqcup_{j=0}^6 \eta_6(O_j)$  and  $G_6/\Gamma_6 \leq \text{Aut}(X_6)$ , it follows that the number of  $\text{Aut}(X_6)$ -orbits of vertices is  $\leq 7$ . This completes the part (6).

Let  $X_7$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_7$ . Let the vertices of  $X_7$  form  $m_7$   $\text{Aut}(X_7)$ -orbits. Let  $K_7$  be as in Section 2. Let  $V_7 = V(K_7)$  be the vertex set of  $K_7$ . Let  $H_7$  be the group of all the translations of  $K_7$ . So,  $H_7 \leq \text{Aut}(K_7)$ .

Since  $X_7$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_7$  (as by Proposition 3.1  $K_7$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_7 : K_7 \rightarrow X_7$  where  $X_7 = K_7/\Gamma_7$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_7 \leq \text{Aut}(K_7)$ . Hence  $\Gamma_7$  consists of translations and glide reflections. Since  $X_7 = K_7/\Gamma_7$  is orientable,  $\Gamma_7$  does not contain any glide reflection. Thus  $\Gamma_7 \leq H_7$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $b_0$  and  $k_0$  of  $K_7$  (see Section 2). Let  $\alpha_7 := a_3 - a_0$ ,  $\beta_7 := d_0 - a_0$  and  $\gamma_7 := d_{-3} - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_7 := \langle x \mapsto x + \alpha_7, x \mapsto x + \beta_7, x \mapsto x + \gamma_7 \rangle$  and

$$G_7 = \{ \alpha : x \mapsto \varepsilon x + m_7 \alpha_7 + n_7 \beta_7 + r_7 \gamma_7 : \varepsilon = \pm 1, m_7, n_7, r_7 \in \mathbb{Z} \} \cong H_7 \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_7 \trianglelefteq G_7$  and the number of  $G_7/\Gamma_7$ -orbits of vertices of  $X_7$  is four. Therefore,  $G_7/\Gamma_7$  acts on  $X_7 = K_7/\Gamma_7$ . Since  $O_1 = \langle a_0 \rangle$ ,  $O_2 = \langle b_0 \rangle$ ,  $O_3 = \langle a_1 \rangle$ ,  $O_4 = \langle k_1 \rangle$  are the  $G_7$ -orbits, it follows that  $O_1 = \langle a_0 \rangle$ ,  $O_2 = \langle b_0 \rangle$ ,  $O_3 = \langle a_1 \rangle$ ,  $O_4 = \langle k_1 \rangle$  are the  $(G_7/\Gamma_7)$ -orbits. Since the vertex set of  $X_7$  is  $\sqcup_{j=1}^4 \eta_7(O_j)$  and  $G_7/\Gamma_7 \leq \text{Aut}(X_7)$ , it follows that the number of  $\text{Aut}(X_7)$ -orbits of vertices is  $\leq 4$ . This completes the part (7).

Let  $X_8$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_8$ . Let the vertices of  $X_8$  form  $m_8$   $\text{Aut}(X_8)$ -orbits. Let  $K_8$  be as in Section 2. Let  $V_8 = V(K_8)$  be the vertex set of  $K_8$ . Let  $H_8$  be the group of all the translations of  $K_8$ . So,  $H_8 \leq \text{Aut}(K_8)$ .

Since  $X_8$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_8$  (as by Proposition 3.1  $K_8$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_8 : K_8 \rightarrow X_8$  where  $X_8 = K_8/\Gamma_8$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_8 \leq \text{Aut}(K_8)$ . Hence  $\Gamma_8$  consists of translations and glide reflections. Since  $X_8 = K_8/\Gamma_8$  is orientable,  $\Gamma_8$  does not contain any glide reflection. Thus  $\Gamma_8 \leq H_8$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $a_0$  and  $e_2$  of  $K_8$  (see Section 2). Let  $\alpha_8 := a_2 - a_0$  and  $\beta_8 := c_1 - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_8 := \langle x \mapsto x + \alpha_8, x \mapsto x + \beta_8 \rangle$  and

$$G_8 = \{ \alpha : x \mapsto \varepsilon x + m_8 \alpha_8 + n_8 \beta_8 : \varepsilon = \pm 1, m_8, n_8 \in \mathbb{Z} \} \cong H_8 \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_8 \trianglelefteq G_8$  and the number of  $G_8/\Gamma_8$ -orbits of vertices of  $X_8$  is three. Therefore,  $G_8/\Gamma_8$  acts on  $X_8 = K_8/\Gamma_8$ . Since  $O_1 = \langle c_1 \rangle$ ,  $O_2 = \langle c_2 \rangle$ ,  $O_3 = \langle c_3 \rangle$  are the  $G_8$ -orbits, it follows that  $\eta_8(O_1)$ ,  $\eta_8(O_2)$ ,  $\eta_8(O_3)$  are the  $(G_8/\Gamma_8)$ -orbits. Since the vertex set of  $X_8$  is  $\eta_8(O_1) \sqcup \eta_8(O_2) \sqcup \eta_8(O_3)$  and  $G_8/\Gamma_8 \leq \text{Aut}(X_8)$ , it follows that the number of  $\text{Aut}(X_8)$ -orbits of vertices is  $\leq 3$ . This completes the part (8).

Let  $X_9$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_9$ . Let the vertices of  $X_9$  form  $m_9$   $\text{Aut}(X_9)$ -orbits. Let  $K_9$  be as in Section 2. Let  $V_9 = V(K_9)$  be the vertex set of  $K_9$ . Let  $H_9$  be the group of all the translations of  $K_9$ . So,  $H_9 \leq \text{Aut}(K_9)$ .

Since  $X_9$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_9$  (as by Proposition 3.1  $K_9$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_9 : K_9 \rightarrow X_9$  where  $X_9 = K_9/\Gamma_9$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_9 \leq \text{Aut}(K_9)$ . Hence  $\Gamma_9$  consists of translations and glide reflections. Since  $X_9 = K_9/\Gamma_9$  is orientable,  $\Gamma_9$  does not contain any glide reflection. Thus  $\Gamma_9 \leq H_9$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $a_0$  and  $a_1$  of  $K_9$  (see Section 2). Let  $\alpha_9 := a_2 - a_0$ ,  $\beta_9 := e_1 - a_0$  and  $\gamma_9 := m_1 - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_9 := \langle x \mapsto x + \alpha_9, x \mapsto x + \beta_9, x \mapsto x + \gamma_9 \rangle$  and

$$G_9 = \{ \alpha : x \mapsto \varepsilon x + m_9 \alpha_9 + n_9 \beta_9 + r_9 \gamma_9 : \varepsilon = \pm 1, m_9, n_9, r_9 \in \mathbb{Z} \} \cong H_9 \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_9 \trianglelefteq G_9$  and the number of  $G_9/\Gamma_9$ -orbits of vertices of  $X_9$  is six. Therefore,  $G_9/\Gamma_9$  acts on  $X_9 = K_9/\Gamma_9$ . Since  $O_1 = \langle a_0 \rangle$ ,  $O_2 = \langle c_1 \rangle$ ,  $O_3 = \langle k_1 \rangle$ ,  $O_4 = \langle b_0 \rangle$ ,  $O_5 = \langle b_1 \rangle$ ,  $O_6 = \langle b_2 \rangle$  are the  $G_9$ -orbits, it follows that  $O_1 = \langle$

$a0 \rangle, O_2 = \langle c-1 \rangle, O_3 = \langle k-1 \rangle, O_4 = \langle b0 \rangle, O_5 = \langle b1 \rangle, O_6 = \langle b2 \rangle$  are the  $(G_9/\Gamma_9)$ -orbits. Since the vertex set of  $X_9$  is  $\sqcup_{j=1}^6 \eta_9(O_j)$  and  $G_9/\Gamma_9 \leq \text{Aut}(X_9)$ , it follows that the number of  $\text{Aut}(X_9)$ -orbits of vertices is  $\leq 6$ . This completes the part (9).

Let  $X_{10}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{10}$ . Let the vertices of  $X_{10}$  form  $m_{10}$   $\text{Aut}(X_{10})$ -orbits. Let  $K_{10}$  be as in Section 2. Let  $V_{10} = V(K_{10})$  be the vertex set of  $K_{10}$ . Let  $H_{10}$  be the group of all the translations of  $K_{10}$ . So,  $H_{10} \leq \text{Aut}(K_{10})$ .

Since  $X_{10}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{10}$  (as by Proposition 3.1  $K_{10}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{10} : K_{10} \rightarrow X_{10}$  where  $X_{10} = K_{10}/\Gamma_{10}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{10} \leq \text{Aut}(K_{10})$ . Hence  $\Gamma_{10}$  consists of translations and glide reflections. Since  $X_{10} = K_{10}/\Gamma_{10}$  is orientable,  $\Gamma_{10}$  does not contain any glide reflection. Thus  $\Gamma_{10} \leq H_{10}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $a1$  and  $b1$  of  $K_{10}$  (see Section 2). Let  $\alpha_{10} := b0 - f0$  and  $\beta_{10} := g2 - f0 \in \mathbb{R}^2$ . Similarly as above, define  $H_{10} := \langle x \mapsto x + \alpha_{10}, x \mapsto x + \beta_{10} \rangle$  and

$$G_{10} = \{ \alpha : x \mapsto \varepsilon x + m_{10}\alpha_{10} + n_{10}\beta_{10} : \varepsilon = \pm 1, m_{10}, n_{10} \in \mathbb{Z} \} \cong H_{10} \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{10} \trianglelefteq G_{10}$  and the number of  $G_{10}/\Gamma_{10}$ -orbits of vertices of  $X_{10}$  is three. Therefore,  $G_{10}/\Gamma_{10}$  acts on  $X_{10} = K_{10}/\Gamma_{10}$ . Since  $O_1 = \langle a0 \rangle, O_2 = \langle a1 \rangle, O_2 = \langle a2 \rangle$  are the  $G_{10}$ -orbits, it follows that  $\eta_{10}(O_1), \eta_{10}(O_2), \eta_{10}(O_3)$  are the  $(G_{10}/\Gamma_{10})$ -orbits. Since the vertex set of  $X_{10}$  is  $\eta_{10}(O_1) \sqcup \eta_{10}(O_2) \sqcup \eta_{10}(O_3)$  and  $G_{10}/\Gamma_{10} \leq \text{Aut}(X_{10})$ , it follows that the number of  $\text{Aut}(X_{10})$ -orbits of vertices is  $\leq 3$ . This completes the part (10).

Let  $X_{11}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{11}$ . Let the vertices of  $X_{11}$  form  $m_{11}$   $\text{Aut}(X_{11})$ -orbits. Let  $K_{11}$  be as in Section 2. Let  $V_{11} = V(K_{11})$  be the vertex set of  $K_{11}$ . Let  $H_{11}$  be the group of all the translations of  $K_{11}$ . So,  $H_{11} \leq \text{Aut}(K_{11})$ .

Since  $X_{11}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{11}$  (as by Proposition 3.1  $K_{11}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{11} : K_{11} \rightarrow X_{11}$  where  $X_{11} = K_{11}/\Gamma_{11}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{11} \leq \text{Aut}(K_{11})$ . Hence  $\Gamma_{11}$  consists of translations and glide reflections. Since  $X_{11} = K_{11}/\Gamma_{11}$  is orientable,  $\Gamma_{11}$  does not contain any glide reflection. Thus  $\Gamma_{11} \leq H_{11}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $a0$  and  $a3$  of  $K_{11}$  (see Section 2). Let  $\alpha_{11} := a17 - a5, \beta_{11} := a24 - a0$  and  $\gamma_{11} := a30 - a0 \in \mathbb{R}^2$ . Similarly as above, define  $H_{11} := \langle x \mapsto x + \alpha_{11}, x \mapsto x + \beta_{11}, x \mapsto x + \gamma_{11} \rangle$  and

$$G_{11} = \{ \alpha : x \mapsto \varepsilon x + m_{11}\alpha_{11} + n_{11}\beta_{11} + r_{11}\gamma_{11} : \varepsilon = \pm 1, m_{11}, n_{11}, r_{11} \in \mathbb{Z} \} \cong H_{11} \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{11} \trianglelefteq G_{11}$  and the number of  $G_{11}/\Gamma_{11}$ -orbits of vertices of  $X_{11}$  is six. Therefore,  $G_{11}/\Gamma_{11}$  acts on  $X_{11} = K_{11}/\Gamma_{11}$ . Since  $O_1 = \langle a1 \rangle, O_2 = \langle a2 \rangle, O_3 = \langle a3 \rangle, O_4 = \langle b2 \rangle, O_5 = \langle b4 \rangle, O_6 = \langle b6 \rangle$  are the  $G_{11}$ -orbits, it follows that  $O_1 = \langle a1 \rangle, O_2 = \langle a2 \rangle, O_3 = \langle a3 \rangle, O_4 = \langle b2 \rangle, O_5 = \langle b4 \rangle, O_6 = \langle b6 \rangle$  are the  $(G_{11}/\Gamma_{11})$ -orbits. Since the vertex set of  $X_{11}$  is  $\sqcup_{j=1}^6 \eta_{11}(O_j)$  and  $G_{11}/\Gamma_{11} \leq \text{Aut}(X_{11})$ , it follows that the number of  $\text{Aut}(X_{11})$ -orbits of vertices is  $\leq 6$ . This completes the part (11).

Let  $X_{12}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{12}$ . Let the vertices of  $X_{12}$  form  $m_{12}$   $\text{Aut}(X_{12})$ -orbits. Let  $K_{12}$  be as in Section 2.

Let  $V_{12} = V(K_{12})$  be the vertex set of  $K_{12}$ . Let  $H_{12}$  be the group of all the translations of  $K_{12}$ . So,  $H_{12} \leq \text{Aut}(K_{12})$ .

Since  $X_{12}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{12}$  (as by Proposition 3.1  $K_{12}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{12} : K_{12} \rightarrow X_{12}$  where  $X_{12} = K_{12}/\Gamma_{12}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{12} \leq \text{Aut}(K_{12})$ . Hence  $\Gamma_{12}$  consists of translations and glide reflections. Since  $X_{12} = K_{12}/\Gamma_{12}$  is orientable,  $\Gamma_{12}$  does not contain any glide reflection. Thus  $\Gamma_{12} \leq H_{12}$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $a_0$  and  $g_0$  of  $K_{12}$  (see Section 2). Let  $\alpha_{12} := a_1 - a_0$  and  $\beta_{12} := d_0 - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_{12} := \langle x \mapsto x + \alpha_{12}, x \mapsto x + \beta_{12} \rangle$  and

$$G_{12} = \{ \alpha : x \mapsto \varepsilon x + m_{12}\alpha_{12} + n_{12}\beta_{12} : \varepsilon = \pm 1, m_{12}, n_{12} \in \mathbb{Z} \} \cong H_{12} \times \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{12} \trianglelefteq G_{12}$  and the number of  $G_{12}/\Gamma_{12}$ -orbits of vertices of  $X_{12}$  is two. Therefore,  $G_{12}/\Gamma_{12}$  acts on  $X_{12} = K_{12}/\Gamma_{12}$ . Since  $O_1 = \langle a_0 \rangle, O_2 = \langle f_0 \rangle$  are the  $G_{12}$ -orbits, it follows that  $\eta_{12}(O_1), \eta_{12}(O_2)$  are the  $(G_{12}/\Gamma_{12})$ -orbits. Since the vertex set of  $X_{12}$  is  $\eta_{12}(O_1) \sqcup \eta_{12}(O_2)$  and  $G_{12}/\Gamma_{12} \leq \text{Aut}(X_{12})$ , it follows that the number of  $\text{Aut}(X_{12})$ -orbits of vertices is  $\leq 2$  and hence,  $m_{12} = 2$ . This completes the part (12).

Let  $X_{13}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{13}$ . Let the vertices of  $X_{13}$  form  $m_{13}$   $\text{Aut}(X_{13})$ -orbits. Let  $K_{13}$  be as in Section 2. Let  $V_{13} = V(K_{13})$  be the vertex set of  $K_{13}$ . Let  $H_{13}$  be the group of all the translations of  $K_{13}$ . So,  $H_{13} \leq \text{Aut}(K_{13})$ .

Since  $X_{13}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{13}$  (as by Proposition 3.1  $K_{13}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{13} : K_{13} \rightarrow X_{13}$  where  $X_{13} = K_{13}/\Gamma_{13}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{13} \leq \text{Aut}(K_{13})$ . Hence  $\Gamma_{13}$  consists of translations and glide reflections. Since  $X_{13} = K_{13}/\Gamma_{13}$  is orientable,  $\Gamma_{13}$  does not contain any glide reflection. Thus  $\Gamma_{13} \leq H_{13}$ .

We take origin  $(0, 0)$  is the middle point of the line segment joining vertices  $b_0$  and  $c_0$  of  $K_{13}$  (see Section 2). Let  $\alpha_{13} := a_1 - a_0$  and  $\beta_{13} := a_1 - k_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_{13} := \langle x \mapsto x + \alpha_{13}, x \mapsto x + \beta_{13} \rangle$  and

$$G_{13} = \{ \alpha : x \mapsto \varepsilon x + m_{13}\alpha_{13} + n_{13}\beta_{13} : \varepsilon = \pm 1, m_{13}, n_{13} \in \mathbb{Z} \} \cong H_{13} \times \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{13} \trianglelefteq G_{13}$  and the number of  $G_{13}/\Gamma_{13}$ -orbits of vertices of  $X_{13}$  is two. Therefore,  $G_{13}/\Gamma_{13}$  acts on  $X_{13} = K_{13}/\Gamma_{13}$ . Since  $O_1 = \langle a_0 \rangle, O_2 = \langle b_0 \rangle$  are the  $G_{13}$ -orbits, it follows that  $\eta_{13}(O_1), \eta_{13}(O_2)$  are the  $(G_{13}/\Gamma_{13})$ -orbits. Since the vertex set of  $X_{13}$  is  $\eta_{13}(O_1) \sqcup \eta_{13}(O_2)$  and  $G_{13}/\Gamma_{13} \leq \text{Aut}(X_{13})$ , it follows that the number of  $\text{Aut}(X_{13})$ -orbits of vertices is  $\leq 2$  and hence,  $m_{13} = 2$ . This completes the part (13).

Let  $X_{14}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{14}$ . Let the vertices of  $X_{14}$  form  $m_{14}$   $\text{Aut}(X_{14})$ -orbits. Let  $K_{14}$  be as in Section 2. Let  $V_{14} = V(K_{14})$  be the vertex set of  $K_{14}$ . Let  $H_{14}$  be the group of all the translations of  $K_{14}$ . So,  $H_{14} \leq \text{Aut}(K_{14})$ .

Since  $X_{14}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{14}$  (as by Proposition 3.1  $K_{14}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{14} : K_{14} \rightarrow X_{14}$  where  $X_{14} = K_{14}/\Gamma_{14}$  for some fixed element

(vertex, edge or face) free subgroup  $\Gamma_{14} \leq \text{Aut}(K_{14})$ . Hence  $\Gamma_{14}$  consists of translations and glide reflections. Since  $X_{14} = K_{14}/\Gamma_{14}$  is orientable,  $\Gamma_{14}$  does not contain any glide reflection. Thus  $\Gamma_{14} \leq H_{14}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $a_0$  and  $a_3$  of  $K_{14}$  (see Section 2). Let  $\alpha_{14} := a_2 - a_0$ ,  $\beta_{14} := a_1 - a_0$  and  $\gamma_{14} := a_3 - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_{14} := \langle x \mapsto x + \alpha_{14}, x \mapsto x + \beta_{14}, x \mapsto x + \gamma_{14} \rangle$  and

$$G_{14} = \{ \alpha : x \mapsto \varepsilon x + m_{14}\alpha_{14} + n_{14}\beta_{14} + r_{14}\gamma_{14} : \varepsilon = \pm 1, m_{14}, n_{14}, r_{14} \in \mathbb{Z} \} \cong H_{14} \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{14} \trianglelefteq G_{14}$  and the number of  $G_{14}/\Gamma_{14}$ -orbits of vertices of  $X_{14}$  is six. Therefore,  $G_{14}/\Gamma_{14}$  acts on  $X_{14} = K_{14}/\Gamma_{14}$ . Since  $O_1 = \langle a_0 \rangle$ ,  $O_2 = \langle a_1 \rangle$ ,  $O_3 = \langle a_2 \rangle$ ,  $O_4 = \langle b_3 \rangle$ ,  $O_5 = \langle b_5 \rangle$ ,  $O_6 = \langle b_7 \rangle$  are the  $G_{14}$ -orbits, it follows that  $O_1 = \langle a_0 \rangle$ ,  $O_2 = \langle a_1 \rangle$ ,  $O_3 = \langle a_2 \rangle$ ,  $O_4 = \langle b_3 \rangle$ ,  $O_5 = \langle b_5 \rangle$ ,  $O_6 = \langle b_7 \rangle$  are the  $(G_{14}/\Gamma_{14})$ -orbits. Since the vertex set of  $X_{14}$  is  $\sqcup_{j=1}^6 \eta_{14}(O_j)$  and  $G_{14}/\Gamma_{14} \leq \text{Aut}(X_{14})$ , it follows that the number of  $\text{Aut}(X_{14})$ -orbits of vertices is  $\leq 6$ . This completes the part (14).

Let  $X_{15}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{15}$ . Let the vertices of  $X_{15}$  form  $m_{15}$   $\text{Aut}(X_{15})$ -orbits. Let  $K_{15}$  be as in Section 2. Let  $V_{15} = V(K_{15})$  be the vertex set of  $K_{15}$ . Let  $H_{15}$  be the group of all the translations of  $K_{15}$ . So,  $H_{15} \leq \text{Aut}(K_{15})$ .

Since  $X_{15}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{15}$  (as by Proposition 3.1  $K_{15}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{15} : K_{15} \rightarrow X_{15}$  where  $X_{15} = K_{15}/\Gamma_{15}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{15} \leq \text{Aut}(K_{15})$ . Hence  $\Gamma_{15}$  consists of translations and glide reflections. Since  $X_{15} = K_{15}/\Gamma_{15}$  is orientable,  $\Gamma_{15}$  does not contain any glide reflection. Thus  $\Gamma_{15} \leq H_{15}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $b_0$  and  $b_1$  of  $K_{15}$  (see Section 2). Let  $\alpha_{15} := b_1 - b_0$  and  $\beta_{15} := c_0 - a_0 \in \mathbb{R}^2$ . Similarly as above, define  $H_{15} := \langle x \mapsto x + \alpha_{15}, x \mapsto x + \beta_{15} \rangle$  and

$$G_{15} = \{ \alpha : x \mapsto \varepsilon x + m_{15}\alpha_{15} + n_{15}\beta_{15} : \varepsilon = \pm 1, m_{15}, n_{15} \in \mathbb{Z} \} \cong H_{15} \rtimes \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{15} \trianglelefteq G_{15}$  and the number of  $G_{15}/\Gamma_{15}$ -orbits of vertices of  $X_{15}$  is two. Therefore,  $G_{15}/\Gamma_{15}$  acts on  $X_{15} = K_{15}/\Gamma_{15}$ . Since  $O_1 = \langle a_0 \rangle$ ,  $O_2 = \langle b_0 \rangle$  are the  $G_{15}$ -orbits, it follows that  $\eta_{15}(O_1), \eta_{15}(O_2)$  are the  $(G_{15}/\Gamma_{15})$ -orbits. Since the vertex set of  $X_{15}$  is  $\eta_{15}(O_1) \sqcup \eta_{15}(O_2)$  and  $G_{15}/\Gamma_{15} \leq \text{Aut}(X_{15})$ , it follows that the number of  $\text{Aut}(X_{15})$ -orbits of vertices is  $\leq 2$  and hence,  $m_{15} = 2$ . This completes the part (15).

Let  $X_{16}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{16}$ . Let the vertices of  $X_{16}$  form  $m_{16}$   $\text{Aut}(X_{16})$ -orbits. Let  $K_{16}$  be as in Section 2. Let  $V_{16} = V(K_{16})$  be the vertex set of  $K_{16}$ . Let  $H_{16}$  be the group of all the translations of  $K_{16}$ . So,  $H_{16} \leq \text{Aut}(K_{16})$ .

Since  $X_{16}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{16}$  (as by Proposition 3.1  $K_{16}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{16} : K_{16} \rightarrow X_{16}$  where  $X_{16} = K_{16}/\Gamma_{16}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{16} \leq \text{Aut}(K_{16})$ . Hence  $\Gamma_{16}$  consists of translations and glide reflections. Since  $X_{16} = K_{16}/\Gamma_{16}$  is orientable,  $\Gamma_{16}$  does not contain any glide reflection. Thus  $\Gamma_{16} \leq H_{16}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $a_0$  and  $a_6$  of  $K_{16}$  (see Section 2). Let  $\alpha_{16} := a_6 - a_1$  and  $\beta_{16} := a_2 - a_9 \in \mathbb{R}^2$ . Similarly as above,

define  $H_{16} := \langle x \mapsto x + \alpha_{16}, x \mapsto x + \beta_{16} \rangle$  and

$$G_{16} = \{ \alpha : x \mapsto \varepsilon x + m_{16}\alpha_{16} + n_{16}\beta_{16} : \varepsilon = \pm 1, m_{16}, n_{16} \in \mathbb{Z} \} \cong H_{16} \times \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{16} \trianglelefteq G_{16}$  and the number of  $G_{16}/\Gamma_{16}$ -orbits of vertices of  $X_{16}$  is three. Therefore,  $G_{16}/\Gamma_{16}$  acts on  $X_{16} = K_{16}/\Gamma_{16}$ . Since  $O_1 = \langle a0 \rangle, O_2 = \langle a2 \rangle, O_3 = \langle a1 \rangle$  are the  $G_{16}$ -orbits, it follows that  $\eta_{16}(O_1), \eta_{16}(O_2), \eta_{16}(O_3)$  are the  $(G_{16}/\Gamma_{16})$ -orbits. Since the vertex set of  $X_{16}$  is  $\eta_{16}(O_1) \sqcup \eta_{16}(O_2) \sqcup \eta_{16}(O_3)$  and  $G_{16}/\Gamma_{16} \leq \text{Aut}(X_{16})$ , it follows that the number of  $\text{Aut}(X_{16})$ -orbits of vertices is  $\leq 2$  and hence,  $m_{16} \leq 3$ . This completes the part (16).

Let  $X_{17}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{17}$ . Let the vertices of  $X_{17}$  form  $m_{17}$   $\text{Aut}(X_{17})$ -orbits. Let  $K_{17}$  be as in Section 2. Let  $V_{17} = V(K_{17})$  be the vertex set of  $K_{17}$ . Let  $H_{17}$  be the group of all the translations of  $K_{17}$ . So,  $H_{17} \leq \text{Aut}(K_{17})$ .

Since  $X_{17}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{17}$  (as by Proposition 3.1  $K_{17}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{17} : K_{17} \rightarrow X_{17}$  where  $X_{17} = K_{17}/\Gamma_{17}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{17} \leq \text{Aut}(K_{17})$ . Hence  $\Gamma_{17}$  consists of translations and glide reflections. Since  $X_{17} = K_{17}/\Gamma_{17}$  is orientable,  $\Gamma_{17}$  does not contain any glide reflection. Thus  $\Gamma_{17} \leq H_{17}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $a6$  and  $a9$  of  $K_{17}$  (see Section 2). Let  $\alpha_{17} := a6 - a31, \beta_{17} := a18 - a31$  and  $\gamma_{17} := a56 - a31 \in \mathbb{R}^2$ . Similarly as above, define  $H_{17} := \langle x \mapsto x + \alpha_{17}, x \mapsto x + \beta_{17}, x \mapsto x + \gamma_{17} \rangle$  and

$$G_{17} = \{ \alpha : x \mapsto \varepsilon x + m_{17}\alpha_{17} + n_{17}\beta_{17} + r_{17}\gamma_{17} : \varepsilon = \pm 1, m_{17}, n_{17}, r_{17} \in \mathbb{Z} \} \cong H_{17} \times \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{17} \trianglelefteq G_{17}$  and the number of  $G_{17}/\Gamma_{17}$ -orbits of vertices of  $X_{17}$  is nine. Therefore,  $G_{17}/\Gamma_{17}$  acts on  $X_{17} = K_{17}/\Gamma_{17}$ . Since  $O_1 = \langle a6 \rangle, O_2 = \langle a7 \rangle, O_3 = \langle a8 \rangle, O_4 = \langle a1 \rangle, O_5 = \langle a12 \rangle, O_6 = \langle a13 \rangle, O_7 = \langle a44 \rangle, O_8 = \langle a43 \rangle, O_9 = \langle a69 \rangle$  are the  $G_{17}$ -orbits, it follows that  $O_1 = \langle a6 \rangle, O_2 = \langle a7 \rangle, O_3 = \langle a8 \rangle, O_4 = \langle a1 \rangle, O_5 = \langle a12 \rangle, O_6 = \langle a13 \rangle, O_7 = \langle a44 \rangle, O_8 = \langle a43 \rangle, O_9 = \langle a69 \rangle$  are the  $(G_{17}/\Gamma_{17})$ -orbits. Since the vertex set of  $X_{17}$  is  $\sqcup_{j=1}^9 \eta_{17}(O_j)$  and  $G_{17}/\Gamma_{17} \leq \text{Aut}(X_{17})$ , it follows that the number of  $\text{Aut}(X_{17})$ -orbits of vertices is  $\leq 9$ . This completes the part (17).

Let  $X_{18}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{18}$ . Let the vertices of  $X_{18}$  form  $m_{18}$   $\text{Aut}(X_{18})$ -orbits. Let  $K_{18}$  be as in Section 2. Let  $V_{18} = V(K_{18})$  be the vertex set of  $K_{18}$ . Let  $H_{18}$  be the group of all the translations of  $K_{18}$ . So,  $H_{18} \leq \text{Aut}(K_{18})$ .

Since  $X_{18}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{18}$  (as by Proposition 3.1  $K_{18}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{18} : K_{18} \rightarrow X_{18}$  where  $X_{18} = K_{18}/\Gamma_{18}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{18} \leq \text{Aut}(K_{18})$ . Hence  $\Gamma_{18}$  consists of translations and glide reflections. Since  $X_{18} = K_{18}/\Gamma_{18}$  is orientable,  $\Gamma_{18}$  does not contain any glide reflection. Thus  $\Gamma_{18} \leq H_{18}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $b0$  and  $b1$  of  $K_{18}$  (see Section 2). Let  $\alpha_{18} := b1 - b0$  and  $\beta_{18} := d1 - a1 \in \mathbb{R}^2$ . Similarly as above, define  $H_{18} := \langle x \mapsto x + \alpha_{18}, x \mapsto x + \beta_{18} \rangle$  and

$$G_{18} = \{ \alpha : x \mapsto \varepsilon x + m_{18}\alpha_{18} + n_{18}\beta_{18} : \varepsilon = \pm 1, m_{18}, n_{18} \in \mathbb{Z} \} \cong H_{18} \times \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{18} \trianglelefteq G_{18}$  and the number of  $G_{18}/\Gamma_{18}$ -orbits of vertices of  $X_{18}$  is three. Therefore,  $G_{18}/\Gamma_{18}$  acts on  $X_{18} = K_{18}/\Gamma_{18}$ . Since  $O_1 = \langle c0 \rangle$ ,  $O_2 = \langle c1 \rangle$ ,  $O_3 = \langle b1 \rangle$  are the  $G_{18}$ -orbits, it follows that  $\eta_{18}(O_1), \eta_{18}(O_2), \eta_{18}(O_3)$  are the  $(G_{18}/\Gamma_{18})$ -orbits. Since the vertex set of  $X_{18}$  is  $\eta_{18}(O_1) \sqcup \eta_{18}(O_2) \sqcup \eta_{18}(O_3)$  and  $G_{18}/\Gamma_{18} \leq \text{Aut}(X_{18})$ , it follows that the number of  $\text{Aut}(X_{18})$ -orbits of vertices is  $\leq 3$ . This completes the part (18).

Let  $X_{19}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{19}$ . Let the vertices of  $X_{19}$  form  $m_{19}$   $\text{Aut}(X_{19})$ -orbits. Let  $K_{19}$  be as in Section 2. Let  $V_{19} = V(K_{19})$  be the vertex set of  $K_{19}$ . Let  $H_{19}$  be the group of all the translations of  $K_{19}$ . So,  $H_{19} \leq \text{Aut}(K_{19})$ .

Since  $X_{19}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{19}$  (as by Proposition 3.1  $K_{19}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{19} : K_{19} \rightarrow X_{19}$  where  $X_{19} = K_{19}/\Gamma_{19}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{19} \leq \text{Aut}(K_{19})$ . Hence  $\Gamma_{19}$  consists of translations and glide reflections. Since  $X_{19} = K_{19}/\Gamma_{19}$  is orientable,  $\Gamma_{19}$  does not contain any glide reflection. Thus  $\Gamma_{19} \leq H_{19}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $b0$  and  $b1$  of  $K_{19}$  (see Section 2). Let  $\alpha_{19} := b1 - b0$  and  $\beta_{19} := d1 - a0 \in \mathbb{R}^2$ . Similarly as above, define  $H_{19} := \langle x \mapsto x + \alpha_{19}, x \mapsto x + \beta_{19} \rangle$  and

$$G_{19} = \{ \alpha : x \mapsto \varepsilon x + m_{19}\alpha_{19} + n_{19}\beta_{19} : \varepsilon = \pm 1, m_{19}, n_{19} \in \mathbb{Z} \} \cong H_{19} \times \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{19} \trianglelefteq G_{19}$  and the number of  $G_{19}/\Gamma_{19}$ -orbits of vertices of  $X_{19}$  is three. Therefore,  $G_{19}/\Gamma_{19}$  acts on  $X_{19} = K_{19}/\Gamma_{19}$ . Since  $O_1 = \langle c0 \rangle$ ,  $O_2 = \langle c1 \rangle$ ,  $O_3 = \langle b1 \rangle$  are the  $G_{19}$ -orbits, it follows that  $\eta_{19}(O_1), \eta_{19}(O_2), \eta_{19}(O_3)$  are the  $(G_{19}/\Gamma_{19})$ -orbits. Since the vertex set of  $X_{19}$  is  $\eta_{19}(O_1) \sqcup \eta_{19}(O_2) \sqcup \eta_{19}(O_3)$  and  $G_{19}/\Gamma_{19} \leq \text{Aut}(X_{19})$ , it follows that the number of  $\text{Aut}(X_{19})$ -orbits of vertices is  $\leq 3$ . This completes the part (19).

Let  $X_{20}$  be a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{20}$ . Let the vertices of  $X_{20}$  form  $m_{20}$   $\text{Aut}(X_{20})$ -orbits. Let  $K_{20}$  be as in Section 2. Let  $V_{20} = V(K_{20})$  be the vertex set of  $K_{20}$ . Let  $H_{20}$  be the group of all the translations of  $K_{20}$ . So,  $H_{20} \leq \text{Aut}(K_{20})$ .

Since  $X_{20}$  is a 2-semiequivelar map on the torus that is the quotient of the plane's 2-uniform lattice  $K_{20}$  (as by Proposition 3.1  $K_{20}$  is unique), so, we can assume, there is a polyhedral covering map  $\eta_{20} : K_{20} \rightarrow X_{20}$  where  $X_{20} = K_{20}/\Gamma_{20}$  for some fixed element (vertex, edge or face) free subgroup  $\Gamma_{20} \leq \text{Aut}(K_{20})$ . Hence  $\Gamma_{20}$  consists of translations and glide reflections. Since  $X_{20} = K_{20}/\Gamma_{20}$  is orientable,  $\Gamma_{20}$  does not contain any glide reflection. Thus  $\Gamma_{20} \leq H_{20}$ .

We take origin  $(0,0)$  is the middle point of the line segment joining vertices  $a0$  and  $a6$  of  $K_{20}$  (see Section 2). Let  $\alpha_{20} := a28 - a6$ ,  $\beta_{20} := a19 - a10$  and  $\gamma_{20} := a16 - a7 \in \mathbb{R}^2$ . Similarly as above, define  $H_{20} := \langle x \mapsto x + \alpha_{20}, x \mapsto x + \beta_{20}, x \mapsto x + \gamma_{20} \rangle$  and

$$G_{20} = \{ \alpha : x \mapsto \varepsilon x + m_{20}\alpha_{20} + n_{20}\beta_{20} + r_{20}\gamma_{20} : \varepsilon = \pm 1, m_{20}, n_{20}, r_{20} \in \mathbb{Z} \} \cong H_{20} \times \mathbb{Z}_2.$$

By the same arguments as above and in Claim 1,  $\Gamma_{20} \trianglelefteq G_{20}$  and the number of  $G_{20}/\Gamma_{20}$ -orbits of vertices of  $X_{20}$  is nine. Therefore,  $G_{20}/\Gamma_{20}$  acts on  $X_{20} = K_{20}/\Gamma_{20}$ . Since  $O_1 = \langle a1 \rangle$ ,  $O_2 = \langle a2 \rangle$ ,  $O_3 = \langle a3 \rangle$ ,  $O_4 = \langle a4 \rangle$ ,  $O_5 = \langle a5 \rangle$ ,  $O_6 = \langle a6 \rangle$ ,  $O_7 = \langle b1 \rangle$ ,  $O_8 = \langle b2 \rangle$ ,  $O_9 = \langle b12 \rangle$  are the  $G_{20}$ -orbits, it follows that  $O_1 = \langle a1 \rangle$ ,  $O_2 = \langle a2 \rangle$ ,  $O_3 = \langle a3 \rangle$ ,  $O_4 = \langle a4 \rangle$ ,  $O_5 = \langle a5 \rangle$ ,  $O_6 = \langle a6 \rangle$ ,  $O_7 = \langle b1 \rangle$ ,  $O_8 = \langle b2 \rangle$ ,  $O_9 = \langle b12 \rangle$  are the



$(G_{20}/\Gamma_{20})$ -orbits. Since the vertex set of  $X_{20}$  is  $\sqcup_{j=1}^9 \eta_{20}(O_j)$  and  $G_{20}/\Gamma_{20} \leq \text{Aut}(X_{20})$ , it follows that the number of  $\text{Aut}(X_{20})$ -orbits of vertices is  $\leq 9$ . This completes the part (20).  $\square$

## References

- [1] L. Babai, *Vertex-transitive graphs and vertex-transitive maps*, J. Graph Theory **15** (1991), 587–627.
- [2] U. Brehm and W. Kühnel, *Equivelar maps on the torus*, European J. Combin. **29** (2008), 1843–1861.
- [3] H. S. M. Coxeter and W. O. J. Moser, *Generators and Relations for Discrete Groups* (4th edition), Springer-Verlag, Berlin-New York, 1980.
- [4] B. Datta, *A note on the existence of  $\{k, k\}$ -equivelar polyhedral maps*, Beiträge Algebra Geom. **46** (2005), 537–544.
- [5] B. Datta and D. Maity, *Semi-equivelar and vertex-transitive maps on the torus*, *Beiträge Algebra Geom.*, **58** (2017), pp. 617–634.
- [6] B. Datta and D. Maity, *Semi-equivelar maps on the torus and the Klein bottle are Archimedean*, Discrete Math. **341** (12) (2018), 329–3309.
- [7] B. Datta and N. Nilakantan, *Equivelar polyhedra with few vertices*, Discrete & Comput Geom. **26** (2001), 429–461.
- [8] B. Datta and A. K. Upadhyay, *Degree-regular triangulations of torus and Klein bottle*, Proc. Indian Acad. Sci. (Math. Sci.) **115** (2005), 279–307.
- [9] Otto Krötenheerdt, *Die homogenen Mosaik n-ter Ordnung in der euklidischen Ebene. I, II, III*. Wiss. Z. Martin-LUther-Univ. Halle-Wittenberg Math.-Natur. Reihe 18 (1969), pp. 273–290, 19 (1970), pp. 19–38 and 97–241.
- [10] L. Fejes Tóth, *Reguläre Figuren*, Akadémiai Kiadó, Budapest, 1965. (English translation: *Regular Figures*, Pergmon Press, Oxford, 1964.
- [11] B. Grünbaum and G. C. Shephard, *Tilings by regular polygons: Patterns in the plane from Kepler to the present, including recent results and unsolved problems*, Math. Mag. **50** (1977), 227–247.
- [12] B. Grünbaum and G. C. Shephard, *The geometry of planar graphs*. Combinatorics (Swansea, 1981), pp. 124–150, London Math. Soc. LNS **52**, Cambridge Univ. Press, Cambridge, 1981.
- [13] D. Maity, D. Bhowmik and M. M. Kharkongor *Quotient maps of n-uniform ( $3 \leq n \leq 6$ ) tilings of the plane on the torus and the Klein bottle*, (preprint).
- [14] D. Maity and A. K. Upadhyay, *On enumeration of a class of maps on Klein bottle*, Contrib. Discrete Math. **13** (2018), no. 1, 79–119..
- [15] E. H. Spanier, *Algebraic Topology*, Springer-Verlag, New York, 1966.
- [16] O. Süch, *Vertex transitive maps on a torus*, Acta Math. Univ. Comenianae **53** (2011), 1–30.
- [17] O. Süch, *Vertex transitive maps on the Klein bottle*, ARS Math. Contemporanea **4** (2011), 363–374.