

A NOTE ON THE COEFFICIENTS OF POWER SUMS OF ARITHMETIC PROGRESSIONS

JOSÉ LUIS CERECEDA

ABSTRACT. In this note we show a simple formula for the coefficients of the polynomial associated with the sums of powers of the terms of an arbitrary arithmetic progression. This formula consists of a double sum involving only ordinary binomial coefficients and binomial powers. Arguably, this is the simplest formula that can probably be found for the said coefficients. Furthermore, as a by-product, we give an explicit formula for the Bernoulli polynomials involving the Stirling numbers of the first and second kind.

1. INTRODUCTION

Consider the sum of k th powers of the terms of an arithmetic progression with first term r and common difference m

$$S_k^{m,r}(n) = r^k + (m+r)^k + (2m+r)^k + \cdots + ((n-1)m+r)^k,$$

where k, m, r , and n are assumed to be integer variables with $m, n \geq 1$ and $k, r \geq 0$. $S_k^{m,r}(n)$ turns out to be a polynomial in n of degree $k+1$ without constant term, that is, it can be expressed in the form $S_k^{m,r}(n) = \sum_{t=1}^{k+1} c_{k,t}^{m,r} n^t$ for certain rational coefficients $c_{k,t}^{m,r}$.

Griffiths [5] obtained the following formula for $c_{k,t}^{m,r}$ (in our notation)

$$c_{k,t}^{m,r} = m^k \sum_{j=t}^{k+1} \sum_{i=j}^{k+1} \frac{1}{i} \left(\frac{r}{m}\right)^{j-t} \binom{j}{t} S_1(i, j) S_2(k, i-1), \quad (1)$$

for $t = 1, \dots, k+1$, where $S_1(i, j)$ are the (signed) Stirling numbers of the first kind and $S_2(k, i-1)$ are the (unsigned) Stirling numbers of the second kind. On the other hand, the authors of [1] gave the following formula for $c_{k,t}^{m,r}$

$$c_{k,t}^{m,r} = \sum_{j=0}^k \frac{m^j W_{m,r}(k, j)}{j+1} S_1(j+1, t), \quad (2)$$

where $S_1(j+1, t)$ are the (signed) Stirling numbers of the first kind and $W_{m,r}(k, j)$ are the so-called r -Whitney numbers of the second kind, which are defined by [8]

$$W_{m,r}(k, j) = \frac{1}{m^j j!} \sum_{i=0}^j (-1)^{j-i} \binom{j}{i} (mi+r)^k.$$

Moreover, from [11, Theorems 5 and 6], it can be shown that

$$c_{k,t}^{m,r} = \frac{(-1)^{k+1-t}}{(k+1)!} \sum_{j=0}^k \sum_{s=t}^{k+1} A_{m,r}(k, k-j) \binom{s}{t} (k-j)^{s-t} S_1(k+1, s), \quad (3)$$

where $S_1(k+1, s)$ are the (signed) Stirling numbers of the first kind, and

$$A_{m,r}(k, j) = \sum_{i=0}^j (-1)^i \binom{k+1}{i} [m(j-i) + r]^k.$$

It is to be noted that the polynomial $S_k^{m,r}(n) = \sum_{t=1}^{k+1} c_{k,t}^{m,r} n^t$ with coefficients given by (1), (2), and (3) can equivalently be expressed, for $k \geq 0$, as

$$S_k^{m,r}(n) = m^k \sum_{j=0}^k j! S_2(k, j) \left[\binom{n + \frac{r}{m}}{j+1} - \binom{\frac{r}{m}}{j+1} \right],$$

$$S_k^{m,r}(n) = \sum_{j=0}^k m^j j! W_{m,r}(k, j) \binom{n}{j+1},$$

and

$$S_k^{m,r}(n) = \sum_{j=0}^k A_{m,r}(k, k-j) \binom{n+j}{k+1},$$

respectively. For other explicit formulas concerning the polynomial $S_k^{m,r}(n)$ as a whole see, for example, [7, 3, 14].

2. A SIMPLER FORMULA FOR $c_{k,t}^{m,r}$

The main purpose of this note is to show that a simpler formula for $c_{k,t}^{m,r}$, as compared to the formulas in (1), (2), and (3), can be derived from the following alternative expression for $S_k^{m,r}(n)$ (see, e.g., [1, Equation 11] and [2, Note 3])

$$S_k^{m,r}(n) = \frac{m^k}{k+1} \sum_{t=1}^{k+1} \binom{k+1}{t} B_{k+1-t} \left(\frac{r}{m} \right) n^t, \quad k \geq 0, \quad (4)$$

where $B_k(x)$ are the Bernoulli polynomials with generating function

$$\frac{te^{xt}}{e^t - 1} = \sum_{k=0}^{\infty} B_k(x) \frac{t^k}{k!}, \quad (|t| < 2\pi).$$

Equation (4) generalizes the well-known formula for the ordinary power sum $S_k^{1,0}(n) = 0^k + 1^k + \dots + (n-1)^k$, namely

$$S_k^{1,0}(n) = \frac{1}{k+1} \sum_{t=1}^{k+1} \binom{k+1}{t} B_{k+1-t} n^t, \quad k \geq 0,$$

where $B_k = B_k(0)$ are the Bernoulli numbers.

Indeed, by using the following specific representation for $B_k(x)$ (see, e.g., [13, Equation (38)] and [12, Equation (18)])

$$B_k(x) = \sum_{j=0}^k \sum_{i=0}^j \frac{(-1)^i}{j+1} \binom{j}{i} (i+x)^k, \quad k \geq 0, \quad (5)$$

we quickly obtain from (4)

$$c_{k,t}^{m,r} = \frac{m^k}{k+1} \binom{k+1}{t} \sum_{j=0}^{k+1-t} \sum_{i=0}^j \frac{(-1)^i}{j+1} \binom{j}{i} \left(i + \frac{r}{m}\right)^{k+1-t}, \quad (6)$$

which gives us $c_{k,t}^{m,r}$ as a double sum involving only ordinary binomial coefficients and binomial powers. Alternatively, we can write (6) as

$$c_{k,t}^{m,r} = \frac{m^{t-1}}{k+1} \binom{k+1}{t} \sum_{j=0}^{k+1-t} (-1)^j \frac{m^j j!}{j+1} W_{m,r}(k+1-t, j). \quad (7)$$

This last formula is simpler than those in (1), (2), and (3) because each of the numbers $S_1(k, j)$, $S_2(k, j)$, and $A_{m,r}(k, j)$, carries at least one extra summation of its own. In particular, the (signed) Stirling numbers of the first kind are given explicitly by

$$S_1(k, j) = \sum_{t=0}^{k-j} \sum_{r=0}^t \frac{(-1)^r}{t!} \binom{k+t-1}{k+t-j} \binom{2k-j}{k-t-j} \binom{t}{r} r^{k-j+t},$$

so that each of the formulas (1), (2), and (3) turns out to be more complex than (7). Indeed, as shall be argued shortly, (6), or its equivalent form (7), can be regarded as the simplest formula attainable for $c_{k,t}^{m,r}$.

Moreover, using (7), one can readily obtain the first few coefficients of highest degree, namely,

$$\begin{aligned} c_{k,k+1}^{m,r} &= \frac{m^k}{k+1}, \quad k \geq 0, \\ c_{k,k}^{m,r} &= m^{k-1} \left(r - \frac{m}{2}\right), \quad k \geq 1, \\ c_{k,k-1}^{m,r} &= \frac{1}{12} k m^{k-2} (m^2 - 6mr + 6r^2), \quad k \geq 2, \\ c_{k,k-2}^{m,r} &= \frac{1}{12} k(k-1) m^{k-3} r (m^2 - 3mr + 2r^2), \quad k \geq 3, \\ c_{k,k-3}^{m,r} &= \frac{1}{720} k(k-1)(k-2) m^{k-4} (m^4 - 30m^2 r^2 + 60mr^3 - 30r^4), \quad k \geq 4, \end{aligned}$$

etc.

3. GOULD'S ASSERTION

Setting $x = 0$ in (5) yields the following well-known explicit formula for the Bernoulli numbers [4, Equation 1]

$$B_k = \sum_{j=0}^k \sum_{i=0}^j \frac{(-1)^i}{j+1} \binom{j}{i} i^k, \quad k \geq 0, \quad (8)$$

or, equivalently,

$$B_k = \sum_{j=0}^k (-1)^j \frac{j!}{j+1} S_2(k, j), \quad k \geq 0. \quad (9)$$

Regarding the structure of explicit formulas for Bernoulli numbers, it is worth recalling Gould's assertion appearing at the end of his definitive paper [4]:

- Gould's assertion: "*the writer has seen no formula for B_k which does not require at least two actual summations.*"

This statement amounts to the conjecture that, in fact, there does not exist any elementary formula for B_k involving just one (finite) summation. Hence, in the spirit of Gould's assertion, we can make the ansatz that the above formula (8) for B_k (or its equivalent form (9)), as well as the above formula (5) for $B_k(x)$, constitutes, in a suitably defined sense, the simplest explicit formula for the Bernoulli numbers and the Bernoulli polynomials, respectively. Consequently, with this proviso, and noting that

$$c_{k,t}^{m,r} = \frac{m^k}{k+1} \binom{k+1}{t} B_{k+1-t} \left(\frac{r}{m} \right),$$

we may conclude that the expression in (6), or its equivalent form (7), constitutes in turn the simplest formula for $c_{k,t}^{m,r}$ one is ever likely to find.

4. CONCLUDING REMARKS

We end this note with the following two remarks.

Remark 1. As a by-product, from (1) and (4) we can readily obtain an explicit formula for the Bernoulli polynomials involving the Stirling numbers of the first and second kind. Indeed, for $t = 1$, (1) becomes

$$c_{k,1}^{m,r} = m^k \sum_{j=1}^{k+1} \sum_{i=j}^{k+1} \frac{j}{i} \left(\frac{r}{m} \right)^{j-1} S_1(i, j) S_2(k, i-1),$$

while, from (4), it follows that

$$c_{k,1}^{m,r} = m^k B_k \left(\frac{r}{m} \right).$$

Therefore, equating the right-hand sides of the last two equations, and replacing the rational $\frac{r}{m}$ by the arbitrary variable x , we find that

$$B_k(x) = \sum_{j=0}^k \sum_{i=j}^k \frac{j+1}{i+1} S_1(i+1, j+1) S_2(k, i) x^j, \quad k \geq 0, \quad (10)$$

which reduces to (9) when $x = 0$. Let us also observe that, by using the property $B_k(1) = (-1)^k B_k$, from (10) we get the identity

$$B_k = (-1)^k \sum_{j=0}^k \sum_{i=j}^k \frac{j+1}{i+1} S_1(i+1, j+1) S_2(k, i), \quad k \geq 0.$$

Moreover, from (10) and $B_k(x) = \sum_{j=0}^k \binom{k}{j} B_{k-j} x^j$, it turns out that

$$\binom{k}{j} B_{k-j} = \sum_{i=j}^k \frac{j+1}{i+1} S_1(i+1, j+1) S_2(k, i), \quad 0 \leq j \leq k,$$

which should be compared with the alternative expression [10, Equation (15.41)]

$$\binom{k}{j} B_{k-j} = k \sum_{i=j}^k \frac{1}{i} S_1(i, j) S_2(k-1, i-1), \quad 1 \leq j \leq k,$$

or, equivalently,

$$\binom{k+1}{j+1} B_{k-j} = (k+1) \sum_{i=j}^k \frac{1}{i+1} S_1(i+1, j+1) S_2(k, i), \quad 0 \leq j \leq k.$$

Remark 2. For $t = 1$, (2), as well as (7), becomes

$$c_{k,1}^{m,r} = \sum_{j=0}^k (-1)^j \frac{m^j j!}{j+1} W_{m,r}(k, j),$$

and then

$$B_k \left(\frac{r}{m} \right) = \sum_{j=0}^k (-1)^j \frac{j!}{j+1} \frac{W_{m,r}(k, j)}{m^{k-j}}, \quad k \geq 0,$$

which was obtained in [9] as a consequence of [9, Theorem 1]. Moreover, when $m = 1$ the last formula reduces to

$$B_k(r) = \sum_{j=0}^k (-1)^j \frac{j!}{j+1} S_{2,r}(k+r, j+r),$$

thus retrieving [6, Theorem 1], where $S_{2,r}(k+r, j+r)$ denotes the r -Stirling numbers of the second kind, namely

$$S_{2,r}(k+r, j+r) = \frac{1}{j!} \sum_{i=0}^j (-1)^{j-i} \binom{j}{i} (i+r)^k.$$

Notice that, by combining the last two equations, we recover (5) with $x = r$.

REFERENCES

- [1] A. Bazsó and I. Mező, On the coefficients of power sums of arithmetic progressions, *J. Number Theory*, 153, 117–123, 2015.
- [2] J. L. Cereceda, Generalized Stirling numbers and sums of powers of arithmetic progressions, *Internat. J. Math. Ed. Sci. Tech.*, 51(6), 954–966, 2020.
- [3] N. Gauthier, Explicit formula for power sums of an arithmetic sequence, *Math. Gaz.*, 91(520), 97–103, 2007.
- [4] H. W. Gould, Explicit formulas for Bernoulli numbers, *Amer. Math. Monthly*, 79(1), 44–51, 1972.
- [5] M. Griffiths, More on sums of powers of an arithmetic progression, *Math. Gaz.*, 93(527), 277–279, 2009.
- [6] B.-N. Guo, I. Mező, and F. Qi, An explicit formula for Bernoulli polynomials in terms of r -Stirling numbers of the second kind, *Rocky Mountain J. Math.*, 46(6), 1919–1923, 2016.
- [7] M. D. Hirschhorn, Evaluating $\sum_{n=1}^N (a+nd)^p$, *Math. Gaz.*, 90(517), 114–116, 2006.
- [8] I. Mező, A new formula for the Bernoulli polynomials, *Results Math.*, 58, 329–335, 2010.
- [9] M. Mihoubi and M. Tiachachat, Some applications of the r -Whitney numbers, *C. R. Acad. Sci. Paris, Ser. I*, 352(12), 965–969, 2014.
- [10] J. Quaintance and H. W. Gould, *Combinatorial Identities for Stirling Numbers. The Unpublished Notes of H. W. Gould*. Singapore: World Scientific Publishing, 2016.
- [11] J. L. Ramírez, S. N. Villamarín, D. Villamizar, Eulerian numbers associated with arithmetical progressions. *The Electronic Journal of Combinatorics*, 25, Article #P1.48, 2018.
- [12] H. M. Srivastava, Some explicit formulas for the Bernoulli and Euler numbers and polynomials, *Internat. J. Math. Ed. Sci. Tech.*, 19(1), 79–82, 1988.

- [13] P. G. Todorov, On the theory of the Bernoulli polynomials and numbers, *J. Math. Anal. Appl.*, 104(2), 309-350, 1984.
- [14] H. P. Tsao, Explicit polynomial expressions for sums of powers of an arithmetic progression, *Math. Gaz.*, 92(523), 87-92, 2008.

COLLADO VILLALBA, 28400 – MADRID, SPAIN
Email address: `j1.cereceda@movistar.es`