

TOGLIATTI SYSTEMS ASSOCIATED TO THE DIHEDRAL GROUP AND THE WEAK LEFSCHETZ PROPERTY.

LIENA COLARTE-GÓMEZ, EMILIA MEZZETTI, ROSA M. MIRÓ-ROIG, AND MARTÍ SALAT-MOLTÓ

ABSTRACT. In this note, we study Togliatti systems generated by invariants of the dihedral group D_{2d} acting on $k[x_0, x_1, x_2]$. This leads to the first family of non monomial Togliatti systems, which we call GT -systems with group D_{2d} . We study their associated varieties $S_{D_{2d}}$, called GT -surfaces with group D_{2d} . We prove that they are arithmetically Cohen-Macaulay surfaces whose homogeneous ideal, $I(S_{D_{2d}})$, is minimally generated by quadrics and we find a minimal free resolution of $I(S_{D_{2d}})$.

CONTENTS

1.	Introduction	1
2.	Preliminaries.	3
3.	The algebra of invariants of the dihedral group.	6
4.	Togliatti systems associated to the dihedral group.	11
	References	15

December 30, 2021

1. INTRODUCTION

Togliatti systems were introduced in [13], where the authors related the existence of homogeneous artinian ideals failing the weak Lefschetz property to the existence of projective varieties satisfying at least one Laplace equation. Precisely, a Togliatti system is an artinian ideal $I_d \subset k[x_0, \dots, x_n]$ generated by $r \leq \binom{n+d-1}{n-1}$ forms F_1, \dots, F_r of degree d which fails the weak Lefschetz property in degree $d-1$. The name is in honour of E. Togliatti who gave a complete classification of rational surfaces parameterized by cubics and satisfying at least one Laplace equation of order 2 (see [22] and [23]). Since then, this topic and related problems have been the focus of attention of many works, as one can see in [1], [2], [4], [5], [6], [11], [12], [14], [15] and [21]. Notwithstanding, most expositions and results deal with *monomial* Togliatti systems, while the non monomial case remains barely known.

Acknowledgments: The first and third authors are partially supported by MTM2016-78623-P. The last author is partially supported by MDM-2014-0445-18-2. The second author is supported by PRIN 2017SSNZAW and by FRA of the University of Trieste, and she is a member of INDAM-GNSAGA.

Recently, in [12] and [5] the authors studied GT -systems, a new family of monomial Togliatti systems having a special geometric property. A GT -system is a Togliatti system I_d whose associated morphism $\varphi_{I_d} : \mathbb{P}^n \rightarrow \mathbb{P}^{\mu_{I_d}-1}$ is a Galois covering with cyclic group $\mathbb{Z}/d\mathbb{Z}$. This geometric property establishes a new link between Togliatti systems and invariant theory. Precisely in [4] and [6], the authors apply invariant theory techniques to investigate both GT -systems and their images $X_d = \varphi_{I_d}(\mathbb{P}^n)$, the so called GT -varieties. These varieties are actually monomial projections of the Veronese variety $\nu_d(\mathbb{P}^n) \subset \mathbb{P}^{\binom{n+d}{d}-1}$ of \mathbb{P}^n from the linear space $\langle I_d^{-1} \rangle$ generated by the Macaulay's inverse system of I_d . Interest in these varieties relies on the following two problems. The first one is to determine whether X_d is an arithmetically Cohen-Macaulay (shortly aCM) monomial projection of $\nu_d(\mathbb{P}^n)$. This contributes to the longstanding problem, posed by Gröbner in [8], to determine when a projection of $\nu_d(\mathbb{P}^n)$ is aCM. The second one is the classical problem of finding a minimal free resolution of the homogeneous ideal of a projective variety. In this note, we extend the notion of GT -systems and GT -varieties, as presented in [4], to the action of any finite group, including non abelian ones. This sheds new light on the study of non monomial Togliatti systems. In this work, we focus on the dihedral group action on $k[x_0, x_1, x_2]$. We show that the invariant theory point of view used in [4], provides enough techniques to study the GT -system I_{2d} associated to the dihedral group and to tackle the geometry of the GT -surfaces $S_{D_{2d}}$ defined by the GT -system I_{2d} .

More precisely, we fix integers $3 \leq d, 0 < a < \frac{d}{2}$ with $\text{GCD}(d, a) = 1$ and ε a $2d$ th primitive root of 1. We set $e = \varepsilon^2$ and let $\rho_a : D_{2d} \rightarrow \text{GL}(3, k)$ be the linear representation of $D_{2d} = \langle \tau, \eta \mid \tau^d = \eta^2 = (\eta\tau)^2 = 1 \rangle$, the dihedral group of order $2d$, defined by:

$$\rho_a(\tau) = M_{d;a,d-a} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^a & 0 \\ 0 & 0 & e^{d-a} \end{pmatrix} \quad \text{and} \quad \rho_a(\eta) = \sigma = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Since $\text{GCD}(a, d) = 1$, the finite cyclic group $\langle \rho_a(\tau) \rangle$ of order d coincides with $\langle M_{d;1,d-1} \rangle \subset \text{GL}(3, k)$. We set C_{2d} the finite cyclic group of order $2d$ generated by εId and let $\overline{D_{2d}} = D_{2d} \times C_{2d} \subset \text{GL}(3, k)$ be the cyclic extension of D_{2d} . We denote by I_{2d} the ideal generated by all forms of degree $2d$ which are invariants of $\overline{D_{2d}}$. Our main goal is to relate the ideal I_{2d} to the ring $R^{\overline{D_{2d}}}$ of invariants of $\overline{D_{2d}}$. Using the structure of the ring $R^{D_{2d}}$, we determine a minimal set of μ_{2d} generators of I_{2d} , formed by monomials and binomials, and we prove that it is a minimal set of generators of $R^{\overline{D_{2d}}}$. This allows us to establish that the ideal I_{2d} is a GT -system with group D_{2d} , once proved in Lemma 4.1 that $\mu_{2d} \leq 2d + 1$ (see Theorem 2.1). As a consequence, we obtain that $R^{\overline{D_{2d}}}$ is the coordinate ring of the surface $S_{D_{2d}} = \varphi_{I_{2d}}(\mathbb{P}^2)$ associated to the GT -system I_{2d} . Through this connection, we prove that $S_{D_{2d}}$ is aCM and we compute a minimal free resolution of $R^{\overline{D_{2d}}}$. In particular, we show

that its homogeneous ideal $I(S_{D_{2d}})$ is minimally generated by quadrics and we determine a minimal set of generators.

Let us explain how this note is organized. We begin establishing in Section 2, all the preliminary results and definitions needed in the sequel. In particular, we define the extended notion of GT -system with respect to any finite group acting on $k[x_0, \dots, x_n]$. Section 3 is devoted to find a set of fundamental invariants of $\overline{D_{2d}}$. We prove that $R^{\overline{D_{2d}}}$ is minimally generated by monomials and homogeneous binomials of degree $2d$, which we completely determine (see Theorem 3.7) using the structure of the ring $R^{D_{2d}}$. We compute the Hilbert function, series and polynomial of $R^{\overline{D_{2d}}}$ and we establish that $R^{\overline{D_{2d}}}$ is a level algebra with Castelnuovo-Mumford regularity three. In Section 4, we introduce a new family of non monomial Togliatti systems, that we call GT -systems with group D_{2d} and we study their associated varieties, we call them GT -surfaces with group D_{2d} . We identify the coordinate ring of any GT -surface with group D_{2d} with the ring $R^{\overline{D_{2d}}}$. This allows us to translate geometrically the results obtained in Section 3. We show that any GT -surface with group D_{2d} is aCM. In addition, the information from the Hilbert series and the regularity allow us to compute a minimal free resolution of the homogeneous ideal $I(S_{D_{2d}})$ of any GT -surface $S_{D_{2d}}$ with group D_{2d} (see Theorem 4.6). In particular, we show that $I(S_{D_{2d}})$ is minimally generated by quadrics. Right after, we focus on determining a minimal set of generators.

Notation. Through this note k denotes an algebraically closed field of characteristic zero and $\text{GL}(n, k)$ denotes the group of invertible $n \times n$ matrices with coefficients in k .

Acknowledgments. The authors are very grateful to the anonymous referee of an earlier version of this work for providing insightful comments and directions which have substantially improved this paper.

2. PRELIMINARIES.

In this section, we collect the main concepts and tools we use in the body of this note. First, we relate the weak Lefschetz property of artinian ideals with varieties satisfying a Laplace equation and we recall the notion of Togliatti system introduced in [13]. Secondly, we see that quotient varieties by finite groups are Galois coverings and we extend the notion of GT -system from [12] to any finite group G . Finally, we review some basic facts on the theory of invariants of finite groups needed in the sequel.

Weak Lefschetz property. Set $R = k[x_0, \dots, x_n]$ the polynomial ring and let $I \subset R$ be a homogeneous artinian ideal. We say that I has the *weak Lefschetz property (WLP)* if there

is a linear form $L \in (R/I)_1$ such that, for all integers j , the multiplication map

$$\times L : (R/I)_{j-1} \rightarrow (R/I)_j$$

has maximal rank, i.e. it is injective or surjective. In [13] Mezzetti, Miró-Roig and Ottaviani proved that the failure of the WLP is related to the existence of varieties satisfying at least one Laplace equation of order greater than 2. More precisely, they proved:

Theorem 2.1. *Let $I \subset R$ be an artinian ideal generated by r forms F_1, \dots, F_r of degree d and let I^{-1} be its Macaulay inverse system. If $r \leq \binom{n+d-1}{n-1}$, then the following conditions are equivalent:*

- (i) I fails the WLP in degree $d - 1$;
- (ii) F_1, \dots, F_r become k -linearly dependent on a general hyperplane H of \mathbb{P}^n ;
- (iii) the n -dimensional variety $X = \overline{\text{Im}(\varphi)}$ where $\varphi : \mathbb{P}^n \dashrightarrow \mathbb{P}^{\binom{n+d}{d}-r-1}$ is the rational map associated to $(I^{-1})_d$, satisfies at least one Laplace equation of order $d - 1$.

Proof. See [13, Theorem 3.2]. □

In view of this result, a *Togliatti system* is defined as an artinian ideal $I \subset R$ generated by $r \leq \binom{n+d-1}{n-1}$ forms of degree d which fails the WLP in degree $d - 1$. This name is in honor of Togliatti who proved that the only smooth Togliatti system of cubics is

$$I = (x_0^3, x_1^3, x_2^3, x_0x_1x_2) \subset k[x_0, x_1, x_2]$$

(see [3], [22] and [23]). The systematic study of Togliatti systems was initiated in [13] and for recent results the reader can see [11], [14], [1], [15], [12], [4] and [5]. In this paper, we will restrict our attention to a particular case of Togliatti systems, the so called GT-systems which we are going to introduce now.

Galois coverings and GT-systems. Let us recall the notion of a Galois covering. A *covering* of a variety X consists of a variety Y and a finite morphism $f : Y \rightarrow X$. The *group of deck transformation* $G := \text{Aut}(f)$ is defined to be the group of automorphisms of Y commuting with f . We say that $f : Y \rightarrow X$ is a covering with group $\text{Aut}(f)$. If the fibre of a covering $f : Y \rightarrow X$ over a general point consists of d points we say that f is a covering of degree d .

Definition 2.2. A covering $f : Y \rightarrow X$ of a variety X is a Galois covering if the group $\text{Aut}(f)$ acts transitively on the fibre $f^{-1}(x)$ for some $x \in X$, and hence for all $x \in X$. We say that $f : Y \rightarrow X$ is a Galois covering with group $\text{Aut}(f)$.

Quotients varieties by finite groups of automorphisms work particularly well with respect to Galois coverings.

Proposition 2.3. *Let X be a projective variety and $G \subset \text{Aut}(X)$ be a finite group. If the quotient variety X/G exists, then $\pi : X \rightarrow X/G$ is a Galois covering.*

Proof. See [4, Proposition 2.3]. □

For further details on quotient varieties see for instance [16]. In [12] and [4], the authors studied a particular class of Togliatti systems arising from actions of the cyclic group over R . In particular, let $d \geq 3$ and $1 \leq a < b \leq d - 1$ be integers such that $\text{GCD}(a, b, d) = 1$. We denote by $M_{d;a,b}$ the matrix $\text{diag}(1, e^a, e^b) \in \text{GL}(3, k)$, where e is a d th root of 1. Then, let $\rho_{a,b} : C_d \rightarrow \text{GL}(3, k)$ be the representation of $C_d = \langle \tau \mid \tau^d = 1 \rangle$, the cyclic group of order d , given by $\rho_{a,b}(\tau) = M_{d;a,b} \in \text{GL}(3, k)$. With this notation, they proved:

Proposition 2.4. *Let $I_d \subset k[x_0, x_1, x_2]$ be the ideal generated by all forms of degree d which are invariants of $\rho_{a,b}(C_d)$. Let $\mu(I_d)$ denote the minimal number of generators of I_d . If $\mu(I_d) \leq d + 1$, then I_d is a Togliatti system. Moreover, the associated morphism $\varphi_{I_d} : \mathbb{P}^2 \rightarrow \mathbb{P}^{\mu(I_d)-1}$ is a Galois covering with cyclic group $\mathbb{Z}/d\mathbb{Z}$.*

Proof. See [4, Corollary 3.5]. □

The above result motivates the following definition.

Definition 2.5. *Let G be a finite group. We say that a Togliatti system $I = (F_1, \dots, F_r) \subset R$ is a GT-system with group G if the associated morphism $\varphi_I : \mathbb{P}^n \rightarrow \mathbb{P}^{r-1}$ is a Galois covering with group G .*

The study of the GT-systems with cyclic group $\mathbb{Z}/d\mathbb{Z}$ is presented in [4], [5], [6] and [12]. In all these papers, the group G is abelian and the GT-system is monomial. In this note we study GT-systems with non-abelian finite group G , more precisely with G the dihedral group, and we get the first examples of non-monomial GT-system.

Invariant theory of finite groups. A finite group of automorphisms of the affine space \mathbb{A}^{n+1} can be regarded as a finite group $G \subset \text{GL}(n+1, k)$ acting on the polynomial ring R . Let us denote by $R^G = \{f \in R \mid g(f) = f, \forall g \in G\}$ the ring of invariants of G . The ring R^G inherits the natural grading of R , that is $R^G = \bigoplus_{t \geq 0} R_t^G$, where $R_t^G := R_t \cap R^G$. We have the following result.

Lemma 2.6. *Fix $t \geq 1$ and let $G \subset \text{GL}(n+1, k)$ be a finite linear group acting on R . Then*

$$\dim_k R_t^G = \frac{1}{|G|} \sum_{g \in G} \text{trace}(g^{(t)})$$

where $g^{(t)}$ is the linear map induced by the action of g on R_t .

Proof. See [20, Lemma 2.2.2]. □

Geometrically, R^G is the coordinate ring of the quotient variety of \mathbb{A}^{n+1} by G . To be more precise, let $\{f_1, \dots, f_t\}$ be a minimal set of generators of the algebra R^G , often called a set of *fundamental invariants*, and let $k[w_1, \dots, w_t]$ be the polynomial ring in the new variables w_1, \dots, w_t . Then, the quotient variety of \mathbb{A}^{n+1} by G is given by the morphism $\pi : \mathbb{A}^{n+1} \rightarrow \pi(\mathbb{A}^{n+1}) \subset \mathbb{A}^t$, such that $\pi(a_0, \dots, a_n) = (f_1(a_0, \dots, a_n), \dots, f_t(a_0, \dots, a_n))$. The ideal $I(\pi(\mathbb{A}^{n+1}))$ of the quotient variety is called the *ideal of syzygies* among the invariants f_1, \dots, f_t ; it is the kernel of the homomorphism from R to $k[w_1, \dots, w_t]$ defined by $w_i \rightarrow f_i$, $i = 1, \dots, t$. We denote it by $\text{syz}(f_1, \dots, f_t)$. We have:

Proposition 2.7. *Let $G \subset \text{GL}(n+1, k)$ be a finite group acting on \mathbb{A}^{n+1} . Let f_1, \dots, f_t be a set of fundamental invariants and let $\pi : \mathbb{A}^{n+1} \rightarrow \mathbb{A}^t$ be the induced morphism. Then,*

- (i) $\pi(\mathbb{A}^{n+1})$ is the affine quotient variety by G with affine coordinate ring R^G .
- (ii) $R^G \cong k[w_1, \dots, w_t] / \text{syz}(f_1, \dots, f_t)$.
- (iii) π is a Galois covering of $\pi(\mathbb{A}^{n+1})$ with group G . The cardinality of a general orbit $G(a)$, $a \in \mathbb{A}^{n+1}$, is called the degree of the covering.

Proof. See [19, Section 6] and Proposition 2.3. □

If we can find a homogeneous set of fundamental invariants $\{f_1, \dots, f_t\}$ such that $\pi : \mathbb{P}^n \rightarrow \mathbb{P}^{t-1}$ is a morphism, then the projective version of 2.7 is true.

3. THE ALGEBRA OF INVARIANTS OF THE DIHEDRAL GROUP.

Through this section, we study the action of the dihedral group on the polynomial ring $R = k[x_0, x_1, x_2]$. We fix integers $3 \leq d$, $0 < a < \frac{d}{2}$ with $\text{GCD}(d, a) = 1$ and ε a $2d$ th primitive root of 1. We set $e = \varepsilon^2$ and let $\rho_a : D_{2d} \rightarrow \text{GL}(3, k)$ be the linear representation of $D_{2d} = \langle \tau, \epsilon \mid \tau^d = \eta^2 = (\eta\tau)^2 = 1 \rangle$, the dihedral group of order $2d$, defined by (see [17]):

$$\rho_a(\tau) = M_{d;a,d-a} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^a & 0 \\ 0 & 0 & e^{d-a} \end{pmatrix} \quad \text{and} \quad \rho_a(\eta) = \sigma = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

It is the direct sum of the trivial representation in $\text{GL}(1, k)$ and a faithful representation in $\text{GL}(2, k)$ of D_{2d} . Therefore, the ring $R^{D_{2d}}$ of invariants is generated by the three algebraically independent invariants x_0 , x_1x_2 and $x_1^d + x_2^d$ of D_{2d} (see [18] and [19]). Thus $R^{D_{2d}} = k[x_0, x_1x_2, x_1^d + x_2^d] = \bigoplus_{t \geq 0} R_t \cap R^{D_{2d}}$ is a non-standard graded polynomial ring.

Since $\text{GCD}(a, d) = 1$, the finite cyclic group $\langle \rho_a(\tau) \rangle$ of order d coincides with $\Gamma := \langle M_{d;1,d-1} \rangle \subset \text{GL}(3, k)$. We set C_{2d} the finite cyclic group of order $2d$ generated by εId and we define $\bar{\Gamma} \subset \text{GL}(3, k)$ to be the cyclic extension of Γ , i.e. $\bar{\Gamma} = \Gamma \times C_{2d}$. Similarly, let $\overline{D_{2d}} = D_{2d} \times C_{2d} \subset \text{GL}(3, k)$ be the cyclic extension of D_{2d} . We see the ring of invariants

$R^{\overline{D_{2d}}}$ of $\overline{D_{2d}}$ as a k -graded subalgebra of R and $R^{D_{2d}}$ as follows:

$$R^{\overline{D_{2d}}} = \bigoplus_{t \geq 0} R_t^{\overline{D_{2d}}}, \quad \text{where} \quad R_t^{\overline{D_{2d}}} := R_{2dt}^{D_{2d}} = R_{2dt} \cap R^{D_{2d}}.$$

We relate $R^{\overline{D_{2d}}}$ to the ring R^Γ studied in [4] and this connection allows to compute the Hilbert function, Hilbert polynomial and Hilbert series of $R^{\overline{D_{2d}}}$. We also provide a complete description of a homogeneous k -basis for each $R_t^{\overline{D_{2d}}}$, $t \geq 1$. Our main result shows that the graded k -algebra $R^{\overline{D_{2d}}}$ is generated in degree 1. Let us begin with the following remarks.

Remark 3.1. The action of σ on a monomial $x_0^{a_0} x_1^{a_1} x_2^{a_2}$ is given by $x_0^{a_0} x_1^{a_2} x_2^{a_1}$. Therefore, the action of $\langle M_{d,1,d-1}^l \sigma \rangle \subset \text{GL}(3, k)$ is the same as the action of $\langle \text{diag}(1, e^{d-1}, e)^l \rangle \subset \text{GL}(3, k)$ for any $0 \leq l \leq d-1$.

Remark 3.2. (i) If a monomial of degree $2dt$ is an invariant of Γ , it is also an invariant of $\langle \text{diag}(1, e^{d-1}, e) \rangle \subset \text{GL}(3, k)$. Actually, a monomial $x_0^{a_0} x_1^{a_1} x_2^{a_2}$ of degree $2dt$ is an invariant of Γ if and only if there exists $r \in \{0, \dots, 2t(d-1)\}$ such that $(a_0, a_1, a_2) \in \mathbb{Z}_{\geq 0}^3$ is a solution of the integer system:

$$\begin{cases} a_0 + a_1 + a_2 = 2dt \\ a_1 + (d-1)a_2 = rd. \end{cases}$$

Now $0 < (d-1)a_1 + a_2 = da_1 - (a_1 - a_2) = d(a_1 + a_2) - rd$ is also a multiple of d . Hence (a_0, a_1, a_2) is also a solution of the system

$$\begin{cases} a_0 + a_1 + a_2 = 2dt \\ (d-1)a_1 + a_2 = (a_1 + a_2 - r)d \end{cases}$$

which implies that $x_0^{a_0} x_1^{a_1} x_2^{a_2}$ is an invariant of $\langle \text{diag}(1, e^{d-1}, e) \rangle$.

(ii) Any monomial of degree $2dt$ of the form $x_0^{2dt-2a_1} x_1^{a_1} x_2^{a_1}$, $a_1 = 0, \dots, td$, is an invariant of Γ . There are exactly $td + 1$ monomials of degree td of such form.

Next, we compute the Hilbert function $\text{HF}(R^{\overline{D_{2d}}}, t)$ of $R^{\overline{D_{2d}}}$. Fix $t \geq 1$. Since $\text{HF}(R^{\overline{D_{2d}}}, t)$ is equal to the Hilbert function $\text{HF}(R^{D_{2d}}, 2dt)$ of $R^{D_{2d}}$, by Lemma 2.6 we have

$$\text{HF}(R^{\overline{D_{2d}}}, t) = \frac{1}{2d} \sum_{g \in \rho(D_{2d})} \text{trace}(g^{(2dt)}) = \frac{1}{2d} \text{trace}\left(\sum_{g \in \rho(D_{2d})} g^{(2dt)} \right),$$

where $g^{(2dt)}$ is the restriction of g to R_{2dt} . We choose the set of all monomials of degree $2dt$ as a basis \mathcal{B} of R_{2dt} , namely $\mathcal{B} = \{m_1, \dots, m_N\}$, where $N = \dim_k R_{2dt} = \binom{2dt+2}{2}$.

Proposition 3.3. *With the above notation,*

$$\text{HF}(R^{\overline{D_{2d}}}, t) = \frac{2dt^2 + (d + \text{GCD}(d, 2) + 2)t + 2}{2}.$$

Proof. Let m_i be a monomial in \mathcal{B} of degree $2dt$. We denote by M the matrix which represents the linear map $\sum_{g \in \rho(D_{2d})} g^{(2dt)}$ in the above basis. We distinguish two cases.

Case 1: $m_i \in R^\Gamma$. Then by Remark 3.2,

$$M_{(i,i)} = \begin{cases} 2d & \text{if } \sigma(m_i) = m_i \\ d & \text{if } \sigma(m_i) \neq m_i \end{cases}$$

Case 2: $m_i \notin R^\Gamma$. If ξ is a d th root of unity, we have the equality $1 + \xi + \dots + \xi^{d-1} = 0$. This, in addition to Remark 3.2, gives $M_{(i,i)} = 0$.

Let μ_{2dt}^c be the number of monomials of degree $2dt$ in R^Γ . Thus, we have obtained that

$$(2d) \text{HF}(R^{\overline{D_{2d}}}, t) = d(\mu_{2dt}^c + td + 1).$$

By [4, Theorem 4.5], μ_{2dt}^c is equal to the Hilbert function $\text{HF}(R^\Gamma, 2t)$ of the ring R^Γ in degree $2t$. By [4, Theorem 4.11] $\mu_{2dt}^c = 2dt^2 + 2t + \text{GCD}(2, d)t + 1$, which completes the proof. \square

From the above result, we directly obtain the Hilbert polynomial $\text{HP}(R^{\overline{D_{2d}}}, t)$ and the Hilbert series $\text{HS}(R^{\overline{D_{2d}}}, z)$ of the ring $R^{\overline{D_{2d}}}$.

Proposition 3.4. *With the above notation, the Hilbert polynomial and the Hilbert series of $R^{\overline{D_{2d}}}$ are given by the following expressions:*

- (i) $\text{HP}(R^{\overline{D_{2d}}}, t) = \frac{1}{2}(2dt^2 + (d + \text{GCD}(d, 2) + 2)t + 2)$
- (ii) $\text{HS}(R^{\overline{D_{2d}}}, z) = \frac{1}{(1-z)^3} \left(\frac{d - \text{GCD}(d, 2)}{2} z^2 + \frac{3d + \text{GCD}(d, 2) - 2}{2} z + 1 \right).$

Next, we use the relation between the two rings $R^{\overline{D_{2d}}}$ and R^Γ to determine a k -basis of any vector space $R_t^{\overline{D_{2d}}}$, $t \geq 1$, and to find a k -algebra basis of $R^{\overline{D_{2d}}}$. Fix $t \geq 1$ and consider $R_t^{\overline{D_{2d}}}$.

Proposition 3.5. *A k -basis \mathcal{B}_{2dt} of the vector space $R_t^{\overline{D_{2d}}}$ is formed by:*

- (i) *the set of $td + 1$ monomial invariants $x_0^{2dt}, x_0^{2dt-2}x_1x_2, x_0^{2dt-4}x_1^2x_2^2, \dots, x_1^{td}x_2^{td}$ of Γ of degree $2dt$; and*
- (ii) *the set of all binomials $x_0^{a_0}x_1^{a_1}x_2^{a_2} + x_0^{a_0}x_1^{a_2}x_2^{a_1}$ of degree $2dt$ such that $a_1 \neq a_2$ and $x_0^{a_0}x_1^{a_1}x_2^{a_2} \in R^\Gamma$.*

Proof. By Remark 3.2(i), there are exactly $\frac{1}{2}(\mu_{2dt}^c - td - 1)$ binomials of the form $x_0^{a_0}x_1^{a_1}x_2^{a_2} + x_0^{a_0}x_1^{a_2}x_2^{a_1} \in R^\Gamma$ of degree $2dt$ with $a_1 \neq a_2$. Since the forms in (i) and (ii) are k -linearly independent, the result follows from Proposition 3.3. \square

We illustrate the above result with a couple of examples.

Example 3.6. (i) For $d = 3$ and $D_{2,3} = \langle M_{3;1,2}, \sigma \rangle$, we have:

$$\mathcal{B}_{2,3} = \{x_0^6, x_0^3x_1^3 + x_0^3x_2^3, x_0^4x_1x_2, x_1^6 + x_2^6, x_0x_1^4x_2 + x_0x_1x_2^4, x_0^2x_1^2x_2^2, x_1^3x_2^3\}, \text{HF}(R^{\overline{D_{2,3}}}, 1) = 7.$$

$$\mathcal{B}_{4,3} = \{x_0^{12}, x_0^9x_1^3 + x_0^9x_2^3, x_0^{10}x_1x_2, x_0^6x_1^6 + x_0^6x_2^6, x_0^7x_1^4x_2 + x_0^7x_1x_2^4, x_0^8x_1^2x_2^2, x_0^3x_1^9 + x_0^3x_2^9, x_0^4x_1^7x_2 + x_0^4x_1x_2^7, x_0^5x_1^5x_2^2 + x_0^5x_1^2x_2^5, x_0^6x_1^3x_2^3, x_1^{12} + x_2^{12}, x_0x_1^{10}x_2 + x_0x_1x_2^{10}, x_0^2x_1^8x_2^2 + x_0^2x_1^2x_2^8, x_0^3x_1^6x_2^3 + x_0^3x_1^3x_2^6, x_0^4x_1^4x_2^4, x_0^9x_1^3 + x_0^9x_2^3, x_0x_1^7x_2^4 + x_0x_1^4x_2^7, x_0^2x_1^5x_2^5, x_1^6x_2^6\}, \text{HF}(R^{\overline{D_{2,3}}}, 2) = 19.$$

(ii) For $d = 4$ and $D_{2,4} = \langle M_{4;1,3}, \sigma \rangle$, we have:

$$\mathcal{B}_{2,4} = \{x_0^8, x_0^4x_1^4 + x_0^4x_2^4, x_0^6x_1x_2, x_1^8 + x_2^8, x_0^2x_1^5x_2 + x_0^2x_1x_2^5, x_0^4x_1^2x_2^2, x_1^6x_2^2 + x_1^2x_2^6, x_0^2x_1^3x_2^3, x_1^4x_2^4\}, \text{HF}(R^{\overline{D_{2,4}}}, 1) = 9.$$

$$\mathcal{B}_{4,4} = \{x_0^{16}, x_0^{12}x_1^4 + x_0^{12}x_2^4, x_0^{14}x_1x_2, x_0^8x_1^8 + x_0^8x_2^8, x_0^{10}x_1^5x_2 + x_0^{10}x_1x_2^5, x_0^{12}x_1^2x_2^2, x_0^4x_1^{12} + x_0^4x_2^{12}, x_0^6x_1^9x_2 + x_0^6x_1x_2^9, x_0^8x_1^6x_2^2 + x_0^8x_1^2x_2^6, x_0^{10}x_1^3x_2^3, x_1^{16} + x_2^{16}, x_0^2x_1^{13}x_2 + x_0^2x_1x_2^{13}, x_0^4x_1^{10}x_2^2 + x_0^4x_1^2x_2^{10}, x_0^6x_1^7x_2^3 + x_0^6x_1^3x_2^7, x_0^8x_1^4x_2^4, x_1^{14}x_2^2 + x_1^2x_2^{14}, x_0^2x_1^{11}x_2^3 + x_0^2x_1^3x_2^{11}, x_0^4x_1^8x_2^4 + x_0^4x_1^4x_2^8, x_0^6x_1^5x_2^5, x_1^{12}x_2^4 + x_1^4x_2^{12}, x_0^2x_1^9x_2^5 + x_0^2x_1^5x_2^9, x_0^4x_1^6x_2^6, x_1^{10}x_2^6 + x_1^6x_2^{10}, x_0^2x_1^7x_2^7, x_1^8x_2^8\}, \text{HF}(R^{\overline{D_{2,4}}}, 2) = 25.$$

The next goal is to prove that \mathcal{B}_{2d} is a set of fundamental invariants of $\overline{D_{2d}}$. To achieve it, we use the natural structure of $R^{\overline{D_{2d}}}$ as a subring of $R^{D_{2d}}$. We set $y_0 = x_0$, $y_1 = x_1x_2$ and $y_2 = x_1^d + x_2^d$, as we have seen $R^{D_{2d}} = k[y_0, y_1, y_2]$ with $\deg(y_0) = 1$, $\deg(y_1) = 2$ and $\deg(y_2) = d$. With this notation, for any $t \geq 1$ we have that $R_t^{D_{2d}} = k[y_0, y_1, y_2]_{2td}$ is the k -vector space with monomial basis $\mathcal{A}_{2dt} = \{y_0^{b_0}y_1^{b_1}y_2^{b_2} \mid b_0 + 2b_1 + db_2 = 2td\}$. In particular, for $t = 1$ we have the change of basis $\rho : k[y_0, y_1, y_2]_{2d} \rightarrow R_1^{\overline{D_{2d}}}$

$$(1) \quad \begin{cases} y_0^{b_0}y_1^{b_1}y_2^{b_2} & \mapsto x_0^{b_0}x_1^{b_1}x_2^{b_1}(x_1^d + x_2^d)^{b_2}, & \text{if } 0 \leq b_2 \leq 1 \\ y_2^2 & \mapsto (x_1^{2d} + x_2^{2d}) + 2x_1^dx_2^d. \end{cases}$$

Theorem 3.7. \mathcal{B}_{2d} is a set of fundamental invariants of $\overline{D_{2d}}$.

Proof. We see that for any $t \geq 2$, any monomial $y_0^{b_0}y_1^{b_1}y_2^{b_2} \in \mathcal{A}_{2dt}$ is divisible by a monomial of \mathcal{A}_{2d} . Then by induction, it follows that \mathcal{A}_{2d} is a set of generators of $R^{\overline{D_{2d}}} \subset R^{D_{2d}}$. Using (1), we obtain that \mathcal{B}_{2d} is a minimal set of generators of $R^{\overline{D_{2d}}}$.

Let $m = y_0^{b_0}y_1^{b_1}y_2^{b_2} \in \mathcal{A}_{2dt}$ be a monomial of degree $b_0 + 2b_1 + db_2 = 2dt$, $t \geq 2$. On one hand, we may suppose that $b_0 < 2d$, $b_1 < d$ and $b_2 < 2$. Otherwise, y_0^{2d} , y_1^d or y_2^2 divide m and the result follows. On the other hand, if $b_2 = 0$, $b_0 < 2d$ and $b_1 < d$, then we have $\deg(m) = 2d$ and $t = 1$. Therefore it only remains to prove the case $b_0 < 2d$, $b_1 < d$ and $b_2 = 1$ with $b_0 + 2b_1 + d = 4d$. Since $b_0 + 2b_1 = 3d$ and $b_1 < d$, this implies that $b_0 \geq d$ and then $y_0^dy_2 \in \mathcal{A}_{2d}$ divides m , as required. \square

As a consequence, $R^{\overline{D_{2d}}} \subset R$ is a graded k -algebra generated in degree 1 and ρ induces an isomorphism of graded k -algebras $\rho : k[\mathcal{A}_{2d}] \rightarrow k[\mathcal{B}_{2d}]$. The following example illustrates Theorem 3.7.

Example 3.8. We express the invariants of $\mathcal{B}_{4,3}$ in terms of \mathcal{B}_{2d} (see Example 3.6(i)). We have

$$\begin{aligned}\mathcal{A}_{2,3} &= \{y_0^6, y_0^3 y_2, y_0^4 y_1, y_2^2, y_0 y_1 y_2, y_0^2 y_1^2, y_1^3\} \\ \mathcal{A}_{4,3} &= \{y_0^{12}, y_0^{10} y_1, y_0^8 y_1^2, y_0^6 y_1^3, y_0^4 y_1^4, y_0^2 y_1^5, y_1^6, y_0^9 y_2, y_0^7 y_1 y_2, y_0^5 y_1^2 y_2, y_0^3 y_1^3 y_2, y_0 y_1^4 y_2, y_0^6 y_2^2, y_0^4 y_1 y_2^2, \\ &\quad y_0^2 y_1^2 y_2^2, y_1^3 y_2^2, y_0^3 y_2^3, y_0 y_1 y_2^3, y_2^4\}.\end{aligned}$$

Expressing all monomials of $\mathcal{A}_{4,3}$ as products of monomials of $\mathcal{A}_{2,3}$ and applying (1) we obtain the following factorizations:

$$\begin{aligned}x_0^{12} &= (x_0^6)(x_0^6) \\ x_0^{10} x_1 x_2 &= (x_0^6)(x_0^4 x_1 x_2) \\ x_0^8 x_1^2 x_2^2 &= (x_0^6)(x_0^2 x_1^2 x_2^2) \\ x_0^6 x_1^3 x_2^3 &= (x_0^6)(x_1^3 x_2^3) \\ x_0^4 x_1^4 x_2^4 &= (x_0^4 x_1 x_2)(x_1^3 x_2^3) \\ x_0^2 x_1^5 x_2^5 &= (x_0^2 x_1^2 x_2^2)(x_1^3 x_2^3) \\ x_1^6 x_2^6 &= (x_1^3 x_2^3)^2 \\ x_0^9 x_1^3 + x_0^9 x_2^3 &= x_0^6 (x_0^3 x_1^3 + x_0^3 x_2^3) \\ x_0^6 x_1^6 + x_0^6 x_2^6 &= x_0^6 (x_1^6 + x_2^6) \\ x_0^7 x_1^4 x_2 + x_0^7 x_1 x_2^4 &= x_0^4 x_1 x_2 (x_0^3 x_1^3 + x_0^3 x_2^3) \\ x_0^4 x_1^7 x_2 + x_0^4 x_1 x_2^7 &= x_0^4 x_1 x_2 (x_1^6 + x_2^6) \\ x_0^5 x_1^5 x_2^2 + x_0^5 x_1^2 x_2^5 &= x_0^2 x_1^2 x_2^2 (x_0^3 x_1^3 + x_0^3 x_2^3) \\ x_0^2 x_1^8 x_2^2 + x_0^2 x_1^2 x_2^8 &= x_0^2 x_1^2 x_2^2 (x_1^6 + x_2^6) \\ x_0^3 x_1^6 x_2^3 + x_0^3 x_1^3 x_2^6 &= x_1^3 x_2^3 (x_0^3 x_1^3 + x_0^3 x_2^3) \\ x_1^9 x_2^3 + x_1^3 x_2^9 &= x_1^3 x_2^3 (x_1^6 + x_2^6) \\ x_0 x_1^7 x_2^4 + x_0 x_1^4 x_2^7 &= x_1^3 x_2^3 (x_0 x_1^4 x_2 + x_0 x_1 x_2^4) \\ x_0^3 x_1^9 + x_0^3 x_2^9 &= (x_1^6 + x_2^6)(x_0^3 x_1^3 + x_0^3 x_2^3) - x_1^3 x_2^3 (x_0^3 x_1^3 + x_0^3 x_2^3) \\ x_1^{12} + x_2^{12} &= (x_1^6 + x_2^6)^2 - 2(x_1^3 x_2^3)^2 \\ x_0 x_1^{10} x_2 + x_0 x_1 x_2^{10} &= (x_0 x_1^4 x_2 + x_0 x_1 x_2^4)(x_1^6 + x_2^6) - x_1^3 x_2^3 (x_0 x_1^4 x_2 + x_0 x_1 x_2^4).\end{aligned}$$

Notice that these decompositions are not unique, for instance $x_0^8 x_1^2 x_2^2$ can also be factored as $(x_0^4 x_1 x_2)^2$.

We end this section with a corollary regarding the Cohen-Macaulayness of $R^{\overline{D_{2d}}}$.

Corollary 3.9. (i) $R^{\overline{D_{2d}}}$ is a Cohen-Macaulay level algebra with Cohen-Macaulay type $\frac{1}{2}(d - \text{GCD}(d, 2))$ and Castelnuovo-Mumford regularity 3.
(ii) $R^{\overline{D_{2d}}}$ is Gorenstein if and only if $d = 3$ or 4.

Proof. Since $R^{\overline{D_{2d}}}$ is the ring of invariants by the action of the linear finite group $\overline{D_{2d}}$ on R , it is Cohen-Macaulay (see [10, Proposition 12]). The other results follow from that and Proposition 3.4. \square

4. TOGLIATTI SYSTEMS ASSOCIATED TO THE DIHEDRAL GROUP.

In this section, we describe Togliatti systems I_{2d} associated to D_{2d} and we study the geometry of their associated varieties. As far as we know, the GT -systems described in previous works were all monomial and it is worthwhile to point out that I_{2d} is the first large class of non monomial GT -systems. Namely, we prove that the ideal generated by the set of fundamental invariants \mathcal{B}_{2d} (see Proposition 3.5 and Theorem 3.7) of $\overline{D_{2d}}$ is a GT -system with group D_{2d} . We connect the ring $R^{\overline{D_{2d}}}$ to the coordinate ring of the associated varieties of these GT -systems.

Let $I_{2d} \subset R$ be the homogeneous ideal generated by \mathcal{B}_{2d} and we set $\mu_{2d} := \text{HF}(R^{\overline{D_{2d}}}, 1)$. We denote by $\varphi_{I_{2d}} : \mathbb{P}^2 \rightarrow \mathbb{P}^{\mu_{2d}-1}$ the morphism induced by I_{2d} . Our first goal is to show that I_{2d} is a non monomial GT -system with group D_{2d} . We set $S_{D_{2d}} := \varphi_{I_{2d}}(\mathbb{P}^2)$ and we call it a GT -surface with group D_{2d} . The ring of invariants $R^{\overline{D_{2d}}}$ is then the coordinate ring of $S_{D_{2d}}$ (see Proposition 2.7 and Theorem 3.7). We study the homogeneous ideal $I(S_{D_{2d}})$ of $S_{D_{2d}}$, which is the prime ideal $\text{syz}(\mathcal{B}_{2d})$ of syzygies among \mathcal{B}_{2d} . We compute a minimal free resolution of $I(S_{D_{2d}})$ and, as a consequence, we prove that $I(S_{D_{2d}})$ is minimally generated by quadrics.

Lemma 4.1. *For all $d \geq 3$, $\mu_{2d} \leq 2d + 1$.*

Proof. From Proposition 3.3,

$$\mu_{2d} = \frac{2d + d + 2 + \text{GCD}(d, 2) + 2}{2} \leq \frac{3d + 6}{2},$$

which is smaller or equal than $2d + 1$ for all $d \geq 3$. \square

Proposition 4.2. *The ideal I_{2d} is a GT -system with group D_{2d} .*

Proof. Firstly, notice that I_{2d} is an artinian non-monomial ideal. Precisely, $x_0^{2d}, x_1^{2d} + x_2^{2d}$ and $x_1^d x_2^d$ form a homogeneous system of parameters of $R^{\overline{D_{2d}}}$. And secondly, since \mathcal{B}_{2d} is a set of fundamental invariants of $\overline{D_{2d}}$ (see Theorem 3.7), by Proposition 2.7 it follows that $\varphi_{I_{2d}}$ is a Galois covering with group D_{2d} . Now, by Lemma 4.1, I_{2d} is generated by $\mu_{2d} \leq 2d + 1$ homogeneous forms of degree $2d$. Hence, by Theorem 2.1 it only remains to see that it fails the weak Lefschetz property from degree $2d - 1$ to $2d$. Let $L \in (R/I_{2d})_1$ be a linear form, and let us consider $F_{2d-1} := \prod_{\substack{g \in D_{2d} \\ g \neq \text{Id}}} g(L)$. By construction, for any element $g \in D_{2d}$ we have that $g(LF_{2d-1}) = LF_{2d-1}$. Thus LF_{2d-1} is an invariant of $\overline{D_{2d}}$ and the result follows from Theorem 3.7. \square

In the rest of this section, we deal with the geometry of $S_{D_{2d}}$. Let us begin with some properties which follow directly from the results we obtained in Section 3.

Proposition 4.3. $S_{D_{2d}}$ is an arithmetically Cohen-Macaulay surface of degree $\deg(S_{D_{2d}}) = 2d$, regularity 3, codimension $\frac{1}{2}(3d + \text{GCD}(d, 2) - 2)$ and Cohen-Macaulay type $\frac{1}{2}(d - \text{GCD}(d, 2))$. In particular, $S_{D_{2d}}$ is Gorenstein if and only if $d = 3, 4$.

Proof. See Corollary 3.9. □

Our next goal is to determine a minimal free resolution of $I(S_{D_{2d}})$. In particular, we obtain that $I(S_{D_{2d}})$ is generated by quadrics. Let us begin introducing some new notation.

We set $\mathcal{W}_d := \{w_{(r,\gamma)} \mid 0 \leq r \leq 2(d-1) \text{ and } \max\{0, \lceil \frac{(r-2)d}{d-2} \rceil\} \leq \gamma \leq r\}$, a set of variables ordered lexicographically. As we will explicitly see in Notation 4.5, each pair (r, γ) as in \mathcal{W}_d uniquely determines the exponents of an element in \mathcal{B}_{2d} (see Remark 3.2 and Proposition 3.5). Hence, the cardinality of \mathcal{W}_d is $\mu_{2d} = d + 2 + \frac{d + \text{GCD}(2,d)}{2}$. We exhibit a few examples.

Example 4.4. (i) For $d = 3$, $\mathcal{B}_{2,3} = \{x_0^6, x_0^3x_1^3 + x_0^3x_2^3, x_0^4x_1x_2, x_1^6 + x_2^6, x_0x_1^4x_2 + x_0x_1x_2^4, x_0^2x_1^2x_2^2, x_1^3x_2^3\}$ and

$$\mathcal{W}_3 = \{w_{(0,0)}, w_{(1,0)}, w_{(1,1)}, w_{(2,0)}, w_{(2,1)}, w_{(2,2)}, w_{(3,3)}\}$$

(ii) For $d = 4$, $\mathcal{B}_{2,4} = \{x_0^8, x_0^4x_1^4 + x_0^4x_2^4, x_0^6x_1x_2, x_1^8 + x_2^8, x_0^2x_1^5x_2 + x_0^2x_1x_2^5, x_0^4x_1^2x_2^2, x_1^6x_2^2 + x_1^2x_2^6, x_0^2x_1^3x_2^3, x_1^4x_2^4\}$ and

$$\mathcal{W}_4 = \{w_{(0,0)}, w_{(1,0)}, w_{(1,1)}, w_{(2,0)}, w_{(2,1)}, w_{(2,2)}, w_{(3,2)}, w_{(3,3)}, w_{(4,4)}\}.$$

(iii) For $d = 5$, $\mathcal{B}_{2,5} = \{x_0^{10}, x_0^5x_1^5 + x_0^5x_2^5, x_0^8x_1x_2, x_1^{10} + x_2^{10}, x_0^3x_1^6x_2 + x_0^3x_1x_2^6, x_0^6x_1^2x_2^2, x_0x_1^7x_2^2 + x_0x_1^2x_2^7, x_0^4x_1^3x_2^3, x_0^2x_1^4x_2^4, x_1^5x_2^5\}$ and

$$\mathcal{W}_5 = \{w_{(0,0)}, w_{(1,0)}, w_{(1,1)}, w_{(2,0)}, w_{(2,1)}, w_{(2,2)}, w_{(3,2)}, w_{(3,3)}, w_{(4,4)}, w_{(5,5)}\}.$$

Notation 4.5. We denote by $S = k[w_{(r,\gamma)}]_{w_{(r,\gamma)} \in \mathcal{W}_d}$ the polynomial ring. The homogeneous ideal $I(S_{D_{2d}})$ is the kernel of the ring homomorphism $\varphi_d : S \rightarrow k[\mathcal{B}_{2d}]$ defined as follows:

$$\varphi_d(w_{(r,\gamma)}) = \begin{cases} x_0^{2d-2\gamma} x_1^\gamma x_2^\gamma =: m_{(r,\gamma)} & \text{if } r = \gamma \\ x_0^{(2-r)d+(d-2)\gamma} (x_1^{rd-(d-1)\gamma} x_2^\gamma + x_1^\gamma x_2^{rd-(d-1)\gamma}) =: m_{(r,\gamma)} + \overline{m_{(r,\gamma)}} & \text{otherwise.} \end{cases}$$

The information from the Hilbert function of $S_{D_{2d}}$ and the regularity allows us to determine a minimal free resolution of $S_{D_{2d}}$. Set

$$C := \text{codim}(S_{D_{2d}}) = \frac{3d + \text{GCD}(d, 2) - 2}{2}, \text{ and}$$

$$h := \text{deg}(S_{D_{2d}}) - C - 2 = \frac{d - \text{GCD}(d, 2) - 2}{2}.$$

Theorem 4.6. *With the above notation, $I(S_{D_{2d}})$ has a minimal free S -resolution*

$$\begin{aligned} 0 \rightarrow S^{bc,2}(-C-2) \rightarrow \bigoplus_{l=1,2} S^{bc-1,l}(-C+1-l) \rightarrow \bigoplus_{l=1,2} S^{bc-2,l}(-C+2-l) \\ \rightarrow \cdots \rightarrow \bigoplus_{l=1,2} S^{bc-h,l}(-C+h-l) \rightarrow S^{bc-h-1,1}(-C+h) \\ \rightarrow \cdots \rightarrow S^{b_{1,1}}(-2) \rightarrow S \rightarrow S/I(S_{2d}) \rightarrow 0 \end{aligned}$$

where

$$b_{i,j-i} := \begin{cases} i \binom{C}{i+1} + (C-i-h) \binom{C}{i-1} & \text{if } 1 \leq i \leq C-h-1, j = i+1 \\ i \binom{r}{i+1} & \text{if } C-h \leq i \leq C, j = i+1 \\ (i-C+h+1) \binom{C}{i} & \text{if } C-h \leq i \leq C, j = i+2 \\ 0 & \text{otherwise} \end{cases}$$

Proof. For $d = 3, 4$ we explicitly compute the resolutions of $S_{D_{2,3}}$ and $S_{D_{2,4}}$ in Example 4.8(i),(ii). For all $d \geq 5$ we check that $C+3 \leq 2d \leq 2C$ and then we apply [24, Corollary 3.4(ii)]. Clearly $2d \leq 3d + \text{GCD}(d, 2) - 2$ for all $d \geq 3$. On the other hand,

$$C+3 = \frac{3d + \text{GCD}(d, 2) + 4}{2} \leq 2d$$

if and only if $3d + \text{GCD}(d, 2) + 4 \leq 4d$ if and only if $\text{GCD}(d, 2) + 4 \leq d$. The last inequality holds for all $d \geq 5$. \square

Corollary 4.7. *$I(S_{D_{2d}})$ is minimally generated by $\frac{9d^2+2d+8}{8}$ quadrics if d is even and by $\frac{9d^2-4d+3}{8}$ quadrics if d is odd.*

Let us illustrate Theorem 4.6 with some examples, which we compute using the software Macaulay2 ([7]).

Example 4.8. (i) For $d = 3$, $S_{D_{2d}}$ has codimension $C = 4$ and degree $\deg(S_{D_{2d}}) = 6$, so $h = 0$. A minimal free resolution of $I(S_{D_{2,3}})$ over $S = k[w_{(r,\gamma)}]_{w_{(r,\gamma)} \in \mathcal{W}_3}$ is

$$0 \rightarrow S(-6) \rightarrow S^9(-4) \rightarrow S^{16}(-3) \rightarrow S(-2)^9 \rightarrow S \rightarrow S/I(S_{D_{2,3}}) \rightarrow 0.$$

(ii) For $d = 4$, $S_{D_{2d}}$ has codimension $C = 6$ and degree $\deg(S_{D_{2d}}) = 8$, so $h = 0$. A minimal free resolution of $I(S_{D_{2,4}})$ over $S = k[w_{(r,\gamma)}]_{w_{(r,\gamma)} \in \mathcal{W}_4}$ is

$$\begin{aligned} 0 \rightarrow S(-8) \rightarrow S^{20}(-6) \rightarrow S^{64}(-5) \rightarrow S^{90}(-4) \rightarrow \\ S^{64}(-3) \rightarrow S^{20}(-2) \rightarrow S \rightarrow S/I(S_{D_{2,4}}) \rightarrow 0. \end{aligned}$$

(iii) For $d = 5$, $S_{D_{2d}}$ has codimension $C = 7$ and degree $\deg(S_{D_{2d}}) = 10$, so we have $h = 1$ and a minimal free resolution of $I(S_{D_{2,5}})$ over $S = k[w_{(r,\gamma)}]_{w_{(r,\gamma)} \in \mathcal{W}_5}$ is

$$\begin{aligned} 0 \rightarrow S^2(-9) \rightarrow S^7(-8) \oplus S^6(-7) \rightarrow S^{70}(-6) \rightarrow S^{154}(-5) \rightarrow S^{168}(-4) \rightarrow \\ S^{98}(-3) \rightarrow S^{26}(-2) \rightarrow S \rightarrow S/I(S_{D_{2,5}}) \rightarrow 0. \end{aligned}$$

Our next aim is to describe a minimal set of generators of $I(S_{D_{2d}})$. We define a new set of indeterminates $z_{(r,\gamma)}$, we set $S' = k[z_{(r,\gamma)}]$ and we consider the linear change of variables induced by ρ (see 1):

$$(2) \quad \begin{cases} z_{(r,\gamma)} = w_{(r,\gamma)}, & \text{if } w_{(r,\gamma)} \neq w_{(2,0)} \\ z_{(2,0)} = w_{(2,0)} + 2w_{(d,d)}, \end{cases}$$

which gives an isomorphism $\tilde{\rho} : k[z_{(r,\gamma)}] \rightarrow S$ of polynomial rings. We have the following commutative diagram

$$\begin{array}{ccc} S' & \xrightarrow{\psi_d} & k[\mathcal{A}_{2d}] \\ \tilde{\rho} \downarrow & & \rho \downarrow \\ S & \xrightarrow{\varphi_d} & k[\mathcal{B}_{2d}] \end{array}$$

where $\psi_d(z_{(r,\gamma)}) = \rho^{-1}(\varphi_d(w_{(r,\gamma)}))$ if $z_{(r,\gamma)} \neq z_{(2,0)}$ (see (1)) and $\psi_d(z_{(2,0)}) = y_2^2$. In particular, ψ_d sends bijectively the variables $z_{(r,\gamma)}$ to the monomials of $\mathcal{A}_{2d} = \{y_0^{b_0} y_1^{b_1} y_2^{b_2} \mid b_0 + 2b_1 + db_2 = 2d\}$ by the formula $\psi_d(z_{(r,\gamma)}) = y_0^{d(2-r)+(d-2)\gamma} y_1^\gamma y_2^{r-\gamma}$. We obtain the following result.

Theorem 4.9. (i) $\ker(\psi_d)$ is a binomial ideal of S' minimally generated by quadrics.
(ii) $I(S_{D_{2d}}) = \tilde{\rho}(\ker(\psi_d))$ and a minimal set of generators of $I(S_{D_{2d}})$ is the following sets of binomials and trinomials:

$$\begin{aligned} & \{w_{(r_1,\gamma_1)}w_{(r_2,\gamma_2)} - w_{(r_3,\gamma_3)}w_{(r_4,\gamma_4)} \mid (r_i, \gamma_i) \neq (2, 0), r_1 + r_2 = r_3 + r_4, \gamma_1 + \gamma_2 = \gamma_3 + \gamma_4\} \\ & \{(w_{(2,0)} + 2w_{(d,d)})w_{(\gamma_1,\gamma_1)} - w_{(r_2,\gamma_2)}w_{(r_3,\gamma_3)} \mid (r_i, \gamma_i) \neq (2, 0), \gamma_1 + 2 = r_2 + r_3, \gamma_1 = \gamma_2 + \gamma_3\}. \end{aligned}$$

Proof. (i) $\ker(\psi_d)$ is generated by the set of binomials of the form $\prod_{i=1}^l z_{(r_{j_i}, \gamma_{j_i})} - \prod_{i=1}^l z_{(r_{m_i}, \gamma_{m_i})}$, $l \geq 2$, such that $\prod_{i=1}^l \psi_d(z_{(r_{j_i}, \gamma_{j_i})}) = \prod_{i=1}^l \psi_d(z_{(r_{m_i}, \gamma_{m_i})})$ (see [9, Theorem 1]). From this and Corollary 4.7, it follows that $\ker(\psi_d)$ is minimally generated by binomials of degree 2. Precisely, since we have $\psi_d(z_{(r,\gamma)}) = y_0^{d(2-r)+(d-2)\gamma} y_1^\gamma y_2^{r-\gamma}$, these binomials are:

$$(3) \quad \{z_{(r_1,\gamma_1)}z_{(r_2,\gamma_2)} - z_{(r_3,\gamma_3)}z_{(r_4,\gamma_4)} \mid r_1 + r_2 = r_3 + r_4, \gamma_1 + \gamma_2 = \gamma_3 + \gamma_4\}.$$

(ii) Since $\tilde{\rho}$ and ρ are isomorphisms of k -algebras, from the above commutative diagram we have that $I(S_{D_{2d}}) = \tilde{\rho}(\ker(\psi_d))$. Applying $\tilde{\rho}$ to (3), we obtain the description of the minimal set of generators in (ii). \square

We end this note with an example.

Example 4.10. Fix $d = 4$. We compute the homogeneous ideal $I(S_{D_{2 \cdot 4}})$ using the software Macaulay2. It is minimally generated by the 15 binomials and 5 trinomials of degree 2 that

we list below.

$$\begin{array}{ll}
w_{(0,0)}w_{(2,2)} - w_{(1,1)}^2 & w_{(1,0)}w_{(3,3)} - w_{(1,1)}w_{(3,2)} \\
w_{(0,0)}w_{(3,3)} - w_{(1,1)}w_{(2,2)} & w_{(1,0)}w_{(3,3)} - w_{(2,1)}w_{(2,2)} \\
w_{(0,0)}w_{(3,2)} - w_{(1,0)}w_{(2,2)} & w_{(1,0)}w_{(4,4)} - w_{(2,1)}w_{(3,3)} \\
w_{(0,0)}w_{(2,1)} - w_{(1,0)}w_{(1,1)} & w_{(1,0)}w_{(4,4)} - w_{(2,2)}w_{(3,2)} \\
w_{(0,0)}w_{(4,4)} - w_{(1,1)}w_{(3,3)} & w_{(1,1)}w_{(4,4)} - w_{(2,2)}w_{(3,3)} \\
w_{(0,0)}w_{(4,4)} - w_{(2,2)}^2 & w_{(2,1)}w_{(4,4)} - w_{(3,2)}w_{(3,3)} \\
w_{(1,0)}w_{(2,2)} - w_{(1,1)}w_{(2,1)} & w_{(2,2)}w_{(4,4)} - w_{(3,3)}^2 \\
w_{(1,0)}w_{(3,2)} - w_{(2,1)}^2 &
\end{array}$$

$$\begin{array}{ll}
w_{(1,0)}^2 & - w_{(0,0)}w_{(2,0)} - 2w_{(0,0)}w_{(4,4)} \\
w_{(1,0)}w_{(2,1)} & - w_{(1,1)}w_{(2,0)} - 2w_{(1,1)}w_{(4,4)} \\
w_{(1,0)}w_{(3,2)} & - w_{(2,0)}w_{(2,2)} - 2w_{(2,2)}w_{(4,4)} \\
w_{(2,1)}w_{(3,2)} & - w_{(2,0)}w_{(3,3)} - 2w_{(3,3)}w_{(4,4)} \\
w_{(3,2)}^2 & - w_{(2,0)}w_{(4,4)} - 2w_{(4,4)}^2
\end{array}$$

REFERENCES

- [1] C. Almeida, A. V. Andrade and R.M. Miró-Roig, *Gaps in the number of generators of monomial Togliatti systems*. J. Pure Appl. Algebra **223** (2019), 1817–1831.
- [2] N. Altafi and M. Boij, *The weak Lefschetz property of equigenerated monomial ideals*. J. Algebra **556** (2020) 136–168.
- [3] H. Brenner and A. Kaid, *Syzygy bundles on \mathbb{P}^2 and the Weak Lefschetz Property*, Illinois J. Math. **51** (2007), 1299–1308.
- [4] L. Colarte, E. Mezzetti and R. M. Miró-Roig, *On the arithmetic Cohen-Macaulayness of varieties parameterized by Togliatti systems*. (To appear in Ann. Mat. Pura. Appl.)
- [5] L. Colarte, E. Mezzetti, R. M. Miró-Roig and M. Salat, *On the coefficients of the permanent and the determinant of a circulant matrix. Applications*. Proc. AMS. **147** (2019), 547–558.
- [6] L. Colarte and R. M. Miró-Roig, *Minimal set of binomial generators for certain Veronese 3-fold projections*. J. Pure Appl. Algebra, **224** (2020), 768–788.
- [7] D.R. Grayson and M.E. Stillman, *Macaulay2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2/>
- [8] W. Gröbner, *Über Veronesische Varietäten und deren Projektionen*, Arch. Math. **16** (1965), 257–264.
- [9] M. Hochster, *Rings of Invariants of Tori, Cohen-Macaulay Rings Generated by Monomials, and Polytopes*. Ann. of Math. **96** (1972), 318–337.
- [10] M. Hochster and J. A. Eagon, *Cohen-Macaulay Rings, Invariant Theory, and the Generic Perfection of Determinantal Loci*, Amer. J. Math. **93** (1971), 1020–1058.
- [11] E. Mezzetti and R.M. Miró-Roig, *The minimal number of generators of a Togliatti system*, Ann. Mat. Pura Appl. (4) **195** (2016), 2077–2098.
- [12] E. Mezzetti and R.M. Miró-Roig, *Togliatti systems and Galois coverings*. J. Algebra **509** (2018), 263–291.

- [13] E. Mezzetti, R.M. Miró-Roig and G. Ottaviani, *Laplace Equations and the Weak Lefschetz Property*, *Canad. J. Math.* **65** (2013), 634–654.
- [14] M. Michałek and R.M. Miró-Roig, *Smooth monomial Togliatti systems of cubics*, *J. Combin. Theory Ser. A* **143** (2016), 66–87.
- [15] R. M. Miró-Roig and M. Salat, *On the classification of Togliatti systems*, *Comm. Algebra* **46** (2018), 2459–2475.
- [16] J. P. Serre, *Groupes algébriques et corps de classes*. Hermann, Paris, 1957.
- [17] J. P. Serre, *Linear Representations of Finite Groups*. Springer, New York, 1977.
- [18] G. Shephard and J. Todd, *Finite unitary reflection groups*. *Can. J. Math.* **6** (1954), 274–304.
- [19] R.P. Stanley, *Invariants of finite groups and their application to combinatorics*. *Bull. Amer. Math. Soc. (N.S.)* **1** (1979), 475–511.
- [20] B. Sturmfels, *Algorithms in Invariant Theory*. Second Edition. Springer, Wien, 2008.
- [21] J. Szpond, *Unexpected curves and Togliatti-type surfaces*. *Math. Nach.* **293:1** (2020) 158–168.
- [22] E. Togliatti, *Alcuni esempi di superficie algebriche degli iperspazi che rappresentano un'equazione di Laplace*, *Comm. Math. Helvetici* **1** (1929), 255–272.
- [23] E. Togliatti, *Alcune osservazioni sulle superficie razionali che rappresentano equazioni di Laplace*, *Ann. Mat. Pura Appl. (4)* **25** (1946) 325–339.
- [24] K. Yanagawa, *Some Generalizations of Castelnuovo's Lemma on Zero-Dimensional Schemes*, *J. Algebra* **170** (1994), 429–439.

DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE BARCELONA, GRAN VIA DE LES CORTS CATALANES 585, 08007 BARCELONA, SPAIN

Email address: `liena.colarte@ub.edu`

DIPARTIMENTO DI MATEMATICA E GEOSCIENZE, UNIVERSITÀ DI TRIESTE, VIA VALERIO 12/1, 34127 TRIESTE, ITALY

Email address: `mezzette@units.it`

DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE BARCELONA, GRAN VIA DE LES CORTS CATALANES 585, 08007 BARCELONA, SPAIN

Email address: `miro@ub.edu`

DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, BARCELONA GRADUATE SCHOOL OF MATHEMATICS (BGS-MATH), UNIVERSITAT DE BARCELONA, GRAN VIA DE LES CORTS CATALANES 585, 08007 BARCELONA, SPAIN

Email address: `marti.salat@ub.edu`