

# A Program Logic for Fresh Name Generation

Harold Pancho Elliott<sup>1</sup> and Martin Berger<sup>1,2</sup>

<sup>1</sup> Department of Informatics, University of Sussex, Brighton, UK.

<sup>2</sup> Turing Core, Huawei 2012 Labs, London, UK.

**Abstract.** We present a program logic for Pitts and Stark’s  $\nu$ -calculus, an extension of the call-by-value simply-typed  $\lambda$ -calculus with a mechanism for the generation of fresh names. Names can be compared for equality and inequality, producing programs with subtle observable properties. Hidden names produced by interactions between generation and abstraction are captured logically with a second-order quantifier over type contexts. We illustrate usage of the logic through reasoning about well-known difficult cases from the literature.

## 1 Introduction

Naming is a long-standing problem in computer science. Most programming languages can define naming constructs, which, when called, yield a fresh name. The  $\pi$ -calculus [11] made naming and the  $\nu$ -operator, a constructor for name creation, a first-class construct, leading to a flurry of research, e.g. [6–8, 14, 15, 18]. Initially it was unclear if the  $\pi$ -calculus approach had purchase beyond process calculi. Pitts and Stark [16] as well as Odersky [12] added the  $\nu$ -operator to the simply-typed  $\lambda$ -calculus (STLC from now on), and showed that the subtleties of naming are already present in the interplay between higher-order functions and fresh name generation. This raises the question of how compositionally to reason about programs that can generate fresh names? There are program logics for ML-like languages that can generate fresh references, such as [4, 19], but, to the best of our knowledge, always in the context of languages with other expressive features such as aliasing, mutable higher-order state or pointer arithmetic, leading to complex logics, where the contribution of fresh name generation to the difficulties of reasoning is not apparent. This is problematic because, while the type  $Nm$  carries the same information as  $Ref(Unit)$  in ML, we are often interested in reasoning about languages that combine fresh name generation with other features, such as meta-programming [3]. Can we study reasoning about fresh names in as simple a programming language as possible?

**Research question.** Is there a Hoare-style program logic for the  $\nu$ -calculus, conservatively extending program logics for the STLC in a natural manner, that allows for compositional reasoning about fresh name generation?

The present paper gives an affirmative answer to the research question, and presents the first program logic for the  $\nu$ -calculus.

**Informal explanation.** By the  $\nu$ -calculus we mean the STLC with a type  $\text{Nm}$  of names, a constructor  $\text{gensym}$  of type  $\text{Unit} \rightarrow \text{Nm}$  and a destructor, in form of equality and inequality on names ( $\text{gensym}$  and  $\nu$  are essentially identical, but the former is more widely used). Immediately we realise that the  $\nu$ -calculus loses extensionality, as  $\text{gensym}() = \text{gensym}()$  evaluates to  $\text{false}$ . While the loss of extensionality is expected in a stateful language, the  $\nu$ -calculus does not have state, at least not in a conventional sense.

A first difficulty is expressing freshness in logic. What does it mean for a name  $x$  to be fresh? A first idea might be to say that  $x$  is guaranteed to be distinct from all existing names. We cannot simply say

$$\{\top\} \text{gensym}() :_u \{\forall x. u \neq x\}$$

since we must prevent  $\forall x. u \neq x$  being instantiated to  $u \neq u$ . We want to say something like:

$$\forall x. \{\top\} \text{gensym}() :_u \{u \neq x\} \quad (1)$$

Unfortunately we cannot quantify over Hoare triples. A second problem is that (1) is not strong enough, in the sense that  $\text{gensym}$  does not just create names that are fresh w.r.t. existing names, but also w.r.t. all future calls to  $\text{gensym}$ . We introduce a new quantifier to deal with both problems at the same time. A third difficulty is that fresh names can be exported or remain hidden, with observable consequences. Consider:

$$\text{let } x = \text{gensym}() \text{ in } \lambda y. x = y \quad (2)$$

of type  $\text{Nm} \rightarrow \text{Bool}$ . It receives a name  $y$  as argument and compares it with fresh name  $x$ . Since  $x$  is never exported to the outside, no context can ever supply this fresh  $x$  to  $\lambda y. x = y$ . Hence (2) must be contextually indistinguishable from  $\lambda y. \text{false}$ . Operationally, this is straightforward. But how can we prove this compositionally? Note that this is not a property of  $\lambda y. x = y$ , but it is also not a consequence of  $x$ 's being freshly generated, for  $x$  is also fresh in this program:

$$\text{let } x = \text{gensym}() \text{ in } \langle x, \lambda y. x = y \rangle \quad (3)$$

But in (3),  $\lambda y. x = y$  can return  $\text{true}$ , for example if we use (3) in this context:

$$\text{let } p = [\cdot] \text{ in } (\pi_2 p)(\pi_1 p)$$

In program logics like [9], the specification of any abstraction  $\lambda y. M$  will be a universally quantified formula  $\forall y. A$ . With fresh names, instantiation of quantification is a core difficulty. Recall that in first-order logic,  $\forall y. A$  always implies  $A[e/y]$ , for all ambient expressions  $e$ . Clearly, in the case of (2) we cannot conclude to  $A[x/y]$  from  $\forall y. A$ , because  $x$  is, in some sense, not available for instantiation. In contrast, in the case of (3) we can infer  $A[x/y]$ . Hence we need to answer the question how to express logically the inability to instantiate a universal quantifier with a fresh and hidden name like  $x$  in (2). We introduce a novel restricted quantifier, limiting the values based on a type context, and a new quantifier over type contexts to extend the reach of restricted quantifiers.

## 2 Programming Language

Our programming language is essentially the  $\nu$ -calculus of [16], with small additions in particular pairs, included for the sake of convenience. We assume a countably infinite set of variables, ranged over by  $x, y, \dots$  and a countably infinite set, disjoint from variables, of names, ranged over by  $r, \dots$ . Constants ranged over by  $c$  are Booleans `true`, `false`, and `Unit` (`()`). For simplicity we also call our language  $\nu$ -calculus. It is given by the following grammar, where  $\alpha$  ranges over types,  $\Gamma$  over *standard type contexts* (STC),  $V$  over values and  $M$  over programs. (Additions over the STLC highlighted.)

$$\begin{aligned} \alpha &::= \text{Unit} \mid \text{Bool} \mid \mathbf{Nm} \mid \alpha \rightarrow \alpha \mid \alpha \times \alpha & \Gamma &::= \emptyset \mid \Gamma, x : \alpha \\ V &::= r \mid \mathbf{gensym} \mid x \mid c \mid \lambda x.M \mid \langle V, V \rangle \\ M &::= V \mid MM \mid \text{let } x = M \text{ in } M \mid M = M \\ &\quad \mid \text{if } M \text{ then } M \text{ else } M \mid \langle M, M \rangle \mid \pi_i(M) \end{aligned}$$

*Free variables* in  $M$ , written  $\text{fv}(M)$  are defined as usual.  $M$  is *closed* if  $\text{fv}(M) = \emptyset$ . There are no binders for names so the set  $\mathfrak{a}(M)$  of *all names* in  $M$ , is given by the obvious rules, including  $\mathfrak{a}(r) = \{r\}$ ,  $\mathfrak{a}(MN) = \mathfrak{a}(M) \cup \mathfrak{a}(N)$ . If  $\mathfrak{a}(M) = \emptyset$  then  $M$  is *compile-time* syntax. The  $\nu n.M$  constructor from the  $\nu$ -calculus [16] is equivalent to `let`  $n = \mathbf{gensym}()$  in  $M$  as  $\mathbf{gensym}()$  generates fresh names. The typing judgements is  $\Gamma \vdash M : \alpha$ , with the STC  $\Gamma$  being an unordered mapping from variables to types. Typing rules are standard [13] with the following extensions:  $\Gamma \vdash \mathbf{gensym} : \text{Unit} \rightarrow \mathbf{Nm}$  and  $\Gamma \vdash r : \mathbf{Nm}$ .

The operational semantics of our  $\nu$ -calculus is straightforward and the same as [16]. A *configuration of type*  $\alpha$  is a pair  $(G, M)$  where  $M$  is a closed term of type  $\alpha$ , and  $G$  a finite set of previously Generated names such that  $\mathfrak{a}(M) \subseteq G$ . The standard call-by-value reduction relation,  $\rightarrow$ , has the following key rules.

$$\begin{aligned} (G, (\lambda x.M)V) &\rightarrow (G, M[V/x]) \\ (G, \mathbf{gensym}()) &\rightarrow (G \cup \{n\}, n) && (n \notin G) \\ (G \cup \{n\}, n = n) &\rightarrow (G \cup \{n\}, \text{true}) \\ (G \cup \{n_1, n_2\}, n_1 = n_2) &\rightarrow (G \cup \{n_1, n_2\}, \text{false}) && (n_1 \neq n_2) \end{aligned}$$

$$(G, M) \rightarrow (G', N) \text{ implies } (G, \mathcal{E}[M]) \rightarrow (G', \mathcal{E}[N])$$

Here  $M[V/x]$  is the usual capture-avoiding substitution, and  $\mathcal{E}[\cdot]$  ranges over the usual reduction contexts of the STLC. Finally,  $\Downarrow$  is short for  $\rightarrow^*$ .

## 3 Logical Language

This section defines the syntax of the logic. As is customary for program logics, ours is an extension of first order logic with equality (alongside axioms for arithmetic). *Expressions*, ranged over by  $e, e', \dots$ , *formulae*, ranged over by  $A, B, C, \dots$

and *Logical Type Contexts* (LTCs), ranged over by  $\mathbb{I}, \mathbb{I}', \mathbb{I}_i, \dots$ , are given by the grammar below. (Extensions over [9] highlighted.)

$$\begin{aligned}
e & ::= x^\alpha \mid c \mid \langle e, e \rangle \mid \pi_i(e) \\
\mathbb{I} & ::= \emptyset \mid \mathbb{I} + x : \alpha \mid \mathbb{I} + \delta : \mathbb{TC} \\
A & ::= e = e \mid \neg A \mid A \wedge A \mid e \bullet e = x^\alpha \{A\} \mid \forall x^\alpha \in (\mathbb{I}).A \mid \forall \delta.A
\end{aligned}$$

Expressions,  $e$ , are standard, where constants,  $c$ , range over Booleans and  $()$ , but do *not* include names or `gensym` as constants. Equality, negation and conjunction are standard. Evaluation formulae  $e \bullet e' = m\{A\}$  internalise triples [9] and express that if the program denoted by  $e$  is executed with argument denoted by  $e'$ , then the result, denoted by  $m$ , satisfies  $A$ . Since the  $\nu$ -calculus has no recursion, all applications terminate and we do not distinguish partial from total correctness. We write  $e_1 \bullet e_2 = e_3$  as shorthand for  $e_1 \bullet e_2 = m\{m = e_3\}$ .

Given variables represent values, ensuring hidden names cannot be revealed in an unsafe manner requires the idea that a value is *derived* from an LTC if a name free term uses the variables in the LTC to evaluate to said value. Specifically define a name as reachable from said LTC if it can be derived from it, and hidden otherwise.

Freshness is not an absolute notion. Instead, a name is fresh with respect to something, in this case names generated in the past, and future of the computation. Formulae refer to names by variables, and variables are tracked in the STC. Freshness is now defined in two steps: (1) First we characterise freshness of a name w.r.t. the current STC, meaning the name cannot be derived from the variables in the STC. Then, (2) we define freshness w.r.t. all future extension of the current STC, details in Sec. 4. The modal operator is used in [19] in order to express “for all future extensions”, but we found modalities inconvenient, since they don’t allow us to name extensions. We introduce a new quantifier  $\forall x^\alpha \in (\mathbb{I}).A$  instead, where  $\mathbb{I}$  ranges over LTCs from which  $x$  can be derived. To make this precise, we need LTCs (explained next), a generalisation of STCs.

**LTCs.** Like STCs, LTCs map variables to types, and are needed for typing expressions, formulae and triples (introduced in Sec. 6), LTCs generalise STCs in two ways: they are *ordered*, and they don’t just contain program variables, but also *type context variables* (TCVs), ranged over by  $\delta$ . TCVs are always mapped to the new type  $\mathbb{TC}$ , short for *type context*. The ordering in LTCs is essential because  $\mathbb{I} + \delta : \mathbb{TC}$  implies  $\delta$  represents an *extension* of the LTC  $\mathbb{I}$ .

**Restricted universal quantification.** The meaning of  $\forall x^\alpha \in (\mathbb{I}).A$  is intuitively simple:  $A$  must be true for all  $x$  that range only over values of type  $\alpha$ , derived from  $\mathbb{I}$  that do *not* reveal hidden names. For example if the model contained the name  $r$  but only as  $\lambda y.y = r$ , then  $r$  was hidden and whatever  $x$  in  $\forall x^\alpha \in (\mathbb{I}).A$  ranged over, it must not reveal  $r$ . Formalising this requirement is subtle.

**Quantification over LTCs.** Below we formalise the axiomatic semantics of `gensym` by saying that the result of each call to this function is fresh w.r.t. all

future extensions of the present state (with the present state being included). The purpose of  $\forall\delta.A$  is to allow us to do so:  $\forall\delta.A$  implies for all future states derived from the current state (included), when the LTC for that state is assigned to the TCV  $\delta$ , then  $A$  holds.

***A convenient shorthand, the freshness predicate.*** We express freshness of the name  $x$  relative to the LTC  $\mathbb{I}$  as  $\forall z \in (\mathbb{I}).x \neq z$ . and, as this predicate is used pervasively, abbreviate it to  $x\#\mathbb{I}$ . Intuitively,  $x\#\mathbb{I}$ , a variant of a similar predicate in [19], states that the name denoted by  $x$  is not derivable, directly or indirectly, from the LTC  $\mathbb{I}$ .

***Typing of expressions, formulae and triples.*** We continue with setting up definitions that allow us to type expressions, formulae and triples. The ordered union of  $\mathbb{I}$  and  $\mathbb{I}'$  with  $\text{dom}(\mathbb{I}) \cap \text{dom}(\mathbb{I}') = \emptyset$  is written  $\mathbb{I} + \mathbb{I}'$ , and should be understood as: every variable from  $\text{dom}(\mathbb{I})$  comes before every variable from  $\text{dom}(\mathbb{I}')$ . Other abbreviations include  $\exists x^\alpha \in (\mathbb{I}).A \stackrel{\text{def}}{=} \neg\forall x^\alpha \in (\mathbb{I}).\neg A$ , and where  $\alpha$  is obvious  $(\mathbb{I} + y) \stackrel{\text{def}}{=} (\mathbb{I} + y : \alpha)$  (respectively  $(\mathbb{I} + \delta) \stackrel{\text{def}}{=} (\mathbb{I} + \delta : \mathbb{TC})$ ). For simplicity, where not explicitly required,  $\mathbb{I} + \delta$  is written  $\delta$ . Functions on LTCs are defined as expected including mapping variables,  $\mathbb{I}(x)$ , and TCVs,  $\mathbb{I}(\delta)$ ; obtaining the domain,  $\text{dom}(\mathbb{I})$ ; ordered removal of a variable,  $\mathbb{I}\backslash x$ ; ordered removal of all TCV,  $\mathbb{I}\backslash_{\text{TCV}}$ ; and removal of TCV to produce a STC,  $\mathbb{I} \downarrow_{\text{TC}}$ . We define free variables of LTC,  $\text{fv}(\mathbb{I}) \stackrel{\text{def}}{=} \text{dom}(\mathbb{I} \downarrow_{\text{TC}}) \stackrel{\text{def}}{=} \text{dom}(\mathbb{I}\backslash_{\text{TCV}})$ , then free variables of formulae defined as expected, with the addition of  $\text{fv}(x\#\mathbb{I}) \stackrel{\text{def}}{=} \text{fv}(\mathbb{I}) \cup \{x\}$ ,  $\text{fv}(\forall x \in (\mathbb{I}).A) \stackrel{\text{def}}{=} (\text{fv}(A)\backslash\{x\}) \cup \text{fv}(\mathbb{I})$ , and  $\text{fv}(\forall\delta.A) \stackrel{\text{def}}{=} \text{fv}(A)$ . Similarly  $\text{ftcv}(\mathbb{I})$  and  $\text{ftcv}(A)$  define all TCV occurring in  $\mathbb{I}$  and unbound by  $\forall\delta$ . in  $A$  respectively, calling  $\mathbb{I}$  *TCV-free* if  $\text{ftcv}(\mathbb{I}) \stackrel{\text{def}}{=} \emptyset$ . The typing judgement for LTCs, written  $\mathbb{I} \Vdash \mathbb{I}'$ , checks that  $\mathbb{I}'$  is an ‘ordered subset’ of  $\mathbb{I}$ . Type checks on expressions, formulae and triples use LTC as the base, written  $\mathbb{I} \Vdash e : \alpha$ ,  $\mathbb{I} \Vdash A$  and  $\mathbb{I} \Vdash \{A\} M :_u \{B\}$  respectively. Fig. 1 gives the rules defining the typing judgements. From now on we adhere to the following convention: *All expressions, formulae and triples are typed*, and we will mostly omit being explicit about typing.

***Advanced substitutions.*** Reasoning with quantifiers requires quantifier instantiation. This is subtle with  $\forall\delta.A$ , and we need to define two substitutions,  $A[e/x]_{\mathbb{I}}$  (substitutes expressions for variables) and  $A[\mathbb{I}_0/\delta]_{\mathbb{I}}$  (LTCs substituted for TCVs). First extend the definition *e is free for  $x^\alpha$  in A* in [10], to ensure if  $e$  contains destructors i.e.  $\pi_i(\ )$  or  $=$ , then all free occurrences of  $x$  in any LTC  $\mathbb{I}_0$  in  $A$  must imply  $\mathbb{I}_0 \Vdash e : \alpha$ . Below, we assume the standard substitution  $e[e'/x]$  of expressions for variables in expressions, simple details omitted.

We define  $A[e/x]_{\mathbb{I}}$ , *logical substitution of e for x in A in the context of  $\mathbb{I}$* , if  $e$  is free for  $x$  in  $A$  and  $e$  is typed by  $\mathbb{I}$ , by the following clauses (simple cases omitted) and the auxiliary operation on LTCs below. We often write  $A[e/x]$  for  $A[e/x]_{\mathbb{I}}$ .

---


$$\begin{array}{c}
\frac{b \in \{\text{true}, \text{false}\}}{\Gamma \Vdash b : \text{Bool}} \quad \frac{-}{\Gamma \Vdash () : \text{Unit}} \quad \frac{\Gamma(x) = \alpha}{\Gamma \Vdash x : \alpha} \quad \frac{\Gamma \Vdash e : \alpha \quad \Gamma \Vdash e' : \beta}{\Gamma \Vdash \langle e, e' \rangle : \alpha \times \beta} \quad \frac{\Gamma \Vdash e : \alpha_1 \times \alpha_2}{\Gamma \Vdash \pi_i(e) : \alpha_i} \\
\\
\frac{-}{\Gamma \Vdash \emptyset} \quad \frac{\Gamma \Vdash \Gamma_0}{\Gamma + x : \alpha \Vdash \Gamma_0 + x : \alpha} \quad \frac{\Gamma \Vdash \Gamma_0}{\Gamma + \delta \Vdash \Gamma_0 + \delta} \quad \frac{\Gamma \Vdash \Gamma_0}{\Gamma + \Gamma' \Vdash \Gamma_0} \\
\\
\frac{\Gamma \Vdash e_1 : \alpha \quad \Gamma \Vdash e_2 : \alpha}{\Gamma \Vdash e_1 = e_2} \quad \frac{\Gamma \Vdash A_1 \quad \Gamma \Vdash A_2}{\Gamma \Vdash A_1 \wedge A_2} \quad \frac{\Gamma \Vdash A}{\Gamma \Vdash \neg A} \quad \frac{\Gamma + \delta : \text{TC} \Vdash A}{\Gamma \Vdash \forall \delta. A} \\
\\
\frac{\Gamma \Vdash e : \alpha \rightarrow \beta \quad \Gamma \Vdash e' : \alpha \quad \Gamma + x : \beta \Vdash A}{\Gamma \Vdash e \bullet e' = x^\beta \{A\}} \quad \frac{\Gamma \Vdash x : \text{Nm} \quad \Gamma \Vdash \Gamma'}{\Gamma \Vdash x \# \Gamma'} \\
\\
\frac{\Gamma \Vdash \Gamma' \quad \Gamma + x : \alpha \Vdash A}{\Gamma \Vdash \forall x^\alpha \in (\Gamma'). A} \quad \frac{\Gamma \Vdash A \quad \Gamma \downarrow_{\text{TC}} \vdash M : \alpha \quad \Gamma + m : \alpha \Vdash B}{\Gamma \Vdash \{A\} M :_m \{B\}}
\end{array}$$


---

**Fig. 1.** Typing rules for LTCs, expressions, formulae and triples (see Sec. 6). Simple cases omitted.  $M$  in the last rule is compile-time syntax.

---

$$\begin{array}{l}
(e_1 \bullet e_2 = m\{A\})[e/x]_{\Gamma} \stackrel{\text{def}}{=} e_1[e/x] \bullet e_2[e/x] = m\{A[e/x]_{\Gamma+m}\} \quad (x \neq m, m \notin \text{fv}(\Gamma)) \\
(y \# \Gamma')[e/x]_{\Gamma} \stackrel{\text{def}}{=} y[e/x] \# (\Gamma'[e/x]_{\Gamma}) \\
(\forall m \in (\Gamma'). A)[e/x]_{\Gamma} \stackrel{\text{def}}{=} \forall m \in (\Gamma'[e/x]_{\Gamma}). (A[e/x]_{\Gamma+m}) \quad (x \neq m, m \notin \text{fv}(\Gamma)) \\
(\forall \delta. A)[e/x]_{\Gamma} \stackrel{\text{def}}{=} \forall \delta. (A[e/x]_{\Gamma+\delta}) \\
\Gamma'[e/x]_{\Gamma} \stackrel{\text{def}}{=} \begin{cases} \Gamma'_e \text{ s.t. } \text{dom}(\Gamma'_e) = \text{fv}(e) \cup \text{dom}(\Gamma' \setminus x), \Gamma \Vdash \Gamma'_e & x \in \text{dom}(\Gamma') \\ \Gamma' & x \notin \text{dom}(\Gamma') \end{cases}
\end{array}$$

*Type context substitution*  $A[\Gamma_0/\delta]_{\Gamma}$  instantiates  $\delta$  with  $\Gamma_0$  in  $A$ , similar to classical substitution. We often write  $[\Gamma_0/\delta]$  for  $[\Gamma_0/\delta]_{\Gamma}$  as  $\Gamma$  is used for ordering and is obvious. As above, the omitted cases are straightforward and the auxiliary operation on LTCs is included.

$$\begin{array}{l}
(x \# \Gamma')[\Gamma_0/\delta]_{\Gamma} \stackrel{\text{def}}{=} x \# (\Gamma'[\Gamma_0/\delta]_{\Gamma}) \\
(e_1 \bullet e_2 = m\{A\})[\Gamma_0/\delta]_{\Gamma} \stackrel{\text{def}}{=} e_1 \bullet e_2 = m\{A[\Gamma_0/\delta]_{\Gamma+m}\} \quad (m \notin \text{dom}(\Gamma_0)) \\
(\forall x \in (\Gamma'). A)[\Gamma_0/\delta]_{\Gamma} \stackrel{\text{def}}{=} \forall x \in (\Gamma'[\Gamma_0/\delta]_{\Gamma}). (A[\Gamma_0/\delta]_{\Gamma+x}) \quad (x \notin \text{dom}(\Gamma_0)) \\
(\forall \delta'. A)[\Gamma_0/\delta]_{\Gamma} \stackrel{\text{def}}{=} \begin{cases} (\forall \delta'. A[\Gamma_0/\delta]_{\Gamma+\delta'}) & \delta \neq \delta' \quad (\delta' \notin \text{dom}(\Gamma_0)) \\ \forall \delta. A & \text{otherwise} \end{cases} \\
\Gamma'[\Gamma_0/\delta]_{\Gamma} \stackrel{\text{def}}{=} \begin{cases} \Gamma_1 \text{ s.t. } \text{dom}(\Gamma_1) = \text{dom}(\Gamma_0, \Gamma'), \Gamma \Vdash \Gamma_1 & \delta \in \text{dom}(\Gamma') \\ \Gamma' & \delta \notin \text{dom}(\Gamma') \end{cases}
\end{array}$$

## 4 Model

We define a *model*  $\xi$  as a finite (possibly empty) map from variables and TCV to closed values and TCV-free LTCs respectively.

$$\xi ::= \emptyset \mid \xi \cdot x : V \mid \xi \cdot \delta : \mathbb{I}'$$

Standard actions on models  $\xi$  are defined as expected and include: variable mappings to values,  $\xi(x)$ , or TCV mapping to LTC,  $\xi(\delta)$ ; removal of variable  $x$  as  $\xi \setminus x$  (with  $(\xi \cdot \delta : \mathbb{I}_1) \setminus x = (\xi \setminus x) \cdot \delta : (\mathbb{I}_1 \setminus x)$ ); removal of TCV  $\delta$  as  $\xi \setminus \delta$ ; removal of all TCVs as  $\xi \setminus_{-TCV}$ ; and defining all names in  $\xi$  as  $\mathfrak{a}(\xi)$  noting that  $\mathfrak{a}(\mathbb{I}) = \emptyset$ .

A model  $\xi$  is typed by a LTC  $\mathbb{I}$  written  $\xi^{\mathbb{I}}$ , if  $\mathbb{I} \Vdash \xi$  as defined below, were  $\mathbb{I}_d = \mathbb{I}_d \setminus_{-TCV}$  formalises that  $\mathbb{I}_d$  is TCV-free.

$$\frac{}{\emptyset \Vdash \emptyset} \quad \frac{\mathbb{I} \Vdash \xi \quad \emptyset \vdash V : \alpha}{\mathbb{I} + x : \alpha \Vdash \xi \cdot x : V} \quad \frac{\mathbb{I} \Vdash \xi \quad \mathbb{I} \Vdash \mathbb{I}_d \quad \mathbb{I}_d = \mathbb{I}_d \setminus_{-TCV}}{\mathbb{I} + \delta \Vdash \xi \cdot \delta : \mathbb{I}_d}$$

The *closure* of a term  $M$  by a model  $\xi$ , written  $M\xi$  is defined as standard with the additions,  $\mathbf{gensym}\xi \stackrel{def}{=} \mathbf{gensym}$  and  $r\xi \stackrel{def}{=} r$ . Noting that  $M\xi \setminus_{-TCV} = M\xi = M\xi \cdot \delta : \mathbb{I}'$  holds for all  $\delta$  and  $\mathbb{I}'$  as  $\mathbb{I} \downarrow_{-TCV} \vdash M : \alpha$ .

The *interpretation of expression*  $e$  in a model  $\xi^{\mathbb{I}}$ , written  $\llbracket e \rrbracket_{\xi}$ , is standard, e.g.  $\llbracket c \rrbracket_{\xi} \stackrel{def}{=} c$ ,  $\llbracket x \rrbracket_{\xi} \stackrel{def}{=} \xi(x)$ ,  $\llbracket \langle e, e' \rangle \rrbracket_{\xi} \stackrel{def}{=} \langle \llbracket e \rrbracket_{\xi}, \llbracket e' \rrbracket_{\xi} \rangle$ , etc.

The *interpretation of LTCs*  $\mathbb{I}_0$  in a model  $\xi^{\mathbb{I}}$ , written  $\llbracket \mathbb{I}_0 \rrbracket_{\xi}$ , outputs a STC. It is assumed  $\mathbb{I} \Vdash$  the LTC in the following definition:

$$\llbracket \emptyset \rrbracket_{\xi} \stackrel{def}{=} \emptyset \quad \llbracket \mathbb{I}_0 + x : \alpha \rrbracket_{\xi} \stackrel{def}{=} \llbracket \mathbb{I}_0 \rrbracket_{\xi}, x : \alpha \quad \llbracket \mathbb{I}_0 + \delta : \mathbb{T}C \rrbracket_{\xi} \stackrel{def}{=} \llbracket \mathbb{I}_0 \rrbracket_{\xi} \cup \llbracket \xi(\delta) \rrbracket_{\xi}$$

Write  $M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V$  as the *derivation of a value*  $V$  from term  $M$  which is typed by the LTC  $\mathbb{I}$  and closed and evaluated in a model  $\xi$ . This ensures names are derived from actual reachable values in  $\xi$  as if they were programs closed by the model, hence not revealing hidden names from  $\xi$ .  $M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V$  holds exactly when:

- $\mathfrak{a}(M) = \emptyset$
- $\llbracket \mathbb{I} \rrbracket_{\xi} \vdash M : \alpha$
- $(\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi) \cup G', V)$

Model extensions aim to capture the fact that models represent real states of execution, by stating a model is only constructed by evaluating terms derivable from the model.

A model  $\xi'$  is a *single step model extension* to another model  $\xi^{\mathbb{I}}$ , written  $\xi \preceq \xi'$ , if the single new value in  $\xi'$  is derived from  $\xi$  or the mapped LTC is  $\mathbb{I}'$  with TCVs removed. Formally  $\xi^{\mathbb{I}} \preceq \xi'$  holds if either of the following hold:

- There is  $M_y^{\alpha}$  such that  $M_y \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V_y$  and  $\xi'^{\mathbb{I}'+y:\alpha} = \xi \cdot y : V_y$ .
- $\xi'^{\mathbb{I}'+\delta} = \xi \cdot \delta : \mathbb{I}' \setminus_{-TCV}$  for some  $\delta$ .

We write  $\preceq^*$  for the transitive, reflexive closure of  $\preceq$ . If  $\xi \preceq^* \xi'$  we say  $\xi'$  is an *extension* of  $\xi$  and  $\xi$  is a *contraction* of  $\xi'$ .

A model  $\xi$  is *constructed by*  $\mathbb{I}$ , written  $\mathbb{I} \triangleright \xi$ , if any TCV represents a model extension. Formally we define  $\mathbb{I} \triangleright \xi$  by the following rules:

$$\frac{-}{\emptyset \triangleright \emptyset} \quad \frac{\mathbb{I} \triangleright \xi \quad \text{exists } M^\alpha.M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V}{\mathbb{I} + x : \alpha \triangleright \xi \cdot x : V} \quad \frac{\mathbb{I} \triangleright \xi_0 \quad \xi_0 \preceq^* \xi^{\mathbb{I}_2} \quad \mathbb{I}_1 = \mathbb{I}_2 \setminus \text{-TCV}}{\mathbb{I} + \delta \triangleright \xi \cdot \delta : \mathbb{I}_1}$$

A model  $\xi^{\mathbb{I}}$  is *well constructed* if there exists an LTC,  $\mathbb{I}'$ , such that  $\mathbb{I}' \triangleright \xi$ , noting that  $\mathbb{I} \Vdash \mathbb{I}'$ .

Model extensions and well constructed models represent models derivable by  $\nu$ -calculus programs, ensuring names cannot be revealed by later programs. Consider the basic model:  $y : \lambda a. \text{if } a = r_1 \text{ then } r_2 \text{ else } r_3$ , if  $r_1$  could be added to the model, this clearly reveals access to  $r_2$  otherwise  $r_2$  is hidden. Hence the assumption that all models are well constructed from here onwards.

*Contextual equivalence* of two terms requires them to be contextually indistinguishable in all variable-closing single holed contexts of Boolean type in any valid configuration, as is standard [1, 17]. When  $M_1$  and  $M_2$  are closed terms of type  $\alpha$  and  $\hat{\mathfrak{a}}(M_1) \cup \hat{\mathfrak{a}}(M_2) \subseteq G$ , we write  $M_1 \cong_\alpha^G M_2$  to be equivalent to  $G, \emptyset \vdash M_1 \equiv M_2 : \alpha$  from [1].

#### 4.1 Semantics

The *satisfaction relation* for formula  $A$  in a well constructed model  $\xi^{\mathbb{I}}$ , written  $\xi \models A$ , assumes  $\mathbb{I} \Vdash A$ , and is defined as follows:

- $\xi \models e = e'$  if  $\llbracket e \rrbracket_\xi \cong_\alpha^{\hat{\mathfrak{a}}(\xi)} \llbracket e' \rrbracket_\xi$ .
- $\xi \models \neg A$  if  $\xi \not\models A$ .
- $\xi \models A \wedge B$  if  $\xi \models A$  and  $\xi \models B$ .
- $\xi \models e \bullet e' = m\{A\}$  if  $\llbracket e \rrbracket_\xi \llbracket e' \rrbracket_\xi \overset{[0, \xi]}{\rightsquigarrow} V$  and  $\xi \cdot m : V \models A$
- $\xi \models \forall x^\alpha \in (\mathbb{I}'). A$  if for all  $M. M \overset{[\mathbb{I}', \xi]}{\rightsquigarrow} V$  implies  $\xi \cdot x : V \models A$
- $\xi \models \forall \delta. A$  if for all  $\xi^{\mathbb{I}'} \cdot \xi \preceq^* \xi'$  implies  $\xi' \cdot \delta : (\mathbb{I}' \setminus \text{-TCV}) \models A$
- $\xi \models x \# \mathbb{I}_0$  if there is no  $M_x$  such that  $M_x \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} \llbracket x \rrbracket_\xi$

In first-order logic, if a formula is satisfied by a model, then it is also satisfied by extensions of that model, and vice-versa (as long as all free variables of the formula remain in the model). This can no longer be taken for granted in our logic. Consider the formula  $\forall \delta. \exists z \in (\delta). (z \# \mathbb{I} \wedge \neg z \# \delta)$ . Validity of this formula depends on how many names exist in the ambient model: it may become invalid under contracting the model. Fortunately, such formulae are rarely needed when reasoning about programs. In order to simplify our soundness proofs we will therefore restrict some of our axioms and rules to formulae that are stable under model extension and contractions. Sometimes we need a weaker property, where formulae preserve their validity when a variable is removed from a model. Both concepts are defined semantically next.

We define formula  $A$  as *model extensions independent*, short **Ext-Ind**, if for all  $\mathbb{I}, \xi^{\mathbb{I}}, \xi'$  such that  $\mathbb{I} \Vdash A$  and  $\xi \preceq^* \xi'$  we have:  $\xi \models A$  iff  $\xi' \models A$ .

We define formula  $A$  as *thin* w.r.t.  $x$ , written  $A$  **thin** w.r.t.  $x^\alpha$ , if for all  $\mathbb{I}$  such that  $\mathbb{I} \setminus x \Vdash A$  and  $x^\alpha \in \text{dom}(\mathbb{I})$  we have for all well constructed models  $\xi^{\mathbb{I}}$  and  $\xi \setminus x$  that:  $\xi \models A$  implies  $\xi \setminus x \models A$ .



## 5 Axioms

Axioms and axiom schemas are similar in intention to those of the logic for the STLC, but expressed within the constraints of our logic. Axiom schemas are indexed by the LTC that types them and the explicit types where noted. We introduce the interesting axioms (schemas) and those used in Sec. 7.

Equality axioms are standard where (*eq1*) allows for substitution. Most axioms for universal quantification over LTCs (*u1*)-(*u5*) are inspired by those of first order logic. The exceptions are (*u2*) which allows for the reduction of LTCs and (*u5*) which holds only on Nm-free types. Axioms for existential quantification over LTCs (*ex1*)-(*ex3*) are new aside from (*ex1*) which is the dual of (*u1*). Axiom (*ex2*) introduces existential quantification from evaluation formulae that produce a fixed result. Reducing  $\mathbb{I}$  in  $\exists x \in (\mathbb{I}).A$  is possible via (*ex3*) for a specific structure. We use base types  $\alpha_b ::= \text{Unit} \mid \text{Bool} \mid \alpha_b \times \alpha_b$  as core lambda calculus types excluding functions. Freshness axioms (*f1*)-(*f2*) show instances LTCs can be extended, whereas (*f3*)-(*f4*) reduce the LTC. Axiom (*f1*) holds due to  $f$  being derived from  $\mathbb{I}+x$ , and the rest are trivial.

$$\begin{array}{llll}
 (\text{eq1}) & \mathbb{I} \Vdash A(x) \wedge x = e & \leftrightarrow & A(x)[e/x]_{\mathbb{I}} \\
 (\text{u1}) & \mathbb{I} \Vdash \forall x^\alpha \in (\mathbb{I}_0).A & \rightarrow & A[e/x]_{\mathbb{I}} \quad \mathbb{I}_0 \vdash e : \alpha \\
 (\text{u2}) & \forall x \in (\mathbb{I}_0 + \mathbb{I}_1).A & \rightarrow & (\forall x \in (\mathbb{I}_0).A) \wedge (\forall x \in (\mathbb{I}_1).A) \\
 (\text{u3}) & A^{-x} & \leftrightarrow & \forall x \in (\mathbb{I}_0).A^{-x} \quad A - \text{Ext-Ind} \\
 (\text{u4}) & \forall x \in (\mathbb{I}_0).(A \wedge B) & \leftrightarrow & (\forall x \in (\mathbb{I}_0).A) \wedge (\forall x \in (\mathbb{I}_0).B) \\
 (\text{u5}) & \forall x^\alpha \in (\mathbb{I}_0).A & \leftrightarrow & \forall x^\alpha \in (\emptyset).A \quad \alpha \text{ is Nm-free} \\
 \\ 
 (\text{ex1}) & \mathbb{I} \Vdash A[e/x]_{\mathbb{I}} & \rightarrow & \exists x' \in (\mathbb{I}_0).A \quad \mathbb{I} \Vdash \mathbb{I}_0 \text{ and } \mathbb{I}_0 \vdash e : \alpha \\
 (\text{ex2}) & \mathbb{I} + x + \mathbb{I}_0 \Vdash a \bullet b = c \{c = x\} & \rightarrow & \exists x' \in (\mathbb{I}_0).x = x' \quad \{a, b\} \subseteq \text{dom}(\mathbb{I}_0) \\
 (\text{ex3}) & \mathbb{I} + x \Vdash \forall y \in (\emptyset).\exists z^{\text{Nm}} \in (\mathbb{I}_0 + y).x = z & \rightarrow & \exists z \in (\mathbb{I}_0).x = z \\
 \\ 
 (\text{f1}) & \mathbb{I} + x + f : \alpha \rightarrow \alpha_b \Vdash x \# \mathbb{I} & \rightarrow & x \# \mathbb{I} + f : \alpha \rightarrow \alpha_b \\
 (\text{f2}) & \mathbb{I} \Vdash x \# \mathbb{I}_0 \wedge \forall y^\alpha \in (\mathbb{I}_0).A & \leftrightarrow & \forall y^\alpha \in (\mathbb{I}_0).(x \# (\mathbb{I}_0 + y) \wedge A) \\
 (\text{f3}) & x \# \mathbb{I}_0 & \rightarrow & x \neq e \quad \mathbb{I}_0 \vdash e : \text{Nm} \\
 (\text{f4}) & x \# (\mathbb{I}_0 + \mathbb{I}_1) & \rightarrow & x \# \mathbb{I}_0 \wedge x \# \mathbb{I}_1
 \end{array}$$

Axioms for quantification over LTCs are also similar to those for the classical universal quantifier except (*utc2*) which extends the restricted quantifier to any future LTC which can only mean adding fresh names.

$$\begin{array}{llll}
 (\text{utc1}) & \mathbb{I} \Vdash \forall \delta.A & \rightarrow & A[\mathbb{I}/\delta]_{\mathbb{I}} \\
 (\text{utc2}) & \mathbb{I} \Vdash \forall x^{\text{Nm}} \in (\mathbb{I}).A^{-\delta} & \leftrightarrow & \forall \delta.\forall x^{\text{Nm}} \in (\mathbb{I} + \delta).A \quad A - \text{Ext-Ind} \\
 (\text{utc3}) & A^{-\delta} & \leftrightarrow & \forall \delta.A^{-\delta} \quad A - \text{Ext-Ind} \\
 (\text{utc4}) & \forall \delta.(A \wedge B) & \leftrightarrow & (\forall \delta.A) \wedge (\forall \delta.B)
 \end{array}$$

Axioms for the evaluation formulae are similar to those of [2]. The interaction between evaluation formulae and the new constructors are shown. All STLC

values are included in the variables of Nm-free type, and if we let  $\text{Ext}(e_2, e_2)$  stand for  $\forall x \in (\emptyset). e_1 \bullet x = m_1 \{e_2 \bullet x = m_2 \{m_1 = m_2\}\}$  then (ext) maintains extensionality in this logic for the STLC terms. Typing restrictions require  $m \notin \text{fv}(A)$  in (e1) and  $\text{fv}(e_1, e_2, m) \cap \text{dom}(\mathbb{I} + x) = \emptyset$  in (e2).

$$\begin{array}{llll}
(e1) & e_1 \bullet e_2 = m\{A \wedge B\} & \leftrightarrow & (A \wedge e_1 \bullet e_2 = m\{B\}) \quad A - \text{Ext-Ind} \\
(e2) & e_1 \bullet e_2 = m\{\forall x \in (\mathbb{I}). A\} & \leftrightarrow & \forall x \in (\mathbb{I}). e_1 \bullet e_2 = m\{A\} \\
(e3) & e_1 \bullet e_2 = m^{\alpha_b}\{\forall \delta. A\} & \leftrightarrow & \forall \delta. e_1 \bullet e_2 = m^{\alpha_b}\{A\} \quad A - \text{Ext-Ind} \\
(ext) & \text{Ext}(e_1, e_2) & \leftrightarrow & e_1 =^{\alpha_1 \rightarrow \alpha_2} e_2 \quad \alpha_1 \rightarrow \alpha_2 \text{ is Nm-free}
\end{array}$$

## 6 Rules

Our logic uses standard *triples*  $\{A\} M :_m \{B\}$  where in this logic, the program  $M$  is restricted to compile-time syntax. Triples are typed by the rule in Fig. 1. Semantics of triples is standard: if the *pre-condition*  $A$  holds and the value derived from  $M$  is assigned to the *anchor*  $m$  then the *post-condition*  $B$  holds. In detail: let  $\xi^{\mathbb{I}}$  be a model.

$$\xi^{\mathbb{I}} \models \{A\} M :_m \{B\} \quad \text{iff} \quad \xi \models A \text{ implies } (M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V \text{ and } \xi \cdot m : V \models B)$$

The triple is *valid*, written  $\models \{A\} M :_m \{B\}$ , if for all  $\mathbb{I}$  and  $\xi_0^{\mathbb{I}_0}$  we have

$$\mathbb{I} \Vdash \{A\} M :_m \{B\} \text{ and } \mathbb{I} \triangleright \xi_0 \text{ together imply: } \xi_0 \models \{A\} M :_m \{B\}$$

From here on we will assume all models are well constructed, noting that the construction of models is the essence of  $\forall \delta$ . as it allows for all possible future names generated. Variables occurring in  $\text{dom}(\xi_0) - \text{dom}(\mathbb{I})$  may never occur directly in the triple, but their mapped values will have an effect.

The rules of inference can be found in Fig. 2 and Fig. 3. We write  $\vdash \{A\} M :_m \{B\}$  to indicate that  $\{A\} M :_m \{B\}$  can be derived from these rules. Our rules are similar to those of [9] for vanilla  $\lambda$ -calculi, but suitably adapted to the effectful nature of the  $\nu$ -calculus. All rules are typed. The typing of rules follows the corresponding typing of the programs occurring in the triples, but with additions to account for auxiliary variables. We have two substantially new rules: [GENSYM] and [LET]. The former lets us reason about fresh name creation by **gensym**, the latter about the **let**  $x = M$  in  $N$ . Operationally, **let**  $x = M$  in  $N$  is often just an abbreviation for  $(\lambda x. N)M$ , but we have been unable to derive [LET] using the remaining rules and axioms. Any syntactic proof of [LET] requires [LAM] and [APP], which requires the postcondition:  $C$  thin w.r.t  $p$  for  $p$  the anchor of the [LAM] rule. We have not been able to prove this thinness for all models of the relevant type.

In comparison with [9, 19], the primary difference with our rules is our substitution. Our changes to substitution only affects [EQ] and [PROJ( $i$ )] which are reduced in strength by the new definition of substitution as more constraints

are placed on the formulae to ensure correct substitution occurs. All other rules remain equally strong. The other difference from [9] is the need for thinness to replace the standard ‘free from’, which is discussed above. Removal of variables via thinness is required in the proof of soundness, for example [APP], which produces  $u$  from  $m$  and  $n$ , hence **Ext-Ind** is insufficient given the order of  $\mathbb{I} + m + n + u \Vdash C$ , i.e.  $u$  introduced after  $m$  and  $n$ . We explain the novelty in the rules in more detail below.

---


$$\begin{array}{c}
 \frac{}{\{A[x/m]\} x :_m \{A\}} \text{[VAR]} \quad \frac{}{\{T\} \text{gensym} :_u \{\forall \delta. u \bullet () = m\{m\#\delta\}\}} \text{[GENSYM]} \\
 \\
 \frac{}{\{A[c/m]\} c :_m \{A\}} \text{[CONST]} \quad \frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{C[m = n/u]\}}{\{A\} M = N :_u \{C\}} \text{[EQ]} \\
 \\
 \frac{A - \text{Ext-Ind} \quad \mathbb{I} + \delta + x : \alpha \vdash \{A^{\neg x} \wedge B\} M :_m \{C\}}{\mathbb{I} \vdash \{A\} \lambda x^\alpha. M :_u \{\forall \delta. \forall x^\alpha \in (\delta). (B \rightarrow u \bullet x = m\{C\})\}} \text{[LAM]} \\
 \\
 \frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{m \bullet n = u\{C\}\}}{\{A\} MN :_u \{C\}} \text{[APP]} \\
 \\
 \frac{\{A\} M :_m \{B\} \quad \{B[b_i/m]\} N_i :_u \{C\} \quad b_1 = \text{true} \quad b_2 = \text{false} \quad i = 1, 2}{\{A\} \text{if } M \text{ then } N_1 \text{ else } N_2 :_u \{C\}} \text{[IF]} \\
 \\
 \frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{C[(m, n)/u]\}}{\{A\} \langle M, N \rangle :_u \{C\}} \text{[PAIR]} \quad \frac{\{A\} M :_m \{C[\pi_i(m)/u]\}}{\{A\} \pi_i(M) :_u \{C\}} \text{[PROJ}(i)] \\
 \\
 \frac{\{A\} M :_m \{B\} \quad \{B\} N :_u \{C\}}{\{A\} \text{let } m^\alpha = M \text{ in } N :_u \{C\}} \text{[LET]}
 \end{array}$$

**Fig. 2.** Rules for the core language, cf. [2,9,19]. We require  $C$  thin w.r.t  $m$  in [PROJ( $i$ )], [LET] and  $C$  thin w.r.t  $m, n$  in [EQ, APP, PAIR]. We omit LTCs where not essential.

---

In [GENSYM],  $u \bullet () = m\{m\#\delta\}$  indicates the name produced by  $u()$  and stored at  $m$  is not derivable from the LTC  $\delta$ . If there were no quantification over LTCs prior to the evaluation we could only say  $m$  is fresh from the current typing context, however we want to say that even if there is a future typing context with new names and we evaluate  $u()$ , this will still produce a fresh name. Hence we introduce the  $\forall \delta$ . to quantify over all future LTCs (and hence all future names). Elsewhere in reasoning it is key that the post-condition of [GENSYM] is **Ext-Ind** and hence holds in all extending and contracting models (assuming the anchor for **gensym** is present), reinforcing the re-applicability of **gensym** in any context.

Rules for  $\lambda$ -abstraction in previous logics for lambda-calculi [9,19] universally quantify over all possible arguments. Our corresponding [LAM] rule refines this and quantifies over current or future values that do not contain hidden names. Comparing the two LTCs typing the assumption and conclusion implies  $\mathbb{I} + \delta + x$  extends  $\mathbb{I}$  to any possible extension assigned to  $\delta$ , and extends to  $x$  a value

derived from  $\mathbb{I} + \delta$ . Hence the typing implies precisely what is conveyed in the post-condition of the conclusion: ‘ $\forall \delta. \forall x \in (\delta)$ .’. Constraints on  $\delta$  and  $x$  are introduced by  $B$ , and  $A - \text{Ext-Ind}$  implies  $A$  still holds in all extensions of  $\mathbb{I}$  including  $\mathbb{I} + \delta + x$ . The rest is trivial when we consider  $((\lambda x. M)x)\xi \cong_{\alpha}^{\hat{\delta}(\xi)} M\xi$ .

The STLC’s [LET] rule introduces  $x$  in the post-condition by means of an ‘ $\exists x. C$ ’. This fails here as  $x$  may be unreachable, hence not derivable from any extending or contracting LTC. The requirement that  $C$  thin w.r.t  $x$  ensures  $x$  is not critical to  $C$  so can either be derived from the current LTC or is hidden. Thinness ensures no reference to the variable  $m$  is somehow hidden under quantification over LTCs.

The [INVAR] rule is standard with the constraint that  $C - \text{Ext-Ind}$  to ensure  $C$  holds in the extension where  $m$  has been assigned. The [LETFRESH] rule is commonly used and hence included for convenience, but it is entirely derivable from the other rules.

$$\frac{A \rightarrow A' \quad \frac{\{A'\} M :_m \{B'\} \quad B' \rightarrow B}{\{A\} M :_m \{B\}} \text{[CONSEQ]} \quad \frac{C - \text{Ext-Ind} \quad \{A\} M :_m \{B\}}{\{A \wedge C\} M :_m \{B \wedge C\}} \text{[INVAR]}}{\frac{A - \text{Ext-Ind} \quad \mathbb{I} + m : \text{Nm} \vdash \{A \wedge m \# \mathbb{I}\} M :_u \{C\}}{\mathbb{I} \vdash \{A\} \text{ let } m = \text{gensym}() \text{ in } M :_u \{C\}} \text{[LETFRESH]}}$$

**Fig. 3.** Key structural rules [CONSEQ] and [INVAR] and for convenience the derived [LETFRESH] rule where  $C$  thin w.r.t  $m$  is required.

**Theorem 1.** *All axioms and rules are sound.*

**Theorem 2.** *The logic for the  $\nu$ -calculus is a conservative extension of the logic [9] for the STLC.*

All proofs can be found in the first author’s forthcoming dissertation [5].

## 7 Reasoning Examples

**Example 1.** We reason about the core construct  $\text{gensym}()$  in an LTC  $\mathbb{I}$ . In Line 2, (*utc1*) instantiates the postcondition to  $b \bullet () = a\{a \# \mathbb{I} + b\}$  and (*f4*) removes the  $b$  from the LTC to ensure the postcondition satisfies the thin w.r.t  $b$  requirement in [APP].

$$\begin{array}{l} 1 \quad \mathbb{I} \Vdash \{\top\} \text{gensym} :_b \{\forall \delta. b \bullet () = a\{a \# \delta\}\} \quad \text{[GENSYM]} \\ \hline 2 \quad \mathbb{I} \Vdash \{\top\} \text{gensym} :_b \{b \bullet () = a\{a \# \mathbb{I}\}\} \quad \text{[CONSEQ], (utc1), (f4), 1} \\ \hline 3 \quad \mathbb{I} + b \Vdash \{b \bullet () = a\{a \# \mathbb{I}\}\} () :_c \{b \bullet c = a\{a \# \mathbb{I}\}\} \quad \text{[CONST]} \\ \hline 4 \quad \mathbb{I} \Vdash \{\top\} \text{gensym}() :_a \{a \# \mathbb{I}\} \quad \text{[APP], 2, 3} \end{array}$$

**Example 2.** We reason about the comparison of two fresh names, clearly returning false, by applying Example 1 in the relevant LTCs.

1	$\mathbb{I} \Vdash \{\mathbb{T}\} \text{gensym}() :_a \{a \# \mathbb{I}\}$	<i>See Example 1</i>
2	$\mathbb{I} + a \Vdash \{\mathbb{T}\} \text{gensym}() :_b \{b \# \mathbb{I} + a\}$	<i>See Example 1</i>
3	$\mathbb{I} + a \Vdash \{a \# \mathbb{I}\} \text{gensym}() :_b \{a \neq b\}$	[CONSEQ], (f3), 2
4	$\mathbb{I} \Vdash \{\mathbb{T}\} \text{gensym}() = \text{gensym}() :_u \{u = \text{false}\}$	[EQ], 3

**Example 3.** Placing name generation inside an abstraction halts the production of fresh names until the function is applied. When  $y$  is of type Unit then this specification is identical to that of `gensym`.

1	$\mathbb{I} + \delta + y \Vdash \{\mathbb{T}\} \text{gensym}() :_m \{m \# \mathbb{I} + \delta + y\}$	<i>See Example 1</i>
2	$\mathbb{I} \Vdash \{\mathbb{T}\} \lambda y. \text{gensym}() :_u \{\forall \delta. \forall y \in (\delta). u \bullet y = m \{m \# \mathbb{I} + \delta + y\}\}$	[LAM], 2

**Example 4.** Generating a name outside an abstraction and returning that same name in the function is often compared to Example 3 [1,17]. We reason as follows: letting  $A_4(p) \stackrel{\text{def}}{=} \forall \delta. \forall y \in (\delta). u \bullet y = m \{m \# \mathbb{I} \wedge p = m\}$ .

1	$\{x \# \mathbb{I}\} x :_m \{m \# \mathbb{I} \wedge x = m\}$	[VAR]
2	$\{x \# \mathbb{I}\} \lambda y. x :_u \{A_4(x)\}$	[LAM], 1
3	$\{x \# \mathbb{I}\} \lambda y. x :_u \{\exists x' \in (u). A_4(x')\}$	[CONSEQ], 2
4	$\mathbb{I} \Vdash \{\mathbb{T}\} \text{let } x = \text{gensym}() \text{ in } \lambda y. x :_u \{\exists x' \in (u). A_4(x')\}$	[LETFRESH], 3

Proof of line 3 above, essentially proves  $x$  is derivable from  $u$ :

5	$A_4(x) \wedge \forall y \in (\emptyset). u \bullet y = m \{x = m\}$	(utc1), (u2), FOL
6	$A_4(x) \wedge \exists x' \in (u). x = x'$	(ex2), (ex3)
7	$\exists x' \in (u). (A_4(x) \wedge x = x')$	(u3), (u4)
8	$\exists x' \in (u). A_4(x')$	(eq1)

**Example 5.** In order to demonstrate the subtlety of hidden names, the Introduction used Program (2), which was  $M \stackrel{\text{def}}{=} \text{let } x = \text{gensym}() \text{ in } \lambda y. x = y$ . We now use our logic to reason about  $M$ .

1	$\mathbb{I}\Gamma + x + \delta + y \Vdash \{\top\} x = y :_m \{m = (x = y)\}$	[EQ]
2	$\mathbb{I}\Gamma + x \Vdash \{\top\} \lambda y. x = y :_u \{\forall \delta. \forall y \in (\delta). u \bullet y = (x = y)\}$	[LAM], 1
3	$\mathbb{I}\Gamma + x \Vdash \{x \# \mathbb{I}\Gamma\} \lambda y. x = y :_u \{x \# \mathbb{I}\Gamma \wedge \forall \delta. \forall y \in (\delta). u \bullet y = (x = y)\}$	[INVAR], 2
4	$\mathbb{I}\Gamma + x \Vdash \{x \# \mathbb{I}\Gamma\} \lambda y. x = y :_u \{\forall y \in (\mathbb{I}\Gamma + u). u \bullet y = \text{false}\}$	[CONSEQ], 3
5	$\mathbb{I}\Gamma \Vdash \{\top\} M :_u \{\forall y \in (\mathbb{I}\Gamma + u). u \bullet y = \text{false}\}$	[LETFRESH]
6	$\mathbb{I}\Gamma \Vdash \{\top\} M :_u \{\forall \delta. \forall y^{Nm} \in (\delta). u \bullet y = \text{false}\}$	(utc2)

To prove line 4 above we apply the axioms as follows:

7	$\mathbb{I}\Gamma + x + u \Vdash x \# \mathbb{I}\Gamma \wedge \forall \delta. \forall y \in (\delta). u \bullet y = (x = y)$	
8	$x \# \mathbb{I}\Gamma \wedge \forall y \in (\mathbb{I}\Gamma + x + u). u \bullet y = (x = y)$	(utc1)
9	$x \# \mathbb{I}\Gamma + u \wedge \forall y \in (\mathbb{I}\Gamma + x + u). u \bullet y = (x = y)$	(f1)
10	$x \# \mathbb{I}\Gamma + u \wedge \forall y \in (\mathbb{I}\Gamma + u). u \bullet y = (x = y)$	(u2)
11	$\forall y \in (\mathbb{I}\Gamma + u). x \# \mathbb{I}\Gamma + u + y \wedge u \bullet y = (x = y)$	(f2)
12	$\forall y \in (\mathbb{I}\Gamma + u). x \neq y \wedge u \bullet y = (x = y)$	(f3)
13	$\forall y \in (\mathbb{I}\Gamma + u). u \bullet y = \text{false}$	(e1)

**Example 6.** To demonstrate the release of a hidden variable using Program (3), which was  $M \stackrel{\text{def}}{=} \text{let } x = \text{gensym}() \text{ in } \langle x, \lambda y. x = y \rangle$ , we reason as follows, with  $A_6(p, q) \stackrel{\text{def}}{=} p \# \mathbb{I}\Gamma \wedge \forall \delta. \forall y \in (\delta). q \bullet y = (p = y)$ :

1	$\{x \# \mathbb{I}\Gamma\} x :_b \{x = b \wedge x \# \mathbb{I}\Gamma\}$	[VAR]
2	$\{\top\} \lambda y. x = y :_c \{\forall \delta. \forall y \in (\delta). c \bullet y = (x = y)\}$	See Example 4, lines 1-2
3	$\{x = b \wedge x \# \mathbb{I}\Gamma\} \lambda y. x = y :_c \{x = \pi_1(\langle b, c \rangle) \wedge C(x, c)\}$	[CONSEQ], [INVAR], 2
4	$\{x = b \wedge x \# \mathbb{I}\Gamma\} \lambda y. x = y :_c \{A_6(\pi_1(a), \pi_2(a))[\langle b, c \rangle/a]\}$	[CONSEQ], (eq1)
5	$\{x \# \mathbb{I}\Gamma\} \langle x, \lambda y. x = y \rangle :_a \{A_6(\pi_1(a), \pi_2(a))\}$	[PAIR], 1, 4
6	$\mathbb{I}\Gamma \Vdash \{\top\} M :_a \{A_6(\pi_1(a), \pi_2(a))\}$	[LETFRESH], 5

## 8 Conclusion

We have presented the first program logic for the  $\nu$ -calculus, a variant of the STLC with names as first class values. Our logic is a conservative extension with two new universal quantifiers of the logic in [9] for the STLC. We provide axioms

and proof rules for the logic, prove their soundness, and show its expressive power by reasoning about well-known difficult examples from the literature.

We are currently unable to reason about this example from [16]:

$$\begin{aligned} &\text{let } F = (\text{let } x, y = \text{gensym}() \text{ in } \lambda f^{\text{Nm} \rightarrow \text{Nm}}.fx = fy) \text{ in} \\ &\text{let } G = \lambda v^{\text{Nm}}.F(\lambda u^{\text{Nm}}.v = u) \text{ in } FG \end{aligned}$$

Is this because our logic is too inexpressive, or did we simply fail to find the right proof? Another open question is whether our logic’s approach to freshness is independent of the  $\nu$ -calculus’s lack of integers and recursion, or not? For both questions we conjecture the former, and leave them as future work.

## References

1. Benton, N., Koutavas, V.: A Mechanized Bisimulation for the Nu-Calculus. Tech. Rep. MSR-TR-2008-129, Microsoft (2008)
2. Berger, M., Tratt, L.: Program Logics for Homogeneous Generative Run-Time Meta-Programming. *Logical Methods in Computer Science (LMCS)* **11**(1:5) (2015)
3. Berger, M., Tratt, L., Urban, C.: Modelling Homogeneous Generative Meta-Programming. In: *Proc. ECOOP*. pp. 5:1–5:23 (2017)
4. Dreyer, D., Neis, G., Birkedal, L.: The impact of higher-order state and control effects on local relational reasoning. In: *Proc. ICFP*. p. 143–156 (2010)
5. Elliott, H.P.: Program Logic for Fresh Name Generation. Ph.D. thesis, University of Sussex (expected 2021), draft.
6. Fernández, M., Gabbay, M.J., Mackie, I.: Nominal Rewriting Systems. In: *Proc. PPDP*. p. 108–119 (2004)
7. Gabbay, M.J., Pitts, A.M.: A New Approach to Abstract Syntax with Variable Binding. *Formal Aspects of Computing* **13**, 341–363 (2001)
8. Honda, K.: Elementary structures in process theory (1): Sets with renaming. *MSCS* **10**(5), 617–663 (2000)
9. Honda, K., Yoshida, N.: A compositional logic for polymorphic higher-order functions. In: *Proc. PPDP’04*. pp. 191–202. ACM Press (2004)
10. Mendelson, E.: *Introduction to Mathematical Logic*. Wadsworth Inc. (1987)
11. Milner, R., Parrow, J., Walker, D.: A Calculus of Mobile Processes, Parts I and II. *Information and Computation* **100**(1) (1992)
12. Odgersky, M.: A Functional Theory of Local Names. In: *Proc. POPL*. pp. 48–59 (1994)
13. Pierce, B.C.: *Types and Programming Languages*. MIT Press (2002)
14. Pitts, A.M.: Nominal logic, a first order theory of names and binding. *Information and Computation* **186**, 165–193 (2003)
15. Pitts, A.M.: *Nominal Sets: Names and Symmetry in Computer Science*. CUP (2013)
16. Pitts, A.M., Stark, I.D.B.: Observable Properties of Higher Order Functions that Dynamically Create Local Names, or What’s new? In: *MFCS* (1993)
17. Stark, I.: *Names and Higher-Order Functions*. Ph.D. thesis, University of Cambridge (1994), technical report 363, Univ. of Cambridge Computer Laboratory
18. Urban, C., Tasson, C.: Nominal Techniques in Isabelle/HOL. In: *Proc. CADE*. p. 38–53 (2005)
19. Yoshida, N., Honda, K., Berger, M.: Logical Reasoning for Higher-Order Functions with Local State. *Logical Methods in Computer Science* **4**(2) (2008)

## Appendix

In the proofs that follow, we will use *blue* coloured to represent the “meta” logic constructors, i.e.  $\forall$  meaning “for all” (in meta language), ...

We prove soundness of the logic by proving that all the rules and axioms are sound, i.e.  $\vdash \{A\} M :_u \{B\}$  implies  $\vDash \{A\} M :_u \{B\}$

The proof of soundness of the logic requires us to prove all axioms and rules sound, this is done in App. C and App. D.

We give a syntactic characterisation of **Ext-Ind** and thinness and prove they hold as such in App. E and App. F.

Conservativity from the STLC is introduced formally and proven in App. G.

### *Structure of appendix.*

- App. A Typing rules of  $\nu$ -calculus program.
- App. B Lemmas used in following proofs
- App. C Proof of soundness of axioms
- App. D Proof of soundness of rules
- App. E Define Syntactic **Ext-Ind** and proof that syntactic **Ext-Ind** implies **Ext-Ind**
- App. F Define Syntactic thinness and proof that syntactic thinness implies thinness
- App. G Definition and proof of conservativity

## A Typing Rules For The $\nu$ -calculus

The typing rules for the  $\nu$ -calculus are included for completeness in Fig. 4

---


$$\begin{array}{c}
 \frac{c \in \{\text{true}, \text{false}\}}{\Gamma \vdash c : \text{Bool}} \quad \frac{-}{\Gamma \vdash () : \text{Unit}} \quad \frac{\Gamma(x) = \alpha}{\Gamma \vdash x : \alpha} \quad \frac{-}{\Gamma \vdash \text{gensym} : \text{Unit} \rightarrow \text{Nm}} \quad \frac{-}{\Gamma \vdash r : \text{Nm}} \\
 \\
 \frac{\Gamma, x : \alpha_1 \vdash M : \alpha_2}{\Gamma \vdash \lambda x^{\alpha_1}. M : \alpha_1 \rightarrow \alpha_2} \quad \frac{\Gamma \vdash M : \alpha_1 \rightarrow \alpha_2 \quad \Gamma \vdash N : \alpha_1}{\Gamma \vdash MN : \alpha_2} \quad \frac{\Gamma \vdash M : \alpha_1 \quad \Gamma, x : \alpha_1 \vdash N : \alpha_2}{\Gamma \vdash \text{let } x^{\alpha_1} = M \text{ in } N : \alpha_2} \\
 \\
 \frac{\Gamma \vdash M : \text{Bool} \quad \Gamma \vdash N_1 : \alpha \quad \Gamma \vdash N_2 : \alpha}{\Gamma \vdash \text{if } M \text{ then } N_1 \text{ else } N_2 : \alpha} \quad \frac{\Gamma \vdash M : \alpha_1 \quad \Gamma \vdash N : \alpha_2}{\Gamma \vdash \langle M, N \rangle : \alpha_1 \times \alpha_2} \quad \frac{\Gamma \vdash M : \alpha_1 \times \alpha_2}{\Gamma \vdash \pi_i(M) : \alpha_i}
 \end{array}$$

**Fig. 4.** Typing rules for the  $\nu$ -calculus programming language.

---

## B Lemmas Used For The Following Proofs

First we introduce some definitions that are either from the literature or derived from other definitions.



**Contextual Equivalence.**

$$\begin{aligned}
 & M_1 \cong_{\alpha}^G M_2 \quad (4) \\
 & \quad \quad \quad \leftrightarrow \\
 & \forall C[\cdot]^{\alpha}, b^{\text{Bool}}. \left( \begin{array}{l} \emptyset \vdash C[\cdot]^{\alpha} : \text{Bool} \\ \wedge \mathfrak{a}(C[\cdot]) \subseteq G \end{array} \right) \rightarrow \left( \begin{array}{l} (\mathfrak{a}(\xi) \cup G, C[M_1]) \Downarrow (G'_1, b) \\ \leftrightarrow \\ (\mathfrak{a}(\xi) \cup G, C[M_2]) \Downarrow (G'_2, b) \end{array} \right)
 \end{aligned}$$

**Derived Semantics of formulae.**

- $\xi^{\mathcal{I}} \models \exists x^{\alpha} \in (\mathcal{I}'). A$  if there exists  $M^{\alpha}. M \xrightarrow{[\mathcal{I}', \xi]} V$  and  $\xi' \cdot x : V \models A$
- $\xi \models A^{-x}(x)[e/x]_{\mathcal{I}}$  if  $x \notin \text{dom}(\xi) \wedge e \xrightarrow{[\mathcal{I}', \xi]} V \wedge \xi \cdot x : V \models A(x)$
- $\xi \models A^{-x}(x)[e/x]_{\mathcal{I}}$  if  $x \in \text{dom}(\xi) \wedge \forall x'. x' \notin \text{dom}(\xi) \rightarrow \xi \models A(x')[e/x']_{\mathcal{I}}$
- $\xi \models A^{-\delta}(\delta)[\mathcal{I}_0/\delta]_{\mathcal{I}}$  if  $\delta \notin \text{dom}(\xi)$  then  $[\mathcal{I}_0]_{\xi} = \mathcal{I}_1$  and  $\xi \cdot \delta : \mathcal{I}_1 \models A(\delta)$
- $\xi \models A^{-\delta}(\delta)[\mathcal{I}_0/\delta]_{\mathcal{I}}$  if  $\delta \in \text{dom}(\xi)$  and for all  $\delta' \notin \text{dom}(\xi). \xi \models A(\delta')[\mathcal{I}_0/\delta']_{\mathcal{I}}$

**Lemmas.** Now we introduce lemmas used to simplify the later proofs in App. C, App. D, App. E, App. F

**Lemma 1 (Adding/removing unused names maintains evaluation).**

$$\forall M, V, G', G_m. \mathfrak{a}(V) \cap G' = \emptyset \rightarrow \left( \begin{array}{l} (\mathfrak{a}(M), G', M) \Downarrow (\mathfrak{a}(M), G', G_m, V) \\ \leftrightarrow \\ (\mathfrak{a}(M), M) \Downarrow (\mathfrak{a}(M), G_m, V) \end{array} \right)$$

*Proof.* Clearly holds as  $M$  does not require any name in  $G'$  if  $\mathfrak{a}(V) \cap G' = \emptyset$ , noting that  $G \cap G_m = \emptyset$  and  $G' \cap G_m = \emptyset$  and hence if this does not hold, the fresh names in  $V$  can be renamed so that it does hold.

*Note:* If  $\mathfrak{a}(V) \cap G' \neq \emptyset$  then this fails as it would imply previously generated names that are not accessible can be reached.

**Lemma 2 (Congruence is unaffected by adding/removing excess names).**

$$\forall G, G'. \mathfrak{a}(M, N) \cap \mathfrak{a}(G') = \emptyset \rightarrow (M \cong_{\alpha}^G N \leftrightarrow M \cong_{\alpha}^{G \cup G'} N)$$

*Proof.* Clearly holds by Sem.  $\cong_{\alpha}^{G \cup G'}$  where  $\mathfrak{a}(M, N) \subseteq G$ .

In the following lemmas we use  $\alpha_b$  as the base types as defined in Sec. 5 i.e.  $\alpha_b ::= \text{Unit} \mid \text{Bool} \mid \alpha_b \times \alpha_b$ . Note  $\alpha_b$  is also always a Nm-free type.

**Lemma 3 (Base type values are name free).**

$$\forall V\text{-value}. \emptyset \vdash V : \alpha_b \rightarrow \mathfrak{a}(V) = \emptyset$$

*Proof.* By induction on the structure of type  $\alpha_b$ , so clearly holds.

- $\alpha = \text{Unit}$  implies  $V = \text{Unit}$  so this clearly holds.
- $\alpha = \text{Bool}$  implies  $V = \text{true}$  or  $V = \text{false}$  so this clearly holds.

- $\alpha = \alpha_{b_1} \times \alpha_{b_2}$  then  $V = \langle V_1, V_2 \rangle$  and by IH assuming
  - $\forall V_1\text{-value}.\emptyset \vdash V_1 : \alpha_{b_1} \rightarrow \hat{\mathfrak{a}}(V_1) = \emptyset$
  - $\forall V_2\text{-value}.\emptyset \vdash V_2 : \alpha_{b_2} \rightarrow \hat{\mathfrak{a}}(V_2) = \emptyset$
  - then clearly  $\hat{\mathfrak{a}}(\langle V_1, V_2 \rangle) = \emptyset$

**Lemma 4 (Base type values can be derived equally from any LTC).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}_1, V^{\alpha_b}. \mathbb{I} \Vdash \mathbb{I}_1 \rightarrow (\exists M_0^{\alpha_b}. M_0 \overset{[\emptyset, \xi]}{\rightsquigarrow} V \leftrightarrow \exists M_1^{\alpha_b}. M_1 \overset{[\mathbb{I}_1, \xi]}{\rightsquigarrow} V)$$

*Proof.* Holds as  $M_0 \equiv V \equiv M_1$  always holds as  $\hat{\mathfrak{a}}(V) = \emptyset \wedge \emptyset \vdash V : \alpha_b$ .

**Lemma 5 (Base type values don't extend reach).**

$$\forall \mathbb{I}_0, \forall \xi^{\mathbb{I}_0}. \mathbb{I}_0 \Vdash \mathbb{I} + y : \alpha_b \rightarrow (\exists M^{\text{Nm}}. M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V \leftrightarrow \exists M^{\text{Nm}}. M \overset{[\mathbb{I} + y, \alpha_b, \xi]}{\rightsquigarrow} V)$$

*Proof.* Clearly holds from Lemmas 3 and 4 so no names are added to the LTC.

**Lemma 6 (Base type values can be added and removed from the  $\delta$  mappings without any harm).**

$$\forall \mathbb{I}. \forall \xi^{\mathbb{I}}. \xi \cdot x : V_x^{\alpha_b} \cdot \delta : \mathbb{I} \setminus \text{-TCV} \models A \leftrightarrow \xi \cdot x : V_x^{\alpha_b} \cdot \delta : (\mathbb{I} + x : \alpha_b) \setminus \text{-TCV} \models A$$

*Proof.* The only two occurrences of  $\delta$  possible in assertions are in  $\mathbb{I}_0$  in  $\forall x \in (\mathbb{I}_0)$ . and  $x \# \mathbb{I}_0$ :

If  $A$  contains  $\forall x \in (\mathbb{I}_0). A'$  and  $\mathbb{I}_0$  contains  $\delta$  then it is unaffected by the addition (or removal) of  $x : \alpha_b$  as  $M \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \leftrightarrow M \overset{[\mathbb{I}_0 + y, \alpha_b, \xi]}{\rightsquigarrow} V$  from Lemma 5

If  $A$  contains  $x \# \mathbb{I}_0$  and  $\mathbb{I}_0$  contains  $\delta$  then it is unaffected by the addition (or removal) of  $x : \alpha_b$  as  $M \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \leftrightarrow M \overset{[\mathbb{I}_0 + y, \alpha_b, \xi]}{\rightsquigarrow} V$  from Lemma 5 hence the Lemma holds.

**Lemma 7 (Base type values can be added/removed and maintain extensions).**

$$\forall \mathbb{I}^{-m}, \xi^{\mathbb{I}}, M^{\alpha_b}, V_m. M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V_m \rightarrow \forall \mathbb{I}', \xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \leftrightarrow \xi \cdot m : V_m \preceq^* \xi' \cdot m : V_m$$

*Proof.*  $\rightarrow$ : from Lemma 28 ( $\hat{\mathfrak{a}}(V_m) = \emptyset$ ).

$\leftarrow$ : This holds as  $V_m : \alpha_b$  means the variable  $m$  can always be replaced by a value  $V_m$  which by Lemma 3,  $\hat{\mathfrak{a}}(V_m) = \emptyset$  and by being a value,  $\emptyset \vdash V_m : \alpha_b$ , so their substitution is not problematic.

**Lemma 8 (Semantics of LTC is equal in model extensions).**

$$\forall \mathbb{I}_1, \mathbb{I}_2, \xi_1^{\mathbb{I}_1}, \xi_2^{\mathbb{I}_2}. (\xi_1 \preceq^* \xi_2 \wedge \mathbb{I}_1 \Vdash \mathbb{I}_0) \rightarrow \llbracket \mathbb{I}_0 \rrbracket_{\xi_1} \equiv \llbracket \mathbb{I}_0 \rrbracket_{\xi_2}$$

*Proof.* Prove by induction on the structure of  $\mathbb{I}_0$ : Induction on  $\mathbb{I}_0$

$$\begin{aligned} \mathbb{I}_0 \equiv \emptyset: \llbracket \emptyset \rrbracket_{\xi_1} &\equiv \llbracket \emptyset \rrbracket_{\xi_2} \\ \mathbb{I}_0 \equiv \mathbb{I}'_0 + x : \alpha: &\text{By IH } \llbracket \mathbb{I}'_0 \rrbracket_{\xi_1} \equiv \llbracket \mathbb{I}'_0 \rrbracket_{\xi_2} \text{ and also by Def } \preceq^*, \mathbb{I}_1(x) \equiv \mathbb{I}_2(x). \end{aligned}$$

$\mathbb{I}'_0 \equiv \mathbb{I}'_0 + \delta : \text{TC}$ : By IH  $[[\mathbb{I}'_0]]_{\xi_1} \equiv [[\mathbb{I}'_0]]_{\xi_2}$  and also by Def  $\preceq^*$ ,  $\mathbb{I}'_1(\delta) \equiv \mathbb{I}'_2(\delta)$  and  $\xi_1(\delta) \equiv \xi_2(\delta)$  gives us that this statement holds.

$\mathbb{I}'_0 \equiv \mathbb{I}'_0 + \mathbb{I}''_0$ : by IH on both  $\mathbb{I}'_0$  and  $\mathbb{I}''_0$  this clearly holds.

**Lemma 9 (Model extensions close terms equally).**

$$\forall \mathbb{I}_1, \xi_1^{\mathbb{I}_1}, \mathbb{I}_2, \xi_2^{\mathbb{I}_2}. (\xi_1 \preceq^* \xi_2 \wedge [[\mathbb{I}_1]]_{\xi_1} \vdash M : \alpha) \rightarrow M\xi_1 \equiv M\xi_2$$

*Proof.* Prove by induction on the structure of  $M$  using the definition of term closure:

$M \equiv x$ :  $x \in \mathbb{I}_1$  implies (Def  $\preceq^*$ )  $x\xi_1 \equiv \xi_1(x) \equiv \xi_2(x) \equiv x\xi_2$   
 $x \notin \mathbb{I}_1$  implies  $x$  is introduced by a  $\lambda$  hence should be left untouched.

$M \equiv c$ :  $c\xi_1 \equiv c \equiv c\xi_2$

$M \equiv \text{gensym}$ :  $\text{gensym}_{\xi_1} \equiv \text{gensym} \equiv \text{gensym}_{\xi_2}$

$M \equiv \lambda x.M'$ :  $x \in \mathbb{I}_1$  implies  $\alpha$ -renaming can be used to substitute  $x$  for some other unused variable,

i.e.  $\lambda x.M \equiv \lambda z.(M[z/x])$  hence the same proof as below holds.

$x \notin \mathbb{I}_1$  implies  $\xi_1^{-x} \equiv \xi_1 \preceq^* \xi_2 \equiv \xi_2^{-x}$

and IH implies  $M'\xi_1^{-x} \equiv M'\xi_2^{-x}$

hence  $(\lambda x.M')\xi_1 \equiv \lambda x.(M'\xi_1^{-x}) \equiv \lambda x.(M'\xi_2^{-x}) \equiv (\lambda x.M')\xi_2$

$M \equiv N_1N_2$ : by IH on both  $N_1$  and  $N_2$  this clearly holds.

$M \equiv \text{let } x = N_1 \text{ in } N_2$  holds by induction on both  $N_1$  and  $N_2$  similar to the above two cases with  $x$   $\alpha$ -renamed if required.

$M \equiv \langle N_1, N_2 \rangle$ : by IH on both  $N_1$  and  $N_2$  this clearly holds.

$M \equiv \pi_i(M')$ : by IH on  $M'$  this clearly holds.

$M \equiv \text{if } M' \text{ then } N_1 \text{ else } N_2$ : by IH on both  $M'$ ,  $N_1$  and  $N_2$  this clearly holds.

**Lemma 10 (LTC derived values are unaffected by the addition/removal of a TCV to model).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}_0, M. \mathbb{I} \Vdash \mathbb{I}_0 \rightarrow (M \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \leftrightarrow M \overset{[\mathbb{I}_0, \xi \cdot \delta : \mathbb{I} \setminus \text{TCV}]}{\rightsquigarrow} V)$$

*Proof.* This holds simply by Def  $\overset{[\cdot]}{\rightsquigarrow}$  and the facts that  $\delta$  cannot appear in  $\mathbb{I}_0$  and  $\hat{\mathfrak{a}}(\xi) \equiv \hat{\mathfrak{a}}(\xi \cdot \delta : \mathbb{I} \setminus \text{TCV})$  and  $M\xi \equiv M(\xi \cdot \delta : \mathbb{I} \setminus \text{TCV})$  and the possibly derivable terms are equivalent.

**Lemma 11 (Evaluation under model extensions are equivalent).**

$$\begin{aligned} & \forall \mathbb{I}_1, \xi_1^{\mathbb{I}_1}, \mathbb{I}_2, \xi_2^{\mathbb{I}_2}, \mathbb{I}_0, M, V. \\ & (\xi_1 \preceq^* \xi_2 \wedge \mathbb{I}_1 \Vdash \mathbb{I}_0 \wedge \hat{\mathfrak{a}}(V) \cap \hat{\mathfrak{a}}(\xi_2) \subseteq \hat{\mathfrak{a}}(\xi_1)) \\ & \rightarrow (M \overset{[\mathbb{I}_0, \xi_1]}{\rightsquigarrow} V \leftrightarrow M \overset{[\mathbb{I}_0, \xi_2]}{\rightsquigarrow} V) \end{aligned}$$

*Proof.* Essentially  $M\xi_1 \equiv M\xi_2$  (Lemma 9) and Lemma 1 prove this in both directions of the  $\leftrightarrow$ .

1	Assume: $\mathbb{I}_1, \mathbb{I}_2, \xi_1^{\mathbb{I}_1}, \xi_2^{\mathbb{I}_2}, \mathbb{I}_0$ s.t	$\forall \mathbb{I}_1, \xi_1^{\mathbb{I}_1}, \mathbb{I}_2, \xi_2^{\mathbb{I}_2}, \mathbb{I}_0.$
2	$\xi_1 \preceq^* \xi_2 \wedge \mathbb{I}_1 \Vdash \mathbb{I}_0 \wedge M \overset{[\mathbb{I}_0, \xi_1]}{\rightsquigarrow} V \wedge \mathring{a}(V) \cap \mathring{a}(\xi_2) \subseteq \mathring{a}(\xi_1)$	Assume
3	$\xi_1 \preceq^* \xi_2 \wedge \mathbb{I}_1 \Vdash \mathbb{I}_0$ $\wedge \mathring{a}(M) = \emptyset \wedge \llbracket \mathbb{I}_0 \rrbracket_{\xi_1} \vdash M : \alpha$ $\wedge (\mathring{a}(\xi_1), M\xi_1) \Downarrow (\mathring{a}(\xi_1), G', V) \leftrightarrow (\mathring{a}(\xi_1), M\xi_2) \Downarrow (\mathring{a}(\xi_1), G', V)$	Lemma 9 $\rightarrow M\xi_1 \equiv M\xi_2$
4	$\xi_1 \preceq^* \xi_2 \wedge \mathbb{I}_1 \Vdash \mathbb{I}_0$ $\wedge \mathring{a}(M) = \emptyset \wedge \llbracket \mathbb{I}_0 \rrbracket_{\xi_1} \vdash M : \alpha$ $\wedge (\mathring{a}(\xi_1), M\xi_2) \Downarrow (\mathring{a}(\xi_1), G', V) \leftrightarrow (\mathring{a}(\xi_2), M\xi_2) \Downarrow (\mathring{a}(\xi_2), G', V)$	$\llbracket \mathbb{I}_0 \rrbracket_{\xi_1} \vdash M : \alpha$ $\rightarrow \mathring{a}(M\xi_1) \subseteq \mathring{a}(\xi_1) \subseteq \mathring{a}(\xi_2)$ Lemma 1 $\mathring{a}(V) \cap \mathring{a}(\xi_2) \subseteq \mathring{a}(\xi_1)$
5	<i>hence:</i> $\forall \mathbb{I}_1, \xi_1^{\mathbb{I}_1}, \mathbb{I}_2, \xi_2^{\mathbb{I}_2}, \mathbb{I}_0, M.$ $(\xi_1 \preceq^* \xi_2 \wedge \mathbb{I}_1 \Vdash \mathbb{I}_0)$ $\rightarrow (M \overset{[\mathbb{I}_0, \xi_1]}{\rightsquigarrow} V \leftrightarrow M \overset{[\mathbb{I}_0, \xi_2]}{\rightsquigarrow} V)$	Lemma 8, $\llbracket \mathbb{I}_0 \rrbracket_{\xi_1} \equiv \llbracket \mathbb{I}_0 \rrbracket_{\xi_2}$

The next few lemmas will be defined for types which are **Nm**-free, these represent the STLC types although the key is there are  $\nu$ -calculus programs that are of this **Nm**-free type but do contain names e.g.  $\lambda x.(r = r)$ . We show that these **Nm**-free typed terms can be equated to a name-free term i.e. a STLC term. One of the key assumptions to make the proof simple is that the initial STLC is simple enough to not contain infinite values of a single type i.e. there are no integers and no recursion, however proving these harder extensions would be harder. To prove a finite number of values of any particular **Nm**-free type (up to equivalence), we first prove that it is possible to define when two functions of **Nm**-free type are equal, this is essentially through comparing each of the finite values of the input type.

**Definition 1.** *Inductively define  $\text{EQ}^\alpha(M, N)$  as the program that equates two functions of type  $\alpha$  as follows:*

$$\begin{aligned}
 \text{EQ}^{\text{Unit}}(M, N) &\stackrel{\text{def}}{=} \text{true} \\
 \text{EQ}^{\text{Bool}}(M, N) &\stackrel{\text{def}}{=} M = N \\
 \text{EQ}^{\alpha \rightarrow \text{Unit}}(M, N) &\stackrel{\text{def}}{=} \text{true} \\
 \text{EQ}^{\alpha_1 \times \alpha_2}(M, N) &\stackrel{\text{def}}{=} \text{if } \neg \text{EQ}^{\alpha_1}(\pi_1(M), \pi_1(N)) \text{ then false else} \\
 &\quad \text{if } \neg \text{EQ}^{\alpha_2}(\pi_2(M), \pi_2(N)) \text{ then false else true} \\
 \text{EQ}^{\alpha_1 \rightarrow \alpha_2}(M, N) &\stackrel{\text{def}}{=} \text{if } \neg \text{EQ}^{\alpha_2}(M\tilde{V}_0, N\tilde{V}_0) \text{ then false else} \\
 &\quad \text{if } \neg \text{EQ}^{\alpha_2}(M\tilde{V}_1, N\tilde{V}_1) \text{ then false else} \\
 &\quad \dots \\
 &\quad \text{if } \neg \text{EQ}^{\alpha_2}(M\tilde{V}_k, N\tilde{V}_k) \text{ then false else true}
 \end{aligned}$$

*where  $\tilde{V}$  is the set of all finite- $k$  number of values of type  $\alpha_1$  (Lemma 12)*

In the following lemmas we will write  $M_\nu$  for a standard  $\nu$ -calculus term and  $M_\lambda$  for a  $\nu$ -calculus term constructed only from STLC terms (i.e. no names and no gensym), we also ignore let  $x = M$  in  $N$  for simplicity.

**Lemma 12 (There are finite STLC values for each type).**

$$\forall \alpha. \exists \tilde{W}_{\lambda \text{ finite}}^\alpha. \forall M_\lambda^\alpha. \exists V_\lambda^\alpha \in \tilde{W}_\lambda. M_\lambda \cong V_\lambda$$

*Proof.* By induction on the structure of  $\alpha$  we create a complete list  $\tilde{W}^\alpha \equiv V[\alpha]$  as follows:

- $\alpha \equiv \text{Unit}$  then this clearly holds as  $(G, M) \Downarrow (G', ())$  must always hold and so  $V[\text{Unit}] \equiv ()$  holds.
- $\alpha \equiv \text{Bool}$  then this clearly holds as  $(G, M) \Downarrow (G', \text{true})$  must always hold or  $(G, M) \Downarrow (G', \text{false})$  must always hold and so  $V[\text{Bool}] \equiv \text{true} \mid \text{false}$  holds.
- $\alpha \equiv \alpha_1 \times \alpha_2$  then clearly  $V[\alpha_1 \times \alpha_2] \equiv \langle V[\alpha_1], V[\alpha_2] \rangle$  where in the RHS  $V[\alpha_1]$  represents every possible value of type  $\alpha_1$ , hence the RHS is the list of every possible combination between  $V[\alpha_1]$  and  $V[\alpha_2]$ .
- $\alpha \equiv \alpha_1 \rightarrow \alpha_2$  then by induction on  $\alpha_1$  we can assume there are finite values of this type i.e. let  $\tilde{W} \equiv V[\alpha_1]$ , we use this to state the values of type  $\alpha_1 \rightarrow \alpha_2$  as

$$\begin{aligned} V[\alpha_1 \rightarrow \alpha_2] \equiv & \lambda x^{\alpha_1}. \text{if EQ}^{\alpha_1}(x, \tilde{W}_0) \text{ then } V[\alpha_2] \text{ else} \\ & \text{if EQ}^{\alpha_1}(x, \tilde{W}_1) \text{ then } V[\alpha_2] \text{ else} \\ & \dots \\ & \text{if EQ}^{\alpha_1}(x, \tilde{W}_k) \text{ then } V[\alpha_2] \text{ else } V[\alpha_2] \end{aligned}$$

hence the RHS is the list of every possible combination between  $\tilde{W}$ , and  $V[\alpha_2]$  in each instance.

The number of values grow exponentially with the size of the type but there are always finite number of values for each type. These values cover all possible cases by definition as no other possible inputs or outputs exist to a function of the given type.

**Lemma 13 (Nm-free terms are equivalent to a name free STLC term).**

If  $\alpha$  is Nm-free.

$$\forall M_\nu^\alpha. \exists N_\lambda. M_\nu \cong_\alpha^{\hat{a}(M_\nu)} N_\lambda$$

I.e. for each term in the  $\nu$ -calculus of a type which is Nm-free then there exists an equivalent term constructed only of core STLC terms.

*Proof.* It is clear that all  $\nu$ -term that are of type **Unit** and **Bool** are equivalent to a constants of that type.

The case for  $\alpha \equiv \alpha_1 \times \alpha_2$  holds trivially by induction on  $\pi_1(M_\nu)^{\alpha_1}$  and  $\pi_2(M_\nu)^{\alpha_2}$ .

The case for  $\alpha \equiv \alpha_1 \rightarrow \alpha_2$  holds as follows:

1	$IH(\alpha) \stackrel{\text{def}}{=} \forall M^\alpha. \exists N_\lambda. M \cong_{\alpha}^{\hat{a}(M)} N$	$IH(\alpha)$
2	Assume: $IH(\alpha_1) \wedge IH(\alpha_2)$	for $\alpha_1, \alpha_2$ being Nm-free
3	Prove: $IH(\alpha_1 \rightarrow \alpha_2)$	for $\alpha_1 \rightarrow \alpha_2$ being Nm-free
4	$\forall \alpha. \exists \tilde{W}_{\text{finite}}^\alpha. \forall M_\lambda^\alpha. \exists V^\alpha \in \tilde{W}. M \cong V$	$\alpha$ is Nm-free Lemma 12
5	$IH(\alpha_1)$ implies for each $\nu$ -term of type $\alpha_1$ there exists an equivalent $\lambda$ -term (which must mean there are finite ones of these) which we call $\tilde{W}$	$IH(\alpha_1)$
6	$\forall M_{1\nu}^{\alpha_1}. \exists N_{1\lambda}^{\alpha_1}. M_{1\nu} \cong_{\alpha_1}^{\hat{a}(M_{1\nu})} N_{1\lambda}$	$IH(\alpha_1)$
7	For each value in $\tilde{W}$ , then $M\tilde{W}_i$ is a term of type $\alpha_2$ which by $IH(\alpha_2)$ implies there is an equivalent $\lambda$ -term we call $\tilde{U}_i$ i.e. $M\tilde{W}_i \cong \tilde{U}_i$	$IH(\alpha_2)$
8	$\forall \tilde{W}_i^{\alpha_1} \in \tilde{W}^{\alpha_1}. \forall M_\nu^{\alpha_1 \rightarrow \alpha_2}. \exists N_\lambda^{\alpha_2}. M\tilde{W}_i \cong_{\alpha_2}^{\hat{a}(M)} N$	$IH(\alpha_2)$
9	Using lines 5-8 we can build an equivalent formula to $M$ by brute force such that for each input case we have the equivalent output case i.e. $N_\lambda \equiv \lambda x^{\alpha_1}. \text{if EQ}^{\alpha_1}(x, \tilde{V}_0) \text{ then } \tilde{U}_0 \text{ else}$ $\quad \text{if EQ}^{\alpha_1}(x, \tilde{V}_1) \text{ then } \tilde{U}_1 \text{ else}$ $\quad \dots$ $\quad \text{if EQ}^{\alpha_1}(x, \tilde{V}_{k-1}) \text{ then } \tilde{U}_{k-1} \text{ else } \tilde{U}_k$	
10	By definition $M_\nu \cong_{\alpha_1 \rightarrow \alpha_2}^{\hat{a}(M_\nu)} N_\lambda$ as any use of $N_\lambda$ behaves identically to $M_\nu$ in any application it is used in.	

**Lemma 14 (Functions that map to base types cannot reveal names in that function).**

$$\forall M^{\text{Nm}}, V^{(\alpha \rightarrow \alpha_b)}, r_x^{\text{Nm}}. (G \equiv \hat{a}(M) \cup \hat{a}(V)) \left( \begin{array}{l} r_x \notin \hat{a}(M) \\ \wedge f : \alpha \rightarrow \alpha_b \vdash M : \text{Nm} \\ \wedge r_x \in \hat{a}(V) \\ \wedge \emptyset \vdash V : (\alpha \rightarrow \alpha_b) \end{array} \right) \rightarrow \neg(G, M[V/f]) \Downarrow (G, G', r_x)$$

*Proof.* Given the semantics,  $\Downarrow \equiv \rightarrow^*$ .

Given  $r_x \notin \hat{a}(M)$  then it is clear  $M \neq r_x$  and given  $\emptyset \vdash V : (\alpha \rightarrow \alpha_b)$  then it is clear  $V \neq r_x$ , hence  $r_x$  must be derived from terms  $M$  and  $V$ . Assume there exists at least one such  $M$  fro which this holds, then take the smallest one  $M$ , then by definition there exists an  $M_k$  such that  $(G, M[V/f]) \Downarrow (G, G', r_x) \leftrightarrow (G, M[V/f]) \Downarrow (G, G', M_k) \rightarrow (G, G', r_x)$  then the following must hold:

- $M_k \equiv \pi_1(\langle r_x, V' \rangle)$  hence  $M[V/f] \equiv \mathcal{E}[\pi_1(\langle M_1, M_2 \rangle)][V/f]$  where  $\mathcal{E}[M_2][V/f] \Downarrow r_x$  so this  $\mathcal{E}[M_2]$  is smaller than  $M$  hence contradiction.
- $M_k \equiv \pi_2(\langle V', r_x \rangle)$  hence  $M[V/f] \equiv \mathcal{E}[\pi_2(\langle M_1, M_2 \rangle)][V/f]$  where  $\mathcal{E}[M_2][V/f] \Downarrow r_x$  so this  $\mathcal{E}[M_2]$  is smaller than  $M$  hence contradiction.
- $M_k \equiv \text{if true then } r_x \text{ else } V'$  hence  $M[V/f] \equiv \mathcal{E}[\text{if } M_b \text{ then } M_1 \text{ else } M_2][V/f]$  where  $\mathcal{E}[M_1][V/f] \Downarrow r_x$  so this  $\mathcal{E}[M_1]$  is smaller than  $M$  hence contradiction.
- $M_k \equiv \text{if false then } V' \text{ else } r_x$  hence  $M[V/f] \equiv \mathcal{E}[\text{if } M_b \text{ then } M_1 \text{ else } M_2][V/f]$  where  $\mathcal{E}[M_2][V/f] \Downarrow r_x$  so this  $\mathcal{E}[M_2]$  is smaller than  $M$  hence contradiction.
- $M_k \equiv (\lambda a.r_x)V'$  hence  $M[V/f] \equiv \mathcal{E}[(\lambda a.M_1)M_2][V/f]$  where  $\mathcal{E}[M_1][V/f] \Downarrow r_x$  so this  $C[M_1]$  is smaller than  $M$  hence contradiction.
- $M_k \equiv (\lambda a.a)r_x$  hence  $M[V/f] \equiv \mathcal{E}[(\lambda a.M_1)M_2][V/f]$  where  $\mathcal{E}[M_2][V/f] \Downarrow r_x$  so this  $C[M_2]$  is smaller than  $M$  hence contradiction.
- $M_k \equiv \text{let } x = V' \text{ in } M$  Similar for let  $x = V'$  in  $M$
- There are no other possible terms that reduce to  $r_x$ .
- $M_k \equiv \text{gensym}()$  fails to produce  $r_x$  as  $r_x \in G$

i.e. Assuming a smallest  $M$  s.t.  $M \Downarrow r_x$  (which is assumed to exist), for each term  $M_k$  which is just one  $\rightarrow$ -step away from  $r_x$  then it can be proven that a smaller  $M$  could be produced hence contradiction.

**Lemma 15 (Adding a name to the model but not to the context means it is fresh).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, r_x. (\exists M_x.M_x \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} r_x \wedge \neg \exists N_x.N_x \overset{[\mathbb{I}, \xi \cdot x : r_x]}{\rightsquigarrow} r_x) \rightarrow r_x \notin \hat{\mathfrak{a}}(\xi)$$

*Proof.* Assume  $r_x \in \hat{\mathfrak{a}}(\xi)$  then clearly there would be a direct contradiction in the assumption as  $\exists N_x.N_x \overset{[\mathbb{I}, \xi \cdot x : r_x]}{\rightsquigarrow} r_x$  as  $\hat{\mathfrak{a}}(\xi \cdot x : r_x) \equiv \hat{\mathfrak{a}}(\xi)$ . Hence  $r_x \notin \hat{\mathfrak{a}}(\xi)$ .

**Lemma 16 (LTC derived terms cannot reveal old names).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}_0, M_y^{\alpha_y}, s^{\text{Nm}}. \left( \begin{array}{l} \mathbb{I} \Vdash \mathbb{I}_0 \\ \wedge M_y \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_y \\ \wedge s \in \hat{\mathfrak{a}}(\xi) \end{array} \right) \rightarrow \begin{array}{l} \exists M_1.M_1 \overset{[\mathbb{I}_0+y, \xi \cdot y : V_y]}{\rightsquigarrow} s \\ \leftrightarrow \\ \exists M_0.M_0 \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} s \end{array}$$

Essentially  $V_y$  cannot reveal any names in  $\xi$  that are not already available to  $\mathbb{I}_0$ .

*Proof.* Proof by contradiction:

assume some  $\mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}_0, M_y^{\alpha_y}, s^{\text{Nm}}$  with  $\mathbb{I} \Vdash \mathbb{I}_0 \wedge M_y \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_y \wedge s \in \hat{\mathfrak{a}}(\xi)$  then show

the following fails  $\neg \exists M_1.M_1 \overset{[\mathbb{I}_0+y, \xi \cdot y : V_y]}{\rightsquigarrow} s \wedge \exists M_0.M_0 \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} s$  but this creates a contradiction simply by Def  $\overset{[\cdot]}{\rightsquigarrow}$ .

and the following fails  $\exists s. \exists M_1.M_1 \overset{[\mathbb{I}_0+y, \xi \cdot y : V_y]}{\rightsquigarrow} s \wedge \neg \exists M_0.M_0 \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} s$

iff  $\exists s. \exists M_1. \llbracket \mathbb{I}_0+y \rrbracket_{\xi \cdot y : V_y} \vdash M_1 : \text{Nm}$

$$\begin{array}{l} \wedge \hat{\mathfrak{a}}(M_1) = \emptyset \\ \wedge (\hat{\mathfrak{a}}(\xi \cdot y : V_y), M_1 \xi \cdot y : V_y) \Downarrow (\hat{\mathfrak{a}}(\xi \cdot y : V_y), G', s) \\ \wedge \neg \exists M_0. \llbracket \mathbb{I}_0 \rrbracket_{\xi} \vdash M_0 : \text{Nm} \\ \wedge \hat{\mathfrak{a}}(M_0) = \emptyset \\ \wedge (\hat{\mathfrak{a}}(\xi), M_1 \xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G', s) \end{array}$$

$$\begin{aligned}
& \text{iff } \exists s. \exists M_1(y). \llbracket \mathbb{I}_0 + y \rrbracket_{\xi \cdot y : V_y} \vdash M_1(y) : \text{Nm} \\
& \quad \wedge \hat{\mathfrak{a}}(M_1(y)) = \emptyset \\
& \quad \wedge (\hat{\mathfrak{a}}(\xi \cdot y : V_y), M_1(y)\xi \cdot y : V_y) \Downarrow (\hat{\mathfrak{a}}(\xi \cdot y : V_y), G', s) \\
& \wedge \neg \exists M_0. \llbracket \mathbb{I}_0 \rrbracket_{\xi} \vdash M_0 : \text{Nm} \\
& \quad \wedge \hat{\mathfrak{a}}(M_0) = \emptyset \\
& \quad \wedge (\hat{\mathfrak{a}}(\xi), M_1\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G', s) \\
& \text{iff } \exists s. \exists M_1(y). \llbracket \mathbb{I}_0 + y \rrbracket_{\xi \cdot y : V_y} \vdash M_1(y) : \text{Nm} \\
& \quad \wedge \hat{\mathfrak{a}}(M_1(y)) = \emptyset \\
& \quad \wedge (\hat{\mathfrak{a}}(\xi \cdot y : V_y), M_1(y)\xi \cdot y : V_y) \Downarrow (\hat{\mathfrak{a}}(\xi \cdot y : V_y), G', s) \\
& \wedge \neg \exists \text{let } y = M_y \text{ in } M_1(y). \llbracket \mathbb{I}_0 \rrbracket_{\xi} \vdash \text{let } y = M_y \text{ in } M_1(y) : \text{Nm} \\
& \quad \wedge \hat{\mathfrak{a}}(\text{let } y = M_y \text{ in } M_1(y)) = \emptyset \\
& \quad \wedge (\hat{\mathfrak{a}}(\xi), (\text{let } y = M_y \text{ in } M_1(y))\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G', s)
\end{aligned}$$

hence contradiction as the name derived by  $M_1(y)$  in the first evaluation, this same name can also be derived by  $\text{let } y = M_y \text{ in } M_1(y)$  in the second evaluation by the semantics of the evaluation of  $\text{let } y = M_y \text{ in } M_1(y)\xi \rightarrow M_1(V_y)\xi \equiv M_1(y)(\xi \cdot y : V_y)$ . Where  $M_y$  reduces to the term equal to  $V_y$  except the fresh names which have no affect on the name produced by  $M_0$  and  $M_1$  hence contradiction.

This works because  $s \in \hat{\mathfrak{a}}(\xi)$  hence the name cannot be fresh (derived from  $V_y$  and any fresh names in  $V_y$  can be replicated by a new generation via  $M_y$  and will be treated equally).

**Lemma 17 (Extensions cannot reveal old names).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}', \xi^{\mathbb{I}'}, s^{\text{Nm}}. \left( \begin{array}{l} \xi \preceq^* \xi' \\ \wedge s \in \hat{\mathfrak{a}}(\xi) \end{array} \right) \rightarrow \begin{array}{l} \exists M'. M' \overset{[\mathbb{I}', \xi']}{\rightsquigarrow} s \\ \leftrightarrow \\ \exists M. M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} s \end{array}$$

Essentially  $\xi'$  cannot reveal any names in  $\xi$  that are not already available to  $\xi$ .

*Proof.*  $\leftarrow$ : clearly holds

$\rightarrow$ : Proof by induction on the structure of  $\xi'$ :

$\xi' \equiv \xi$  then this clearly holds.

$\xi' \equiv \xi'_0 \cdot \delta : \mathbb{I}_0 \setminus \text{-TCV}$  then this holds as no new names are reachable by  $\delta$  and IH on  $\xi'_0$ .

$\xi' \equiv \xi'_0 \cdot y : V_y$  then  $\exists M_y. M_y \overset{[\mathbb{I}'_0, \xi'_0]}{\rightsquigarrow} V_y$  and using Lemma 16 and IH on  $\xi'_0$  this clearly holds.

In the next few lemmas we treat expressions as potential terms as each expression,  $c$ ,  $x$ ,  $\pi_i(e)$ , and  $\langle e, e \rangle$  is also a method of constructing a term and hence  $\llbracket e \rrbracket_{\xi} \equiv e\xi$  (closure) implies we can treat  $e$  as a term even though it is an expression, i.e. we write  $e$  even though we mean “the term constructed using the equivalent expression-constructors used to construct the expression  $e$ ”.

**Lemma 18 (Expressions cannot create new names).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, e. \mathbb{I} \Vdash e : \alpha \rightarrow \exists V. (\hat{\mathfrak{a}}(\xi), \llbracket e \rrbracket_{\xi}) \Downarrow (\hat{\mathfrak{a}}(\xi), V)$$



Clearly guaranteed termination implies  $\exists V'.(\hat{a}(\xi), \llbracket e \rrbracket_\xi) \Downarrow (\hat{a}(\xi), G', V')$ , however this Lemma proves that no new names are produced in such an evaluation.

*Proof.* By induction on structure of  $e \in \{c, x, \pi_i(e), \langle e, e \rangle\}$   
let  $G = \hat{a}(\xi)$

- $e = c$  constants then clearly  $(G, \llbracket e \rrbracket_\xi) \equiv (G, c) \Downarrow (G, c)$
- $e = x$  clearly  $(G, \llbracket e \rrbracket_\xi) \equiv (G, \xi(x)) \Downarrow (G, \xi(x))$  as  $\xi(x)$  is a value by definition.
- $e = \pi_i(e')$   
clearly  $(G, \llbracket \pi_i(e') \rrbracket_\xi) \equiv (G, \pi_i(\llbracket e' \rrbracket_\xi))$   
and by IH on  $e'$  then  $\exists V'.(G, \llbracket e' \rrbracket_\xi) \Downarrow (G, V')$  with  $V' : \alpha \times \beta$   
hence  $V' \equiv \langle V_1, V_2 \rangle$   
hence  $\pi_i(V_i)$  is a value and hence  $(G, \llbracket \pi_i(e) \rrbracket_\xi) \equiv (G, \pi_i(\llbracket e \rrbracket_\xi)) \Downarrow (G, V_i)$
- $e = \langle e_1, e_2 \rangle$   
by IH on  $e_1$  and  $e_2$ :  
 $\exists V_1.(G, \llbracket e_1 \rrbracket_\xi) \Downarrow (G, V_1)$   
and  $\exists V_2.(G, \llbracket e_2 \rrbracket_\xi) \Downarrow (G, V_2)$   
hence from operational semantics:  
 $\exists V (= \langle V_1, V_2 \rangle).(G, \llbracket \langle e_1, e_2 \rangle \rrbracket_\xi) \equiv (G, \langle \llbracket e_1 \rrbracket_\xi, \llbracket e_2 \rrbracket_\xi \rangle) \Downarrow (G, V)$

**Lemma 19 (Expressions are name free).**

$$\forall e. \mathbb{I} \Vdash e : \alpha \rightarrow \hat{a}(e) = \emptyset$$

*Proof.* By IH on the structure of  $e$  knowing that  $e \in \{c, x, \pi_i(e), \langle e_1, e_2 \rangle\}$ , i.e. no built in names, names only come from closing with a model via the semantics of  $\llbracket e \rrbracket_\xi$ .

**Lemma 20 (Expressions are fresh name free).**

$$\forall e. \mathbb{I} \Vdash e : \alpha \wedge e \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_e \rightarrow \hat{a}(V_e) \subseteq \hat{a}(\xi)$$

*Proof.* By IH on the structure of  $e$  knowing that  $e \in \{c, x, \pi_i(e), \langle e_1, e_2 \rangle\}$ , i.e. no built in names, names only come from closing with a model via the semantics of  $\llbracket e \rrbracket_\xi$ , and hence no evaluation of  $\text{gensym}()$  can occur, just producing old names.

**Lemma 21 (Expressions are congruent to their evaluation).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, e. \mathbb{I} \Vdash e : \alpha \wedge e \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_e \rightarrow e\xi \cong_{\alpha}^{\hat{a}(\xi)} V_e$$

*Proof.* By Lemma 20 we know all values contain names in the model.

By induction on the structure of  $e$ :

- $e \equiv c$  clearly holds.
- $e \equiv x$  clearly holds as  $x\xi \equiv \xi(x) \equiv V_e$

$e \equiv \pi_i(e')$  holds as by induction  $e'$ ,  
*i.e.*  $\mathbb{I} \Vdash e' : \alpha_1 \times \alpha_2 \wedge e' \xrightarrow{[\mathbb{I}, \xi]} \langle V_1, V_2 \rangle \rightarrow e' \xi \cong_{\alpha}^{\hat{a}(\xi)} \langle V_1, V_2 \rangle$   
 hence  $\pi_i(e') \xi \cong_{\alpha}^{\hat{a}(\xi)} V_i$   
 $e \equiv \langle e_1, e_2 \rangle$  holds by IH on  $e_1$  and  $e_2$   
*i.e.*  $\mathbb{I} \Vdash e_1 : \alpha_1 \wedge e_1 \xrightarrow{[\mathbb{I}, \xi]} V_1 \rightarrow e_1 \xi \cong_{\alpha}^{\hat{a}(\xi)} V_1$   
 and  $\mathbb{I} \Vdash e_2 : \alpha_2 \wedge e_2 \xrightarrow{[\mathbb{I}, \xi]} V_2 \rightarrow e_2 \xi \cong_{\alpha}^{\hat{a}(\xi)} V_2$   
 implies  $\langle e_1, e_2 \rangle \xi \cong_{\alpha}^{\hat{a}(\xi)} \langle V_1, V_2 \rangle$

**Lemma 22 (Names fresh in term imply name fresh in value).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, M, r. r \notin \hat{a}(M) \wedge (\hat{a}(M), r, G_0, M) \Downarrow (\hat{a}(\xi), r, G_0, G', V) \rightarrow r \notin \hat{a}(V)$$

*Proof.* By induction on the structure of  $M$ , most cases are trivial, the only non-trivial cases are:

$M \equiv M_1 M_2$ :

clearly  $M_1 \Downarrow V_1$  and  $M_2 \Downarrow V_2$  with  $r \notin V_1 V_2$  (by IH) hence 2 cases occur:

$V_1 \equiv \lambda x. M'_1$  then this evaluates to  $M'_1[V_2/x]$  which by assumptions and Op semantics  $r \notin M'_1[V_2/x]$

$V_1 \equiv \text{gensym}$  and  $V_2 \equiv ()$  then by Op. Sem. this generates a fresh name ( $\neq r$ ).

**Lemma 23 (Fresh names are underivable from model plus that name).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, r. r \notin \hat{a}(\xi) \rightarrow \neg \exists M^{\text{Nm}}. M \xrightarrow{[\mathbb{I}, \xi; m:r]} r$$

*Proof.* This holds as  $r$  must be a freshly generated name not appearing in  $\xi$  hence no term can be derived from the model  $\xi$  to generate such a name.

1 Assume:  $\mathbb{I}, \xi^{\mathbb{I}}$  s.t.

2 prove by contradiction:

*i.e.* assume  $\neg(r \notin \hat{a}(\xi) \rightarrow \neg \exists M^{\text{Nm}}. M \xrightarrow{[\mathbb{I}, \xi; m:r]} r)$  prove contradiction

3  $\leftrightarrow (r \notin \hat{a}(\xi) \wedge \exists M^{\text{Nm}}. M \xrightarrow{[\mathbb{I}, \xi; m:r]} r)$  FOL

4  $\leftrightarrow r \notin \hat{a}(\xi) \wedge \exists M^{\text{Nm}}. \hat{a}(M) = \emptyset \wedge \mathbb{I} \vdash M : \alpha$  Sem.  $\xrightarrow{[\cdot]}$   
 $\wedge (\hat{a}(\xi), r, M(\xi \cdot m : r)) \Downarrow (\hat{a}(\xi), r, G', r)$

5  $\leftrightarrow r \notin \hat{a}(\xi) \wedge \exists M^{\text{Nm}}. \hat{a}(M) = \emptyset \wedge \mathbb{I} \vdash M : \alpha$   $m \notin \text{dom}(\mathbb{I}) \wedge \mathbb{I} \vdash M : \alpha$   
 $\wedge (\hat{a}(\xi), r, M\xi) \Downarrow (\hat{a}(\xi), r, G', r)$

6  $\leftrightarrow r \notin \hat{a}(\xi) \wedge \exists M^{\text{Nm}}. \hat{a}(M) = \emptyset \wedge \mathbb{I} \vdash M : \alpha$   $r \notin \hat{a}(M\xi)$  and  
 $\wedge r \notin \hat{a}(M\xi)$  Lemma 22  
 $\wedge (\hat{a}(\xi), r, M\xi) \Downarrow (\hat{a}(\xi), r, G', r)$  imply contradiction

7  $\leftrightarrow r \notin \hat{a}(\xi) \rightarrow \neg \exists M^{\text{Nm}}. M \xrightarrow{[\mathbb{I}, \xi; m:r]} r$

**Lemma 24 (Extensions give equal semantics of expressions).**

$$\forall \mathbb{I}_1, \xi_1^{\mathbb{I}_1}, \mathbb{I}_2, \xi_2^{\mathbb{I}_2}, e. \xi_1 \preceq^* \xi_2 \wedge \mathbb{I}_1 \Vdash e : \alpha \rightarrow [e]_{\xi_1} \equiv [e]_{\xi_2}$$

*Proof.* Prove by induction on the structure of  $e$  using semantics of expressions:

$$\begin{aligned}
 e \equiv c: & \llbracket c \rrbracket_{\xi_1} \equiv c \equiv \llbracket c \rrbracket_{\xi_2} \\
 e \equiv x: & x \in \mathbb{I}_1 \text{ implies } (Def \preceq^*) \llbracket x \rrbracket_{\xi_1} \equiv \xi_1(x) \equiv \xi_2(x) \equiv \llbracket c \rrbracket_{\xi_2} \\
 e \equiv (e_1, e_2): & \text{ by IH on both } e_1 \text{ and } e_2 \text{ this clearly holds.} \\
 e \equiv \pi_i(e'): & \text{ by IH on } e' \text{ this clearly holds.}
 \end{aligned}$$

**Lemma 25 (Sub LTC's implies subtype-contexts).**

$$\forall \mathbb{I}, \mathbb{I}_0, \xi^{\mathbb{I}}. \mathbb{I} \Vdash \mathbb{I}_0 \rightarrow \llbracket \mathbb{I}_0 \rrbracket_{\xi} \subseteq \llbracket \mathbb{I} \rrbracket_{\xi}$$

*Proof:* clearly holds through induction on structure of  $\mathbb{I}$  and  $\mathbb{I}_0$  and using the typing rules in Fig. 1.

**Lemma 26 (Values derivable from subset of LTC is also derivable from superset of LTC).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}_0 + \mathbb{I}_1, M. \mathbb{I} \Vdash \mathbb{I}_0 \wedge \mathbb{I} \Vdash \mathbb{I}_0 + \mathbb{I}_1 \wedge M \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \rightarrow M \overset{[\mathbb{I}_0 + \mathbb{I}_1, \xi]}{\rightsquigarrow} V$$

*Proof.* By Lemma 25,  
 $\mathbb{I} \Vdash \mathbb{I}_0 \wedge \mathbb{I} \Vdash \mathbb{I}_0 + \mathbb{I}_1$   
 $\rightarrow \mathbb{I}_0 + \mathbb{I}_1 \Vdash \mathbb{I}_0$   
 $\rightarrow \llbracket \mathbb{I}_0 \rrbracket_{\xi} \subseteq \llbracket \mathbb{I}_0 + \mathbb{I}_1 \rrbracket_{\xi}$   
 hence this must hold.

**Lemma 27 (Values derived from  $\mathbb{I}$  maintain extension (single) when added to models which are extensions).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}', \xi^{\mathbb{I}'}, M^\alpha. \xi \preceq \xi' \wedge M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V \wedge \hat{a}(V) \cap \hat{a}(\xi') \subseteq \hat{a}(\xi) \rightarrow \xi \cdot x : V \preceq \xi' \cdot x : V$$

*Proof.*

- |   |  |   |
|---|--|---|
| 1 | Assume: $\mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}', \xi^{\mathbb{I}'}, M^\alpha$ s.t. $\xi \preceq \xi' \wedge M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V$  |   |
| 2 | The case for $\xi' \equiv \xi \cdot \delta : \mathbb{I}_0 \setminus_{-TCV}$ is proven below:   |   |
| 3 | assume some $\mathbb{I}_0$ s.t. $\mathbb{I} \Vdash \mathbb{I}_0$ with $\xi \preceq \xi \cdot \delta : \mathbb{I}_0 \setminus_{-TCV}$<br>then $\mathbb{I} + x \Vdash \mathbb{I}_0$ and hence $\xi \cdot x : V \preceq \xi \cdot \delta : \mathbb{I}_0 \setminus_{-TCV} \cdot x : V$ | Sem. $\preceq$  |
| 4 | The case for $\xi' \equiv \xi \cdot y : V'$ is proven below:   |   |
| 5 | $(\xi \preceq \xi' \wedge M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V) \rightarrow \exists M'. M' \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V' \wedge \xi' \equiv \xi \cdot y' : V' \wedge M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V$                            | Sem. $\preceq$  |
| 6 | $\rightarrow \exists M'. M' \overset{[\mathbb{I}, \xi \cdot x : V]}{\rightsquigarrow} V' \wedge \xi' \cdot x : V \equiv \xi \cdot x : V \cdot y' : V'$   | Lemma 11 ( $\hat{a}(V) \cap \hat{a}(V') \subseteq \hat{a}(\xi)$ ) |
| 7 | $\rightarrow \exists M'. M' \overset{[\mathbb{I} + x : \alpha_x, \xi \cdot x : V]}{\rightsquigarrow} V' \wedge \xi' \cdot x : V \equiv \xi \cdot x : V \cdot y' : V'$  | Lemma 26  |
| 8 | $\rightarrow \xi \cdot x : V \preceq \xi' \cdot x : V$   | Sem. $\preceq$  |

**Lemma 28 (Values derived from  $\mathbb{I}$  maintain extension (star) when added to models which are extensions).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}', \xi'^{\mathbb{I}'}, M. \xi \preceq^* \xi' \wedge M \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V \wedge \tilde{a}(V) \cap \tilde{a}(\xi') \subseteq \tilde{a}(\xi) \rightarrow \xi \cdot x : V \preceq^* \xi' \cdot x : V$$

*Proof.* Use Lemma 27 with Def  $\preceq^*$  to see this clearly holds.

**Lemma 29 (Names produced from an extension can be added to an extension and still hold).**

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}', \xi'^{\mathbb{I}'}, M, r. M \stackrel{[\mathbb{I}', \xi']}{\rightsquigarrow} r \wedge \xi \preceq^* \xi' \rightarrow \xi \cdot x : r \preceq^* \xi' \cdot x : r$$

*Proof.* Assume a sequence of terms  $[\tilde{z} : \tilde{N}]$  that derive the (non-LTC) values of  $\xi'$  (the LTC are trivial), then three cases exist:

- $r \in \tilde{a}(\xi)$  then clearly this holds as the same sequence of terms  $[\tilde{z} : \tilde{N}]$  derives  $\xi' \cdot x : r$  with the new LTC  $\mathbb{I} + x : \text{Nm}$ . Hence  $\xi \cdot x : r \preceq^* \xi' \cdot x : r$
- $r \notin \tilde{a}(\xi) \wedge r \notin \tilde{a}(\xi')$  then clearly  $r$  is fresh and the same reasoning as above holds.
- $r \notin \tilde{a}(\xi) \wedge r \in \tilde{a}(\xi')$  then  $r$  is derived from  $\xi'$  but not from  $\xi$  and hence the sequence  $[\tilde{z} : \tilde{N}[x/\text{gensym}()]]$  also derives  $\xi'$  proves  $\xi \cdot x : r \preceq^* \xi' \cdot x : r$  where we write  $\tilde{N}[x/\text{gensym}()]$  for the term  $\tilde{N}_i$  that initially produce  $r$  to substitute  $x$  for that  $\text{gensym}()$  which clearly must produce the same results. All other fresh names are irrelevant.

**Lemma 30 (Extending the model by  $\delta$  maintains models).** (*A-Ext-Ind not required*)

$$\forall \mathbb{I}, \xi^{\mathbb{I}}, \mathbb{I}', \xi'^{\mathbb{I}'}, \mathbb{I}_a. \mathbb{I} \Vdash \mathbb{I}_a \rightarrow (\xi \cdot \xi' \models A^{-\delta} \leftrightarrow \xi \cdot \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi' \models A^{-\delta})$$

where  $A^{-\delta}$  means  $\delta$  does not occur syntactically in  $A$ .

*Proof.* By definition of  $\preceq$ ,  $\xi \preceq \xi \cdot \delta : \mathbb{I}_a \setminus \text{-TCV}$  holds and implies via Def  $\preceq^*$  that  $\xi \cdot \xi' \preceq^* \xi \cdot \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi'$  which allows the use of Lemma 24 and Lemma 11 ( $\tilde{a}(\xi \cdot \xi') \equiv \tilde{a}(\xi \cdot \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi')$ ).

The proof follows by IH on structure of  $A$ :

- $e_1 = e_2$  proven by noting that  $\llbracket e_i \rrbracket_{\xi, \xi'} \equiv \llbracket e_i \rrbracket_{\xi, \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi'}$  (Lemma 24).
- $A_1 \wedge A_2, A_1 \vee A_2, A_1 \rightarrow A_2$ : IH on  $A_1, A_2$ .
- $\neg A_1$ , proof by IH on  $A_1$ , assume  $\xi \cdot \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi' \models \neg A_1$  then clearly  $\xi \cdot \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi' \not\models A_1$  and by IH on  $A_1$  then  $\xi \cdot \xi' \not\models A_1$  i.e.  $\xi \cdot \xi' \models \neg A_1$ .
- $x \# \mathbb{I}'$  holds by Lemma 24 and Lemma 11 ( $\tilde{a}(\xi \cdot \xi') \equiv \tilde{a}(\xi \cdot \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi')$ ) which ensure  $\llbracket x \rrbracket_{\xi, \xi'} \equiv \llbracket x \rrbracket_{\xi, \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi'}$  and
  - $\exists M_x. M_x \stackrel{[\mathbb{I}', \xi, \xi']}{\rightsquigarrow} V \leftrightarrow \exists M_x. M_x \stackrel{[\mathbb{I}', \xi, \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi']}{\rightsquigarrow} V$  respectively, hence clearly
  - $\neg \exists M_x. M_x \stackrel{[\mathbb{I}', \xi, \xi']}{\rightsquigarrow} \llbracket x \rrbracket_{\xi, \xi'} \leftrightarrow \neg \exists M_x. M_x \stackrel{[\mathbb{I}', \xi, \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi']}{\rightsquigarrow} \llbracket x \rrbracket_{\xi, \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi'}$ .
- $u \bullet e = m\{A_1\}$  by IH on  $A_1$  with  $\llbracket u \rrbracket_{\xi, \xi'} \equiv \llbracket u \rrbracket_{\xi, \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi'}$  and  $\llbracket e \rrbracket_{\xi, \xi'} \equiv \llbracket e \rrbracket_{\xi, \delta : \mathbb{I}_a \setminus \text{-TCV} \cdot \xi'}$  (Lemma 24) hence evaluation is equivalent and this holds.

- $\forall x \in (\mathbb{I}').A_1$  knowing that  $\delta \notin \mathbb{I}'$  then the same possible terms are quantified over given Lemma 11 ( $M_x \stackrel{[\mathbb{I}', \xi, \xi']}{\rightsquigarrow} V \leftrightarrow M_x \stackrel{[\mathbb{I}', \xi, \delta: \mathbb{I}'_a \setminus \text{TCV} \cdot \xi']}{\rightsquigarrow} V$ ) and by IH on  $A_1$  the case holds.
- $\forall \delta'.A_1$  by IH on  $A_1$  this holds for any model containing  $\delta$  and  $\delta'$  then by the semantics the lemma holds.

The following 2 definitions will be used in some of the proofs using equality, but are not required otherwise.

**Definition 2 (Similar extensions (single)).** *Two model extensions are congruent model extensions (single) iff there exists an equivalent derivation for both models:*

$$\begin{aligned}
 & \forall \mathbb{I}'_i, \xi_1^{\mathbb{I}'_i}, \xi_2^{\mathbb{I}'_i}, \mathbb{I}'_i, \xi_1^{\mathbb{I}'_i}, \xi_2^{\mathbb{I}'_i}. \\
 & (\xi_1 \preceq \xi'_1) \sim^{\preceq} (\xi_2 \preceq \xi'_2) \leftrightarrow \mathbb{I}'_i \equiv \mathbb{I}_i + \delta_j \wedge \xi'_1(\delta_j) \equiv \xi'_2(\delta_j) \\
 & \quad \quad \quad \downarrow \\
 & \mathbb{I}'_i \equiv \mathbb{I}_i + x_j : \alpha_j \wedge \exists M_j^{\alpha_j}. M_j \stackrel{[\mathbb{I}'_i, \xi_1]}{\rightsquigarrow} \xi'_1(x_j) \wedge M_j \stackrel{[\mathbb{I}'_i, \xi_2]}{\rightsquigarrow} \xi'_2(x_j)
 \end{aligned}$$

**Definition 3 (Similar extensions (star)).** *Two model extensions are similarly derived extensions (star) iff there exists an equivalent derivation for both models:*

$$\begin{aligned}
 & \forall \mathbb{I}'_i, \xi_1^{\mathbb{I}'_i}, \xi_2^{\mathbb{I}'_i}, \mathbb{I}'_i, \xi_1^{\mathbb{I}'_i}, \xi_2^{\mathbb{I}'_i}. \\
 & (\xi_1 \preceq^* \xi'_1) \sim^{\preceq^*} (\xi_2 \preceq^* \xi'_2) \leftrightarrow \xi_1 \equiv \xi'_1 \wedge \xi_2 \equiv \xi'_2 \\
 & \quad \quad \quad \downarrow \\
 & (\xi_1 \preceq \xi'_1) \sim^{\preceq} (\xi_2 \preceq \xi'_2) \\
 & \quad \quad \quad \downarrow \\
 & \exists \xi_{1j}^{\mathbb{I}'_j}, \xi_{2j}^{\mathbb{I}'_j}. (\xi_1 \preceq^* \xi_{1j}) \sim^{\preceq^*} (\xi_2 \preceq^* \xi_{2j}) \\
 & \quad \quad \quad \wedge (\xi_{1j} \preceq^* \xi'_1) \sim^{\preceq^*} (\xi_{2j} \preceq^* \xi'_2)
 \end{aligned}$$

**Lemma 31 (Congruent extensions implies their evaluation added to a model, model equivalence equivalently).**

$$\begin{aligned}
 & \forall \mathbb{I}', \xi^{\mathbb{I}'}, M_1^\alpha, M_2^\alpha. \\
 & M_1 \xi \cong_\alpha^{\tilde{a}(\xi)} M_2 \xi \wedge M_1 \stackrel{[\mathbb{I}', \xi]}{\rightsquigarrow} V_1 \wedge M_2 \stackrel{[\mathbb{I}', \xi]}{\rightsquigarrow} V_2 \\
 & \rightarrow \\
 & \forall \mathbb{I}'_i, \xi_1^{\mathbb{I}'_i}, \xi_2^{\mathbb{I}'_i}, e = e'. (\xi \cdot x : V_1 \preceq^* \xi'_1) \sim^{\preceq^*} (\xi \cdot x : V_2 \preceq^* \xi'_2) \wedge \mathbb{I}'_i \Vdash e = e' \\
 & \rightarrow \\
 & \xi'_1 \models e = e' \leftrightarrow \xi'_2 \models e = e'
 \end{aligned}$$

As an example consider  $M_1 \equiv \text{gensym}() \equiv M_2$  clearly any use of the names produced must be indistinguishable in any future similarly derived models as the names can be switched asimilar to  $\alpha$ -equivalence.

*Proof.* Required to prove  $\llbracket e \rrbracket_{\xi'_1} \cong_{\alpha}^{\tilde{a}(\xi'_1)} \llbracket e' \rrbracket_{\xi'_1} \rightarrow \llbracket e \rrbracket_{\xi'_2} \cong_{\alpha}^{\tilde{a}(\xi'_2)} \llbracket e' \rrbracket_{\xi'_2}$

1	Making the relevant assumptions about $\mathbb{I}, \xi^{\mathbb{I}}, M_1, M_2, \mathbb{I}_i, \xi_1^{\mathbb{I}_i}, \xi_2^{\mathbb{I}_i}, e = e', s.t. \dots$	
2	$M_1 \xi \cong_{\alpha}^{\tilde{a}(\xi)} M_2 \xi$	
3	$\Leftrightarrow \forall C[\cdot]^{\alpha}, b^{\text{Bool}}. \left( \begin{array}{l} \emptyset \vdash C[\cdot]^{\alpha} : \text{Bool} \\ \wedge \tilde{a}(C[\cdot]) \subseteq \tilde{a}(\xi) \end{array} \right) \rightarrow \left( \begin{array}{l} (\tilde{a}(\xi), C[M_1 \xi]) \Downarrow (\tilde{a}(\xi), G_1, b) \\ \Leftrightarrow \\ (\tilde{a}(\xi), C[M_2 \xi]) \Downarrow (\tilde{a}(\xi), G_2, b) \end{array} \right)$	Sem. $\cong$
4	Let $\xi'_j[\cdot]$ be derived from the terms as follows $\tilde{z} : \tilde{N}$ and $x : V_i$ noting that $\tilde{N}_{i+1}$ may contain the variable $\tilde{z}_i$ and ignoring the inconsequential TCV-parts, then if $\mathbb{I}_i \Vdash e : \alpha_e$ a subset of $C[\cdot]$ can be set for any $C'[\cdot]^{\alpha_e}$ as follows:	
5	$\rightarrow \forall C[\cdot] \equiv (\text{let } x = [\cdot]_{\emptyset}^{\alpha} \text{ in let } \tilde{z} = \tilde{N}\xi \text{ in } C'[e\xi] = C'[e'\xi]), b^{\text{Bool}}.$ $\emptyset \vdash C[\cdot]^{\alpha} : \text{Bool} \wedge \tilde{a}(C[\cdot]) \subseteq \tilde{a}(\xi)$ $\rightarrow$ $\left( \begin{array}{l} (\tilde{a}(\xi), \text{let } x = [M_1 \xi]_{\emptyset}^{\alpha} \text{ in let } \tilde{z} = \tilde{N}\xi \text{ in } C'[e\xi] = C'[e'\xi]) \Downarrow (\tilde{a}(\xi), G_1, b) \\ \Leftrightarrow \\ (\tilde{a}(\xi), \text{let } x = [M_2 \xi]_{\emptyset}^{\alpha} \text{ in let } \tilde{z} = \tilde{N}\xi \text{ in } C'[e\xi] = C'[e'\xi]) \Downarrow (\tilde{a}(\xi), G_2, b) \end{array} \right)$	Subset $C[\cdot] \equiv \dots$ $C'[\cdot]$ -capture-avoiding variables in $\text{dom}(\mathbb{I}_i)$
6	$\rightarrow \forall b^{\text{Bool}}. \left( \begin{array}{l} \forall C[\cdot] \equiv (\text{let } x = [\cdot]_{\emptyset}^{\alpha} \text{ in let } \tilde{z} = \tilde{N}\xi \text{ in } C'[e\xi] = C'[e'\xi]), b^{\text{Bool}}. \\ \emptyset \vdash C[\cdot] : \text{Bool} \wedge \tilde{a}(C[\cdot]) \subseteq \tilde{a}(\xi) \\ \rightarrow (\tilde{a}(\xi), C[M_1 \xi]) \Downarrow (\tilde{a}(\xi), G_1, b) \end{array} \right) \rightarrow (\forall x. A(x) \rightarrow (B(x) \Leftrightarrow C(x)))$ $\Leftrightarrow$ $\left( \begin{array}{l} \forall C[\cdot] \equiv (\text{let } x = [\cdot]_{\emptyset}^{\alpha} \text{ in let } \tilde{z} = \tilde{N}\xi \text{ in } C'[e\xi] = C'[e'\xi]), b^{\text{Bool}}. \\ \emptyset \vdash C[\cdot] : \text{Bool} \wedge \tilde{a}(C[\cdot]) \subseteq \tilde{a}(\xi) \\ \rightarrow (\tilde{a}(\xi), C[M_2 \xi]) \Downarrow (\tilde{a}(\xi), G_2, b) \end{array} \right) \rightarrow (\forall x. A(x) \rightarrow B(x))$ $\Leftrightarrow (\forall x. A(x) \rightarrow C(x))$ FOL	
7	$\rightarrow \forall b^{\text{Bool}}. \left( \begin{array}{l} \forall \text{let } \tilde{z} = \tilde{N} \text{ in } C'[e\xi_1] = C'[e'\xi_1]. \\ \left( \begin{array}{l} \emptyset \vdash \text{let } \tilde{z} = \tilde{N}\xi_1 \text{ in } C'[e\xi_1] = C'[e'\xi_1] : \text{Bool} \\ \wedge \tilde{a}(\text{let } \tilde{z} = \tilde{N}\xi_1 \text{ in } C'[e\xi_1] = C'[e'\xi_1]) \subseteq \tilde{a}(\xi_1) \end{array} \right) \\ \rightarrow (\tilde{a}(\xi_1), \text{let } \tilde{z} = \tilde{N}\xi_1 \text{ in } C'[e\xi_1] = C'[e'\xi_1]) \Downarrow (\tilde{a}(\xi_1), G_1, b) \end{array} \right) \rightarrow$ $\Leftrightarrow$ $\left( \begin{array}{l} \forall \text{let } \tilde{z} = \tilde{N}\xi_2 \text{ in } C'[e\xi_2] = C'[e'\xi_2]. \\ \left( \begin{array}{l} \emptyset \vdash \text{let } \tilde{z} = \tilde{N} \text{ in } C'[e\xi_2] = C'[e'\xi_2] : \text{Bool} \\ \wedge \tilde{a}(\text{let } \tilde{z} = \tilde{N}\xi_2 \text{ in } C'[e\xi_2] = C'[e'\xi_2]) \subseteq \tilde{a}(\xi_2) \end{array} \right) \\ \rightarrow (\tilde{a}(\xi_2), \text{let } \tilde{z} = \tilde{N}\xi_2 \text{ in } C'[e\xi_2] = C'[e'\xi_2]) \Downarrow (\tilde{a}(\xi_2), G_2, b) \end{array} \right) \rightarrow$	$M_j \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_j$ then $\xi_j \equiv \xi \cdot x : V_j$ ( $j = 1, 2$ ) let $x = M_j$ in $M\xi \rightarrow M\xi_j$
8	$\rightarrow \forall b^{\text{Bool}}. \left( \begin{array}{l} \forall C'[\cdot]^{\alpha}. \left( \begin{array}{l} \emptyset \vdash C'[\cdot]^{\alpha_e} : \text{Bool} \\ \wedge \tilde{a}(C'[\cdot]) \subseteq \tilde{a}(\xi'_1) \end{array} \right) \\ \rightarrow (\tilde{a}(\xi'_1), C'[e\xi'_1] = C'[e'\xi'_1]) \Downarrow (\tilde{a}(\xi'_1), G_1, b) \end{array} \right) \rightarrow$ $\Leftrightarrow$ $\left( \begin{array}{l} \forall C'[\cdot]^{\alpha_e}. \left( \begin{array}{l} \emptyset \vdash C'[\cdot]^{\alpha_e} : \text{Bool} \\ \wedge \tilde{a}(C'[\cdot]) \subseteq \tilde{a}(\xi'_2) \end{array} \right) \\ \rightarrow (\tilde{a}(\xi'_2), C'[e\xi'_2] = C'[e'\xi'_2]) \Downarrow (\tilde{a}(\xi'_2), G_2, b) \end{array} \right) \rightarrow$	let $\tilde{z} = \tilde{N}\xi_j$ in $M\xi_j \rightarrow M\xi'_j$ $C[M_j \xi] \rightarrow C'[e\xi_j]^{\alpha} = C'[e'\xi_j]^{\alpha}$
9	$\rightarrow \left( \begin{array}{l} \forall C'[\cdot]^{\alpha_e}, b^{\text{Bool}}. \left( \begin{array}{l} \emptyset \vdash C'[\cdot]^{\alpha_e} : \text{Bool} \\ \wedge \tilde{a}(C'[\cdot]) \subseteq \tilde{a}(\xi'_1) \end{array} \right) \\ \rightarrow \left( \begin{array}{l} (\tilde{a}(\xi'_1), C'[e\xi'_1]) \Downarrow (\tilde{a}(\xi'_1), G_1, b') \\ \Leftrightarrow (\tilde{a}(\xi'_1), C'[e'\xi'_1]) \Downarrow (\tilde{a}(\xi'_1), G_1, b') \end{array} \right) \end{array} \right) \rightarrow$ $\Leftrightarrow$ $\left( \begin{array}{l} \forall C'[\cdot]^{\alpha_e}, b^{\text{Bool}}. \left( \begin{array}{l} \emptyset \vdash C'[\cdot]^{\alpha_e} : \text{Bool} \\ \wedge \tilde{a}(C'[\cdot]) \subseteq \tilde{a}(\xi'_2) \end{array} \right) \\ \rightarrow \left( \begin{array}{l} (\tilde{a}(\xi'_2), C'[e\xi'_2]) \Downarrow (\tilde{a}(\xi'_2), G_1, b') \\ \Leftrightarrow (\tilde{a}(\xi'_2), C'[e'\xi'_2]) \Downarrow (\tilde{a}(\xi'_2), G_1, b') \end{array} \right) \end{array} \right) \rightarrow$	Hence given $C'[e\xi'_i]^{\alpha_e} : \text{Bool}$ this implies: (To think about this set $b \equiv \text{true}$ or $b = \text{false}$ then clearly this holds) $b = \text{true}$ : implies the $=$ case of $\Leftrightarrow$ $b = \text{false}$ : implies the $\neq$ case of $\Leftrightarrow$
10	$\xi'_1 \models e = e' \Leftrightarrow \xi'_2 \models e = e'$	Sem. $=$

**Lemma 32 (Congruent extensions implies their evaluation added to a model, model equivalently).**

$$\begin{aligned}
 & \forall \Gamma, \xi^T, M_1^\alpha, M_2^\alpha. \\
 & M_1 \xi \cong_a^{\dot{\beta}(\xi)} M_2 \xi \wedge M_1 \xrightarrow{[\Gamma, \xi]} V_1 \wedge M_2 \xrightarrow{[\Gamma, \xi]} V_2 \\
 & \rightarrow \\
 & \forall \Gamma_i, \xi_1^{\Gamma_i}, \xi_2^{\Gamma_i}, A. (\xi \cdot x : V_1 \preceq^* \xi_1') \sim^{\preceq^*} (\xi \cdot x : V_2 \preceq^* \xi_2') \wedge \Gamma_i \Vdash A \\
 & \rightarrow \\
 & \forall \Gamma_0. \Gamma_i \Vdash \Gamma_0 \rightarrow (\exists M_u. M_u \xrightarrow{[\Gamma_0, \xi_1']} \llbracket u \rrbracket_{\xi_1'} \leftrightarrow \exists N_u. N_u \xrightarrow{[\Gamma_0, \xi_2']} \llbracket u \rrbracket_{\xi_2'})
 \end{aligned}$$

If  $u \in \text{dom}(\xi)$  then clearly this holds.  
 If  $u \notin \text{dom}(\xi)$  then clearly the  $u$  must be in  $\xi_j'$  and hence is derived from a series of extensions, which are characterised by  $\tilde{z} : \tilde{N}$  therefore the same use in  $M_u$  and  $N_u$  of these variables derives the same  $\llbracket u \rrbracket_{\xi_j'}$ . This relies on the fact that if a name is used in  $\xi_1'$  then it is the same name used in  $\xi_2'$  or it is fresh from  $\xi$  in both.

*Proof.* This is proven using the fact that  $\xi_1'$  and  $\xi_2'$  are similar and hence any  $M_u$  that produces  $\llbracket u \rrbracket_{\xi_1'}$  can be used to derive  $\llbracket u \rrbracket_{\xi_2'}$  as follows:

1	Let the $\sim^{\approx^*}$ -extension part be derived from by $\tilde{z} : \tilde{M}$ where $\tilde{M}_{i+1}$ may contain the variable $\tilde{z}_i$ .	
2	Assume some $\mathbb{I}, \xi^{\mathbb{I}}, M_1^\alpha, M_2^\alpha$ s.t. $M_1\xi \cong_\alpha^{\tilde{a}(\xi)} M_2\xi \wedge M_1 \stackrel{[\mathbb{I}; \xi]}{\rightsquigarrow} V_1 \wedge M_2 \stackrel{[\mathbb{I}; \xi]}{\rightsquigarrow} V_2$	
3	also assume some $\mathbb{I}_i, \xi_1^{\mathbb{I}_i}, \xi_2^{\mathbb{I}_i}, A$ s.t. $(\xi \cdot x : V_1 \preceq^* \xi_1') \sim^{\approx^*} (\xi \cdot x : V_2 \preceq^* \xi_2') \wedge \mathbb{I}_i \Vdash A$	
4	also assume some $\mathbb{I}_0$ s.t. $\mathbb{I}_i \Vdash \mathbb{I}_0$ , prove $\rightarrow$ to prove $\leftrightarrow$ -by symmetry	
5	Assume there exists $M_u$ s.t. $M_u \stackrel{[\mathbb{I}_0, \xi_1']}{\rightsquigarrow} \llbracket u \rrbracket_{\xi_1'}$	
6	then show that there exists $N_u$ s.t. $N_u \stackrel{[\mathbb{I}_0, \xi_2']}{\rightsquigarrow} \llbracket u \rrbracket_{\xi_2'}$	
7	$M_1\xi \cong_\alpha^{\tilde{a}(\xi)} M_2\xi$	
8	$\leftrightarrow \forall C[\cdot]^\alpha, b^{\text{Bool}}. \left( \begin{array}{l} \emptyset \vdash C[\cdot]^\alpha : \text{Bool} \\ \wedge \tilde{a}(C[\cdot]) \subseteq \tilde{a}(\xi) \end{array} \right) \rightarrow \left( \begin{array}{l} (\tilde{a}(\xi), C[M_1\xi]) \Downarrow (\tilde{a}(\xi), G_1, b) \\ \leftrightarrow \\ (\tilde{a}(\xi), C[M_2\xi]) \Downarrow (\tilde{a}(\xi), G_2, b) \end{array} \right)$	Sem. $\cong$
9	$\forall b^{\text{Bool}}. \forall C[\cdot]^\alpha. \left( \begin{array}{l} \emptyset \vdash C[\cdot]^\alpha : \text{Bool} \\ \wedge \tilde{a}(C[\cdot]) \subseteq \tilde{a}(\xi) \end{array} \right) \rightarrow (\tilde{a}(\xi), C[M_1\xi]) \Downarrow (\tilde{a}(\xi), G_1, b)$ $\leftrightarrow$ $\forall C[\cdot]^\alpha. \left( \begin{array}{l} \emptyset \vdash C[\cdot]^\alpha : \text{Bool} \\ \wedge \tilde{a}(C[\cdot]) \subseteq \tilde{a}(\xi) \end{array} \right) \rightarrow (\tilde{a}(\xi), C[M_2\xi]) \Downarrow (\tilde{a}(\xi), G_2, b)$	$(\forall x. A(x) \rightarrow (B(x) \leftrightarrow C(x))) \rightarrow$ $((\forall x. A(x) \rightarrow B(x)))$ $\leftrightarrow (\forall x. A(x) \rightarrow C(x))$
10	Assume $\xi_1'$ and $\xi_2'$ are both derived from $\xi$ via $\tilde{z} : \tilde{N}$ and $x : M_j$ (from $\sim^{\approx^*}$ ) Select $C[\cdot]$ as let $x = [\cdot]^\alpha$ in let $\tilde{z} = \tilde{N}\xi$ in $(C'[M_u\xi]^{\text{Nm}} = C'[u\xi]^{\text{Nm}})$ with $C'[\cdot]$ not capturing $x, \tilde{z}$ $M_j \stackrel{[\mathbb{I}; \xi]}{\rightsquigarrow} V_j$ implies $C[M_j]\xi \rightarrow \text{let } \tilde{z} = \tilde{N}(\xi \cdot x : V_j) \text{ in } C'[M_u(\xi \cdot x : V_j)]^{\text{Nm}} = C'[u(\xi \cdot x : V_j)]^{\text{Nm}}$ $\dots \rightarrow C'[M_u\xi_1']^{\text{Nm}} = C'[u\xi_1']^{\text{Nm}}$	
11	$\rightarrow \forall b^{\text{Bool}}. \left( \begin{array}{l} \emptyset \vdash C'[M_u\xi_1']^{\text{Nm}} = C'[u\xi_1']^{\text{Nm}} : \alpha \\ \wedge \tilde{a}(C'[M_u\xi_1']^{\text{Nm}} = C'[u\xi_1']^{\text{Nm}}) \subseteq \tilde{a}(\xi_1') \end{array} \right) \rightarrow (\tilde{a}(\xi_1'), (C'[M_u\xi_1'] = C'[u\xi_1']) \Downarrow (\tilde{a}(\xi_1'), G_1, b)$ $\leftrightarrow$ $\forall C'[M_u\xi_2']^{\text{Nm}} = C'[u\xi_2']^{\text{Nm}}. \left( \begin{array}{l} \emptyset \vdash C'[M_u\xi_2']^{\text{Nm}} = C'[u\xi_2']^{\text{Nm}} : \alpha \\ \wedge \tilde{a}(C'[M_u\xi_2']^{\text{Nm}} = C'[u\xi_2']^{\text{Nm}}) \subseteq \tilde{a}(\xi_2') \end{array} \right) \rightarrow (\tilde{a}(\xi_2'), (C'[M_u\xi_2'] = C'[u\xi_2']) \Downarrow (\tilde{a}(\xi_2'), G_1, b)$	Subset of $C[\cdot]$ from line 10
12	$\rightarrow \forall C'[\cdot]^{\text{Nm}}, b'^{\text{Bool}}. \left( \begin{array}{l} \emptyset \vdash C'[\cdot]^{\text{Nm}} : \text{Bool} \\ \wedge \tilde{a}(C'[\cdot]^{\text{Nm}}) \subseteq \tilde{a}(\xi_1') \end{array} \right) \rightarrow \left( \begin{array}{l} (\tilde{a}(\xi_1'), C'[M_u\xi_1']) \Downarrow (\tilde{a}(\xi_1'), G_2, b') \\ \leftrightarrow \\ (\tilde{a}(\xi_1'), C'[u\xi_1']) \Downarrow (\tilde{a}(\xi_1'), G_2, b') \end{array} \right)$ $\leftrightarrow$ $\forall C'[\cdot]^{\text{Nm}}, b'^{\text{Bool}}. \left( \begin{array}{l} \emptyset \vdash C'[\cdot]^{\text{Nm}} : \text{Bool} \\ \wedge \tilde{a}(C'[\cdot]^{\text{Nm}}) \subseteq \tilde{a}(\xi_2') \end{array} \right) \rightarrow \left( \begin{array}{l} (\tilde{a}(\xi_2'), C'[M_u\xi_2']) \Downarrow (\tilde{a}(\xi_2'), G_2, b') \\ \leftrightarrow \\ (\tilde{a}(\xi_2'), C'[u\xi_2']) \Downarrow (\tilde{a}(\xi_2'), G_2, b') \end{array} \right)$	$b = \text{true}$ : clearly holds. $b = \text{false}$ : holds as $(C'[M_u\xi_j'] = C'[u\xi_j']) \Downarrow \text{false}$ $\leftrightarrow$ $C'[M_u\xi_j'] \Downarrow b' \leftrightarrow C'[u\xi_j'] \Downarrow \neg b'$ implies $(\forall C'_{M_u, u}[\cdot]. \text{T} \rightarrow \text{F}) \leftrightarrow (\forall C'_{M_u, u}[\cdot]. \text{T} \rightarrow \text{F})$ and hence the more general $\text{F} \leftrightarrow \text{F}$
13	$\rightarrow (\exists M_u. M_u \stackrel{[\mathbb{I}_0, \xi_1']}{\rightsquigarrow} \llbracket u \rrbracket_{\xi_1'} \leftrightarrow \exists N_u. N_u \stackrel{[\mathbb{I}_0, \xi_2']}{\rightsquigarrow} \llbracket u \rrbracket_{\xi_2'})$	with precisely the same $M_u \equiv N_u$ .



**Lemma 33 (Congruent extensions implies their evaluation added to a model, model equivalently).**

$$\begin{aligned}
 & \forall \mathbb{I}, \xi^{\mathbb{I}}, M_1^\alpha, M_2^\alpha. \\
 & M_1 \xi \cong_\alpha^{\hat{a}(\xi)} M_2 \xi \wedge M_1 \xrightarrow{[\mathbb{I}, \xi]} V_1 \wedge M_2 \xrightarrow{[\mathbb{I}, \xi]} V_2 \\
 & \rightarrow \\
 & \forall \mathbb{I}_i, \xi_1^{\mathbb{I}_i}, \xi_2^{\mathbb{I}_i}, A. (\xi \cdot x : V_1 \preceq^* \xi_1') \sim^{\preceq^*} (\xi \cdot x : V_2 \preceq^* \xi_2') \wedge \mathbb{I}_i \Vdash A \\
 & \rightarrow \\
 & \xi_1' \models A \leftrightarrow \xi_2' \models A
 \end{aligned}$$

Use IH as the additions are all of the same form when proving  $A$ :

*Proof.* Proof by IH on the structure of  $A$ :

- |   |   |
|---|---|
| 1 | Assume: $\mathbb{I}, \xi^{\mathbb{I}}, M_1^\alpha, M_2^\alpha$ s.t. $M_1 \xi \cong_\alpha^{\hat{a}(\xi)} M_2 \xi \wedge M_1 \xrightarrow{[\mathbb{I}, \xi]} V_1 \wedge M_2 \xrightarrow{[\mathbb{I}, \xi]} V_2$ |
| 2 | Assume: $\mathbb{I}_i, \xi_1^{\mathbb{I}_i}, \xi_2^{\mathbb{I}_i}, A$ s.t. $(\xi \cdot x : V_1 \preceq^* \xi_1') \sim^{\preceq^*} (\xi \cdot x : V_2 \preceq^* \xi_2') \wedge \mathbb{I}_i \Vdash A$            |
| 3 | Assume: $\xi_1' \models A$ to prove $\xi_2' \models A$  |

Noting that  $\forall \mathbb{I}_0. \mathbb{I}_i \Vdash \mathbb{I}_0 \rightarrow \llbracket \mathbb{I}_0 \rrbracket_{\xi_1'} \equiv \llbracket \mathbb{I}_0 \rrbracket_{\xi_2'}$  as  $\delta$ 's are unaffected by  $\sim^{\preceq^*}$ , and the standard mappings are also unaffected.

Next prove this final step by IH on the structure of  $A$ :

- $A \equiv e = e'$  Lemma 31.  
 $A \equiv \neg A'$  By IH on  $A'$ .  
 $A \equiv A_1 \wedge A_2$  By IH on  $A_1$  and  $A_2$ .  
 $A \equiv A_1 \vee A_2$  By IH on  $A_1$  and  $A_2$ .  
 $A \equiv A_1 \rightarrow A_2$  By IH on  $A_1$  and  $A_2$ .
- $A \equiv u \bullet e = m\{A'\}$  Assuming  $\xi_1' \models u \bullet e = m\{A'\}$  iff  $(\hat{a}(\xi_1'), ue \xrightarrow{[\mathbb{I}, \xi_1']} V_m \wedge \xi_1' \cdot m : V_m \models A'$   
 then it is clear there exists  $W_m$  s.t.  $ue \xrightarrow{[\mathbb{I}, \xi_2']} W_m$   
 thus  $(\xi \cdot x : V_1 \preceq^* \xi_1' \cdot m : V_m) \sim^{\preceq^*} (\xi \cdot x : V_2 \preceq^* \xi_2' \cdot m : W_m)$   
 hence by IH on  $A'$ :  $\xi_2' \cdot m : W_m \models A'$   
 hence  $\xi_2' \models u \bullet e = m\{A'\}$
- $A \equiv u \# \mathbb{I}_0$  Prove by direct use of Lemma 32.
- $A \equiv \forall u \in (\mathbb{I}_0). A'$  Assume  $\xi_1' \models \forall u \in (\mathbb{I}_0). A'$  iff  $\forall M_u. M_u \xrightarrow{[\mathbb{I}_0, \xi_1']} V_u \rightarrow \xi_1' \cdot u : V_u \models A'$   
 Prove  $\xi_2' \models \forall u \in (\mathbb{I}_0). A'$  iff  $\forall M_u. M_u \xrightarrow{[\mathbb{I}_0, \xi_2']} W_u \rightarrow \xi_2' \cdot u : W_u \models A'$   
 Assume some  $M_u$  in the  $\xi_2'$  such that  $M_u \xrightarrow{[\mathbb{I}_0, \xi_2']} W_u$  then given  $\llbracket \mathbb{I}_0 \rrbracket_{\xi_2'} \equiv \llbracket \mathbb{I}_0 \rrbracket_{\xi_1'}$  and the assumption then: for that  $M_u$  then  $M_u \xrightarrow{[\mathbb{I}_0, \xi_1']} V_u$  and hence  $\xi_1' \cdot u : V_u \models A'$  and as  $W_u$  and  $V_u$  are derived from the same term then  $(\xi \cdot x : V_1 \preceq^* \xi_1' \cdot u : V_u) \sim^{\preceq^*} (\xi \cdot x : V_2 \preceq^* \xi_2' \cdot u : W_u)$   
 by induction on  $A'$  this implies  $\xi_2' \cdot u : W_u \models A'$ .  
 Hence  $\forall M_u. M_u \xrightarrow{[\mathbb{I}_0, \xi_2']} W_u \rightarrow \xi_2' \cdot u : W_u \models A'$  hence  $\xi_2' \models \forall u \in (\mathbb{I}_0). A'$
- $A \equiv \exists u \in (\mathbb{I}_0). A'$  Similar proof as  $\forall u \in (\mathbb{I}_0). A'$  above.

$A \equiv \forall \delta. A'$  Assume  $\xi'_1 \models \forall \delta. A'$  iff  $\forall \xi_1'' \xi_1' \preceq^* \xi_1'' \rightarrow \xi_1' \cdot \delta : \mathbb{I}_i'' \setminus \text{-TCV} \models A'$   
 Prove  $\xi'_2 \models \forall \delta. A'$  iff  $\forall \xi_2'' \xi_2' \preceq^* \xi_2'' \rightarrow \xi_2' \cdot \delta : \mathbb{I}_i'' \setminus \text{-TCV} \models A'$   
 i.e. assume some  $\xi_2'' \xi_2'$  such that  $\xi_2' \preceq^* \xi_2''$  then for the same derivation  
 required for  $\xi_2''$  from  $\xi_2'$  there is an similarly derived model  $\xi_1''$  derived from  
 $\xi_1'$   
 and hence  $(\xi \cdot x : V_1 \preceq^* \xi_1'') \sim^{\preceq^*} (\xi \cdot x : V_2 \preceq^* \xi_2'')$  (By assumption)  
 and hence  $(\xi \cdot x : V_1 \preceq^* \xi_1' \cdot \delta : \mathbb{I}_i'' \setminus \text{-TCV}) \sim^{\preceq^*} (\xi \cdot x : V_2 \preceq^* \xi_2' \cdot \delta : \mathbb{I}_i'' \setminus \text{-TCV})$   
 (Def 3)  
 and IH on  $A'$  can now be used to show that given  $\xi_1'' \cdot \delta : \mathbb{I}_i'' \setminus \text{-TCV} \models A'$   
 then  $\xi_2'' \cdot \delta : \mathbb{I}_i'' \setminus \text{-TCV} \models A'$  hence the case holds.

**Lemma 34 (Two extensions combine to make extensions of each other).**

$$\begin{aligned}
 & \forall \mathbb{I}, \mathbb{I}_1, \mathbb{I}_2, \xi^{\mathbb{I}}, \xi_1^{\mathbb{I}_1}, \xi_2^{\mathbb{I}_2}. \\
 & \xi \preceq^* \xi \cdot \xi_1 \wedge \xi \preceq^* \xi \cdot \xi_2 \\
 & \rightarrow (\xi \cdot \xi_1 \preceq^* \xi \cdot \xi_1 \cdot \xi_2 \leftrightarrow \xi \cdot \xi_2 \preceq^* \xi \cdot \xi_1 \cdot \xi_2)
 \end{aligned}$$

i.e.

$$\begin{aligned}
 & \xi \preceq^* \xi \cdot \tilde{X} : \tilde{V} \wedge \xi \preceq^* \xi \cdot \tilde{Y} : \tilde{W} \\
 & \rightarrow (\xi \cdot \tilde{X} : \tilde{V} \preceq^* \xi \cdot \tilde{X} : \tilde{V} \cdot \tilde{Y} : \tilde{W} \leftrightarrow \xi \cdot \tilde{Y} : \tilde{W} \preceq^* \xi \cdot \tilde{X} : \tilde{V} \cdot \tilde{Y} : \tilde{W})
 \end{aligned}$$

*Proof.*

1 By symmetry this only needs to be proven in one direction, which is as follows:

2 Assume:  $\xi \preceq^* \xi \cdot \xi_1 \wedge \xi \preceq^* \xi \cdot \xi_2 \wedge \xi \cdot \xi_1 \preceq^* \xi \cdot \xi_1 \cdot \xi_2$

3 i.e. assume:  $\xi \preceq^* \xi \cdot \tilde{X} : \tilde{V}^{\tilde{\alpha}} \wedge \xi \preceq^* \xi \cdot \xi_2 \wedge \xi \cdot \tilde{X} : \tilde{V}^{\tilde{\alpha}} \preceq^* \xi \cdot \tilde{X} : \tilde{V}^{\tilde{\alpha}} \cdot \xi_2$

4 From Def  $\preceq^*$ ,  $\preceq$ , and Lemma 30, the TCV can be discarded in the reasoning, hence only standard variables are considered

5 Line 3  $\leftrightarrow \exists \tilde{M}. \tilde{M}_1 \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} \tilde{V}_1 \wedge \tilde{M}_2 \overset{[\mathbb{I} + \tilde{X}_1, \xi \cdot \tilde{X}_1 : \tilde{V}_1]}{\rightsquigarrow} \tilde{V}_2 \wedge \dots$   
 $\wedge \tilde{M}_{i+1} \overset{[\mathbb{I} + \tilde{X}_1 + \dots + \tilde{X}_i, \xi \cdot \tilde{X}_1 : \tilde{V}_1 \dots \tilde{X}_i : \tilde{V}_i]}{\rightsquigarrow} \tilde{V}_{i+1} \wedge \dots$   
 $\wedge \xi \preceq^* \xi \cdot \xi_2$   
 $\wedge \xi \cdot \tilde{X} : \tilde{V}^{\tilde{\alpha}} \preceq^* \xi \cdot \tilde{X} : \tilde{V}^{\tilde{\alpha}} \cdot \xi_2$

6  $\rightarrow \exists \tilde{M}. \tilde{M}_1 \overset{[\mathbb{I}, \xi \cdot \xi_2]}{\rightsquigarrow} \tilde{V}_1 \wedge \tilde{M}_2 \overset{[\mathbb{I} + \tilde{X}_1, \xi \cdot \xi_2 \cdot \tilde{X}_1 : \tilde{V}_1]}{\rightsquigarrow} \tilde{V}_2 \wedge \dots$  Lemma 11  $\xi \preceq^* \xi \cdot \xi_2$   
 $\wedge \tilde{M}_{i+1} \overset{[\mathbb{I} + \tilde{X}_1 + \dots + \tilde{X}_i, \xi \cdot \xi_2 \cdot \tilde{X}_1 : \tilde{V}_1 \dots \tilde{X}_i : \tilde{V}_i]}{\rightsquigarrow} \tilde{V}_{i+1} \wedge \dots$   $(\mathfrak{a}(\tilde{V}_2) \cap \mathfrak{a}(\xi_2) \subseteq \mathfrak{a}(\xi \cdot \tilde{X}_1 : \tilde{V}_1))$   
 $\xi \cdot \tilde{X}_1 : \tilde{V}_1 \preceq^* \xi \cdot \tilde{X}_1 : \tilde{V}_1 \cdot \xi_2$

7  $\rightarrow \exists \tilde{M}. \tilde{M}_1 \overset{[\mathbb{I} + \mathbb{I}_2, \xi \cdot \xi_2]}{\rightsquigarrow} \tilde{V}_1$   $[[\mathbb{I}]]_{\xi \cdot \xi_2} \subseteq [[\mathbb{I} + \mathbb{I}_2]]_{\xi \cdot \xi_2}$   
 $\wedge \tilde{M}_2 \overset{[\mathbb{I} + \mathbb{I}_2 + \tilde{X}_1, \xi \cdot \xi_2 \cdot \tilde{X}_1 : \tilde{V}_1]}{\rightsquigarrow} \tilde{V}_2 \wedge \dots$   $[[\mathbb{I} + \tilde{X}_1]]_{\xi \cdot \xi_2 \cdot \tilde{X}_1 : \tilde{V}_1} \subseteq [[\mathbb{I} + \tilde{X}_1 + \mathbb{I}_2]]_{\xi \cdot \tilde{X}_1 : \tilde{V}_1 \cdot \xi_2}$   
 $\wedge \tilde{M}_{i+1} \overset{[\mathbb{I} + \mathbb{I}_2 + \tilde{X}_1 + \dots + \tilde{X}_i, \xi \cdot \xi_2 \cdot \tilde{X}_1 : \tilde{V}_1 \dots \tilde{X}_i : \tilde{V}_i]}{\rightsquigarrow} \tilde{V}_{i+1} \wedge \dots$   $\dots$

8  $\leftrightarrow \xi \cdot \tilde{Y} : \tilde{W}^{\tilde{\beta}} \preceq^* \xi \cdot \tilde{X} : \tilde{V}^{\tilde{\alpha}} \cdot \tilde{Y} : \tilde{W}^{\tilde{\beta}}$

## C Soundness of Axioms

We now prove the soundness of the axioms in Sec. 5 split up into sections for each type of axiom.

### C.1 Soundness of Axioms for $e = e'$

The following axioms are proven trivially:

$$(eq2) \equiv x = x, (eq3) \equiv x = y \leftrightarrow y = x, (eq4) \equiv x = y \wedge y = z \rightarrow x = z$$

#### Soundness proof of Axiom (eq1):

$$A(x) \wedge x = e \rightarrow A(x)[e/x]$$

*Proof.* Syntactically: if  $x$  occurs in an LTC and  $e$  contains a destructor as in Sec. 3, then  $\mathbb{I}_0$  in  $A$  then  $\mathbb{I}_0 \Vdash e : \alpha$  holds, otherwise we just add all variables in  $e$  to  $\mathbb{I}_0$  s.t. the order is maintained by the global LTC.

1 Assume  $\mathbb{I}, \xi^{\mathbb{I}}$  s.t.  $\mathbb{I} \Vdash A(x) \wedge x = e \rightarrow A(x)[e/x]$  then:

2	$\xi^{\mathbb{I}} \models A(x) \wedge x = e$	<i>Assume</i>
3	$e$ free for $x$ in $A(x)$	<i>Assume</i>
4	$\xi \models A(x) \wedge \xi \models x = e$	<i>Sem. <math>\wedge</math></i>
5	$\xi \models A(x) \wedge \llbracket x \rrbracket_{\xi} \cong_{\alpha}^{\hat{\mathbf{a}}(\xi)} \llbracket e \rrbracket_{\xi}$	<i>Sem. =</i>
6	$\xi \models A(x) \wedge \llbracket x \rrbracket_{\xi} \cong_{\alpha}^{\hat{\mathbf{a}}(\xi)} \llbracket e \rrbracket_{\xi} \wedge (\hat{\mathbf{a}}(\xi), \llbracket e \rrbracket_{\xi}) \Downarrow (G', V_e)$	
7	$\xi \models A(x) \wedge \llbracket x \rrbracket_{\xi} \cong_{\alpha}^{\hat{\mathbf{a}}(\xi)} \llbracket e \rrbracket_{\xi} \wedge \llbracket e \rrbracket_{\xi} \cong_{\alpha}^{\hat{\mathbf{a}}(\xi)} V_e \wedge (\hat{\mathbf{a}}(\xi), \llbracket e \rrbracket_{\xi}) \Downarrow (G', V_e)$	<i>Lemma 21</i>
8	$\xi \models A(x) \wedge \llbracket x \rrbracket_{\xi} \cong_{\alpha}^{\hat{\mathbf{a}}(\xi)} V_e \wedge (\hat{\mathbf{a}}(\xi), \llbracket e \rrbracket_{\xi}) \Downarrow (G', V_e)$	$e \cong e' \wedge e' \cong e'' \rightarrow e \cong e''$
9	$\rightarrow \xi \models A(x) \wedge (\hat{\mathbf{a}}(\xi), \llbracket e \rrbracket_{\xi}) \Downarrow (G', V_e)$ $\wedge \forall x'. A_0(x'). x' \notin \text{dom}(\mathbb{I}) \wedge \mathbb{I} \cdot x' : \mathbb{I}(x) \Vdash A_0(x')$ $\rightarrow (\xi \cdot x' : \xi(x) \models A_0(x)[x'/x] \leftrightarrow \xi \cdot x' : V_e \models A_0(x)[x'/x])$	<i>Lemma 33</i> $B^{-x, x'}(x') \equiv B(x)[x'/x]$ $(\llbracket e \rrbracket_{\xi} \equiv e\xi)$
10	$\leftrightarrow \xi \models A(x) \wedge (\hat{\mathbf{a}}(\xi), \llbracket e \rrbracket_{\xi}) \Downarrow (G', V_e)$ $\wedge \forall x'. x' \notin \text{dom}(\mathbb{I}) \rightarrow (\xi \models A(x) \leftrightarrow \xi \cdot x' : V_e \models A(x'))$	<i>Let <math>A_0 \equiv A</math></i> $\xi \cdot x' : \xi(x) \models A(x') \leftrightarrow \xi \models A(x)$
11	$\rightarrow (\hat{\mathbf{a}}(\xi), \llbracket e \rrbracket_{\xi}) \Downarrow (G', V_e) \wedge \forall x'. x' \notin \text{dom}(\mathbb{I}) \rightarrow \xi \cdot x' : V_e \models A(x')$	$B \wedge (B \leftrightarrow C) \rightarrow C$
12	$\rightarrow \forall x'. x' \notin \text{dom}(\mathbb{I}) \rightarrow ((\hat{\mathbf{a}}(\xi), \llbracket e \rrbracket_{\xi}) \Downarrow (G', V_e) \wedge \xi \cdot x' : V_e \models A(x'))$	$A \wedge \forall x. (B \rightarrow C) \rightarrow \forall x. (B \rightarrow A \wedge C)$
13	$\leftrightarrow \forall x'. x' \notin \text{dom}(\mathbb{I}) \rightarrow \xi \models A(x')[e/x']$	<i>Sem. <math>A[e/x']</math> (<math>x' \notin \text{fv}(\xi)</math>)</i>
14	$\leftrightarrow \xi \models A(x)[e/x]$	<i>Sem. <math>A[e/x]</math> (<math>x \in \text{fv}(\xi)</math>)</i>
15	Hence: $\forall \xi^{\mathbb{I}}. \mathbb{I} \Vdash A(x) \wedge x = e \rightarrow A(x)[e/x] \rightarrow \xi \models A(x) \wedge x = e \rightarrow A(x)[e/x]$	<i>lines 1-14</i>

### C.2 Soundness of Axioms for $\forall x \in (\Gamma). A$

**Soundness proof of Axiom (u1):**

$$\forall x^\alpha \in (\mathbb{I}_0).A \wedge \mathbb{I}_0 \Vdash e : \alpha \rightarrow A[e/x]$$

*Proof.*

1	Assume: $\mathbb{I}, \xi^{\mathbb{I}^d}$ s.t. $\mathbb{I} \triangleright \xi \wedge \mathbb{I} \Vdash \forall x^\alpha \in (\mathbb{I}_0).A \rightarrow A[e/x]$ and $\mathbb{I}_0 \Vdash e : \alpha$	
2	Assume: $\xi \models \forall x^\alpha \in (\mathbb{I}_0).A$ and $\mathbb{I}_0 \vdash e : \alpha$ Note $e \in \{x, c, Op(\tilde{e}), \langle e_1, e_2 \rangle \pi_i(e_1)\}$	
3	$\forall M_x.M_x \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_x \rightarrow \xi \cdot x : V_x \models A$	<i>Sem. <math>\forall \in ()</math>, 2</i>
4	$e$ – expression $\rightarrow e$ – term $\rightarrow \hat{a}(e) = \emptyset$	<i>Lemma 19</i>
5	$\mathbb{I} \Vdash \mathbb{I}_0 \rightarrow (\mathbb{I}_0 \Vdash e : \alpha \rightarrow \mathbb{I} \Vdash e : \alpha)$	<i>Typing rules</i>
6	$\mathbb{I}_0 \Vdash e : \alpha \rightarrow \exists V_e.(\hat{a}(\xi), \llbracket e \rrbracket_\xi) \Downarrow (\hat{a}(\xi), V_e)$	<i>Lemma 18, 6</i>
7	$e \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_e$	<i>Sem. <math>\overset{[\cdot]}{\rightsquigarrow}</math>, 4, 5, 6, <math>\llbracket e \rrbracket_\xi \equiv e\xi</math></i>
8	$e \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_e \rightarrow \xi \cdot x : V_e \models A$	<i>Instantiate <math>\forall M_x</math> to <math>e</math>, 3</i>
9	$\rightarrow e \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V_e \wedge \xi \cdot x : V_e \models A$	<i>MP, 7, 8</i>
10	$\xi \models A[e/x]$	<i>Sem. <math>[e/x]</math>, <math>x \notin \text{dom}(\xi)</math></i>
11	Hence $\xi \models \forall x^\alpha \in (\mathbb{I}_0).A$ and $\mathbb{I}_0 \vdash e : \alpha$ implies $\xi \models A[e/x]$	<i>lines 2-10</i>
12	Hence $\forall \mathbb{I}, \mathbb{I}_0, A, e. \mathbb{I} \Vdash \forall x^\alpha \in (\mathbb{I}_0).A \rightarrow A[e/x] \wedge \mathbb{I}_0 \Vdash e : \alpha$ $\rightarrow \forall \xi. \mathbb{I} \triangleright \xi \rightarrow \xi \models ((\forall x^\alpha \in (\mathbb{I}_0).A) \rightarrow (A[e/x]))$	<i>lines 1-11</i>

**Soundness proof of Axiom (u2):**

$$\forall x^\alpha \in (\mathbb{I}_0 + \mathbb{I}_1).A \rightarrow (\forall x^\alpha \in (\mathbb{I}_0).A) \wedge (\forall x^\alpha \in (\mathbb{I}_1).A)$$

*Proof.*

1	Assume: $\mathbb{I}, \xi^{\mathbb{I}^d}$ s.t. $\mathbb{I} \triangleright \xi \wedge \mathbb{I} \Vdash \forall x^\alpha \in (\mathbb{I} + \mathbb{I}').A \rightarrow (\forall x^\alpha \in (\mathbb{I}).A) \wedge (\forall x^\alpha \in (\mathbb{I}').A)$	
2	Assume: $\xi \models \forall x^\alpha \in (\mathbb{I}_0, \mathbb{I}_1).A$	
3	Prove: $\xi \models (\forall x^\alpha \in (\mathbb{I}_0).A) \wedge (\forall x^\alpha \in (\mathbb{I}_1).A)$	
4	$\forall M.M \overset{[\mathbb{I}_0 + \mathbb{I}_1, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models A$	<i>Sem. <math>\forall x \in (\mathbb{I}_0, \mathbb{I}_1)</math>, 1</i>
5	$\rightarrow \forall M.M \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models A$	$\forall V. \mathbb{I}_i \vdash V : \alpha \dots \subseteq \forall V. \mathbb{I}_0 + \mathbb{I}_1 \vdash V : \alpha \dots$
6	$\xi \models \forall x^\alpha \in (\mathbb{I}_0).A$	<i>Sem. <math>\forall \in ()</math>.</i>
7	$\xi \models \forall x^\alpha \in (\mathbb{I}_1).A$	<i>Same as 3-5 but with <math>\mathbb{I}_1</math></i>
8	$\rightarrow \xi \models (\forall x^\alpha \in (\mathbb{I}_0).A) \wedge (\forall x^\alpha \in (\mathbb{I}_1).A)$	<i>5, 6</i>
9	Hence: $\forall \mathbb{I}. \mathbb{I} \Vdash \forall x^\alpha \in (\mathbb{I}_0 + \mathbb{I}_1).A \rightarrow (\forall x^\alpha \in (\mathbb{I}_0).A) \wedge (\forall x^\alpha \in (\mathbb{I}_1).A)$ $\rightarrow \forall \xi. \mathbb{I} \triangleright \xi \rightarrow \xi \models (\forall x^\alpha \in (\mathbb{I}_0).A) \wedge (\forall x^\alpha \in (\mathbb{I}_1).A)$	<i>lines 1-8</i>

**Soundness proof of Axiom (u3):**

$$A^{-x} \leftrightarrow \forall x^\alpha \in (\mathbb{I}_0).A$$

*Proof.* Clearly  $\forall x \in (\mathbb{I}_0).A \rightarrow A$  applying (u1) given  $A^{-x}[e/x] \equiv A$  and the  $x$  derived from  $\mathbb{I}_0$  (say  $V$ ) must imply an extension  $\xi \preceq^* \xi \cdot x : V$  which given  $A$ -Ext-Ind implies this direction.

Then to prove  $A^{-x} \rightarrow \forall x \in (\mathbb{I}_0).A$ :

1	Assume: $\mathbb{I}, \mathbb{I}_0, A$ s.t. $\mathbb{I} \Vdash A^{-x} \rightarrow \forall x^\alpha \in (\mathbb{I}_0).A \wedge A$ – Ext-Ind	
2	Assume: $\xi^{\mathbb{I}_d}$ s.t. $\mathbb{I} \triangleright \xi$ and $\xi \models A^{-x}$	
3	$\rightarrow \forall M. M \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \models A$	<i>Tautology</i>
4	$\rightarrow \forall M. M \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models A$	<i>Sem. <math>\preceq^*</math>, Lemmas 26 and A-Ext-Ind</i>
5	$\xi \models \forall x^\alpha \in (\mathbb{I}_0).A$	<i>Sem. <math>\forall \in ()</math>.</i>
6	$\xi \models A \rightarrow \xi \models \forall x^\alpha \in (\mathbb{I}_0).A$	<i>2-9</i>
7	$\xi \models A \rightarrow \forall x^\alpha \in (\mathbb{I}_0).A$	<i>Sem. <math>\rightarrow</math></i>
8	Hence: $\forall \mathbb{I}. \mathbb{I} \Vdash A \rightarrow \forall x^\alpha \in (\mathbb{I}_0).A$ $\wedge \forall \xi^{\mathbb{I}_d}. \mathbb{I} \triangleright \xi \rightarrow \xi \models A \rightarrow \forall x^\alpha \in (\mathbb{I}_0).A$	<i>lines 1-7</i>

**Soundness proof of Axiom (u4):**

$$(\forall x^\alpha \in (\mathcal{I}_0).(A \wedge B)) \leftrightarrow (\forall x^\alpha \in (\mathcal{I}_0).A) \wedge (\forall x^\alpha \in (\mathcal{I}_0).B)$$

*Proof.* Assume  $\mathcal{I}, \xi^{\mathcal{I}^d}, A$

such that  $\mathcal{I} \triangleright \xi \wedge \mathcal{I} \Vdash (\forall x^\alpha \in (\mathcal{I}_0).(A \wedge B)) \leftrightarrow (\forall x^\alpha \in (\mathcal{I}_0).A) \wedge (\forall x^\alpha \in (\mathcal{I}_0).B)$

$\rightarrow$ :

1	Assume: $\xi \models (\forall x^\alpha \in (\mathcal{I}_0).(A \wedge B))$	
2	$\forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \left( \begin{array}{l} \xi \cdot x : V \models A \\ \wedge \xi \cdot x : V \models B \end{array} \right)$	<i>Sem.</i> $\forall \in (), \wedge$
3	$\forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \left( \begin{array}{l} \xi \cdot x : V \models A \\ \wedge \xi \cdot x : V \models B \end{array} \right)$	<i>FOL</i> $A \rightarrow A \wedge A$
	$\wedge$	
	$\forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \left( \begin{array}{l} \xi \cdot x : V \models A \\ \wedge \xi \cdot x : V \models B \end{array} \right)$	
4	$\rightarrow \forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models A$	<i>FOL</i> $\wedge$ -elim
	$\wedge$	
	$\forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models B$	
5	$\xi \models \forall x^\alpha \in (\mathcal{I}_0).A \wedge \xi \models \forall x^\alpha \in (\mathcal{I}_0).B$	<i>Sem.</i> $\forall \in ()$ .
6	$\xi \models (\forall x^\alpha \in (\mathcal{I}_0).A) \wedge (\forall x^\alpha \in (\mathcal{I}_0).B)$	<i>Sem.</i> $\wedge$

$\leftarrow$ :

1	Assume: $\xi \models (\forall x^\alpha \in (\mathcal{I}_0).A) \wedge (\forall x^\alpha \in (\mathcal{I}_0).B)$	
2	$\forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models A$	<i>Sem.</i> $\forall \in (), \wedge, 1$
	$\wedge$	
	$\forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models B$	
3	$\forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models A$	<i>extract the common</i> $\forall M$
	$\wedge$	
	$M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \xi \cdot x : V \models B$	
4	$\forall M.M \stackrel{[\mathcal{I}_0, \xi]}{\rightsquigarrow} V \rightarrow \left( \begin{array}{l} \xi \cdot x : V \models A \\ \wedge \xi \cdot x : V \models B \end{array} \right)$	<i>FOL</i> $((A \rightarrow B) \wedge (A \rightarrow C))$ $\rightarrow (A \rightarrow (B \wedge C))$
5	$\xi \models \forall x^\alpha \in (\mathcal{I}_0).(A \wedge B)$	<i>Sem.</i> $\wedge, \forall x \in (\mathcal{I})$ .

**Soundness proof of Axiom (u5):**

$$\forall x^\alpha \in (\mathbb{I}_0).A \leftrightarrow \forall x^\alpha \in (\emptyset).A \quad \text{iff } \alpha \text{ is Nm-free}$$

*Proof.* Trivial using Lemma 13 as they both quantify over the same set of values.

**Soundness proof of Axiom (ex1):**

$$(\mathbb{I} \Vdash \mathbb{I}_0 \wedge \mathbb{I}_0 \Vdash e : \alpha) \rightarrow A[e/x] \rightarrow \exists x \in (\mathbb{I}_0).x = e \wedge A$$

*Proof.* Assume  $\mathbb{I}^{-x}, \xi^{\mathbb{I}^a}, \mathbb{I}_0^{-x}, e, A$   
such that  $\mathbb{I} \Vdash A[e/x] \rightarrow \exists x \in (\mathbb{I}).A$ ,  $\mathbb{I} \Vdash \mathbb{I}_0$  and  $\mathbb{I}_0 \Vdash e : \alpha$

1	Assume: $\mathbb{I}$ s.t. $\mathbb{I} \Vdash A[e/x] \rightarrow \exists x \in (\mathbb{I}_0).x = e \wedge A$	
2	Assume: $\mathbb{I}_0, e$ s.t. $\mathbb{I} \Vdash \mathbb{I}_0$ and $\mathbb{I}_0 \Vdash e : \alpha$	
3	Assume: $\xi^{\mathbb{I}^a}$ s.t. $\mathbb{I} \triangleright \xi$ and $\xi \models A[e/x]$	
4	$\leftrightarrow e \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V \wedge \xi \cdot x : V \models A$	<i>Sem.</i> $A[e/x]$ , $x \notin \text{dom}(\xi)$
5	$\leftrightarrow e \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \wedge \xi \cdot x : V \models A \wedge e(\xi \cdot x : V) \overset{\hat{\alpha}(\xi \cdot x : V)}{\cong_\alpha} x(\xi \cdot x : V)$	<i>Lemma 21</i>
6	$\leftrightarrow e \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \wedge \xi \cdot x : V \models x = e \wedge A$	<i>Sem.</i> $=, \wedge$
7	$\rightarrow \exists M_e.M_e \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V \wedge \xi \cdot x : V \models x = e \wedge A$	<i>Clearly holds for</i> $M_e = e$
8	$\leftrightarrow \xi \models \exists x \in (\mathbb{I}_0).x = e \wedge A$	<i>Sem.</i> $\exists x \in (\mathbb{I}_0)$ .

**Soundness proof of Axiom (ex2):** Assuming  $\{a, b\} \subseteq \text{dom}(\mathbb{I}_0)$ 

$$\mathbb{I} + x + \mathbb{I}_0 \Vdash a \bullet b = c\{c = x\} \rightarrow \exists x' \in (\mathbb{I}_0).x = x'$$

*Proof.*

1	Assume: $\xi^{\mathbb{I}^a}$ s.t. $\mathbb{I} + x + \mathbb{I}_0 \triangleright \xi$	
2	Assume: $\xi \models a \bullet b = c\{c = x\}$	
3	$\rightarrow ab \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_c \wedge \xi \cdot c : V_c \models x = c$	<i>Sem.</i> $\bullet = \{\}$
4	$\rightarrow \exists M'_x.M'_x \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_c \wedge \xi \cdot c : V_c \models x = c$	$M'_x \equiv ab$ ( $\{a, b\} \subseteq \text{dom}(\mathbb{I}_0)$ )
5	$\rightarrow \xi \models \exists x' \in (\mathbb{I}_0).x = x'$	<i>Sem.</i> $\exists \in ()$ .
6	Hence: $\forall \mathbb{I} + x + \mathbb{I}_0. \mathbb{I} + x + \mathbb{I}_0 \Vdash a \bullet b = c\{c = x\} \rightarrow \exists x' \in (\mathbb{I}_0).x = x'$ $\rightarrow \forall \xi^{\mathbb{I}^a}. \mathbb{I} + x + \mathbb{I}_0 \triangleright \xi \rightarrow \xi \models a \bullet b = c\{c = x\} \rightarrow \exists x' \in (\mathbb{I}_0).x = x'$	

**Soundness proof of Axiom (ex3):**

$$\mathbb{I}\Gamma + x \Vdash \forall y^{\text{Nm}} \in (\emptyset). \exists z^{\text{Nm}} \in (\mathbb{I}\Gamma_0 + y). x = z \rightarrow \exists z \in (\mathbb{I}\Gamma_0). x = z$$

*Proof.*

1	Assume some model $\xi^{\mathbb{I}\Gamma_d}$ s.t. $\mathbb{I}\Gamma + x \triangleright \xi$ and $\xi \models \forall y^{\text{Nm}} \in (\emptyset). \exists z^{\text{Nm}} \in (\mathbb{I}\Gamma_0 + y). x = z$	
2	$\rightarrow \forall M_y^{\text{Nm}}. M_y \xrightarrow{[\emptyset, \xi]} V_y \rightarrow \exists M_z^{\text{Nm}}. M_z \xrightarrow{[\mathbb{I}\Gamma_0 + y, \xi \cdot y : V_y]} V_z \wedge \xi \cdot y : V_y \cdot z : V_z \models x = z$	
3	$\rightarrow \forall M_y^{\text{Nm}}. M_y \xrightarrow{[\emptyset, \xi]} V_y \rightarrow \exists M_z^{\text{Nm}}. M_z \xrightarrow{[\mathbb{I}\Gamma_0, \xi \cdot y : V_y]} V_z \wedge \xi \cdot y : V_y \cdot z : V_z \models x = z$	<i>replace y with M<sub>y</sub></i>
4	$\rightarrow \forall M_y^{\text{Nm}}. M_y \xrightarrow{[\emptyset, \xi]} V_y \rightarrow \exists M_z^{\text{Nm}}. M_z \xrightarrow{[\mathbb{I}\Gamma_0, \xi]} V_z \wedge \llbracket x \rrbracket_{\xi \cdot y : V_y \cdot z : V_z} \cong_{\text{Nm}}^{\hat{a}(\xi \cdot y : V_y \cdot z : V_z)} \llbracket z \rrbracket_{\xi \cdot y : V_y \cdot z : V_z}$	<i>Lemma 1</i>
5	$\rightarrow \forall M_y^{\text{Nm}}. M_y \xrightarrow{[\emptyset, \xi]} V_y \rightarrow \exists M_z^{\text{Nm}}. M_z \xrightarrow{[\mathbb{I}\Gamma_0, \xi]} V_z \wedge \llbracket x \rrbracket_{\xi \cdot z : V_z} \cong_{\text{Nm}}^{\hat{a}(\xi \cdot y : V_y \cdot z : V_z)} \llbracket z \rrbracket_{\xi \cdot z : V_z}$	$\llbracket x \rrbracket_{\xi \cdot y : V_y} = \llbracket x \rrbracket_{\xi}$
6	$\rightarrow \forall M_y^{\text{Nm}}. M_y \xrightarrow{[\emptyset, \xi]} V_y \rightarrow \exists M_z^{\text{Nm}}. M_z \xrightarrow{[\mathbb{I}\Gamma_0, \xi]} V_z \wedge \llbracket x \rrbracket_{\xi \cdot z : V_z} \cong_{\text{Nm}}^{\hat{a}(\xi \cdot z : V_z)} \llbracket z \rrbracket_{\xi \cdot z : V_z}$	<i>Lemma 2</i>
7	$\rightarrow \exists M_z^{\text{Nm}}. M_z \xrightarrow{[\mathbb{I}\Gamma_0, \xi]} V_z \wedge \llbracket x \rrbracket_{\xi \cdot z : V_z} \cong_{\text{Nm}}^{\hat{a}(\xi \cdot z : V_z)} \llbracket z \rrbracket_{\xi \cdot z : V_z}$	<i>FOL</i>
8	$\rightarrow \xi \models \exists z \in (\mathbb{I}\Gamma_0). x = z$	<i>Sem. <math>\exists \in ()</math>.</i>
9	Hence: $\forall \mathbb{I}\Gamma + x. \mathbb{I}\Gamma + x \Vdash \forall y^{\text{Nm}} \in (\emptyset). \exists z^{\text{Nm}} \in (\mathbb{I}\Gamma_0 + y). x = z \rightarrow \exists z \in (\mathbb{I}\Gamma_0). x = z$ $\rightarrow \forall \xi^{\mathbb{I}\Gamma_d}. \mathbb{I}\Gamma + x \triangleright \xi \rightarrow \xi \models \forall y^{\text{Nm}} \in (\emptyset). \exists z^{\text{Nm}} \in (\mathbb{I}\Gamma_0 + y). x = z \rightarrow \exists z \in (\mathbb{I}\Gamma_0). x = z$	

**C.3 Soundness of Axioms for  $x \# \Gamma$** **Soundness proof of Axiom (f1):**

$$\mathbb{I}\Gamma + x : \text{Nm} + f : \alpha \rightarrow \alpha_{\mathfrak{b}} \Vdash x \# \mathbb{I}\Gamma \rightarrow x \# \mathbb{I}\Gamma + f : \alpha \rightarrow \alpha_{\mathfrak{b}}$$



*Proof.* We use Lemma 14 so show that  $f$  cannot help produce  $x$  if  $f : \alpha \rightarrow \alpha_b$ .

1	Assume: $\xi_{xf}$ s.t. $\mathbb{I}\Gamma + x : \text{Nm} + f : \alpha \rightarrow \alpha_b \triangleright (\xi^{\mathbb{I}\Gamma_0} \cdot x : r_x \cdot f : V_f)^{\mathbb{I}\Gamma_0 + x : \text{Nm} + f : \alpha \rightarrow \alpha_b} \equiv \xi_{xf}$	
2	$\mathbb{I}\Gamma + x + f \triangleright \xi_{xf} \rightarrow \exists M_x.M_x \overset{[\mathbb{I}\Gamma, \xi]}{\rightsquigarrow} r_x \wedge \exists M_f.M_f \overset{[\mathbb{I}\Gamma + x, \xi \cdot x : r_x]}{\rightsquigarrow} V_f$	
3	Assume: $\xi_{xf} \models x \# \mathbb{I}\Gamma$	<i>Assumption</i>
4	Hence: $\exists M_x.M_x \overset{[\mathbb{I}\Gamma, \xi]}{\rightsquigarrow} r_x \wedge \neg \exists N_x.N_x \overset{[\mathbb{I}\Gamma, \xi_{xf}]}{\rightsquigarrow} r_x$	<i>Sem. <math>x \# \mathbb{I}\Gamma</math>, 2</i>
5	Hence: $\exists M_x.M_x \overset{[\mathbb{I}\Gamma, \xi]}{\rightsquigarrow} r_x \wedge \neg \exists N_x.N_x \overset{[\mathbb{I}\Gamma, \xi \cdot x : r_x]}{\rightsquigarrow} r_x$	<i>Lemma 11</i>
6	$\rightarrow r_x \notin \mathfrak{a}(\xi)$	<i>Lemma 15</i>
7	Assume: $r_x \notin \mathfrak{a}(V_f)$	<i>Then trivial as <math>x</math>-fresh and <math>f</math> cannot output <math>r_x</math>, Lemma 22</i>
8	Assume: $r_x \in \mathfrak{a}(V_f)$	<i>(proof by contradiction)</i>
9	Assume: $\exists P_x.P_x \overset{[\mathbb{I}\Gamma + f, \xi_{xf}]}{\rightsquigarrow} r_x$	
10	$\leftrightarrow \exists P_x.\mathfrak{a}(P_x) = \emptyset$ $\wedge \llbracket \mathbb{I}\Gamma \rrbracket_{\xi}, f : \alpha \rightarrow \alpha_b \vdash P_x : \text{Nm}$ $\wedge (\mathfrak{a}(\xi_{xf}), P_x \xi_{xf}) \Downarrow (\mathfrak{a}(\xi_{xf}), G', r_x)$	<i>Sem. <math>\overset{[\cdot]}{\rightsquigarrow}</math></i>
11	$\leftrightarrow \exists P_x.\mathfrak{a}(P_x) = \emptyset$ $\wedge \llbracket \mathbb{I}\Gamma \rrbracket_{\xi}, f : \alpha \rightarrow \alpha_b \vdash P_x : \text{Nm}$ $\wedge (\mathfrak{a}(\xi_{xf}), P_x[V_f/f]\xi) \Downarrow (\mathfrak{a}(\xi_{xf}), G', r_x)$	<i>Def <math>[V/x]</math>, <math>P_x^{-x}</math></i>
12	$\leftrightarrow \exists P_x.\mathfrak{a}(P_x) = \emptyset$ $\wedge \llbracket \mathbb{I}\Gamma \rrbracket_{\xi}, f : \alpha \rightarrow \alpha_b \vdash P_x : \text{Nm}$ $\wedge (\mathfrak{a}(\xi_{xf}), (P_x \xi)[V_f/f]) \Downarrow (\mathfrak{a}(\xi_{xf}), G', r_x)$	<i>Def closure, <math>V_f</math>-value</i>
13	$\leftrightarrow \exists P_x.\mathfrak{a}(P_x) = \emptyset$ $\wedge \llbracket \mathbb{I}\Gamma \rrbracket_{\xi}, f : \alpha \rightarrow \alpha_b \vdash P_x : \text{Nm}$ $\wedge (\mathfrak{a}(\xi_{xf}), (P_x \xi)[V_f/f]) \Downarrow (\mathfrak{a}(\xi_{xf}), G', r_x)$ $\wedge \neg(\mathfrak{a}(\xi_{xf}), (P_x \xi)[V_f/f]) \Downarrow (\mathfrak{a}(\xi_{xf}), G', r_x)$	$r_x \notin \xi \rightarrow r_x \notin \mathfrak{a}(P_x \xi),$ $r_x \in \mathfrak{a}(V_f)$ <i>Lemma 14 <math>\rightarrow \neg(\dots \Downarrow r_x)</math></i>
14	Contradiction, hence: $\neg \exists P_x.P_x \overset{[\mathbb{I}\Gamma + f, \xi_{xf}]}{\rightsquigarrow} r_x$	
15	Hence: $\forall \xi_{xf}.\mathbb{I}\Gamma + x + f \triangleright \xi_{xf} \rightarrow \xi_{xf} \models x \# \mathbb{I}\Gamma \rightarrow x \# \mathbb{I}\Gamma + f$	

**Soundness proof of Axiom (f2):**

$$(x\#\mathbb{I}_0 \wedge \forall y^{\alpha_y} \in (\mathbb{I}_0).A) \leftrightarrow \forall y^{\alpha_y} \in (\mathbb{I}_0).(x\#(\mathbb{I}_0+y : \alpha_y) \wedge A)$$

*Proof.* The  $\leftarrow$  direction is elementary.

The  $\rightarrow$  direction is as follows:

1	Assume some $\mathbb{I}_1, \xi^{\mathbb{I}_1}$ s.t. $\mathbb{I}_1 \Vdash (x\#\mathbb{I}_0 \wedge \forall y^{\alpha_y} \in (\mathbb{I}_0).A)$	
	$\rightarrow \forall y^{\alpha_y} \in (\mathbb{I}_0).(x\#(\mathbb{I}_0+y : \alpha_y) \wedge A)$	
2	Assume: $\xi \models x\#\mathbb{I}_0 \wedge (\forall y \in (\mathbb{I}_0).A)$	
3	$\neg \exists M_x.M_x \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} \llbracket x \rrbracket_\xi$ $\wedge \forall M_y.M_y \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_y$ $\rightarrow \xi \cdot y : V_y \models A$	<i>Sem.</i> $x\#\mathbb{I}_0$ $\forall y \in (\mathbb{I}_0).$ $A$
4	$\forall M_y.M_y \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_y$ $\rightarrow \neg \exists M_x.M_x \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} \llbracket x \rrbracket_\xi$ $\wedge \xi \cdot y : V_y \models A$	$A^{-M_y} \wedge \forall M_y. B \rightarrow C$ $\rightarrow$ $\forall M_y. B \rightarrow (A \wedge C)$
5	$\forall M_y.M_y \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_y$ $\rightarrow \neg \exists M_x.M_x \overset{[\mathbb{I}_0+y, \xi \cdot y : V_y]}{\rightsquigarrow} \llbracket x \rrbracket_{\xi \cdot y : V_y}$ $\wedge \xi \cdot y : V_y \models A$	<i>Lemma 16</i> $\overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} s \leftrightarrow \overset{[\mathbb{I}_0+y, \xi \cdot y]}{\rightsquigarrow} s,$ <i>Lemma 24,</i> $\llbracket x \rrbracket_\xi \equiv \llbracket x \rrbracket_{\xi \cdot y : V_y}$
6	$\forall M_y.M_y \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_y$ $\rightarrow \xi \cdot y : V_y \models x\#\mathbb{I}_0+y$ $\wedge \xi \cdot y : V_y \models A$	<i>Sem.</i> #
7	$\xi \models \forall y \in (\mathbb{I}_0).x\#(\mathbb{I}_0+y) \wedge A$	<i>Sem.</i> $\forall \in (\cdot), \wedge$

**Soundness proof of Axiom (f3):**

$$x\#\mathbb{I}_0 \wedge \mathbb{I}_0 \Vdash e : \text{Nm} \rightarrow x \neq e$$

*Proof.* Provable directly from syntactic definition of  $x\#\mathbb{I}_0$  and (u1).

**Soundness proof of Axiom (f4)**

$$x\#(\mathbb{I}_0+\mathbb{I}_1) \rightarrow x\#\mathbb{I}_0 \wedge x\#\mathbb{I}_1$$

*Proof.* Provable directly from syntactic definition of  $x\#\mathbb{I}_0$ , and (u2).

**C.4 Soundness of Axioms for  $\forall\delta.A$** **Soundness proof of Axiom (utc1):**

$$\mathbb{I} \Vdash (\forall\delta.A) \rightarrow A[\mathbb{I}/\delta]$$

*Proof.*

1	Assume: $\mathbb{I}\Gamma$ s.t. $\mathbb{I}\Gamma \Vdash (\forall\delta.A) \rightarrow A[\mathbb{I}\Gamma/\delta]$	
2	Assume: $\xi^{\mathbb{I}d}$ s.t. $\mathbb{I}\Gamma \triangleright \xi$ and $\xi \models \forall\delta.A$	
3	$\leftrightarrow \forall \xi_0^{\mathbb{I}0}. \xi \preceq^* \xi_0 \rightarrow \xi_0 \cdot \delta : \mathbb{I}0 \models A$	<i>Sem.</i> $\forall\delta$ .
4	$\rightarrow \xi \preceq^* \xi \rightarrow \xi \cdot \delta : \mathbb{I}d \models A$	<i>Instantiate</i> $\forall \xi_0^{\mathbb{I}0}$ with $\xi$
5	$\rightarrow \xi \cdot \delta : \mathbb{I}d \models A$	<i>FOL</i> , $(\xi \preceq^* \xi) \equiv \top$
6	$\rightarrow \xi \cdot \delta : \mathbb{I}\Gamma \models A$	$\mathbb{I}\Gamma \triangleright \xi \rightarrow \llbracket \mathbb{I}\Gamma \rrbracket_\xi \equiv \llbracket \mathbb{I}d \rrbracket_\xi$
7	$\rightarrow \xi \models A[\mathbb{I}\Gamma/\delta]$	<i>Sem.</i> $[\mathbb{I}\Gamma/\delta]$

**Soundness proof of Axiom (*utc2*):**

$$\mathbb{I}\Gamma \Vdash \forall x^{\text{Nm}} \in (\mathbb{I}\Gamma).A^{-\delta} \leftrightarrow \forall \delta. \forall x^{\text{Nm}} \in (\mathbb{I}\Gamma + \delta).A^{-\delta} \text{ if } A\text{-Ext-Ind}$$

*Proof.*  $\leftarrow$ : Holds through (*utc1*)

$\rightarrow$ :

1	Assume: $\mathbb{I}\Gamma$ s.t. $\mathbb{I}\Gamma \Vdash \forall x \in (\mathbb{I}\Gamma).A \leftrightarrow \forall \delta. \forall x \in (\mathbb{I}\Gamma + \delta).A^{-\delta}$	
2	Assume: $\xi^{\mathbb{I}d}$ s.t. $\mathbb{I}\Gamma \triangleright \xi$ and $\xi \models \forall x^{\text{Nm}} \in (\mathbb{I}\Gamma).A$	
3	$\leftrightarrow \forall M.M \overset{[\mathbb{I}\Gamma, \xi]}{\rightsquigarrow} r_0 \rightarrow \xi \cdot x : r_0 \models A$	
4	Assume arbitrary $\xi'^{\mathbb{I}d}$ , $M'$ s.t. $\xi \preceq^* \xi'$ and $M' \overset{[\mathbb{I}\Gamma + \delta, \xi' \cdot \delta : \mathbb{I}\Gamma' \setminus \text{TCV}]}{\rightsquigarrow} r_1$	‘ $\forall \delta. \forall x \in (\delta)$ .’
5	$\leftrightarrow M' \overset{[\mathbb{I}\Gamma', \xi']}{\rightsquigarrow} r_1$	$\llbracket \mathbb{I}\Gamma + \delta \rrbracket_{\xi' \cdot \delta : \mathbb{I}\Gamma' \setminus \text{TCV}} \equiv \mathbb{I}\Gamma'$ , <i>Lemma 10</i>
6	$r_1 \in \hat{\mathfrak{a}}(\xi) \rightarrow r_0 \equiv r_1 \rightarrow \xi \cdot x : r_1 \models A$	<i>Lemma 17</i> $\rightarrow r_0 \equiv r_1$ obtainable
7	$r_1 \notin \hat{\mathfrak{a}}(\xi) \rightarrow \text{fresh-}r_0 \equiv r_1 \rightarrow \xi \cdot x : r_1 \models A$	<i>Let</i> $r_0 \equiv r_1$ as $r_1 \notin \hat{\mathfrak{a}}(\xi)$ (fresh names can be swapped)
8	$\rightarrow \xi' \cdot x : r_1 \models A$	<i>Lemma 29</i> , <i>A – Ext-Ind</i>
9	$\rightarrow \xi' \cdot \delta : \mathbb{I}\Gamma' \setminus \text{TCV} \cdot x : r_1 \models A$	<i>Lemma 30</i> , $A^{-\delta}$
10	$\rightarrow \xi \models \forall \delta. \forall x^{\text{Nm}} \in (\mathbb{I}\Gamma + \delta).A^{-\delta}$	<i>lines 4-9</i>
11	Hence: $\forall \mathbb{I}\Gamma. \mathbb{I}\Gamma \Vdash \forall x \in (\mathbb{I}\Gamma).A \leftrightarrow \forall \delta. \forall x \in (\mathbb{I}\Gamma + \delta).A^{-\delta}$	<i>lines 1-10</i>
	$\rightarrow \forall \xi^{\mathbb{I}d}. \mathbb{I}\Gamma \triangleright \xi \rightarrow \xi \models \forall x^{\text{Nm}} \in (\mathbb{I}\Gamma).A^{-\delta} \rightarrow \forall \delta. \forall x^{\text{Nm}} \in (\mathbb{I}\Gamma + \delta).A^{-\delta}$	
	(Assuming <i>A – Ext-Ind</i> )	

Essentially in line 6: if  $r_1$  is in  $\hat{\mathfrak{a}}(\xi)$  then it can be derived from  $\xi$ , and in line 7 if  $r_1$  is not in  $\hat{\mathfrak{a}}(\xi)$  then instantiating  $M \equiv \text{gensym}()$  produces a fresh name which can easily be set as (or swapped for)  $r_1$ .

**Soundness proof of Axiom (utc3):**

$$A - \text{Ext-Ind} \rightarrow A^{-\delta} \leftrightarrow \forall \delta. A$$

*Proof.* Assume:  $\mathbb{I}, \xi^{\mathbb{I}}$  s.t.  $\mathbb{I} \Vdash A^{-\delta} \leftrightarrow \forall \delta. A$  and assume some  $A^{-\delta}$ -Ext-Ind, then:

$\leftarrow$ : Use (utc1) knowing that  $A^{-\delta}[\mathbb{I}/\delta] \equiv A$ .

$\rightarrow$ :

1	Assume: $\xi \models A^{-\delta}$	
2	Prove: $\xi \models \forall \delta. A$	
3	$\xi \models A^{-\delta} \rightarrow \forall \xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow \xi \models A^{-\delta}$	<i>trivial addition</i>
4	$\xi \models A^{-\delta} \rightarrow \forall \xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow \xi' \models A^{-\delta}$	<i>A-Ext-Ind</i>
5	$\xi \models A^{-\delta} \rightarrow \forall \xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \wedge \xi' \cdot \delta : \mathbb{I}' \models A^{-\delta}$	<i>Lemma 30</i>
6	$\xi \models A^{-\delta} \rightarrow \xi \models \forall \delta. A$	<i>Sem. <math>\forall \delta</math>.</i>

**Soundness proof of Axiom (utc4):**

$$\forall \delta. (A \wedge B) \leftrightarrow (\forall \delta. A) \wedge (\forall \delta. B)$$

*Proof.* Assume  $\mathbb{I}, \xi^{\mathbb{I}}$  s.t.  $\mathbb{I} \Vdash \forall \delta. (A \wedge B) \leftrightarrow (\forall \delta. A) \wedge (\forall \delta. B)$

$\rightarrow$ :

1	Assume: $\xi \models \forall \delta. (A \wedge B)$	
2	Prove: $\xi \models (\forall \delta. A) \wedge (\forall \delta. B)$	
3	$\forall \xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow \xi' \cdot \delta : \mathbb{I}' \models A \wedge B$	<i>Sem. <math>\forall</math>.</i>
4	$\leftrightarrow \forall \xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow \xi' \cdot \delta : \mathbb{I}' \models A \wedge B$	<i>FOL <math>A \rightarrow (A \wedge A)</math></i>
5	$\rightarrow \forall \xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow \xi' \cdot \delta : \mathbb{I}' \models A$	<i><math>\wedge</math>-elim</i>
6	$\leftrightarrow \xi \models (\forall \delta. A) \wedge (\forall \delta. B)$	<i><math>\wedge</math>-elim</i>

←:

1	Assume: $\xi \models (\forall\delta.A) \wedge (\forall\delta.B)$	
2	Prove: $\xi \models \forall\delta.(A \wedge B)$	
3	$\Leftrightarrow \forall\xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow \xi' \cdot \delta : \mathbb{I}' \models A$	<i>Sem. <math>\wedge, \forall</math>.</i>
	$\wedge \forall\xi'^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow \xi' \cdot \delta : \mathbb{I}' \models B$	
4	$\rightarrow \forall\xi'^{\mathbb{I}'}. \xi \preceq^* \xi'$	<i><math>\forall\xi'</math> unifying</i>
	$\rightarrow \left( \begin{array}{l} \xi' \cdot \delta : \mathbb{I}' \models A \\ \wedge \xi' \cdot \delta : \mathbb{I}' \models B \end{array} \right)$	
5	$\Leftrightarrow \forall\xi'^{\mathbb{I}'}. \xi \preceq^* \xi'$	<i>Sem. <math>\wedge</math></i>
	$\rightarrow \xi' \cdot \delta : \mathbb{I}' \models A \wedge B$	
6	$\Leftrightarrow \xi \models \forall\delta.(A \wedge B)$	<i><math>\wedge</math>-elim</i>

### C.5 Soundness of Axioms for $u \bullet e = m\{A\}$

**Soundness proof of Axiom (e1)**

$$e_1 \bullet e_2 = m\{A^{-m} \wedge B\} \leftrightarrow (A \wedge e_1 \bullet e_2 = m\{B\}) \quad \text{iff } m \notin \text{fv}(A) \text{ and } A\text{-Ext-Ind.}$$

1	Assume: $\mathbb{I}, \xi^{\mathbb{I}}$ s.t.		
2	Assume: $\mathbb{I} \Vdash e_1 \bullet e_2 = m\{A^{-m} \wedge B\} \leftrightarrow (A \wedge e_1 \bullet e_2 = m\{B\})$ with $m \notin \text{fv}(A)$ and $A\text{-Ext-Ind.}$		
3	$\xi \models e_1 \bullet e_2 = m\{A^{-m} \wedge B\}$		$m \notin \text{dom}(\mathbb{I})$
4	$\leftrightarrow e_1 e_2 \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V \wedge \xi \cdot m : V \models A \wedge B$		$\text{Sem. } \bullet = \{\}$
5	$\leftrightarrow e_1 e_2 \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V \wedge \xi \cdot m : V \models A \wedge \xi \cdot m : V \models B$		$\text{Sem. } \wedge$
6	$\leftrightarrow e_1 e_2 \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V \wedge \xi \models A \wedge \xi \cdot m : V \models B$	$\text{Sem. } \preceq^*, \text{ Lemma 26 and } A\text{-Ext-Ind}$	
7	$\leftrightarrow \xi \models A \wedge \xi \models e_1 \bullet e_2 = m\{B\}$		$\text{Sem. } \bullet = \{\}$
8	$\leftrightarrow \xi \models (A \wedge e_1 \bullet e_2 = m\{B\})$		$\text{Sem. } \wedge$

**Soundness proof of Axiom (e2)**

$$e_1 \bullet e_2 = m\{\forall x \in (\mathbb{I}_0).A\} \leftrightarrow \forall x \in (\mathbb{I}_0).e_1 \bullet e_2 = m\{A\} \text{ iff } x \neq e_1, e_2, m \wedge e_1, e_2, m \notin \mathbb{I}_0$$

1	Assume: $\mathbb{I}, \xi^{\mathbb{I}}$ s.t.		
2	Assume: $\mathbb{I} \Vdash e_1 \bullet e_2 = m\{\forall x \in (\mathbb{I}_0).A\} \leftrightarrow \forall x \in (\mathbb{I}_0).e_1 \bullet e_2 = m\{A\}$ with $x \neq e_1, e_2, m \wedge e_1, e_2, m \notin \mathbb{I}_0$		
3	Assume: $\xi \models e_1 \bullet e_2 = m\{\forall x \in (\mathbb{I}_0).A\}$		
4	$\leftrightarrow e_1 e_2 \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_m \wedge \forall M_x. M_x \stackrel{[\mathbb{I}_0, \xi; m:V_m]}{\rightsquigarrow} V_x \rightarrow \xi \cdot m : V_m \cdot x : V_x \models A$		$\text{Sem. } \bullet = \{\}, \forall \in ()$
5	$\leftrightarrow e_1 e_2 \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_m \wedge \forall M_x. M_x \stackrel{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_x \rightarrow \xi \cdot m : V_m \cdot x : V_x \models A$	$m \notin \mathbb{I}_0 \wedge \xi \preceq^* \xi \cdot m : V_m$ $\text{Lemma 11 } (\hat{a}(V_x) \cap \hat{a}(V_m) \subseteq \hat{a}(\xi))$	
6	$\leftrightarrow \forall M_x. e_1 e_2 \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_m \wedge (M_x \stackrel{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_x \rightarrow \xi \cdot m : V_m \cdot x : V_x \models A)$	$A \wedge \forall X. B$ $\leftrightarrow \forall X. A^{-X} \wedge B$	
7	$\leftrightarrow \forall M_x. M_x \stackrel{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_x \rightarrow (e_1 e_2 \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_m \wedge \xi \cdot m : V_m \cdot x : V_x \models A) \rightarrow: A \wedge (B \rightarrow C) \rightarrow B \rightarrow (C \wedge A)$	$\leftarrow: \llbracket e_1 e_2 \rrbracket \Downarrow V_m \text{ terminates}$	
8	$\leftrightarrow \forall M_x. M_x \stackrel{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_x \rightarrow (e_1 e_2 \stackrel{[\mathbb{I}, \xi; x:V_x]}{\rightsquigarrow} V_m \wedge \xi \cdot m : V_m \cdot x : V_x \models A)$	$x \neq e_1, e_2,$ $\text{Lemma 11}$ $(\hat{a}(V_x) \cap \hat{a}(V_m) \subseteq \hat{a}(\xi \cdot x))$	
9	$\leftrightarrow \xi \models \forall x \in (\mathbb{I}_0).e_1 \bullet e_2 = m\{A\}$		$\text{Sem. } \forall \in (), \bullet = \{\}$

**Soundness proof of Axiom (e3)**

$$e_1 \bullet e_2 = m^{\alpha_b}\{\forall\delta.A\} \leftrightarrow \forall\delta.e_1 \bullet e_2 = m^{\alpha_b}\{A\} \text{ iff } A - \text{Ext-Ind}$$

1	Assume: $\mathbb{I}, \xi^{\mathbb{I}}$ s.t.	
2	Assume: $\mathbb{I} \Vdash e_1 \bullet e_2 = m^{\alpha_b}\{\forall\delta.A\} \leftrightarrow \forall\delta.e_1 \bullet e_2 = m^{\alpha_b}\{A\}$ with $A$ -Ext-Ind	
3	Assume: $\xi \models e_1 \bullet e_2 = m^{\alpha_b}\{\forall\delta.A\}$	
4	$\leftrightarrow e_1 e_2 \xrightarrow{[\mathbb{I}, \xi]} V_m \wedge \forall \xi_m^{\mathbb{I}'}. \xi \cdot m : V_m \preceq^* \xi_m \rightarrow \xi_m \cdot \delta : \mathbb{I}' \setminus \text{-TCV} \models A$	<i>Sem. <math>\bullet = \{\}, \forall\delta.</math></i>
5	$\leftrightarrow e_1 e_2 \xrightarrow{[\mathbb{I}, \xi]} V_m \wedge \forall \xi^{\mathbb{I}'}. \xi \cdot m : V_m \preceq^* \xi' \cdot m : V_m \rightarrow \xi' \cdot m : V_m \cdot \delta : \mathbb{I}' \setminus \text{-TCV} \models A$	<i>Rewrite <math>\xi_m</math> Lemma 6</i>
6	$\leftrightarrow e_1 e_2 \xrightarrow{[\mathbb{I}, \xi]} V_m \wedge \forall \xi^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow \xi' \cdot m : V_m \cdot \delta : \mathbb{I}' \setminus \text{-TCV} \models A$	<i>Lemma 7</i>
7	$\leftrightarrow \forall \xi^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow (e_1 e_2 \xrightarrow{[\mathbb{I}, \xi]} V_m \wedge \xi' \cdot m : V_m \cdot \delta : \mathbb{I}' \setminus \text{-TCV} \models A)$	<i><math>[[e_1 e_2]]_\xi \Downarrow V_m</math> terminates</i>
8	$\leftrightarrow \forall \xi^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow (e_1 e_2 \xrightarrow{[\mathbb{I}, \xi]} V_m \wedge \xi' \cdot m : V_m \cdot \delta : \mathbb{I}' \setminus \text{-TCV} \models A)$	<i>Lemmas 3, 11</i>
9	$\leftrightarrow \forall \xi^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow (e_1 e_2 \xrightarrow{[\mathbb{I}, \xi', \delta : \mathbb{I}' \setminus \text{-TCV}]} V_m \wedge \xi' \cdot \delta : \mathbb{I}' \setminus \text{-TCV} \cdot m : V_m \models A)$	<i>Lemma 10</i>
10	$\leftrightarrow \forall \xi^{\mathbb{I}'}. \xi \preceq^* \xi' \rightarrow (e_1 e_2 \xrightarrow{[\mathbb{I}' + \delta, \xi', \delta : \mathbb{I}' \setminus \text{-TCV}]} V_m \wedge \xi' \cdot \delta : \mathbb{I}' \setminus \text{-TCV} \cdot m : V_m \models A)$	<i><math>\mathbb{I} \Vdash e_1 e_2 : \alpha_b</math></i>
11	$\leftrightarrow \xi \models \forall\delta.e_1 \bullet e_2 = m^{\alpha_b}\{A\}$	<i>Sem. <math>\bullet = \{\}, \forall\delta.</math></i>

**Soundness proof of Axiom (ext)** For all  $e_1, e_2 : \alpha_1 \rightarrow \alpha_2$  s.t.  $\alpha_1$  and  $\alpha_2$  are both Nm-free:

$$(\forall x^{\alpha_1} \in (\emptyset). e_1 \bullet x = m_1^{\alpha_2}\{e_2 \bullet x = m_2^{\alpha_2}\{m_1 = m_2\}\}) \leftrightarrow e_1 =^{\alpha_1 \rightarrow \alpha_2} e_2$$

*Proof.* Use Lemma 13 and the proof for (ext) in STLC to see this holds

## D Soundness of Rules

In this section soundness proofs of the rules are introduced.

Given the logic is limited to compile-time syntax, the condition then for any term s.t.  $\mathbb{I}\Gamma \Vdash M\alpha$  where  $\mathfrak{a}(M) = \emptyset$  is implied then clearly  $(\mathfrak{a}(\xi), M\xi) \Downarrow$   $(\mathfrak{a}(\xi), G', V) \leftrightarrow M \xrightarrow{[\mathbb{I}, \xi]} V$  hence this will be used equivalently for brevity.

### D.1 Soundness of [VAR]

$\overline{\{A[x/m]\} x :_m \{A\}}$  <sup>[VAR]</sup> Proof:

- |   |  |                            |
|---|--|----------------------------|
| 1 | Assume: $\mathbb{I}\Gamma$ s.t. $\mathbb{I}\Gamma \Vdash \{A[x/m]\} x :_m \{A\}$ s.t.  |                            |
| 2 | Assume $\xi$ s.t. $\mathbb{I}\Gamma \triangleright \xi$ and $\xi \models A[x/m]$   |                            |
| 3 | $\rightarrow x \xrightarrow{[\mathbb{I}, \xi]} \xi(x)$   | $x\xi \equiv \xi(x)$       |
| 4 | $\xi \models A[x/m] \leftrightarrow \xi \cdot m : \xi(x) \models A$  | <i>Sem. [x/m], m-fresh</i> |
| 5 | $\xi \models A[x/m] \rightarrow x \xrightarrow{[\mathbb{I}, \xi]} \xi(x) \wedge \xi \cdot m : \xi(x) \models A$  | $\xi(x)$ a value           |
| 6 | $\forall \xi. \mathbb{I}\Gamma \triangleright \xi \rightarrow \xi \models A[x/m] \rightarrow x \xrightarrow{[\mathbb{I}, \xi]} \xi(x) \wedge \xi \cdot m : \xi(x) \models A$ | 2, 5                       |
| 7 | Hence: $\overline{\{A[x/m]\} x :_m \{A\}}$   | <i>Sem. valid triple</i>   |

### D.2 Soundness of [CONST]

$\overline{\{A[c/m]\} c :_m \{A\}}$  <sup>[CONST]</sup>

Proof:

- |   |  |                          |
|---|--|--------------------------|
| 1 | Assume: $\mathbb{I}\Gamma$ s.t. $\mathbb{I}\Gamma \Vdash \{A[c/m]\} c :_m \{A\}$   |                          |
| 2 | Assume $\xi$ s.t. $\mathbb{I}\Gamma \triangleright \xi$ and $\xi \models A[c/m]$   |                          |
| 3 | $\rightarrow c \xrightarrow{[\mathbb{I}, \xi]} c$  | $c\xi \equiv c$          |
| 4 | $\xi \models A[c/m] \leftrightarrow \xi \cdot m : c \models A$   | <i>Sem. [c/m]</i>        |
| 5 | $\xi \models A[c/m] \rightarrow c \xrightarrow{[\mathbb{I}, \xi]} c \wedge \xi \cdot m : c \models A$  | $c$ a value              |
| 6 | $\forall \xi. \mathbb{I}\Gamma \triangleright \xi \rightarrow \xi \models A[c/m] \rightarrow c \xrightarrow{[\mathbb{I}, \xi]} c \wedge \xi \cdot m : c \models A$ | 1, 5                     |
| 7 | Hence: $\overline{\{A[c/m]\} c :_m \{A\}}$   | <i>Sem. valid triple</i> |



**D.3 Soundness of [Eq]**

$$\frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{C[m = n/u]\} \quad C \text{ thin w.r.t } m, n}{\{A\} M = N :_u \{C\}} \text{ [Eq]}$$

Proof:

1	Assume: $\forall \mathbb{I}_0, \xi_1^{\mathbb{I}_1}. \mathbb{I}_0 \Vdash \{A\} M :_m \{B\} \wedge \mathbb{I}_0 \triangleright \xi_1 \rightarrow \xi_1 \models \{A\} M :_m \{B\}$	<i>IH(1)</i>
2	Assume: $\forall \mathbb{I}_0, \xi_1^{\mathbb{I}_1}. \mathbb{I}_0 \Vdash \{B\} N :_n \{C[m = n/u]\} \wedge \mathbb{I}_0 \triangleright \xi_1 \rightarrow \xi_1 \models \{B\} N :_n \{C[m = n/u]\}$	<i>IH(2)</i>
3	Assume: $\mathbb{I}$ s.t. $\mathbb{I} \Vdash \{A\} M = N :_u \{C\}$	
4	Assuming the typing constraints of this rule hold:	
5	Assume $\xi$ s.t., $\mathbb{I} \triangleright \xi \wedge \xi \models A$	
6	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G_m, V_m) \wedge \xi \cdot m : V_m \models B$	<i>IH(1)</i>
7	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G_m, V_m) \wedge (\hat{\mathfrak{a}}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\hat{\mathfrak{a}}(\xi \cdot m : V_m), G_n, V_n) \wedge \xi \cdot m : V_m \cdot n : V_n \models C[m = n/u]$	<i>IH(2)</i>
8	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G_m, V_m) \wedge (\hat{\mathfrak{a}}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\hat{\mathfrak{a}}(\xi \cdot m : V_m), G_n, V_n) \wedge (m = n) \overset{[\mathbb{I}+m+n, \xi \cdot m : V_m \cdot n : V_n]}{\rightsquigarrow} V_u \wedge \xi_{mn} \cdot u : V_u \models C$	<i>Sem. C[m = n/u]</i>
9	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G_m, V_m) \wedge (\hat{\mathfrak{a}}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\hat{\mathfrak{a}}(\xi \cdot m : V_m), G_n, V_n) \wedge (m = n) \overset{[\mathbb{I}+m+n, \xi \cdot m : V_m \cdot n : V_n]}{\rightsquigarrow} V_u \wedge \xi \cdot u : V_u \models C$	<i>C thin w.r.t m, n</i>
10	$\rightarrow (\hat{\mathfrak{a}}(\xi), (M = N)\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G', V_u) \wedge \xi \cdot u : V_u \models C$	<i>Op. Sem.(=)</i>
11	$\rightarrow \forall \xi. \mathbb{I} \triangleright \xi \rightarrow \xi \models \{A\} M = N :_u \{C\}$	<i>Assumption, 5-10</i>
12	Hence: $\frac{\models \{A\} M :_m \{B\} \quad \models \{B\} N :_n \{C^{-m,n}[m = n/u]\} \quad C \text{ thin w.r.t } m, n}{\models \{A\} M = N :_u \{C\}}$	<i>lines 1-11</i>

**D.4 Soundness of [GENSYM]**

$$\overline{\{\mathsf{T}\} \text{gensym} :_u \{\forall \delta. u \bullet () = m\{m\#\delta\}\}}^{\text{[GENSYM]}}$$

Soundness proof:

1	Assume: $\mathbb{I}\Gamma$ s.t. $\mathbb{I}\Gamma \Vdash \{\mathsf{T}\} \text{gensym} :_u \{\forall \delta. u \bullet () = m\{m\#\delta\}\}$	
2	Assume: $\xi$ s.t. $\mathbb{I}\Gamma \triangleright \xi \wedge \xi \models \mathsf{T}$	
3	Let: $\xi_u^{\mathbb{I}\Gamma} \equiv \xi \cdot u : \text{gensym}$	<i>gensym-value and <math>u \notin \text{dom}(\xi)</math></i>
4	$\forall \xi_x^{\mathbb{I}\Gamma}. (\hat{\mathbf{a}}(\xi_x), \text{gensym}()) \Downarrow ((\hat{\mathbf{a}}(\xi_x), r), r) \wedge r \notin \hat{\mathbf{a}}(\xi_x)$	<i>Op. Sem. gensym()</i>
5	$\forall \xi_1^{\mathbb{I}\Gamma_1}. \xi \cdot u : \text{gensym} \preceq^* \xi_1$ $\rightarrow (\hat{\mathbf{a}}(\xi_1), \llbracket u \rrbracket_{\xi_1}()) \Downarrow ((\hat{\mathbf{a}}(\xi_1), r), r)$ $\wedge r \notin \hat{\mathbf{a}}(\xi_1)$	<i>restrict the quantification, 4</i> $\llbracket u \rrbracket_{\xi_1} \equiv \text{gensym}$
6	$\rightarrow \forall \xi_1^{\mathbb{I}\Gamma_1}. \xi \cdot u : \text{gensym} \preceq^* \xi_1$ $\rightarrow (\hat{\mathbf{a}}(\xi_1), \llbracket u \rrbracket_{\xi_1}()) \Downarrow ((\hat{\mathbf{a}}(\xi_1), r), r)$ $\wedge \neg \exists M_m^{\text{Nm}}. M_m \overset{[\mathbb{I}\Gamma_1, \xi_1 \cdot m : r]}{\rightsquigarrow} r$	<i>Lemma 23</i>
7	$\rightarrow \forall \xi_1^{\mathbb{I}\Gamma_1}. \xi \cdot u : \text{gensym} \preceq^* \xi_1$ $\rightarrow (\hat{\mathbf{a}}(\xi_1), \llbracket u \rrbracket_{\xi_1}()) \Downarrow ((\hat{\mathbf{a}}(\xi_1), r), r)$ $\wedge \neg \exists M_m^{\text{Nm}}. M_m \overset{[\mathbb{I}\Gamma_1, \xi_1 \cdot m : r]}{\rightsquigarrow} \llbracket m \rrbracket_{\xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r}$	$r \equiv \llbracket m \rrbracket_{\xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r}$
8	$\rightarrow \forall \xi_1^{\mathbb{I}\Gamma_1}. \xi \cdot u : \text{gensym} \preceq^* \xi_1$ $\rightarrow (\hat{\mathbf{a}}(\xi_1), \llbracket u \rrbracket_{\xi_1}()) \Downarrow ((\hat{\mathbf{a}}(\xi_1), r), r)$ $\wedge \neg \exists M_m^{\text{Nm}}. M_m \overset{[\mathbb{I}\Gamma_1, \xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r]}{\rightsquigarrow} \llbracket m \rrbracket_{\xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r}$	<i>Lemma 10</i>
9	$\rightarrow \forall \xi_1^{\mathbb{I}\Gamma_1}. \xi \cdot u : \text{gensym} \preceq^* \xi_1$ $\rightarrow (\hat{\mathbf{a}}(\xi_1), \llbracket u \rrbracket_{\xi_1}()) \Downarrow ((\hat{\mathbf{a}}(\xi_1), r), r)$ $\wedge \neg \exists M_m^{\text{Nm}}. M_m \overset{[\mathbb{I}\Gamma_1 + \delta, \xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r]}{\rightsquigarrow} \llbracket m \rrbracket_{\xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r}$	$\llbracket \mathbb{I}\Gamma_1 \rrbracket_{\xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r} \equiv \llbracket \mathbb{I}\Gamma_1 + \delta \rrbracket_{\xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r}$
10	$\forall \xi_1^{\mathbb{I}\Gamma_1}. \xi \cdot u : \text{gensym} \preceq^* \xi_1$ $\rightarrow (\hat{\mathbf{a}}(\xi_1), \llbracket u \rrbracket_{\xi_1}()) \Downarrow ((\hat{\mathbf{a}}(\xi_1), r), r)$ $\wedge \xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV} \cdot m : r \models m\#\delta$	<i>Sem. #</i>
11	$\forall \xi_1^{\mathbb{I}\Gamma_1}. \xi \cdot u : \text{gensym} \preceq^* \xi_1 \wedge \xi_d \equiv \xi_1 \cdot \delta : \mathbb{I}\Gamma_1 \setminus \text{TCV}$ $\rightarrow (\hat{\mathbf{a}}(\xi_d), \llbracket u \rrbracket_{\xi_d}()) \Downarrow ((\hat{\mathbf{a}}(\xi_d), r), r)$ $\wedge \xi_d \cdot m : r \models m\#\delta$	<i>Lemma 11</i> $\hat{\mathbf{a}}(\xi_d) \equiv \hat{\mathbf{a}}(\xi)$ <i>(Shorthand <math>\delta</math>)</i>
12	$\xi \cdot u : \text{gensym} \models \forall \delta. u \bullet () = m\{m\#\delta\}$	<i>Sem. <math>\forall \delta. u \bullet () = m\{m\#\delta\}</math></i>
13	$\forall \xi^{\mathbb{I}\Gamma}. \mathbb{I}\Gamma \triangleright \xi \rightarrow \xi \models \mathsf{T} \rightarrow (\hat{\mathbf{a}}(\xi), \text{gensym}()) \Downarrow (\hat{\mathbf{a}}(\xi), \text{gensym})$ $\wedge \xi \cdot u : \text{gensym} \models \forall \delta. u \bullet () = m\{m\#\mathbb{I}\Gamma + u + \delta\}$	<i>2-12</i>
14	Hence: $\overline{\{\mathsf{T}\} \text{gensym} :_u \{\forall \delta. u \bullet () = m\{m\#\delta\}\}}$	<i>Sem. valid triple</i>

**D.5 Soundness of [LAM]**

	$\frac{A - \text{Ext-Ind} \quad (\mathbb{I}\Gamma + \delta + x : \alpha) \Vdash \{A^{-x} \wedge B\} M :_m \{C\}}{\mathbb{I}\Gamma \Vdash \{A\} \lambda x^\alpha . M :_u \{\forall \delta . \forall x^\alpha \in (\delta) . (B \rightarrow u \bullet x = m\{C\})\}} \text{[LAM]}$	
1	Assume: $\mathbb{I}\Gamma$ s.t. $\mathbb{I}\Gamma \Vdash \{A\} \lambda x^\alpha . M :_u \{\forall \delta . \forall x^\alpha \in (\delta) . (B \rightarrow u \bullet x = m\{C\})\}$	
2	and $\mathbb{I}\Gamma + \delta + x \Vdash \{A^{-x} \wedge B\} M :_m \{C\}$	
3	Assume: $\forall \xi_0^{\mathbb{I}\Gamma_1 + \delta + x} . \mathbb{I}\Gamma + \delta + x \triangleright \xi_0 \wedge \xi_0 \models A^{-x} \wedge B \rightarrow (\hat{\mathfrak{a}}(\xi_0), M\xi_0) \Downarrow (G', V_m)$ $\wedge \xi_0 \cdot m : V_m \models C$	<i>IH(1)</i>
4	$\forall \xi^{\mathbb{I}\Gamma_1} . \mathbb{I}\Gamma \triangleright \xi \rightarrow \xi \models A \rightarrow (G, (\lambda x^\alpha . M)\xi) \Downarrow (G, G', V_u)$ $\wedge \xi \cdot u : V_u \models \forall \delta . \forall x^\alpha \in (\delta) . (B \rightarrow u \bullet x = m\{C\})$	<i>Need to Prove</i>
5	Assume: $\xi^{\mathbb{I}\Gamma_1}$ s.t. $\mathbb{I}\Gamma \triangleright \xi \wedge \xi \models A$	<i>Assume</i>
6	$\rightarrow (\hat{\mathfrak{a}}(\xi), (\lambda x . M)\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), \lambda x . (M\xi \setminus x))$	<i>Op. Sem. <math>\lambda x . M</math></i>
7	For brevity write: $\xi_{uddx} \equiv \wedge \xi_{ud} \cdot \delta : \mathbb{I}\Gamma_{ud} \setminus \text{-TCV} \cdot x : V_x$	
8	$\rightarrow (\hat{\mathfrak{a}}(\xi), (\lambda x . M)\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), \lambda x . (M\xi \setminus x))$ $\wedge \forall \xi_{ud}^{\mathbb{I}\Gamma_{ud}} . \xi \cdot u : \lambda x . (M\xi \setminus x) \preceq^* \xi_{ud}$ $\wedge \forall P_x . P_x \overset{[\mathbb{I}\Gamma_{ud} + \delta, \xi_{ud} \cdot \delta : \mathbb{I}\Gamma_{ud} \setminus \text{-TCV}]}{\rightsquigarrow} V_x$ $\wedge \xi_{uddx} \models B \rightarrow \xi_{uddx} \models B$	<i>Assume some <math>u, \delta, x</math>-extension that models <math>B \rightarrow B</math></i> <i>Tautology</i>
9	$\rightarrow (\hat{\mathfrak{a}}(\xi), (\lambda x . M)\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), \lambda x . (M\xi \setminus x))$ $\wedge \forall \xi_{ud}^{\mathbb{I}\Gamma_{ud}} . \xi \cdot u : \lambda x . (M\xi \setminus x) \preceq^* \xi_{ud}$ $\wedge \forall P_x . P_x \overset{[\mathbb{I}\Gamma_{ud} + \delta, \xi_{ud} \cdot \delta : \mathbb{I}\Gamma_{ud} \setminus \text{-TCV}]}{\rightsquigarrow} V_x$ $\wedge \xi_{uddx} \models B \rightarrow \xi_{uddx} \models A \wedge B$	<i>line 5 (<math>\xi \models A</math>)</i> $\xi \preceq^* \xi_{uddx}$ <i>A-Ext-Ind Sem. <math>\wedge</math></i>
10	$\rightarrow (\hat{\mathfrak{a}}(\xi), (\lambda x . M)\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), \lambda x . (M\xi \setminus x))$ $\wedge \forall \xi_{ud}^{\mathbb{I}\Gamma_{ud}} . \xi \cdot u : \lambda x . (M\xi \setminus x) \preceq^* \xi_{ud}$ $\wedge \forall P_x . P_x \overset{[\mathbb{I}\Gamma_{ud} + \delta, \xi_{ud} \cdot \delta : \mathbb{I}\Gamma_{ud} \setminus \text{-TCV}]}{\rightsquigarrow} V_x$ $\wedge \xi_{uddx} \models B \rightarrow (\hat{\mathfrak{a}}(\xi_{uddx}), M\xi_{uddx}) \Downarrow (\hat{\mathfrak{a}}(\xi_{uddx}), G_m, V_m)$ $\wedge \xi_{uddx} \cdot m : V_m \models C$	<i>IH(1)</i> $\mathbb{I}\Gamma + u + \delta + x \triangleright \xi_{uddx}$
11	$\rightarrow (\hat{\mathfrak{a}}(\xi), (\lambda x . M)\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), \lambda x . (M\xi \setminus x))$ $\wedge \forall \xi_{ud}^{\mathbb{I}\Gamma_{ud}} . \xi \cdot u : \lambda x . (M\xi \setminus x) \preceq^* \xi_{ud}$ $\wedge \forall P_x . P_x \overset{[\mathbb{I}\Gamma_{ud} + \delta, \xi_{ud} \cdot \delta : \mathbb{I}\Gamma_{ud} \setminus \text{-TCV}]}{\rightsquigarrow} V_x$ $\wedge \xi_{uddx} \models B \rightarrow (\hat{\mathfrak{a}}(\xi_{uddx}), \llbracket u \rrbracket_{\xi_{uddx}} \llbracket x \rrbracket_{\xi_{uddx}}) \Downarrow (\hat{\mathfrak{a}}(\xi_{uddx}), G_m, V_m)$ $\wedge \xi_{uddx} \cdot m : V_m \models C$	$M\xi \equiv ((\lambda x . M)x)\xi \equiv \llbracket u \rrbracket_{\xi_{uddx}} \llbracket x \rrbracket_{\xi_{uddx}}$
12	$\rightarrow (\hat{\mathfrak{a}}(\xi), (\lambda x . M)\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), \lambda x . (M\xi \setminus x))$ $\wedge \forall \xi_{ud}^{\mathbb{I}\Gamma_{ud}} . \xi \cdot u : \lambda x . (M\xi \setminus x) \preceq^* \xi_{ud}$ $\wedge \forall P_x . P_x \overset{[\mathbb{I}\Gamma_{ud} + \delta, \xi_{ud} \cdot \delta : \mathbb{I}\Gamma_{ud} \setminus \text{-TCV}]}{\rightsquigarrow} V_x \rightarrow \xi_{uddx} \models B \rightarrow u \bullet x = m\{C\}$	<i>Sem. <math>u \bullet x = m\{C\}, \rightarrow</math></i> $\wedge$ <i>implies <math>\rightarrow</math></i>
13	$\rightarrow (\hat{\mathfrak{a}}(\xi), (\lambda x . M)\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), V_u) \wedge \xi \cdot u : V_u \models \forall \delta . \forall x \in (\delta) . B \rightarrow u \bullet x = m\{C\}$	<i>Sem. <math>\forall \delta . \forall x \in (\delta)</math>.</i>
14	$\models \{A\} \lambda x . M :_u \{\forall \delta . \forall x \in (\mathbb{I}\Gamma + u + \delta) . (B \rightarrow u \bullet x = m\{C\})\}$	<i>line 5, Sem. <math>\{A\} M :_u \{B\}</math></i>
15	Hence: $\frac{A - \text{Ext-Ind} \quad \models \{A \wedge B\} M :_m \{C\}}{\models \{A\} \lambda x . M :_u \{\forall \delta . \forall x \in (\mathbb{I}\Gamma + u + \delta) . (B \rightarrow u \bullet x = m\{C\})\}}$	<i>lines 1-13</i>

**D.6 Soundness of [APP]**

$$\frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{m \bullet n = u\{C\}\} \quad C - \text{thin w.r.t } m, n}{\{A\} MN :_u \{C\}} \text{ [APP]}$$

Proof:

1	Assume: $\mathbb{I}$ s.t. $\mathbb{I} \Vdash \{A\} MN :_u \{C\}$	
2	$\rightarrow \mathbb{I} \Vdash \{A\} M :_m \{B\} \wedge \mathbb{I} + m \Vdash \{B\} N :_n \{m \bullet n = u\{C\}\}$	
3	$\mathbb{I} \downarrow_{-TC} \vdash MN : \beta \rightarrow \mathbb{I} \downarrow_{-TC} \vdash M : \alpha \rightarrow \beta \wedge \mathbb{I} \downarrow_{-TC} \vdash N : \alpha$	
4	$\forall \xi_1^{\mathbb{I}^1}. \mathbb{I} \triangleright \xi_1 \rightarrow \xi_1 \models A \rightarrow (\mathfrak{a}(\xi_1), M\xi_1) \Downarrow (\mathfrak{a}(\xi_1), G_m, V_m) \wedge \xi_1 \cdot m : V_m \models B$	<i>IH(1)</i>
5	$\forall \xi_2^{\mathbb{I}^2}. \mathbb{I} + m \triangleright \xi_2 \rightarrow \xi_2 \models B \rightarrow (\mathfrak{a}(\xi_2), N\xi_2) \Downarrow (\mathfrak{a}(\xi_2), G_n, V_n) \wedge \xi_2 \cdot n : V_n \models m \bullet n = u\{C\}$	<i>IH(2)</i>
6	Assume: $\xi^{\mathbb{I}^0}$ s.t. $\mathbb{I} \triangleright \xi \wedge \xi \models A$	
7	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge \xi \cdot m : V_m \models B$	<i>IH(1)</i>
8	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge (\mathfrak{a}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\mathfrak{a}(\xi \cdot m : V_m), G_n, V_n) \wedge \xi \cdot m : V_m \cdot n : V_n \models m \bullet n = u\{C\}$	<i>IH(2)</i>
9	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge (\mathfrak{a}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\mathfrak{a}(\xi \cdot m : V_m), G_n, V_n) \wedge (\mathfrak{a}(\xi \cdot m : V_m \cdot n : V_n), \llbracket m \rrbracket_{\xi \cdot m : V_m \cdot n : V_n} \llbracket n \rrbracket_{\xi \cdot m : V_m \cdot n : V_n}) \Downarrow (G''', V_u) \wedge \xi \cdot m : V_m \cdot n : V_n \cdot u : V_u \models C$	<i>Sem. <math>\bullet = \{\}</math></i>
10	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge (\mathfrak{a}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\mathfrak{a}(\xi \cdot m : V_m), G_n, V_n) \wedge (\mathfrak{a}(\xi \cdot m : V_m \cdot n : V_n), V_m V_n) \Downarrow (G''', V_u) \wedge \xi \cdot m : V_m \cdot n : V_n \cdot u : V_u \models C$	<i>Sem. <math>\llbracket x \rrbracket_\xi</math></i>
11	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge (\mathfrak{a}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\mathfrak{a}(\xi \cdot m : V_m), G_n, V_n) \wedge (\mathfrak{a}(\xi \cdot m : V_m \cdot n : V_n), V_m V_n) \Downarrow (G''', V_u) \wedge \xi \cdot u : V_u \models C$	<i>C - thin w.r.t m, n</i>
12	$\rightarrow (\mathfrak{a}(\xi), (MN)\xi) \Downarrow (G''', V_u) \wedge \xi \cdot u : V_u \models C$	<i>Op. Sem. (App)</i>
13	$\rightarrow \forall \xi^{\mathbb{I}^0}. \mathbb{I} \triangleright \xi \rightarrow \xi \models A \rightarrow (\mathfrak{a}(\xi), (MN)\xi) \Downarrow (G''', V_u) \wedge \xi \cdot u : V_u \models C$	<i>Assumption, line 6</i>
14	Hence: $\frac{\models \{A\} M :_m \{B\} \quad \models \{B\} N :_n \{m \bullet n = u\{C\}\} \quad C - \text{thin w.r.t } m, n}{\models \{A\} MN :_u \{C\}}$	<i>lines 1-13</i>

**D.7 Soundness of [PAIR]**

$$\frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{C[\langle m, n \rangle / u]_{\mathbb{I}+m+n}\} \quad C \text{ thin w.r.t } x}{\{A\} \langle M, N \rangle :_u \{C\}} \text{ [PAIR]}$$

Proof:

1	Assume: $\mathbb{I} \text{ s.t. } \mathbb{I} \Vdash \{A\} \langle M, N \rangle :_u \{C[\langle m, n \rangle / u]_{\mathbb{I}+m+n}\}$	
2	$\rightarrow \mathbb{I} \Vdash \{A\} M :_m \{B\} \wedge \mathbb{I} + m \Vdash \{B\} N :_n \{C[\langle m, n \rangle / u]_{\mathbb{I}+m+n}\} \wedge \mathbb{I} \vdash N : \alpha$	
3	$\forall \mathbb{I}', \xi^{\mathbb{I}'_0}. \mathbb{I}' \triangleright \xi \rightarrow \xi \models A \rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge \xi \cdot m : V_m \models B$	<i>IH(1)</i>
4	$\forall \mathbb{I}', \xi^{\mathbb{I}'_0+m}. \mathbb{I}' + m \triangleright \xi_m \rightarrow \xi_m \models B \rightarrow (\mathfrak{a}(\xi_m), N\xi_m) \Downarrow (\mathfrak{a}(\xi_m), G_n, V_n) \wedge \xi_n \cdot n : V_n \models C[\langle m, n \rangle / u]_{\mathbb{I}+m+n}$	<i>IH(2)</i>
5	Assume: $\xi^{\mathbb{I}'_0} \text{ s.t. } \mathbb{I}' \triangleright \xi \wedge \xi \models A$	
6	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge \xi \cdot m : V_m \models B$	<i>IH(1)</i>
7	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge (\mathfrak{a}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\mathfrak{a}(\xi \cdot m : V_m), G_n, V_n) \wedge \xi \cdot m : V_m \cdot n : V_n \models C[\langle m, n \rangle / u]_{\mathbb{I}+m+n}$	<i>IH(2)</i>
8	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge (\mathfrak{a}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\mathfrak{a}(\xi \cdot m : V_m), G_n, V_n) \wedge (\mathfrak{a}(\xi \cdot m : V_m \cdot n : V_n), \llbracket \langle m, n \rangle \rrbracket_{\xi \cdot m : V_m \cdot n : V_n}) \Downarrow (G''', V_u) \wedge \xi \cdot m : V_m \cdot n : V_n \cdot u : V_u \models C$	<i>Sem. [e/x], u ∉ dom(ξ · m · n)</i>
9	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge (\mathfrak{a}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\mathfrak{a}(\xi \cdot m : V_m), G_n, V_n) \wedge (\mathfrak{a}(\xi \cdot m : V_m \cdot n : V_n), \langle V_m, V_n \rangle) \Downarrow (G''', V_u) \wedge \xi \cdot m : V_m \cdot n : V_n \cdot u : V_u \models C$	<i>Sem. [e]<sub>ξ</sub></i>
10	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge (\mathfrak{a}(\xi \cdot m : V_m), N(\xi \cdot m : V_m)) \Downarrow (\mathfrak{a}(\xi \cdot m : V_m), G_n, V_n) \wedge (\mathfrak{a}(\xi \cdot m : V_m \cdot n : V_n), \langle V_m, V_n \rangle) \Downarrow (G''', V_u) \wedge \xi \cdot u : V_u \models C$	<i>C thin w.r.t m, n</i>
11	$\rightarrow (\mathfrak{a}(\xi), \langle M, N \rangle \xi) \Downarrow (G''', V_u) \wedge \xi \cdot u : V_u \models C$	<i>Op. Sem. (Pair)</i>
12	$\rightarrow \forall \xi^{\mathbb{I}'_0}. \mathbb{I}' \triangleright \xi \rightarrow \xi \models A \rightarrow (\mathfrak{a}(\xi), \langle M, N \rangle \xi) \Downarrow (G''', V_u) \wedge \xi \cdot u : V_u \models C$	<i>Assumption, (line 5)</i>
13	Hence: $\frac{\models \{A\} M :_m \{B\} \quad \models \{B\} N :_n \{C[\langle m, n \rangle / u]_{\mathbb{I}+m+n}\} \quad C \text{ thin w.r.t } m, n}{\models \{A\} \langle M, N \rangle :_u \{C\}}$	<i>lines 1- 12</i>

**D.8 Soundness of [PROJ(i)]**

$$\frac{\{A\} M :_m \{C[\pi_i(m)/u]_{\mathcal{I}+m}\} \quad C \text{ thin w.r.t } m}{\{A\} \pi_i(M) :_u \{C\}} \text{ [PROJ(i)]}$$

Proof:

1	Assume: $\mathcal{I} \text{ s.t. } \mathcal{I} \Vdash \{A\} \pi_i(M) :_u \{C\}$	
2	$\rightarrow \mathcal{I} \Vdash \{A\} M :_m \{C[\pi_i(m)/u]_{\mathcal{I}+m}\}$	<i>Typing rules</i>
3	$\forall \xi^{\mathcal{I}_0}. \mathcal{I} \triangleright \xi \rightarrow \xi \models A \rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge \xi \cdot m : V_m \models C[\pi_i(m)/u]_{\mathcal{I}+m}$	<i>IH(1)</i>
4	Assume: $\xi^{\mathcal{I}_0} \text{ s.t. } \mathcal{I} \triangleright \xi \wedge \xi \models A$	
5	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m) \wedge \xi \cdot m : V_m \models C[\pi_i(m)/u]_{\mathcal{I}+m}$	<i>IH(1)</i>
6	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m)$ $\wedge (\mathfrak{a}(\xi \cdot m : V_m), \llbracket \pi_i(m) \rrbracket_{\xi \cdot m : V_m}) \Downarrow (G''', V_u)$ $\wedge \xi \cdot m : V_m \cdot u : V_u \models C$	<i>Sem. [e/x],</i> $u \notin \text{dom}(\xi \cdot m \cdot n)$
7	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m)$ $\wedge (\mathfrak{a}(\xi \cdot m : V_m), \pi_i(V_m)) \Downarrow (G''', V_u)$ $\wedge \xi \cdot m : V_m \cdot u : V_u \models C$	<i>Sem. [e]<math>_{\xi}</math></i>
8	$\rightarrow (\mathfrak{a}(\xi), M\xi) \Downarrow (\mathfrak{a}(\xi), G_m, V_m)$ $\wedge (\mathfrak{a}(\xi \cdot m : V_m), \pi_i(V_m)) \Downarrow (G''', V_u)$ $\wedge \xi \cdot u : V_u \models C$	<i>C thin w.r.t m</i>
9	$\rightarrow (\mathfrak{a}(\xi), \pi_i(M)\xi) \Downarrow (G''', V_u) \wedge \xi \cdot u : V_u \models C$	<i>Op. Sem. (Pair)</i>
10	$\rightarrow \forall \xi^{\mathcal{I}_0}. \mathcal{I} \triangleright \xi \rightarrow \xi \models A \rightarrow (\mathfrak{a}(\xi), \langle M, N \rangle \xi) \Downarrow (G''', V_u)$ $\wedge \xi \cdot u : V_u \models C$	<i>Assumption, (line 4)</i>
11	Hence: $\frac{\models \{A\} M :_m \{C[\pi_i(m)/u]_{\mathcal{I}+m}\} \quad C \text{ thin w.r.t } m}{\models \{A\} \pi_i(M) :_u \{C\}}$	<i>lines 1-10</i>

**D.9 Soundness of [IF]**

$$\frac{\{A\} M :_m \{B\} \quad \{B[b_i/m]\} N_i :_u \{C\} \quad b_1 = \text{true} \quad b_2 = \text{false} \quad i = 1, 2}{\{A\} \text{if } M \text{ then } N_1 \text{ else } N_2 :_u \{C\}} \text{ [IF]}$$

Proof: standard. Note that substitution is equivalent to standard substitution for  $b_i$ -values of type Bool.

**D.10 Soundness of [LET]**

$$\frac{\{A\} M :_x \{B\} \quad \{B\} N :_u \{C\} \quad C \text{ thin w.r.t } x}{\{A\} \text{ let } x = M \text{ in } N :_u \{C\}} \text{ [LET]}$$

1	Assume: $\mathbb{I} \text{ s.t. } \mathbb{I} \Vdash \{A\} \text{ let } x = M \text{ in } N :_u \{C\}$	
2	$\rightarrow \mathbb{I} \downarrow_{-TC} \vdash M : \alpha \wedge \mathbb{I} \downarrow_{-TC}, x : \alpha \vdash N : \beta$	<i>Typing terms</i>
3	Assume: $\forall \xi^{\mathbb{I}^0}. \mathbb{I} \triangleright \xi \rightarrow \xi \models A \rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (G'_m, V_m) \wedge \xi \cdot x : V_m \models B$	<i>IH(1)</i>
4	Assume: $\forall \xi_x^{\mathbb{I}^0+x:\alpha}. \mathbb{I} + x \triangleright \xi_x \rightarrow \xi_x \models B \rightarrow (\hat{\mathfrak{a}}(\xi_x), N\xi_x) \Downarrow (G'_n, V_n) \wedge \xi_x \cdot u : V_n \models C$	<i>IH(2)</i>
5	Assume: $\xi^{\mathbb{I}^0} \text{ s.t. } \mathbb{I} \triangleright \xi \wedge \xi \models A$	
6	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (G', V_m) \wedge \xi \cdot x : V_m \models B$	<i>IH(1)</i>
7	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (G', V_m)$ $\wedge (\hat{\mathfrak{a}}(\xi \cdot x : V_m), N(\xi \cdot x : V_m)) \Downarrow (G'_n, V_n)$ $\wedge \xi \cdot x : V_m \cdot u : V_n \models C$	<i>IH(2)</i>
8	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (G', V_m)$ $\wedge (\hat{\mathfrak{a}}(\xi \cdot x : V_m), N(\xi \cdot x : V_m)) \Downarrow (G'_n, V_n)$ $\wedge \xi \cdot u : V_n \models C$	<i>C thin w.r.t x</i>
9	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (G', V_m)$ $\wedge (\hat{\mathfrak{a}}(\xi \cdot x : V_m), N[V_m/x]\xi) \Downarrow (G'_n, V_n)$ $\wedge \xi \cdot u : V_n \models C$	<i>Sem. [V<sub>m</sub>/x]</i>
10	$\rightarrow (\hat{\mathfrak{a}}(\xi), (\text{let } x = M \text{ in } N)\xi) \Downarrow (G'', V_n) \wedge \xi \cdot u : V_n \models C$	<i>Op. Sem. let x = M in N</i>
11	$\forall \xi. \mathbb{I} \triangleright \xi \rightarrow \xi \models \{A\} \text{ let } x = M \text{ in } N :_m \{C\}$	<i>Sem. valid triple</i>
12	Hence: $\frac{\models \{A\} M :_x \{B\} \quad \models \{B\} N :_u \{C\} \quad C \text{ thin w.r.t } x}{\models \{A\} \text{ let } x = M \text{ in } N :_u \{C\}}$	<i>lines 1-11</i>

**D.11 Soundness of derived rule [LETFRESH]**

$$\frac{A - \text{Ext-Ind} \quad \Gamma + x \Vdash \{A \wedge x \# \Gamma\} \quad M :_m \{C\} \quad C \text{ thin w.r.t } x}{\Gamma \Vdash \{A\} \text{ let } x = \text{gensym } () \text{ in } M :_m \{C\}} \text{[LETFRESH]}$$

1	Assume: $\Gamma$ s.t. $\Gamma \Vdash \{A\} \text{ let } x = \text{gensym } () \text{ in } N :_u \{C\}$	
2	$\Gamma \Vdash \{\top\} \text{ gensym } () :_x \{x \# \Gamma\}$	<i>See Example 1</i>
3	$\Gamma \Vdash \{A\} \text{ gensym } () :_x \{A \wedge x \# \Gamma\}$	<i>[INVAR], 7, A – Ext-Ind</i>
4	$\Gamma + x \Vdash \{A \wedge x \# \Gamma\} \quad M :_m \{C\}$	<i>Assumption</i>
5	$C - \text{ thin w.r.t } x$	<i>Assumption</i>
6	$\Gamma \Vdash \{A\} \text{ let } x = \text{gensym } () \text{ in } M :_m \{C\}$	<i>[LET], 3, 4, 5</i>
7	Hence: $\frac{A - \text{Ext-Ind} \quad \Vdash \{A \wedge x \# \Gamma\} \quad N :_u \{C\} \quad C \text{ thin w.r.t } x}{\Vdash \{A\} \text{ let } x = M \text{ in } N :_u \{C\}}$	<i>lines 1-6</i>



**D.12 Soundness of Key structural rules:**

**D.13 Soundness of structural rule [CONSEQ]**

$$\frac{A \rightarrow A' \quad \{A'\} M :_m \{B'\} \quad B' \rightarrow B}{\{A\} M :_m \{B\}} \text{ [CONSEQ]}$$

Proof, simply by application of the assumptions:

1	Assume $\mathbb{I}\Gamma$ , s.t. $\mathbb{I}\Gamma \Vdash \{A\} M :_m \{B\}$	
2	Assume typing holds for assumptions i.e. $\mathbb{I}\Gamma \Vdash A \rightarrow A' \wedge \mathbb{I}\Gamma \Vdash \{A'\} M^\alpha :_m \{B'\} \wedge \mathbb{I}\Gamma + m : \alpha \Vdash B' \rightarrow B$	
3	Assume: $\forall \xi_0^{\mathbb{I}\Gamma_0}. \mathbb{I}\Gamma \triangleright \xi_0 \rightarrow \xi_0 \Vdash A \rightarrow A'$	<i>IH(1)</i>
4	Assume: $\forall \xi_0^{\mathbb{I}\Gamma_0}. \mathbb{I}\Gamma \triangleright \xi_0 \rightarrow \xi_0 \Vdash \{A'\} M :_m \{B'\}$	<i>IH(2)</i>
5	Assume: $\forall \xi_1^{\mathbb{I}\Gamma_1}. \mathbb{I}\Gamma + m \triangleright \xi_1 \rightarrow \xi_1 \Vdash B' \rightarrow B$	<i>IH(3)</i>
6	Assume: $\xi^{\mathbb{I}\Gamma'}$ s.t. $\mathbb{I}\Gamma \triangleright \xi \wedge \xi \Vdash A$	
7	$\rightarrow \xi \Vdash A'$	<i>MP, IH(1)</i>
8	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G', V) \wedge \xi \cdot m : V \Vdash B'$	<i>MP, IH(2)</i>
9	$\rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G', V) \wedge \xi \cdot m : V \Vdash B$	<i>MP, IH(3)</i>
10	$\rightarrow \xi \Vdash A \rightarrow (\hat{\mathfrak{a}}(\xi), M\xi) \Downarrow (\hat{\mathfrak{a}}(\xi), G', V) \wedge \xi \cdot m : V \Vdash B$	<i>Assumption 6-9</i>
11	$\rightarrow \forall \xi^{\mathbb{I}\Gamma'}. \mathbb{I}\Gamma \triangleright \xi \rightarrow \xi \Vdash \{A\} M :_m \{B\}$	<i>Assumption 3-10</i>
12	Hence: $\frac{A \rightarrow A' \quad \Vdash \{A'\} M :_m \{B'\} \quad B' \rightarrow B}{\Vdash \{A\} M :_m \{B\}}$	<i>Assumption 1-11</i>

**D.14 Soundness of structural rule [INVAR]**

$$\frac{C - \text{Ext-Ind} \quad \{A\} M :_m \{B\}}{\{A \wedge C\} M :_m \{B \wedge C\}} \text{[INVAR]}$$

Proof:

1	Assume: $\mathbb{I}$ s.t. $\mathbb{I} \Vdash \{A \wedge C\} M :_m \{B \wedge C\}$	
2	$\rightarrow \mathbb{I} \Vdash \{A\} M :_m \{B\}$	
3	Assume: $\forall \xi^{\mathbb{I}^0}. \mathbb{I} \triangleright \xi \rightarrow \xi \models A \rightarrow (G, M\xi) \Downarrow (G, G', V) \wedge \xi \cdot m : V \models B$	<i>IH(1)</i>
4	Assume for some model $\xi^{\mathbb{I}^0}$ s.t. $\mathbb{I} \triangleright \xi \wedge \xi \models A \wedge C$	
5	$\xi \models A \wedge \xi \models C$	<i>Sem. <math>\wedge</math></i>
6	$(G, M\xi) \Downarrow (G, G', V) \wedge \xi \models C$ $\wedge \xi \cdot m : V \models B$	<i>IH(1)</i>
7	$(G, M\xi) \Downarrow (G, G', V)$ $\wedge \xi \cdot m : V \models B$ $\wedge \xi \cdot m : V \models C$	<i>C - Ext-Ind</i> <i>Sem. <math>\Downarrow</math> <math>\rightarrow M^{\{\mathbb{I}, \xi\}} V</math></i> <i>Sem. <math>\preceq^* \rightarrow \xi \preceq^* \xi \cdot m : V</math></i>
8	$(G, M\xi) \Downarrow (G, G', V) \wedge \xi \cdot m : V \models B \wedge C$	<i>Sem. <math>\wedge</math></i>
9	$\forall \xi^{\mathbb{I}^0}. \mathbb{I} \triangleright \xi \rightarrow \xi \models \{A \wedge C\} M :_m \{B \wedge C\}$	<i>lines 5-8</i>
10	Hence: $\frac{C - \text{Ext-Ind} \quad \models \{A\} M :_m \{B\}}{\models \{A \wedge C\} M :_m \{B \wedge C\}}$	<i>lines 1-9</i>

## E Ext-Ind Formulae Construction Lemmas

Syntactically define Ext-Ind formulae and how to construct them:

Some are not core constructions but specific instances used in the reasoning examples in Sec. 7.

A subset of all possible Ext-Ind formulae are defined as SYN-EXT-IND below and are proven as Ext-Ind in the lemma that follows.

**Definition 4 (A Syntactic classification of Ext-Ind formulae, written SYN-EXT-IND).** Define SYN-EXT-IND inductively as follows:

1.  $\top, \text{F}, e = e'$  and  $x \# \mathbb{I}'$  are all SYN-EXT-IND.
2. If  $C_1$  and  $C_2$  are Ext-Ind then  $\neg C_1, C_1 \wedge C_2, C_1 \vee C_2, C_1 \rightarrow C_2, u \bullet e = m^\alpha\{C_1\}, \exists x \in (\mathbb{I}').C_1, \forall x \in (\mathbb{I}').C_1$  are all SYN-EXT-IND.
3. If  $C_1$  is Ext-Ind and contains no reference to  $\delta$  then  $\forall \delta.C_1$  and  $\forall \delta.\forall x \in (\delta).C_1$  are SYN-EXT-IND.
4. Specific cases:  $\forall \delta.f \bullet () = b\{b\#\delta\}$  and  $\forall \delta.\forall x \in (\delta).f \bullet x = b\{b\#\delta+x\}$  are SYN-EXT-IND.

This is an incomplete characterisation of all Ext-Ind formulae, but covers all cases required for the proofs.

We introduce the following 2 lemmas for the specific cases in Def. 4

**Lemma 35 (Constructing Ext-Ind formulae from  $\forall \delta.\forall x \in (\delta).$ ).**

$$A^{-\delta} - \text{Ext-Ind} \rightarrow \forall \delta.\forall x \in (\delta).A^{-\delta} - \text{Ext-Ind}$$

1	Assume: $A^{-\delta} - \text{Ext-Ind}$	
2	$\leftrightarrow \forall \mathbb{I}, \xi_x^{\mathbb{I}+x:\alpha}, \xi_x^{\mathbb{I}',x:\alpha}. (\mathbb{I}+x : \alpha \Vdash A \wedge \xi_x \preceq^* \xi'_x) \rightarrow (\xi_x \models A \leftrightarrow \xi'_x \models A)$	
3	Show: $\forall \mathbb{I}, \xi^{\mathbb{I}}, \xi^{\mathbb{I}'}. \mathbb{I} \Vdash \forall \delta.\forall x \in (\delta).A^{-\delta} \wedge \xi \preceq^* \xi' \rightarrow \left( \begin{array}{l} \xi \models \forall \delta.\forall x \in (\delta).A \\ \leftrightarrow \xi' \models \forall \delta.\forall x \in (\delta).A \end{array} \right)$	
4	Hence assume: $\mathbb{I}, \xi^{\mathbb{I}}, \xi^{\mathbb{I}'}$ s.t. $\mathbb{I} \Vdash \forall \delta.\forall x \in (\delta).A \wedge \xi \preceq^* \xi'$	
5	$\rightarrow$ : Show that: $\xi \models \forall \delta.\forall x \in (\delta).A \rightarrow \xi' \models \forall \delta.\forall x \in (\delta).A$	line 7 below
6	$\leftarrow$ : Show that: $\xi' \models \forall \delta.\forall x \in (\delta).A \rightarrow \xi \models \forall \delta.\forall x \in (\delta).A$	line 13 below
7	$\rightarrow$ :	
7	Let: $\xi_{1d} \equiv \xi_1 \cdot \delta : \mathbb{I}_1 \setminus \text{-TCV}$ and $\xi'_{2d} \equiv \xi'_2 \cdot \delta : \mathbb{I}'_2 \setminus \text{-TCV}$	
8	$\xi \preceq^* \xi' \wedge \xi \models \forall \delta.\forall x \in (\delta).A$	Ignore Typing Constraints as these hold
9	$\leftrightarrow \xi \preceq^* \xi' \wedge \forall \xi_1^{\mathbb{I}_1}. \xi \preceq^* \xi_1 \rightarrow \forall M. M^{\left[\delta, \overset{\sim}{\rightsquigarrow} \xi_{1d}\right]} V \rightarrow \xi_{1d} \cdot x : V \models A$	Sem. $\forall \delta.\forall x \in (\delta).$
10	$\leftrightarrow \xi \preceq^* \xi' \wedge \forall \xi_1^{\mathbb{I}_1}. \xi \preceq^* \xi_1 \rightarrow \forall M. M^{\left[\delta, \overset{\sim}{\rightsquigarrow} \xi_{1d}\right]} V \rightarrow \xi_1 \cdot x : V \models A$	Lemma 30
11	$\rightarrow \xi \preceq^* \xi' \wedge \forall \xi_2^{\mathbb{I}'_2}. \xi' \preceq^* \xi_2 \rightarrow \forall M. M^{\left[\delta, \overset{\sim}{\rightsquigarrow} \xi'_{2d}\right]} V \rightarrow \xi_2 \cdot x : V \models A$	select $\xi'_2$ s.t. $\xi' \preceq^* \xi'_2$
12	$\leftrightarrow \xi' \models \forall \delta.\forall x \in (\delta).A$	Lemma 30, Sem. $\forall \delta.\forall x \in (\delta).$

←:

13	Let: $\xi'_{2d} \equiv \xi'_2 \cdot \delta : \mathbb{I}_2 \setminus \text{-TCV}$ and $\xi_{1d} \equiv \xi_1 \cdot \delta : \mathbb{I}_1 \setminus \text{-TCV}$ and $\xi'_{1d} \equiv \xi_1 \cdot \delta : \mathbb{I}'_1 \setminus \text{-TCV}$	
14	$\xi \preceq^* \xi' \wedge \xi' \models \forall \delta. \forall x \in (\delta). A$	Ignore Typing Constraints as these hold
15	$\leftrightarrow \xi \preceq^* \xi' \wedge \forall \xi'_2 \in \mathbb{I}'_2. \xi' \preceq^* \xi'_2 \rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi'_{2d} \cdot x : V \models A$	Sem. $\forall \delta. \forall x \in (\delta)$ .
16	$\leftrightarrow \xi \preceq^* \xi' \wedge \forall \xi'_2 \in \mathbb{I}'_2. \xi' \preceq^* \xi'_2 \rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi'_2 \cdot x : V \models A$	Lemma 30
17	$\leftrightarrow \xi \preceq^* \xi' \wedge \forall \xi_1 \in \mathbb{I}_1. \xi \preceq^* \xi_1$	Intro $\xi_1$
	$\rightarrow \forall \xi'_2 \in \mathbb{I}'_2. \xi' \preceq^* \xi'_2 \rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi'_2 \cdot x : V \models A$	
18	$\leftrightarrow \xi \preceq^* \xi \cdot \tilde{\xi}' \wedge \forall (\xi \cdot \tilde{\xi}_1)^{\mathbb{I}_1}. \xi \preceq^* \xi \cdot \tilde{\xi}_1$	write $\xi \equiv \xi \cdot \tilde{\xi}'$
	$\rightarrow \forall (\xi \cdot \tilde{\xi}')^{\mathbb{I}'_2}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}'_2$	write $\xi_1 \equiv \xi \cdot \tilde{\xi}_1$
	$\rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi'_{2d} \cdot x : V \models A$	write $\xi'_2 \equiv \xi' \cdot \tilde{\xi}'_2$
19	$\leftrightarrow \xi \preceq^* \xi \cdot \tilde{\xi}' \wedge \forall \tilde{\xi}_1. \xi \preceq^* (\xi \cdot \tilde{\xi}_1)^{\mathbb{I}_1} \wedge \xi \cdot \tilde{\xi}' \preceq^* (\xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1)^{\mathbb{I}'_1}$	Select $\tilde{\xi}'_2$ as $\tilde{\xi}_1$ $\mathbb{I}'_2 \equiv \mathbb{I}'_1$ FOL
	$\rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot x : V \models A$	
20	$\leftrightarrow \xi \preceq^* \xi \cdot \tilde{\xi}' \wedge \forall \tilde{\xi}_1. \xi \preceq^* (\xi \cdot \tilde{\xi}_1)^{\mathbb{I}_1} \wedge \xi \cdot \tilde{\xi}_1 \preceq^* (\xi \cdot \tilde{\xi}_1 \cdot \tilde{\xi}')^{\mathbb{I}'_1}$	Lemma 34
	$\rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi \cdot \tilde{\xi}_1 \cdot \tilde{\xi}' \cdot x : V \models A$	
21	$\rightarrow \xi \preceq^* \xi \cdot \tilde{\xi}' \wedge \forall \tilde{\xi}_1. \xi \preceq^* (\xi \cdot \tilde{\xi}_1)^{\mathbb{I}_1} \wedge \xi \cdot \tilde{\xi}_1 \preceq^* (\xi \cdot \tilde{\xi}_1 \cdot \tilde{\xi}')^{\mathbb{I}'_1}$	Subset of possible $M$ 's $\mathbb{I}'_1 \Vdash \mathbb{I}_1$
	$\rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi \cdot \tilde{\xi}_1 \cdot \tilde{\xi}' \cdot x : V \models A$	Lemma 11
22	$\rightarrow \forall \xi_1. \xi \preceq^* \xi_1^{\mathbb{I}_1} \wedge \xi_1 \preceq^* (\xi_1 \cdot \tilde{\xi}')^{\mathbb{I}'_1}$	Lemma 28 $A$ -Ext-Ind
	$\rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi_1 \cdot x : V \models A$	
23	$\rightarrow \forall \xi_1. \xi \preceq^* \xi_1^{\mathbb{I}_1} \rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi_1 \cdot x : V \models A$	$\hat{a}(\tilde{\xi}_1) \cap \hat{a}(\tilde{\xi}') \subseteq \hat{a}(\xi)$ guarantees constraint
24	$\leftrightarrow \forall \xi_1. \xi \preceq^* \xi_1^{\mathbb{I}_1} \rightarrow \forall M. M^{[\delta, \xi'_{2d}]} V \rightarrow \xi_{1d} \models A$	Lemma 30, $A^{-\delta}$
25	$\leftrightarrow \xi \models \forall \delta. \forall x \in (\delta). A$	Sem. $\forall \delta. \forall x \in (\mathbb{I} + \delta)$ .

**Lemma 36** (The formula in the postcondition of [GENSYM] is **Ext-Ind**).

$$\forall \delta. f \bullet () = b\{b\#\delta\} - \text{Ext-Ind}$$

*Proof:*

$$\forall \xi^{\mathbb{I}}, \xi'. \xi \preceq^* \xi' \rightarrow (\xi \models \forall \delta. f \bullet () = b\{b\#\delta\}) \leftrightarrow \xi' \models \forall \delta. f \bullet () = b\{b\#\delta\})$$

Extending  $(\xi \models \forall \delta. f \bullet () = b\{b\#\delta\} \rightarrow \xi' \models \forall \delta. f \bullet () = b\{b\#\delta\})$ :

1	<i>Assume: <math>\mathbb{I}</math>, s.t. <math>\mathbb{I} \Vdash \forall \delta. f \bullet () = b\{b\#\mathbb{I}\}</math></i>	
2	<i>Assume some <math>\xi^{\mathbb{I}^a}, \xi'^{\mathbb{I}'}</math> s.t. <math>\mathbb{I} \triangleright \xi \xi \preceq^* \xi'</math> and <math>\xi \models \forall \delta. f \bullet () = b\{b\#\delta\}</math></i>	
3	$\leftrightarrow \forall \xi_1^{\mathbb{I}_1}. \xi \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}_1 \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	Sem. $\forall \delta$ .
4	$\rightarrow \forall \xi_1^{\mathbb{I}_1}. \xi \preceq^* \xi' \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}_1 \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	Subset $\forall \xi_1$
5	$\rightarrow \forall \xi_1^{\mathbb{I}_1}. \xi' \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}_1 \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	Remove $\xi \preceq^*$
6	$\rightarrow \xi' \models \forall \delta. f \bullet () = b\{b\#\delta\}$	

Contracting  $(\xi \models \forall \delta. f \bullet () = b\{b\#\delta\} \leftarrow \xi' \models \forall \delta. f \bullet () = b\{b\#\delta\})$ :

7	<i>Assume: <math>\mathbb{I}</math>, s.t. <math>\mathbb{I} \Vdash \forall \delta. f \bullet () = b\{b\#\mathbb{I}\}</math></i>	
8	<i>Assume some <math>\xi^{\mathbb{I}^a}, \xi'</math> s.t. <math>\mathbb{I} \triangleright \xi</math> and <math>\xi \preceq^* \xi'</math> and <math>\xi' \models \forall \delta. f \bullet () = b\{b\#\delta\}</math></i>	
9	$\leftrightarrow \forall \xi_1^{\mathbb{I}_1}. \xi' \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}_1 \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	Sem. $\forall \delta$ .
10	$\leftrightarrow \forall \xi_1^{\tilde{\mathbb{I}}_1}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1$ $\rightarrow \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1) \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	$\xi'^{\mathbb{I}'} \equiv \xi^{\mathbb{I}'} \cdot \tilde{\xi}'^{\tilde{\mathbb{I}}'}$ $\xi_1^{\mathbb{I}_1} \equiv \xi^{\mathbb{I}'} \cdot \tilde{\xi}'^{\tilde{\mathbb{I}}'} \cdot \tilde{\xi}_1^{\tilde{\mathbb{I}}_1}$
11	$\rightarrow \forall \xi_1^{\tilde{\mathbb{I}}_1}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1$ $\rightarrow \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1) \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	Subset $\forall \tilde{\xi}_1$ only $\tilde{\xi}_1$ s.t. $\xi \preceq^* \xi \cdot \tilde{\xi}_1$
12	$\rightarrow \forall \xi_1^{\tilde{\mathbb{I}}_1}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1$ $\rightarrow \xi \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}_1) \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	See below: Line 16 (reducing $\delta$ holds)
13	$\rightarrow \forall \xi_1^{\tilde{\mathbb{I}}_1}. \xi \preceq^* \xi \cdot \tilde{\xi}_1 \rightarrow \xi \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}_1) \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	Lemma 34, $\mathfrak{a}(\tilde{\xi}_1) \cap \mathfrak{a}(\tilde{\xi}') \subseteq \mathfrak{a}(\xi)$ guarantees constraints
14	$\rightarrow \forall \xi_1^{\mathbb{I}_1}. \xi \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}_1 \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}$	$\mathbb{I}_1 \equiv \mathbb{I} + \tilde{\mathbb{I}}_1, \xi_1 \equiv \xi \cdot \tilde{\xi}_1$
15	$\rightarrow \xi \models \forall \delta. f \bullet () = b\{b\#\delta\}$	Sem. $\forall \delta$ .

*Proof of:*

$$\begin{aligned}
 & \forall \xi^{\mathbb{I}}, \tilde{\xi}_1^{\tilde{\mathbb{I}}_1}, \tilde{\xi}'^{\tilde{\mathbb{I}}'}, V_x. \\
 & \xi \preceq^* \xi \cdot \tilde{\xi}' \wedge \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1 \\
 & \wedge \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1) \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\} \\
 & \rightarrow \\
 & \xi \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}_1) \setminus \neg_{TCV} \models f \bullet () = b\{b\#\delta\}
 \end{aligned}$$

*Proof:*

16	$\xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \leftrightarrow \xi \cdot \tilde{\xi}_1 \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1$	Lemma 34
17	$\rightarrow \xi \cdot \tilde{\xi}_1 \cdot b : r_b \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot b : r_b$	Lemma 28
18	<i>Let</i> $\xi_3 \equiv \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1) \setminus \neg_{TCV}$ <i>and</i> $\xi_2 \equiv \xi \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}_1) \setminus \neg_{TCV}$	
19	<i>Assume:</i> $\xi_3 \models f \bullet () = b\{b\#\delta\}$	
20	$\leftrightarrow f() \overset{[\mathbb{I}_3, \xi_3]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\delta, \xi_3 \cdot b:r_b]}{\rightsquigarrow} r_b$	Sem. $\bullet = \{\}, \#$
21	$\rightarrow f() \overset{[\mathbb{I}_2, \xi_2]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\delta, \xi_3 \cdot b:r_b]}{\rightsquigarrow} r_b$	$f \in \llbracket \mathbb{I} \rrbracket_{\xi} \subseteq \llbracket \mathbb{I}_2 \rrbracket_{\xi_2} \subseteq \llbracket \mathbb{I}_3 \rrbracket_{\xi_3}$ , Lemmas 11
22	$\rightarrow f() \overset{[\mathbb{I}_2, \xi_2]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1, \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot b:r_b]}{\rightsquigarrow} r_b$	$M \overset{[\delta, \xi \cdot \delta : \mathbb{I} \setminus \neg_{TCV}]}{\rightsquigarrow} V \equiv M \overset{[\mathbb{I}, \xi]}{\rightsquigarrow} V$
23	$\rightarrow f() \overset{[\mathbb{I}_2, \xi_2]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\mathbb{I} + \tilde{\mathbb{I}}_1, \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot b:r_b]}{\rightsquigarrow} r_b$	Subset, $\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1 \Vdash \mathbb{I} + \tilde{\mathbb{I}}_1$
24	$\rightarrow f() \overset{[\mathbb{I}_2, \xi_2]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\mathbb{I} + \tilde{\mathbb{I}}_1, \xi \cdot \tilde{\xi}_1 \cdot b:r_b]}{\rightsquigarrow} r_b$	Lemma 11, line 17
25	$\rightarrow f() \overset{[\mathbb{I}_2, \xi_2]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\delta, \xi_2 \cdot b:r_b]}{\rightsquigarrow} r_b$	$M \overset{[\delta, \xi \cdot \delta : \mathbb{I} \setminus \neg_{TCV} \cdot b:r_b]}{\rightsquigarrow} V \equiv M \overset{[\mathbb{I}, \xi \cdot b:r_b]}{\rightsquigarrow} V$
26	$\rightarrow \xi_2 \models f \bullet () = b\{b\#\delta\}$	

**Lemma 37 (The formula used in the reasoning for  $\lambda x.\text{gensym}()$  is **Ext-Ind**).**

$$\forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\} - \text{Ext-Ind}$$

(Similar to proof above)

*Proof:*

$$\forall \xi^{\mathbb{I}}, \xi'. \xi \preceq^* \xi' \rightarrow \xi \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\} \leftrightarrow \xi' \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$$

*Extending* ( $\xi \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\} \rightarrow \xi' \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$ ):

1	Assume: $\mathbb{I}'$ s.t. $\mathbb{I}' \Vdash \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	
2	Assume some $\xi^{\mathbb{I}'_a}$ , $\xi'$ s.t. $\mathbb{I}' \triangleright \xi$ and $\xi \preceq^* \xi'$ and $\xi \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	
3	$\leftrightarrow \forall \xi_1^{\mathbb{I}'_1}. \xi \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}'_1 \setminus \neg_{TCV} \models \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	Sem. $\forall \delta$ .
4	$\rightarrow \forall \xi_1^{\mathbb{I}'_1}. \xi \preceq^* \xi' \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}'_1 \setminus \neg_{TCV} \models \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	Subset $\forall \xi_1$
5	$\rightarrow \forall \xi_1^{\mathbb{I}'_1}. \xi' \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}'_1 \setminus \neg_{TCV} \models \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	Remove $\xi \preceq^*$
6	$\rightarrow \xi' \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	

Contracting ( $\xi \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\} \leftarrow \xi' \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$ ):

7	Assume: $\mathbb{I}, \xi^{\mathbb{I}}$ s.t. $\mathbb{I} \Vdash \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	
8	Assume some $\xi'$ s.t. $\xi \preceq^* \xi'$ and $\xi' \models \forall \delta. \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	
9	Let: $\xi'_{1d} \equiv \xi_1^{\mathbb{I}'} \cdot \delta : \mathbb{I}'_1 \setminus \text{-TCV}$ ( $\equiv \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1) \setminus \text{-TCV}$ ) and $\xi'_{1d}$	
10	$\leftrightarrow \forall \xi_1^{\mathbb{I}'}. \xi' \preceq^* \xi_1 \rightarrow \forall M_x. M_x \xrightarrow{[\delta, \xi'_{1d}]} V_x \rightarrow \xi'_{1d} \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\}$	Sem. $\forall \delta, \forall x \in (\delta)$ .
11	$\leftrightarrow \forall \tilde{\xi}_1^{\tilde{\mathbb{I}}'}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1$ $\rightarrow \forall M_x. M_x \xrightarrow{[\delta, \xi'_{1d}]} V_x \rightarrow \xi'_{1d} \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\}$	$\xi^{\mathbb{I}'} \equiv \xi^{\mathbb{I}} \cdot \tilde{\xi}'^{\tilde{\mathbb{I}}'}$ $\xi_1^{\mathbb{I}'} \equiv \xi^{\mathbb{I}} \cdot \tilde{\xi}'^{\tilde{\mathbb{I}}'} \cdot \tilde{\xi}_1^{\tilde{\mathbb{I}}_1}$
12	$\rightarrow \forall \tilde{\xi}_1^{\tilde{\mathbb{I}}'}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1$ $\rightarrow \forall M_x. M_x \xrightarrow{[\delta, \xi'_{1d}]} V_x \rightarrow \xi'_{1d} \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\}$	Subset $\forall \tilde{\xi}_1$ only $\tilde{\xi}_1$ s.t. $\xi \preceq^* \xi \cdot \tilde{\xi}_1$
13	$\rightarrow \forall \tilde{\xi}_1^{\tilde{\mathbb{I}}'}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1$ $\rightarrow \forall M_x. M_x \xrightarrow{[\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1, \xi'_{1d} \setminus \delta]} V_x \rightarrow \xi'_{1d} \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\}$	$[\delta]_{\xi_{1d}} \equiv \mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1$ Lemma 10
14	Let: $\xi_2 \equiv \xi \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}_1) \setminus \text{-TCV}$	
15	$\rightarrow \forall \tilde{\xi}_1^{\tilde{\mathbb{I}}'}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1$ $(\rightarrow \forall M_x. M_x \xrightarrow{[\mathbb{I} + \tilde{\mathbb{I}}_1, \xi \cdot \tilde{\xi}_1]} V_x \rightarrow \xi'_{1d} \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\})$ $\rightarrow \forall M_x. M_x \xrightarrow{[\delta, \xi_2]} V_x \rightarrow \xi'_{1d} \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\}$	Subset $\forall M_x$ $\xi_2 \setminus \delta \subseteq \xi_{1d} \setminus \delta$
16	$\rightarrow \forall \tilde{\xi}_1^{\tilde{\mathbb{I}}'}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1$ $\rightarrow \forall M_x. M_x \xrightarrow{[\delta, \xi_2]} V_x \rightarrow \xi_2 \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\}$	See below, Line 19
17	$\rightarrow \forall \tilde{\xi}_1^{\tilde{\mathbb{I}}'}. \xi \preceq^* \xi \cdot \tilde{\xi}_1$ $\rightarrow \xi \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}_1) \setminus \text{-TCV} \models \forall x \in (\delta). f \bullet x = b\{b\#\delta+x\}$	Lemma 34, $\mathfrak{a}(\tilde{\xi}_1) \cap \mathfrak{a}(\tilde{\xi}') \subseteq \mathfrak{a}(\xi)$ guarantees constraints
18	$\rightarrow \forall \xi_1^{\mathbb{I}'}. \xi \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : \mathbb{I}'_1 \setminus \text{-TCV} \models f \bullet x = b\{b\#\delta+x\}$	$\xi_1^{\mathbb{I}'} \equiv \xi^{\mathbb{I}} \cdot \tilde{\xi}_1^{\tilde{\mathbb{I}}_1}$
19	$\rightarrow \xi \models \forall \delta. f \bullet x = b\{b\#\delta+x\}$	Sem. $\forall \delta$ .

Proof of

$$\begin{aligned}
 & \forall \tilde{\xi}_1^{\tilde{\mathbb{I}}'}. \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1 \\
 & \rightarrow \xi_{1d} \equiv \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1) \setminus \text{-TCV} \\
 & \wedge \xi_2 \equiv \xi \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I} + \tilde{\mathbb{I}}_1) \setminus \text{-TCV} \\
 & \wedge \forall M_x. M_x \xrightarrow{[\mathbb{I} + \tilde{\mathbb{I}}_1, \xi \cdot \tilde{\xi}_1]} V_x \\
 & \rightarrow \\
 & (\xi_{1d} \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\} \rightarrow \xi_2 \cdot x : V_x \models f \bullet x = b\{b\#\delta+x\})
 \end{aligned}$$

19	$\xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \leftrightarrow \xi \cdot \tilde{\xi}_1 \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1$	Lemma 34
20	$\rightarrow \xi \cdot \tilde{\xi}_1 \cdot b : r_b \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot b : r_b$	Lemma 28
21	Let $\xi_{3x}^{\mathbb{I}^3x} \equiv \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I}^+ + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1) \setminus \text{-TCV} \cdot x : V_x$	
22	Let $\xi_{2x}^{\mathbb{I}^2x} \equiv \xi^{\mathbb{I}^+} \cdot \tilde{\xi}_1 \cdot \delta : (\mathbb{I}^+ + \tilde{\mathbb{I}}_1) \setminus \text{-TCV} \cdot x : V_x$	
23	Assume: $\xi \preceq^* \xi \cdot \tilde{\xi}' \wedge \xi \cdot \tilde{\xi}' \preceq^* \xi \cdot \tilde{\xi}' \cdot \tilde{\xi}_1 \wedge \xi \preceq^* \xi \cdot \tilde{\xi}_1$	
24	Assume: $\xi_{3x} \models f \bullet x = b\{b\#\delta+x\}$	
25	$\leftrightarrow fx \overset{[\mathbb{I}^3x, \xi_{3x}]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\delta+x, \xi_{3x} \cdot b:r_b]}{\rightsquigarrow} r_b$	Sem. $\bullet = \{\}, \#$
26	$\leftrightarrow fx \overset{[\mathbb{I}^2x, \xi_{2x}]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\delta+x, \xi_{3x} \cdot b:r_b]}{\rightsquigarrow} r_b$	$f, x \in [\mathbb{I}^+]\xi \subseteq [\mathbb{I}^2x]_{\xi_{2x}} \subseteq [\mathbb{I}^3x]_{\xi_{3x}}$ , Lemma 11
27	$\leftrightarrow fx \overset{[\mathbb{I}^2x, \xi_{2x}]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\mathbb{I}^3x, (\xi_{3x} \setminus \delta) \cdot b:r_b]}{\rightsquigarrow} r_b$	$[\delta]_{\xi_{3x}} = \mathbb{I}^3x$ , Lemma 10
28	$\rightarrow fx \overset{[\mathbb{I}^2x, \xi_{2x}]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\mathbb{I}^2x, (\xi_{3x} \setminus \delta) \cdot b:r_b]}{\rightsquigarrow} r_b$	Subset, $\mathbb{I}^+ + \tilde{\mathbb{I}}' + \tilde{\mathbb{I}}_1 \Vdash \mathbb{I}^+ + \tilde{\mathbb{I}}_1$
29	$\rightarrow fx \overset{[\mathbb{I}^2x, \xi_{2x}]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\mathbb{I}^2x, (\xi_{2x} \setminus \delta) \cdot b:r_b]}{\rightsquigarrow} r_b$	Lemma 17, line 20
30	$\rightarrow fx \overset{[\mathbb{I}^2x, \xi_{2x}]}{\rightsquigarrow} r_b \wedge \neg \exists M_b.M_b \overset{[\delta+x, \xi_{2x} \cdot b:r_b]}{\rightsquigarrow} r_b$	$[\delta]_{\xi_{2x}} = \mathbb{I}^2x$ , Lemma 10
31	$\rightarrow \xi_{2x} \models f \bullet x = b\{b\#\delta+x\}$	

**Lemma 38 (SYN-EXT-IND implies Ext-Ind).**

$A$  is SYN-EXT-IND  $\rightarrow A$  is Ext-Ind

*Proof.* i.e.  $A - \text{SYN-EXT-IND}$

$$\rightarrow \forall \mathbb{I}^+, \xi^{\mathbb{I}^+}, \mathbb{I}', \xi'^{\mathbb{I}'} \cdot \mathbb{I}^+ \Vdash A \wedge \xi \preceq^* \xi' \rightarrow (\xi \models A \leftrightarrow \xi' \models A)$$

*Proof by induction on the structure of  $A$ , using Lemma 8 (i.e.  $\mathbb{I}_0 \subseteq \mathbb{I}^+ \rightarrow [\mathbb{I}_0]_{\xi} \equiv [\mathbb{I}_0]_{\xi'}$ ) and Lemma 24 (i.e.  $[e]_{\xi} \equiv [e]_{\xi'}$ )*

1. Base Formulas:

$A \equiv \top, \text{F}$  Clearly hold.

$A \equiv x\#\mathbb{I}_0$   $\mathbb{I}^+ \Vdash x\#\mathbb{I}_0$  implies  $\mathbb{I}^+ \Vdash x : \text{Nm}$  and  $\mathbb{I}^+ \Vdash \mathbb{I}_0$ .

Lemma 24 implies  $[x]_{\xi} \equiv [x]_{\xi'}$ .

Lemma 8 implies  $[\mathbb{I}_0]_{\xi} \equiv [\mathbb{I}_0]_{\xi'}$ .

Lemma 11 implies  $\exists M. M \overset{[\mathbb{I}_0, \xi]}{\rightsquigarrow} [x]_{\xi} \leftrightarrow \exists M. M \overset{[\mathbb{I}_0, \xi']}{\rightsquigarrow} [x]_{\xi'}$

2. Core Inductive Cases:

$A \equiv \neg C_1$  Holds from IH on  $C_1$  as  $\xi \models C_1 \leftrightarrow \xi' \models C_1$

hence  $\xi \not\models C_1 \leftrightarrow \xi' \not\models C_1$ , hence  $\xi \models \neg C_1 \leftrightarrow \xi' \models \neg C_1$ .

$A \equiv C_1 \wedge C_2$  Holds from IH on  $C_1$  and  $C_2$ , as  $\xi \models C_i \leftrightarrow \xi' \models C_i$  implies

$\xi \models C_1 \wedge C_2 \leftrightarrow \xi \models C_1 \wedge \xi \models C_2 \leftrightarrow \xi' \models C_1 \wedge \xi' \models C_2 \leftrightarrow \xi' \models C_1 \wedge C_2$ .



$$\begin{aligned}
 & A \equiv C_1 \vee C_2 \text{ Derivable given } A \vee B \equiv \neg(\neg A \wedge \neg B) \\
 & A \equiv C_1 \rightarrow C_2 \text{ Derivable given } A \rightarrow B \equiv \neg(A \wedge \neg B) \\
 & A \equiv u \bullet e = m\{C_1\} \text{ Lemma 11 implies } ue \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_m \leftrightarrow ue \stackrel{[\mathbb{I}, \xi']}{\rightsquigarrow} V_m. \\
 & \text{Lemma 28 implies } \xi \preceq^* \xi' \wedge ue \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_m \rightarrow \xi \cdot m : V_m \preceq^* \xi' \cdot m : V_m. \\
 & \text{Induction on } C_1 \text{ implies } \xi \cdot m : V_m \models C_1 \leftrightarrow \xi' \cdot m : V_m \models C_1. \\
 & \text{Hence } \xi \models u \bullet e = m\{C_1\} \\
 & \quad \leftrightarrow ue \stackrel{[\mathbb{I}, \xi]}{\rightsquigarrow} V_m \wedge \xi \cdot m : V_m \models C_1 \\
 & \quad \leftrightarrow ue \stackrel{[\mathbb{I}, \xi']}{\rightsquigarrow} V_m \wedge \xi' \cdot m : V_m \models C_1 \\
 & \quad \leftrightarrow \xi' \models u \bullet e = m\{C_1\} \\
 & A \equiv \forall x \in (\mathbb{I}_0).C_1 \text{ Lemma 8 implies } \llbracket \mathbb{I}_0 \rrbracket_\xi \equiv \llbracket \mathbb{I}_0 \rrbracket_{\xi'} \\
 & \text{Lemma 11 implies } \exists M. M \stackrel{[\mathbb{I}_0, \xi]}{\rightsquigarrow} \llbracket x \rrbracket_\xi \leftrightarrow \exists M. M \stackrel{[\mathbb{I}_0, \xi']}{\rightsquigarrow} \llbracket x \rrbracket_{\xi'} \\
 & \text{Lemma 28 implies } \xi \preceq^* \xi' \wedge M_x \stackrel{[\mathbb{I}_0, \xi]}{\rightsquigarrow} V_x \rightarrow \xi \cdot x : V_x \preceq^* \xi' \cdot x : V_x \\
 & \text{Induction on } C_1 \text{ implies } \xi \cdot x : V_x \models C_1 \leftrightarrow \xi' \cdot x : V_x \models C_1 \\
 & \text{hence } \xi \models \forall x \in (\mathbb{I}_0).C_1 \leftrightarrow \forall M.M \stackrel{[\mathbb{I}_0, \xi]}{\rightsquigarrow} W \rightarrow \xi \cdot x : W \models C_1 \quad \text{Sem. } \forall x \in (\mathbb{I}_0). \\
 & \quad \leftrightarrow \forall M.M \stackrel{[\mathbb{I}_0, \xi']}{\rightsquigarrow} W \rightarrow \xi \cdot x : W \models C_1 \quad \llbracket \mathbb{I}_0 \rrbracket_\xi \equiv \llbracket \mathbb{I}_0 \rrbracket_{\xi'} \\
 & \quad \quad \quad \text{Lemma 11} \\
 & \quad \leftrightarrow \forall M.M \stackrel{[\mathbb{I}_0, \xi']}{\rightsquigarrow} W \rightarrow \xi' \cdot x : W \models C_1 \quad \text{IH on } C_1 \\
 & \quad \leftrightarrow \xi' \models \forall x \in (\mathbb{I}_0).C_1 \quad \text{Sem. } \forall x \in (\mathbb{I}_0). \\
 & A \equiv \exists x \in (\mathbb{I}_0).C_1 \text{ Derivable from } \exists x \in (\mathbb{I}_0).C_1 \equiv \neg \forall x \in (\mathbb{I}_0).\neg C_1
 \end{aligned}$$

### 3. $\forall\delta$ -Inductive Cases:

$A \equiv \forall\delta.C_1$  holds by IH on  $C_1$ , Knowing that  $C_1$  is  $\delta$ -free then:

$$\begin{aligned}
 \xi \models \forall\delta.C_1 & \leftrightarrow \forall \xi_1^{I_1}. \xi \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : I_1 \models C_1 & \text{Sem. } \forall\delta. \\
 & \rightarrow \forall \xi_1^{I_1}. \xi \preceq^* \xi' \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : I_1 \models C_1 & \text{Subset } \forall \\
 & \rightarrow \forall \xi_1^{I_1}. \xi' \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : I_1 \models C_1 & \xi \preceq^* \xi' \text{ clearly holds} \\
 & \rightarrow \xi' \models \forall\delta.C_1 & \text{Sem. } \forall\delta. \\
 \\ 
 \xi' \models \forall\delta.C_1 & \leftrightarrow \forall \xi_1^{I_1}. \xi' \preceq^* \xi_1 \rightarrow \xi_1 \cdot \delta : I_1 \models C_1 & \text{Sem. } \forall\delta. \\
 & \rightarrow \forall \xi_1^{I_1}. \xi' \preceq^* \xi_1 \rightarrow \xi_1 \models C_1 & \text{Lemma 30 } C_1^{-\delta} \\
 & \rightarrow \xi' \models C_1 & \forall \xi_1 \rightarrow \xi' \\
 & \rightarrow \xi \models C_1 & \text{IH} \\
 & \rightarrow \forall \xi_2^{I_2}. \xi \preceq^* \xi_2 \rightarrow \xi_2 \models C_1 & \text{IH} \\
 & \rightarrow \forall \xi_2^{I_2}. \xi \preceq^* \xi_2 \rightarrow \xi_2 \cdot \delta : I_2 \models C_1 & \text{Lemma 30 } C_1^{-\delta} \\
 & \rightarrow \xi \models \forall\delta.C_1 & \text{Sem. } \forall\delta.
 \end{aligned}$$

$A \equiv \forall\delta.\forall x \in (\delta).C_1$  holds by IH on  $C_1^{-\delta}$ , Lemma 35

### 4. Two specific cases:

$\forall\delta.f \bullet () = b\{b\#\delta\}$  Lemma 36

$\forall\delta.\forall x \in (\delta).f \bullet x = b\{b\#\delta+x\}$  Lemma 37

## F Syntactic Thinness Construction Lemmas

**Definition 5.** *Syntactically define thin formulae as follows:*

1. If  $\mathbb{I} \Vdash A$  and  $\mathbb{I} \Vdash x : \alpha_b$  then  $A$  thin w.r.t  $x$   
Base types can always be removed if they do not occur in the assertion.
2. For  $A \equiv \top, \text{F}, e = e', e \neq e', y \# \mathbb{I}_1 \setminus \text{-TCV}$  and  $x \notin \text{fv}(A)$  then  $A$  thin w.r.t  $x$   
The base assertions are thin w.r.t  $x$  under basic assumptions.
3. If  $(\mathbb{I} + x + \mathbb{I}') \setminus x \Vdash y \# \mathbb{I}_0 \wedge \mathbb{I} \Vdash \mathbb{I}_0$  then  $y \# \mathbb{I}_0$  thin w.r.t  $x$   
If  $\mathbb{I} + x$  and  $\mathbb{I} \Vdash \mathbb{I}_0$  then  $x$  is not included in  $\mathbb{I}_0$  and cannot be included by any TCV occurring in  $\mathbb{I}_0$ .
4. If  $(\mathbb{I} + x : \text{Nm} + \mathbb{I}') \setminus x \Vdash y \# \mathbb{I}_0 + b \wedge \mathbb{I} \Vdash \mathbb{I}_0 \wedge \mathbb{I}' \Vdash b : \text{Nm}$  then  $y \# \mathbb{I}_0 + b$  thin w.r.t  $x$   
Similar to 3, but two possibilities:  $b = x \vee b \neq x$  both of which hold.
5. If  $A_1$  thin w.r.t  $x$  and  $A_2$  thin w.r.t  $x$  then  $A_1 \wedge A_2, A_1 \vee A_2, u \bullet e = m\{A_1\}, \forall y \in (\mathbb{I}_1).A_1, \exists y^{\alpha_b} \in (\mathbb{I}_1).A_1, \exists y \in (\mathbb{I}_1 \setminus \text{-TCV}).A_1$  are all thin w.r.t  $x$ .  
Constructing thin assertions with thin assertions holds in most cases.
6. If  $A$  thin w.r.t  $x$  and  $\delta \notin \text{ftcv}(A)$  then  $\forall \delta. A$  thin w.r.t  $x$   
If  $\delta$  is unused in  $A$  then thinness is maintained.
7. If  $A$  thin w.r.t  $x$  and  $\delta \notin \text{ftcv}(A)$  then  $\forall \delta. \forall y^{\alpha_b} \in (\mathbb{I} + \delta). A$  thin w.r.t  $x$   
If  $\delta$  is only used to derive  $y$ , then thinness is maintained as it reduces the potentially derived values.

We now define some lemmas that help prove that syntactic thinness implies thinness.

**Lemma 39 (Syntactically thin implies thin for Freshness).**

$$(\mathbb{I} + y + \mathbb{I}') \setminus y \Vdash x \# \mathbb{I}_0 \wedge \mathbb{I} \Vdash \mathbb{I}_0 \rightarrow x \# \mathbb{I}_0 \text{ thin w.r.t } y$$

*Proof:*

1	Assume: $(\mathbb{I}\Gamma + y + \mathbb{I}\Gamma') \setminus y \Vdash x \# \mathbb{I}\Gamma_0 \wedge \mathbb{I}\Gamma \Vdash \mathbb{I}\Gamma_0$	
2	$\rightarrow \mathbb{I}\Gamma + \mathbb{I}\Gamma' \Vdash x \# \mathbb{I}\Gamma_0 \wedge \mathbb{I}\Gamma \Vdash \mathbb{I}\Gamma_0$	
3	Assume: $\xi^{\mathbb{I}\Gamma + y + \mathbb{I}\Gamma'}$ such that $\xi \models x \# \mathbb{I}\Gamma_0$	
4	$\rightarrow \neg \exists M_x. M_x \overset{[\mathbb{I}\Gamma_0, \xi]}{\rightsquigarrow} \llbracket x \rrbracket_\xi$	
5	$\rightarrow \neg \exists M_x. M_x \overset{[\mathbb{I}\Gamma_0, \xi]}{\rightsquigarrow} \llbracket x \rrbracket_{\xi \setminus y}$	$\llbracket x \rrbracket_\xi \equiv \llbracket x \rrbracket_{\xi \setminus y}$
6	$\rightarrow \neg \exists M_x. \llbracket \mathbb{I}\Gamma_0 \rrbracket_\xi \vdash M_x : \text{Nm} \wedge \hat{a}(M_x) = \emptyset$	Sem. $\overset{[\cdot]}{\rightsquigarrow}$
7	$\rightarrow \neg \exists M_x. \llbracket \mathbb{I}\Gamma_0 \rrbracket_{\xi \setminus y} \vdash M_x : \text{Nm} \wedge \hat{a}(M_x) = \emptyset$	$\mathbb{I}\Gamma \Vdash \mathbb{I}\Gamma_0$
8	$\rightarrow \neg \exists M_x. \llbracket \mathbb{I}\Gamma_0 \rrbracket_{\xi \setminus y} \vdash M_x : \text{Nm} \wedge \hat{a}(M_x) = \emptyset$	$\mathbb{I}\Gamma \Vdash \mathbb{I}\Gamma_0 \wedge \llbracket \mathbb{I}\Gamma_0 \rrbracket_{\xi \setminus y} \vdash M_x : \text{Nm}$
9	$\rightarrow \neg \exists M_x. \llbracket \mathbb{I}\Gamma_0 \rrbracket_{\xi \setminus y} \vdash M_x : \text{Nm} \wedge \hat{a}(M_x) = \emptyset$	$\hat{a}(\xi) \equiv \hat{a}(\xi \setminus y) \cup \hat{a}(\xi(y))$
10	$\rightarrow \neg \exists M_x. \llbracket \mathbb{I}\Gamma_0 \rrbracket_{\xi \setminus y} \vdash M_x : \text{Nm} \wedge \hat{a}(M_x) = \emptyset$	$y \notin \text{fv}(M_x)$ , Lemma 1
11	$\rightarrow \xi \setminus y \models y \# \mathbb{I}\Gamma_0$	Sem. #

**Lemma 40 (Syntactically Thin formulae implies Thin).** *Define a method to syntactically define thin formulae as follows:*

1. If  $\mathbb{I}\Gamma \Vdash A$  and  $\mathbb{I}\Gamma \Vdash x : \alpha_b$  then  $A$  thin w.r.t  $x$
2. The assertions  $A \equiv \top, F, e = e', e \neq e', y \# \mathbb{I}\Gamma_1 \setminus \text{-TCV}$  and are all free from  $x$  ( $x \notin \text{fv}(A)$ ) then  $A$  thin w.r.t  $x$
3. If  $(\mathbb{I}\Gamma + y + \mathbb{I}\Gamma') \setminus y \Vdash x \# \mathbb{I}\Gamma_0 \wedge \mathbb{I}\Gamma \Vdash \mathbb{I}\Gamma_0$  then  $x \# \mathbb{I}\Gamma_0$  thin w.r.t  $y$
4. If  $(\mathbb{I}\Gamma + y : \text{Nm} + \mathbb{I}\Gamma') \setminus y \Vdash x \# \mathbb{I}\Gamma_0 + b \wedge \mathbb{I}\Gamma \Vdash \mathbb{I}\Gamma_0 \wedge \mathbb{I}\Gamma' \Vdash b : \text{Nm}$  then  $x \# \mathbb{I}\Gamma_0 + b$  thin w.r.t  $y$
5. If  $A_1$  thin w.r.t  $x$  and  $A_2$  thin w.r.t  $x$  then  $A_1 \wedge A_2, A_1 \vee A_2, u \bullet e = m\{A_1\}, \forall y \in (\mathbb{I}\Gamma_1).A_1, \exists y^{\alpha_b} \in (\mathbb{I}\Gamma_1).A_1, \exists y \in (\mathbb{I}\Gamma_1 \setminus \text{-TCV}).A_1$  are all thin w.r.t  $x$ .
6. If  $A_1$  thin w.r.t  $x$  and  $\delta \notin \text{ftcv}(A_1)$  then  $\forall \delta. A_1$  thin w.r.t  $x$
7. If  $A_1$  thin w.r.t  $x$  then  $\forall \delta. \forall y^{\alpha_b} \in (\mathbb{I}\Gamma + \delta). A_1^{-\delta}$  thin w.r.t  $x$

*Proof.* This proof assumes that  $\xi \setminus x$  (or  $\xi \setminus y$ ) is a well-constructed models.

1. If  $\mathbb{I}\Gamma \setminus x \Vdash A$  and  $\mathbb{I}\Gamma \Vdash x : \alpha_b$  then  $A$  thin w.r.t  $x$ :  
Given  $\llbracket e \rrbracket_\xi \equiv \llbracket e \rrbracket_{\xi \cdot x : V_x^{\alpha_b}}$  and  $\llbracket \mathbb{I}\Gamma_1 \rrbracket_\xi \equiv \llbracket \mathbb{I}\Gamma_1 \rrbracket_{\xi \cdot x : V_x^{\alpha_b}}$  the proof holds easily.  
Lemmas 5, 6, 7.
2. The assertions  $A \equiv \top, F, e = e', e \neq e', y \# \mathbb{I}\Gamma_1 \setminus \text{-TCV}$  and are all free from  $x$  ( $x \notin \text{fv}(A)$ ) then  $A$  thin w.r.t  $x$ :

$\top, F$  clearly hold.

$$e = e'$$

1	$\xi \models e = e'$	
2	$\leftrightarrow \llbracket e \rrbracket_\xi \cong_\alpha^{\hat{\alpha}(\xi)} \llbracket e' \rrbracket_\xi$	Sem. =
3	$\leftrightarrow \llbracket e \rrbracket_{\xi \setminus x} \cong_\alpha^{\hat{\alpha}(\xi)} \llbracket e' \rrbracket_{\xi \setminus x}$	$\mathbb{I} \setminus x \Vdash e : \alpha \rightarrow \llbracket e \rrbracket_\xi \equiv \llbracket e \rrbracket_{\xi \setminus x}$
4	$\leftrightarrow \llbracket e \rrbracket_{\xi \setminus x} \cong_\alpha^{\hat{\alpha}(\xi \setminus x)} \llbracket e' \rrbracket_{\xi \setminus x}$	Lemma 2
5	$\leftrightarrow \xi \setminus x \models e = e'$	

$e \neq e'$  See above.

$$y \# \mathbb{I}_1 \setminus \text{-TCV}$$

1	$\xi \models y \# \mathbb{I}_1 \setminus \text{-TCV}$	
2	$\leftrightarrow \neg \exists M_y. M_y \overset{[\mathbb{I}_1 \setminus \text{-TCV}, \xi]}{\rightsquigarrow} \llbracket y \rrbracket_\xi$	Sem. #
3	$\leftrightarrow \neg \exists M_y. M_y \overset{[\mathbb{I}_1 \setminus \text{-TCV}, \xi]}{\rightsquigarrow} \llbracket y \rrbracket_{\xi \setminus x}$	$\mathbb{I} \setminus x \Vdash y : \text{Nm} \rightarrow \llbracket y \rrbracket_\xi \equiv \llbracket y \rrbracket_{\xi \setminus x}$
4	$\leftrightarrow \neg \exists M_y. M_y \overset{[\mathbb{I}_1 \setminus \text{-TCV}, \xi]}{\rightsquigarrow} \llbracket y \rrbracket_{\xi \setminus x}$	$\llbracket \mathbb{I}_1 \rrbracket_\xi \equiv \llbracket \mathbb{I}_1 \rrbracket_{\xi \setminus x}$
5	$\rightarrow \xi \setminus x \models y \# \mathbb{I}_1 \setminus \text{-TCV}$	

3. If  $(\mathbb{I} + y + \mathbb{I}') \setminus y \Vdash x \# \mathbb{I}_0 \wedge \mathbb{I} \Vdash \mathbb{I}_0$  then  $x \# \mathbb{I}_0$  thin w.r.t  $y$ :  
Lemma 39

4. If  $(\mathbb{I} + y : \text{Nm} + \mathbb{I}') \setminus y \Vdash x \# \mathbb{I}_0 + b \wedge \mathbb{I} \Vdash \mathbb{I}_0 \wedge \mathbb{I}' \Vdash b : \text{Nm}$  then  
 $x \# \mathbb{I}_0 + b$  thin w.r.t  $y$   
Essentially relies on  $b : \text{Nm}$  must either be equal to  $y$  in which case no issue,  
or different in which case no issue.

1	$\xi^{\mathbb{I}\Gamma+y+\mathbb{I}\Gamma'} \models x\#\mathbb{I}\Gamma_0+b$	
2	$\leftrightarrow \neg\exists M_x.M_x \overset{[\mathbb{I}\Gamma_0+b, \xi]}{\rightsquigarrow} \llbracket x \rrbracket_\xi$	Sem. #
3	$\leftrightarrow \neg\exists M_x.M_x \overset{[\mathbb{I}\Gamma_0+b, \xi]}{\rightsquigarrow} \llbracket x \rrbracket_{\xi\setminus y} \quad \mathbb{I}\Gamma\setminus y \Vdash x : \text{Nm} \rightarrow \llbracket x \rrbracket_\xi \equiv \llbracket x \rrbracket_{\xi\setminus y}$	
4	$\leftrightarrow \neg\exists M_x. \llbracket \mathbb{I}\Gamma_0+b \rrbracket_\xi \vdash M_x : \text{Nm} \wedge \mathfrak{a}(M_x) = \emptyset$ $\wedge (\mathfrak{a}(\xi), M_x\xi) \Downarrow (\mathfrak{a}(\xi), G, \llbracket x \rrbracket_{\xi\setminus y})$	Sem. $\overset{[\cdot]}{\rightsquigarrow}$
5	$\rightarrow \neg\exists M_x. \llbracket \mathbb{I}\Gamma_0+b \rrbracket_{\xi\setminus y} \vdash M_x : \text{Nm} \wedge \mathfrak{a}(M_x) = \emptyset$ $\wedge (\mathfrak{a}(\xi), M_x(\xi\setminus y)) \Downarrow (\mathfrak{a}(\xi), G, \llbracket x \rrbracket_{\xi\setminus y}) \quad \llbracket \mathbb{I}\Gamma_0+b \rrbracket_{\xi\setminus y} \subseteq \llbracket \mathbb{I}\Gamma_0+b \rrbracket_\xi$	Subset $M_x$
6	<i>Case <math>\xi(y) \in \mathfrak{a}(\xi\setminus y)</math> implies <math>\mathfrak{a}(\xi) \equiv \mathfrak{a}(\xi\setminus y)</math></i>	
7	$\rightarrow \neg\exists M_x. \llbracket \mathbb{I}\Gamma_0+b \rrbracket_{\xi\setminus y} \vdash M_x : \text{Nm} \wedge \mathfrak{a}(M_x) = \emptyset$ $\wedge (\mathfrak{a}(\xi\setminus y), M_x(\xi\setminus y)) \Downarrow (\mathfrak{a}(\xi\setminus y), G, \llbracket x \rrbracket_{\xi\setminus y})$	$\mathfrak{a}(\xi) \equiv \mathfrak{a}(\xi\setminus y)$
8	<i>Case <math>\xi(y) \notin \mathfrak{a}(\xi\setminus y)</math> implies <math>\xi(y) \notin \mathfrak{a}(M_1(\xi\setminus y))</math></i>	
9	$\rightarrow \neg\exists M_x. \llbracket \mathbb{I}\Gamma_0+b \rrbracket_{\xi\setminus y} \vdash M_x : \text{Nm} \wedge \mathfrak{a}(M_x) = \emptyset$ $\wedge (\mathfrak{a}(\xi\setminus y), M_x(\xi\setminus y)) \Downarrow (\mathfrak{a}(\xi\setminus y), G, \llbracket x \rrbracket_{\xi\setminus y})$	Lemma 1
10	$\rightarrow \neg\exists M_x.M_x \overset{[\mathbb{I}\Gamma_0+b, \xi\setminus y]}{\rightsquigarrow} \llbracket x \rrbracket_{\xi\setminus y}$	Sem. $\overset{[\cdot]}{\rightsquigarrow}$
11	$\rightarrow \xi\setminus y \models x\#\mathbb{I}\Gamma_0+b$	

5. If  $A_1$  thin w.r.t  $x$  and  $A_2$  thin w.r.t  $x$  then  $A_1 \wedge A_2$ ,  $A_1 \vee A_2$ ,  $u \bullet e = m\{A_1\}$ ,  $\forall y \in (\mathbb{I}\Gamma_1).A_1$ ,  $\exists y^{\alpha_b} \in (\mathbb{I}\Gamma_1).A_1$ ,  $\exists y \in (\mathbb{I}\Gamma_1 \setminus \text{TCV}).A_1$  are all thin w.r.t  $x$ :

Elementary cases: obvious cases are  $A_1 \wedge A_2$ ,  $A_1 \vee A_2$  where the proofs are omitted.

$e \bullet e' = m\{A_1\}$ : holds by IH on  $A_1$  and

1	$\xi \models e \bullet e' = m\{A_1\}$	
2	$\leftrightarrow ee' \overset{[\mathbb{I}\Gamma, \xi]}{\rightsquigarrow} V_m \wedge \xi \cdot m : V_m \models A_1$	IH
3	$\rightarrow ee' \overset{[\mathbb{I}\Gamma, \xi\setminus x]}{\rightsquigarrow} V_m \wedge \xi\setminus x \cdot m : V_m \models A_1 \quad \llbracket ee' \rrbracket_\xi \equiv \llbracket ee' \rrbracket_{\xi\setminus x}, \text{ Lemma 1}$ $(\mathfrak{a}(V_m) \cap \mathfrak{a}(V_x)) \subseteq \mathfrak{a}(\xi\setminus x)$	
4	$\rightarrow \xi\setminus x \models e \bullet e' = m\{A_1\}$	

$\forall y \in (\mathbb{I}_1).A_1$ : holds assuming  $A_1$  thin w.r.t  $x$  then

1	$\xi \models \forall y \in (\mathbb{I}_1).A_1$		
2	$\leftrightarrow \forall M_y.M_y \overset{[\mathbb{I}_1, \xi]}{\rightsquigarrow} V_y \rightarrow \xi \cdot y : V_y \models A_1$		Sem. $\forall y \in (\mathbb{I}_1).A_1$
3	$\rightarrow \forall M_y. \hat{a}(M_y) = \emptyset \wedge [\mathbb{I}_1]_\xi \vdash M_y : \alpha \rightarrow \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi), M_y \xi) \Downarrow (\hat{a}(\xi), G', V_y)$		Sem. $\overset{[\cdot]}{\rightsquigarrow}$
4	$\rightarrow \forall M_y. \hat{a}(M_y) = \emptyset \wedge [\mathbb{I}_1]_{\xi \setminus x} \vdash M_y : \alpha \rightarrow \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi), M_y \xi) \Downarrow (\hat{a}(\xi), G', V_y)$		$[\mathbb{I}_1]_{\xi \setminus x} \subseteq [\mathbb{I}_1]_\xi$
5	$\rightarrow \forall M_y. \hat{a}(M_y) = \emptyset \wedge [\mathbb{I}_1]_{\xi \setminus x} \vdash M_y : \alpha \rightarrow \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi \setminus x), M_y \xi \setminus x) \Downarrow (\hat{a}(\xi \setminus x), G', V_y)$		$x \notin \text{fv}([\mathbb{I}_1]_{\xi \setminus x})$ , Lemma 1 $(\hat{a}(V_y) \cap \hat{a}(V_x) \subseteq \hat{a}(\xi \setminus x))$
6	$\rightarrow \forall M_y.M_y \overset{[\mathbb{I}_1, \xi \setminus x]}{\rightsquigarrow} V_y \rightarrow (\xi \cdot y : V_y) \models A_1$		Sem. $\overset{[\cdot]}{\rightsquigarrow}$
7	$\rightarrow \forall M_y.M_y \overset{[\mathbb{I}_1, \xi \setminus x]}{\rightsquigarrow} V_y \rightarrow (\xi \cdot y : V_y) \setminus x \models A_1$		IH
8	$\rightarrow \forall M_y.M_y \overset{[\mathbb{I}_1, \xi \setminus x]}{\rightsquigarrow} V_y \rightarrow \xi \setminus x \cdot y : V_y \models A_1$		Def $\xi \setminus x$
9	$\rightarrow \xi \setminus x \models \forall y \in (\mathbb{I}_1).A_1$		Sem. $\forall y \in (\mathbb{I}_1).A_1$

$\exists y^{\text{ob}} \in (\mathbb{I}_1).A_1$ : holds as follows:

1	$\xi \models \exists y^{\text{ob}} \in (\mathbb{I}_1).A_1$		
2	$\leftrightarrow \exists M_y.M_y \overset{[\mathbb{I}_1, \xi]}{\rightsquigarrow} V_y \wedge \xi \cdot y : V_y \models A_1$		Sem. $\exists y \in (\mathbb{I}_1).A_1$
3	$\rightarrow \exists M_y. \hat{a}(M_y) = \emptyset \wedge [\mathbb{I}_1]_\xi \vdash M_y : \alpha \wedge \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi), M_y \xi) \Downarrow (\hat{a}(\xi), G', V_y)$		Sem. $\overset{[\cdot]}{\rightsquigarrow}$
4	$\rightarrow \exists M_y. \hat{a}(M_y) = \emptyset \wedge [\mathbb{I}_1]_{\xi \setminus x} \vdash M_y : \alpha \wedge \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi), M_y \xi) \Downarrow (\hat{a}(\xi), G', V_y)$		$V_y^{\text{ob}}$ equally derivable from any TC, Lemma 4
5	$\rightarrow \exists M_y. \hat{a}(M_y) = \emptyset \wedge [\mathbb{I}_1]_{\xi \setminus x} \vdash M_y : \alpha \wedge \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi \setminus x), M_y \xi \setminus x) \Downarrow (\hat{a}(\xi \setminus x), G', V_y)$		$x \notin \text{fv}([\mathbb{I}_1]_{\xi \setminus x})$ , Lemma 1 $(\hat{a}(V_y) \cap \hat{a}(V_x) \subseteq \hat{a}(\xi \setminus x))$
6	$\rightarrow \exists M_y.M_y \overset{[\mathbb{I}_1, \xi \setminus x]}{\rightsquigarrow} V_y \wedge \xi \cdot y : V_y \models A_1$		Sem. $\overset{[\cdot]}{\rightsquigarrow}$
7	$\rightarrow \exists M_y.M_y \overset{[\mathbb{I}_1, \xi]}{\rightsquigarrow} V_y \wedge (\xi \cdot y : V_y) \setminus x \models A_1$		IH
8	$\rightarrow \exists M_y.M_y \overset{[\mathbb{I}_1, \xi]}{\rightsquigarrow} V_y \wedge \xi \setminus x \cdot y : V_y \models A_1$		Def $\xi \setminus x$
9	$\rightarrow \xi \setminus x \models \exists y \in (\mathbb{I}_1).A_1$		Sem. $\exists y \in (\mathbb{I}_1).A_1$

$\exists y \in (\mathbb{I}_1 \setminus \text{-TCV}). A_1$ : holds as follows:

1	$\xi \models \exists y \in (\mathbb{I}_1 \setminus \text{-TCV}). A_1$	
2	$\leftrightarrow \exists M_y. M_y \overset{[\mathbb{I}_1 \setminus \text{-TCV}, \xi]}{\rightsquigarrow} V_y \wedge \xi \cdot y : V_y \models A_1$	Sem. $\exists y \in (\mathbb{I}_1 \setminus \text{-TCV}). A_1$
3	$\rightarrow \exists M_y. \hat{a}(M_y) = \emptyset \wedge \llbracket \mathbb{I}_1 \setminus \text{-TCV} \rrbracket_\xi \vdash M_y : \alpha \wedge \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi), M_y \xi) \Downarrow (\hat{a}(\xi), G', V_y)$	Sem. $\llbracket \cdot \rrbracket$
4	$\rightarrow \exists M_y. \hat{a}(M_y) = \emptyset \wedge \llbracket \mathbb{I}_1 \setminus \text{-TCV} \rrbracket_{\xi \setminus x} \vdash M_y : \alpha \wedge \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi), M_y \xi) \Downarrow (\hat{a}(\xi), G', V_y)$	$\llbracket \mathbb{I}_0 \rrbracket_{\xi \setminus x} \subseteq \llbracket \mathbb{I}_0 \rrbracket_\xi$
5	$\rightarrow \exists M_y. \hat{a}(M_y) = \emptyset \wedge \llbracket \mathbb{I}_1 \setminus \text{-TCV} \rrbracket_{\xi \setminus x} \vdash M_y : \alpha \wedge \xi \cdot y : V_y \models A_1$ $\wedge (\hat{a}(\xi \setminus x), M_y \xi \setminus x) \Downarrow (\hat{a}(\xi \setminus x), G', V_y)$	$x \notin \text{fv}(\llbracket \mathbb{I}_1 \rrbracket_{\xi \setminus x}),$ Lemma 1 $(\hat{a}(V_y) \cap \hat{a}(V_x) \subseteq \hat{a}(\xi \setminus x))$
6	$\rightarrow \exists M_y. M_y \overset{[\mathbb{I}_1 \setminus \text{-TCV}, \xi]}{\rightsquigarrow} V_y \wedge (\xi \cdot y : V_y) \setminus x \models A_1$	IH
7	$\rightarrow \exists M_y. M_y \overset{[\mathbb{I}_1 \setminus \text{-TCV}, \xi]}{\rightsquigarrow} V_y \wedge \xi \setminus x \cdot y : V_y \models A_1$	Def $\xi \setminus x$
8	$\rightarrow \exists M_y. M_y \overset{[\mathbb{I}_1 \setminus \text{-TCV}, \xi \setminus x]}{\rightsquigarrow} V_y \wedge (\xi \cdot y : V_y) \setminus x \models A_1$	Sem. $\llbracket \cdot \rrbracket$
9	$\rightarrow \xi \setminus x \models \exists y \in (\mathbb{I}_1 \setminus \text{-TCV}). A_1$	Sem. $\exists y \in (\mathbb{I}_1 \setminus \text{-TCV}). A_1$

6 If  $A_1$  thin w.r.t  $x$  and  $\delta \notin \text{ftcv}(A_1)$  then  $\forall \delta. A_1$  thin w.r.t  $x$ :  
Holds by IH on  $A_1$  as follows

1	$\xi \models \forall \delta. A_1$	
2	$\leftrightarrow \forall \xi' \mathbb{I}' . \xi \preceq^* \xi' \rightarrow \xi' \cdot \delta : \mathbb{I}' \models A_1$	Sem. $\forall \delta$ .
3	$\leftrightarrow \forall \xi' \mathbb{I}' . \xi \preceq^* \xi' \rightarrow \xi' \models A_1^{-\delta}$	Lemma 30
4	$\leftrightarrow \forall \xi' \mathbb{I}' . \xi \preceq^* \xi' \rightarrow \xi' \setminus x \models A_1^{-\delta}$	IH
5	$\rightarrow \forall \xi' \setminus x \mathbb{I}' \setminus x . \xi \setminus x \preceq^* \xi' \setminus x \rightarrow \xi' \setminus x \models A_1^{-\delta}$	Subset of $\forall \xi' \mathbb{I}'$ as $\llbracket \mathbb{I}' \setminus x \rrbracket_{\xi \setminus x} \subseteq \llbracket \mathbb{I}' \rrbracket_\xi$ and $\xi \setminus x \subseteq \xi$
6	$\rightarrow \forall \xi'' \mathbb{I}'' . \xi \setminus x \preceq^* \xi'' \rightarrow \xi'' \models A_1^{-\delta}$	Rename $\xi'' \equiv \xi' \setminus x$
7	$\rightarrow \forall \xi'' \mathbb{I}'' . \xi \setminus x \preceq^* \xi'' \rightarrow \xi'' \cdot \delta : \mathbb{I}'' \models A_1^{-\delta}$	Lemma 30
8	$\rightarrow \xi \setminus x \models \forall \delta. A_1^{-\delta}$	Sem. $\forall \delta$ .

7 If  $A_1$  thin w.r.t  $x$  then  $\forall\delta.\forall y^{\alpha_y} \in (\mathbb{I}\Gamma+\delta).A_1^{-\delta}$  thin w.r.t  $x$   
Holds by IH on  $A_1$  as follows:

1	Assume: $A$ thin w.r.t $x$	
2	i.e. assume $\forall\mathbb{I}\Gamma, \delta, y^{\alpha_y}. \mathbb{I}\Gamma+\delta+y : \alpha_y \backslash x \Vdash A \rightarrow \forall\xi_{dy}^{\mathbb{I}\Gamma+\delta+y:\alpha_y}. \xi_{dy} \Vdash A \rightarrow \xi_{dy} \backslash x \Vdash A$	
3	Assume: $\mathbb{I}\Gamma, \xi^{\mathbb{I}\Gamma}$ s.t. $\mathbb{I}\Gamma \backslash x \Vdash \forall\delta.\forall y \in (\mathbb{I}\Gamma+\delta).A$ and $\xi \Vdash \forall\delta.\forall y \in (\mathbb{I}\Gamma+\delta).A$	
4	$\rightarrow \mathbb{I}\Gamma \backslash x + \delta + y : \alpha_y \Vdash A$ $\wedge \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow \forall M_y^{\alpha_y}. M_y \xrightarrow{[\mathbb{I}\Gamma+\delta, \xi_d^{\delta:\mathbb{I}\Gamma_d}]} V_y \rightarrow \xi_d \cdot \delta : \mathbb{I}\Gamma_d \cdot y : V_y \Vdash A$	Typing rules, Sem. $\forall, \forall \in ()$ .
5	$\rightarrow \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow \forall M_y^{\alpha_y}. M_y \xrightarrow{[\mathbb{I}\Gamma+\delta, \xi_d^{\delta:\mathbb{I}\Gamma_d}]} V_y \rightarrow \xi_d \cdot y : V_y \Vdash A$	$A^{-\delta}$ , Lemma 30
6	$\rightarrow \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow \forall M_y^{\alpha_y}. M_y \xrightarrow{[\mathbb{I}\Gamma+\delta, \xi_d^{\delta:\mathbb{I}\Gamma_d}]} V_y \rightarrow (\xi_d \cdot y : V_y) \backslash x \Vdash A$	Assumption, 2
7	$\rightarrow \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow \forall M_y^{\alpha_y}. M_y \xrightarrow{[\mathbb{I}\Gamma+\delta, \xi_d^{\delta:\mathbb{I}\Gamma_d}]} V_y \rightarrow (\xi_d \cdot \delta : \mathbb{I}\Gamma_d \cdot y : V_y) \backslash x \Vdash A$	$A^{-\delta}$ , Lemma 30 (Ignoring other TCV's)
8	$\rightarrow \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow \forall M_y^{\alpha_y}. \llbracket \mathbb{I}\Gamma+\delta \rrbracket_{\xi_d \cdot \delta : \mathbb{I}\Gamma_d} \vdash M_y : \alpha_y \wedge \hat{\mathfrak{a}}(M_y) = \emptyset$ $\wedge (\hat{\mathfrak{a}}(\xi_d \cdot \delta : \mathbb{I}\Gamma_d), M_y(\xi_d \cdot \delta : \mathbb{I}\Gamma_d)) \Downarrow (\hat{\mathfrak{a}}(\xi_d \cdot \delta : \mathbb{I}\Gamma_d), G', V_y)$ $\rightarrow (\xi_d \cdot \delta : \mathbb{I}\Gamma_d \cdot y : V_y) \backslash x \Vdash A$	Sem. $\xrightarrow{[\cdot]}$
9	$\rightarrow \wedge \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow \forall M_y^{\alpha_y}. \xi_{dd} \equiv \xi_d \cdot \delta : \mathbb{I}\Gamma_d$ $\wedge \llbracket (\mathbb{I}\Gamma+\delta) \backslash x \rrbracket_{\xi_{dd} \backslash x} \vdash M_y : \alpha_y \wedge \hat{\mathfrak{a}}(M_y) = \emptyset$ $\wedge (\hat{\mathfrak{a}}(\xi_{dd}), M_y \xi_{dd}) \Downarrow (\hat{\mathfrak{a}}(\xi_{dd}), G', V_y)$ $\rightarrow (\xi_{dd} \cdot y : V_y) \backslash x \Vdash A$	Subset of Type Context $\llbracket \mathbb{I}\Gamma+\delta \rrbracket_{\xi_{dd} \backslash x} \subseteq \llbracket \mathbb{I}\Gamma+\delta \rrbracket_{\xi_{dd}}$ $\llbracket \mathbb{I}\Gamma_0 \rrbracket_{\xi_0} \backslash x \equiv \llbracket \mathbb{I}\Gamma_0 \backslash x \rrbracket_{\xi_0}$
10	$\rightarrow \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow \forall M_y^{\alpha_y}. \xi_{dd-x} \equiv (\xi_d \cdot \delta : \mathbb{I}\Gamma_d) \backslash x$ $\wedge \llbracket (\mathbb{I}\Gamma+\delta) \backslash x \rrbracket_{\xi_{dd-x}} \vdash M_y : \alpha_y \wedge \hat{\mathfrak{a}}(M_y) = \emptyset$ $\wedge (\hat{\mathfrak{a}}(\xi_d \cdot \delta : \mathbb{I}\Gamma_d), M_y \xi_{dd-x}) \Downarrow (\hat{\mathfrak{a}}(\xi_d \cdot \delta : \mathbb{I}\Gamma_d), G', V_y)$ $\rightarrow \xi_{dd-x} \cdot y : V_y \Vdash A$	$x \notin \text{fv}(M) \rightarrow M\xi \equiv M(\xi \backslash x)$
11	$\rightarrow \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow \forall M_y^{\alpha_y}. \xi_{dd-x} \equiv (\xi_d \cdot \delta : \mathbb{I}\Gamma_d) \backslash x$ $\wedge \llbracket (\mathbb{I}\Gamma+\delta) \backslash x \rrbracket_{\xi_{dd-x}} \vdash M_y : \alpha_y \wedge \hat{\mathfrak{a}}(M_y) = \emptyset$ $\wedge (\hat{\mathfrak{a}}(\xi_{dd-x}), M_y \xi_{dd-x}) \Downarrow (\hat{\mathfrak{a}}(\xi_{dd-x}), G', V_y)$ $\rightarrow \xi_{dd-x} \cdot y : V_y \Vdash A$	Lemma 1
12	$\rightarrow \forall\xi_d^{\mathbb{I}\Gamma_d}. \xi \preceq^* \xi_d \rightarrow (\xi_d \cdot \delta : \mathbb{I}\Gamma_d) \backslash x \Vdash \forall y \in (\mathbb{I}\Gamma+\delta).A$	Sem. $\forall \in ()$ .
13	$\rightarrow \forall\xi_{d-x}^{\mathbb{I}\Gamma_{d-x}}. \xi \preceq^* \xi_{d-x} \cdot x : V_x \rightarrow (\xi_{d-x} \cdot x : V_x \cdot \delta : \mathbb{I}\Gamma_d) \backslash x \Vdash \forall y \in (\mathbb{I}\Gamma+\delta).A$	$\xi_d \equiv \xi_{d-x} \cdot x : V_x$
14	$\rightarrow \forall\xi_{d-x}^{\mathbb{I}\Gamma_{d-x}}. \xi \backslash x \preceq^* \xi_{d-x} \rightarrow \xi_{d-x} \cdot \delta : \mathbb{I}\Gamma_{d-x} \Vdash \forall y \in (\mathbb{I}\Gamma+\delta).A$	Subset of $\forall\xi_{d-x}$
15	$\rightarrow \xi \backslash x \Vdash \forall\delta.\forall y \in (\mathbb{I}\Gamma+\delta).A$	Sem. $\forall\delta$ .
16	Hence: $\forall\mathbb{I}\Gamma. \mathbb{I}\Gamma \backslash x \Vdash \forall\delta.\forall y \in (\mathbb{I}\Gamma+\delta).A$ $\rightarrow \forall\xi^{\mathbb{I}\Gamma}. \xi \Vdash \forall\delta.\forall y \in (\mathbb{I}\Gamma+\delta).A \rightarrow \xi \backslash x \Vdash \forall\delta.\forall y \in (\mathbb{I}\Gamma+\delta).A$	lines 3-15
17	Hence: $\forall\delta.\forall y \in (\mathbb{I}\Gamma+\delta).A$ thin w.r.t $x$	



## G Conservativity

We sketch the proof that our logic is an extension of the STLC logic as defined in [9], and do not explain any of the trivial details. We define the logic introduced in this paper as the  $\nu$ -logic for brevity. We ignore the other possible extensions of  $\text{Int}$  and operations on  $\text{Int}$ ,  $\text{Bool}$ ,...

**The language for STLC.**

$$\begin{aligned}
\alpha &::= \text{Unit} \mid \text{Bool} \mid \alpha \rightarrow \alpha \mid \alpha \times \alpha \\
\Gamma &::= \emptyset \mid \Gamma, x : \alpha \\
V &::= x \mid c \mid \lambda x.M \mid \langle V, V \rangle \\
M &::= V \mid MM \mid \text{let } x = M \text{ in } M \mid M = M \\
&\quad \mid \text{if } M \text{ then } M \text{ else } M \mid \langle M, M \rangle \mid \pi_i(M)
\end{aligned}$$

**The logic for STLC.**

$$\begin{aligned}
\alpha &::= \text{Unit} \mid \text{Bool} \mid \alpha \rightarrow \alpha \mid \alpha \times \alpha \\
e &::= c \mid x \mid \pi_i(e) \mid \langle e, e \rangle \\
A &::= e = e \mid \neg A \mid A \wedge A \mid e \bullet e = x^\alpha \{A\} \mid \forall x^\alpha.A
\end{aligned}$$

**Axioms for STLC.** The axioms are:

- STLC terms can only be reasoned about through types which are  $\text{Nm}$ -free, for which we introduce two axioms:

$$\begin{aligned}
(\text{ext}) \quad \text{Ext}(e_1, e_2) &\leftrightarrow e_1 = e_2 \\
(u5) \quad \forall x^\alpha \in (\emptyset).A &\leftrightarrow \forall x^\alpha \in (\mathbb{I}).A \quad \alpha \text{ is Nm-free}
\end{aligned}$$

Axiom  $(\text{ext})$  is similar to that of [9], but extended to all  $\text{Nm}$ -free types terms. Axiom  $(u5)$  ensures names in a  $\text{Nm}$ -free terms can be swapped for any fresh name.

- The axioms for predicate logic, which are equivalent in the  $\nu$ -logic.
- The axioms for evaluation formulae ( $\bullet = \{\}$ ) which are all lifted directly from the STLC-logic [2, 9] but with the added assumption that all terms terminate and new definition of  $\forall x^\alpha \in (\mathbb{I}).A$ . which behaves identically to  $\forall x$ . when  $\alpha$  is  $\text{Nm}$ -free.
- The axioms for  $\forall x.A$  which are those of FOL, are represented by their translation to the  $\nu$ -logic and axioms for  $\forall x \in (\mathbb{I}).A$ .
- $(\text{ext})$  this axiom holds in the STLC-logic because functions must always produce the same result if applied twice. This still holds in  $\nu$ -logic as seen in Sec. 5, but only for  $\text{Nm}$ -free types. A simple example of why this fails for other types is  $\text{gensym} : \text{Unit} \rightarrow \text{Nm}$  which clearly produces different names each time it is applied.

**Rules for STLC.** Can be seen in Fig. 5.

$$\begin{array}{c}
\frac{}{\{A[x/m]\} x :_m \{A\}} \text{[VAR]} \quad \frac{}{\{A[c/m]\} c :_m \{A\}} \text{[CONST]} \quad \frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{C[m = n/u]\}}{\{A\} M = N :_u \{C\}} \text{[Eq]} \\
\frac{\{A^{\neg x} \wedge B\} M :_m \{C\}}{\{A\} \lambda x^\alpha . M :_u \{\forall x^\alpha . (B \rightarrow u \bullet x = m\{C\})\}} \text{[LAM]} \quad \frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{m \bullet n = u\{C\}\}}{\{A\} MN :_u \{C\}} \text{[APP]} \\
\frac{\{A\} M :_m \{B\} \quad \{B[b_i/m]\} N_i :_u \{C\} \quad b_1 = \text{true} \quad b_2 = \text{false} \quad i = 1, 2}{\{A\} \text{if } M \text{ then } N_1 \text{ else } N_2 :_u \{C\}} \text{[IF]} \\
\frac{\{A\} M :_m \{B\} \quad \{B\} N :_n \{C[(m, n)/u]\}}{\{A\} \langle M, N \rangle :_u \{C\}} \text{[PAIR]} \quad \frac{\{A\} M :_m \{B[\pi_i(m)/u]\}}{\{A\} \pi_i(M) :_u \{B\}} \text{[PROJ}(i)\text{]}
\end{array}$$

Fig. 5. Rules for the STLC

**Definition 6.** *The logic for the  $\nu$ -calculus is a conservative extension of the logic for the simply typed  $\lambda$ -calculus, provided:*

- If  $A$  is a formula in the  $\lambda$ -logic derivable from the axioms for the STLC above, then the translation of  $A$  to the  $\nu$ -logic is also derivable from the axioms in the logic of the  $\nu$ -calculus.
- If  $\{A\} M :_m \{B\}$  is a triple in the logic of the STLC, derivable from the rules for the STLC then  $\{A\} M :_m \{B\}$  is also derivable from the rules in the logic of the  $\nu$ -calculus.

**Definition 7.** *Define a translation of STLC assertions (and triples) into  $\nu$ -calculus assertions (and triples) as follows:*

$$\begin{aligned}
\langle\langle e = e' \rangle\rangle_{\lambda \rightarrow \nu} &= e = e' \\
\langle\langle \neg A \rangle\rangle_{\lambda \rightarrow \nu} &= \neg \langle\langle A \rangle\rangle_{\lambda \rightarrow \nu} \\
\langle\langle A \wedge B \rangle\rangle_{\lambda \rightarrow \nu} &= \langle\langle A \rangle\rangle_{\lambda \rightarrow \nu} \wedge \langle\langle B \rangle\rangle_{\lambda \rightarrow \nu} \\
\langle\langle e \bullet e' = m\{A\} \rangle\rangle_{\lambda \rightarrow \nu} &= e \bullet e' = m\{\langle\langle A \rangle\rangle_{\lambda \rightarrow \nu}\} \\
\langle\langle \forall x^\alpha . A \rangle\rangle_{\lambda \rightarrow \nu} &= \forall x^\alpha \in (\emptyset). \langle\langle A \rangle\rangle_{\lambda \rightarrow \nu}
\end{aligned} \tag{5}$$

$$\langle\langle \{A\} M :_m \{B\} \rangle\rangle_{\lambda \rightarrow \nu} = \{\langle\langle A \rangle\rangle_{\lambda \rightarrow \nu}\} M :_m \{\langle\langle B \rangle\rangle_{\lambda \rightarrow \nu}\}$$

**Theorem 3.** *The logic for the  $\nu$ -calculus is a conservative extension of the logic for the simply typed  $\lambda$ -calculus above.*

*Proof.* Most **axioms** are simple extensions of the STLC logic, and thus no proof is required. Note that for all  $A_\lambda. \langle\langle A \rangle\rangle_{\lambda \rightarrow \nu}$  – **Ext-Ind** as  $\langle\langle A \rangle\rangle_{\lambda \rightarrow \nu}$  is TCV free. Axioms that should hold easily are: (eq1), (u3), (u4), (e1 – 3), as these are copied from the STLC logic. For (u1) an adaption (u5) for all types which are name free is required to instantiate to variables derived from  $\emptyset$ .

See Sec C.5 for soundness of (ext) for the limited case of name free types in the LTC and formulae.

All **rules** for  $\nu$ -calculus are an extension of the rules for the STLC, with a few exceptions:

- Substitution: given all LTCs in the logic are reduced to  $\emptyset$  (u5) then substitution in all rules becomes standard as seen in the STLC as we never need to perform  $\mathit{I}[e/x]$ .
- [LAM] introduces  $\forall\delta$ . however this can just be instantiated. It also introduces  $\forall x \in (\delta)$ . which can be reduced to  $\forall x \in (\emptyset)$ . via (u2).
- **thin w.r.t  $\tilde{x}$** : Given names are not present then thinness simply implies non-existence of such a variable, i.e.  $x \notin \text{fv}(A)$  or  $A^{-x}$
- **Ext-Ind** holds for all assertions that are free from  $\delta$  hence holds for all STLC logical assertions.