

On Graded 2-Absorbing Coprimary Submodules

Malik BATAINEH and Rashid ABU-DAWWAS

ABSTRACT. The aim of this article is to introduce the concept of graded 2-absorbing coprimary submodules as a generalization of graded strongly 2-absorbing second submodules, and explore some properties of this class. A non-zero graded R -submodule N of a graded R -module M is called a graded 2-absorbing coprimary R -submodule if whenever x, y are homogeneous elements of R and K is a graded R -submodule of M such that $xyN \subseteq K$, then either x or y is in the graded radical of $(K :_R N)$ or $xy \in \text{Ann}_R(N)$. Several results have been achieved.

1. Introduction

Throughout this article, G will be a group with identity e and R a commutative ring with nonzero unity 1. Then R is said to be G -graded if $R = \bigoplus_{g \in G} R_g$ with $R_g R_h \subseteq R_{gh}$ for all $g, h \in G$ where R_g is an additive subgroup of R for all $g \in G$. The elements of R_g are called homogeneous of degree g . If $x \in R$, then x can be written uniquely as $\sum_{g \in G} x_g$, where x_g is the component of x in R_g . It is known that R_e is a subring of R and $1 \in R_e$. The set of all homogeneous elements of R is $h(R) = \bigcup_{g \in G} R_g$. Let I be an ideal of a graded ring R . Then I is said to be graded ideal if $I = \bigoplus_{g \in G} (I \cap R_g)$, i.e., for $x \in I$, $x = \sum_{g \in G} x_g$ where $x_g \in I$ for all $g \in G$. An ideal of a graded ring need not be graded. Assume that M is a left unital R -module. Then M is said to be G -graded if $M = \bigoplus_{g \in G} M_g$ with $R_g M_h \subseteq M_{gh}$ for all $g, h \in G$ where M_g is an additive subgroup of M for all $g \in G$. The elements of M_g are called homogeneous of degree g . It is clear that M_g is an R_e -submodule of M for all $g \in G$. If $x \in M$, then x can be written uniquely as $\sum_{g \in G} x_g$, where x_g is the component of x in M_g . The set of all homogeneous elements of M is $h(M) = \bigcup_{g \in G} M_g$. Let N be an R -submodule of a graded R -module M . Then N is said to be graded R -submodule if $N = \bigoplus_{g \in G} (N \cap M_g)$, i.e., for $x \in N$, $x = \sum_{g \in G} x_g$ where $x_g \in N$ for all $g \in G$.

2010 *Mathematics Subject Classification.* Primary 16W50; Secondary 13A02.

Key words and phrases. Graded strongly 2-absorbing second submodules; graded 2-absorbing primary submodules; graded strongly 2-absorbing secondary submodules.

An R -submodule of a graded R -module need not be graded. For more details and terminology, see [9, 10].

LEMMA 1.1. ([8], Lemma 2.1) *Let R be a G -graded ring and M be a G -graded R -module.*

- (1) *If I and J are graded ideals of R , then $I + J$ and $I \cap J$ are graded ideals of R .*
- (2) *If N and K are graded R -submodules of M , then $N + K$ and $N \cap K$ are graded R -submodules of M .*
- (3) *If N is a graded R -submodule of M , $r \in h(R)$, $x \in h(M)$ and I is a graded ideal of R , then Rx , IN and rN are graded R -submodules of M . Moreover, $(N :_R M) = \{r \in R : rM \subseteq N\}$ is a graded ideal of R .*

In particular, $Ann_R(M) = (0_M :_R M)$ is a graded ideal of R . Let M be a graded R -module, N be a graded R -submodule of M and $x \in h(R)$. Then $(N :_M x) = \{m \in M : xm \in N\}$ is a graded R -submodule of M , see ([11], Lemma 2.9). Let P be a proper graded ideal of R . Then the graded radical of P is $Grad(P)$, and is defined to be the set of all $r \in R$ such that for each $g \in G$, there exists a positive integer n_g for which $r_g^{n_g} \in P$. One can see that if $r \in h(R)$, then $r \in Grad(P)$ if and only if $r^n \in P$ for some positive integer n . In fact, $Grad(P)$ is a graded ideal of R , see [12].

In [2], graded 2-absorbing ideals as a generalization of graded prime ideals have been defined. A proper graded ideal P of R is said to be graded 2-absorbing if whenever $a, b, c \in h(R)$ such that $abc \in P$, then either $ab \in P$ or $ac \in P$ or $bc \in P$. Al-Zoubi and Sharafat in [3], introduced the concept of graded 2-absorbing primary ideals. This concept generalizes both graded primary ideals and graded 2-absorbing ideals. A proper graded ideal P of R is called a graded 2-absorbing primary ideal of R if whenever $a, b, c \in h(R)$ with $abc \in P$, then $ab \in P$ or $ac \in Grad(P)$ or $bc \in Grad(P)$. Graded second submodules of graded modules over commutative rings were introduced by Ansari-Toroghy and Farshadifar in [5] as follows: A non-zero graded R -submodule N of a graded R -module M is said to be a graded second R -submodule if for all $a \in h(R)$, either $aN = \{0\}$ or $aN = N$. In this case, $P = Ann_R(N)$ is a graded prime ideal of R , and N is called a graded P -second R -submodule of M . In [11], a generalization of graded second submodules has been given as follows: A non-zero graded R -submodule N of a graded R -module M is said to be a graded strongly 2-absorbing second R -submodule of M if whenever $x, y \in h(R)$ and K is a graded R -submodule of M such that $xyN \subseteq K$, then $xN \subseteq K$ or $yN \subseteq K$ or $xy \in Ann_R(N)$.

In this article, we act in accordance with [7] to introduce the concept of graded 2-absorbing coprimary R -submodules of a graded R -module M as a generalization of graded strongly 2-absorbing second submodules, and explore some properties of this class. A non-zero graded R -submodule N of a graded R -module M is called a graded 2-absorbing coprimary R -submodule if whenever $x, y \in h(R)$ and K is a graded R -submodule of M such that $xyN \subseteq K$, then $x \in Grad((K :_R N))$ or $y \in Grad((K :_R N))$ or $xy \in Ann_R(N)$.

2. Graded 2-Absorbing Coprimary Submodules

In this section, we introduce and study the concept of graded 2-absorbing coprimary submodules.

DEFINITION 2.1. *Let M be a graded R -module and N be a non-zero graded R -submodule of M . Then N is said to be a graded 2-absorbing coprimary R -submodule of M if whenever $x, y \in h(R)$ and K is a graded R -submodule of M such that $xyN \subseteq K$, then $x \in \text{Grad}((K :_R N))$ or $y \in \text{Grad}((K :_R N))$ or $xy \in \text{Ann}_R(N)$.*

Clearly, every graded strongly 2-absorbing second submodule is graded 2-absorbing coprimary. The next example shows that the converse is not true in general.

EXAMPLE 2.2. *Consider the \mathbb{Z} -module $M = \mathbb{Z}_4 \oplus \mathbb{Z}_9 \oplus \mathbb{Z}_5$. Then by ([7], Example 2.2), $N = \mathbb{Z}_4 \oplus \mathbb{Z}_9 \oplus \{0\}$ is a 2-absorbing coprimary submodule of M . So, if we consider the trivial graduation for M , N will be a graded 2-absorbing coprimary submodule of M . On the other hand, N is not graded strongly 2-absorbing second submodule of M by ([11], Corollary 3.4).*

PROPOSITION 2.3. *Let M be a graded R -module and N be a graded 2-absorbing coprimary R -submodule of M . If K is a graded R -submodule of M such that $N \not\subseteq K$, then $(K :_R N)$ is a graded 2-absorbing primary ideal of R .*

PROOF. By Lemma 1.1, $(K :_R N)$ is a graded ideal of R . Let $x, y, z \in h(R)$ such that $xyz \in (K :_R N)$. Then $xyzN = z(xyN) \subseteq K$, and then $xyN \subseteq (K :_M z)$. Since N is a graded 2-absorbing coprimary R -submodule of M , $x \in \text{Grad}((K :_M z) :_R N)$ or $y \in \text{Grad}((K :_M z) :_R N)$ or $xy \in \text{Ann}_R(N)$. If $xy \in \text{Ann}_R(N)$, then $xyN = \{0\} \subseteq K$, and then $xy \in (K :_R N)$. If $x \in \text{Grad}((K :_M z) :_R N)$, then $x^n zN \subseteq K$ for some positive integer n , and then $(xz)^n N \subseteq K$, which implies that $xz \in \text{Grad}((K :_R N))$. Similarly, if $y \in \text{Grad}((K :_M z) :_R N)$, then $yz \in \text{Grad}((K :_R N))$. Hence, $(K :_R N)$ is a graded 2-absorbing primary ideal of R . \square

As a direct consequence of Proposition 2.3, we have the following corollary.

COROLLARY 2.4. *Let M be a graded R -module and N be a graded 2-absorbing coprimary R -submodule of M . Then $\text{Ann}_R(N)$ is a graded 2-absorbing primary ideal of R .*

COROLLARY 2.5. *Let M be a graded R -module and N be a graded 2-absorbing coprimary R -submodule of M . Then $\text{Grad}(\text{Ann}_R(N))$ is a graded 2-absorbing ideal of R .*

PROOF. The result follows by Corollary 2.4 and ([3], Theorem 2.3). \square

PROPOSITION 2.6. *Let M be a graded R -module and N be a graded 2-absorbing coprimary R -submodule of M . Then aN is a graded 2-absorbing coprimary R -submodule of M for all $a \in h(R) - \text{Ann}_R(N)$.*

PROOF. Let $a \in h(R) - \text{Ann}_R(N)$. Then by Lemma 1.1, aN is a graded R -submodule of M . Suppose that $x, y \in h(R)$ and K is a graded R -submodule of M such that $xy(aN) \subseteq K$. Then $xyN \subseteq (K :_M a)$. Since N is a graded 2-absorbing coprimary R -submodule of M , $x^n N \subseteq (K :_M a)$ for some positive integer n or $y^m N \subseteq (K :_M a)$ for some positive integer m or $xy \in \text{Ann}_R(N)$, and then $x^n(aN) \subseteq K$ or $y^m(aN) \subseteq K$ or $xy \in \text{Ann}_R(N) \subseteq \text{Ann}_R(aN)$, as needed. \square

Let M and S be two G -graded R -modules. An R -homomorphism $f : M \rightarrow S$ is said to be a graded R -homomorphism if $f(M_g) \subseteq S_g$ for all $g \in G$.

LEMMA 2.7. ([1], Lemma 2.16) *Suppose that $f : M \rightarrow S$ is a graded R -homomorphism of graded R -modules. If K is a graded R -submodule of S , then $f^{-1}(K)$ is a graded R -submodule of M .*

LEMMA 2.8. ([6], Lemma 4.8) Suppose that $f : M \rightarrow S$ is a graded R -homomorphism of graded R -modules. If L is a graded R -submodule of M , then $f(L)$ is a graded R -submodule of $f(M)$.

PROPOSITION 2.9. Suppose that $f : M \rightarrow S$ is a graded R -homomorphism of graded R -modules.

- (1) If N is a graded 2-absorbing coprimary R -submodule of M such that $N \not\subseteq \text{Ker}(f)$, then $f(N)$ is a graded 2-absorbing coprimary R -submodule of S .
- (2) If K is a graded 2-absorbing coprimary R -submodule of S and $K \subseteq f(M)$, then $f^{-1}(K)$ is a graded 2-absorbing coprimary R -submodule of M .

PROOF. (1) By Lemma 2.8, $f(N)$ is a graded R -submodule of S . Let $x, y \in h(R)$ and K be a graded R -submodule of S such that $xyf(N) \subseteq K$. Then $xyN \subseteq f^{-1}(K)$. Since $f^{-1}(K)$ is a graded R -submodule of M by Lemma 2.7 and N is a graded 2-absorbing coprimary R -submodule of M , we have $x^nN \subseteq f^{-1}(K)$ for some positive integer n or $y^mN \subseteq f^{-1}(K)$ for some positive integer m or $xy \in \text{Ann}_R(N) \subseteq \text{Ann}_R(f(N))$, and then $x^n f(N) \subseteq K$ or $y^m f(N) \subseteq K$ or $xy \in \text{Ann}_R(f(N))$, as required.

- (2) By Lemma 2.7, $f^{-1}(K)$ is a graded R -submodule of M . If $f^{-1}(K) = \{0\}$, then since $K \subseteq f(M)$, we have $f(f^{-1}(K)) = K = \{0\}$, which is a contradiction. So, $f^{-1}(K) \neq \{0\}$. Let $x, y \in h(R)$ and L be a graded R -submodule of M such that $xyf^{-1}(K) \subseteq L$. Since $K \subseteq f(M)$, $xyK \subseteq f(L)$. Since $f(L)$ is a graded R -submodule of S by Lemma 2.8 and K is a graded 2-absorbing coprimary R -submodule of S , $x^nK \subseteq f(L)$ for some positive integer n or $y^mK \subseteq f(L)$ for some positive integer m or $xy \in \text{Ann}_R(K)$, and then $x^n f^{-1}(K) = f^{-1}(x^n K) \subseteq L$ or $y^m f^{-1}(K) = f^{-1}(y^m K) \subseteq L$ or $xy \in \text{Ann}_R(K) = \text{Ann}_R(f^{-1}(K))$, as needed. □

LEMMA 2.10. Let M be a graded R -module and N be a non-zero graded R -submodule of M . Then N is a graded 2-absorbing coprimary R -submodule of M if and only if for any $x, y \in h(R)$, either $x^nN \subseteq xyN$ for some positive integer n or $y^mN \subseteq xyN$ for some positive integer m or $xy \in \text{Ann}_R(N)$.

PROOF. Suppose that N is a graded 2-absorbing coprimary R -submodule of M . Let $x, y \in h(R)$. Then $xy \in h(R)$, and then by Lemma 1.1, xyN is a graded R -submodule of M such that $xyN \subseteq xyN$. Since N is a graded 2-absorbing coprimary R -submodule of M , $x^nN \subseteq xyN$ for some positive integer n or $y^mN \subseteq xyN$ for some positive integer m or $xy \in \text{Ann}_R(N)$. The converse is clear. □

Let M be a graded R -module and S be a multiplicative subset of $h(R)$. Then $S^{-1}M$ is a graded $S^{-1}R$ -module with

$$(S^{-1}M)_g = \left\{ \frac{m}{s}, m \in M_h, s \in S \cap R_{hg^{-1}} \right\},$$

$$(S^{-1}R)_g = \left\{ \frac{a}{s}, a \in R_h, s \in S \cap R_{hg^{-1}} \right\}.$$

PROPOSITION 2.11. Let M be a graded R -module, S be a multiplicative subset of $h(R)$ and N be a graded R -submodule of M such that $S^{-1}N \neq \{0\}$. If N is a graded 2-absorbing coprimary R -submodule of M , then $S^{-1}N$ is a graded 2-absorbing coprimary $S^{-1}R$ -submodule of $S^{-1}M$.

PROOF. Let $x, y \in h(R)$ and $s, t \in S$. Since N is a graded 2-absorbing copri-
 mary R -submodule of M , by Lemma 2.10, $x^n N \subseteq xyN$ for some positive integer n
 or $y^m N \subseteq xyN$ for some positive integer m or $xy \in \text{Ann}_R(N)$. If $xy \in \text{Ann}_R(N)$,
 then $\frac{x}{s}\frac{y}{t}(S^{-1}N) = \{0\}$. If $x^n N \subseteq xyN$, then $(\frac{x}{s})^n(S^{-1}N) = (\frac{x^n}{s^n})(S^{-1}N) =$
 $\frac{1}{s^n}S^{-1}(x^n N) \subseteq \frac{1}{s^n}S^{-1}(xyN) = S^{-1}(xyN) = (\frac{1}{s}\frac{1}{t})(S^{-1}(xyN)) = (\frac{x}{s}\frac{y}{t})(S^{-1}N)$. Sim-
 ilarly, if $y^m N \subseteq xyN$, then $(\frac{x}{s})^m(S^{-1}N) \subseteq (\frac{x}{s}\frac{y}{t})(S^{-1}N)$. Hence, by Lemma 2.10,
 $S^{-1}N$ is a graded 2-absorbing copri- $S^{-1}R$ -submodule of M . \square

DEFINITION 2.12. Let M be a G -graded R -module, $g \in G$ and N be a non-zero
 graded R -submodule of M . Then N is said to be a g -2-absorbing copri- R -
 submodule of M if whenever $x, y \in R_g$ and K is a graded R -submodule of M such that
 $xyN \subseteq K$, then $x \in \text{Grad}((K :_R N))$ or $y \in \text{Grad}((K :_R N))$ or $xy \in \text{Ann}_R(N)$.

LEMMA 2.13. Let M be a G -graded R -module, $g \in G$, I be a graded ideal of
 R and N be a g -2-absorbing copri- R -submodule of M . If $x \in R_g$ and K is
 a graded R -submodule of M such that $IxN \subseteq K$, then $x \in \text{Grad}((K :_R N))$ or
 $I_g \subseteq \text{Grad}((K :_R N))$ or $I_g x \subseteq \text{Ann}_R(N)$.

PROOF. Suppose that $x \notin \text{Grad}((K :_R N))$ and $I_g x \not\subseteq \text{Ann}_R(N)$. Then there
 exists $y \in I_g$ such that $xy \notin \text{Ann}_R(N)$, and then $xyN \subseteq K$. Since N is g -2-
 absorbing copri- R -submodule, $y \in \text{Grad}((K :_R N))$. Let $z \in I_g$. Then $(y+z)xN \subseteq K$, and
 then $y+z \in \text{Grad}((K :_R N))$ or $(y+z)x \in \text{Ann}_R(N)$. If $y+z \in \text{Grad}((K :_R N))$,
 then $z \in \text{Grad}((K :_R N))$. If $(y+z)x \in \text{Ann}_R(N)$, then $xz \notin \text{Ann}_R(N)$. So,
 $zxN \subseteq K$ implies that $z \in \text{Grad}((K :_R N))$. Hence, $I_g \subseteq \text{Grad}((K :_R N))$. \square

THEOREM 2.14. Let M be a G -graded R -module, $g \in G$, I, J be two graded
 ideals of R and N be a g -2-absorbing copri- R -submodule of M . If K is a
 graded R -submodule of M such that $IJN \subseteq K$, then $I_g \subseteq \text{Grad}((K :_R N))$ or
 $J_g \subseteq \text{Grad}((K :_R N))$ or $I_g J_g \subseteq \text{Ann}_R(N)$.

PROOF. Suppose that $I_g \not\subseteq \text{Grad}((K :_R N))$ and $J_g \not\subseteq \text{Grad}((K :_R N))$. Then
 there exist $x \in I_g - \text{Grad}((K :_R N))$ and $y \in J_g - \text{Grad}((K :_R N))$. Let $z \in$
 I_g and $w \in J_g$. Now, $xJN \subseteq K$, so by Lemma 2.13, $xJ_g \subseteq \text{Ann}_R(N)$, and so
 $(I_g - \text{Grad}((K :_R N)))J_g \subseteq \text{Ann}_R(N)$. Similarly, $I_g y \subseteq \text{Ann}_R(N)$, and so $I_g(J_g -$
 $\text{Grad}((K :_R N))) \subseteq \text{Ann}_R(N)$. So, we conclude that $xy, xw, zy \in \text{Ann}_R(N)$. Now,
 $(x+z)(y+w)N \subseteq K$. Therefore, $x+z \in \text{Grad}((K :_R N))$ or $y+w \in \text{Grad}((K :_R N))$
 or $(x+z)(y+w) \in \text{Ann}_R(N)$. If $x+z \in \text{Grad}((K :_R N))$, then $z \notin \text{Grad}((K :_R N))$.
 Thus $z \in I_g - \text{Grad}((K :_R N))$, which implies that $zw \in \text{Ann}_R(N)$. Similarly, if
 $y+w \in \text{Grad}((K :_R N))$, then $zw \in \text{Ann}_R(N)$. If $(x+z)(y+w) = xy+xw+yz+zw \in$
 $\text{Ann}_R(N)$, then $zw \in \text{Ann}_R(N)$. Hence, $I_g J_g \subseteq \text{Ann}_R(N)$. \square

Graded comultiplication modules have been introduced by Toroghy and Far-
 shadifar in [4]; a graded R -module M is said to be graded comultiplication if for
 every graded R -submodule N of M , $N = (0 :_M I)$ for some graded ideal I of R , or
 equivalently, $N = (0 :_M \text{Ann}_R(N))$.

PROPOSITION 2.15. Let M be a graded comultiplication R -module. If N is
 a graded 2-absorbing copri- R -submodule of M such that $\text{Grad}(\text{Ann}_R(N)) =$
 $\text{Ann}_R(N)$, then N is a graded strongly 2-absorbing second R -submodule of M . In
 particular, N is a graded 2-absorbing second R -submodule of M .

PROOF. The result follows by Corollary 2.5 and ([11], Proposition 3.7). \square

Let R_1 and R_2 be two G -graded rings, M_1 be a G -graded R_1 -module, M_2 be a G -graded R_2 -module, $R = R_1 \times R_2$ and $M = M_1 \times M_2$. Then M is a G -graded R -module by $M_g = (M_1)_g \times (M_2)_g$ for all $g \in G$.

THEOREM 2.16. *Let R_1 and R_2 be two G -graded rings, M_1 be a G -graded R_1 -module, M_2 be a G -graded R_2 -module, $R = R_1 \times R_2$ and $M = M_1 \times M_2$.*

- (1) *If $N = N_1 \times N_2$ is a graded 2-absorbing coprimary R -submodule of M , then $\text{Ann}_{R_1}(N_1)$ is a graded primary ideal of R_1 and $\text{Ann}_{R_2}(N_2)$ is a graded primary ideal of R_2 .*
- (2) *If N_1 is a graded R_1 -submodule of M_1 , N_2 is a graded R_2 -submodule of M_2 such that $\text{Ann}_{R_1}(N_1)$ is a graded primary ideal of R_1 , $\text{Ann}_{R_2}(N_2)$ is a graded primary ideal of R_2 and $N = N_1 \times N_2$, then $\text{Ann}_R(N)$ is a graded 2-absorbing primary ideal of R .*
- (3) *If N_1 is a graded 2-absorbing coprimary R_1 -submodule of M_1 and $N = N_1 \times \{0_{M_2}\}$, then $\text{Ann}_R(N)$ is a graded 2-absorbing primary ideal of R .*
- (4) *If N_2 is a graded 2-absorbing coprimary R_2 -submodule of M_2 and $N = \{0_{M_1}\} \times N_2$, then $\text{Ann}_R(N)$ is a graded 2-absorbing primary ideal of R .*

PROOF. (1) By ([13], Lemma 3.12), N_1 is a graded R_1 -submodule of M_1 and N_2 is a graded R_2 -submodule of M_2 , and then $\text{Ann}_{R_1}(N_1)$ is a graded ideal of R_1 and $\text{Ann}_{R_2}(N_2)$ is a graded ideal of R_2 . Since N is graded 2-absorbing coprimary, by Corollary 2.4, $\text{Ann}_R(N) = \text{Ann}_{R_1}(N_1) \times \text{Ann}_{R_2}(N_2)$ is a graded 2-absorbing primary ideal of R , and then by ([3], Theorem 2.10), $\text{Ann}_{R_1}(N_1)$ is a graded primary ideal of R_1 and $\text{Ann}_{R_2}(N_2)$ is a graded primary ideal of R_2 .

- (2) By ([13], Lemma 3.12), N is a graded R -submodule of M , and then $\text{Ann}_R(N)$ is a graded ideal of R . Since $\text{Ann}_{R_1}(N_1)$ is a graded primary ideal of R_1 , $\text{Ann}_{R_1}(N_1) \times R_2$ is a graded primary ideal of R . Similarly, $R_1 \times \text{Ann}_{R_2}(N_2)$ is a graded primary ideal of R . So, by ([3], Lemma 2.9), $\text{Ann}_{R_1}(N_1) \cap \text{Ann}_{R_2}(N_2) = \text{Ann}_{R_1}(N_1) \times \text{Ann}_{R_2}(N_2) = \text{Ann}_R(N)$ is a graded 2-absorbing primary ideal of R .
- (3) By ([13], Lemma 3.12), N is a graded R -submodule of M , and then $\text{Ann}_R(N)$ is a graded ideal of R . Since N_1 is graded 2-absorbing coprimary, by Corollary 2.4, $\text{Ann}_{R_1}(N_1)$ is a graded 2-absorbing primary ideal of R_1 , and then $\text{Ann}_{R_1}(N_1) \times R_2 = \text{Ann}_R(N)$ is a graded 2-absorbing primary ideal of R .
- (4) This result holds similar to part (3). □

References

- [1] R. Abu-Dawwas, M. Bataineh and H. Shashan, Graded generalized 2-absorbing submodules, *Beiträge zur Algebra und Geometrie / Contributions to Algebra and Geometry*, (2020), <https://doi.org/10.1007/s13366-020-00544-1>.
- [2] K. Al-Zoubi, R. Abu-Dawwas and S. Çeken, On graded 2-absorbing and graded weakly 2-absorbing ideals, *Hacettepe Journal of Mathematics and Statistics*, 48 (3) (2019), 724-731.
- [3] K. Al-Zoubi and N. Sharafat, On graded 2-absorbing primary and graded weakly 2-absorbing primary ideals, *Journal of the Korean Mathematical Society*, 54 (2) (2017), 675-684.
- [4] H. Ansari-Toroghy and F. Farshadifar, Graded comultiplication modules, *Chiang Mai Journal of Science*, 38 (3) (2011), 311-320.
- [5] H. Ansari-Toroghy and F. Farshadifar, On graded second modules, *Tamkang Journal of Mathematics*, 43 (4) (2012), 499-505.
- [6] S. E. Atani and F. E. K. Saraei, Graded modules which satisfy the gr-radical formula, *Thai Journal of Mathematics*, 8 (1) (2010), 161-170.

- [7] S. Çeken, 2-Absorbing coprimary submodules, Beiträge zur Algebra und Geometrie / Contributions to Algebra and Geometry, (2021), <https://doi.org/10.1007/s13366-020-00554-z>.
- [8] F. Farzalipour, P. Ghiasvand, On the union of graded prime submodules, Thai Journal of Mathematics, 9 (1) (2011), 49-55.
- [9] R. Hazrat, Graded rings and graded Grothendieck groups, Cambridge University press, 2016.
- [10] C. Nastasescu and F. Oystaeyen, Methods of graded rings, Lecture Notes in Mathematics, 1836, Springer-Verlag, Berlin, 2004.
- [11] M. Refai and R. Abu-Dawwas, On generalizations of graded second submodules, Proyecciones Journal of Mathematics, 39 (6) (2020), 1547-1564.
- [12] M. Refai, M. Hailat and S. Obiedat, Graded radicals and graded prime spectra, Far East Journal of Mathematical Sciences, (2000), 59-73.
- [13] H. Saber, T. Alraqad and R. Abu-Dawwas, On graded s -prime submodules, AIMS Mathematics, 6 (3) (2020), 2510-2524.

DEPARTMENT OF MATHEMATICS AND STATISTICS, JORDAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, IRBID, JORDAN

Email address: msbataineh@just.edu.jo

DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN

Email address: rrashid@yu.edu.jo