

Hadamard Extensions and the Identification of Mixtures of Product Distributions

Spencer L. Gordon* Leonard J. Schulman†

February 16, 2021

Abstract

The Hadamard Extension of a matrix is the matrix consisting of all Hadamard products of subsets of its rows. This construction arises in the context of identifying a mixture of product distributions on binary random variables: full column rank of such extensions is a necessary ingredient of identification algorithms. We provide several results concerning when a Hadamard Extension has full column rank.

1 Introduction

The Hadamard product for row vectors $u = (u_1, \dots, u_k)$, $v = (v_1, \dots, v_k)$ is the mapping $\odot : \mathbb{R}^k \times \mathbb{R}^k \rightarrow \mathbb{R}^k$ given by

$$u \odot v := (u_1 v_1, \dots, u_k v_k)$$

The identity for this product is the all-ones vector $\mathbb{1}$. We associate with vector v the linear operator $v_\odot = \text{diag}(v)$, a $k \times k$ diagonal matrix, so that

$$u \cdot v_\odot = v \odot u.$$

Throughout this paper \mathbf{m} is a real matrix with row set $[n] := \{1, \dots, n\}$ and column set $[k]$; write \mathbf{m}_i for a row and \mathbf{m}^j for a column.

As a matter of notation, for a matrix Q and nonempty sets R of rows and C of columns, let $Q|_R^C$ be the restriction of Q to those columns and rows (with either index omitted if all rows or columns are retained).

Definition 1. The Hadamard Extension of \mathbf{m} , written $\mathbb{H}(\mathbf{m})$, is the $2^n \times k$ matrix with rows \mathbf{m}_S for all $S \subseteq [n]$, where, for $S = \{i_1, \dots, i_\ell\}$, $\mathbf{m}_S = \mathbf{m}_{i_1} \odot \dots \odot \mathbf{m}_{i_\ell}$; equivalently $\mathbf{m}_S^j = \prod_{i \in S} \mathbf{m}_i^j$. (In particular $\mathbf{m}_\emptyset = \mathbb{1}$.)

This construction has arisen recently in learning theory [3, 8] where it is essential to source identification for a mixture of product distributions on binary random variables. We explain the connection further in Section 5. Motivated by this application, we are interested in the following two questions:

- (1) If $\mathbb{H}(\mathbf{m})$ has full column rank, must there exist a subset R of the rows, of bounded size, such that $\mathbb{H}(\mathbf{m}|_R)$ has full column rank?
- (2) In each row of \mathbf{m} , assign distinct colors to the distinct real values. Is there a condition on the coloring that ensures $\mathbb{H}(\mathbf{m})$ has full column rank?

In answer to the first question we show in Section 2:

Theorem 2. *If $\mathbb{H}(\mathbf{m})$ has full column rank then there is a set R of no more than $k - 1$ of the rows of \mathbf{m} , such that $\mathbb{H}(\mathbf{m}|_R)$ has full column rank.*

*Engineering and Applied Science, California Institute of Technology, slgordon@caltech.edu.

†Engineering and Applied Science, California Institute of Technology, schulman@caltech.edu. Research supported in part by NSF grant CCF-1909972.

Considering the more combinatorial second question, observe that if \mathbf{m} possesses two identical columns then the same is true of $\mathbb{H}(\mathbf{m})$, and so it cannot be full rank. Extending this further, suppose there are three columns C in which only one row r has more than one color. Then $\text{Rowspace } \mathbb{H}(\mathbf{m}|^C)$ is spanned by $\mathbb{1}|^C$ and $r|^C$, so again $\mathbb{H}(\mathbf{m})$ cannot be full rank. Motivated by these necessary conditions, set:

Definition 3. For a matrix Q let $\text{NAE}(Q)$ be the set of nonconstant rows of Q (NAE=“not all equal”); let $\varepsilon(Q|^C) = |\text{NAE}(Q|^C)| - |C|$; and let $\bar{\varepsilon}(Q) = \min_{C \neq \emptyset} \varepsilon(Q|^C)$. If $\bar{\varepsilon}(Q) \geq -1$ we say Q satisfies the NAE condition.

In answer to the second question we have the following:

Theorem 4. *If \mathbf{m} satisfies the NAE condition then*

- (a) *There is a restriction of \mathbf{m} to some $k - 1$ rows R such that $\bar{\varepsilon}(\mathbf{m}|_R) = -1$.*
- (b) *$\mathbb{H}(\mathbf{m})$ is full column rank.*

(As a consequence also $\mathbb{H}(\mathbf{m}|_R)$ is full column rank.)

Apparently the only well-known example of the NAE condition is when \mathbf{m} contains $k - 1$ rows which are identical and whose entries are all distinct. Then the vectors $\mathbf{m}_\emptyset, \mathbf{m}_{\{1\}}, \mathbf{m}_{\{1,2\}}, \dots, \mathbf{m}_{\{1,\dots,k-1\}}$ form a nonsingular Vandermonde matrix. This example shows that the bound of $k - 1$ in (a) is best possible.

For another example in which the NAE condition ensures that $\text{rank } \mathbb{H}(\mathbf{m}) = k$, take the $(k - 1)$ -row matrix with $\mathbf{m}_i^j = 1$ for $i \leq j$ and $\mathbf{m}_i^j = 1/2$ for $i > j$. Here the NAE condition is only minimally satisfied, in that for every $\ell \leq k$ there are ℓ columns C s.t. $\varepsilon(\mathbf{m}|^C) = -1$.

For $k > 3$ the NAE condition is no longer necessary for $\mathbb{H}(\mathbf{m})$ to have full column rank. E.g., for $k = 2^\ell$, the $\ell \times k$ “Hamming matrix” $\mathbf{m}_i^j = (-1)^{j_i}$ where j is an ℓ -bit string $j = (j_1, \dots, j_\ell)$, forms $\mathbb{H}(\mathbf{m}) =$ the Fourier transform for the group $(\mathbb{Z}/2)^\ell$ (often called a Hadamard matrix), which is invertible. Furthermore, almost all (in the sense of Lebesgue measure) $\lceil \lg k \rceil \times k$ matrices \mathbf{m} form a full-rank $\mathbb{H}(\mathbf{m})$. (This is because $\det \mathbb{H}(\mathbf{m})$ is a polynomial in the entries of \mathbf{m} , and the previous example shows the polynomial is nonzero.) Despite this observation, the Vandermonde case, in which $k - 1$ rows are required, is very typical, as it is what arises in $\mathbb{H}(\mathbf{m})$ for a mixture model of observables X_i that are iid conditional on a hidden variable.

2 Some Theory for Hadamard Products, and a Proof of Theorem 2

For $v \in \mathbb{R}^k$ and U a subspace, extend the definition v_\odot to

$$v_\odot(U) = \{u \cdot v_\odot : u \in U\}$$

and introduce the notation

$$v_{\odot}(U) = \text{span}\{U \cup v_\odot(U)\}.$$

We want to understand which subspaces U are invariant under v_{\odot} . Let v have distinct values $\lambda_1 > \dots > \lambda_\ell$ for $\ell \leq k$. Let the polynomials $p_{v,i}$ ($i = 1, \dots, \ell$) of degree $\ell - 1$ be the Lagrange interpolation polynomials for these values, so $p_{v,i}(\lambda_j) = \delta_{ij}$ (Kronecker delta). Let $B(v)$ denote the partition of $[k]$ into blocks $B(v)_{(i)} = \{j : v_j = \lambda_i\}$. Let $V_{(i)}$ be the space spanned by the elementary basis vectors in $B(v)_{(i)}$, and $P_{(i)}$ the projection onto $V_{(i)}$ w.r.t. standard inner product. We have the matrix equation

$$p_{v,i}(v_\odot) = P_{(i)}.$$

The collection of all linear combinations of the matrices $P_{(i)}$ is a commutative algebra, the $B(v)$ projection algebra, which we denote $A_{B(v)}$. The identity of the algebra is $I = \sum P_{(i)}$.

Definition 5. A subspace of \mathbb{R}^k respects $B(v)$ if it is spanned by vectors each of which lies in some $V_{(i)}$.

For U respecting $B(v)$ write $U = \text{span}(\bigcup U_{(i)})$ for $U_{(i)} \subseteq V_{(i)}$. Let $D_{(i)} = (U_{(i)})^\perp \cap V_{(i)}$. Then $(U_{(i)})^\perp = D_{(i)} \oplus \bigoplus_{j \neq i} V_{(j)}$.

Lemma 6. *Subspace U^\perp respects $B(v)$ if U does.*

Proof. In general, $(\text{span}(W \cup W'))^\perp = W^\perp \cap W'^\perp$. So $U^\perp = \bigcap (U_{(i)})^\perp = \bigoplus D_{(i)}$. \square

Lemma 7. *Subspace U respects $B(v)$ iff $U = \bigoplus (P_{(i)}U)$.*

Proof. (\Leftarrow): Because this gives an explicit representation of U as a direct sum of subspaces each restricted to some $V_{(i)}$. (\Rightarrow): By definition U is spanned by some collection of subspaces $V'_{(i)} \subseteq V_{(i)}$; since these subspaces are necessarily orthogonal, $U = \bigoplus V'_{(i)}$. Moreover, since $P_{(i)}$ annihilates $V_{(j)}, j \neq i$, and is the identity on $V_{(i)}$, it follows that each $V'_{(i)} = P_{(i)}U$. \square

Theorem 8. *Subspace U is invariant under v_\odot iff U respects $B(v)$.*

Proof. (\Leftarrow): It suffices to show U^\perp is invariant under v_\odot . By the previous lemma, it is equivalent to suppose that U^\perp respects $B(v)$. So let $d \in U^\perp$ and write $d = \sum d_i, d_i \in D_{(i)}$. Then $v \odot d_i = \lambda_i d_i \in D_{(i)}$. So $v \odot d = \sum v \odot d_i \in \bigoplus D_{(i)} = U^\perp$.

(\Rightarrow): If $U = v_\odot(U)$ then these also equal $v_\odot(v_\odot(U))$, etc., so U is an invariant space of $A_{B(v)}$, meaning, $aU \subseteq U$ for any $a \in A_{B(v)}$. In particular for $a = P_{(i)}$. So $U \supseteq \bigoplus (P_{(i)}U)$. On the other hand, since $\sum P_{(i)} = I$, $U = (\sum P_{(i)})U \subseteq \bigoplus (P_{(i)}U)$. So $U = \bigoplus (P_{(i)}U)$. Now apply Lemma 7. \square

The symbol \subset is reserved for strict inclusion.

Lemma 9. *If $S, T \subseteq [n]$ and $\text{Rowspace } \mathbb{H}(\mathbf{m}|_S) \subset \text{Rowspace } \mathbb{H}(\mathbf{m}|_{S \cup T})$, then there is a row $t \in T$ such that $\text{Rowspace } \mathbb{H}(\mathbf{m}|_S) \subset \text{Rowspace } \mathbb{H}(\mathbf{m}|_{S \cup \{t\}})$.*

Proof. Without loss of generality S, T are disjoint. Let $T' \subseteq T$ be a smallest set s.t. $\exists S' \subseteq S$ s.t. $\mathbf{m}_{S'} \odot \mathbf{m}_{T'} \notin \text{Rowspace } \mathbb{H}(\mathbf{m}|_S)$. Select any $t \in T'$ and write $\mathbf{m}_{S'} \odot \mathbf{m}_{T'} = \mathbf{m}_{S'} \odot \mathbf{m}_{T' - \{t\}} \odot \mathbf{m}_t$. By minimality of T' , $\mathbf{m}_{S'} \odot \mathbf{m}_{T' - \{t\}} \in \text{Rowspace } \mathbb{H}(\mathbf{m}|_S)$. But then $\mathbf{m}_{S'} \odot \mathbf{m}_{T'} \in \text{Rowspace } \mathbb{H}(\mathbf{m}|_{S \cup \{t\}})$, so $\text{Rowspace } \mathbb{H}(\mathbf{m}|_S) \subset \text{Rowspace } \mathbb{H}(\mathbf{m}|_{S \cup \{t\}})$. \square

Theorem 2 is now a consequence of Lemma 9. \square

It follows from Theorem 2 that we can check whether $\text{rank } \mathbb{H}(\mathbf{m}) = k$ in time $O(n)^k$ by computing $\text{rank } \mathbb{H}(\mathbf{m}|_S)$ for each $S \in \binom{[n]}{k-1}$.

3 Combinatorics of the NAE Condition: Proof of Theorem 4 (a)

Recall we are to show: 4 (a): *If $\bar{\varepsilon}(\mathbf{m}) \geq -1$ then \mathbf{m} has a restriction to some $k-1$ rows on which $\bar{\varepsilon} = -1$.*

Proof. We induct on k . The (vacuous) base-case is $k = 1$. For $k > 1$, we induct on n , with base-case $n = k - 1$.

Supposing the Theorem fails for $k, k > 1$, let \mathbf{m} be a k -column counterexample with least n . Necessarily every row is in $\text{NAE}(\mathbf{m})$, and $n > k - 1 \geq 1$. We will show \mathbf{m} has a restriction \mathbf{m}' to $n - 1$ rows, for which $\bar{\varepsilon}(\mathbf{m}') \geq -1$; this will imply a contradiction because, by minimality of \mathbf{m} , \mathbf{m}' has a restriction to $k - 1$ rows on which $\bar{\varepsilon} = -1$.

If $\bar{\varepsilon}(\mathbf{m}) \geq 0$ then we can remove any single row of \mathbf{m} and still satisfy $\bar{\varepsilon} \geq -1$.

Otherwise, $\bar{\varepsilon}(\mathbf{m}) = -1$, so there is a nonempty S such that $|\text{NAE}(\mathbf{m}|^S)| = |S| - 1$; choose a largest such S . It cannot be that $S = [k]$ (as then $n = k - 1$). Arrange the rows $\text{NAE}(\mathbf{m}|^S)$ as the bottom $|S| - 1$ rows of the matrix. As discussed earlier, for the NAE condition one may regard the distinct real values in each row of \mathbf{m} simply as distinct colors; relabel the colors in each row above $\text{NAE}(\mathbf{m}|^S)$ so the color above S is called ‘‘white.’’ (There need be no consistency among the real numbers called white in different rows.) See Figure 1.

Due to the maximality of $|S|$, there is no white rectangle on ℓ columns and $n - |S| - \ell + 1$ rows inside $\mathbf{m}|_{[n] - \text{NAE}(\mathbf{m}|^S)}^{[k] - S}$ for any $\ell \geq 1$. That is to say, if we form a bipartite graph on right vertices corresponding to the columns $[k] - S$, and left vertices corresponding to the rows $[n] - \text{NAE}(\mathbf{m}|^S)$, with non-white cells being edges, then any subset of the right vertices of size $\ell \geq 1$ has at least $\ell + 1$ neighbors within the left vertices.




Figure 1: Argument for Theorem 4 (a). Upper-left region is white. Entries $(t, f(t))$ are not white.

By the induction on k (since $S \neq \emptyset$), for the set of columns $[k] - S$ there is a set R'' of $k - |S| - 1$ rows such that $\bar{\varepsilon}(\mathbf{m}|_{R''}^{[k]-S}) = -1$. Together with the rows of $\text{NAE}(\mathbf{m}|^S)$ this amounts to at most $k - 2$ rows, so since $n \geq k$, we can find two rows outside this union; delete either one of them, leaving a matrix \mathbf{m}' with $n - 1$ rows. This matrix has the rows $\text{NAE}(\mathbf{m}|^S)$ at the bottom, and $n - |S|$ remaining rows which we call R' . The lemma will follow by showing that $\bar{\varepsilon}(\mathbf{m}') \geq -1$.

In \mathbf{m}' , the induced bipartite graph on right vertices $[k] - S$ and left vertices R' has the property that any right subset of size $\ell \geq 1$ has a neighborhood of size at least ℓ in R' . Applying Hall's Marriage Theorem, there is an injective $f : [k] - S \rightarrow R'$ employing only edges of the graph.

Now consider any set of columns T , $T = T_1 \cup T_2, T_1 \subseteq [k] - S, T_2 \subseteq S$. We need to show that $\varepsilon(\mathbf{m}|^T) \geq -1$. Let $R_1 = \text{NAE}(\mathbf{m}|^{T_1}) \cap R''$, $R_2 = \text{NAE}(\mathbf{m}|^{T_2}) \subseteq \text{NAE}(\mathbf{m}|^S)$, and note that $|R_1| \geq |T_1| - 1$, $|R_2| \geq |T_2| - 1$. If $T_2 = \emptyset$ we simply use R_1 . Likewise if $T_1 = \emptyset$, we use R_2 .

If both T_1 and T_2 are nonempty, $\text{NAE}(\mathbf{m}|^{T_2}) \subseteq \text{NAE}(\mathbf{m}|^S)$, and $|\text{NAE}(\mathbf{m}|^{T_2})| \geq |T_2| - 1$. Now use the matching f . The set of rows $f(T_1)$ lies in R' and is therefore disjoint from $\text{NAE}(\mathbf{m}|^{T_2})$. Moreover since $T_2 \neq \emptyset$, every entry (t, j) for $t \in T_2, j \in R'$ is white. On the other hand due to the construction of f , for every $t \in T_1$ the entry $(t, f(t))$ is non-white. Therefore every row in $f(T_1)$ is in $\text{NAE}(\mathbf{m}|^{T_1 \cup T_2})$. So $|\text{NAE}(\mathbf{m}|^{T_1 \cup T_2})| \geq |T_2| - 1 + |T_1|$. Thus $\bar{\varepsilon}(\mathbf{m}') \geq -1$. \square

4 From NAE to Rank: Proof of Theorem 4 (b)

Recall we are to show: 4 (b): $\mathbb{H}(\mathbf{m})$ has full column rank if $\bar{\varepsilon}(\mathbf{m}) \geq -1$.


Proof. The case $k = 1$ is trivial. Now suppose $k \geq 2$ and that Theorem 4 (b) holds for all $k' < k$. Any constant rows of \mathbf{m} affect neither the hypothesis nor the conclusion, so remove them, leaving \mathbf{m} with at least $k - 1$ rows. Now pick any set, C , of $k - 1$ columns of \mathbf{m} . By Theorem 4 (a) there are some $k - 2$ rows of \mathbf{m} , call them R' , on which $\bar{\varepsilon}(\mathbf{m}|_{R'}^C) = -1$. Let v be a row of \mathbf{m} outside R' . Call the rows of \mathbf{m} apart from v , R'' . Since R'' contains

R' , by induction $\dim \text{Rowspace } \mathbb{H}(\mathbf{m}|_{R'})^C = k - 1$. Therefore $U := \text{Rowspace } \mathbb{H}(\mathbf{m}|_{R'}) \subseteq \mathbb{R}^k$ is of dimension at least $k - 1$. We claim now that $\dim U = k$.

Suppose to the contrary that $\dim U = k - 1$. If $v_\odot(U) \subseteq U$ then as proven earlier in Theorem 8, U respects $B(v)$. Since v is nonconstant, $B(v)$ is a partition of $[k]$ into $\ell \geq 2$ nonempty blocks $B(v)_{(i)}$, and $U = \bigoplus_{i=1}^{\ell} U_{(i)}$ with $U_{(i)} = P_{(i)}U_{(i)}$. So there is some i_0 for which $U_{(i_0)} \subset V_{(i_0)}$; specifically, $U_{(i)} = V_{(i)}$ for all $i \neq i_0$, and $\dim U_{(i_0)} = \dim V_{(i_0)} - 1$. Since $|B(v)_{(i_0)}| < k$, we know by induction that the rows of $\mathbb{H}(\mathbf{m})$ span $V_{(i_0)}$. Thus in fact $U = \mathbb{R}^k$. (Further detail for the last step: let $w \in \mathbb{R}^k$. Since the rows of $\mathbb{H}(\mathbf{m})$ span $V_{(i_0)}$, there is a $w' \in \text{Rowspace } \mathbb{H}(\mathbf{m})$ s.t. $P_{(i_0)}w' = P_{(i_0)}w$. Moreover since $U_{(i)} = V_{(i)}$ for all $i \neq i_0$, there is a $w'' \in U$ s.t. $w'' = (I - P_{(i_0)})(w - w')$. Then $w' + w'' \in \text{Rowspace } \mathbb{H}(\mathbf{m})$, and $w' + w'' = w$.) \square

5 Motivation

Consider *observable* random variables X_1, \dots, X_n that are statistically independent conditional on H , a *hidden* random variable H supported on $\{1, \dots, k\}$. (See causal diagram.)



The most fundamental case is that the X_i are binary. Then we denote $\mathbf{m}_i^j = \Pr(X_i = 1 | H = j)$. The model parameters are \mathbf{m} along with a probability distribution (the *mixture* distribution) $\pi = (\pi_1, \dots, \pi_k)$ on H .

Finite mixture models were pioneered in the late 1800s in [13, 14]. The problem of learning such distributions has drawn a great deal of attention. For surveys see, e.g., [5, 17, 11, 12]. For some algorithmic papers on discrete X_i , see [9, 4, 7, 2, 6, 1, 15, 10, 3, 8]. The source identification problem is that of computing (\mathbf{m}, π) from the joint statistics of the X_i . Put another way, the problem is to invert the multilinear moment map

$$\begin{aligned} \mu : (\mathbf{m}, \pi) &\rightarrow \mathbb{R}^{2^{[n]}} \\ \mu(\mathbf{m}, \pi)_S &= \Pr(X_S = 1) \quad \text{where } S \subseteq [n], X_S = \prod_{i \in S} X_i \\ &= \mathbf{m}_S \cdot \pi^\top \end{aligned}$$

The last line shows the significance of $\mathbb{H}(\mathbf{m})$ to mixture model identification, since $\mathbf{m}_S^j = \Pr(X_S = 1 | H = j)$.

Connection to rank $\mathbb{H}(\mathbf{m})$. In general μ is not injective (even allowing for permutation among the values of H). For instance it is clearly not injective if \mathbf{m} has two identical columns (unless π places no weight on those). More generally, and assuming all $\pi_j > 0$, it cannot be injective unless $\mathbb{H}(\mathbf{m})$ has full column rank.

One sufficient condition for injectivity, due to [16], is that there be $2k - 1$ “separated” observables X_i ; X_i is separated if all \mathbf{m}_i^j are distinct, or in our terminology, if no color recurs in \mathbf{m}_i . (Further [8], one can lower bound the distance between $\mu(\mathbf{m}, \pi)$ and any $\mu(\mathbf{m}', \pi')$ in terms of $\min_{i,j} |\mathbf{m}_i^j - \mathbf{m}_i^{j'}|$ and the distance between (\mathbf{m}, π) and (\mathbf{m}', π') .)

A weaker sufficient condition for injectivity of μ , due to [8], is that for every $i \in [n]$ there exist two disjoint sets $A, B \subseteq [n] - \{i\}$ such that $\mathbb{H}(\mathbf{m}|_A)$ and $\mathbb{H}(\mathbf{m}|_B)$ have full column rank. (It is not known whether two disjoint such A, B are strictly necessary, but the implied $n \leq 2k - 1$ is in general best possible [15].)

References

- [1] A. Anandkumar, D. Hsu, and S. M. Kakade. A method of moments for mixture models and hidden Markov models. In *Proc. 25th Ann. Conf. on Computational Learning Theory*, pages 33.1–33.34, 2012.
- [2] K. Chaudhuri and S. Rao. Learning mixtures of product distributions using correlations and independence. In *Proc. 21st Ann. Conf. on Computational Learning Theory*, pages 9–20, 2008.
- [3] S. Chen and A. Moitra. Beyond the low-degree algorithm: mixtures of subcubes and their applications. In *Proc. 51st Ann. ACM Symp. on Theory of Computing*, pages 869–880, 2019.
- [4] M. Cryan, L. Goldberg, and P. Goldberg. Evolutionary trees can be learned in polynomial time in the two state general Markov model. *SIAM J. Comput.*, 31(2):375–397, 2001. Prev. FOCS '98.
- [5] B. S. Everitt and D. J. Hand. *Mixtures of discrete distributions*, pages 89–105. Springer Netherlands, Dordrecht, 1981.
- [6] J. Feldman, R. O’Donnell, and R. A. Servedio. Learning mixtures of product distributions over discrete domains. *SIAM J. Comput.*, 37(5):1536–1564, 2008.
- [7] Y. Freund and Y. Mansour. Estimating a mixture of two product distributions. In *Proc. 12th Ann. Conf. on Computational Learning Theory*, pages 183–192, July 1999.
- [8] S. L. Gordon, B. Mazaheri, Y. Rabani, and L. J. Schulman. Source identification for mixtures of product distributions. arXiv:2012.14540, 2020.
- [9] M. Kearns, Y. Mansour, D. Ron, R. Rubinfeld, R. Schapire, and L. Sellie. On the learnability of discrete distributions. In *Proc. 26th Ann. ACM Symp. on Theory of Computing*, pages 273–282, 1994.
- [10] J. Li, Y. Rabani, L. J. Schulman, and C. Swamy. Learning arbitrary statistical mixtures of discrete distributions. In *Proc. 47th Ann. ACM Symp. on Theory of Computing*, pages 743–752, 2015.
- [11] B. G. Lindsay. Mixture models: theory, geometry and applications. In *NSF-CBMS regional conference series in probability and statistics*, pages i–163. JSTOR, 1995.
- [12] G. J. McLachlan, S. X. Lee, and S. I. Rathnayake. Finite mixture models. *Annual Review of Statistics and Its Application*, 6(1):355–378, 2019.
- [13] S. Newcomb. A generalized theory of the combination of observations so as to obtain the best result. *American Journal of Mathematics*, 8(4):343–366, 1886.
- [14] K. Pearson. Contributions to the mathematical theory of evolution III. *Philosophical Transactions of the Royal Society of London (A.)*, 185:71–110, 1894.
- [15] Y. Rabani, L. J. Schulman, and C. Swamy. Learning mixtures of arbitrary distributions over large discrete domains. In *Proc. 5th Conf. on Innovations in Theoretical Computer Science*, pages 207–224, 2014.
- [16] B. Tahmasebi, S. A. Motahari, and M. A. Maddah-Ali. On the identifiability of finite mixtures of finite product measures. IEEE International Symposium on Information Theory (ISIT) 2018 and arXiv:1807.05444v1, 2018.
- [17] D. M. Titterton, A. F. M. Smith, and U. E. Makov. *Statistical Analysis of Finite Mixture Distributions*. John Wiley and Sons, Inc., 1985.