

Continuous One-Counter Automata

Michael Blondin*, Tim Leys† Filip Mazowiecki‡, Philip Offtermatt*‡, and Guillermo A. Pérez†

*Université de Sherbrooke, Canada

†University of Antwerp, Belgium

‡Max Planck Institute for Software Systems, Germany

Abstract

We study the reachability problem for continuous one-counter automata, COCA for short. In such automata, transitions are guarded by upper and lower bound tests against the counter value. Additionally, the counter updates associated with taking transitions can be (non-deterministically) scaled down by a nonzero factor between zero and one. Our three main results are as follows: (1) We prove that the reachability problem for COCA with global upper and lower bound tests is in NC²; (2) that, in general, the problem is decidable in polynomial time; and (3) that it is decidable in the polynomial hierarchy for COCA with parametric counter updates and bound tests.

I. INTRODUCTION

Counter machines form a fundamental computational model which captures the behavior of infinite-state systems. Unfortunately, their central decision problems, like the (*configuration*) *reachability problem*, are undecidable as the model is Turing-complete [1], [2]. To circumvent this issue, numerous restrictions of counter machines have been studied in the literature. For instance, vector addition systems with states (VASS) arise from restricting the type of tests that can be used to guard transitions [3], [4], [5], [6]. One-counter automata are yet another well-studied model [7], [8], [9], in this case arising from the restriction to a single counter, hence the name.

We consider one-counter automata that can use tests of the form “ $\leq c$ ” and “ $\geq d$ ” — where c and d are constants — to guard their transitions. As a natural extension of finite-state automata, one-counter automata allow for better conservative approximations of classical static-analysis problems like instruction reachability (see, e.g., *program graphs* as defined in [10]). They also enable the verification of programs with lists [11] and XML-stream validation [12]. Furthermore, their reachability problem seems intrinsically connected to that of timed automata (TA). The reachability problem for two-clock TA is known to be logspace-equivalent to the same problem for succinct one-counter automata (SOCA), that is, where constants used in counter updates and tests are encoded in binary [13]. An analogue of this connection holds when SOCA are enriched with parameters that can be used on updates: reachability for two-parametric-clock TA reduces to the (existential) reachability problem for parametric one-counter automata [14]. Interestingly, Alur et al. observe [15] the former subsumes a long-standing open problem of Ibarra [16] concerning “simple programs”.

All of the above connections from interesting problems to reachability for SOCA and parametric SOCA indicate that efficient algorithms for the problem are very much desirable. Unfortunately, it is known that reachability for SOCA (with upper and lower-bound tests) is PSPACE-complete [8]. For parametric SOCA, the situation is even worse as the general problem is not even known to be decidable. In this work, we study *continuous relaxations* of these problems and show that their complexities belong in tractable complexity classes. We thus give the first efficient conservative approximation for the reachability problem for SOCA and parametric SOCA.

The continuous relaxations: We observe that the model considered by Fearnley and Jurdziński [8] is not precisely our SOCA, rather they consider *bounded 1-dimensional VASS*. In such VASS, the counter is not allowed to take negative values. Additionally, it is not allowed to take values greater than some global upper bound. Note that inequality tests against constants can be added to such VASS as “syntactic sugar” since these can be implemented making use of the upper and lower bounds. These observations allow us to adapt Blondin and Haase’s definition of continuous VASS [17] to introduce (*bounded*) *continuous one-counter automata* (COCA) which have *global* upper and lower-bound tests: Transitions are allowed to be “partially taken” in the sense that the respective counter updates can be scaled by some factor $\alpha \in (0, 1]$.

In contrast to the situation in the discrete world, because of the continuous semantics, adding arbitrary upper and lower-bound tests to COCA does result in the more expressive model of *guarded COCA*. Importantly, guarded COCA are a “tighter” relaxation of SOCA than COCA are (via the translation to bounded 1-VASS). Finally, we also study the reachability problem for *parametric COCA*. These are guarded COCA where counter updates can be variables $x \in X$ whose values range over the rationals; bound tests can also be against variables from X . The resulting model can be seen as a continuous relaxation of Ibarra’s simple programs [16], [14].

Contributions: Our main contributions are three-fold (see [Theorem 1](#)). First, we show that the reachability problem for COCA is decidable in NC². Second, we give a polynomial-time algorithm for the same problem for guarded COCA. Finally, we show that the reachability problem for parametric COCA belongs to Σ_2^P and is NP-hard.

On the way, we prove that the reachability problem for COCA enriched with equality tests is in NC^2 ; that the reachability problem for parametric COCA where only counter updates are allowed to be parametric is equivalent to the integer-valuation restriction of the problem; and that the reachability problem for acyclic parametric COCA is NP -complete.

Other related work: To complete a full circle of connections between timed and counter automata, we note that the closest model to ours is that of one-clock TA. The value of the clock in such automata evolves (continuously) at a fixed positive rate and can be reset by some transitions. Guarded COCA can simulate clock delays using $+1$ self-loops and resets using -1 self-loops and bound tests “ ≤ 0 ” and “ ≥ 0 ”. Our model thus generalises one-clock TA.

The reachability problem for (non-parametric) one-clock TA is NL -complete [18]. The NL membership proof from [18] relies on the fact that clock delays can always occur and do so without changing the state. This does not hold in the more general framework of COCA. Consequently, the proof does not extend directly to COCA.

The reachability problem for parametric one-clock TA with integer-valued parameters is known to be decidable in NEXP [14]. Since non-parametric clocks can be removed at the cost of an exponential blow-up [15], it is also argued in [14] that the problem belongs to N2EXP if an arbitrary number of non-parametric clocks is allowed [14]. For the latter problem, the authors also prove that it is NEXP -hard. Our Σ_2^P upper bound for update-parametric COCA with integer-valued parameters improves the latter two bounds.

II. PRELIMINARIES

We write $\mathbb{Q}_{\geq 0}$ for the set of nonnegative rationals, and $\mathbb{Q}_{> 0}$ for the set of positive rationals. We use symbols “[” and “]” for closed intervals, and “(” and “)” for open intervals of rational numbers. For example, $[a, b]$ denotes $\{q \in \mathbb{Q} \mid a \leq q \leq b\}$. Intervals do not have to be bounded, e.g. we allow $[3, +\infty)$. We denote the set of all intervals over \mathbb{Q} by \mathcal{I} . We write \bar{X} to denote the *closure* of a set $X \subseteq \mathbb{Q}$, i.e. X enlarged with its limit points. For example, $\overline{(3, 5)} = [3, 5]$, $\overline{[1, 4) \cup (4, 5]} = [1, 5]$ and $\overline{(-\infty, +\infty)} = (-\infty, +\infty)$. Throughout the paper, numbers are encoded in *binary* and we assume intervals to be encoded as pairs of endpoints, together with binary flags indicating whether the endpoints are contained or not.

A. One-counter automata

A *continuous one-counter automaton* (COCA) is a triple $\mathcal{V} = (Q, T, \tau)$, where Q and $T \subseteq Q \times \mathbb{Z} \times Q$ are finite sets of *states* and *transitions*, and $\tau \in \mathcal{I}$. A *configuration* of \mathcal{V} is a pair $(q, a) \in Q \times \mathbb{Q}$, denoted $q(a)$. A *run* from $p(a)$ to $q(b)$ in \mathcal{V} is a sequence $\alpha_1 t_1 \cdots \alpha_n t_n$, where $\alpha_i \in (0, 1]$ and $t_i = (q_{i-1}, z_i, q_i) \in T$, for which there exist configurations $q_0(a_0), \dots, q_n(a_n)$ such that $q_0(a_0) = p(a)$, $q_n(a_n) = q(b)$ and $a_i = a_{i-1} + \alpha_i \cdot z_i$ for all $i \in \{1, \dots, n\}$. We say that such a run is *admissible* if $a_0, \dots, a_n \in \tau$. For readers familiar with one-counter automata, note that the model of one-counter nets is obtained by setting $\tau = [0, +\infty)$ and $\alpha_i = 1$ for all i .

A *guarded COCA* is a triple $\mathcal{W} = (Q, T, \tau)$, where (Q, T) is as for a COCA and $\tau : Q \rightarrow \mathcal{I}$ assigns intervals to states. *Configurations and runs* of \mathcal{W} are defined as for a COCA. A run is *admissible* if each of its configurations $q_i(a_i)$ satisfies $a_i \in \tau(q_i)$. Hence, a COCA can be seen as a guarded COCA where $\tau(q)$ is the same for all $q \in Q$.

The set \mathcal{I}_X of *parameterised intervals* over a set X is the set of intervals whose endpoints belong either to $\mathbb{Q} \cup \{-\infty, +\infty\}$ or X . A *parametric COCA* is a tuple $\mathcal{P} = (Q, T, \tau, X)$, where Q , X and $T \subseteq Q \times (\mathbb{Z} \cup X) \times Q$ are finite sets of *states*, *parameters* and *transitions*; and where $\tau : Q \rightarrow \mathcal{I}_X$. A *valuation* of X is a function $\mu : X \rightarrow \mathbb{Q}$. We write $\mathcal{P}^\mu = (Q, T^\mu, \tau^\mu)$ to denote the guarded COCA obtained from \mathcal{P} by replacing each parameter $x \in X$, occurring in T and τ , with $\mu(x)$. We say that there is a run from $p(a)$ to $q(b)$ in \mathcal{P} if there exists a valuation μ such that \mathcal{P}^μ has a run from $p(a)$ to $q(b)$. In particular, \mathcal{P} is a guarded COCA if $X = \emptyset$. Otherwise, the notion of run only makes sense w.r.t. a valuation μ , i.e. in the guarded COCA \mathcal{P}^μ .

In summary, we deal with three increasingly richer models: $\text{COCA} \subseteq \text{guarded COCA} \subseteq \text{parametric COCA}$. In all variants, the *size* of the automaton is $|Q| + |T| \cdot s$, where s is the maximal number of bits required to encode a number in T and τ .

B. Runs, paths and cycles

Let $\mathcal{W} = (Q, T, \tau)$ be a guarded COCA. We write $\text{Paths}_{p,q}$ to denote the set of paths from state $p \in Q$ to state $q \in Q$ in the graph induced by T . Let $\rho = \alpha_1 t_1 \cdots \alpha_n t_n$ be a run where each $t_i = (q_{i-1}, z_i, q_i)$. The *underlying path* of ρ is $\text{path}(\rho) := t_1 \cdots t_n \in \text{Paths}_{q_0, q_n}$. We further define $\rho[i..j] := \alpha_i t_i \cdots \alpha_j t_j$, $\rho_i := \rho[i..i]$, $\text{in}(\rho) := q_0$, $\text{out}(\rho) := q_n$ and $\Delta(\rho) := \sum_{i=1}^n \alpha_i z_i$. By convention, $\rho[i..j] := \varepsilon$ if $j < i$, and $\Delta(\varepsilon) := 0$. We write $p(a) \rightarrow_\rho q(b)$ to denote the fact that ρ is admissible from $p(a)$ to $q(b)$. Since states p and q are determined by ρ , we may omit them and simply write $a \rightarrow_\rho b$. For every $\beta \in (0, 1]$, we define $\beta\rho := (\beta\alpha_1)t_1 \cdots (\beta\alpha_n)t_n$. Note that $\beta\rho$ is a run, but it may not preserve admissibility.

Let $\pi = t_1 \cdots t_n \in \text{Paths}_{p,q}$ be such that each $t_i = (q_{i-1}, z_i, q_i)$. We say that π is a *cycle* if $p = q$, and *simple* if π does not repeat any state. Let $\Delta(\pi) := z_1 + \dots + z_n$, $\Delta^+(\pi) := \sum_{i=1}^n \max(0, z_i)$ and $\Delta^-(\pi) := \sum_{i=1}^n \min(0, z_i)$, with $\Delta(\varepsilon) = \Delta^+(\varepsilon) = \Delta^-(\varepsilon) := 0$. In particular, $\Delta^-(\pi) \leq 0 \leq \Delta^+(\pi)$. Moreover, scaling the positive or negative transitions of a path π arbitrarily close to zero yields a run of effect arbitrarily close to $\Delta^-(\pi)$ or $\Delta^+(\pi)$.




Fig. 1. A guarded COCA; each state s is labeled with the interval $\tau(s)$.

We write $p(a) \rightarrow_{\pi} q(b)$ to denote the existence of a run ρ such that $p(a) \rightarrow_{\rho} q(b)$ and $\text{path}(\rho) = \pi$. As for runs, we may omit states and simply write $a \rightarrow_{\pi} b$. The *reachability function* given by π is defined as $\text{Post}_{\pi}(a) := \{b \in \mathbb{Q} \mid p(a) \rightarrow_{\pi} q(b)\}$. We generalise this notion to sets of paths and numbers:

$$\text{Post}_S(A) := \bigcup_{\pi \in S} \bigcup_{a \in A} \text{Post}_{\pi}(a).$$

If $S = \text{Paths}_{p,q}$, we write $\text{Post}_{p,q}(a)$ and $\text{Post}_{p,q}(A)$. For example, for the guarded COCA of Figure 1, the following holds: $\text{Post}_{p,q}(a) = (10, 18) \cup [19, 100]$ if $a = 15$; $(a - 5, a + 3)$ if $a \in [-5, 15]$; and \emptyset otherwise.

Finally, we define the set of starting points as $\text{enab}(\pi) := \{a \in \mathbb{Q} \mid \text{Post}_{\pi}(a) \neq \emptyset\}$ and $\text{enab}(S) := \bigcup_{\pi \in S} \text{enab}(\pi)$.

C. Our contribution

In this work, we study the *reachability problem* that asks the following question: Given a COCA or a guarded COCA \mathcal{W} with configurations $p(a)$ and $q(b)$, is there an admissible run from $p(a)$ to $q(b)$? In other words, by abbreviating “ $\text{Paths}_{p,q}$ ” with “ $*$ ”, the problem asks whether $p(a) \rightarrow_* q(b)$ holds. For parametric COCAs, the (*existential*) *reachability problem* asks whether $p(a) \rightarrow_* q(b)$ for some parameter valuation.

We will establish the following complexity results:

Theorem 1. *The reachability problem is:*

- 1) in NC^2 for COCAs;
- 2) in P for guarded COCAs; and
- 3) NP -hard and in Σ_2^{P} for parametric COCAs.

Recall that Σ_2^{P} is the level of the polynomial hierarchy that corresponds to NP^{NP} . Moreover, NC is the class of problems solvable in polylogarithmic parallel time, i.e. $\text{NC} = \bigcup_{i \geq 0} \text{NC}^i$ where NC^i is the class of problems decidable by logspace-uniform families of circuits of polynomial size, depth $\mathcal{O}(\log^i n)$ and bounded fan-in (e.g., see [19], [20] for a more thorough definition). It is well-known that $\text{NL} \subseteq \text{NC}^2 \subseteq \text{P}$. We also refer to the functional variant of NC^i as NC^i .

The two first results of Theorem 1 are obtained by characterising reachability functions and by showing how to efficiently compute their representation. More precisely, we show:

Proposition 2. *Let \mathcal{W} be a COCA or a guarded COCA. It is the case that:*

- 1) If \mathcal{W} is a COCA, then $\text{Post}_{p,q}(a)$ consists of at most two intervals whose representations are computable in NC^2 ;
- 2) $\text{Post}_{p,q}(a)$ is made of $|\mathcal{W}|^{\mathcal{O}(1)}$ intervals, and a representation of $\text{Post}_{p,q}(a)$ is computable in polynomial time.

To derive the third result, i.e. the Σ_2^{P} upper bound, we borrow ideas from the above technical results to reduce reachability to determining the truth value of linear arithmetic Σ_2 -sentences.

III. COCA REACHABILITY

In this section, we prove that the reachability problem for COCAs belongs in NC^2 by showing how to compute a representation of $\text{Post}_{p,q}(a)$ from some a . In the remainder, we fix a COCA $\mathcal{V} = (Q, T, \tau)$.

A. Testing emptiness

We first aim to show that deciding whether $\text{Post}_{p,q}(a) \neq \emptyset$ can be checked in NC^2 . To this end, we first state some simple graph properties checkable in NC^2 . For a path π , let us write $\text{first}(\pi)$ (resp. $\text{last}(\pi)$) to denote the first (resp. last) index such that $\Delta(\pi_i) \neq 0$ if any, and $\text{first}(\pi) = \text{last}(\pi) := \infty$ if none. We naturally extend the notations $\Delta^+(\pi)$, $\Delta^-(\pi)$, $\Delta(\pi)$, $\text{first}(\pi)$ and $\text{last}(\pi)$ to weighted multigraphs.

This lemma follows from standard results on NC^2 :

Lemma 3. *Let $G = (Q, E)$ be a weighted multigraph whose weights are encoded in binary, and let $p, q \in Q$ be nodes. Deciding whether $S = \emptyset$ is in NC^2 , where S is the set of paths $\pi \in \text{Paths}_{p,q}$ that satisfy a fixed subset of these conditions¹:*

¹The set of conditions may be empty, in which case $S = \text{Paths}_{p,q}$.

- (a) $\Delta^+(\pi) \neq 0$ (resp. $\Delta^-(\pi) \neq 0$);
- (b) $\Delta^+(\pi) = 0$ (resp. $\Delta^-(\pi) = 0$);
- (c) $\Delta(\text{first}(\pi)) < 0$ (resp. $\Delta(\text{first}(\pi)) > 0$);
- (d) $\Delta(\text{last}(\pi)) < 0$ (resp. $\Delta(\text{last}(\pi)) > 0$).

Furthermore, for any such set S , the following value can be computed in NC^2 : $\text{opt}\{w(\pi) \mid \pi \in S \text{ and } |\pi| \leq |Q|\}$, where $\text{opt} \in \{\min, \max\}$ and $w \in \{\Delta^+, \Delta^-\}$.

In essence, conditions (a)–(d) above can be checked through preprocessing and standard graph reachability, e.g. for (c), we make a copy of G restricted to zero transitions, which branches into a full copy of G via negative (resp. positive) transitions. Optimising the value of such paths can be done by a typical “divide-and-conquer procedure” that optimizes paths of length $1, 2, 4, \dots, |Q|$ (hence $\log |Q|$ levels of NC^1 computations).

Lemma 4. *Let $a \in \mathbb{Q}$, $p, q \in Q$, and $\pi \in \text{Paths}_{p,q}$. We have $a \in \text{enab}(\pi)$ iff $a \in \tau$ and any of these conditions hold:*

- (a) $a \notin \{\inf \tau, \sup \tau\}$;
- (b) $a = \inf \tau = \sup \tau$, $\text{first}(\pi) = \infty$;
- (c) $a = \inf \tau < \sup \tau$, $\text{first}(\pi) \neq \infty \implies \Delta(\pi_{\text{first}(\pi)}) > 0$;
- (d) $a = \sup \tau > \inf \tau$, $\text{first}(\pi) \neq \infty \implies \Delta(\pi_{\text{first}(\pi)}) < 0$.

Proof. Having $a \in \tau$ is obviously necessary, so we assume it holds throughout the proof.

\Leftarrow We proceed by induction on $|\pi|$. If $|\pi| = 0$, then the claim is trivial as the empty path is admissible from a . Assume $|\pi| = n > 0$ and π satisfies a condition. Let $t := \pi_1$ and $\sigma := \pi[2..n]$. If (a) holds, then $a \rightarrow_{\beta t} a'$ for some $a' \in \tau \setminus \{\inf \tau, \sup \tau\}$ and sufficiently small $\beta \in (0, 1]$. If (b) holds, then $a \rightarrow_t a' = a$ as $\Delta(t) = 0$. If (c) or (d) holds, then either $a \rightarrow_t a' = a$ if $\Delta(t) = 0$, or $a \rightarrow_{\beta t} a'$ for some $a' \in \tau \setminus \{\inf \tau, \sup \tau\}$ and sufficiently small $\beta \in (0, 1]$ otherwise. In all cases, σ satisfies one of the conditions w.r.t. value a' . Thus, we are done by the induction hypothesis.

\Rightarrow Towards a contradiction, let us assume that $a \in \text{enab}(\pi)$ and that no condition is satisfied. If $a = \inf \tau = \sup \tau$ and $\text{first}(\pi) \neq \infty$, then there is obviously a contradiction. Otherwise, either (i) $a = \inf \tau$ and $\Delta(\pi_{\text{first}(\pi)}) < 0$; or (ii) $a = \sup \tau$ and $\Delta(\pi_{\text{first}(\pi)}) > 0$. We only consider (ii) as (i) is symmetric. Let $a \rightarrow_{\pi[1..\text{first}(\pi)-1]} a'$. We have $a' = a$ by definition of $\text{first}(\cdot)$. Moreover, $a' + \beta \cdot \Delta(\pi_{\text{first}(\pi)}) > a' = a = \sup \tau$ for any $\beta \in (0, 1]$. Since exceeding $\sup \tau$ is forbidden, we obtain the contradiction $a \notin \text{enab}(\pi[1..\text{first}(\pi)]) \supseteq \text{enab}(\pi)$. \square

Corollary 5. *Given $a \in \mathbb{Z}$ and $p, q \in Q$, deciding whether $a \in \text{enab}(\text{Paths}_{p,q})$, or equivalently $\text{Post}_{p,q}(a) \neq \emptyset$, is in NC^2 .*

Proof. We report “empty” if $a \notin \tau$. Otherwise, let

$$\begin{aligned} S_0 &:= \{\pi \in \text{Paths}_{p,q} \mid \Delta^+(\pi) = \Delta^-(\pi) = 0\}, \\ S_+ &:= \{\pi \in \text{Paths}_{p,q} \mid \text{first}(\pi) \neq \infty \implies \Delta(\pi_{\text{first}(\pi)}) > 0\}, \\ S_- &:= \{\pi \in \text{Paths}_{p,q} \mid \text{first}(\pi) \neq \infty \implies \Delta(\pi_{\text{first}(\pi)}) < 0\}. \end{aligned}$$

By Lemma 4, it suffices if one of the following holds:

- (a) $a \notin \{\inf \tau, \sup \tau\}$ and $\text{Paths}_{p,q} \neq \emptyset$;
- (b) $a = \inf \tau = \sup \tau$ and $S_0 \neq \emptyset$;
- (c) $a = \inf \tau < \sup \tau$ and $S_+ \neq \emptyset$;
- (d) $a = \sup \tau > \inf \tau$ and $S_- \neq \emptyset$.

All of the above can be checked in NC^2 by Lemma 3. \square

B. Characterisation of reachability sets

As a step towards computing a representation of $\text{Post}_{p,q}(a)$, we characterise $\text{Post}_{p,q}(a)$ in terms of its closure. To this end, we note that admissible runs remain admissible whenever they are scaled down. Consequently, $\overline{\text{Post}_{p,q}(a)}$ is a closed interval that differs from $\text{Post}_{p,q}(a)$ in at most three points.

Proposition 6 (Adapted from [17, Lemma 4.2(c)]). *Let $\beta \in (0, 1]$ and let ρ be an admissible run from configuration $p(a)$. It is the case that run $\beta\rho$ is also admissible from $p(a)$.*

Lemma 7. *For every $b \in \overline{\text{Post}_{p,q}(a)}$, it is the case that $(a, b) \subseteq \text{Post}_{p,q}(a)$ and $(b, a) \subseteq \text{Post}_{p,q}(a)$.*

The above lemma holds as a run from a to b can be scaled to reach an arbitrary value from (a, b) and remain admissible.

Corollary 8. *Set $\overline{\text{Post}_{p,q}(a)}$ is a closed interval. Moreover, $\overline{\text{Post}_{p,q}(a)} \setminus \text{Post}_{p,q}(a) \subseteq \{\inf \overline{\text{Post}_{p,q}(a)}, a, \sup \overline{\text{Post}_{p,q}(a)}\}$.*

Proof. Let $b := \inf \overline{\text{Post}_{p,q}(a)}$ and $c := \sup \overline{\text{Post}_{p,q}(a)}$. For the sake of contradiction, suppose there is some $v \in \overline{\text{Post}_{p,q}(a)} \setminus \text{Post}_{p,q}(a)$ such that $v \notin \{b, a, c\}$. By Lemma 7, we have $(a, b) \cup (a, c) \cup (b, a) \cup (c, a) \subseteq \text{Post}_{p,q}(a) \subseteq \overline{\text{Post}_{p,q}(a)}$. Since $v \in (b, c) \setminus \{a\}$, we obtain $v \in \text{Post}_{p,q}(a)$, which is a contradiction. \square

C. Identifying the endpoints

We now show that a representation of the interval $\overline{\text{Post}_{p,q}(a)}$ can be obtained by identifying its endpoints in NC^2 . Some simple observations follow from [Proposition 6](#) and [Lemma 7](#):

Proposition 9. *The following statements hold:*

- (a) *If $\text{Post}_{p,q}(\inf \tau) \neq \emptyset$, then $\inf \overline{\text{Post}_{p,q}(\inf \tau)} = \inf \tau$.*
- (b) *If $\text{Post}_{p,q}(\sup \tau) \neq \emptyset$, then $\sup \overline{\text{Post}_{p,q}(\sup \tau)} = \sup \tau$.*
- (c) *Let $v \in \tau \setminus \{\inf \tau, \sup \tau\}$ and let ρ be a run. There exists $\varepsilon \in (0, 1]$ such that for all $\beta \in (0, \varepsilon]$ there exists $v_\beta > 0$ such that $v \rightarrow_{\beta\rho} v_\beta$. Moreover, $\lim_{\beta \rightarrow 0} v_\beta = v$.*

The two forthcoming lemmas characterise the endpoints of $\overline{\text{Post}_{p,q}(a)}$ through so-called admissible cycles. We say that a cycle θ is (a, p, q) -admissible if its first transition t satisfies $\Delta(t) \neq 0$, $a \in \text{enab}(\text{Paths}_{p,\text{in}(t)})$ and $\text{Paths}_{\text{in}(t),q} \neq \emptyset$. We say that such an admissible cycle is *positive* if $\Delta(t) > 0$, and *negative* if $\Delta(t) < 0$. Such cycles can be iterated to approach the endpoints of τ , by scaling all transitions but t arbitrarily close to zero.

Lemma 10. *If $\text{Post}_{p,q}(a) \neq \emptyset$ and \mathcal{V} has an (a, p, q) -admissible cycle θ , then the following holds:*

- (a) *$\inf \overline{\text{Post}_{p,q}(a)} = \inf \tau$, if θ is negative;*
- (b) *$\sup \overline{\text{Post}_{p,q}(a)} = \sup \tau$, if θ is positive.*

Lemma 11. *Let $\text{Post}_{p,q}(a) \neq \emptyset$, $b := \inf \overline{\text{Post}_{p,q}(a)}$ and $c := \sup \overline{\text{Post}_{p,q}(a)}$. If \mathcal{V} has no (a, p, q) -admissible cycle which is:*

- (a) *negative, then $b \neq -\infty$ and $b = \max(\inf \tau, a + \min\{\Delta^-(\pi) \mid \pi \in \text{Paths}_{p,q}, a \in \text{enab}(\pi)\})$;*
- (b) *positive, then $c \neq +\infty$ and $c = \min(\sup \tau, a + \max\{\Delta^+(\pi) \mid \pi \in \text{Paths}_{p,q}, a \in \text{enab}(\pi)\})$.*

Proof. We only prove (b) as (a) is symmetric. Assume \mathcal{V} has no positive (a, p, q) -admissible cycle. Let $D^+ := \{\Delta^+(\pi) \mid \pi \in \text{Paths}_{p,q}, a \in \text{enab}(\pi)\}$. We show that $\max D^+$ is well-defined. For the sake of contradiction, suppose that D^+ is infinite. By a pigeonhole argument, we obtain a run ρ admissible from a and such that ρ contains at least two occurrences of a transition t with $\Delta(t) > 0$. Let $\text{path}(\rho) = \pi t \pi' t \pi''$ where π, π', π'' are paths. The cycle $\theta := t \pi'$ is a positive admissible cycle, which yields a contradiction.

Note that $c \leq \min(\sup \tau, a + \max D^+)$, so $c \neq +\infty$. It remains to show that $c = \min(\sup \tau, a + \max D^+)$. Let $\pi \in \text{Paths}_{p,q}$ be such that $a \in \text{enab}(\pi)$ and $\Delta^+(\pi) = \max D^+$. By definition, there exists a run $\rho = \alpha_1 t_1 \cdots \alpha_n t_n$ admissible from $p(a)$ and such that $\text{path}(\rho) = \pi$. Since $a \in \tau$, there exists $\lambda \in (0, 1]$ such that $a + \lambda \cdot \max D^+ = \min(\sup \tau, a + \max D^+)$. For all $\varepsilon \in (0, 1)$, let $\rho_\varepsilon := \alpha'_1 t_1 \cdots \alpha'_n t_n$ be the run such that

$$\alpha'_i := \begin{cases} (1 - \varepsilon) \cdot \lambda & \text{if } \Delta(t_i) \geq 0, \\ \varepsilon \cdot (1/|\Delta(t_i)|) \cdot (1/n) & \text{otherwise.} \end{cases}$$

Informally, if were allowed to scale transitions by 0, then we would be done by using ρ_0 from a , as it would never decrease and reach exactly $a + \lambda \cdot \max D^+ = \min(\sup \tau, a + \max D^+)$.

Formally, we choose a small $\varepsilon \in (0, 1]$ as follows. If $a > \inf \tau$, then we pick ε so that $a - \varepsilon \geq \inf \tau$. Otherwise, we pick ε so that $(1 - \varepsilon) \cdot \lambda \geq \varepsilon$. We claim that the run ρ_δ is admissible from a for every $\delta \in (0, \varepsilon]$. First note that the top guard is never exceeded since $a + \Delta^+(\rho_\delta) = a + (1 - \delta) \cdot \lambda \cdot \max D^+ \leq a + \lambda \cdot \max D^+ \leq \sup \tau$. Let us now consider the bottom guard.

If $a > \inf \tau$, then $a + \Delta^-(\rho_\delta) \geq a - \delta \geq a - \varepsilon \geq \inf \tau$. Otherwise, if $a = \inf \tau$, then either $\Delta^-(\rho_\delta) = \Delta^+(\rho_\delta) = 0$, in which case admissibility is trivial, or the first transition t_i such that $\Delta(t_i) \neq 0$ is such that $\Delta(t_i) \geq 1$. In that case, the following holds for every $j \geq i$:

$$\begin{aligned} a + \Delta(\rho_\delta[1..j]) &\geq a + (1 - \delta) \cdot \lambda \cdot \Delta(t_i) + \Delta^-(\rho_\delta[i + 1..j]) \\ &\geq a + (1 - \varepsilon) \cdot \lambda + \Delta^-(\rho_\delta[i + 1..j]) \\ &\geq a + (1 - \varepsilon) \cdot \lambda - \varepsilon \geq a = \inf \tau. \end{aligned}$$

This shows the admissibility of ρ_δ . Thus, for all $\delta \in (0, \varepsilon]$, we have $a \rightarrow_{\rho_\delta} a_\delta$ where $a_\delta \geq \min(\sup \tau, a + \max D^+) - \delta \cdot \lambda \cdot \max D^+$. We are done since $\lim_{\delta \rightarrow 0} a_\delta = \min(\sup \tau, a + \max D^+)$. \square

In the forthcoming propositions, we show how the previous characterisations can be turned into NC^2 procedures.

Lemma 12. *On input $a \in \mathbb{Z}$ and $p, q \in Q$, deciding if \mathcal{V} has a positive or negative (a, p, q) -admissible cycle is in NC^2 .*

Proof. We consider the positive case; the negative one is symmetric. Testing whether there is a positive (a, p, q) -admissible cycle beginning with a transition t with $\Delta(t) > 0$ amounts to testing whether (i) $a \in \text{enab}(\text{Paths}_{p,\text{in}(t)})$, (ii) $\text{Paths}_{\text{in}(t),q} \neq \emptyset$, and (iii) $\text{Paths}_{\text{out}(t),\text{in}(t)} \neq \emptyset$. Condition (i) can be checked in NC^2 by [Corollary 5](#). Conditions (ii) and (iii) are graph reachability

queries which can be tested in $\mathbf{NL} \subseteq \mathbf{NC}^2$. There are at most $|T|$ transitions with positive effect, so the conditions can be tested in parallel for each $t \in T$. \square

Proposition 13. *On input $a \in \mathbb{Z}$ and states p, q , the values $\inf \overline{\text{Post}_{p,q}(a)}$ and $\sup \overline{\text{Post}_{p,q}(a)}$ can be computed in \mathbf{NC}^2 .*

Proof. We explain how to compute $c := \sup \overline{\text{Post}_{p,q}(a)}$ in \mathbf{NC}^2 . The procedure for $\inf \overline{\text{Post}_{p,q}(a)}$ is symmetric. By [Corollary 5](#), testing whether $\text{Post}_{p,q}(a) = \emptyset$ is in \mathbf{NC}^2 . If it holds, then trivially $c = -\infty$. Otherwise, assume that $\text{Post}_{p,q}(a) \neq \emptyset$, and hence $a \in \tau$. Additionally, if $a = \sup \tau$, then $c = \sup \tau$ by [Proposition 9\(b\)](#). So we assume $a < \sup \tau$.

By [Lemma 12](#), it can be decided in \mathbf{NC}^2 whether there exists a positive (a, p, q) -admissible cycle. If such a cycle exists, then $c = \sup \tau$ by [Lemma 10](#). Otherwise, by [Lemma 11](#), we have $c = \min(\sup \tau, a + \max D^+)$ where $D^+ := \{\Delta^+(\pi) \mid \pi \in \text{Paths}_{p,q}, a \in \text{enab}(\pi)\}$.

Consider a path $\pi \in \text{Paths}_{p,q}$ that satisfies $a \in \text{enab}(\pi)$ and which can be decomposed as $\pi = \sigma\theta\sigma'$ where θ is a cycle. As θ is (a, p, q) -admissible by definition, it cannot contain a positive transition. Otherwise, \mathcal{V} would admit a positive (a, p, q) -admissible cycle, which is a contradiction. Hence $\Delta^+(\pi) = \Delta^+(\sigma\sigma')$. We show that $a \in \text{enab}(\sigma\sigma')$. Recall that $a \neq \sup \tau$. Hence $a \in \text{enab}(\pi)$ follows from [Lemma 4\(a\)](#) or [\(c\)](#). Note that neither condition can be violated by removing a nonpositive transition from π , so $a \in \text{enab}(\sigma\sigma')$.

The above shows that there is a simple path π_{max} such that $\max D^+ = \Delta^+(\pi_{max})$ and $a \in \text{enab}(\pi_{max})$. Therefore, to obtain $\max D^+$, it is sufficient to compute $\max E^+$ where $E^+ := \{\Delta^+(\pi) \mid \pi \in \text{Paths}_{p,q}, a \in \text{enab}(\pi), |\pi| \leq |Q|\}$.

Now, let us make a case distinction based on whether $a = \inf \tau$. Assume this is true. By [Lemma 4](#), $\max E^+$ equals $\max(\$

$$\begin{aligned} & \max\{\Delta^+(\pi) \mid \pi \in \text{Paths}_{p,q}, \text{first}(\pi) = \infty, |\pi| \leq |Q|\}, \\ & \max\{\Delta^+(\pi) \mid \pi \in \text{Paths}_{p,q}, \Delta(\pi_{\text{first}(\pi)}) > 0, |\pi| \leq |Q|\} \end{aligned}$$

$\left. \right)$. By [Lemma 3](#), the above is the maximum of two values that can be computed in \mathbf{NC}^2 . If $a \neq \inf \tau$, then we have $a \in \tau \setminus \{\inf \tau, \sup \tau\}$, so by [Lemma 4\(a\)](#) all paths are admissible from a , and hence $\max E^+ = \max\{\Delta^+(\pi) \mid \pi \in \text{Paths}_{p,q} \wedge |\pi| \leq |Q|\}$. This value can be computed in \mathbf{NC}^2 by [Lemma 3](#). \square

D. Computing the representation

To obtain a representation of $\text{Post}_{p,q}(a)$, it remains to explain how to check in \mathbf{NC}^2 which of the three limit elements $\inf \overline{\text{Post}_{p,q}(a)}$, $\sup \overline{\text{Post}_{p,q}(a)}$ and a belong to $\text{Post}_{p,q}(a)$.

Proposition 14. *Testing whether $a \in \text{Post}_{p,q}(a)$ is in \mathbf{NC}^2 .*

Proof. By [Corollary 5](#), $\text{Post}_{p,q}(a) \neq \emptyset$ can be tested in \mathbf{NC}^2 . Thus, we assume that it is nonempty. It is easy to show that $a \in \text{Post}_{p,q}(a)$ iff at least one of these conditions holds:

- (a) there exists a path $\pi \in \text{Paths}_{p,q}$ whose transitions are all zero, i.e. $\Delta(\pi) = \Delta^+(\pi) = \Delta^-(\pi) = 0$;
- (b) there exist $\pi \in \text{Paths}_{p,q}$ and i, j such that $\Delta(\pi_i) > 0$ and $\Delta(\pi_j) < 0$. If $a = \inf \tau$, then we also require $\Delta(\pi_k) = 0$ for all $k < i$ and $k > j$. Similarly, if $a = \sup \tau$, then we also require $\Delta(\pi_k) = 0$ for all $k < j$ and $k > i$.

It remains to argue that both conditions can be checked in \mathbf{NC}^2 . We can check condition [\(a\)](#) in \mathbf{NC}^2 via [Lemma 3\(b\)](#). If $a \notin \{\inf \tau, \sup \tau\}$, then condition [\(b\)](#) can be checked in \mathbf{NC}^2 via [Lemma 3\(a\)](#). If $a \in \{\inf \tau, \sup \tau\}$, then condition [\(b\)](#) can be checked in \mathbf{NC}^2 via [Lemma 3\(d\)](#) and [Lemma 3\(c\)](#). \square

Proposition 15. *On input $a \in \mathbb{Z}$ and $p, q \in Q$, computing $\text{Post}_{p,q}(a) \cap \{\inf \overline{\text{Post}_{p,q}(a)}, \sup \overline{\text{Post}_{p,q}(a)}\}$ is in \mathbf{NC}^2 .*

Proof. By [Corollary 5](#), we can check if $\text{Post}_{p,q}(a) \neq \emptyset$ in \mathbf{NC}^2 . Thus, we assume it is nonempty. Let $b := \inf \overline{\text{Post}_{p,q}(a)}$ and $c := \sup \overline{\text{Post}_{p,q}(a)}$, which can be computed in \mathbf{NC}^2 by [Proposition 13](#). We check whether $b = c$. If it is, we return $\{b, c\}$ since $\text{Post}_{p,q}(a) \neq \emptyset$. Otherwise, we explain how to check whether $c \in \text{Post}_{p,q}(a)$; the case of b can be handled symmetrically. We assume that $b < a < c$, as [Proposition 14](#) handles the case $a \in \{b, c\}$ when checking membership of a .

If $c \notin \tau$, then $c \notin \text{Post}_{p,q}(a)$. Otherwise, it can be shown that $c \in \text{Post}_{p,q}(a)$ iff there is a state $r \in Q$ and a path $\sigma \in \text{Paths}_{r,q}$ that satisfy $\Delta^+(\sigma) > 0$, $\Delta^-(\sigma) = 0$ and either of the following holds:

- (i) there exists a path $\sigma' \in \text{Paths}_{p,r}$ such that $|\sigma|, |\sigma'| \leq |Q|$, $\Delta^-(\sigma') = 0$ and $\Delta^+(\pi) \geq c - a$ where $\pi := \sigma'\sigma$;
- (ii) there exists a path $\sigma' \in \text{Paths}_{p,r}$ such that $|\sigma|, |\sigma'| \leq |Q|$ and $\Delta^+(\pi) > c - a$ where $\pi := \sigma'\sigma$;
- (iii) there is a positive (a, p, r) -admissible cycle θ .

It remains to show that the conditions can be tested in \mathbf{NC}^2 . There are $|Q|$ choices for state r , so we can test the conditions for all choices in parallel. Let $S := \{\sigma \in \text{Paths}_{r,q} \mid \Delta^+(\sigma) > 0 \text{ and } \Delta^-(\sigma) = 0\}$. We first check whether $S \neq \emptyset$, which can be done in \mathbf{NC}^2 by [Lemma 3\(a\)–\(b\)](#). Moreover, condition [\(iii\)](#) can be checked in \mathbf{NC}^2 by [Lemma 12](#). We proceed as follows to check [\(i\)](#). Let

$$\begin{aligned} W & := \{\Delta^+(\sigma) \mid \sigma \in S, |\sigma| \leq |Q|\}, \\ W' & := \{\Delta^+(\sigma') \mid \sigma' \in \text{Paths}_{p,r}, |\sigma'| \leq |Q|, \Delta^-(\sigma') = 0\}. \end{aligned}$$

By Lemma 3, we can compute $m := \max W + \max W'$ in NC^2 and check that $m \geq c - a$. Lastly, we define $W'' := \{\Delta^+(\sigma') \mid \sigma' \in \text{Paths}_{p,r}, |\sigma'| \leq |Q|\}$ and test whether $\max W + \max W'' > c - a$ to verify condition (ii). \square

Theorem 16. *Given $a, a' \in \mathbb{Z}$ and $p, q \in Q$, the following can be done in NC^2 : obtaining a representation of $\text{Post}_{p,q}(a)$ and testing whether $a' \in \text{Post}_{p,q}(a)$.*

Proof. By Proposition 13, we can compute $b := \inf \overline{\text{Post}_{p,q}(a)}$ and $c := \sup \overline{\text{Post}_{p,q}(a)}$ in NC^2 . By Corollary 8, Proposition 14 and Proposition 15, the set $S := \overline{\text{Post}_{p,q}(a)} \setminus \text{Post}_{p,q}(a)$, of size at most three, can be computed in NC^2 . By Corollary 8, this yields the representation $\text{Post}_{p,q}(a) = [b, c] \setminus S$. Thus, $a' \in \text{Post}_{p,q}(a)$ iff $b \leq a' \leq c$ and $a' \notin S$. \square

E. Equality tests

A COCA with equality tests is a tuple $\mathcal{V} = (Q, T, \tau, \phi)$, where (Q, T, τ) is a COCA and $\phi: Q \rightarrow \{[z, z] \mid z \in \mathbb{Z}\} \cup \mathbb{Q}$. We say that a run of \mathcal{V} is *admissible* if each of its configurations $q(a)$ satisfies $a \in \tau \cap \phi(q)$.

Using the previous results, we can extend the NC^2 membership of the reachability problem $p(a) \rightarrow_* q(b)$ to COCA with equality tests. The proof relies on the fact that each equality test is passed by exactly one configuration. For this reason, we can construct a reachability graph between equality tests using a quadratic number of COCA reachability queries.

Let us assume that p has no incoming edges; if it does, we can simply add a new initial state p' and a single transition $(p', 0, p)$. Similarly, we can assume q has no outgoing edges.

We will reason about reachability in \mathcal{V} where we avoid all equality tests. For every states $p', q' \in Q$, let us define the COCA $\mathcal{V}_{p',q'} := (Q_{p',q'}, T_{p',q'})$ where $Q_{p',q'} := \{s \in Q \mid \phi(s) = \mathbb{Q}\} \cup \{p', q'\}$. We treat p' as a dedicated input state, and q' as a dedicated output state. That is,

$$T_{p',q'} := \{t \in T \mid \text{in}(t) \in Q_{p',q'} \setminus \{q'\}, \text{out}(t) \in Q_{p',q'} \setminus \{p'\}\}.$$

Let us define a graph $\mathcal{G} := (V, E)$, where $V := \{p(a), q(b)\} \cup \{r(z) \mid r \in Q, \phi(r) = [z, z], z \in \tau\}$. If $a \notin \tau \cap \phi(p)$ or $b \notin \tau \cap \phi(q)$, then we trivially conclude that $p(a)$ cannot reach $q(b)$. Hence, $|V| \leq |Q|$ holds. Intuitively, the nodes of \mathcal{G} correspond to the initial and final configurations, plus, for each equality test, the configuration that passes this test. Let us define $E := \{(p'(x), q'(y)) \mid p'(x) \rightarrow_* q'(y) \text{ in } \mathcal{V}_{p',q'}\}$.

Lemma 17. *It is the case that $p(a) \rightarrow_* q(b)$ in \mathcal{V} if and only if there is a path from $p(a)$ to $q(b)$ in \mathcal{G} .*

An edge between two nodes $p'(a')$ and $q'(b')$ in \mathcal{G} implies reachability between $p'(a')$ and $q'(b')$ in $\mathcal{V}_{p',q'}$, from which reachability between $p'(a')$ and $q'(b')$ in \mathcal{V} follows. Hence the “if” direction follows by the fact that reachability is transitive.

Further, a run from $p(a)$ to $q(b)$ in \mathcal{V} can be split into runs that each witness reachability between two equality tests (or the initial/target configuration), but avoid all equality tests in intermediate configurations. Therefore, these runs are also runs in $\mathcal{V}_{p',q'}$ for some p', q' , so the corresponding edges exist in \mathcal{G} . Hence the “only if” direction follows.

Theorem 18. *The reachability problem for COCA with equality tests is in NC^2 .*

Proof. Let us first argue that the graph \mathcal{G} can be constructed in NC^2 . There are at most $|Q|$ nodes in \mathcal{G} , and hence at most $|Q|^2$ edges. Note that an edge $(p'(x), q'(y))$ is present iff $p'(x) \rightarrow_* q'(y)$ in $\mathcal{V}_{p',q'}$, which can be decided in NC^2 by Theorem 16, as $\mathcal{V}_{p',q'}$ is a COCA. By running these $|Q|^2$ queries in parallel, it follows that \mathcal{G} can be obtained in NC^2 .

Once the graph \mathcal{G} has been constructed, by Lemma 17, it suffices to test reachability from $p(a)$ to $q(b)$ in \mathcal{G} . Since graph reachability is in $\text{NL} \subseteq \text{NC}^2$, we are done. \square

IV. GUARDED COCA REACHABILITY

We now turn to the reachability problem $p(a) \rightarrow_\rho q(b)$ for a guarded COCA $\mathcal{W} = (Q, T, \tau)$.

In contrast to COCAs, for guarded COCAs $\text{Post}_{p,q}(a)$ does not necessarily admit a decomposition into a constant number of intervals. Nevertheless, we show that it can always be decomposed into a linear number of intervals with respect to the number of states. Throughout this section, we write $\mathcal{I}(R)$ to denote the unique decomposition of a set $R \subseteq \mathbb{Q}$ into maximal disjoint nonempty intervals. For example, $\mathcal{I}([3, 4] \cup (4, 5) \cup (5, +\infty)) = \{[3, 5), (5, +\infty)\}$ and $\mathcal{I}(\emptyset) = \emptyset$.

A. Controlling the number of intervals

We will prove that, for every k , the set $\{b \in \mathbb{Q} \mid p(a) \rightarrow_\rho q(b), |\rho| = k\}$ decomposes into a polynomial number of intervals. To do so, we will bound the size of the decomposition of sets obtained by updating $A = [a, a]$ with operations which suffice to implement continuous counter updates and guard tests. More precisely, the following operations: Minkowski sums, intersections (with elements of $\mathcal{L} = \{\tau(q) \mid q \in Q\}$), and unions (with sets constructed similarly). For technical reasons, we also consider a fourth operation.

Let us fix a bounded interval $A \in \mathcal{I}$ and $\mathcal{L} \subseteq \mathcal{I}$. We write $P_{\mathcal{L}} := \{\inf I, \sup I \mid I \in \mathcal{L}\}$ to denote the set of endpoints within \mathcal{L} . We define the *MIUN-closure* (short for Minkowski sum, Intersection, Union, and New), of interval A w.r.t. \mathcal{L} , as the smallest collection $\mathcal{C} \subseteq 2^{\mathbb{Q}}$ such that $A \in \mathcal{C}$ and:

- M: if $B \in \mathcal{C}$ and $z \in \mathbb{Q}_{>0}$ then $B + (0, z], B + [-z, 0) \in \mathcal{C}$;
- I: if $B \in \mathcal{C}$ and $L \in \mathcal{L}$ then $B \cap L \in \mathcal{C}$;
- U: if $B, B' \in \mathcal{C}$ then $B \cup B' \in \mathcal{C}$;
- N: if $B \in \mathcal{C}$ and $I \in \mathcal{I}$ s.t. $\bar{I} \cap P_{\mathcal{L}} \neq \emptyset$ then $B \cup I \in \mathcal{C}$.

The forthcoming lemma forms the basis of our bound. It is based on so-called *indicator functions* which give us, for every interval I , the set of endpoints of \mathcal{L} and A that belong to the closure of I . As we will see later, the set of endpoints needed to analyse COCA is small. Furthermore, all MIUN-operations are such that sets of \mathcal{C} decompose into intervals whose closure contains at least one such endpoint.

More formally, for all $B \in \mathcal{C}$, let $\phi_B: \mathcal{I}(B) \rightarrow 2^{P_{\mathcal{L}} \cup P_A}$ be the function defined as $\phi_B(I) := \bar{I} \cap (P_{\mathcal{L}} \cup P_A)$.

Lemma 19. *We have $\phi_B(I) \neq \emptyset$ for all $B \in \mathcal{C}$ and $I \in \mathcal{I}(B)$.*

Proof. We proceed by induction on the definition of MIUN-closures. We define $\mathcal{C}_0 := \{A\}$ and \mathcal{C}_{i+1} as \mathcal{C}_i extended with all sets obtained by applying the MIUN-operations applied to any $B, B' \in \mathcal{C}_i$. We will show that the lemma holds for all \mathcal{C}_i , which will conclude the proof since $\mathcal{C} = \bigcup_{i \in \mathbb{N}} \mathcal{C}_i$.

We have $\mathcal{I}(A) = \{A\}$ and the claim holds since $P_A \subseteq \phi_A(A)$. For the induction step, we suppose the claim holds for \mathcal{C}_i . We have to prove that for all $C \in \mathcal{C}_{i+1}$ and all $I \in \mathcal{I}(C)$ it holds that $\phi_C(I) \neq \emptyset$. Notice that this is trivial if C is obtained from $B \in \mathcal{C}_i$ by application of the New operation.

First, we consider the Minkowski sum. Consider some $B \in \mathcal{C}_i$ with the function ϕ_B and let $I \in \{(0, z], [-z, 0)\}$ for some $z \in \mathbb{Q}_{>0}$. Let $C := B + I$. For all $J \in \mathcal{I}(C)$ there exists $K_B \in \mathcal{I}(B)$ such that $K_B \subseteq J$. Thus, $\phi_B(K_B) \subseteq \phi_C(J)$ and the claim holds by the inductive hypothesis for ϕ_B .

Second, we consider intersections. We only deal with intervals of the form $[\ell, +\infty)$, $(\ell, +\infty)$, $(-\infty, \ell]$, or $(-\infty, \ell]$, since intersection with any interval can be expressed by at most two consecutive intersections with intervals of this form. Let $B \in \mathcal{C}_i$ and $L \in \mathcal{L}$. Suppose that $\bar{L} = [\ell, +\infty)$ and let $C := B \cap L$. Recall that $\ell \in P_{\mathcal{L}}$. Observe that $\mathcal{I}(B)$ contains at most one interval I such that $\ell \in \bar{I}$. If such an I exists, then $\ell \in \phi_C(\bar{I} \cap \bar{L})$. For all other intervals $J \in \mathcal{I}(B)$, we have that $J \cap L$ is either J or \emptyset . If the intersection is nonempty, then $\phi_C(J) = \phi_B(J)$ and the claim holds by inductive hypothesis. If \bar{L} is instead of the form $(-\infty, \ell]$, then we proceed similarly.

Finally, we consider unions. Let $B, B' \in \mathcal{C}_i$ and $I \in \mathcal{I}(B \cup B')$. By definition, there exists $J \in \mathcal{I}(B) \cup \mathcal{I}(B')$ with $J \subseteq I$. Therefore, either $\phi_B(J)$ or $\phi_{B'}(J)$ is nonempty and contained in $\phi_{B \cup B'}(I)$. \square

Lemma 20. *For every set $B \subseteq \mathbb{Q}$ and every pairwise distinct intervals $I_1, I_2, I_3 \in \mathcal{I}(B)$, it is the case that $\bar{I}_1 \cap \bar{I}_2 \cap \bar{I}_3 = \emptyset$.*

A point can belong to at most one interval among disjoint intervals. Moreover, a point can belong to at most two closures, e.g. consider $[0, 1)$ and $(1, 2]$. This is no longer possible for three intervals due to maximality of intervals in $\mathcal{I}(B)$. Thus, the proof of Lemma 20 follows from a simple case analysis.

Now, we show that if \mathcal{L} is finite, then there is a polynomial bound on the number of intervals within the decomposition of any set from the MIUN-closure \mathcal{C} . More formally:

Lemma 21. *If \mathcal{L} is finite, then $\mathcal{I}(B)$ consists of at most $4(|\mathcal{L}| + 1)$ intervals, for every $B \in \mathcal{C}$*

Proof. By Lemma 20, there are at most two pairwise disjoint intervals that share a point in their closure. By Lemma 19, the indicator function guarantees that $\bar{J} \cap (P_{\mathcal{L}} \cup P_A) \neq \emptyset$ for all $J \in \mathcal{I}(B)$. Thus, $\mathcal{I}(B)$ has at most $2(2|\mathcal{L}| + 2)$ intervals. Otherwise, by the pigeonhole principle, a point of $P_{\mathcal{L}} \cup P_A$ would belong to at least three closures of intervals from $\mathcal{I}(B)$. \square

B. Approximations of the reachability function

It will be convenient to manipulate mappings from states to (under-approximations of) their reachability functions. We consider the mappings $\mathcal{R}_Q := \{R: Q \rightarrow 2^Q\}$. An example of such a mapping is $\text{Reach}_{p(a)}$ defined as $\text{Reach}_{p(a)}(q) := \text{Post}_{p,q}(a)$. Given $R, R' \in \mathcal{R}_Q$, we write $R \preceq R'$ iff $R(q) \subseteq R'(q)$ for all $q \in Q$. We seek to define a sequence of mappings $R_0 \preceq R_1 \preceq \dots$ such that $R_n = \text{Reach}_{p(a)}$ for some $n \in \mathbb{N}$.

For all state $q \in Q$, we define the *successor mapping-update function* $\text{Succ}_q: \mathcal{R}_Q \rightarrow 2^Q$ as follows:

$$\begin{aligned} \text{Succ}_q(R) := & R(q) \\ & \cup \bigcup \{(R(r) + (0, z]) \cap \tau(q) \mid (r, z, q) \in T, z > 0\} \\ & \cup \bigcup \{(R(r) + [z, 0)) \cap \tau(q) \mid (r, z, q) \in T, z < 0\} \\ & \cup \bigcup \{R(r) \cap \tau(q) \mid (r, 0, q) \in T\}. \end{aligned}$$

Let $\text{Succ}: \mathcal{R}_Q \rightarrow \mathcal{R}_Q$ be defined as $\text{Succ}(R)(q) := \text{Succ}_q(R)$. Below, we state the key property enjoyed by Succ . In words, its i -fold composition coincides with the set of configurations reachable via runs of length at most i . It can be easily proved by induction on the definition of Succ .

Lemma 22. Let $S_0 \in \mathcal{R}_Q$ and $S_i := \text{Succ}(S_{i-1})$ for all $i \geq 1$. The following holds:

$$S_i(q) = \bigcup_{p \in Q} \{b \in \mathbb{Q} \mid a \in S_0(p), p(a) \rightarrow_\rho q(b) \text{ and } |\rho| \leq i\}.$$

Now we can state a proposition that shows how the previous section relates to these definitions. Let us fix a configuration $p(a)$. We will focus on the MIUN-closure \mathcal{C} of $A := [a, a]$ with respect to $\mathcal{L} := \{\tau(q) \mid q \in Q\}$. We say that a mapping $S \in \mathcal{R}_Q$ is \mathcal{C} -valid if $S(q) \in \mathcal{C}$ for all $q \in Q$.

Proposition 23. Let $S \in \mathcal{R}_Q$ be a \mathcal{C} -valid mapping. We have $S \preceq \text{Succ}(S)$ and $\text{Succ}(S)$ is a \mathcal{C} -valid mapping.

Proof. We have $S \preceq \text{Succ}(S)$ directly from the definition of Succ . To prove that $\text{Succ}(S)(q) \in \mathcal{C}$, it suffices to observe that $\text{Succ}_q(S)$ is defined using Minkowski sums, intersections, and unions which are building blocks of MIUN-closures. \square

C. Accelerations

Unfortunately, applying Succ might not give us $\text{Reach}_{p(a)}$ in a small or even finite number of steps, e.g. if $\text{Reach}_{p(a)}(q)$ is unbounded for some $q \in Q$. We introduce another operation on mappings to resolve this. We start by defining some special form of cycles.

Let us fix a mapping $S_0 \in \mathcal{R}_Q$ and let $S_{i+1} := \text{Succ}(S_i)$ for every $i \geq 0$. We say that a run $\rho = \alpha_1 t_1 \cdots \alpha_n t_n$ is a *positively expanding cycle* from S_0 if it is admissible and there exist configurations $p_0(a_0), p_1(a_1), \dots, p_n(a_n)$ such that:

- (1) $p_0 = p_n$ and $\Delta(\rho) > 0$;
- (2) $a_0 \in S_0(p_0)$ and $p_0(a_0) \rightarrow_{\rho_i} p_i(a_i)$ for all $i \geq 1$; and
- (3) $a_i \in S_i(p_i) \setminus S_{i-1}(p_i)$ for all $i \geq 1$.

Moreover, letting I_0, \dots, I_n be the sequence of intervals such that $a_i \in I_i \in \mathcal{I}(S_i(p_i))$ for all $i \in \{0, \dots, n\}$, we require:

- (4) $I_0 \subseteq I_n$;
- (5) for all $i \geq 1$, there is a unique interval $I'_i \in \mathcal{I}(S_{i-1}(p_i))$ such that $I'_i \subseteq I_i$; and
- (6) $a_i \geq \sup(I'_i)$ for every $i \geq 1$.

Intuitively, the third condition states that each a_i is a “new” value, and the fifth and sixth conditions state that a_i expands some interval towards the top.

Similarly, we say that ρ is a *negatively expanding cycle* from S_0 if in the first item we replace $\Delta(\rho) > 0$ with $\Delta(\rho) < 0$; and in the last item we replace $a_i \geq \sup(I'_i)$ with $a_i \leq \inf(I'_i)$.

The following property follows from the definitions:

Lemma 24. It holds that $a_0, a_n \in I_n \subseteq \text{Reach}_{p_0(a_0)}(p_0)$.

It transpires that the Succ function always yields expanding cycles after a polynomial number of applications. The proof of this claim relies on our bounds for interval decompositions of sets from the MIUN-closure. The computational part is a simple backward construction of a run containing a cycle. A full proof is given in [subsection B](#) of the appendix.

Proposition 25. For some n polynomially bounded in $|Q|$, at least one of the following holds: $S_n = S_{n+1}$,

- there is a positively expanding cycle ρ from S_n , or
- there is a negatively expanding cycle ρ from S_n .

Moreover, it can be determined in time $|Q|^{\mathcal{O}(1)}$ whether the second or third case hold, and then ρ and its witnessing configurations can be computed in time $|Q|^{\mathcal{O}(1)}$.

We are ready to define the *acceleration* operation. Let ρ be a positively or negatively expanding cycle from $S \in \mathcal{R}_Q$ and let $p_0(a_0), p_1(a_1), \dots, p_n(a_n)$ be the configurations witnessing the run. Let I_0, \dots, I_n be the intervals given by the definition of expanding cycles. If ρ is positively expanding, then we define $\delta_i^+ := \sup \tau(p_i) - a_i$ for all $i \in \{1, \dots, n\}$. If ρ is a negatively expanding cycle, then we define $\delta_i^- := a_i - \inf \tau(p_i)$. Let $j \in \{1, \dots, n\}$ be such that:

$$\delta_j^+ = \min\{\delta_i^+ \mid 1 \leq i \leq n\} \text{ or } \delta_j^- = \min\{\delta_i^- \mid 1 \leq i \leq n\}.$$

We define Acc so that, given ρ and the mapping S , it outputs a new mapping $\text{Acc}(S, \rho) = S'$. If ρ is positively expanding from S then $S'(q) := S(q)$ for all $q \neq p_j$ and:

$$S'(p_j) := S(p_j) \cup \underbrace{I_j \cup (\tau(p_j) \cap [a_j, +\infty))}_{=K \subseteq \mathbb{Q}}.$$

Recall that $a_j \in I_j$, and $a_j \in \tau(p_j)$ since $a_0 \rightarrow_{\rho[1..j]} a_j$, so K is an interval. Also, since ρ is positively expanding and $j \geq 1$, we have $a_j \notin S(p_j)$ and $S'(p_j) \setminus S(p_j) \neq \emptyset$. Similarly, if ρ is negatively expanding then $S'(q) := S(q)$ for all $q \neq p_j$ and:

$$S'(p_j) := S(p_j) \cup I_j \cup ((-\infty, a_j] \cap \tau(p_j)).$$

Lemma 26. *Let $S \in \mathcal{R}_Q$ be a \mathcal{C} -valid mapping such that $S \preceq \text{Reach}_{p(a)}$ and let ρ be a positively or negatively expanding cycle from S . If $S' = \text{Acc}(S, \rho)$, then $S \preceq S'$, S' is a \mathcal{C} -valid mapping, and $S' \preceq \text{Reach}_{p(a)}$.*

Proof. We have $S \preceq S'$ directly from the definition of Acc . Similarly, S' is \mathcal{C} -valid because the operation to define $S'(p_j)$ is the ‘‘New’’ operation since the closure of the added interval always contains one of the endpoints from $\tau(p_j)$. It remains to prove that $S' \preceq \text{Reach}_{p(a)}$.

We assume that $\Delta(\rho) > 0$; the proof is similar for the other case. Let $\rho = \alpha_1 t_1 \cdots \alpha_n t_n$. Let $p_0(a_0), p_1(a_1), \dots, p_n(a_n)$ and I_0, \dots, I_n be the configurations and intervals given by the definition of positively expanding cycles. Let j be the index minimising δ_j^+ . We must prove that $S'(p_j) \preceq \text{Reach}_{p(a)}$. Since $S \preceq \text{Reach}_{p(a)}$ and $a_0 \in S(p_0)$ by definition of positively expanding cycles, it suffices to show that for every $b \in S'(p_j) \setminus S(p_j)$ there is an admissible run from $p_0(a_0)$ to $p_j(b)$.

If $\delta_j^+ = +\infty$, then $\sup \tau(p_i) = +\infty$ for all $i \in \{1, \dots, n\}$. Since $\Delta(\rho) > 0$, for all $\alpha, \beta \in (0, 1]$ and $m \in \mathbb{N}$ the run $\rho' := (\beta\rho)^m \alpha \rho[1..j]$ is admissible from any $p_n(a')$ with $a' \geq a_n$, to state p_j . Note that $\Delta(\rho') = m\beta\Delta(\rho) + \alpha\Delta(\rho[1..j])$, which can be any positive rational number by properly choosing α, β and m . Thus, $b \in \text{Reach}_{p(a)}(p_j)$ for every $b > a_n$.

It remains to consider the case $a_j \leq a_n$ to prove the claim for every $b \in [a_j, a_n]$. Let $\varepsilon \in (0, a_n - a_0]$. Since $a_0 < a_0 + \varepsilon \leq a_n$, we have $a_0 + \varepsilon \in I_n \subseteq \text{Reach}_{p(a)}(p_0)$ where the latter follows from Lemma 24. Recall that ρ' is admissible from all $p_n(a')$ with $a' \geq a_n$, and hence from $a_n + \varepsilon$. Thus, $p(a) \rightarrow_* p_0(a_0 + \varepsilon) \rightarrow_\rho p_n(a_n + \varepsilon)$, and $p_0(a_0 + \varepsilon) \rightarrow_{\rho[1..j]} p_j(a_j + \varepsilon)$. This shows that $b \in \text{Reach}_{p(a)}(p_j)$ for all $b \in [a_j, a_n]$.

Now, suppose $\delta_j^+ < +\infty$. If $a_j = \sup \tau(p_j)$, then we are done because $a_j \in I_j \subseteq \text{Reach}_{p(a)}(p_j)$. Otherwise, let $b \in [a_j, +\infty) \cap \tau(p_j)$. We need to prove that $b \in \text{Reach}_{p(a)}(p_j)$. Note that, by definition, we have $0 \leq b - a_j \leq \delta_j^+$.

Let $m \in \mathbb{N}$ and $c \in \mathbb{Q}_{\geq 0}$ be the unique numbers that satisfy $b - a_j = m \cdot \Delta(\rho) + c$ and $c < \Delta(\rho)$. Since $a_0 \leq a_0 + c \leq a_0 + \Delta(\rho) = a_n$, then by Lemma 24 we conclude that $a_0 + c \in \text{Reach}_{p(a)}(p_0)$. Notice that $a_j + c + m \cdot \Delta(\rho) = b$. It thus remains to prove that $p_0(a_0 + c) \rightarrow_{\rho^m \rho[1..j]} p_j(b)$. We prove something stronger, namely that $p_0(a_0 + c) \rightarrow_{\rho^{m+1}} p_n(a_n + b - a_j)$. Since $\Delta(\rho) > 0$, for the bottom guards it suffices to check whether the configurations are large enough when ρ is applied the first time. Indeed, since ρ is admissible from $p_0(a_0)$, we get $a_i + c + \Delta(\rho_i) \geq a_i + \Delta(\rho_i) \geq \inf \tau(q_i)$. Similarly, for the top guards, it suffices to check whether the configurations are small enough when ρ is applied last.

Indeed, since $b - a_j \leq \delta_j^+$, we have $a_i + c + m \cdot \Delta(\rho) = a_i + b - a_j \leq a_i + \delta_j^+ \leq a_i + \delta_i^+ = \sup \tau(p_i)$. If $\sup \tau(p_j) \notin \tau(p_j)$, then $b - a_j < \delta_j^+$ and the previous inequalities are strict. \square

D. Polynomial time algorithm

We summarise how to obtain the polynomial time algorithm for deciding $p(a) \rightarrow_* q(b)$. We begin with the mapping $R_0 \in \mathcal{R}_Q$ defined as $R_0(p) := [a, a]$ and $R_0(r) := \emptyset$ for every $r \neq p$. Clearly, $R_0 \preceq \text{Reach}_{p(a)}$. The next mappings R_1, R_2, \dots are defined as follows. Suppose we have defined R_0, \dots, R_i . Let $S_0^i := R_i$ and $S_{j+1}^i := \text{Succ}(S_j^i)$ for all $j \geq 0$.

By Proposition 25, we will either find an expanding cycle ρ from some S_n^j , where n is bounded polynomially, or we will find some $S_n^j = S_{n+1}^j$, again for n bounded polynomially. If there is an expanding cycle — a fact which, by Proposition 25, we can check in polynomial time — then we define $R_{i+j} := S_j^i$ for $1 \leq j < n$ and $R_{i+n} := \text{Acc}(R_{i+n-1}, \rho)$. Otherwise, we define $R_{i+j} := S_j^i$ for $j \in \{1, \dots, n\}$ and the algorithm returns R_{i+n} . By Lemma 22 and Lemma 26, we have $R_i \preceq \text{Reach}_{p(a)}$ for all defined R_i . Hence, if the algorithm terminates then by Lemma 22 it returns $\text{Reach}_{p(a)}$.

The rest of this section is devoted to proving that the above-described algorithm has a polynomial worst-case running time. It suffices to argue that expanding cycles can only be found some polynomial number of times.

Proposition 27. *The algorithm computes a representation of $\text{Reach}_{p(a)}$ in time $|Q|^{\mathcal{O}(1)}$.*

By Proposition 25, it suffices to show that Acc can be applied at most polynomially many times. To do so, we introduce the notion of progressing extensions. Let \mathcal{C} be the MIUN-closure of A w.r.t. \mathcal{L} , and let $B, B' \in \mathcal{C}$ be such that $B \subseteq B'$. We say that B' is a *progressing extension* of B if:

(1) there is $I' \in \mathcal{I}(B')$ such that $B \cap I' = \emptyset$;

or if there are $I \in \mathcal{I}(B)$ and $I' \in \mathcal{I}(B')$ such that $I \subseteq I'$ and at least one of the following holds:

(2) either $\phi_{B'}(I') \setminus \phi_B(I) \neq \emptyset$, or

(3) there exists $\ell \in \phi_B(I)$ such that $\ell \notin I$ and $\ell \in I'$.

See Figure 2 for a pictorial description of progressing extensions. Observe that in case (3) we necessarily have that $\ell \in \bar{I}$.

We show that accelerating leads to a progressing extension. Then, in the sequel, we provide a polynomial bound on the number of progressing extensions in a \subseteq -increasing sequence.

Lemma 28. *Let ρ be an expanding cycle from R and let $R' := \text{Acc}(R, \rho)$. There is some state p_j such that $R(p_j) \subseteq R'(p_j)$ is a progressing extension.*

Proof. Let $p_j \in Q$ be such that $R'(p_j) = R(p_j) \cup I_j \cup J$ for some interval J . In the proof, we write $\alpha_i t_i, p_i(a_i)$, and I_i , as in the definition of expanding cycles. We will assume that ρ is positively expanding; the other case is similar. Thus,




Fig. 2. Left: A set B such that $\mathcal{I}(B) = \{(-\infty, 3], [4, 5]\}$. Right: Example of the three possible types of progressing extensions of B . Dashed lines denote open interval borders; $\ell_1 = 4, \ell_2 = 5$ and $\ell_3 = 6$ denote values in $P_{\mathcal{L}} \cup P_A$.

$J \in \{[a_j, g], [a_j, g), [a_j, +\infty)\}$, where $g := \sup \tau(p_j)$. Recall that by definition $a_j \notin R(p_j)$ and $a_j \in I_j \in \mathcal{I}(\text{Succ}^k(R)(p_j))$ for some k . Thus, $I_j \cup J$ is an interval.

If $(I_j \cup J) \cap R(p_j) = \emptyset$, then $R'(p_j)$ is a progressing extension due to (1). For the remaining case, let $b \in (I_j \cup J) \cap R(p_j)$ and $K \in \mathcal{I}(R(p_j))$ be such that $b \in K$. Note that $K \cup I_j \cup J$ is an interval. If $b < a_j$, then, because $a_j \notin R(p_j)$, either $g \notin K$ or K has an upper bound if $J = [a_j, +\infty)$. Thus, $R'(p_j)$ is a progressing extension due to (2) or (3). Finally, suppose that $b > a_j$. By definition of ρ , there is a unique interval $I'_j \in \mathcal{I}(\text{Succ}^{k-1}(R)(p_j))$ such that $I'_j \subseteq I_j$. Moreover, $a_j \notin I'_j$ and $a_j \geq \sup I'_j$. Thus, $K \cap I'_j$ is empty and $g \notin I'_j$ or I'_j has an upper bound if $J = [a_j, +\infty)$. Since $K \cup I_j \cup J \in \mathcal{I}$, $R'(p_j)$ is a progressing extension due to (2) or (3). \square

To conclude, we show the following bound.

Lemma 29. *Let $B_0, B_1, B_2, \dots \in \mathcal{C}$ be a sequence such that $B_i \subseteq B_{i+1}$ for all $i \in \mathbb{N}$. The set of $i \in \mathbb{N}$ such that B_{i+1} is a progressing extension of B_i has cardinality at most $|\mathcal{L}|^{\mathcal{O}(1)}$.*

Proof. Let $P := P_{\mathcal{L}} \cup P_A$. First, observe that (3) can happen only if there exists some $\ell \in P$ such that $\ell \in B_{i+1} \setminus B_i$, and thus at most $|P|$ times.

Let $\phi_{B_i}(\mathcal{I}(B_i)) \subseteq P$ be the image of all intervals of $\mathcal{I}(B_i)$. Note that $\phi_{B_i}(\mathcal{I}(B_i)) \subseteq \phi_{B_{i+1}}(\mathcal{I}(B_{i+1}))$. A strict inclusion can happen at most $|P|$ times. Thus, we can assume that $\phi_{B_i}(\mathcal{I}(B_i)) = \phi_{B_{i+1}}(\mathcal{I}(B_{i+1}))$. Note that (1) can happen at most $|P|$ times due to Lemma 20 and because $\phi_{B_{i+1}}(I') \neq \emptyset$ for all $I' \in \mathcal{I}(B_{i+1})$. Indeed, for all $\ell \in P$, Lemma 20 tells us there are no pairwise distinct intervals $I_1, I_2, I_3 \in \mathcal{I}(B_{i+1})$ such that $\ell \in \phi_{B_{i+1}}(I_1)$, $\ell \in \phi_{B_{i+1}}(I_2)$, and $\ell \in \phi_{B_{i+1}}(I_3)$.

Now, assume that (1) and (3) are not the case and that $\phi_{B_i}(\mathcal{I}(B_i)) = \phi_{B_{i+1}}(\mathcal{I}(B_{i+1}))$. Since (1) does not hold, we have $|\mathcal{I}(B_{i+1})| \leq |\mathcal{I}(B_i)|$. Note that a strict inequality can happen at most $|P|$ times, so we can assume that $|\mathcal{I}(B_{i+1})| = |\mathcal{I}(B_i)|$. We define a function $f: \mathcal{I}(B_i) \rightarrow \mathcal{I}(B_{i+1})$. Recall that $B_i \subseteq B_{i+1}$. So, for every $I \in \mathcal{I}(B_i)$ there exists a unique $f(I) \in \mathcal{I}(B_{i+1})$ such that $I \subseteq f(I)$, and hence $\phi_{B_i}(I) \subseteq \phi_{B_{i+1}}(f(I))$. Uniqueness follows from maximality of intervals within $\mathcal{I}(B_{i+1})$. Since (1) does not hold, the function f is a surjection. Moreover, $\ell \in \phi_{B_i}(I)$ iff $\ell \in \phi_{B_{i+1}}(f(I))$. Thus, (2) can happen at most $|P|$ times by Lemma 20. \square

As a corollary of how we computed $\text{Reach}_{p(a)}$ we get the following result that will be needed in section V.

Corollary 30. *The mapping $\text{Reach}_{p(a)}$ is \mathcal{C} -valid.*

V. PARAMETRIC COCA REACHABILITY

Finally, we consider the reachability problem $p(a) \rightarrow_* q(b)$ for parametric COCAs. We build on the proof of the last section for guarded COCA. However, we “guess” under-approximations of reachability functions by efficiently encoding them in the $\exists\forall$ -fragment of $\text{FO}(\mathbb{Q}, +, <)$ instead of using iterative computation. We then invoke the following result:

Theorem 31 ([21], [22]). *The language of sentences in the $\exists\forall$ -fragment of $\text{FO}(\mathbb{Q}, +, <)$ is decidable and in Σ_2^P .*

In most of the section, we will work with a guarded COCA \mathcal{W} rather than a parametric COCA $\mathcal{P} = (Q, T, \tau, X)$. One should think of \mathcal{W} as \mathcal{P}^μ for some valuation μ . This valuation will be guessed later on in our final $\text{FO}(\mathbb{Q}, +, <)$ formula.

A. Characterisation of the reachability function

Recall the mapping $\text{Reach}_{p(a)} \in \mathcal{R}_Q$ from section IV defined by $\text{Reach}_{p(a)}(r) := \text{Post}_{p,r}(a)$. Note that $p(a) \rightarrow_* q(b)$ holds iff $b \in \text{Reach}_{p(a)}(q)$.

Let $R \in \mathcal{R}_Q$ and recall the Succ function from section IV. We say that mapping R is a *reachability candidate* if $a \in R(p)$ and $\text{Succ}(R) \preceq R$. For example, the mapping U defined by $U(r) := \mathbb{Q}$, for all $r \in Q$, is (trivially) a reachability candidate. As we will show, $\text{Reach}_{p(a)}$ is also a reachability candidate.

The following provides a characterisation of $\text{Reach}_{p(a)}$ that will allow us to encode it in $\text{FO}(\mathbb{Q}, +, <)$:

Proposition 32. *Let $R \in \mathcal{R}_Q$. It is the case that $\text{Reach}_{p(a)} = R$ iff the following two conditions hold:*

- R is a reachability candidate, and
- $R \preceq R'$ for every reachability candidate $R' \in \mathcal{R}_Q$.

Proof. \Rightarrow Let $R := \text{Reach}_{p(a)}$. By definition, we have $a \in R$ and $\text{Succ}(R) = R$. Hence, R is a reachability candidate. Let $R' \in \mathcal{R}_Q$ be a reachability candidate. Let $q \in Q$ and $b \in R(q)$. We must show that $b \in R'(q)$. By definition of R , we have $p(a) \rightarrow_\rho q(b)$ for some run ρ . Note that $\text{Succ}(R') \preceq R'$. Thus, by induction, we have $\text{Succ}^{|\rho|}(R') \preceq R'$. Since $a \in R'(p)$, by [Lemma 22](#) b belongs to $\text{Succ}^{|\rho|}(R')(q)$ and hence to $R'(q)$.

\Leftarrow Let $S_0 \in \mathcal{R}_Q$ be the mapping defined as $S_0(p) := \{a\}$ and $S_0(r) := \emptyset$ for every $r \neq p$. Let $S_i := \text{Succ}(S_{i-1})$ for every $i \geq 1$. Since R is a reachability candidate, we have $a \in R(p)$ and $\text{Succ}(R) \preceq R$. In particular, we have $S_0 \preceq R$ and hence $S_1 = \text{Succ}(S_0) \preceq \text{Succ}(R) \preceq R$.

By induction, we obtain $\bigcup_{i \in \mathbb{N}} S_i \preceq R$. By [Lemma 22](#), $\text{Reach}_{p(a)} = \bigcup_{i \in \mathbb{N}} \text{Succ}(S_i)$. Thus, $\text{Reach}_{p(a)} \preceq R$. Since $\text{Reach}_{p(a)}$ is a reachability candidate, we have $R \preceq \text{Reach}_{p(a)}$ by assumption. This shows that $R = \text{Reach}_{p(a)}$. \square

By [Lemma 21](#) and [Corollary 30](#), $\text{Reach}_{p(a)}(q)$ consists of at most n intervals for every $q \in Q$, where n is polynomial in $|Q|$. We define $\mathcal{R}_Q^n \subseteq \mathcal{R}_Q$ as the following set of mappings:

$$\mathcal{R}_Q^n := \{R : Q \rightarrow 2^{\mathbb{Q}} \mid |\mathcal{I}(R(q))| \leq n \text{ for all } q \in Q\}.$$

In particular, we have $\text{Reach}_{p(a)} \in \mathcal{R}_Q^n$. Hence, we may rewrite [Proposition 32](#) as follows:

Corollary 33. *Let $R \in \mathcal{R}_Q^n$. It is the case that $\text{Reach}_{p(a)} = R$ iff the following two conditions hold:*

- R is a reachability candidate, and
- $R \preceq R'$ for every reachability candidate $R' \in \mathcal{R}_Q^n$.

B. Encoding intervals

We now describe our logical encoding of intervals. An interval I is uniquely determined by its endpoints and their membership in I . We represent intervals as tuples (b, t, \perp, \top) , where $b, t \in \mathbb{Q}$ and $\perp, \top \in \{0, 1, 2\}$. Such a tuple represents \emptyset if $(\perp \neq 2 \wedge \top \neq 2) \wedge (b > t \vee (b = t \wedge \neg(\top = \perp = 1)))$ holds, and otherwise the unique interval $I \in \mathcal{I}$ such that

$$\perp = \begin{cases} 0 & \text{if } \inf I = b \notin I, \\ 1 & \text{if } \inf I = b \in I, \\ 2 & \text{if } \inf I = -\infty, \end{cases} \quad \top = \begin{cases} 0 & \text{if } \sup I = t \notin I, \\ 1 & \text{if } \sup I = t \in I, \\ 2 & \text{if } \sup I = +\infty. \end{cases}$$

We write $\text{int}(I)$ to denote the interval represented by the tuple $I = (b, t, \perp, \top)$, e.g. $\text{int}(2, 4, 0, 1) = (2, 4]$, $\text{int}(1, 5, 2, 0) = (-\infty, 5)$ and $\text{int}(3, -2, 1, 0) = \emptyset$. Note that some intervals are not encoded uniquely, e.g. $\text{int}(0, 5, 2, 0) = (-\infty, 5)$ as well.

Let $\mathcal{E} := \mathbb{Q}^2 \times \{0, 1, 2\}^2$ be the set of all interval encodings. We will represent each such encoding with four variables from $\text{FO}(\mathbb{Q}, +, <)$, i.e. $I_x = (x_b, x_t, x_\perp, x_\top)$. When I_x appears in a formula, we will assume that it is conjoined with $\bigvee_{i=0}^2 (x_\perp = i) \wedge \bigvee_{i=0}^2 (x_\top = i)$. Hence, variables x_\top and x_\perp evaluate only to $\{0, 1, 2\}$. From now on, we can define $\text{FO}(\mathbb{Q}, +, <)$ formulas that operate on variables of the form I_x . As usual, outputs can be represented with an extra variable, e.g. $z = \min(x, y)$ can be expressed as a quantifier-free formula.

Given two interval encodings $I = (b, t, \perp, \top) \in \mathcal{E}$ and $I' = (b', t', \perp', \top') \in \mathcal{E}$, we can construct an encoding $I'' \in \mathcal{E}$ such that $\text{int}(I'') = \text{int}(I) + \text{int}(I')$:

$$I'' := (b + b', t + t', \text{msum}(\perp, \perp'), \text{msum}(\top, \top'));$$

where $\text{msum} : \{0, 1, 2\}^2 \rightarrow \{0, 1, 2\}$ is defined as:

$$\text{msum}(i, j) = \begin{cases} 2 & \text{if } \max(i, j) = 2; \\ \min(i, j) & \text{otherwise.} \end{cases}$$

Note that I'' can be written as a quantifier-free linear formula as it involves addition and comparisons. By a slightly tedious case distinction, it is further possible to construct a quantifier-free linear formula that specifies an encoding $I'' \in \mathcal{E}$ such that $\text{int}(I'') = \text{int}(I) \cap \text{int}(I')$ (see [subsection C](#) of the appendix).

Formally, for every variable valuation μ , we write $\mu(I_x)$ for $(\mu(x_b), \mu(x_t), \mu(x_\perp), \mu(x_\top))$. We can construct quantifier-free formulas φ_+ and φ_\cap such that for every valuation μ :

- $\mu(\varphi_\cap(I_x, I_y, I_z))$ holds iff $\mu(I_z) = \mu(I_x) \cap \mu(I_y)$,
- $\mu(\varphi_+(I_x, I_y, I_z))$ holds iff $\mu(I_z) = \mu(I_x) + \mu(I_y)$.

For example, the latter is defined as:

$$\varphi_+(I_x, I_y, I_z) := (z_b = x_b + y_b) \wedge (z_t = x_t + y_t) \wedge z_{\perp} = \text{msum}(x_{\perp}, y_{\perp}) \wedge z_{\top} = \text{msum}(x_{\top}, y_{\top}).$$

We will further use the following quantifier-free formulas for membership, interval inclusion and emptiness checks:

$$\begin{aligned} \varphi_{\in}(x, I_y) &:= (y_b < x) \vee (y_b = x \wedge y_{\perp} = 1) \vee (y_{\perp} = 2) \\ &\quad \wedge (x < y_t) \vee (y_t = x \wedge y_{\top} = 1) \vee (y_{\top} = 2); \\ \varphi_{\subseteq}(I_x, I_y) &:= (x_{\perp} = 2 \rightarrow y_{\perp} = 2) \wedge (x_{\top} = 2 \rightarrow y_{\top} = 2) \\ &\quad \wedge (y_b < x_b) \vee (y_b = x_b \wedge x_{\perp} \leq y_{\perp}) \vee (y_{\perp} = 2) \\ &\quad \wedge (x_t < y_t) \vee (y_t = x_t \wedge x_{\top} \leq y_{\top}) \vee (y_{\top} = 2); \\ \varphi_{\emptyset}(I_x) &:= (x_{\perp} \neq 2) \wedge (x_{\top} \neq 2) \\ &\quad \wedge ((x_b > x_t) \vee (x_b = x_t \wedge \neg(x_{\top} = x_{\perp} = 1))). \end{aligned}$$

For every valuation μ , the following holds by definition:

- $\mu(\varphi_{\in}(x, I_y))$ holds iff $\mu(x) \in \text{int}(\mu(I_y))$;
- $\mu(\varphi_{\subseteq}(I_x, I_y))$ holds iff $\text{int}(\mu(I_x)) \subseteq \text{int}(\mu(I_y))$;
- $\mu(\varphi_{\emptyset}(I_x))$ holds iff $\text{int}(\mu(I_x)) = \emptyset$.

C. Unions of intervals

We will consider a union of intervals to represent $\text{Reach}_{p(a)}$. We will represent them as vectors of variables. We write $\mathbf{I}_x = (I_{x_1}, \dots, I_{x_n})$ to denote a vector of $4n$ variables, where n is the bound from [Corollary 33](#). Given a valuation μ , we write $\mu(\mathbf{I}_x)$ for the valuation of all variables in \mathbf{I}_x , and we write $\text{int}(\mu(\mathbf{I}_x)) := \bigcup_{i=1}^n \text{int}(\mu(I_{x_i}))$.

We define formulas dealing with union of intervals \mathbf{I}_x and generalising the formulas on I_x . For membership, we define:

$$\psi_{\in}(x, \mathbf{I}_y) := \bigvee_{i=1}^n \varphi_{\in}(x, I_{y_i}).$$

For Minkowski sums, we will only need to sum a union of intervals with a single interval:

$$\psi_+(\mathbf{I}_x, I_y, \mathbf{I}_z) := \bigwedge_{i=1}^n \varphi_+(I_{x_i}, I_y, I_{z_i}).$$

Similarly, we will intersect a union of intervals with a single interval. Using $I \cap \bigcup_{a \in A} I_a = \bigcup_{a \in A} I \cap I_a$, we can define:

$$\psi_{\cap}(\mathbf{I}_x, I_y, \mathbf{I}_z) := \bigwedge_{i=1}^n \varphi_{\cap}(I_{x_i}, I_y, I_{z_i}).$$

The formula for interval inclusion is defined as:

$$\psi_{\subseteq}(\mathbf{I}_x, \mathbf{I}_y) := \bigwedge_{i=1}^n \bigvee_{j=1}^n \varphi_{\subseteq}(I_{x_i}, I_{y_j}).$$

Note that this formula does not exactly express inclusion. For example, let $n := 2$. Suppose \mathbf{I}_x encodes $(1, 5) \cup (4, 9)$ and \mathbf{I}_y encodes $[1, 4] \cup [4, 10]$. We have $(1, 5) \cup (4, 9) \subseteq [1, 4] \cup [4, 10]$ even though $(1, 5)$ is not a subset of any of the latter intervals.

Therefore, we need a formula to express that \mathbf{I}_x is decomposed into maximal intervals. Let us start with a formula to express that an interval does not ‘‘prolong’’ another one (e.g. as opposed to $[4, 10]$ which prolongs $[1, 4]$):

$$\varphi_{dis}(I_x, I_y) := (x_t = y_b) \rightarrow (x_{\top} = y_{\perp} = 0).$$

Notice that this formula alone does not express that I_x and I_y are disjoint. The following formula expresses that \mathbf{I}_x consists of maximal intervals, using some auxiliary intervals $I_{y_{i,j}}$:

$$\psi_{max}(\mathbf{I}_x) := \bigwedge_{1 \leq i, j \leq n} (\varphi_{\cap}(I_{x_i}, I_{x_j}, I_{y_{i,j}}) \rightarrow \varphi_{\emptyset}(I_{y_{i,j}})) \wedge \varphi_{dis}(I_{x_i}, I_{x_j}).$$

The above formula has a quadratic number of variables, but we will omit variables $I_{y_{i,j}}$ in $\psi_{max}(\mathbf{I}_x)$ for readability. For every valuation μ , it is readily seen that $\mu(\psi_{max}(\mathbf{I}_x))$ holds iff $\text{int}(\mu(I_{x_1})), \dots, \text{int}(\mu(I_{x_n}))$ are the maximal intervals from $\mathcal{I}(\text{int}(\mu(\mathbf{I}_x)))$, plus possibly some empty intervals. Thus, we say that $\mu(\psi_{max}(\mathbf{I}_x))$ is decomposed into maximal intervals.

More formally, the following holds for every valuation μ :

- $\mu(\psi_{\in}(x, \mathbf{I}_y))$ holds iff $\mu(x) \in \text{int}(\mu(\mathbf{I}_y))$;
- $\mu(\psi_{+}(\mathbf{I}_x, I_y, \mathbf{I}_z))$ holds iff $\text{int}(\mu(\mathbf{I}_x)) + \text{int}(\mu(I_y)) = \text{int}(\mu(\mathbf{I}_z))$;
- $\mu(\psi_{\cap}(\mathbf{I}_x, I_y, \mathbf{I}_z))$ holds iff $\text{int}(\mu(\mathbf{I}_x)) \cap \text{int}(\mu(I_y)) = \bigcup_{i=1}^n \text{int}(I_{z_i})$;
- $\mu(\psi_{\subseteq}(\mathbf{I}_x, \mathbf{I}_y) \cap \psi_{max}(\mathbf{I}_y))$ holds iff \mathbf{I}_y is decomposed into maximal intervals and $\text{int}(\mathbf{I}_x) \subseteq \text{int}(\mathbf{I}_y)$.

D. Encoding the reachability function

For every $z \in \mathbb{Z} \cup X$, let $I_z^+, I_z^-, I_z^0 \in \mathcal{E}$ be defined as $I_z^+ := (0, z, 0, 1)$, $I_z^- := (z, 0, 1, 0)$, and $I_z^0 = (0, 0, 1, 1)$. These encode $(0, z]$, $[z, 0)$ and $[0, 0]$ which are the possible scalings of z depending on its sign. For every $q \in Q$, let $I_q \in \mathcal{E}$ be such that $I_q := (q_b, q_t, q_{\perp}, q_{\top})$ encodes $\tau(q)$. For example, if $\tau(q) = (x, 10]$, then we can take $I_q := (x, 10, 0, 1)$. Note that q_b and q_t can be variables corresponding to parameters.

Given \mathbf{I}_x , for all transition $t \in T$, we define \mathbf{I}_t through:

$$\begin{aligned} \psi_t(\mathbf{I}_x, \mathbf{I}_t) &:= (\Delta(t) > 0) \rightarrow \psi_{+}(\mathbf{I}_x, I_{\Delta(t)}^+, \mathbf{I}_y) \\ &\quad \wedge (\Delta(t) < 0) \rightarrow \psi_{+}(\mathbf{I}_x, I_{\Delta(t)}^-, \mathbf{I}_y) \\ &\quad \wedge (\Delta(t) = 0) \rightarrow \psi_{+}(\mathbf{I}_x, I_{\Delta(t)}^0, \mathbf{I}_y) \\ &\quad \wedge \psi_{\cap}(\mathbf{I}_y, I_{\text{out}(t)}, \mathbf{I}_t). \end{aligned}$$

Note that the above formula uses auxiliary variables \mathbf{I}_y , $I_{\Delta(t)}^+$, $I_{\Delta(t)}^-$, $I_{\Delta(t)}^0$, and $I_{\text{out}(t)}$.

Let $\mathbf{I}_Q = (\mathbf{I}_r)_{r \in Q}$ be the vector of $|Q| \cdot 4n$ variables, where each \mathbf{I}_r consists of n intervals. Similarly, we define \mathbf{I}_T as a vector of $|T| \cdot 4n$ variables. We define the formula

$$\psi_{succ}(\mathbf{I}_Q, \mathbf{I}_T) := \bigwedge_{t \in T} \psi_t(\mathbf{I}_{\text{in}(t)}, \mathbf{I}_t).$$

Given a valuation μ and a vector \mathbf{I}_Q , we define the mapping $\mu(\mathbf{I}_Q)$ as $q \mapsto \text{int}(\mu(\mathbf{I}_q))$. The following holds by definition:

Lemma 34. *For every valuation μ , $\psi_{succ}(\mathbf{I}_Q, \mathbf{I}_T)$ holds iff for every $r \in Q$ the following holds:*

$$\text{Succ}_r(\mu(\mathbf{I}_Q)) = \mu(\mathbf{I}_Q)(r) \cup \bigcup_{\substack{t \in T, \\ \text{out}(t)=r}} \mu(\mathbf{I}_t).$$

This allows us to define the following formula:

$$\begin{aligned} \psi_{cand}(\mathbf{I}_Q) &:= \psi_{succ}(\mathbf{I}_Q, \mathbf{I}_T) \\ &\quad \wedge \bigwedge_{r \in Q} \bigwedge_{\substack{t \in T \\ \text{out}(t)=r}} \psi_{\subseteq}(\mathbf{I}_t, \mathbf{I}_r) \cap \psi_{max}(\mathbf{I}_r) \wedge \psi_{\in}(a, \mathbf{I}_p). \end{aligned}$$

Again, by definition we obtain:

Lemma 35. *For all valuation μ , it is the case that $\psi_{cand}(\mathbf{I}_Q)$ holds iff $\mu(\mathbf{I}_Q)$ is a reachability candidate.*

Let Q' be a disjoint copy of Q . We write \mathbf{X} for the vector of variables corresponding to parameters of the COCA, \mathbf{Y} for the vector of all remaining variables used in the quantifier-free formula $\psi_{cand}(\mathbf{I}_Q)$ (not just \mathbf{I}_Q), and \mathbf{Z} for the vector of remaining variables used in the quantifier-free formula $\psi_{cand}(\mathbf{I}_{Q'})$. Further, recall I_p encodes $\tau(p)$. We define:

$$\begin{aligned} \psi &:= \exists \mathbf{X} \exists \mathbf{Y} \forall \mathbf{Z} \psi_{cand}(\mathbf{I}_Q) \wedge \psi_{cand}(\mathbf{I}_{Q'}) \wedge \psi_{\in}(a, I_p) \\ &\quad \wedge \bigwedge_{r \in Q} \psi_{\subseteq}(\mathbf{I}_r, \mathbf{I}_{r'}) \wedge \psi_{max}(\mathbf{I}_r) \wedge \psi_{\in}(b, \mathbf{I}_q). \end{aligned}$$

Since ψ holds iff $p(a) \rightarrow_* q(b)$, by [Corollary 33](#) we obtain:

Theorem 36. *The (existential) reachability problem for parametric COCAs belongs in Σ_2^P .*

E. Integer valuations

We briefly consider parametric COCAs where only updates can be parameterised. In this setting, a rational valuation that witnesses reachability can be turned into an *integer* valuation witnessing reachability. This follows by rescaling the factors of the witnessing run so that it remains admissible.

Lemma 37. *Let μ be a valuation under which $p(a) \rightarrow_* q(b)$. For any valuation μ' such that $\mu'(x) = \lambda\mu(x)$ with $\lambda \in \mathbb{N}_{\geq 1}$, it is the case that $p(a) \rightarrow_* q(b)$ under μ' .*

Proof. Let $\lambda \in \mathbb{N}_{\geq 1}$ and let μ' be defined w.r.t. μ and λ . Let $s(v) \rightarrow_{\alpha t} s'(v')$ be consecutive configurations from the run ρ witnessing $p(a) \rightarrow_{\rho} q(b)$ under valuation μ . If $\Delta(t) \in \mathbb{Z}$, then no rescaling is needed as the update of t is nonparametric. Otherwise, we have $v' - v = \alpha \cdot \mu(\Delta(t))$. Let $\beta := \alpha/\lambda$. Since $\alpha \in (0, 1]$ and $\lambda \geq 1$, we have $\beta \in (0, 1]$. Therefore, we have $s(v) \rightarrow_{\beta t} s'(v')$. Hence, by rescaling each transition, we obtain a run ρ' such that $p(a) \rightarrow_{\rho'} q(b)$ under μ' . \square

Now, consider a rational valuation μ witnessing $p(a) \rightarrow_* q(b)$. Since μ is rational, each parameter value $\mu(x)$ can be represented as a fraction a_x/b_x . By Lemma 37, we know that valuation $\mu'(x) = \lambda\mu(x)$, where $\lambda := \prod_{x \in X} b_x$, also witnesses reachability. Moreover, it is integral, hence:

Corollary 38. *The (existential) reachability problem for parametric COCAs, where valuations must be integral, is equivalent to the rational variant if guards are nonparametric.*

F. Hardness result and acyclic parametric COCA

To conclude our treatment of parametric COCAs, we establish **NP**-hardness of the reachability problem, and a matching **NP** upper bound for the special case of acyclic COCAs.

Theorem 39. *The reachability problem for parametric COCAs is **NP**-hard. Moreover, this holds if the underlying graph is acyclic, and if parameters occur on transitions, guards or both. In the latter special cases, the problem is **NP**-complete.*

Proof. We first show membership in **NP** for the acyclic case. To determine whether $p(a) \rightarrow_* q(b)$ under some valuation μ , we encode the problem into an existential FO($\mathbb{Q}, +, <$) sentence. Since the language of such sentences is known to be in **NP** [22] the result will follow.

We start with the observation that $\text{Post}_{\pi}(a)$ is an interval for every path π since it can be described as a chain of Minkowski sums and intersections starting from $[a, a]$. Recall the formula $\psi_t(\mathbf{I}_x, \mathbf{I}_t)$ from subsection V-D. We assume there is a bijection f between T and $\{1, \dots, |T|\}$ and identify $t \in T$ with $f(t)$. Let $\varphi_{f(t)}(I_x, I_t)$ be defined in the same way as $\psi_t(\mathbf{I}_x, \mathbf{I}_t)$ but replacing vectors of intervals with intervals and ψ_+ , ψ_{\cap} with φ_+ , φ_{\cap} . Intuitively, $\varphi_{f(t)}(I_x, I_t)$ means that $I_t = \text{Post}_t(I_x)$.

Given two transitions $t, t' \in T$ we also need a formula that checks $\text{out}(t) = \text{in}(t')$. We define this by $\varphi_{\text{next}}(x, y) := \bigwedge_{i,j=1}^{|T|} (x = i \wedge y = j) \rightarrow t_{ij}$, where t_{ij} is a Boolean constant that is true iff $\text{out}(f^{-1}(i)) = \text{in}(f^{-1}(j))$. Similarly, we define $\varphi_{\text{init}}(x)$, $\varphi_{\text{fin}}(x)$ as the formulas which are true iff x is a transition that starts in state p and finishes in state q , respectively.

Let $\mathbf{T} = (t_1, \dots, t_{|Q|})$ and $\mathbf{I} = (I_1, \dots, I_{|Q|})$ be vectors of variables representing $|Q|$ transitions and $|Q|$ intervals. We write I_0 for the vector $(a, a, 1, 1)$ encoding $[a, a]$. Finally, let \mathbf{X} be the vector of variables encoding the parameters of the COCA. We define the final formula

$$\begin{aligned} \exists \mathbf{X} \exists \mathbf{T} \exists \mathbf{I} \quad & \varphi_{\text{init}}(t_1) \wedge \bigwedge_{i=1}^{|Q|} \bigvee_{j=1}^{|T|} t_i = j \wedge \varphi_j(I_{i-1}, I_i) \\ & \wedge \bigvee_{m=1}^{|Q|} \varphi_{\text{fin}}(t_m) \wedge \varphi_{\in}(b, I_m) \wedge \bigwedge_{i=1}^{m-1} \varphi_{\text{next}}(t_i, t_{i+1}). \end{aligned}$$

By definition, this sentence holds iff $p(a) \rightarrow_* q(b)$. Indeed, since the COCA is acyclic all runs are of length at most $|Q|$. Thus, in the first line we “guess” transitions per state and “compute” their reachability set; then, in the second line we check whether they form a path π and whether $p(a) \rightarrow_{\pi} q(b)$.

It remains to show **NP**-hardness for the acyclic case, where parameters either only occur on transitions or on guards. In both cases, we give a reduction from 3-SAT. Let $\varphi = \bigwedge_{1 \leq j \leq m} C_j$ be a 3-CNF formula over variables $X = \{x_1, \dots, x_n\}$.

Let us give two acyclic parametric COCAs \mathcal{P} and \mathcal{P}' , both with parameters X . Each one will guess an assignment to X , and check whether it satisfies φ . Additionally, \mathcal{P} uses parameters only on guards; \mathcal{P}' , only on updates. We only sketch \mathcal{P} . While the exact gadgets used in the construction for \mathcal{P}' differ, they perform the same functions (see the appendix).

The first part is done by sequentially composing n copies of the left gadget depicted in Figure 3. The gadget functions as follows: (1) state p_i is entered with counter value 0; (2) the counter is set to x_i ; (3) membership of the counter value in $\{0, 1\}$ is checked; and (4) the counter is reset to zero upon leaving to q_i . The only way to traverse the chain of n such gadgets from p_1 to q_n is to have $x_i \in \{0, 1\}$ for each $x_i \in X$.




Fig. 3. Gadgets of the reduction from 3-SAT, where parameters occur only on guards, for: variable x_i (left) and clause $C_j = (x_1 \vee x_2 \vee \neg x_4)$ (right).

The second part is achieved by chaining a gadget for each clause similar to the one depicted on the right-hand side of Figure 3 for $C_j = (x_1 \vee x_2 \vee \neg x_4)$. In words, it (1) enters state r_j with the counter value set to 0; (2) nondeterministically picks a variable x_i of some literal of C_j and increments the counter by x_i ; (3) checks whether the counter holds the right value w.r.t. the literal polarity; and (4) resets the counter to zero upon leaving to state s_j . Thus, the chain of gadgets can be traversed from r_1 to s_m iff φ is satisfied by the assignment.

Altogether, these statements are equivalent: (1) formula φ is satisfiable; (2) there exists a valuation $\mu: X \rightarrow \mathbb{Q}$ such that $p_1(0) \rightarrow_* s_m(0)$ holds in \mathcal{P}^μ ; and (3) there exists a valuation $\mu': X \rightarrow \mathbb{Q}$ such that $p_1(0) \rightarrow_* s_m(0)$ holds in $\mathcal{P}^{\mu'}$ \square

VI. CONCLUSION

In this work, we have introduced COCA and guarded COCA as over-approximations of SOCA, and we have given efficient algorithms for their reachability problems. For both models, the only lower bound we are aware of is the NL-hardness that follows trivially from the directed-graph reachability problem. It thus remains open whether our algorithms are computationally optimal. Additionally, we have shown that the reachability problem lies in the polynomial hierarchy for parametric COCA, in contrast to the discrete variant whose decidability is unknown. We leave open whether the reachability problem for parametric COCA with integer-valued parameters is decidable in general (i.e., when guards can also be parametric).

REFERENCES

- [1] M. L. Minsky, "Recursive unsolvability of Post's problem of "tag" and other topics in theory of Turing machines," *Annals of Mathematics*, vol. 74, no. 3, pp. 437–455, 1961.
- [2] —, *Computation: Finite and Infinite Machines*. Prentice-Hall, Inc., 1967.
- [3] R. J. Lipton, "The reachability problem requires exponential space," Yale University, Tech. Rep. 62, 1976. [Online]. Available: <http://cpsc.yale.edu/sites/default/files/files/tr63.pdf>
- [4] E. W. Mayr, "An algorithm for the general Petri net reachability problem," in *Proc. 13th Annual ACM Symposium on Theory of Computing (STOC)*. ACM, 1981, pp. 238–246.
- [5] J. Leroux and S. Schmitz, "Reachability in vector addition systems is primitive-recursive in fixed dimension," in *Proc. 34th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, 2019, pp. 1–13.
- [6] W. Czerwinski, S. Lasota, R. Lazic, J. Leroux, and F. Mazowiecki, "The reachability problem for Petri nets is not elementary," in *Proc. 51st Annual ACM SIGACT Symposium on Theory of Computing (STOC)*. ACM, 2019, pp. 24–33.
- [7] C. Haase, S. Kreutzer, J. Ouaknine, and J. Worrell, "Reachability in succinct and parametric one-counter automata," in *Proc. 20th International Conference on Concurrency Theory (CONCUR)*. Springer, 2009, pp. 369–383.
- [8] J. Fearnley and M. Jurdziński, "Reachability in two-clock timed automata is PSPACE-complete," *Information and Computation*, vol. 243, pp. 26–36, 2015.
- [9] B. Bollig, K. Quaas, and A. Sangnier, "The complexity of flat freeze LTL," *Logical Methods in Computer Science*, vol. 15, no. 3, 2019.
- [10] C. Baier and J. Katoen, *Principles of model checking*. MIT Press, 2008.
- [11] A. Bouajjani, M. Bozga, P. Habermehl, R. Iosif, P. Moro, and T. Vojnar, "Programs with lists are counter automata," *Formal Methods in System Design*, vol. 38, no. 2, pp. 158–192, 2011.
- [12] C. Chitic and D. Rosu, "On validation of XML streams using finite state machines," in *Proc. 7th International Workshop on the Web and Databases (WebDB)*. ACM, 2004, pp. 85–90.

- [13] C. Haase, J. Ouaknine, and J. Worrell, "On the relationship between reachability problems in timed and counter automata," in *Proc. 6th International Workshop Reachability Problems (RP)*. Springer, 2012, pp. 54–65.
- [14] D. Bundala and J. Ouaknine, "On parametric timed automata and one-counter machines," *Information and Computation*, vol. 253, pp. 272–303, 2017.
- [15] R. Alur, T. A. Henzinger, and M. Y. Vardi, "Parametric real-time reasoning," in *Proc. Twenty-Fifth Annual ACM Symposium on Theory of Computing (STOC)*. ACM, 1993, pp. 592–601.
- [16] O. H. Ibarra, T. Jiang, N. Q. Tran, and H. Wang, "New decidability results concerning two-way counter machines and applications," in *Proc. 20th International Colloquium on Automata, Languages and Programming (ICALP)*. Springer, 1993, pp. 313–324.
- [17] M. Blondin and C. Haase, "Logics for continuous reachability in Petri nets and vector addition systems with states," in *Proc. 32nd Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*. IEEE Computer Society, 2017, pp. 1–12.
- [18] F. Laroussinie, N. Markey, and P. Schnoebelen, "Model checking timed automata with one or two clocks," in *Proc. 15th International Conference on Concurrency Theory (CONCUR)*. Springer, 2004, pp. 387–401.
- [19] C. H. Papadimitriou, *Computational complexity*. Addison-Wesley, 1994.
- [20] S. Arora and B. Barak, *Computational Complexity - A Modern Approach*. Cambridge University Press, 2009.
- [21] J. Ferrante and C. Rackoff, "A decision procedure for the first order theory of real addition with order," *SIAM Journal on Computing*, vol. 4, no. 1, pp. 69–76, 1975.
- [22] E. D. Sontag, "Real addition and the polynomial hierarchy," *Information Processing Letters*, vol. 20, no. 3, pp. 115–120, 1985.
- [23] D. P. Bovet and P. Crescenzi, *Introduction to the Theory of Complexity*. Prentice Hall London, 1994.

A. Missing proofs of section III

Lemma 3. Let $G = (Q, E)$ be a weighted multigraph whose weights are encoded in binary, and let $p, q \in Q$ be nodes. Deciding whether $S = \emptyset$ is in \mathbf{NC}^2 , where S is the set of paths $\pi \in \text{Paths}_{p,q}$ that satisfy a fixed subset of these conditions²:

- (a) $\Delta^+(\pi) \neq 0$ (resp. $\Delta^-(\pi) \neq 0$);
- (b) $\Delta^+(\pi) = 0$ (resp. $\Delta^-(\pi) = 0$);
- (c) $\Delta(\text{first}(\pi)) < 0$ (resp. $\Delta(\text{first}(\pi)) > 0$);
- (d) $\Delta(\text{last}(\pi)) < 0$ (resp. $\Delta(\text{last}(\pi)) > 0$).

Furthermore, for any such set S , the following value can be computed in \mathbf{NC}^2 : $\text{opt}\{w(\pi) \mid \pi \in S \text{ and } |\pi| \leq |Q|\}$, where $\text{opt} \in \{\min, \max\}$ and $w \in \{\Delta^+, \Delta^-\}$.

Proof. Let us explain how one can check emptiness of S when only one of the conditions is required by reducing the problem to standard graph reachability, which is in $\mathbf{NL} \subseteq \mathbf{NC}^2$.

For each condition, we focus only on one of the two stated cases. The other case will follow similarly. Each time, we will give a graph H such that reachability between two fixed nodes of H corresponds to reachability from p to q in G via a path that satisfies the condition.

(a) Let us define H as joining G with a modified copy \overline{G} which remains in \overline{G} with nonpositive edges, and moves into G with positive edges. More formally, let $H := (Q', E')$, where $Q' := Q \cup \{\overline{q} \mid q \in Q\}$. Further, $E' := E \cup \{(\overline{p}, z, \overline{q}) \mid (p, z, q) \in E, z \leq 0\} \cup \{(\overline{p}, z, q) \mid (p, z, q) \in E, z > 0\}$. It is easy to see any path $\overline{\pi}$ from \overline{p} to \overline{q} in H corresponds to a path π from p to q in G such that $\Delta^+(\pi) \neq 0$.

(b) We set $H := (Q, E')$, where $E' := \{(p', z, q') \in E \mid z \leq 0\}$. Clearly, reachability from p to q in H is equivalent to reachability from p to q in G via a path π with $\Delta^+(\pi) = 0$.

(c) We again define H as joining G with a modified copy \overline{G} . In \overline{G} , we omit all positive edges, while edges with weight zero remain in \overline{G} and negative edges lead to G . Formally, let $H := (Q', E')$, where $Q' := Q \cup \{\overline{q} \mid q \in Q\}$ and $E' := E \cup \{(\overline{p}, 0, \overline{q}) \mid (p, 0, q) \in E\} \cup \{(\overline{p}, z, q) \mid (p, z, q) \in E \wedge z < 0\}$. A path $\overline{\pi}$ from \overline{p} to \overline{q} in H corresponds to a path π from p to q in G such that $\Delta(\text{first}(\pi)) < 0$.

(d) We again join G with a modified copy \overline{G} . Now, \overline{G} omits all positive edges, and for each negative edge in G , we add a copy that leads from G to \overline{G} . We define $H := (Q', E')$ with $Q' := Q \cup \{\overline{q} \mid q \in Q\}$ and $E' := E \cup \{(p, z, \overline{q}) \mid (p, z, q) \in E \wedge z < 0\} \cup \{(\overline{p}, z, \overline{q}) \mid (p, z, q) \in E \wedge z \leq 0\}$. A path $\overline{\pi}$ from \overline{p} to \overline{q} in H corresponds to a path π from p to q in G such that $\Delta(\text{last}(\pi)) < 0$.

When we wish to require several conditions at once, note that for each condition, we constructed a graph H from a given input graph G . To require many conditions at once, we simply apply the transformations for each condition sequentially, and obtain a graph H' such that paths of H' satisfy all imposed conditions and correspond to paths in the original graph G . Observe that H' is of polynomial size.

Finally, let us argue that the following value can be computed in \mathbf{NC}^2 : $\text{opt}\{w(\pi) \mid \pi \in S \text{ and } |\pi| \leq |Q|\}$, where $\text{opt} \in \{\min, \max\}$ and $w \in \{\Delta^+, \Delta^-\}$. Let us first deal with $w = \Delta^+$, for which it suffices to treat edges with negative weight as having zero weight.

The problem of finding a shortest weighted path in a graph with edges of nonnegative weights is in \mathbf{NC}^2 (e.g., see [23, Example 12.4]). The procedure relies on the fact that there must be an acyclic shortest path, and hence that it suffices to consider paths of length at most $|Q|$. We can easily adapt the standard procedure for maximisation. Indeed, it successively minimises paths of length 1, 2, 4, 8, \dots , $|Q|$. By maximising rather than minimising, it follows that the claim holds for $\text{opt} \in \{\min, \max\}$. Note that finding a longest simple path is \mathbf{NP} -complete, while we obtain \mathbf{NC}^2 because we may find a nonsimple path (of length at most $|Q|$).

Since each of the conditions (a)–(d) is achieved by transforming the input graph into another graph of polynomial size, this holds also if we require any subset of these conditions.

The case of Δ^- can be handled similarly, e.g. by flipping the sign of the weights and the optimisation type (max/min). \square

Proposition 6 (Adapted from [17, Lemma 4.2(c)]). Let $\beta \in (0, 1]$ and let ρ be an admissible run from configuration $p(a)$. It is the case that run $\beta\rho$ is also admissible from $p(a)$.

Proof. We will show that for all $i \in \{0, \dots, |\rho|\}$, either $a \leq a + \Delta(\beta\rho[1..i]) \leq a + \Delta(\rho[1..i])$ or $a \geq a + \Delta(\beta\rho[1..i]) \geq a + \Delta(\rho[1..i])$. Since $a + \Delta(\rho[1..i]) \in \tau$ holds by the admissibility of ρ from a , it follows that $a + \Delta(\beta\rho[1..i]) \in \tau$, and so that $\beta\rho$ is admissible.

²The set of conditions may be empty, in which case $S = \text{Paths}_{p,q}$.

By definition, we have $a + \Delta(\beta\rho[1..i]) = a + \beta\Delta(\rho[1..i])$. Additionally, $\beta \in (0, 1]$. Hence, if $\Delta(\rho[1..i]) \geq 0$, then we have $a + \Delta(\rho[1..i]) \geq a + \Delta(\beta\rho[1..i]) \geq a$. If $\Delta(\rho[1..i]) < 0$, then $a + \Delta(\rho[1..i]) \leq a + \Delta(\beta\rho[1..i]) < a$, so we are done. \square

Lemma 7. For every $b \in \overline{\text{Post}_{p,q}(a)}$, it is the case that $(a, b) \subseteq \text{Post}_{p,q}(a)$ and $(b, a) \subseteq \text{Post}_{p,q}(a)$.

Proof. We only prove $(a, b) \subseteq \text{Post}_{p,q}(a)$ as the other inclusion is symmetric. We assume that $a < b$ as we are otherwise done. Let $c \in (a, b)$. Since $b \in \overline{\text{Post}_{p,q}(a)}$, there exists $b' \in \text{Post}_{p,q}(a)$ such that $b' \in [c, b]$. Let ρ be an admissible run from $p(a)$ to $q(b')$. By definition, $\Delta(\rho) = b' - a$. Let $\beta := (c - a)/(b' - a) \in (0, 1]$. By [Proposition 6](#), $\beta\rho$ is admissible from $p(a)$. Since $\Delta(\beta\rho) = c - a$, this concludes the proof of the main claim. \square

Proposition 9. The following statements hold:

- (a) If $\text{Post}_{p,q}(\inf \tau) \neq \emptyset$, then $\inf \overline{\text{Post}_{p,q}(\inf \tau)} = \inf \tau$.
- (b) If $\text{Post}_{p,q}(\sup \tau) \neq \emptyset$, then $\sup \overline{\text{Post}_{p,q}(\sup \tau)} = \sup \tau$.
- (c) Let $v \in \tau \setminus \{\inf \tau, \sup \tau\}$ and let ρ be a run. There exists $\varepsilon \in (0, 1]$ such that for all $\beta \in (0, \varepsilon]$ there exists $v_\beta > 0$ such that $v \rightarrow_{\beta\rho} v_\beta$. Moreover, $\lim_{\beta \rightarrow 0} v_\beta = v$.

Proof.

- (a) Let $a := \inf \tau$ and $c := \sup \overline{\text{Post}_{p,q}(a)}$. By [Lemma 7](#), we have $(a, c) \subseteq \overline{\text{Post}_{p,q}(a)}$. Since the latter is closed by definition, we have $\inf \overline{\text{Post}_{p,q}(a)} = \inf \tau$.
- (b) The proof is symmetric to (a).
- (c) Since $v \notin \{\inf \tau, \sup \tau\}$, there is a small enough $\varepsilon \in (0, 1]$ such that $v + |\varepsilon \cdot \Delta(\rho[1..i])| \in \tau$ for all $i \in \{1, \dots, |\rho|\}$. By definition, $\varepsilon\rho$ is admissible from v . Let $v_\beta := v + \beta \cdot \Delta(\rho)$. By [Proposition 6](#), $v \rightarrow_{\beta\rho} v_\beta$ is admissible for every $\beta \in (0, \varepsilon]$. Moreover, $\lim_{\beta \rightarrow 0} v_\beta = \lim_{\beta \rightarrow 0} v + \beta \cdot \Delta(\rho) = v$. \square

Lemma 10. If $\text{Post}_{p,q}(a) \neq \emptyset$ and \mathcal{V} has an (a, p, q) -admissible cycle θ , then the following holds:

- (a) $\inf \overline{\text{Post}_{p,q}(a)} = \inf \tau$, if θ is negative;
- (b) $\sup \overline{\text{Post}_{p,q}(a)} = \sup \tau$, if θ is positive.

Proof. Let $\theta = t\pi$ where t is the first transition of θ and π is the remaining path. Let $r := \text{in}(t)$. Since $a \in \text{enab}(\text{Paths}_{p,r})$, there is an admissible run ρ_1 from $p(a)$ that ends in state r . Similarly, since $\text{Paths}_{r,q} \neq \emptyset$, there is a run ρ_3 from r to q .

We only show (b) as (a) is symmetric. We assume that $a < \sup \tau$, as otherwise we are done by [Proposition 9\(b\)](#). We make a case distinction on whether $\sup \tau = \infty$.

Case $\sup \tau \neq \infty$. We must show that we can reach values arbitrarily close to $\sup \tau$, i.e. that for every $\varepsilon \in (0, 1]$, there exists a value $b \in [\sup \tau - \varepsilon, \sup \tau)$ and a run $p(a) \rightarrow_\rho q(b)$.

By [Proposition 6](#) and $a < \sup \tau$, we have $p(a) \rightarrow_{(1/2)\rho_1} r(a')$ for some $a' < \sup \tau$. Let:

$$m := \sum_{i=1}^{|\pi|} |\Delta(\pi_i)|, \alpha_t := \frac{\varepsilon}{4|\Delta(t)|} \text{ and } \alpha_\pi := \frac{\varepsilon}{4m+1}.$$

Let $\rho_2 := \alpha_t t \alpha_\pi \pi$. By definition, we have $\Delta(\alpha_t t) = \varepsilon/4$ and $|\Delta(\alpha_\pi \pi[1..i])| < \varepsilon/4$ for all $i \in \{1, \dots, |\pi|\}$. Consequently, it is the case that $\Delta(\rho_2[1..i]) \in (0, \varepsilon/2)$ for all $i \in \{2, \dots, |\rho_2|\}$.

Hence, there exists $k \geq 0$ such that ρ_2^k is admissible from a' and $\sup \tau - \varepsilon/2 \leq a' + \Delta(\rho_2^k) < \sup \tau$. Therefore, we have:

$$r(a') \rightarrow_{\rho_2^k} r(b') \text{ where } b' \in [\sup \tau - \varepsilon/2, \sup \tau).$$

By [Proposition 9\(c\)](#), we can scale the run ρ_3 so that it is admissible from $r(b')$ and reaches a value arbitrarily close to b' in state q . More formally, there exists $\beta \in (0, 1]$ such that

$$r(b') \rightarrow_{\beta\rho_3} q(b) \text{ where } b \in [b' - \varepsilon/2, \sup \tau).$$

We are done since $p(a) \rightarrow_{(1/2)\rho_1} r(a') \rightarrow_{\rho_2^k} r(b') \rightarrow_{\beta\rho_3} q(b)$ and $b \in [\sup \tau - \varepsilon, \sup \tau)$.

Case $\sup \tau = \infty$. We must show that we can reach arbitrarily large values. Let $b \geq a$. For all $\ell \geq 0$, the run $\rho'_\ell := (1/2)\rho_1 \rho_2^\ell$ is admissible from a , and such that $\Delta(\rho'_\ell) > 0$. Thus, there exists $\ell \geq 0$ such that $\Delta(\rho'_\ell) \geq (b - a) + \Delta^-(\rho_3)$. We are done since

$$a \rightarrow_{\rho'_\ell} b' \rightarrow_{\rho_3} b'' \text{ where } b' \geq b + \Delta^-(\rho_3) \text{ and } b'' \geq b. \quad \square$$

We show the characterisation of “ $a \in \text{Post}_{p,q}(a)$ ” stated, but left unproven, within the proof of [Proposition 14](#):

Proposition 40. Let $\text{Post}_{p,q}(a) \neq \emptyset$. It is the case that $a \in \text{Post}_{p,q}(a)$ iff at least one of these conditions holds:

- (a) there exists a path $\pi \in \text{Paths}_{p,q}$ whose transitions are all zero, i.e. $\Delta(\pi) = \Delta^+(\pi) = \Delta^-(\pi) = 0$;

(b) there exist $\pi \in \text{Paths}_{p,q}$ and indices i, j such that $\Delta(\pi_i) > 0$ and $\Delta(\pi_j) < 0$. If $a = \inf \tau$, then we also require that $\Delta(\pi_k) = 0$ for all $k < i$ and $k > j$. Similarly, if $a = \sup \tau$, then we also require $\Delta(\pi_k) = 0$ for all $k < j$ and $k > i$.

Proof. \Leftarrow) If (a) holds, then trivially $a \in \text{Post}_{p,q}(a)$. Assume (b) holds. Let $\rho := 1t_1 \cdots 1t_n$ where $\pi = t_1 \cdots t_n$. Suppose $a \notin \{\inf \tau, \sup \tau\}$. By Proposition 9(c), for all β small enough, it is the case that $a \rightarrow_{\beta\rho} a_\beta$, where $|a - a_\beta| < 1/2$. Let $p(a) = q_0(a_0), \dots, q_n(a_n) = q(a_\beta)$ be the sequence of configurations witnessing $a \rightarrow_{\beta\rho} a_\beta$. Since n is fixed we can choose $\beta < 1/2$ small enough so that $|a_i - a| < 1/2$ for all i . If $a_\beta > a$, then we enlarge the coefficient of t_j to $\alpha_j > \beta$ so that $(\alpha_j - \beta) \cdot \Delta(t_j) = a - a_\beta$. By the choice of β , we get an admissible run $\rho' := \beta t_1 \dots \beta t_{j-1} \alpha_j t_j \beta t_{j+1} \dots \beta t_n$ that satisfies $a \rightarrow_{\rho'} a$. If $a_\beta < a$, then we proceed analogously with index i .

It remains to prove the case where $a = \inf \tau$; the case where $a = \sup \tau$ is symmetric. By assumption, we have $\Delta(t_k) = 0$ for all $k < i$ and $k > j$. For the sake of simplicity, assume $\Delta(t_1) > 0$ and $\Delta(t_n) < 0$. Let $\alpha_1 \in (0, 1)$ be such that $\alpha_1 \cdot \Delta(t_1) < 1/2$. Let $\rho_1 := 1t_2 \cdots 1t_{n-1}$. By Proposition 9(c), there exists $\beta \in (0, 1]$ such that $\inf \tau \rightarrow_{\alpha_1 t_1 \beta \rho_1} \delta$, where $\delta < 1$. Since $\Delta(t_n) < 0$, there exists $\alpha_n \in (0, 1)$ such that $\alpha_n \cdot \Delta(t_n) = -\delta$. Thus, we have $p(\inf \tau) \rightarrow_{\alpha_1 t_1 \beta \rho_1 \alpha_n t_n} q(\inf \tau)$.

\Rightarrow) Let $p(a) \rightarrow_\rho q(a)$ and $\pi := \text{path}(\rho)$. Suppose (a) does not hold. If all transitions of π were positive, then we would obtain the contradiction $p(a) \rightarrow_\rho q(a')$ with $a' > a$. Similarly, all transitions cannot be negative. For the specific case where $a = \inf \tau$, observe that if the first nonzero transition is negative, then ρ cannot be admissible. Similarly, if the last nonzero transition is positive then $p(\inf \tau) \rightarrow_\rho q(\delta)$ for some $\delta > \inf \tau$. The reasoning for the case $a = \sup \tau$ is symmetric. \square

We show the characterisation of “ $c \in \text{Post}_{p,q}(a)$ ” stated, but left unproven, within the proof of Proposition 15:

Proposition 41. Let $\text{Post}_{p,q}(a) \neq \emptyset$, $b := \inf \overline{\text{Post}_{p,q}(a)}$ and $c := \sup \overline{\text{Post}_{p,q}(a)}$. If $b < a < c$ and $c \in \tau$, then $c \in \text{Post}_{p,q}(a)$ iff there is a state r and a path $\sigma \in \text{Paths}_{r,q}$ that satisfy $\Delta^+(\sigma) > 0$, $\Delta^-(\sigma) = 0$ and either of the following:

- (i) there exists a path $\sigma' \in \text{Paths}_{p,r}$ such that $|\sigma|, |\sigma'| \leq |Q|$, $\Delta^-(\sigma') = 0$ and $\Delta^+(\pi) \geq c - a$ where $\pi := \sigma' \sigma$;
- (ii) there exists a path $\sigma' \in \text{Paths}_{p,r}$ such that $|\sigma|, |\sigma'| \leq |Q|$ and $\Delta^+(\pi) > c - a$ where $\pi := \sigma' \sigma$;
- (iii) there is a positive (a, p, r) -admissible cycle θ .

Proof. \Rightarrow) Assume $c \in \text{Post}_{p,q}(a)$. There is a run ρ such that $p(a) \rightarrow_\rho q(c)$. Let ρ' be the run obtained from ρ by repeatedly removing a cycle θ with $\Delta^+(\theta) = 0$, until no further possible. Let $\pi := \text{path}(\rho')$. We have $\Delta^+(\pi) \geq \Delta(\rho') \geq \Delta(\rho) = c - a$. Since $c > a$, there is a maximal index i such that $\Delta(\pi_i) > 0$. Let $r := \text{in}(\pi_i)$, $\sigma' := \pi[1..i - 1]$ and $\sigma := \pi[i..|\pi|]$. Note that $\sigma' \in \text{Paths}_{p,r}$ and $\sigma \in \text{Paths}_{r,q}$. Moreover, $\Delta^+(\sigma) > 0$ holds by maximality of i . It must also be the case that $\Delta^-(\sigma) = 0$. Indeed, otherwise the last nonzero transition t of σ , and consequently of ρ , would be negative. Hence, this would contradict $c = \sup \overline{\text{Post}_{p,r}(a)}$ as we could reach values arbitrarily close to $c + \varepsilon$ for some $\varepsilon > 0$ by scaling t arbitrarily close to zero. Observe that if $\Delta^+(\pi) = c - a$, then $\Delta(\rho) = \Delta^+(\pi) = c - a$ which implies $\Delta^-(\rho) = \Delta^-(\pi) = 0$. Therefore, if $|\sigma|, |\sigma'| \leq |Q|$, we have shown (i) or (ii).

Otherwise, π is a nonsimple path. So, by our past cycle elimination, π contains a cycle θ with $\Delta^+(\theta) > 0$. Let us reorder θ into θ' so that the first transition t of θ' satisfies $\Delta(t) > 0$. We have $a \in \text{enab}(\text{Paths}_{p,\text{in}(t)})$ as state $\text{in}(t)$ occurs on the original run ρ that leads to state q . Moreover, $\text{Paths}_{\text{in}(t),r} \neq \emptyset$ holds by maximality of i . Thus, θ' is a positive (a, p, r) -admissible cycle. Hence, we have shown that (iii) holds.

\Leftarrow) If (i) holds, then $\Delta^+(\pi) = c - a$ or $\Delta^+(\pi) > c - a$. The latter case is subsumed by (ii), and in the former case we are done as $a \rightarrow_\pi c$ due to $\Delta^-(\pi) = 0$. If (iii) holds, then since θ is a positive (a, p, r) -admissible cycle — and hence (a, p, q) -admissible — Lemma 10(b) yields $\sup \overline{\text{Post}_{p,r}(a)} = c = \sup \tau$. Thus, there exists $\varepsilon \in [0, 1]$ such that $c - \varepsilon \in \text{Post}_{p,r}(a)$. By $\Delta^+(\sigma) \geq 1$ and $\Delta^-(\sigma) = 0$, we have

$$p(a) \rightarrow_* r(c - \varepsilon) \rightarrow_{\beta\sigma} q(c) \text{ where } \beta := \varepsilon / \Delta^+(\sigma).$$

If (ii) holds, then we proceed as follows. Recall that $b < a < c$. Therefore, $a \notin \{\inf \tau, \sup \tau\}$, since $\inf \tau \leq b$ and $c \leq \sup \tau$ by definition of b and c . Due to $a \notin \{\inf \tau, \sup \tau\}$, we can scale the negative transitions of σ' arbitrarily close to zero and scale its positive transitions so that either $a + \Delta^+(\sigma') - \varepsilon \in \text{Post}_{p,r}(a)$ or $\sup \tau - \varepsilon \in \text{Post}_{p,r}(a)$ for some $\varepsilon \in (0, 1]$. Since $\Delta^+(\sigma) \geq c - a - \Delta^+(\sigma') + 1$, $\Delta^+(\sigma) \geq 1$ and $\Delta^-(\sigma) = 0$, we can derive either $c \in \text{Post}_{r,q}(a)$ or $\sup \tau \in \text{Post}_{r,q}(a)$. As the latter implies $c = \sup \tau$, we are done proving the claim. \square

Lemma 17. It is the case that $p(a) \rightarrow_* q(b)$ in \mathcal{V} if and only if there is a path from $p(a)$ to $q(b)$ in \mathcal{G} .

Proof. We show the “if” direction first. Assume that $p(a) \rightarrow_\rho q(b)$ for some run ρ . Without loss of generality, we assume that no configuration repeats when starting at $p(a)$ with ρ ; otherwise, we can simply shorten ρ . Let $\sigma_1 \sigma_2 \cdots \sigma_n$ be the unique maximal decomposition of ρ into runs such that $\phi(\text{out}(\sigma_i)) \neq \mathbb{Q}$ for all $1 \leq i < n$. For ease of notation, let $q_i := \text{out}(\sigma_i)$. It holds that $p(a) \rightarrow_{\sigma_1} q_1(a_1) \rightarrow_{\sigma_2} q_2(a_2) \cdots \rightarrow_{\sigma_n} q(b)$, where $\phi(q_i) = [a_i, a_i]$ for all i . Since $\sigma_1 \sigma_2 \cdots \sigma_n$ is the maximal decomposition, $\phi(\text{out}(\sigma_i)) = \mathbb{Q}$ holds that for all $j < |\sigma_i|$.

Additionally, recall that p has no incoming edges and q has no outgoing edges. Hence, the following holds:

$$\begin{aligned} p(a) &\rightarrow_{\sigma_1} q_1(a_1) \text{ in } \mathcal{V}_{p,q_1}, \\ q_{i-1}(a_{i-1}) &\rightarrow_{\sigma_i} q_i(a_i) \text{ in } \mathcal{V}_{q_{i-1},q_i} \text{ for all } 1 < i < n, \\ q_{n-1}(a_{n-1}) &\rightarrow_{\sigma_n} q(b) \text{ in } \mathcal{V}_{q_{n-1},q}. \end{aligned}$$

We are done as $p(a)q_1(a_1) \cdots q_{n-1}(a_{n-1})q(b)$ is a path of \mathcal{G} .

It remains to show the ‘‘only if’’ direction. There exists a Suppose there is a path $p(a)q_1(a_1) \cdots q_{n-1}(a_{n-1})q(b)$ in \mathcal{G} . Note that if $p'(a') \rightarrow_* q'(b')$ in $\mathcal{V}_{p',q'}$, then by definition we also have $p'(a') \rightarrow_* q'(b')$ in \mathcal{V} . So we have

$$p(a) \rightarrow_* q_1(a_1) \rightarrow_* \cdots \rightarrow_* q_{n-1}(a_{n-1}) \rightarrow_* q(b) \text{ in } \mathcal{V}. \quad \square$$

B. Missing proofs from [section IV](#)

The following lemma will be useful to prove [Proposition 25](#).

Lemma 42. *Let $B, B' \in \mathcal{C}$ be such that $B \subseteq B'$ where B' is not a progressing extension of B . There is a bijection $f: \mathcal{I}(B) \rightarrow \mathcal{I}(B')$ s.t. $\phi_B(I) = \phi_{B'}(f(I))$ for all $I \in \mathcal{I}(B)$.*

Proof. Since $B \subseteq B'$, for all $I \in \mathcal{I}(B)$ there is a unique $f(I) \in \mathcal{I}(B')$ such that $I \subseteq f(I)$. We show that if f is not a bijection then it will contradict that $B \subseteq B'$ is not a progressing extension.

First, we prove that f is an injection. Suppose it is not and that $f(I) = f(J)$ for some $I, J \in \mathcal{I}(B)$. Then $\phi_B(I) \cup \phi_B(J) \subseteq \phi_{B'}(f(I))$. If there exists $\ell \in \phi_B(I) \cap \phi_B(J)$, then it must be the case that $\ell \notin I, \ell \notin J$ and $\ell \in f(I)$. This is a contradiction because the extension is progressing due to [\(3\)](#). Otherwise, there is $\ell \in \phi_B(I) \setminus \phi_B(J)$. Since $\phi_B(I) \subseteq \phi_B(I) \cup \phi_B(J) \subseteq \phi_{B'}(f(I))$ we get a contradiction because the extension is progressing due to [\(2\)](#).

Now, we prove that f is a surjection. Suppose it is not. Then there is an interval $I' \in \mathcal{I}(B')$ such that $f^{-1}(I') = \emptyset$. Thus $I' \cap B = \emptyset$, which is a contradiction because of [\(1\)](#).

To conclude, we note that by definition we have $\phi_B(I) \subseteq \phi_{B'}(f(I))$ for all $I \in \mathcal{I}(B)$. If the inclusion is strict for some I , then we get a contradiction because the extension is progressing due to [\(2\)](#). \square

Proposition 25. *For some n polynomially bounded in $|Q|$, at least one of the following holds: $S_n = S_{n+1}$,*

- *there is a positively expanding cycle ρ from S_n , or*
- *there is a negatively expanding cycle ρ from S_n .*

Moreover, it can be determined in time $|Q|^{\mathcal{O}(1)}$ whether the second or third case hold, and then ρ and its witnessing configurations can be computed in time $|Q|^{\mathcal{O}(1)}$.

Proof. The value of the polynomially bounded number n will be determined by the proof. By definition, we have $S_i \preceq S_{i+1}$ for all $i \in \mathbb{N}$. By [Lemma 29](#), there is a polynomial number of indices i (w.r.t. $|Q|$) such that $S_i(q) \subseteq S_{i+1}(q)$ is a progressing extension for some $q \in Q$. Thus, there exists an index j polynomial in $|Q|$ such that the extensions $S_i(q) \subseteq S_{i+1}(q)$ are not progressing for all $j \leq i \leq j+k$, where k is sufficiently large (but polynomially bounded in $|Q|$). To simplify the notation we will assume that $j=0$ and consider $S_0 \preceq S_1 \preceq \cdots \preceq S_k$.

By [Lemma 42](#), there is a bijection $f_i: \mathcal{I}(S_i(q)) \rightarrow \mathcal{I}(S_{i+1}(q))$ for every $q \in Q$ and $0 \leq i < k$. Thus, for every $q \in Q$, the sets $\mathcal{I}(S_1(q)), \dots, \mathcal{I}(S_k(q))$ have the same number m_q of intervals. By [Lemma 21](#), m_q is polynomially bounded in $|Q|$. Let us denote the intervals $I_1^{q,i}, \dots, I_{m_q}^{q,i}$ where $f(I_j^{q,i}) = I_j^{q,i+1}$ for all $0 \leq i < k, q \in Q$ and $1 \leq j \leq m_q$. By [Lemma 42](#), $\phi_{S_i(q)}(I_j^{q,i}) = \phi_{S_{i+1}(q)}(I_j^{q,i+1})$. Since $S_i \preceq S_{i+1}$, we conclude that $I_j^{q,1} \subseteq I_j^{q,2} \subseteq \cdots \subseteq I_j^{q,k}$ for all $q \in Q$ and $1 \leq j \leq m_q$.

Recall that T is the set of transitions of the guarded COCA. Let $\text{active}(q, i, j) := \{t \in T \mid I_j^{q,i} \cap \text{enab}(t) \neq \emptyset\}$. Since $I_j^{q,i} \subseteq I_j^{q,i+1}$, we have $\text{active}(q, i, j) \subseteq \text{active}(q, i+1, j)$, and a strict inclusion can occur at most $|T|$ times. Since it is polynomially bounded in $|Q|$ and $|T|$, we can assume that $\text{active}(q, i, j) = \text{active}(q, i+1, j)$ for all $q \in Q, 0 \leq i < k$, and $1 \leq j \leq m_q$. Suppose $S_i \neq S_{i+1}$ for every $0 \leq i < k$. Then we prove that there is a positively or negatively expanding cycle from S_0 . Since $S_i \neq S_{i+1}$, there exists $v_k \in S_k(q_k)$ for some $q_k \in Q$ such that $v_k \notin S_{k-1}(q_k)$. Tracing back, by definition of Succ, we can find a sequence of transitions t_1, \dots, t_k , scalars $\alpha_1, \dots, \alpha_k \in (0, 1]$, and configurations $q_0(v_0), \dots, q_k(v_k)$ such that: $v_i \in S_i(q_i), v_i \notin S_{i-1}(q_i)$, and $q_{i-1}(v_{i-1}) \rightarrow_{\alpha_i t_i} q_i(v_i)$ for all $0 < i \leq k$. We show that an infix of this sequence defines a positively or negatively expanding cycle.

Let j_k be the unique index with $v_k \in I_{j_k}^{q_k, k}$. By definition, we have $v_k \notin I_{j_k}^{q_k, k-1}$. If $v_k \geq \sup(I_{j_k}^{q_k, k-1})$, then we construct a positively expanding cycle, and otherwise we construct a negatively expanding one. We will prove only the former case; the latter case follows the same steps. We claim that $v_i \geq \sup(I_{j_i}^{q_i, i-1})$ for all $0 < i \leq k$.

Let us argue that the claim allows us to conclude. For a large enough k , we can find an infix $\rho := q_a(v_a), \dots, q_b(v_b)$ such that: $a < b, q_a = q_b$, and $j_a = j_b$. Thus, $v_b \geq \sup(I_{j_b}^{q_b, b-1}) \geq \sup(I_{j_a}^{q_a, a}) \geq v_a$. Since $v_b \in I_{j_b}^{q_b, b}$ and $v_a \notin I_{j_b}^{q_b, b}$, we obtain $\Delta(\rho) > 0$. The remaining conditions of the positively expanding cycle follow directly from the definition.

It remains to prove the claim, i.e. that $v_i \geq \sup(I_{j_i}^{q_i, i-1})$ for every $0 < i \leq k$. We proceed by induction going from $i = k$ down to $i = 1$. The base case follows by assumption. For the inductive step, towards a contradiction, suppose that $v_i \leq \inf(I_{j_i}^{q_i, i-1})$ and $v_i \notin I_{j_i}^{q_i, i-1}$. Recall that $\text{active}(q_i, i-1, j_i) = \text{active}(q_i, i, j_i)$. Since $v_i \rightarrow_{\alpha_{i+1}t_{i+1}} v_{i+1}$, there exists $\bar{v} \in I_{j_i}^{q_i, i-1}$ and $\beta \in (0, 1]$ such that $\bar{v} \rightarrow_{\beta t_{i+1}} \bar{w}$ for some $\bar{w} \in S_i(q_{i+1})$.

We show that $v_{i+1} < \bar{w}$. If $v_{i+1} = \bar{w}$, then $v_{i+1} \in S_i(q_{i+1})$, which contradicts the definition of v_{i+1} . Suppose that $v_{i+1} \neq \bar{w}$ for any choice of β . Let $t_{i+1} = (q_i, z_{i+1}, q_i + 1)$. Since $v_i \leq \inf(I_{j_i}^{q_i, i-1})$, notice that $\bar{v} + \alpha_{i+1}z_{i+1} > v_i + \alpha_{i+1}z_{i+1} = v_{i+1}$. Thus, it must be the case that $v_{i+1} < \bar{w}$.

We prove that $\bar{w} \in I_{j_{i+1}}^{q_{i+1}, i+1}$. Recall that $v_i, \bar{v} \in I_{j_i}^{q_i, i}$ and thus $[v_i, \bar{v}] \subseteq I_{j_i}^{q_i, i}$. Also $[v_i + \alpha_{i+1}z_{i+1}, \bar{v} + \beta z_{i+1}] = [v_{i+1}, \bar{w}]$. Thus, for every $v_{i+1} \leq w' \leq \bar{w}$ there exists $v_i \leq v' \leq \bar{v}$ and $\gamma \in (0, 1]$ such that $v' \rightarrow_{\gamma t_{i+1}} w'$. Thus, \bar{w} and v_{i+1} belong to the same interval in $S_{i+1}(q_{i+1})$ as required.

Since $\bar{w} \in I_{j_{i+1}}^{q_{i+1}, i+1}$ and $\bar{w} \in S_i(q_{i+1})$, we have $\bar{w} \in I_{j_{i+1}}^{q_{i+1}, i}$. We have reached a contradiction, since by the inductive hypothesis we have:

$$v_{i+1} \geq \sup(I_{j_{i+1}}^{q_{i+1}, i}).$$

Observe that since Succ is easily computable in polynomial time, one can also find $q_0(v_0), \dots, q_k(v_k)$ and their corresponding intervals in polynomial time. \square

C. Missing formulas from [section V](#)

Intersection of intervals: Let χ_x be the auxiliary function defined by $\chi_x(y, z) := y$ if $x \neq 2$ and z otherwise. Let $I = (b, t, \perp, \top) \in \mathcal{E}$ and $I' = (b', t', \perp', \top') \in \mathcal{E}$. The intersection $I \cap I' := (b'', t'', \perp'', \top'')$ can be defined by

$$\begin{aligned} b'' &:= \max(\chi_{\perp}(b, b'), \chi_{\perp'}(b', b)), \\ t'' &:= \min(\chi_{\top}(t, t'), \chi_{\top'}(t', t)), \\ \perp'' &:= \begin{cases} \chi_{\perp}(\perp, \perp') & \text{if } b > b', \\ \chi_{\perp'}(\perp', \perp) & \text{if } b < b', \\ \min(\perp, \perp') & \text{otherwise,} \end{cases} \\ \top'' &:= \begin{cases} \chi_{\top}(\top, \top') & \text{if } t < t', \\ \chi_{\top'}(\top', \top) & \text{if } t > t', \\ \min(\top, \top') & \text{otherwise.} \end{cases} \end{aligned}$$

To explain the above, let us first look at the case where both intervals are bounded, that is, $\perp, \top, \perp', \top' \in \{0, 1\}$. Each invocation of χ will return its first argument. Let us examine the lower endpoint. It is the case that $b'' = \max(b, b')$. As expected, the intersection operation uses the larger lower endpoint among the two intervals. Further, whether the lower endpoint of the interval is included depends on whether the value that was used was included in the interval it originated from, i.e. consider $(3, 5] \cap [4, 6] = [4, 5]$. The only remaining nuance is when both intervals have the same endpoint, then it is only included if it is included in both intervals. This is why we choose $\min(\perp, \perp')$ in the case where $b = b'$.

Now, let us consider the case where either interval may be unbounded. Then it is not sufficient to compare endpoints, e.g. the encoding $(0, 1, 2, 2)$ does not encode an interval with endpoints 0 and 1, but rather $(-\infty, +\infty)$. Intuitively, the χ function is used in order to handle such unbounded intervals. When intersecting two intervals, where one interval has no finite lower endpoint, while the other does, the lower endpoint should only depend on the interval that indeed has a lower endpoint. So for determining the lower endpoint and whether it is included, the χ function will check whether the interval is bounded, and only use its endpoint if it exists. Otherwise, it defaults to the endpoint of the other interval.

The reasoning behind the upper endpoint is symmetric.

D. Missing gadgets from the proof of [Theorem 39](#)

Recall that the proof of [Theorem 39](#) gives the gadgets of \mathcal{P} , but not of \mathcal{P}' . [Figure 4](#) depicts the missing gadgets of \mathcal{P}' .




Fig. 4. Gadgets used in the reduction from 3-SAT to parametric COCA reachability where parameters occur only on updates. *Left*: Gadget for variable x_i ; *Right*: Gadget used for clause $C_j = (x_1 \vee x_2 \vee \neg x_4)$.