

THE LEAST GENERATING DEGREE OF SYMBOLIC POWERS OF IDEALS OF FERMAT CONFIGURATION OF POINTS

THÁI THÀNH NGUYỄN

ABSTRACT. Let $n \geq 2$ be an integer and consider the defining ideal of the Fermat configuration of points in \mathbb{P}^2 : $I_n = (x(y^n - z^n), y(z^n - x^n), z(x^n - y^n)) \subset R = \mathbb{C}[x, y, z]$. In this paper, we compute explicitly the least degree of generators of its symbolic powers in all unknown cases. As direct applications, we easily verify Chudnovsky's Conjecture, Demailly's Conjecture and Harbourne-Huneke Containment problem as well as calculating explicitly the Waldschmidt constant and (asymptotic) resurgence number.

1. INTRODUCTION

Let $n \geq 2$ be an integer and consider the *Fermat ideal*

$$I_n = (x(y^n - z^n), y(z^n - x^n), z(x^n - y^n)) \subset R = \mathbb{C}[x, y, z]$$

This ideal corresponds to Fermat arrangement of lines (or Ceva arrangement in some literature) in \mathbb{P}^2 , more precisely, the variety of I_n is a reduced set of $n^2 + 3$ points in \mathbb{P}^2 [HS15], where n^2 of these points form the intersection locus of the pencil of curves spanned by $x^n - y^n$ and $x^n - z^n$, while the other 3 are the coordinate points $[1 : 0 : 0]$, $[0 : 1 : 0]$ and $[0 : 0 : 1]$. This set of points is said to be the Fermat configuration of points, justifying the terminology of Fermat ideal. Fermat ideal has attracted a lot of attention recently in commutative algebra research since it appeared as the first example of the non-containment between the third symbolic power and the second ordinary power of a defining ideal of a set of points in \mathbb{P}^2 , in the work of Dumnicki, Szemberg and Tutaj-Gasńska [DSTG13] (when $n = 3$) and were generalized by Harbourne and Seceleanu [HS15]. It is worth to emphasize that this is a quite surprising relation between ordinary and symbolic powers of ideal of points in \mathbb{P}^2 , since I^2 always contains $I^{(3)}$ where I is the ideal of a general set of points [BH10].

Since then, much has become known about Fermat ideals for $n \geq 3$. Fermat ideals can also be thought of the ideals determining the singular loci of the arrangements of lines given by the monomial groups $G(n, n, 3)$, see [DS20] or [Szp19], *Waldschmidt constant* and *(asymptotic) resurgence number* of Fermat ideals have been computed in [DHN⁺15] for $n \geq 3$. Nagel and Seceleanu in [NS16] studied Rees algebra and symbolic Rees Algebra of Fermat ideals as well as the minimal generators, minimal free resolutions of all their ordinary powers and many symbolic powers. Specifically, it was shown that symbolic Rees algebra of I_n is Noetherian; Castelnuovo-Mumford regularity of powers of I_n and their reduction ideals were provided.

2010 *Mathematics Subject Classification.* 14N20, 13F20, 14C20.

Key words and phrases. Fermat Ideals, Fermat Points Configuration, Resurgence Number, Waldschmidt Constant, Ideals of Points, Symbolic Powers, Containment problem, Stable Harbourne–Huneke Conjecture, Interpolation Problem.

In term of minimal generators, when $n \geq 3$, they provided the minimal generators for all ordinary powers as well all multiple of n symbolic powers. All known results are:

- $\alpha(I_n^{(3k)}) = 3nk$ for all $k \geq 1$ by [DHN⁺15, Theorem 2.1].
- $\alpha(I_n^{(nk)}) = n^2k$ for all $k \geq 1$ by [NS16, Theorem 3.6].
- $\alpha(I_3^{(3m+2)}) = 9m + 8$ for all $m \geq 0$ by [MSS18, Example 4.4].

In this paper, we will compute explicitly the least degree of generators of symbolic powers of I_n for *all* remaining cases as a contribution to provide a more complete picture of Fermat ideals. Our results are the following:

Theorem 1.1. *Let $n \geq 2$ be an integer and $I_n = (x(y^n - z^n), y(z^n - x^n), z(x^n - y^n))$ in $\mathbb{C}[x, y, z]$ be a Fermat ideal. Then:*

- (1) For $n \geq 5$, $\alpha(I_n^{(m)}) = mn$ for all $m \geq 3$.
- (2) $\alpha(I_4^{(m)}) = 4m$ for all $m \geq 3$ and $m \neq 5$.
- (3) $\alpha(I_3^{(3k+1)}) = 9k + 4$ for all $k \geq 1$.
- (4) For $n \geq 2$, $\alpha(I_n^{(2)}) = \alpha(I_n^2) = 2(n + 1)$.
- (5) $\alpha(I_2^{(2k)}) = 5k$ for all $k \geq 2$.
- (6) $\alpha(I_2^{(2k+1)}) = 5k + 3$ for all $k \geq 0$.

The above results give complete answer to the question of the least generating degree all symbolic powers of I_n . That includes the calculation for the ideal I_2 , which is less considered in the aforementioned works. Note that the ideal I_2 is the ideal determining the singular locus of the arrangement of lines given by the pseudoreflection group D_3 , see [DS20].

The *containment problem* for ideal I is to determine the set of pairs (m, r) for which $I^{(m)} \subseteq I^r$. The deep results in [ELS01] and [HH02] show that $I^{(m)} \subseteq I^r$ whenever $m \geq Nr$. In order to characterize the pairs (r, m) numerically, the *resurgence* $\rho(I)$ is introduced in [BH10], and the *asymptotic resurgence* $\widehat{\rho}(I)$ is introduced in [GHVT13]. It is known that for $n \geq 3$, $\rho(I_n) = \frac{3}{2}$, $\widehat{\rho}(I_n) = \frac{n+1}{n}$ and $\widehat{\alpha}(I_n) = n$ by [DHN⁺15, Theorem 2.1], where $\widehat{\alpha}(I)$ is the *Waldschmidt constant* of I . We will compute the Waldschmidt constant and (asymptotic) resurgence number of I_2 .

Theorem 1.2. *For the ideal I_2 , the Waldschmidt constant is $\widehat{\alpha}(I_2) = \frac{5}{2}$; the resurgence number and asymptotic resurgence number are $\rho(I_2) = \widehat{\rho}(I_2) = \frac{6}{5}$.*

In an effort to improve the containment $I^{(m)} \subseteq I^r$ for $m \geq Nr$ as well as to deduce Chudnovsky's Conjecture, Harbourne and Huneke in [HH13] conjectured that the defining ideal I for any set of points in \mathbb{P}^N satisfies some stronger containment, namely, $I^{(Nm)} \subseteq \mathfrak{m}^{m(N-1)} I^m$ and $I^{(Nm-N+1)} \subseteq \mathfrak{m}^{(m-1)(N-1)} I^m$, for all $m \geq 1$. An interesting fact about the Fermat ideals is the verification and failure of the containment can be checked purely by their numerical invariants including the least degree of generators of symbolic powers, the regularity, or the resurgence number and the maximal degree of generators. As a consequence of the above computations, we will easily deduce that all Fermat ideals verify *Harbourne-Huneke Containment*, *stable Harbourne Containment* and some stronger containment. In [BGHN20a, Example 3.5], we showed the stronger containment (which implies both Harbourne-Huneke

containment) $I_n^{(2r-2)} \subseteq \mathfrak{m}^r I_n^r$ for $n \geq 3$ and for all $r \gg 0$. Here, we show the containment for all possible cases (and show that they fail for the rest).

Corollary 1.3. *For every $n \geq 2$, Fermat configuration ideal verifies the following containment:*

(1) *Harbourne-Huneke containment (see [HH13, Conjecture 2.1])*

$$I_n^{(2r)} \subseteq \mathfrak{m}^r I_n^r, \quad \forall r$$

(2) *Harbourne-Huneke containment (see [HH13, Conjecture 2.1])*

$$I_n^{(2r-1)} \subseteq \mathfrak{m}^{r-1} I_n^r$$

$\forall r \geq 3$ if $n \geq 3$ and $\forall r \geq 1$ if $n = 2$.

(3) *A stronger containment $I_n^{(2r-2)} \subseteq \mathfrak{m}^r I_n^r$, $\forall r \geq 5$*

This work can also be thought of a modest contribution to the theory of Hermite interpolation. Specifically, given a set of points $Z = \{P_1, \dots, P_s\}$ in a projective space \mathbb{P}^n and positive integers m_1, \dots, m_s , a fundamental problem in the theory of Hermite interpolation is to determine the least degree of a homogeneous polynomial that vanishes to order m_i at the point P_i for every $i = 1, \dots, s$. This is a very hard problem even when $m_1 = \dots = m_s$. In this case when all m_i are the same, thanks to the Zariski-Nagata theorem, the above interpolation problem is the same as asking for the least degree of a nonzero homogeneous polynomial in the k -th symbolic power of the defining ideal of Z where $k = m_1 = \dots = m_s$. Hence, this work combined with many previous works provide a complete answer to the above question for Z be a Fermat configuration of points in \mathbb{P}^2 .

n	2			3				4			$n \geq 5$	
m	2	$2k$	$2k+1$	2	$3k$	$3k+1$	$3k+2$	2	5	$\geq 3, \neq 5$	2	≥ 3
$\alpha(I_n^{(m)})$	6	$5k$	$5k+3$	8	$9k$	$9k+4$	$9k+8$	10	21	$4m$	$2(n+1)$	nm
$\widehat{\alpha}(I_n)$	$5/2$			3				4			n	
$\rho(I_n) =$ $\widehat{\rho}(I_n)$	$\frac{6}{5}$							$\frac{3}{2}$				

TABLE 1. Least degree of generators and other invariants related to symbolic powers of I_n

In order to understand the generating degrees of symbolic powers of I_n , we discuss the maximal degree of a set of minimal generators of symbolic powers of I_n , denoted by $\omega(I_n^{(m)})$. From the description of generating sets in [NS16], it can be seen that for $n \geq 3$, $\omega(I_n^{(m)}) = m(n+1)$, for $m = kn$ or $m = n - 1$. By relating with another invariant $\beta(I_n^{(m)})$ that is defined in [HH13, Definition 2.2], we show that $\beta(J) \leq \omega(J)$ for an ideal of points J in general and use this to show that for $n \geq 3$, $\omega(I_n^{(m)}) = m(n+1)$ for $m \geq n^2 - 3n + 2$.

It is worth to remark that in the works [MS17], [MS18], Malara and Szpond also studied the generalization of Fermat configuration in higher dimension to provide more counterexamples to the containment $I^{(3)} \subseteq I^2$. It turns out that these ideals share some similar properties to Fermat ideals. We will investigate these ideals in 3-dimensional space with

the same questions in the continuation paper [Ngu20] in order to keep this paper more concise and focused. We work over the field of complex numbers but our results hold over any algebraically closed field of characteristic 0.

Acknowledgements. The author would like to thank his advisor, Dr. Tàì Hà, for introducing him this subject and giving many helpful suggestions and comments.

2. PRELIMINARY

Let $R = \mathbb{C}[x_0, \dots, x_N]$ be the homogeneous coordinate ring of \mathbb{P}^N , and let \mathfrak{m} be its maximal homogeneous ideal. For a homogeneous ideal $I \subseteq R$, let $\alpha(I)$ denote the least degree of a homogeneous polynomial in I , and let

$$I^{(m)} := \bigcap_{\mathfrak{p} \in \text{Ass}(R/I)} I^m R_{\mathfrak{p}} \cap R$$

denote its m -th *symbolic power*.

Geometrically, given a set of distinct points $\mathbb{X} \subseteq \mathbb{P}^N$ and an integer $m \geq 1$, by the Zariski–Nagata Theorem [Zar49, Nag62, EH79] (cf. [DDSG⁺18, Proposition 2.14]), the least degree of a homogeneous polynomial in the homogeneous coordinate ring $\mathbb{C}[x_0, \dots, x_N]$ that vanishes at each point in \mathbb{X} of order at least m is $\alpha(I_{\mathbb{X}}^{(m)})$, where $I_{\mathbb{X}} \subseteq \mathbb{C}[x_0, \dots, x_N]$ is the defining ideal of \mathbb{X} .

The *Waldschmidt constant* of I is defined to be the limit and turned out to be the infimum

$$\widehat{\alpha}(I) := \lim_{m \rightarrow \infty} \frac{\alpha(I^{(m)})}{m} = \inf_{m \rightarrow \infty} \frac{\alpha(I^{(m)})}{m}$$

There is a tight connection between Waldschmidt constant and an algebraic manifestation of the Seshadri constant, especially for a set of general points, see [BDRH⁺09, Section 8].

In studying the lower bound for the least degree of a homogeneous polynomial vanishing at a given set of points in \mathbb{P}^N with a prescribed order, Chudnovsky [Chu81] made the following conjecture.

Conjecture 2.1 (Chudnovsky). Let I be the defining ideal of a set of points $\mathbb{X} \subseteq \mathbb{P}_{\mathbb{C}}^N$. Then, for all $n \geq 1$,

$$\frac{\alpha(I^{(n)})}{n} \geq \frac{\alpha(I) + N - 1}{N}$$

Chudnovsky’s Conjecture has been investigated extensively, for example, in [EV83, BH10, HH13, GHM13, Dum15, DTG17, FMX18, BGHN20a]. Recently, the conjecture was proved for a *very general* set of points [DTG17, FMX18] and for a *general* set of sufficiently many points [BGHN20a]. The conjecture was also generalized by Demailly [Dem82] as follows.

Conjecture 2.2 (Demailly). Let I be the defining ideal of a set of points $\mathbb{X} \subseteq \mathbb{P}_{\mathbb{C}}^N$ and let $m \in \mathbb{N}$ be any integer. Then, for all $n \geq 1$,

$$\frac{\alpha(I^{(n)})}{n} \geq \frac{\alpha(I^{(m)}) + N - 1}{m + N - 1}$$

Demailly's Conjecture for $N = 2$ was proved by Esnault and Viehweg [EV83]. Recent work of Malara, Szemberg and Szpond [MSS18], and of Chang and Jow [CJ20], showed that for a fixed integer m , Demailly's Conjecture holds for a *very general* set of sufficiently many points and for a *general* set of k^N points. In [BGHN20b], the results was extended for a *general* set of sufficiently many points.

The *containment problem* for ideal I is to determine the set S_I of pairs (r, m) for which $I^{(m)} \subseteq I^r$. The deep results in [ELS01] and [HH02] show that $I^{(m)} \subseteq I^r$ whenever $m \geq Nr$, hence, $\{(m, r) : m \geq Nr\} \subseteq S_I$. In order to characterize S_I numerically, the *resurgence number* $\rho(I)$ is introduced in [BH10] as

$$\rho(I) := \sup\left\{\frac{m}{r} : I^{(m)} \not\subseteq I^r\right\}$$

and the *asymptotic resurgence number* $\widehat{\rho}(I)$ is introduced in [GHVT13] as

$$\widehat{\rho}(I) := \sup\left\{\frac{m}{r} : I^{(mt)} \not\subseteq I^{rt} \text{ for } t \gg 0\right\}$$

There are only few cases for which S_I is known completely or the resurgence number have been determined. In general, $\rho(I)$ and $\widehat{\rho}(I)$ are different.

In an effort to improve the containment $I^{(m)} \subseteq I^r$ for $m \geq Nr$, Harbourne and Huneke in [HH13] conjectured that the defining ideal I for any set of points in \mathbb{P}^N satisfies some stronger containment, namely, $I^{(Nm)} \subseteq \mathfrak{m}^{m(N-1)}I^m$ and $I^{(Nm-N+1)} \subseteq \mathfrak{m}^{(m-1)(N-1)}I^m$, for all $m \geq 1$. Knowing these containment clearly helps us to know the set S_I and the numbers $\rho(I), \widehat{\rho}(I)$. Conversely, knowledge about $S_I, \rho(I), \widehat{\rho}(I)$ can be helpful to prove the containment. One useful result that we will use lately is the following: If $\frac{m}{r} > \rho(I)$ then by definition $I^{(m)} \subseteq I^r$ and suppose in addition, $\alpha(I^{(m)}) \geq a + \omega(I^r)$ for some integer a , where $\omega(I)$ is the maximum degree of generators in a set of minimal generators of I , then $I^{(m)} \subseteq \mathfrak{m}^a I^r$. We refer interested readers to [CHHVT20] for more information about the Waldschmidt constant, resurgence number, containment between symbolic and ordinary powers of ideals.

3. FERMAT IDEALS FOR $n \geq 3$

In this section, we focus on the Fermat ideals $I_n = (x(y^n - z^n), y(z^n - x^n), z(x^n - y^n))$ for $n \geq 3$. Let us first recall some known results about degree of generators, Waldschmidt constants and (asymptotic) resurgence numbers of I_n .

Known Results:

- $\alpha(I_n^{(3k)}) = 3nk$ for all $k \geq 1$ by [DHN⁺15, Theorem 2.1].
- $\alpha(I_n^{(nk)}) = n^2k$ for all $k \geq 1$ by [NS16, Theorem 3.6].
- $\alpha(I_3^{(3m+2)}) = 9m + 8$ for all $m \geq 0$ by [MSS18, Example 4.4].
- $\rho(I_n) = \frac{3}{2}$ and $\widehat{\rho}(I_n) = \frac{n+1}{n}$ by [DHN⁺15, Theorem 2.1]

Let $f = y^n - z^n, g = z^n - x^n, h = x^n - y^n$. Then

$$I_n = (xf, yg, zh) = (f, g) \cap (x, y) \cap (y, z) \cap (z, x)$$

It is well-known that, since f and g form a regular sequence, for any $m \geq 1$ we have

$$I_n^{(m)} = (f, g)^m \cap (x, y)^m \cap (y, z)^m \cap (z, x)^m$$

We will use this description to compute explicitly the least degree of generators of $I_n^{(m)}$ in all remaining unknown cases, provided the knowledge about the Waldschmidt constant of I_n . This is our main strategy in this paper and the continuation paper [Ngu20], we study a subsequence of $\alpha(I^{(m)})$, which gives us information about $\widehat{\alpha}(I)$, then use this to calculate other $\alpha(I^{(m)})$.

The following 4 theorems provide us all remaining unknown least degree of generators of symbolic powers of I_n :

Theorem 3.1. *For $n \geq 5$, we have*

$$\alpha(I_n^{(m)}) = nm$$

for all $m \geq 3$.

Proof. First, for $3 \leq m \leq n$, we observe that

$$fgh(f, g)^{m-3} \subseteq I_n^{(m)} = (f, g)^m \cap (x, y)^m \cap (y, z)^m \cap (z, x)^m$$

In fact, it is clear that $fgh(f, g)^{m-3} \subseteq (f, g)^m$. Also, it is clear that the sum of degree w.r.t x and degree w.r.t y of any monomials of fgh is at least $n \geq m$ so $fgh \in (x, y)^m$. Similarly, $fgh \in (y, z)^m \cap (z, x)^m$.

Thus, for $3 \leq m \leq n$, $\alpha(I_n^{(m)}) \leq nm$. Since $\widehat{\alpha}(I_n) = n$, we have $\alpha(I_n^{(m)}) \geq nm$, therefore $\alpha(I_n^{(m)}) = nm$.

Now for any $k \geq 2$, we claim that for $0 \leq a \leq n - 1$

$$(fgh)^k (f, g)^{k(n-3)-a} \subseteq I^{(kn-a)}$$

The argument is identical to that of the above, the only notice here is that for $n \geq 5$,

$$k(n-3) \geq 2(n-3) \geq n-1 \geq a$$

Thus, for all $k \geq 2$ and $0 \leq a \leq n - 1$,

$$\alpha(I_n^{(kn-a)}) \leq (kn-a)n$$

and hence $\alpha(I_n^{(kn-a)}) = (kn-a)n$. Since for $m > n$, there are unique $k \geq 2$ and $0 \leq a \leq n-1$ such that $m = kn - a$, we have that $\alpha(I_n^{(m)}) = mn$ for all $m > n$. \square

Theorem 3.2. *When $n = 4$, for all $m \geq 3$ but $m \neq 5$,*

$$\alpha(I_4^{(m)}) = 4m$$

Proof. With the same argument we have that:

- (1) For $3 \leq m \leq 4$, $fgh(f, g)^{m-3} \subseteq I_4^{(m)}$. The statement is true for $m = 3, 4$.

(2) For $k = 2$ and $0 \leq a \leq 2$,

$$(fgh)^k(f, g)^{k(n-3)-a} \subseteq I_4^{(kn-a)}$$

i.e.,

$$(fgh)^2(f, g)^{2-a} \subseteq I_4^{(8-a)}$$

for $0 \leq a \leq 2$. Thus, the statement is true for $m = 6, 7, 8$.

(3) For $k \geq 3$, we have $k(4-3) \geq 3$ so for $0 \leq a \leq 3$,

$$(fgh)^k(f, g)^{k(n-3)-a} \subseteq I_4^{(kn-a)}$$

therefore, the statement is true for all $m \geq 9$.

In case 2, the argument does not work for $m = 5$. However, by the same argument, we can check that $zf^2gh^2 \in I_4^{(5)}$. In fact, Macaulay2 computation [GS] suggests that in this only case $\alpha(I_4^{(5)}) = 21 \neq 20$. \square

Theorem 3.3. For all $n \geq 3$, $\alpha(I_n^{(2)}) = 2(n+1)$.

Proof. We know that $\alpha(I_n^{(2)}) \leq \alpha(I_n^2) = 2(n+1)$. Now suppose that $\alpha(I_n^{(2)}) \leq 2n+1$. Then there is a divisor D of degree $2n+1$ vanishing to order at least 2 at every P_i in the Fermat configuration. Recall that the Fermat configuration consists of n^2+3 points which are all points having each coordinate equal to a n^{th} root of 1 and the points $[0:0:1]$, $[0:1:0]$, $[1:0:0]$. Furthermore, these points are intersections of $3n$ lines L_j which have equations:

$$x - \epsilon^k y = 0, y - \epsilon^k z = 0, z - \epsilon^k x = 0$$

for $k = 0, 1, \dots, n-1$. Each of these lines contains exactly $n+1$ points P_i , and each of the points $[0:0:1]$, $[0:1:0]$, $[1:0:0]$ is on exactly n lines while each of the other n^2 points is on exactly 3 lines.

Since the intersection of D and any line L_j consists of $n+1$ points to order at least 2, by Bezout theorem, each L_j is a component of D because $\deg(D) \cdot \deg(L_j) = 2n+1 < 2(n+1)$. This is a contradiction since there are $3n$ lines and $3n > 2n+1 = \deg(D)$ when $n \geq 3$. Therefore, $\alpha(I_n^{(2)}) = 2(n+1)$ for all $n \geq 3$. \square

It is known that $\alpha(I_3^{(3m+2)}) = 9m+8$ for all $m \geq 0$ from [MSS18, Example 4.4] and $\alpha(I_3^{(3m)}) = 9m$ for $m \geq 1$ from [DHN⁺15, Theorem 2.1] when $n = 3$. Now we compute $\alpha(I_3^{(3m+1)})$.

Theorem 3.4. $\alpha(I_3^{(3m+1)}) = 9m+4$ for all m .

Proof. Proceed by the argument in [MSS18, Example 4.4], suppose that there is $m \geq 1$ such that $\alpha(I_3^{(3m+1)}) \leq 9m+3$. Then there is a divisor D of degree $9m+3$ vanishing to order at least $3m+1$ at every point of 12 points P_i in the Fermat configuration. Intersecting D with any of the 9 lines L_j , since each lines L_j contains exactly 4 points and $9m+3 < 4(3m+1)$, we conclude by Bezout Theorem that each L_j is a component of D . Hence, there exists a divisor $D' = D - \sum_{j=1}^9 L_j$ of degree $9(m-1)+3$ vanishing to order at least $3(m-1)+1$ at every point of P_i . Repeating this argument m times we get a contradiction with $\alpha(I_3) = 4$. Thus $\alpha(I_3^{(3m+1)}) \geq 9m+4$ for all m .

On the other hand, by degree argument, $f^m g^m h^{m+1} z \in I_3^{(3m+1)}$ so $\alpha(I_3^{(3m+1)}) \leq 9m + 4$ for all m . Therefore, $\alpha(I_3^{(3m+1)}) = 9m + 4$ for all m . \square

Example 3.5. It is worth to point out that the first immediate application of the above calculations combining with the already known cases is the verification of Chudnovsky's Conjecture and Demailly's Conjecture, although the general case is already known from [EV83]. For any $n \geq 3$, Fermat ideals verify

- (1) Chudnovsky's Conjecture $\widehat{\alpha}(I_n) \geq \frac{\alpha(I_n) + 1}{2}$.
- (2) Demailly's Conjecture $\widehat{\alpha}(I_n) \geq \frac{\alpha(I_n^{(m)}) + 1}{m + 1}$ for all m .

Proof. Directly from the formulae of $\widehat{\alpha}(I_n)$ and $\alpha(I_n^{(m)})$. \square

The following containment are also direct consequences of the above calculations about $\alpha(I_n^{(m)})$. Note that these containment (and in fact, the stable version of them, i.e, the containment for $r \gg 0$ imply Chudnovsky's Conjecture). First, in [BGHN20a, Example 3.5], we showed the stronger containment (which implies both Harbourne-Huneke containment)

$$I_n^{(2r-2)} \subseteq \mathfrak{m}^r I_n^r$$

for $r = 6$ and thus for all $r \gg 0$. In particular, from the proof of [BGHN20a, Theorem 3.1], the containment hold for $r \geq 12$. Here we show that the containment hold for all $r \geq 5$. Notice that for $r \leq 4$, since the resurgence number $\rho(I_n) = \frac{3}{2}$, we know that $I_n^{(2r-2)} \not\subseteq I_n^r$.

Corollary 3.6. *For every $n \geq 3$, Fermat configuration ideal verifies the following containment*

$$I_n^{(2r-2)} \subseteq \mathfrak{m}^r I_n^r, \quad \forall r \geq 5$$

Proof. As before, since for all $r \geq 5$, $I_n^{(2r-2)} \subseteq I_n^r$, it suffices to check the inequalities

$$\alpha(I_n^{(2r-2)}) \geq r + \omega(I_n^r)$$

case by case.

- (1) For $n = 3$, we have $r + \omega(I_n^r) = 5r$ and $\alpha(I_n^{(2r-2)}) = \begin{cases} 9m, & \text{if } 2r - 2 = 3m. \\ 9m + 4, & \text{if } 2r - 2 = 3m + 1 \\ 9m + 8, & \text{if } 2r - 2 = 3m + 2. \end{cases}$
 - (a) If $2r - 2 = 3m$, the inequality becomes $9m \geq \frac{5}{2}(3m + 2)$ which is equivalent to $3m \geq 10$. Since $r \geq 5$, we have $3m \geq 8$. Moreover, $3m = 2r - 2$ can't be 8 or 9.
 - (b) If $2r - 2 = 3m + 1$, the inequality becomes $9m + 4 \geq \frac{5}{2}(3m + 3)$ which is equivalent to $3m \geq 7$ (which is true because $3m = 2r - 3 \geq 7$).
 - (c) If $2r - 2 = 3m + 2$, the inequality becomes $9m + 8 \geq \frac{5}{2}(3m + 4)$ which is equivalent to $3m \geq 4$.
- (2) For $n \geq 4$, since $r \geq 5$, $2r - 2 \geq 8$ so we have $\alpha(I_n^{(2r-2)}) = (2r - 2)n$, and the inequalities $(2r - 2)n \geq r(n + 1) + r \Leftrightarrow (r - 2)n \geq 2r$, which is true since we have $(r - 2)n \geq 4(r - 2) \geq 2r$ for all $r \geq 5$.

□

Although the above containment imply the Harbourne-Huneke containment for $r \geq 5$, we can check easily that Harbourne-Huneke containment hold for all possible r by our computations.

Corollary 3.7. *For every $n \geq 3$, Fermat configuration ideal verifies Harbourne-Huneke containment (see [HH13, Conjecture 2.1])*

$$I_n^{(2r)} \subseteq \mathfrak{m}^r I_n^r, \quad \forall r$$

Proof. Since $I_n^{(2r)} \subseteq I_n^r, \forall r \geq 1$ the above containment come from the fact that

$$\alpha(I_n^{(2r)}) \geq r + \omega(I_n^r)$$

for all $n \geq 3$ and $r \geq 1$ (the case $r = 0$ is trivial). Indeed, we check case by case

- (1) For $n \geq 3$ and $r = 1$, we have $2(n+1) \geq n+1+1$
- (2) For $n = 3$, we have $r + \omega(I_n^r) = 5r$ and $\alpha(I_n^{(2r)}) = \begin{cases} 9m, & \text{if } 2r = 3m \\ 9m + 4, & \text{if } 2r = 3m + 1 \\ 9m + 8, & \text{if } 2r = 3m + 2. \end{cases}$
- (3) For $n \geq 4$ and $r \geq 2$, we have $r + \omega(I_n^r) = r(n+2)$ and $\alpha(I_n^{(2r)}) = 2rn$.

□

Corollary 3.8. *For every $n \geq 3$, Fermat configuration ideal verifies Harbourne-Huneke containment (see [HH13, Conjecture 4.1.5])*

$$I_n^{(2r-1)} \subseteq \mathfrak{m}^{r-1} I_n^r, \quad \forall r \geq 3$$

Proof. Since $\rho(I_n) = \frac{3}{2}$, for all $r \geq 3$, $I_n^{(2r-1)} \subseteq I_n^r$, the above containment comes from the fact that

$$\alpha(I_n^{(2r-1)}) \geq r - 1 + \omega(I_n^r)$$

for all $n \geq 3$ and $r \geq 3$. Notice that when $r = 1, r = 2$, containment $I_n^{(2r-1)} \subseteq I_n^r$ fail. We check case by case

- (1) For $n = 3$, we have $r-1+\omega(I_n^r) = 5r-1$ and $\alpha(I_n^{(2r-1)}) = \begin{cases} 9m, & \text{if } 2r-1 = 3m. \\ 9m + 4, & \text{if } 2r-1 = 3m + 1 \\ 9m + 8, & \text{if } 2r-1 = 3m + 2. \end{cases}$
- (2) For $n = 4, r = 3$, we have $\alpha(I_n^{(5)}) = 21 > 3 + 5 \cdot 3 = r + \omega(I_n^3)$.
- (3) For $n \geq 4, r \neq 3$, we have $\alpha(I_n^{(2r-1)}) = (2r-1)n$, and the inequalities $(2r-1)n \geq r(n+1) + r - 1 \Leftrightarrow (r-1)n \geq 2r-1$, which is true since $(r-1)n \geq 4(r-1) \geq 2r-1$ for all $r \geq 3$.

□

We end this section by calculating another invariant related to generating degree of symbolic powers of I_n . As introduced in [HH13, Definition 2.2], for a homogeneous ideal $J \subset \mathbb{C}[\mathbb{P}^N]$, define $\beta(J)$ to be the smallest integer t such that $[J]_t$ contains a regular sequence of length two where $[J]_t$ is the graded component of degree t of J . It turns out that when $N = 2$ and J is a defining ideal of a finite set of (fat) points, $\beta(J)$ is in fact the least degree t such that the zero locus of $[J]_t$ is 0-dimensional since the condition that $[J]_t$

contains a regular sequence of length two is equivalent to the condition that all elements of $[J]_t$ does not have a non-constant common factor, c.f [HS15]. This invariant is related to the maximum degree of generators in a set of minimal generators of J as follows:

Proposition 3.9. *Let $J \subset \mathbb{C}[x, y, z]$ is a defining ideal of a set of finite (fat) points. Then $\beta(J) \leq \omega(J)$.*

Proof. Suppose that $J = \langle g_1, \dots, g_k \rangle$ and $\omega(J) < \beta(J)$, then the zero locus of $[J]_{\omega(J)}$ is not 0-dimensional by definition of $\beta(J)$. For $j = 1, \dots, k$, consider the set A consists of all forms $g_j x^{d_j}, g_j y^{d_j}, g_j z^{d_j}$ where $d_j = \omega(J) - \deg(g_j)$. Then $A \subset [J]_{\omega(J)}$, hence the zero locus of A is not 0-dimensional. This is a contradiction since the zero locus of A is also the zero locus of J . \square

In the following result, we calculate $\beta(I_n^{(m)})$ and get an immediate bound for $\omega(I_n^{(m)})$.

Proposition 3.10. *For $n \geq 3$ and $m \geq 1$, $\beta(I_n^{(m)}) = m(n+1)$.*

Proof. First, for any $n \geq 3$, note that I_n^m is generated by all generators of the same degree $m(n+1)$, hence, $\beta(I_n^m) = m(n+1)$ as $[I_n^m]_{m(n+1)}$ contains a regular sequence of length two. Since $I_n^m \subseteq I_n^{(m)}$, we have that $\beta(I_n^{(m)}) \geq m(n+1)$ for all $m \geq 1$. On the other hand, for any $m \geq 1$, let $f \in [I_n^{(m)}]_t$ be any element where $t < m(n+1)$. Recall that each line L_j in the Fermat configuration contains exactly $n+1$ points of the configuration. Intersecting any line L_j with the variety defined by f , since f vanishes at every point in the configuration to order at least m , by Bezout's theorem, since $\deg(f) \deg(L_j) < m(n+1)$, L_j is a component of the variety of f . Therefore, for any $t < m(n+1)$, L_j is a component of the zero locus of $[I_n^{(m)}]_t$, i.e, $\beta(I_n^{(m)}) \leq m(n+1)$. This ends the proof. \square

Corollary 3.11. *For $n \geq 3$ and $m \geq 1$, $\omega(I_n^{(m)}) \geq m(n+1)$. Moreover, for each $n \geq 3$, $\omega(I_n^{(m)}) = m(n+1)$ for all $m \geq n^2 - 3n + 2$.*

Proof. $\omega(I_n^{(m)}) \geq m(n+1)$ for all m is immediate. On the other hand, by [NS16, Theorem 3.10, Remark 3.11], $\text{reg}(I_n^{(m)}) = m(n+1)$ for all $n \geq 3$ and $m \geq n^2 - 3n + 2$. Thus $\omega(I_n^{(m)}) \leq \text{reg}(I_n^{(m)}) = m(n+1)$ for all $n \geq 3$ and $m \geq n^2 - 3n + 2$. \square

Remark 3.12. It is known that for any k and $n \geq 3$, $\omega(I_n^{(kn)}) = \beta(I_n^{(kn)}) = kn(n+1)$, since any generators of $I_n^{(kn)}$ with degree less than $kn(n+1)$ must be divisible by fgh and hence, none of two elements in degree less than $kn(n+1)$ of it form a regular sequence (because they always share the common factor fgh), see [NS16, Theorem 3.7]. In the same paper, it is also known that $\omega(I_n^{(n-1)}) = \beta(I_n^{(n-1)}) = (n-1)(n+1)$ with the same reason. As in above corollary, $\omega(I_n^{(m)}) = \beta(I_n^{(m)}) = m(n+1)$ for $m \gg 0$. It is reasonable to ask if this is the case for all m . It is suggested by Macaulay2 [GS] that in fact $\omega(I_n^{(m)}) = m(n+1)$ for all m .

4. FERMAT IDEAL I_2

In this section, we will deal with the ideal $I_2 = (x(y^2 - z^2), y(z^2 - x^2), z(x^2 - y^2))$. Unlike the ideals I_n for $n \geq 3$, this ideal satisfies the Harbourne containment $I_2^{(3)} \subseteq I_2^2$. This is probably one reason why it is less considered in the literature. We will see later that this ideal is very different to Fermat ideals when $n \geq 3$ in term of generating degree of symbolic

powers and hence, in term of Waldschmidt constant and (asymptotic) resurgence number.

In the following, we will compute the least degree of generators of symbolic powers of I_2 as well as its Waldschmidt constant, (asymptotic) resurgence number in order to complete the picture of Fermat ideals. This ideal I_2 is also known to be the ideal determining the singular locus of the arrangement of lines given by the pseudoreflection group D_3 , see [DS20].

First, recall that, geometrically, I_2 is the defining ideal of the singular locus of the line arrangement in \mathbb{P}^2 that consists of 6 lines L_j whose equations are

$$x = \pm y, y = \pm z, z = \pm x$$

These 6 lines intersect at 7 points P_i which are $[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1], [1 : 1 : -1], [1 : -1 : 1], [1 : -1 : -1]$ and $[1 : 1 : 1]$ such that the first 3 points lie on 2 lines each and the rest lie on 3 lines each; and each line contains exactly 3 points.

Similar to I_n for $n \geq 3$, we can write

$$I_2 = (x^2 - y^2, y^2 - z^2) \cap (x, y) \cap (y, z) \cap (z, x)$$


so that

$$I_2^{(m)} = (x^2 - y^2, y^2 - z^2)^m \cap (x, y)^m \cap (y, z)^m \cap (z, x)^m \quad \forall m$$

Theorem 4.1. *For ideal I_2 we have the following*

- (1) $\widehat{\alpha}(I_2) = \frac{5}{2}$
- (2) $\alpha(I_2^{(2k)}) = 5k$, for all $k \geq 2$.
- (3) $\alpha(I_2^{(2k+1)}) = 5k + 3$, for all $k \geq 0$.
- (4) $\alpha(I_2^{(2)}) = 6$

Proof. By [FGH⁺17, Theorem 2.3], we can check that $\widehat{\alpha}(I_2) \geq \frac{5}{2}$.



In particular, we have that $\alpha(I_2^{(2k)}) \geq 5k$ and $\alpha(I_2^{(2k+1)}) \geq 5k + 3$, for all $k \geq 1$. We will show the reverse by showing there exists some element with the desired degree in the symbolic powers.

Indeed, we consider case by case. Denote $K = (x^2 - y^2, y^2 - z^2)$, we have

- (1) Case 1: when $m = 4k$. Consider $F = (x^2 - y^2)^{2k}(y^2 - z^2)^k(z^2 - x^2)^k z^{2k}$ that has degree $10k$. Since $(x^2 - y^2)^{2k}(y^2 - z^2)^k(z^2 - x^2)^k \in K^{4k}$, $(x^2 - y^2)^{2k} \in (x, y)^{4k}$, $z^{2k}(y^2 - z^2)^k \in (y, z)^{4k}$ and $z^{2k}(z^2 - x^2)^k \in (z, x)^{4k}$ it follows that

$$F \in K^{4k} \cap (x, y)^{4k} \cap (y, z)^{4k} \cap (z, x)^{4k} = I_2^{(4k)}, \quad \forall k \geq 1$$

By similar argument we see that

- (2) Case 2: when $m = 4k + 2$. $F = (x^2 - y^2)^{2k}(y^2 - z^2)^{k+1}(z^2 - x^2)^{k+1}xyz^{2k-1}$ has degree $10k + 5$ and

$$F \in K^{4k+2} \cap (x, y)^{4k+2} \cap (y, z)^{4k+2} \cap (z, x)^{4k+2} = I_2^{(4k+2)}, \quad \forall k \geq 1$$

- (3) Case 3: when $m = 4k + 1$. $F = (x^2 - y^2)^{2k+1}(y^2 - z^2)^k(z^2 - x^2)^k z^{2k+1}$ has degree $10k + 3$ and

$$F \in K^{4k+1} \cap (x, y)^{4k+1} \cap (y, z)^{4k+1} \cap (z, x)^{4k+1} = I_2^{(4k+1)}, \quad \forall k \geq 0$$

- (4) Case 4: when $m = 4k + 3$. $F = (x^2 - y^2)^{2k+1}(y^2 - z^2)^{k+1}(z^2 - x^2)^{k+1}xyz^{2k}$ has degree $10k + 8$ and

$$F \in K^{4k+3} \cap (x, y)^{4k+3} \cap (y, z)^{4k+3} \cap (z, x)^{4k+3} = I_2^{(4k+3)}, \quad \forall k \geq 0$$

Thus, $\alpha(I_2^{(2k)}) \leq 5k$, for all $k \geq 2$ and $\alpha(I_2^{(2k+1)}) \leq 5k + 3$, for all $k \geq 0$. It follows that statements (2) and (3) are true and by taking limits as k goes to ∞ , (1) follows as well. Part (4) can be checked directly by Macaulay2 or by Bezout theorem argument as follows: We know that $\alpha(I_2^{(2)}) \leq \alpha(I_2^2) = 6$. Now suppose that $\alpha(I_2^{(2)}) \leq 5$. Then there is a divisor D of degree 5 vanishing to order at least 2 at every point P_i . Since the intersection of D and any line L_j consists of 3 points to order at least 2, we get a contradiction to Bezout theorem because $\deg(D) \cdot \deg(L_j) = 5 < 2 \cdot 3$. \square

For I_2 , the asymptotic resurgence number and the resurgence number turn out to be the same.

Theorem 4.2. *The resurgence number and asymptotic resurgence number of I_2 are:*

$$\rho(I_2) = \widehat{\rho}(I_2) = \frac{6}{5}$$

Proof. The asymptotic resurgence number $\widehat{\rho}(I_2) = \frac{6}{5}$ follows from [GHVT13, Theorem 1.2] that

$$\frac{6}{5} = \frac{\alpha(I_2)}{\widehat{\alpha}(I_2)} \leq \widehat{\rho}(I_2) \leq \frac{\omega(I_2)}{\widehat{\alpha}(I_2)} = \frac{6}{5}$$

By [NS16, Theorem 2.5], since I_2 is a strict almost complete intersection ideal with minimal generators of degree 3 and its module syzygy is generated in degree 1 and 2, the minimal free resolution of I_2^r is:

$$0 \rightarrow R(-3(r+1))^{\binom{r}{2}} \xrightarrow{\psi} \begin{array}{c} R(-3r-1)^{\binom{r+1}{2}} \\ \oplus \\ R(-3r-2)^{\binom{r+1}{2}} \end{array} \xrightarrow{\varphi} R(-3r)^{\binom{r+2}{2}} \rightarrow I_2^r \rightarrow 0$$

for any $r \geq 2$, in particular, $\text{reg}(I_2^r) = 3r + 1$ for all $r \geq 2$, where $\text{reg}(I)$ denotes the Castelnuovo-Mumford regularity of I . We also have $\text{reg}(I_2) = 4$. By [GHVT13, Theorem 1.2] again, we have $\rho(I_2) \geq \frac{\alpha(I_2)}{\widehat{\alpha}(I_2)} = \frac{6}{5}$. Conversely, for any $\frac{m}{r} > \frac{6}{5}$, we have:

- If $m = 2k$ then $10k > 6r$ implies that $10k \geq 6r + 2$ or $5k \geq 3r + 1$ since both numbers are even. It follows that $\alpha(I_2^{(2k)}) \geq 5k \geq 3r + 1 = \text{reg}(I_2^r)$ and hence, $I_2^{(2k)} \subseteq I_2^r$.
- If $m = 2k + 1$ then $10k + 2 > 6r$ implies that $10k + 6 \geq 6r + 2$ or $5k + 3 \geq 3r + 1$. It follows that $\alpha(I_2^{(2k+1)}) = 5k + 3 \geq 3r + 1 = \text{reg}(I_2^r)$ and hence, $I_2^{(2k+1)} \subseteq I_2^r$.

Thus, for any $\frac{m}{r} > \frac{6}{5}$, we have $I_2^{(m)} \subseteq I_2^r$, i.e., $\rho(I_2) = \frac{6}{5}$. □

Example 4.3. It is worth to point out that the first direct consequence of the above calculation is the verification of I_2 to Chudnovsky's Conjecture and Demailly's Conjecture, although the general case is already known from [EV83]. Ideal I_2 verifies

- (1) Chudnovsky's Conjecture $\widehat{\alpha}(I_2) \geq \frac{\alpha(I_2) + 1}{2}$.
- (2) Demailly's Conjecture $\widehat{\alpha}(I_2) \geq \frac{\alpha(I_2^{(m)}) + 1}{m + 1}$ for all $m \geq 1$.

Proof. Directly from the formulae of $\widehat{\alpha}(I_2)$ and $\alpha(I_2^{(m)})$. □

Example 4.4. Another difference between I_2 and I_n when $n \geq 3$ is that while $I_n^{(nk)} = (I_n^{(n)})^k$ for all k and $n \geq 3$ by [NS16], $I_2^{(4)} \neq (I_2^{(2)})^2$. In fact, as in the proof of theorem 4.1,

$$F = (x^2 - y^2)^2(y^2 - z^2)(z^2 - x^2)z^2 \in I_2^{(4)}$$

whereas, we can check that $F \notin (I_2^{(2)})^2$. Moreover, we can also check that $I_2^{(6)} \neq (I_2^{(3)})^2$ since

$$G = (x^2 - y^2)^2(y^2 - z^2)^2(z^2 - x^2)^2xyz \in I_2^{(6)} \setminus (I_2^{(3)})^2$$

It is suggested by Macaulay2 [GS] that $I_2^{(8)} = (I_2^{(4)})^2$. It is interesting to know if $I_2^{(4k)} = (I_2^{(4)})^k$ for all k .

As all other Fermat ideals, I_2 also satisfies the following containment. In [BGHN20a, Example 3.7], we showed the stronger containment (which implies both Harbourne-Huneke) containment

$$I_2^{(2r-2)} \subseteq \mathfrak{m}^r I_2^r$$

for $r = 5$ (by Macaulay2) and thus for all $r \gg 0$ by our method. In particular, from the proof of [BGHN20a, Theorem 3.1], the containment hold for $r \geq 10$. Here we show that the containment hold for all $r \geq 5$.

Corollary 4.5. *For every $n \geq 3$, ideal I_2 verifies the following stronger containment*

$$I_2^{(2r-2)} \subseteq \mathfrak{m}^r I_2^r, \quad \forall r \geq 5$$

Proof. As before, since $\rho(I_2) = \frac{6}{5}$, we know that $I_2^{(2r-2)} \subseteq I_2^r$ for $r \geq 3$. Hence, the containment follows from the inequalities

$$\alpha(I_2^{(2r-2)}) = 5r - 5 \geq r + 3r = r + \omega(I_2^r)$$

for all $r \geq 5$. □

We can also detect the failure of the containment in the remaining cases by only using formulae for $\alpha(I_2^{(m)})$ as follows.

Remark 4.6. For $r \leq 4$, the above containment fail. In fact, notice that for $r \leq 2$, since $\rho(I_2) = \frac{6}{5}$, we know that $I_2^{(2r-2)} \not\subseteq I_2^r$. When $r = 3$, since $\alpha(I_2^{(4)}) = 10 < 3 + 9 = 3 + \alpha(I_2^3)$, we see that the containment $I_2^{(4)} \subseteq \mathfrak{m}^3 I_2^3$ fails. Similarly, for $r = 4$, since $\alpha(I_2^{(6)}) = 15 < 4 + 12 = 4 + \alpha(I_2^4)$, we see that the containment $I_2^{(6)} \subseteq \mathfrak{m}^4 I_2^4$ also fails.

Although the above containment imply the Harbourne-Huneke containment for $r \geq 5$, we can check easily that Harbourne-Huneke containment hold for all possible r by our computations.

Corollary 4.7. *Ideal I_2 verifies Harbourne-Huneke containment (see [HH13, Conjecture 2.1])*

$$I_2^{(2r)} \subseteq \mathfrak{m}^r I_2^r, \quad \forall r$$

Proof. Since for all r , $I_2^{(2r)} \subseteq I_2^r$, the containment follow from the fact that

$$\alpha(I_2^{(2r)}) \geq 5r \geq r + 3r = r + \omega(I_2^r)$$

for all r . □

Corollary 4.8. *Ideal I_2 verifies Harbourne-Huneke containment (see [HH13, Conjecture 4.1.5])*

$$I_2^{(2r-1)} \subseteq \mathfrak{m}^{r-1} I_2^r, \quad \forall r \geq 1$$

Proof. Since $\rho(I_2) = \frac{6}{5}$, for all $r \geq 2$, $I_2^{(2r-1)} \subseteq I_2^r$, the above containment comes from the fact that

$$\alpha(I_2^{(2r-1)}) = 5r - 2 \geq r - 1 + 3r = r - 1 + \omega(I_2^r)$$

for all $r \geq 2$. The case $r = 1$ is obvious. □

Remark 4.9. The above corollary gives a proof for the case D_3 in [DS20, Proposition 6.3].

We end this section by calculating $\beta(I_2^{(m)})$.

Proposition 4.10. *For all $m \geq 1$, $\beta(I_2^{(m)}) = 3m$ and $\omega(I_2^{(m)}) \geq 3m$.*

Proof. The proof is the same as that of the case where $n \geq 3$. First, since $I_2^m \subseteq I_2^{(m)}$, we have that $\beta(I_2^{(m)}) \geq 3m$ for all $m \geq 1$. On the other hand, recall that each line L_j in the configuration contains exactly 3 points of the configuration. Thus, for any $m \geq 1$ and for any $f \in [I_n^{(m)}]_t$ where $t < 3m$, intersecting any line L_j with the variety defined by f , by Bezout's theorem, since $\deg(f) \deg(L_j) < 3m$, L_j is a component of the variety of f . Therefore, for any $t < 3m$, L_j is a component of the zero locus of $[I_2^{(m)}]_t$, i.e., $\beta(I_2^{(m)}) \leq 3m$. □

Remark 4.11. As in the case where $n \geq 3$, Macaulay2 [GS] suggests that $\omega(I_2^{(m)}) = 3m$. It is interesting to know if $\omega(J^{(m)}) = \beta(J^{(m)})$ hold for what radical ideal of points J in general.

REFERENCES

- [BDRH⁺09] Thomas Bauer, Sandra Di Rocco, Brian Harbourne, Michał Kapustka, Andreas Knutsen, Wioletta Syzdek, and Tomasz Szemberg. A primer on Seshadri constants. In *Interactions of classical and numerical algebraic geometry*, volume 496 of *Contemp. Math.*, pages 33–70. Amer. Math. Soc., Providence, RI, 2009.
- [BGHN20a] Sankhaneel Bisui, Eloísa Grifo, Huy Tài Hà, and Thái Thành Nguyễn. Chudnovsky’s conjecture and the stable Harbourne-Huneke containment. *arXiv:2004.11213*, 2020.
- [BGHN20b] Sankhaneel Bisui, Eloísa Grifo, Huy Tài Hà, and Thái Thành Nguyễn. Demailly’s conjecture and the containment problem. *arXiv:2009.05022*, 2020.
- [BH10] Cristiano Bocci and Brian Harbourne. Comparing powers and symbolic powers of ideals. *J. Algebraic Geom.*, 19(3):399–417, 2010.
- [CHHVT20] Enrico Carlini, Huy Tài Hà, Brian Harbourne, and Adam Van Tuyl. *Ideals of powers and powers of ideals: Intersecting Algebra, Geometry and Combinatorics*, volume 27 of *Lecture Notes of the Unione Matematica Italiana*. Springer International Publishing, 2020.
- [Chu81] Gregory V. Chudnovsky. Singular points on complex hypersurfaces and multidimensional schwarz lemma. In *Séminaire de Théorie des Nombres, Paris 1979-80, Séminaire Delange-Pisot-Poitou*, volume 12 of *Progress in Math.*, pages 29–69. Birkhäuser, Boston, Basel, Stuttgart, 1981.
- [CJ20] Yu-Lin Chang and Shin-Yao Jow. Demailly’s conjecture on Waldschmidt constants for sufficiently many very general points in \mathbb{P}^n . *J. Number Theory*, 207:138–144, 2020.
- [DDSG⁺18] Hailong Dao, Alessandro De Stefani, Eloísa Grifo, Craig Huneke, and Luis Núñez Betancourt. Symbolic powers of ideals. In *Singularities and foliations. geometry, topology and applications*, volume 222 of *Springer Proc. Math. Stat.*, pages 387–432. Springer, Cham, 2018.
- [Dem82] J.-P. Demailly. Formules de jensen en plusieurs variables et applications arithmétiques. *Bull. Soc. Math. France*, 110(1):75–102, 1982.
- [DHN⁺15] M. Dumnicki, B. Harbourne, U. Nagel, A. Secoleanu, T. Szemberg, and H. Tutaj-Gasińska. Resurgences for ideals of special point configurations in \mathbf{P}^N coming from hyperplane arrangements. *J. Algebra*, 443:383–394, 2015.
- [DS20] Ben Drabkin and Alexandra Secoleanu. Singular loci of reflection arrangements and the containment problem. *arXiv:2002.05353*, 2020.
- [DSTG13] Marcin Dumnicki, Tomasz Szemberg, and Halszka Tutaj-Gasińska. Counterexamples to the $I^{(3)} \subseteq I^2$ containment. *J. Alg.*, 393:24–29, 2013.
- [DTG17] Marcin Dumnicki and Halszka Tutaj-Gasińska. A containment result in P^n and the Chudnovsky conjecture. *Proc. Amer. Math. Soc.*, 145(9):3689–3694, 2017.
- [Dum15] Marcin Dumnicki. Containments of symbolic powers of ideals of generic points in \mathbb{P}^3 . *Proc. Amer. Math. Soc.*, 143(2):513–530, 2015.
- [EH79] David Eisenbud and Melvin Hochster. A Nullstellensatz with nilpotents and Zariski’s main lemma on holomorphic functions. *J. Algebra*, 58(1):157–161, 1979.
- [ELS01] Lawrence Ein, Robert Lazarsfeld, and Karen E. Smith. Uniform bounds and symbolic powers on smooth varieties. *Invent. Math.*, 144 (2):241–25, 2001.
- [EV83] Hélène Esnault and Eckart Viehweg. Sur une minoration du degré d’hypersurfaces s’annulant en certains points. *Math. Ann.*, 263(1):75–86, 1983.
- [FGH⁺17] L. Farnik, J. Gwoździewicz, B. Hejmej, M. Lampa-Baczyńska, G. Malara, and J. Szpond. Initial sequences and Waldschmidt constants of planar point configurations. *Int. J. Algebra Comput.*, 27 (06):717–729, 2017.
- [FMX18] Louiza Fouli, Paolo Mantero, and Yu Xie. Chudnovsky’s conjecture for very general points in \mathbb{P}_k^N . *J. Algebra*, 498:211–227, 2018.
- [GHM13] Anthony V. Geramita, Brian Harbourne, and Juan Migliore. Star configurations in \mathbb{P}^n . *J. Algebra*, 376:279–299, 2013.
- [GHVT13] Elena Guardo, Brian Harbourne, and Adam Van Tuyl. Asymptotic resurgences for ideals of positive dimensional subschemes of projective space. *Adv. Math.*, 246:114–127, 2013.

- [GS] D. R. Grayson and M. E. Stillman. *Macaulay2, a software system for research in algebraic geometry*.
- [HH02] Melvin Hochster and Craig Huneke. Comparison of symbolic and ordinary powers of ideals. *Invent. Math.*, 147, 2002.
- [HH13] Brian Harbourne and Craig Huneke. Are symbolic powers highly evolved? *J. Ramanujan Math. Soc.*, 28A:247–266, 2013.
- [HS15] Brian Harbourne and Alexandra Seceleanu. Containment counterexamples for ideals of various configurations of points in \mathbf{P}^N . *J. Pure Appl. Algebra*, 219(4):1062–1072, 2015.
- [MS17] G. Malara and J. Szpond. Fermat-type configurations of lines in \mathbf{p}^3 and the containment problem. *Journal of Pure and Applied Algebra*, 222:2323–2329, 2017.
- [MS18] Grzegorz Malara and Justyna Szpond. On codimension two flats in Fermat-type arrangements. In *Multigraded algebra and applications*, volume 238 of *Springer Proc. Math. Stat.*, pages 95–109. Springer, Cham, 2018.
- [MSS18] Grzegorz Malara, Tomasz Szemberg, and Justyna Szpond. On a conjecture of Demailly and new bounds on Waldschmidt constants in \mathbb{P}^N . *J. Number Theory*, 189:211–219, 2018.
- [Nag62] Masayoshi Nagata. *Local rings*. Interscience, 1962.
- [Ngu20] Thái Thành Nguyễn. The least generating degree of symbolic powers of fermat-like ideals of planes and lines arrangements. *Preprint*, 2020.
- [NS16] Uwe Nagel and Alexandra Seceleanu. Ordinary and symbolic Rees algebras for ideals of Fermat point configurations. *J. Algebra*, 468:80–102, 2016.
- [Szp19] Justyna Szpond. Fermat-type arrangements. *arXiv:1909.04089*, 2019.
- [Zar49] Oscar Zariski. A fundamental lemma from the theory of holomorphic functions on an algebraic variety. *Ann. Mat. Pura Appl. (4)*, 29:187–198, 1949.

TULANE UNIVERSITY, DEPARTMENT OF MATHEMATICS, 6823 ST. CHARLES AVE., NEW ORLEANS, LA 70118, USA AND HUE UNIVERSITY, COLLEGE OF EDUCATION, VIETNAM.

Email address: `tnguyen11@tulane.edu`

URL: <https://sites.google.com/view/thainguyenmath>