

PROJECTIVE CLOSURES OF AFFINE MONOMIAL CURVES

JOYDIP SAHA, INDRANATH SENGUPTA, AND PRANJAL SRIVASTAVA

ABSTRACT. Our aim in this paper is to study the projective closures of three important families of affine monomial curves in dimension 4, in order to explore possible connections between syzygies and the arithmetic Cohen-Macaulay property.

1. INTRODUCTION

Let $r \geq 3$ and $\underline{n} = (n_1, \dots, n_r)$ be a sequence of r distinct positive integers with $\gcd(\underline{n}) = 1$. Let us assume that the numbers n_1, \dots, n_r generate the numerical semigroup $\Gamma(n_1, \dots, n_r)$ in the set of all non-negative integers \mathbb{N} :

$$\Gamma(n_1, \dots, n_r) = \left\{ \sum_{j=1}^r z_j n_j \mid z_j \text{ nonnegative integers} \right\}$$

minimally, that is, if $n_i = \sum_{j=1}^r z_j n_j$ for some non-negative integers z_j , then $z_j = 0$ for all $j \neq i$ and $z_i = 1$. Let $\eta : k[x_1, \dots, x_r] \rightarrow k[t]$ be the mapping defined by $\eta(x_i) = t^{n_i}$, $1 \leq i \leq r$. Let $\mathfrak{p}(n_1, \dots, n_r) = \ker(\eta)$. Let $\beta_i(\mathfrak{p}(n_1, \dots, n_r))$ denote the i -th Betti number of the ideal $\mathfrak{p}(n_1, \dots, n_r)$. Therefore, $\beta_1(\mathfrak{p}(n_1, \dots, n_r))$ denotes the minimal number of generators $\mathfrak{p}(n_1, \dots, n_r)$. For a given $r \geq 3$, let $\beta_i(r) = \text{lub}(\beta_i(\mathfrak{p}(n_1, \dots, n_r)))$, where lub is taken over all the sequences of positive integers n_1, \dots, n_r . Herzog [9] proved that $\beta_1(3)$ is 3 and it follows easily that $\beta_2(3)$ is a finite integer as well. Bresinsky [5] (and [6]) defined a class of monomial curves in \mathbb{A}^4 and proved that $\beta_1(4) = \infty$. He used this observation to prove that $\beta_1(r) = \infty$, for every $r \geq 4$. We call this family the *Bresinsky curves*. Subsequently, it has been proved in [10] that all the higher Betti numbers of Bresinsky curves are also not bounded above by a fixed integer. Subsequent to Bresinsky curves came the examples of *Backelin curves* [2] and

2010 *Mathematics Subject Classification.* Primary 13D02, 13F05, 13P10, 13P20.

Key words and phrases. Monomial curves, Gröbner bases, Betti numbers.

The first author thanks SERB, Government of India for NPDF at ISI kolkata, through the research project PDF/2019/001074.

The second author is the corresponding author.

the examples of *Arslan curves* [1]. All three families have one strong resemblance that all are monomial curves in the affine 4 space and there is no upper bound on the minimal generating set of the defining ideals of these curves.

Let us assume that $n_r > n_i$ for all $i < r$, for the sequence $\underline{n} = (n_1, \dots, n_r)$. We fix $n_0 = 0$. We define the semigroup $\overline{\Gamma}(n_1, \dots, n_r) \subset \mathbb{N}^2$, (\mathbb{N} is the set of all non negative integers) generated by $\{(n_i, n_r - n_i) \mid 0 \leq i \leq r\}$. Let us denote $\overline{\mathfrak{p}}(n_1, \dots, n_r)$ be the kernel of k -algebra map $\eta^H : k[x_0, \dots, x_r] \rightarrow k[s, t]$, $\eta^H(x_i) = t^{n_i} s^{n_r - n_i}$, $0 \leq r$. Then homogenization of the ideal $\mathfrak{p}(n_1, \dots, n_r)$ w.r.t. the variable x_0 is $\overline{\mathfrak{p}}(n_1, \dots, n_r)$. Thus the projective curve $\{[(a^{n_r} : a^{n_r - n_1} b^{n_1} : \dots : b^{n_r})] \in \mathbb{P}_k^n \mid a, b \in k\}$ is the projective closure of the affine curve $C(n_1, \dots, n_r) := \{(b^{n_1}, \dots, b^{n_r}) \in \mathbb{A}_k^n \mid b \in k\}$ and we denote it by $\overline{C}(n_1, \dots, n_r)$. We say that the Projective curve $\overline{C}(n_1, \dots, n_r)$ is arithmetically Cohen-Macaulay if the vanishing ideal $\overline{\mathfrak{p}}(n_1, \dots, n_r)$ is a Cohen-Macaulay ideal.

This paper is devoted to a study of the three families: Backelin, Bresinsky and Arslan curves and their projective closures. First we study the affine Backelin curves, its Gröbner basis and use that to prove that its projective closure is arithmetically Cohen-Macaulay. We also compute the Hilbert series of the affine Backelin curves. Next we compute the syzygies of the projective closure of Bresinsky curves and finally the syzygies of the projective closure of Arslan curves. Our aim in this paper is to find suitable conditions on the affine monomial curves which ensure the arithmetic Cohen-Macaulayness of its projective closure. We suspect that syzygies is likely to act as a determining factor. This paper is a case study with the help of three very important and interesting classes of curves.

2. BACKELIN CURVES AND ITS GRÖBNER BASIS

Let us first begin with a description of Bakelin's example of monomial curves in the affine space \mathbb{A}^4 . Beckelin defines the following semigroups $\langle s, s+3, s+3n+1, s+3n+2 \rangle$, for $n \geq 2$, $r \geq 3n+2$ and $s = r(3n+2)+3$. In [2], it has been shown that the type of such semigroups are not bounded by the embedding dimension.

Notations 2.1. We fix some notations. For $n \geq 2$, $r \geq 3n+2$ and $s = r(3n+2)+3$,

- $\Gamma_{nr} := \Gamma(s, s+3, s+3n+1, s+3n+2)$;
- $\overline{\Gamma}_{nr} := \overline{\Gamma}(s, s+3, s+3n+1, s+3n+2)$;
- $\mathfrak{P}_{nr} := \mathfrak{p}(s, s+3, s+3n+1, s+3n+2)$;

- $\overline{\mathfrak{P}}_{nr} := \overline{\mathfrak{p}(s, s+3, s+3n+1, s+3n+2)}$, therefore $\overline{\mathfrak{P}}_{nr} = (\mathfrak{P}_{nr})^H$ with respect to the variable x_0 ;
- B_{nr} denotes the affine Backelin curves $C(s, s+3, s+3n+1, s+3n+2)$;
- \overline{B}_{nr} denotes the projective closures of Backelin curves;
- $G(I)$ denotes the unique minimal generating set of a monomial I in a polynomial ring over a field.

Theorem 2.1. (Gastinger) *Let $A = k[x_1, \dots, x_r]$ be the polynomial ring, $I \subset A$ the defining ideal of a monomial curve defined by natural numbers a_1, \dots, a_r , whose greatest common divisor is 1. Let J be an ideal contained in I . Then $J = I$ if and only if $\dim_k A/\langle J + (x_i) \rangle = a_i$, for some i ; equivalently $\dim_k A/\langle J + (x_i) \rangle = a_i$ for any i .*

Proof. See in [3]. □

Let $I \subset k[x_1, \dots, x_r]$ be a monomial ideal, then it has unique minimal generating set. We denote the minimal generating set by $G(I)$.

Theorem 2.2. *The defining ideal \mathfrak{P}_{nr} of monomial curve associated to Γ_{nr} is minimally generated by following binomials*

$$\begin{aligned} f_1 &= x_2 x_3^3 - x_1 x_4^3 \\ f_{2i} &= x_1^{n-i} x_3^{3i-1} - x_2^{n-i+1} x_4^{3i-2}, 1 \leq i \leq n \\ f_{3j} &= x_1^{r-n+3+j} x_2^{n-1-j} - x_3^{2+3j} x_4^{r-1-3j}, 0 \leq j \leq n-1 \\ f_{4j} &= x_1^{r-2n+3+j} x_2^{2n-j} - x_3^{3j+1} x_4^{r+1-3j}, 0 \leq j \leq n-1 \\ f_5 &= x_1^{r-n+2} x_2^n x_3 - x_4^{r+2} \\ f_6 &= x_2^{n+1} x_3 - x_1^n x_4^2 \\ f_7 &= x_2^{2n+1} - x_1^{2n-1} x_3 x_4 \end{aligned}$$

Proof. We proceed by Gastinger's theorem stated as in 2.1.

Let $J_{nr} = \langle \{f_1, f_{2i}, f_{3j}, f_{4j}, f_5, f_6 \mid 1 \leq i \leq n, 0 \leq j \leq n-1\} \rangle$ and $\mathfrak{A}_{nr} = \{x_1 x_4^3, x_2^{n-i+1} x_4^{3i-2}, x_1^{r-n+3+j} x_2^{n-1-j}, x_1^{r-2n+3+j} x_2^{2n-j}, x_4^{r+2}, x_1^n x_4^2, x_2^{2n+1} \mid 1 \leq i \leq n, 0 \leq j \leq n-1\}$. Then $J_{nr} + (x_3) = \langle \mathfrak{A}_{nr} \rangle$. At first we note that $J_{nr} \subset \mathfrak{P}_{nr}$.

$A/J_{nr} + \langle x_3 \rangle$ is the vector space over k whose basis consists of the images of monomials are listed bellow

- $S_1 = \{x_1^\alpha : 0 \leq \alpha \leq r+1\}$
- $S_2 = \{x_2^\beta : 0 \leq \beta \leq 2n\}$
- $S_3 = \{x_3^\gamma : 0 \leq \gamma \leq r+1\}$
- $S_4 = \{x_1^\alpha x_2^\beta : 1 \leq \alpha \leq r-\beta+1, 1 \leq \beta \leq n-1\}$
- $S_5 = \{x_1^\alpha x_2^\beta : 1 \leq \alpha \leq r-\beta+2, n+1 \leq \beta \leq 2n\}$
- $S_6 = \{x_1^\alpha x_3 : 1 \leq \alpha \leq r+1\}$
- $S_7 = \{x_1^\alpha x_3^2 : 1 \leq \alpha \leq n-1\}$

- $S_8 = \{x_2^\beta x_3^\gamma : 1 \leq \beta \leq n-1, 1 \leq \gamma \leq 3(n-\beta)\}$
- $S_9 = \{x_1^\alpha x_2^\beta x_3^\gamma : 1 \leq \alpha \leq n-1, 1 \leq \beta \leq n-1, 1 \leq \gamma \leq 2\}$
- $S_{10} = \{x_1^\alpha x_\beta x_3 : n \leq \alpha \leq r-\beta+1, 1 \leq \beta \leq n-1\}$
- $S_{11} = \{x_1^\alpha x_2^n : 1 \leq \alpha \leq r-n+2\}$

Then cardinality of this basis is,

$$\begin{aligned} \sum_{i=1}^{11} |S_i| &= 2n + 2r + 3 + \frac{(n-1)}{2}(2r-n+2) + \frac{n}{2}(2r-3n+3) + (r+1) \\ &+ (n-1) + \frac{3}{2}n(n-1) + 2(n-1)^2 + (r-n+2) + \frac{n-1}{2}(2r-3n+4) \\ &= 3nr + 3n + 2r + 4. \end{aligned}$$

Hence $\dim_k(A/J_{nr} + \langle x_3 \rangle) = 3nr + 3n + 2r + 4 = s + 3n + 1$. So by Gastinger Theorem we have $J_{nr} = \mathfrak{P}_{nr}$. \square

Let us denote the above generating set of \mathfrak{P}_{nr} by \mathfrak{S}_{nr} , thus $\mathfrak{S}_{nr} = \{f_1, f_{2i}, f_{3j}, f_{4j}, f_5, f_6 \mid 1 \leq i \leq n, 0 \leq j \leq n-1\}$.

Theorem 2.3. *Let $g = x_1^{r+2} - x_2 x_4^r$ be a polynomial in $k[x_1, \dots, x_4]$. Suppose $G_{nr} = (\mathfrak{S}_{nr} \setminus \{f_{3(n-1)}\}) \cup \{g\}$. Then G_{nr} also a generating set for the defining ideal \mathfrak{P}_{nr} .*

Proof. Follows from the relation $g = f_{3,(n-1)} + x_4^{r-3n+2} \cdot f_{2,n}$. \square

Theorem 2.4. *Let us consider the degree reverse lexicographic monomial order on $k[x_1, \dots, x_4]$ induced by $x_1 > x_2 > x_3 > x_4$ then G_{nr} is Gröbner basis of defining ideal \mathfrak{P}_{nr} with respect to this order.*

Proof. We proceed by Buchberger's algorithm,

(1)

$$\begin{aligned} S(f_1, f_{(2,i)}) &= x_2^{n+1} x_4 x_3 - x_1^n x_4^3 = x_4 f_6, \quad \text{for } i = 1 \\ &= x_2^{n-i+2} x_4^{3i-2} - x_1^{n-i+1} x_3^{3i-4} x_4^3 = x_4^3 f_{(2,i-1)}, \quad \text{for } 2 \leq i \leq n. \end{aligned}$$

(2)

$$\begin{aligned} S(f_1, f_{(3,i)}) &= x_3^{3n-1} x_4^{r-3n+5} - x_1^{r+2} x_4^3 = f_{(2,0)} x_4^{r-3n+5} - g x_4^3, \quad \text{for } i = n-2 \\ &= x_3^{5+3i} x_4^{r-1-3i} - x_1^{r-n+4+i} x_2^{n-2-i} x_4^3 = -f_{(3,i+1)} x_4^3, \quad \text{for } 2 \leq i \leq n-3. \end{aligned}$$

(3)

$$\begin{aligned} S(f_1, f_{(4,i)}) &= x_3^{3n+1} x_4^{r+1-3n+3} - x_1^{r-n+3} x_2^n x_4^3 = f_{(2,n)} x_3^2 x_4^{r-3n+4} - f_{(3,0)} x_2 x_4^3 \quad \text{for } i = n-1 \\ &= x_3^{3i+4} x_4^{r+1-3i} - x_1^{r-2n+4+i} x_2^{2n-i-1} x_4^3 = -f_{(4,i+1)} x_4^3, \quad \text{for } 0 \leq i \leq n-2. \end{aligned}$$

(4) $S(f_1, g) = x_2^2 x_3^3 x_4^r - x_1^{r+3}$. Since $\gcd(Lm(f_1), Lm(g)) = 1$, so by lemma ?? $S(f_1, g) \rightarrow_{G_{nr}} 0$

- (5) $S(f_1, f_5) = -x_1^{r-n+3}x_2^{n-1}x_4^3 + x_4^{r+2}x_3^2 = -f_{(3,0)}(x_4^3)$
(6) $S(f_1, f_6) = x_1^n x_3^2 x_4^2 - x_1 x_2^n x_4^3 = f_{(2,1)}(x_1 x_4^2)$
(7) $S(f_1, f_7) = -x_1 x_2^{2n} x_4^3 + x_1^{2n-1} x_3^4 x_4 = f_{(2,1)}(x_1^n x_3^2 x_4 + x_1 x_4^2 x_2^n)$
(8) for $i < j$, we have

$$\begin{aligned} S(f_{(2,i)}, f_{(2,j)}) &= x_1^{j-i} x_2^{n-j+1} x_4^{3j-2} - x_2^{n-i+1} x_3^{3(j-i)} x_4^{3i-2} \\ &= f_1 \left(\sum_{l=0}^{j-i-1} (x_1^l x_2^{(n-i)-l} x_3^{3(j-i)-3(l+1)} x_4^{3i+3(l-2)}) \right) \end{aligned}$$

- (9) $S(f_{(2,i)}, f_{(3,j)}) = x_3^{3(i+j)+1} x_4^{r-1-3j} - x_1^{r-2n+3+i+j} x_2^{2n-i-j} x_4^{3i-2}$, Now we have,

$$\begin{aligned} S(f_{2i}, f_{3j}) &= f_{(3,0)}(-x_2 x_4^{3i-2}) + f_{2n}(x_3^2 x_4^{r-1-3j}), \quad \text{for } i+j = n \\ &= f_{(3,(i+j)-n-1)}(-x_1 x_4^{3i-2}) + f_{(2,n)}(x_3^{3(i+j)-3n+2} x_4^{(r-1-3j)}) \\ &\quad + f_1(x_3^{3(i+j)-3n-1} x_4^{r+3n-3j-3}), \quad \text{for } i+j > n \\ &= f_{(4,i+j)} x_4^{3i-2}, \quad \text{for } i+j < n \end{aligned}$$

- (10) $S(g, f_{(2,i)}) = x_1^{r-n+2+i} x_2^{n-i+1} x_4^{3i-2} - x_2 x_3^{3i-1} x_4^r = f_{(3,i-1)} x_2 x_4^{3i-2}$
for $1 \leq i \leq n-1$
for $i = n$, Since $\gcd(Lm(g), Lm(f_{(2,n)})) = 1$, we have $S(f_{(2,n)}, g) \rightarrow_{G_{nr}} 0$

- (11) $S(f_{(2,i)}, f_{(4,j)}) = x_3^{3(i+j)} x_4^{r+1-3j} - x_1^{r-3n+3+i+j} x_2^{3n-i-j+1} x_4^{3i-2}$, We have,

$$\begin{aligned} S(f_{(2,i)}, f_{(4,j)}) &= f_{(3,l-1)}(-x_3 x_4^{3i-1}) + f_7(-x_1^{r-3n+l+3} x_2^{n-l} x_4^{3i-2}) \quad l = i+j \leq n-2 \\ &= f_7(-x_1^{r-2n+2} x_2 x_4^{3i-2}) + f_{(3,n-2)}(-x_3 x_4^{3i-1}) \quad l = i+j = n-1 \\ &= g(-x_3 x_4^{3i-1}) + f_7(-x_1^{r-2n+3} x_4^{3i-2}) + f_{(2,n)}(x_3 x_4^{r+1-3j}) \quad i+j = n \\ &= f_{(4,(i+j)-n-1)}(-x_1 x_4^{3i-2}) + f_{(2,n)}(x_3^{3(i+j)-3n+1} x_4^{r+1-3j}) \\ &\quad + f_1(x_3^{3l-3n-2} x_4^{r-1-3j+3n}) \quad i+j \geq n+1 \end{aligned}$$

- (12) $S(f_{(2,i)}, f_5) = x_3^{3i-2} x_4^{r+2} - x_1^{r-2n+2+i} x_2^{2n-i+1} x_4^{3i-2} = -f_{(4,i-1)} x_4^{3i+1}$, $1 \leq i \leq n$.

(13)

$$\begin{aligned} S(f_{(2,i)}, f_6) &= x_2^{2n-i+2} x_4^{3i-2} - x_1^{2n-i} x_4^2 x_3^{3i-2} \\ &= f_{(2,1)}(-x_1^{n-i+1} x_4^2 x_3^{3i-4} - x_2^{n-i+2} x_4^{3i-3}) \\ &\quad - (x_2 x_3^3 - x_1 x_4^3) \left(\sum_{l=1}^{i-2} (x_1^{n-i+l} x_2^{(n-l)} x_3^{3i-(7+3(l-1))} x_4^{3l}) \right), \quad 3 \leq i \leq n \end{aligned}$$

$$\text{for } i = 2, S(f_{(2,2)}, f_6) = f_{(2,1)}(x_1^{n-i+1} x_4^2 x_3^{3i-4} + x_2^{n-i+2} x_4^{3i-3})$$

for $i = 1$, $S(f_{(2,1)}, f_6) = f_7 x_4$

(14) $S(f_{(2,i)}, f_7) = -x_1^{n-i} x_2^{3n-i+2} x_4^{3i-2} + x_1^{3n-i-1} x_4 x^{3i}$, $1 \leq i \leq n$.

Since $\gcd(Lm(f_{(2,i)}), Lm(f_7)) = 1$, we have $S(f_{(2,i)}, f_7) \xrightarrow{G_{nr}}$

0

(15)

$$S(f_{(3,i)}, f_{(3,j)}) = -x_3^{2+3i} x_4^{r-1-3i} x_1^{j-i} + x_2^{j-i} x_3^{2+3j} x_4^{r-1-3j}$$

$$= x_2 x_3^3 - x_1 x_4^3 \left(\sum_{l=0}^{j-i-1} (x_1^l x_2^{(j-i)-(l+1)} x_3^{3j-(1+3l)} x_4^{r-3j+(-1+3l)}) \right), \quad i < j$$

(16) $S(f_{(3,i)}, g) = x_2^{n-i} x_4^r - x_1^{n-1-i} x_3^{2+3i} x_4^{r-1-3i} = -f_{(2,i+1)} x_4^{r-1-3i}$

(17) $S(f_{(3,i)}, f_{(4,j)}) = x_1^{n+1-j} x_3^{3j+1} x_4^{r+1-3j} - x_2^{n-j+1+i} x_3^{2+3i} x_4^{r-1-3i}$, Now we have,

$$S(f_{(3,i)}, f_{(4,j)}) = f_1 (-x_2^n x_3^{3i-1} x_4^{r-1-3i} + (x_1^{n-1} x_3^2 - x_2^n x_4) x_1 x_3^{3i-1} x_4^{r-3i+1}), \quad i = j$$

$$= f_{(2,l)} x_3^{3i+2} x_4^{r+1-3j}, \quad i < j, l = j - i$$

$$= f_{(2,1)} x_1^{(i-j)+1} x_3^{3j-1} x_4^{r+1-3j} + (-f_1) \sum_{l=0}^{i-j} x_1^l x_2^{n-j+i-l} x_3^{3i-1-3l} x_4^{r-1-3i+3l}, \quad i > j$$

(18) $S(f_{(3,i)}, f_5) = x_1^{1+i} x_4^{r+2} - x_2^{1+i} x_3^{3+3i} x_4^{r-1-3i} = -f_1 (\sum_{l=0}^i x_1^l x_2^{i-l} x_3^{3i-3l} x_4^{r-3i+3l-1})$,
 $0 \leq i \leq n - 2$

(19)

$$S(f_{(3,i)}, f_6) = x_1^{r+3+i} x_4^2 - x_2^{2+i} x_3^{3+3i} x_4^{r-1-3i}$$

$$= g x_1^{1+i} x_4^2 - f_1 \left(\sum_{l=0}^i x_1^l x_2^{1+(i-l)} x_3^{3(i-l)} x_4^{r-3i+3l-1} \right)$$

(20)

$$S(f_{(3,i)}, f_7) = x_1^{r+n+2+i} x_3 x_4 - x_2^{n+i+2} x_3^{2+3i} x_4^{r-1-3i}$$

$$= g(x_1^{n+i} x_3 x_4) + f_6(-x_1^i x_2 x_3 x_4^{r-1})$$

$$- f_1 \left(\sum_{l=0}^{i-2} (x_1^l x_2^{n+i+(1-l)} x_3^{3i-(3l+1)} x_4^{r-3i-1+3l}) \right), \quad 0 \leq i < n - 1$$

(21)

$$\begin{aligned}
S(g, f_{(4,i)}) &= -f_7 x_4^r, \quad i = 0 \\
&= f_{(2,1)}(x_1^{n-i} x_3^{3i-1} x_4^{r+1-3i} + x_2 x_4^{r-1}), \quad i = 1 \\
&= f_{(2,1)}(x_1^{n-i} x_3^{3i-1} x_4^{r+1-3i} + x_2^{n-i+1} x_4^{r-1}) \\
&\quad + f_1 \left(\sum_{l=0}^{i-2} x_1^{n-i+l} x_2^{n-1-l} x_3^{3i-4-3l} x_4^{r+2-3i+3l} \right), \quad i \geq 2.
\end{aligned}$$

(22) $S(g, f_5) = x_1^n x_4^{r+2} - x_2^{n+1} x_3 x_4^r = f_6(-x_4^r)$

(23) $S(g, f_6) = -x_2^{n+2} x_3 x_4^r + x_1^{r+n+2} x_4^2$

Since $\gcd(Lm(g), Lm(f_6)) = 1$, therefore $S(g, f_6) \rightarrow_G 0$

(24) Since $\gcd(Lm(g), Lm(f_7)) = 1$, therefore $S(g, f_7) \rightarrow_G 0$

(25)

$$\begin{aligned}
S(f_{(4,i)}, f_{(4,j)}) &= x_2^{j-i} x_3^{3j+1} x_4^{r+1-3j} - x_1^{j-i} x_3^{3i+1} x_4^{r+1-3i} \\
&= f_1 \left(\sum_{l=0}^{j-i-1} x_1^l x_2^{(j-i)-(l+1)} x_3^{3j-(2+3l)} x_4^{r-3j+(1+3l)} \right), \quad i < j
\end{aligned}$$

(26) $S(f_{(4,i)}, f_5) = x_2^{n-i} x_4^{r+2} - x_1^{n-1-i} x_3^{3i+2} x_4^{r+1-3i} = f_{(2,i+1)}(-x_4^{r+1-3i}), \quad 0 \leq i \leq n-1$

(27) $S(f_{(4,i)}, f_6) = x_1^{r-n+3+i} x_2^{n-i-1} x_4^2 - x_3^{3i+2} x_4^{r+1-3i} = x_4^2(f_{(3,i)}), \quad 0 \leq i < n-1$

for $i = n-1$, $S(f_{(4,i)}, f_6) = g x_4^2 - f_{(2,n)} x_4^{r-3n+4}$

(28)

$$\begin{aligned}
S(f_{(4,i)}, f_7) &= x_1^{r+2+i} x_3 x_4 - x_2^{1+i} x_3^{3i+1} x_4^{r+1-3i} \\
&= g(x_1^i x_3 x_4) - f_1 \left(\sum_{l=0}^{i-1} x_1^l x_2^{i-l} x_3^{3i+(-3l-2)} x_4^{r-3i+1+3l} \right), \quad 0 \leq i \leq n-1
\end{aligned}$$

(29) $S(f_5, f_6) = x_1^{r+2} x_4^2 - x_2 x_4^{r+2} = g x_4^2$

(30) $S(f_5, f_7) = -x_2^{n+1} x_4^{r+2} + x_1^{r+n+1} x_3^2 x_4 = f_{(2,1)}(x_1^{r+2} x_4 + x_2 x_4^{r+1} + f_{(3,0)} x_1^{n-1} x_2 x_4^2)$

(31) $S(f_6, f_7) = x_1^{2n-1} x_3^2 x_4 - x_1^n x_2^n x_4^2 = f_{(2,1)} x_1^n x_4$

By Buchberger's Criterion S-polynomials of all generator reduces to zero, so G_{nr} is Gröbner basis of \mathfrak{P}_{nr} with respect to degree reverse lexicographic monomial order $>$ induced by $x_1 > x_2 > x_3 > x_4$

□

Corollary 2.5. *Let us consider the degree reverse lexicographic monomial order on $k[x_1, \dots, x_4]$ induced by $x_1 > x_2 > x_3 > x_4$ then with respect to this order $G(\text{in}_{<}(\mathfrak{P}_{nr}))$ is the set*

$\{x_2x_3^3, x_1^{n-i}x_3^{3i-1}, x_1^{r-n+3+j}x_2^{n-1-j}, x_1^{r-2n+3+j}x_2^{2n-j}, x_1^{r-n+2}x_2^n x_3, x_2^{n+1}x_3, x_2^{2n+1} \mid 1 \leq i \leq n, 0 \leq j \leq n-1\}$.

Proof. Follows from the theorem 2.4. \square

We will take two theorems from [4].

Lemma 2.6. *Let I be an ideal in $A = k[x_1, \dots, x_r]$ and $I^H \subset A[x_0]$ its homogenization w.r.t. the variable x_0 . Let $<$ be any reverse lexicographic monomial order on A and $<_0$ the reverse lexicographic monomial order on $A[x_0]$ extended from A such that $x_i > x_0$ is the least variable.*

If $\{f_1, \dots, f_n\}$ is the reduced Gröbner basis for I w.r.t $<$, then $\{f_1^H, \dots, f_n^H\}$ is the reduced Gröbner basis for I^H w.r.t $<_0$, and $\text{in}_{<_0}(I^H) = (\text{in}_{<}(I))A[x_0]$.

Proof. See lemma 2.1 in [4]. \square

Theorem 2.7. *Let $\mathbf{n} : n_1, \dots, n_r$ be a sequence of positive integers with $n_r > n_i$ for all $i < n$. Let $<$ any reverse lexicographic order on $A = k[x_1, \dots, x_r]$ such that $x_i > x_r$ for all $1 \leq i < r$. and $<_0$ the induced reverse lexicographic order on $A[x_0]$, where $x_n > x_0$. Then the following conditions are equivalent:*

- (i) *The projective monomial curve $\overline{C(n_1, \dots, n_r)}$ is arithmetically Cohen-Macaulay.*
- (ii) *$\text{in}_{<_0}((\mathfrak{p}(n_1, \dots, n_r))^H)$ (homogenization w.r.t. x_0) is a Cohen-Macaulay ideal.*
- (iii) *$\text{in}_{<}(\mathfrak{p}(n_1, \dots, n_r))$ is a Cohen-Macaulay ideal.*
- (iv) *x_r does not divide any element of $G(\text{in}_{<}(\mathfrak{p}(n_1, \dots, n_r)))$.*

Proof. See theorem 2.2 in [4]. \square

Theorem 2.8. *The projective closure \overline{B}_{nr} of Bakelin's curve is arithmetically Cohen-Macaulay.*

Proof. From the corollary 2.5, we see that x_4 does not divide any element of $G(\text{in}_{<}(\mathfrak{P}_{nr}))$. The proof follows from the theorem 2.7. \square

Lemma 2.9. *The reduced Gröbner basis of the ideal \overline{P}_{nr} w.r.t. reverse lexicographic monomial order on $k[x_0, \dots, x_4]$ induced by $x_1 > x_2 > x_3 > x_4 > x_0$ is $\overline{G}_{nr} = \{f^H \mid f \in G_{nr}\}$. Where f^H is the homogenization of f w.r.t. the variable x_0 and G_{nr} is same as in 2.3.*

Proof. Follows from the lemma 2.6 \square

3. HILBERT SERIES OF BAKELIN'S CURVE

Lemma 3.1. *$I \subset S$ be a graded ideal and $<$ a monomial order on S . Then S/I and $S/in_{<}(I)$ have the same Hilbert function, i.e. $H(S/I, i) = H(S/in_{<}(I), i)$ for all i .*

Lemma 3.2. *Let $I = (x^{A_1}, \dots, x^{A_r}) \subset K[x_1, \dots, x_n]$ be a monomial ideal. Let $p(I)$ denote the numerator of the Hilbert series of $K[x_1, x_2, \dots, x_n]/I$, and let $|A|$ denote the total degree of the monomial x^A . Then*

$$p(I) = p(x^{A_1}) - \sum_{i=2}^{i=r} t^{|A_i|} p(x^{A_1}, \dots, x^{A_{i-1}} : x^{A_i}).$$

Theorem 3.3. *The numerator of the Hilbert series of the B_{nr} is*

$$1 - nt^{r+2} - 2t^{r+3} + (3n+4)t^{r+4} - (2n+2)t^{r+5} - t^{2n+1} - t^{n+2} + 2t^{n+2} - t^{n+1} + t^{n+3} + t^{n+4} - t^{2n+3} - \sum_{i=2}^n (t^{n+2i-1} + t^{n+2i+1} - 2t^{n+2i})$$

Proof. : From Corollary 3.5, $in_{<}(\mathfrak{P}_{nr})$ is generated by the set $\{x_2x_3^3, x_1^{n-i}x_3^{3i-1}, x_1^{r-n+3+j}x_2^{n-1-j}, x_1^{r-2n+3+j}x_2^{2n-j}, x_1^{r-n+2}x_2^n x_3, x_2^{n+1}x_3, x_2^{2n+1} \mid 1 \leq i \leq n, 0 \leq j \leq n-1\}$. We will use algorithm 2.6 of [11]. Rearrange $in_{<}(\mathfrak{P}_{nr})$ so that they are in ascending lexicographic order on the reversed set of variables $x_4 > x_3 > x_2 > x_1$. We have $in_{<}(\mathfrak{P}_{nr}) = \{x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n}, x_2^{2n+1}, x_1^{r-n+2}x_2^n x_3, x_2^{n+1}x_3, x_1^{n-1}x_3^2, x_1x_3^3, x_1^{n-2}x_3^5, \dots, x_3^{3n-1}\}$.

By Lemma 3.10, Hilbert function of $k[x_1, x_2, x_3, x_4]/\mathfrak{P}_{nr}$ is equal to the Hilbert function of $k[x_1, x_2, x_3, x_4]/in_{<}(\mathfrak{P}_{nr})$, it is sufficient to compute the Hilbert function of the latter. Let I denote the monomial ideal $in_{<}(\mathfrak{P}_{nr})$. We apply Lemma 3.2 to the ideal I , then

$$\begin{aligned}
p(I) &= p(x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n}, x_2^{2n+1}, \\
&\quad , x_1^{r-n+2}x_2^n x_3, x_2^{n+1}x_3, x_1^{n-1}x_3^2, x_2x_3^3, x_1^{n-2}x_3^5, \dots, x_3^{3n-1}) \\
&= p(x_1^{r+2}) - t^{r+2}p((x_1^{r+2} : x_1^{r+1}x_2)) - \dots - t^{r+2}p((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+4}x_2^{n-2} : x_1^{r-n+3}x_2^{n-1})) \\
&\quad t^{r+3}p((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1} : x_1^{r-n+2}x_2^{n+1})) \\
&\quad - \dots - t^{r+3}p((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+4}x_2^{2n-1} : x_1^{r-2n+3}x_2^{2n})) \\
&\quad - t^{2n+1}p(((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n} : x_2^{2n+1})) \\
&\quad - t^{r+3}p((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n}, x_2^{2n+1} : x_1^{r-n+2}x_2^n x_3)) \\
&\quad - t^{n+2}p(((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n}, x_2^{2n+1}, \\
&\quad x_1^{r-n+2}x_2^n x_3 : x_2^{n+1}x_3) \\
&\quad - t^{n+1}p(((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n}, x_2^{2n+1}, x_1^{r-n+2}x_2^n x_3, \\
&\quad x_2^{n+1}x_3 : x_1^{n-1}x_3^2)) \\
&\quad - t^4p(((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n}, x_2^{2n+1}, x_1^{r-n+2}x_2^n x_3 \\
&\quad , x_2^{n+1}x_3, x_1^{n-1}x_3^2 : x_2x_3^3)) \\
&\quad - t^{n+3}p((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n}, x_2^{2n+1}, x_1^{r-n+2}x_2^n x_3, \\
&\quad x_2^{n+1}x_3, x_1^{n-1}x_3^2, x_2x_3^3 : x_1^{n-2}x_3^5)) \\
&\quad - \dots - t^{3n-1}p((x_1^{r+2}, x_1^{r+1}x_2, \dots, x_1^{r-n+3}x_2^{n-1}, x_1^{r-n+2}x_2^{n+1}, \dots, x_1^{r-2n+3}x_2^{2n}, x_2^{2n+1}, x_1^{r-n+2}x_2^n x_3, \\
&\quad x_2^{n+1}x_3, x_1^{n-1}x_3^2, x_2x_3^3, x_1^{n-2}x_3^5, \dots, x_1x_3^{3n-4} : x_3^{3n-1})) \\
&= (1 - t^{r+2}) - (n-1)t^{r+2}(1-t) - nt^{r+3}(1-t) - t^{2n+1}(1 - t^{r-2n+3} - t^{r+3}(1+t^2-2t) \\
&\quad - t^{n+2}(1-t^n - nt^{r-n+2} + nt^{r-n+3}) - t^{n+1}(1-t^{n+1}) - (n+1)t^{r-n+3} + (n+1)t^{r-n+4}) \\
&\quad - t^4(1-t^{n-1} - t^n + t^{2n-1}) - (1+t^2-2t)\left(\sum_{i=2}^n t^{n+2i-1}\right) \\
&= 1 - t^{r+2} - (n-1)t^{r+2} + (n-1)t^{r+3} - nt^{r+3} + nt^{r+4} - t^{2n+1} + t^{r+4} - t^{r+3} - t^{r+5} + 2t^{r+4} \\
&\quad - t^{n+2} + t^{2n+2} + nt^{r+4} - nt^{r+5} - t^{n+1} + t^{2n+2} + (n+1)t^{r+4} - (n+1)t^{r+5} - t^4 + t^{n+3} \\
&\quad + t^{n+4} - t^{2n+3} - \sum_{i=2}^n (t^{n+2i-1} + t^{n+2i+1} - 2t^{n+2i}) \\
&= 1 - nt^{r+2} - 2t^{r+3} + (3n+4)t^{r+4} - (2n+2)t^{r+5} - t^{2n+3} + 2t^{2n+2} - t^{2n+1} + t^{n+4} + t^{n+3} \\
&\quad - t^{n+2} - t^{n+1} - t^4 - \sum_{i=2}^n (t^{n+2i-1} + t^{n+2i+1} - 2t^{n+2i}).
\end{aligned}$$

Hence the Hilbert Series of the ring $k[x_1, x_2, x_3, x_4]/I$ is $\frac{p(I)}{(1-t)^4}$. \square

4. SYZYGIES OF PROJECTIVE CLOSURE OF BRESINSKY'S CURVE

In [5], Bresinsky defined the following family of curves, for $h \geq 2$,

$$C((2h-1)2h, (2h-1)(2h+1), 2h(2h+1), 2h(2h+1) + 2h-1).$$

He showed that minimal number of generators of the defining ideal, that is $\mu(\mathfrak{p}((2h-1)2h, (2h-1)(2h+1), 2h(2h+1), 2h(2h+1) + 2h-1))$ is an unbounded function of h . Gröbner basis and syzygies of Bresinsky's curves has been studied in [10]. It has been proved in [10] that all betti numbers are unbounded function of h . Recently Herzog and Stamate proved in [4] that Bresinsky's curves are not arithmetically Cohen-Macaulay using Gröbner basis criterion. Here we will study syzygies of projective closure of Bresinsky's curves and will show that all betti numbers are not unbounded function of h , infact type is 1.

Notations 4.1. For $h \geq 2$, we define the following notations,

- let $B_h = C((2h-1)2h, (2h-1)(2h+1), 2h(2h+1), 2h(2h+1) + 2h-1)$
- $\mathfrak{Q}_h = \mathfrak{p}((2h-1)2h, (2h-1)(2h+1), 2h(2h+1), 2h(2h+1) + 2h-1)$.

Theorem 4.1. Consider the degree reverse lexicographic monomial order induced by $x_1 > x_2 > x_3 > x_4$ in the polynomial ring $k[x_1, x_2, x_3, x_4]$. For $h \geq 2$, let us consider the following polynomials

- $p_1 = x_2x_3 - x_1x_4$
- $p_2 = x_2^{2h} - x_3^{2h-1}$
- $p_{(3,j)} = x_1^{j+1}x_3^{2h-l} - x_2^jx_4^{2h-l}, 0 \leq j \leq 2h-1$
- $p_{(4,i)} = x_1^{i+1}x_2^{2h-i} - x_3^{i-1}x_4^{2h-i}, 1 \leq i \leq 2h$
- $p_5 = x_3^{4h} - x_2^{2h-1}x_4^{2h+1}$
- $p_{(6,i)} = x_1^{2+i}x_2^{2h-2-i}x_4^{2+i} - x_3^{2h+1+i}, 0 \leq i \leq 2h-3$
- $p_7 = x_1x_2^{2h-1}x_4 - x_3^{2h}$
- $p_8 = x_1^{2h}x_4^{2h} - x_3^{4h-1}$.

Let $\mathfrak{G}_h = \{p_1, p_2, p_5, p_7, p_8\} \cup \{p_{(3,j)} \mid 0 \leq j \leq 2h-1\} \cup \{p_{(4,i)} \mid 1 \leq i \leq 2h\}$, then \mathfrak{G}_h is Gröbner basis of \mathfrak{Q}_h with respect to the given monomial order.

Proof. We proceed by Buchberger's algorithm. We compute S -polynomials and show that it reduces to zero upon division by \mathfrak{G}_h .

$$(1) S(p_1, p_2) = x_3^{2h} - x_1x_2x_2^{2h-1}x_4 = -p_7$$

(2)

$$\begin{aligned} S(p_1, p_{(3,j)}) &= x_2^{j+1}x_4^{2h-l} - x_1^{2+j}x_4x_3^{2h-j-1} = -x_4p_{(3,l+1)}, \text{ for } \leq j < 2h-1 \\ &= x_2^{2h}x_4 - x_1^{2h+1}x_4 = x_4p_2 - x_4p_{(4,2h)}, \text{ when } i = 2h-1. \end{aligned}$$

$$(3) \ S(p_1, p_{(4,i)}) = x_3^i x_4^{2h-i} - x_1^{i+2} x_2^{2h-i-1} x_4 = -x_4 p_{(4,i+1)}, \text{ for } 1 \leq i < 2h$$

$$S(p_1, p_{(4,2h)}) = x_2 x_3^{2h} - x_1^{2h+2} x_4, \text{ since } \gcd(\text{Lt}(p_1), \text{Lt}(p_{(4,2h)})) = 1, S(p_1, p_{(4,2h)}) \longrightarrow_{\mathfrak{G}_h} 0$$

$$(4) \ S(p_1, p_5) = x_2^{2h} x_4^{2h+1} - x_1 x_3^{4h-1} x_4 = p_2 x_4^{2h+1} - p_{(3,0)} x_3^{2h-1} x_4$$

$$(5)$$

$$S(p_1, p_{(6,i)}) = x_3^{2h+2+i} - x_1^{3+i} x_2^{2h-3-i} x_4^{3+i} = -p_{(6,i+1)}, 0 \leq i \leq 2h-4 \\ = -p_8, i = 2h-3$$

$$(6) \ S(p_1, p_7) = x_3^{2h+1} - x_1^2 x_2^{2h-2} x_4^2 = -p_{(6,0)}$$

$$(7) \ S(p_1, p_8) = x_2 x_3^{4h} - x_1^{2h+1} x_4^{2h+1} = x_3^{4h-1} p_1 + p_{(3,0)} x_3^{2h-1} x_4 - x_4^{2h+1} p_{(4,2h)}$$

$$(8) \ \text{since } \gcd(\text{Lt}(p_2), \text{Lt}(p_{(3,l)})) = 1, S(p_2, p_{(3,l)}) \longrightarrow_{\mathfrak{G}_h} 0$$

$$(9)$$

$$S(p_2, p_{(4,i)}) = x_2^i x_3^{i-1} x_4^{2h-1} - x_1^{i+1} x_3^{2h-1} \\ = p_{(3,1)} (-x_1^{i-1}) + p_1 \left(\sum_{l=0}^{i-2} x_1^l x_2^{i-1-l} x_3^{i-2-l} x_4^{2h-i+l} \right), \quad 2 \leq i \leq 2h \\ = -p_{(3,1)}, \quad i = 1.$$

$$(10) \ \text{Since } \gcd(\text{Lt}(p_2), \text{Lt}(p_{(5)})) = 1, S(p_2, p_5) \longrightarrow_{\mathfrak{G}_h} 0$$

$$(11)$$

$$S(p_2, p_{(6,i)}) = x_2^{2+i} x_3^{2h+1+i} - x_1^{2+i} x_3^{2h-1} x_4^{2+i} \\ = -p_1 \left(\sum_{l=0}^{i+1} x_1^l x_2^{1+i-l} x_3^{2h+i-l} x_4^l \right), \quad 0 \leq i \leq 2h-3.$$

$$(12) \ S(p_2, p_7) = x_2 x_3^{2h} - x_1 x_3^{2h-1} x_4 = p_1 x_3^{2h-1}$$

$$(13) \ S(p_2, p_8) = x_2^{2h} x_3^{4h-1} - x_1^{2h} x_3^{2h-1} x_4^{2h} = p_1 \left(\sum_{l=0}^{2h-1} x_1^l x_2^{2h-1-l} x_3^{4h-2-l} x_4^l \right)$$

$$(14)$$

$$S(p_{(3,i)}, p_{(3,j)}) = x_2^j x_3^{j-i} x_4^{2h-j} - x_1^{j-i} x_2^i x_4^{2h-i} \\ = p_1 \left(\sum_{r=0}^{j-i-1} x_1^r x_2^{j-1-r} x_3^{j-i-2-r} x_4^{2h-j+r} \right).$$

(15)

$$\begin{aligned}
S(p_{(3,j)}, p_{(4,i)}) &= -x_2^{2h} x_4^{2h-i} + x_3^{2h-1} x_4^{2h-i} = p_2(-x_4^{2h-i}), \text{ for } j = i \\
&= x_3^{2h-j+i-1} x_4^{2h-i} - x_1^{i-j} x_2^{2h-i+j} x_4^{2h-j} = -x_4^{2h-i} p_{(6,i-j-2)}, \text{ for } j < i \\
&= x_1^{j-i} x_3^{2h+i-j-1} x_4^{2h-i} - x_2^{2h-i+j} x_4^{2h-i} = p_2(-x_2^{j-i} x_4^{2h-j}) \\
&\quad + p_1\left(-\sum_{k=0}^{j-i-1} x_1^k x_2^{j-i-1-k} x_3^{2h-2-k} x_4^{2h-j+k}\right), \text{ for } j > i.
\end{aligned}$$

(16)

$$\begin{aligned}
S(p_{(3,j)}, p_5) &= x_1^{1+j} x_2^{2h-1} x_4^{2h+1} - x_2^j x_3^{2h+j} x_4^{2h-j} \\
&= p_7 x_1^j x_4^{2h} - p_1\left(\sum_{k=0}^{j-1} x_1^k x_2^{j-1-k} x_3^{2h+j-1-k} x_4^{2h-j+k}\right)
\end{aligned}$$

(17)

$$\begin{aligned}
S(p_{(3,j)}, p_{(6,i)}) &= x_3^{4h} - x_1 x_2^{2h-2} x_4^{2h+2} = p_5 x_3 + p_1 x_2^{2h-2} x_4^{2h+1} \text{ for } j = i \\
&= x_1^{j-i-1} x_3^{4h+1+i-j} - x_2^{2h-2-i+j} x_4^{2h-j+i-2} = p_5 \text{ for } j = i + 1 \\
&= p_{(3,0)}(x_1^{j-i-2} x_3^{2h+1+i-j}) - p_2(x_2^{j-i-2} x_4^{2h-j+i+2}) \\
&\quad - p_1\left(\sum_{k=0}^{j-i-3} x_1^k x_2^{j-i-3-k} x_3^{2h-2-k} x_4^{2h-j+i+2+k}\right), \text{ for } j > i + 1 \\
&= p_5 x_3^{1+i-j} + p_1\left(\sum_{k=0}^{i-j} (x_1^k x_2^{2h-2-k} x_3^{i-j-k} x_4^{2h+1+k})\right), \text{ for } j < i.
\end{aligned}$$

(18)

$$\begin{aligned}
S(p_{(3,j)}, p_7) &= -x_2^{2h+j-1} x_4^{2h-j+1} + x_1^j x_3^{4h-j} \\
&= p_{(3,0)} x_1^{j-1} x_3^{2h-l} + p_2(-x_2^{j-1} x_4^{2h-j+1}) \\
&\quad - p_1\left(\sum_{k=0}^{j-2} x_1^k x_2^{j-2-k} x_3^{2h-2-k} x_4^{2h-j+1+k}\right), \text{ for } 0 < j \leq 2h - 1 \\
&= p_5, \text{ for } j = 0
\end{aligned}$$

(19)

$$\begin{aligned}
S(p_{(3,j)}, p_8) &= x_3^{6h-j-1} - x_1^{2h-j-1} x_2^j x_4^{4h-l} \\
&= p_5 x_3^{2h-j-1} + p_1\left(\sum_{k=0}^{2h-j-2} x_1^k x_2^{2h-2-k} x_3^{2h-j-2-k} x_4^{2h+1+k}\right).
\end{aligned}$$

$$(20) \quad S(p_{(4,i)}, p_{(4,j)}) = x_2^{j-i} x_3^{j-i} x_4^{2h-j} - x_1^{j-i} x_3^{i-1} x_4^{2h-i} = p_1 \left(\sum_{k=0}^{j-i-1} x_1^k x_2^{j-i-1-k} x_3^{j-2-k} x_4^{2h-j+k} \right).$$

$$(21) \quad \text{Since } \gcd(\text{Lt}(p_{(4,i)}), \text{Lt}(p_5)) = 1, \quad S(p_{(4,i)}, p_5) \longrightarrow_{\mathfrak{G}_h} 0.$$

$$(22)$$

$$\begin{aligned} S(p_{(4,i)}, p_{(6,j)}) &= x_2^2 x_3^{2h+1+i} - x_1 x_4^{2+i} x_3^{i-1} x_4^{2h-i} \\ &= p_1 (x_2 x_3^{2h+i} + x_1 x_3^{2h+i-1} x_4 + p_{(3,0)} x_1 x_3^{i-1} x_4^2, \text{ for } i = j) \\ &= x_2^{j+2-i} x_3^{2h+1+j} - x_1^{1+j-i} x_3^{i-1} x_4^{2h-i+j+2} \\ &= p_1 \left(\sum_{l=0}^{j-i+1} x_1^l x_2^{j+1-i-l} x_3^{2h+j-l} x_4^l \right) + p_{(3,0)} x_1^{1+j-i} x_3^{i-1} x_4^{j-i+2}, \text{ for } i < j \\ &= x_1^{i-j-1} x_3^{2h+1+j} - x_2^{i-j-2} x_3^{i-1} x_4^{2h-i+j+2} \\ &= p_{(3,0)} x_1^{i-j-2} x_3^{1+j} - p_1 \left(\sum_{l=0}^{i-j-3} x_1^l x_2^{i-j-3-l} x_3^{i-2-l} x_4^{2h-i+j+2+l} \right), \text{ for } i \geq j+2 \\ &= x_2 x_3^{2h+1+j} - x_3^j x_4^{2h+1} \\ &= p_1 x_3^{2h+j} - p_{(3,0)} x_3^j x_4, \text{ for } i = j+1. \end{aligned}$$

$$(23)$$

$$\begin{aligned} S(p_{(4,i)}, p_7) &= x_3^{2h} x_1^i - x_2^{i-1} x_3^{i-1} x_4^{2h-i+1} = p_{(3,0)}, \text{ for } i = 1 \\ &= p_{(3,0)} x_1^{i-1} + p_1 \left(- \sum_{l=0}^{i-2} x_1^l x_2^{i-2-l} x_3^{i-2-l} x_4^{2h-i+1+l} \right), \text{ for } 1 < i \leq 2h. \end{aligned}$$

$$(24)$$

$$\begin{aligned} S(p_{(4,i)}, p_8) &= x_2^{2h-i} x_3^{4h-1} - x_1^{2h-i-1} x_3^{i-1} x_4^{4h-i} \\ &= p_1 \left(\sum_{l=0}^{2h-i-1} x_1^l x_2^{2h-i-1-l} x_3^{4h-2-l} x_4^l \right) + p_{(3,0)} x_1^{2h-i-1} x_3^{i-1} x_4^{2h-i}, \text{ for } 1 \leq i < 2h \\ &= x_1 x_3^{4h-1} - x_3^{2h-1} x_4^{2h} = p_{(3,0)} x_3^{2h-1}, \text{ for } i = 2h. \end{aligned}$$

$$(25)$$

$$\begin{aligned} S(p_{(6,i)}, p_{(6,j)}) &= x_2^{j-i} x_3^{2h+1+j} - x_1^{j-i} x_3^{2h+1+i} x_4^{j-i} \\ &= p_1 \left(\sum_{l=0}^{j-i-1} x_1^l x_2^{j-i-1-l} x_3^{2h+j-l} x_4^l \right), \text{ for } i < j. \end{aligned}$$

$$(26) \quad S(p_{(6,i)}, p_7) = x_1^{1+i} x_4^{1+i} x_3^{2h} - x_2^{1+i} x_3^{2h+1+i} = -p_1 \left(\sum_{l=0}^i x_1^l x_2^{i-l} x_3^{2h+i-l} x_4^l \right)$$

(27)

$$\begin{aligned} S(p_{(6,i)}, p_8) &= x_2^{2h-2-i} x_3^{4h-1} - x_1^{2h-2-i} x_4^{2h-2-i} x_3^{2h+1+i} \\ &= p_1 \left(\sum_{l=0}^{2h-3-i} x_1^l x_2^{2h-3-i-l} x_3^{4h-2-l} x_4^l \right). \end{aligned}$$

$$(28) \quad S(p_7, p_8) = x_2^{2h-1} x_3^{4h-1} - x_1^{2h-1} x_3^{2h} x_4^{2h-1} = p_1 \left(\sum_{l=0}^{2h-2} x_1^l x_2^{2h-2-l} x_3^{4h-2-l} x_4^l \right)$$

We see that all S-polynomials reduce to zero upon division by \mathfrak{G}_h , therefore \mathfrak{G}_h is Gröbner basis of $\overline{\Omega}_h$ with respect to degree reverse lexicographic monomial order on $k[x_1, x_2, x_3, x_4]$ induced by $x_1 > x_2 > x_3 > x_4$. \square

Suppose for any $g \in k[x_1, x_2, x_3, x_4]$, we denote $g^H \in k[x_0, x_1, x_2, x_3, x_4]$ the homogenization of g w.r.t. the indeterminates x_0 .

Lemma 4.2. *For $h \geq 2$, let $\overline{\mathfrak{G}}_h := \{g^H \mid g \in \mathfrak{G}_h\}$. Then $\overline{\mathfrak{G}}_h$ is a Gröbner basis of the homogenized ideal $\overline{\Omega}_h$ with respect to degree reverse lexicographic monomial order on $k[x_0, x_1, x_2, x_3, x_4]$ induced by $x_1 > x_2 > x_3 > x_4 > x_0$.*

Proof. Follows from theorems 4.1 and 2.6. \square

Lemma 4.3. *The ideal $\overline{\Omega}_h$, for $h \geq 2$ is generated by following polynomials*

- $p_1^H = x_2 x_3 - x_1 x_4$
- $p_2^H = x_2^{2h} - x_0 x_3^{2h-1}$
- $p_{(3,j)}^H = x_1^{j+1} x_3^{2h-l} - x_0 x_2^j x_4^{2h-j}; 0 \leq j \leq 2h-1$
- $p_{(4,i)}^H = x_1^{i+1} x_2^{2h-i} - x_0^2 x_3^{i-1} x_4^{2h-i}; 1 \leq i \leq 2h$
- $p_5^H = x_3^{4h} - x_2^{2h-1} x_4^{2h+1}$

Proof. Let $\overline{\mathfrak{T}}_h = \{p_1^H, p_2^H, p_{(3,j)}^H, p_{(4,i)}^H, p_5^H \mid 0 \leq j \leq 2h-1, 1 \leq i \leq 2h\}$, for $h \geq 2$. Then $\overline{\mathfrak{T}}_h \subset \overline{\mathfrak{G}}_h$, for $h \geq 2$. We have \square

Notations 4.2. *For $h \geq 2$ we define the following matrices,*

$$(1) \quad \mathfrak{B}_h^1 := \left(p_1^H \quad p_2^H \quad p_{(3,0)}^H \quad \cdots \quad p_{(3,2h-1)}^H \quad p_{(4,1)}^H \quad \cdots \quad p_{(4,2h)}^H \quad p_5^H \right)_{1 \times (4h+3)}$$

$$(2) \quad \mathfrak{B}_h^2 := \left(\Omega_{ij} \right)_{(4h+3) \times (8h+4)}, \text{ where } \Omega_{ij}, 1 \leq i \leq 4h+3, 1 \leq j \leq 8h+4 \text{ are defined as follows,}$$

- $\Omega_{(2,1)} = x_0 x_4, \Omega_{(2h+2,1)} = x_2, \Omega_{(4h+1,1)} = -x_3$ and $\Omega_{(i,1)} = 0$, for $i \in [4h+3] \setminus \{2, 2h+2, 4h+1\}$.
- $\Omega_{(2,2)} = x_1^2, \Omega_{(4,2)} = x_2, \Omega_{(2h+3,2)} = -x_2$ and $\Omega_{(i,2)} = 0$, for $i \in [4h+3] \setminus \{2, 4, 2h+3\}$.

- $\Omega_{(1,3+l)} = x_0 x_2^l x_4^{2h-1-l}$, $\Omega_{(3+l,3+l)} = -x_1$, $\Omega_{(4+l,3+l)} = x_3$ and $\Omega_{(i,3+l)} = 0$, for $i \in [4h+3] \setminus \{1, 3+l, 4+l\}$, $0 \leq l \leq 2h-2$.
 - $\Omega_{(1,2h+1+l)} = x_0^2 x_3^{l-1} x_4^{2h-1-l}$, $\Omega_{(2h+2+l,2h+1+l)} = -x_1$, $\Omega_{(2h+3+l,2h+1+l)} = x_2$ and $\Omega_{(i,2h+1+l)} = 0$, for $i \in [4h+3] \setminus \{1, 2h+2+l, 2h+3+l\}$, $1 \leq l \leq 2h-1$.
 - $\Omega_{(1,4h+1+l)} = x_1^{l+1} x_3^{2h-1-l}$, $\Omega_{(3+l,4h+1+l)} = -x_2$, $\Omega_{(4+l,4h+1+l)} = x_4$ and $\Omega_{(i,4h+1+l)} = 0$, for $i \in [4h+3] \setminus \{1, 3+l, 4+l\}$, $0 \leq l \leq 2h-2$.
 - $\Omega_{(1,6h-1+l)} = x_1^{l+1} x_2^{2h-1-l}$, $\Omega_{(2h+2+l,6h-1+l)} = -x_3$, $\Omega_{(2h+3+l,6h-1+l)} = x_4$ and $\Omega_{(i,6h-1+l)} = 0$, for $i \in [4h+3] \setminus \{1, 2h+2+l, 2h+3+l\}$, $1 \leq l \leq 2h-1$.
 - $\Omega_{(1,8h-1)} = x_2^{2h} - x_0 x_3^{2h-1}$, $\Omega_{(2,8h-1)} = -x_2 x_3 + x_1 x_4$, and $\Omega_{(i,8h-1)} = 0$, for $i \in [4h+3] \setminus \{1, 2\}$.
 - $\Omega_{(1,8h)} = x_1 x_2^{2h-1}$, $\Omega_{(2,8h)} = -x_1 x_3$, $\Omega_{(3,8h)} = -x_0$, $\Omega_{(2h+3,8h)} = x_4$, and $\Omega_{(i,8h)} = 0$, for $i \in [4h+3] \setminus \{1, 2, 3, 2h+3\}$.
 - $\Omega_{(1,8h+1)} = x_0 x_2^{2h-1}$, $\Omega_{(2,8h+1)} = -x_0 x_3$, $\Omega_{(2h+2,8h+1)} = -x_1$, $\Omega_{(4h+2,8h+1)} = x_3$, and $\Omega_{(i,8h+1)} = 0$, for $i \in [4h+3] \setminus \{1, 2, 2h+2, 4h+2\}$.
 - $\Omega_{(1,8h+2)} = x_2^{2h-1} x_4^{2h}$, $\Omega_{(2,8h+2)} = -x_3 x_4$, $\Omega_{(3,8h+2)} = x_3^{2h}$, $\Omega_{(4h+3,8h+2)} = -x_1$, and $\Omega_{(i,8h+2)} = 0$, for $i \in [4h+3] \setminus \{1, 2, 3, 4h+3\}$.
 - $\Omega_{(1,8h+3)} = x_3^{4h-1}$, $\Omega_{(2,8h+3)} = -x_4^{2h+1}$, $\Omega_{(3,8h+3)} = x_3^{2h-1} x_4$, $\Omega_{(4h+3,8h+3)} = -x_2$, and $\Omega_{(i,8h+3)} = 0$, for $i \in [4h+3] \setminus \{1, 2, 3, 4h+3\}$.
 - $\Omega_{(1,8h+4)} = x_2^{2h-1} x_3^{2h}$, $\Omega_{(2,8h+4)} = -x_3^{2h+1}$, $\Omega_{(3,8h+4)} = x_2^{2h-1} x_4$, $\Omega_{(4h+3,8h+3)} = -x_0$, and $\Omega_{(i,8h+3)} = 0$, for $i \in [4h+3] \setminus \{1, 2, 3, 4h+3\}$.
- (3) $\mathfrak{B}_h^3 := (\Delta_{ij})_{(8h+4) \times (4h+3)}$, where Δ_{ij} , $1 \leq i \leq 8h+4$, $1 \leq j \leq 4h+3$ are defined as follows
- $\Delta_{(4h+l,l+1)} = x_1$, $\Delta_{(3+l,l+1)} = -x_2$, $\Delta_{(4h+l+1,l+1)} = -x_3$, $\Delta_{(4+l,l+1)} = x_4$, and $\Delta_{(i,l+1)} = 0$, for $i \in [8h+4] \setminus \{4h+l, 3+l, 4h+l+1, 4+l\}$, $0 \leq l \leq 2h-3$.
 - $\Delta_{(6h-1,2h-1)} = x_1$, $\Delta_{(2h+1,2h-1)} = -x_2$, $\Delta_{(8h-2,2h-1)} = -x_3$, $\Delta_{(1,2h-1)} = x_3$, $\Delta_{(8h+1,2h-1)} = x_4$ and $\Delta_{(i,2h-1)} = 0$, for $i \in [8h+4] \setminus \{6h-1, 2h+1, 8h-2, 1, 8h+1\}$
 - $\Delta_{(6h-1+l,2h-1+l)} = x_1$, $\Delta_{(6h+l,2h-1+l)} = -x_2$, $\Delta_{(2h+1+l,2h-1+l)} = -x_3$, $\Delta_{(2h+2+l,2h-1+l)} = x_4$, and $\Delta_{(i,2h-1+l)} = 0$, for $i \in [8h+4] \setminus \{6h-1+l, 6h+l, 2h+1+l, 2h+2+l\}$ $1 \leq l \leq 2h-2$
 - $\Delta_{(3,4h-2)} = x_0$, $\Delta_{(8h,4h-2)} = -x_1$, $\Delta_{(6h,4h-2)} = x_2$, $\Delta_{(2,4h-2)} = -x_3$, $\Delta_{(2h+2,4h-2)} = -x_4$ and $\Delta_{(i,4h-2)} = 0$, for $i \in [8h+4] \setminus \{3, 8h, 6h, 2, 2h+2\}$

- $\Delta_{(4h+1,4h-1)} = x_0, \Delta_{(8h-1,4h-1)} = x_1, \Delta_{(8h,4h-1)} = -x_2, \Delta_{(2,4h-1)} = -x_4, \text{ and } \Delta_{(i,4h-1)} = 0, \text{ for } i \in [8h+4] \setminus \{4h+1, 8h-1, 8h, 2\}$
- $\Delta_{(8h-1,4h)} = x_0, \Delta_{(1,4h)} = -x_1, \Delta_{(8h+1,4h)} = -x_2, \Delta_{(4h,4h)} = x_3, \text{ and } \Delta_{(i,4h)} = 0, \text{ for } i \in [8h+4] \setminus \{1, 8h-1, 8h+1, 4h\}$
- $\Delta_{(4h+1,4h+1)} = x_3^{2h}, \Delta_{(3,4h+1)} = -x_3^{2h-1}x_4, \Delta_{(8h-1,4h+1)} = -x_4^{2h}, \Delta_{(8h+3,4h+1)} = -x_1, \Delta_{(8h+2,4h+1)} = x_2 \text{ and } \Delta_{(i,4h+1)} = 0, \text{ for } i \in [8h+4] \setminus \{4h+1, 3, 8h-1, 8h+3, 8h+2\}$
- $\Delta_{(4h+2,8h-1)} = x_3^{2h}, \Delta_{(4h+1+i,4h+2)} = -x_2^{2h-1-i}x_4^{i+1}, \Delta_{(8h,4h+2)} = x_3^{2h-1}x_4, \Delta_{(6h-1+j,4h+2)} = x_3^{(2h-1-j)}x_4^{j+1}, \Delta_{(1,4h+2)} = x_4^{2h}, \Delta_{(8h+3,4h+2)} = x_0, \Delta_{(8h+4,4h+2)} = -x_2, \text{ and } \Delta_{(i,4h+2)} = 0, \text{ for } i \in [8h+4] \setminus \{8h-1, 4h+1+i, 8h, 6h-1+j, 1, 8h+3, 8h+4\}, 0 \leq i \leq 2h-2, 1 \leq j \leq 2h-2$
- $\Delta_{(8h,4h+3)} = x_3^{2h}, \Delta_{(3,4h+3)} = -x_2^{2h-1}x_4, \Delta_{(4h+1+i,4h+3)} = -x_2^{2h-1-i}x_3x_4^i, \Delta_{(6h-1+i,4h+3)} = x_3^{2h-i}x_4^i, \Delta_{(1,4h+3)} = x_3x_4^{2h-1}, \Delta_{(8h+2,4h+3)} = x_0, \Delta_{(8h+4,4h+3)} = -x_1, \text{ and } \Delta_{(i,4h+3)} = 0, \text{ for } i \in [8h+4] \setminus \{8h, 3, 4h+1+i, 6h-1+i, 1, 8h+2, 8h+4\}$

(4) $\mathfrak{B}_h^4 := (\Gamma_{ij})_{(4h+3) \times 1}$, where $\Gamma_{i1}, 1 \leq i \leq 4h+3$ are defined as follows

- $\Gamma_{(4h-1,1)} = x_3^{2h},$
- $\Gamma_{(m,1)} = -x_2^{2h-m}x_4^m, \text{ for } 1 \leq m \leq 2h-1$
- $\Gamma_{(4h-2,1)} = -x_3^{(2h-1)}x_4,$
- $\Gamma_{(2h-1+n,1)} = x_3^{2h-1-n}x_4^{n+1}, \text{ for } 1 \leq n \leq 2h-2$
- $\Gamma_{(4h,1)} = -x_4^{2h},$
- $\Gamma_{(4h+1,1)} = -x_0,$
- $\Gamma_{(4h+2,1)} = -x_1,$
- $\Gamma_{(4h+3,1)} = x_2,$
- $\Gamma_{(i,4h+1)} = 0, \text{ for } i \in \{1, \dots, 8h+4\} \setminus \{4h-1, i, 4h-2, 2h-1+j, 4h, 4h+1, 4h+2, 4h+3\}, 1 \leq i \leq 2h-1, 1 \leq j \leq 2h-2.$

Theorem 4.4. For $h \geq 2$, we consider the homogenized ideal $\overline{\mathfrak{Q}}_h$ (defining ideal of projective closure of Bresinsky's curve) in the polynomial ring $k[x_0, x_1, x_2, x_3, x_4]$. Then a minimal graded free resolution of $\overline{\mathfrak{Q}}_h$ is given by,

$$\mathfrak{B}_h : 0 \longrightarrow R \xrightarrow{\mathfrak{B}_h^4} R^{4h+3} \xrightarrow{\mathfrak{B}_h^3} R^{8h+4} \xrightarrow{\mathfrak{B}_h^2} R^{4h+3} \xrightarrow{\mathfrak{B}_h^1} R \longrightarrow R/\overline{\mathfrak{Q}}_h \longrightarrow 0$$

Where matrices $\mathfrak{B}^i, 1 \leq i \leq 4$ are defined in 4.2.

Proof. One can check that

$$\mathfrak{B}_h : 0 \longrightarrow R \xrightarrow{\mathfrak{B}_h^4} R^{4h+3} \xrightarrow{\mathfrak{B}_h^3} R^{8h+4} \xrightarrow{\mathfrak{B}_h^2} R^{4h+3} \xrightarrow{\mathfrak{B}_h^1} R \longrightarrow R/\overline{\mathfrak{Q}_h} \longrightarrow 0$$

is a chain complex by calculating $\mathfrak{B}_h^i \mathfrak{B}_h^{i+1} = 0, 1 \leq i \leq 3$. We proceed by Buchsbaum Eisenbud acyclicity criterion. Let r_i be the i th expected rank of \mathfrak{B}_h . Then $r_1 = (4h+3) - (8h+4) + (4h+3) - 1 = 1, r_2 = 8h+4 - (4h+3) + 1 = 4h+2, r_3 = 4h+3 - 1 = 4h+2, r_4 = 1$. We need to show that $\text{grade}(I_{r_i}(\mathfrak{B}_h^i)) \geq i, 1 \leq i \leq 4$ (where $I_{r_i}(\mathfrak{B}_h^i)$ denotes the ideal generated by $r_i \times r_i$ minors of the matrix \mathfrak{B}_h^i)

(1) We take $p_1^H \in I_{r_1}(\mathfrak{B}^1)$ and we have $I_{r_1}(\mathfrak{B}^1) \geq 1$.

(2) We take

- $\mathfrak{L}_h^{[21]} := [134 \dots (4h+3)|(4h+1) \dots (8h-2)8h(8h+2)(8h+3)(8h+4)] = x_4^{4h-1}(-x_2^{2h} + x_3^{2h-1}x_0)(x_3^{4h} - x_2^{2h-1}x_4^{2h+1})$
- $\mathfrak{L}_h^{[22]} := [2 \dots (4h+3)|13 \dots 4h(8h+1)(8h+2)(8h+4)] = -x_1^{2h-2}(-x_2x_3 + x_1x_4)(-x_1x_2^{2h-1}x_4 + x_3^{2h}x_0)(x_0x_3^{2h})$.

Since gcd of any two prime factors of $\mathfrak{L}_h^{[21]}$ and $\mathfrak{L}_h^{[22]}$ is 1, $\mathfrak{L}_h^{[21]}, \mathfrak{L}_h^{[22]}$ forms a regular sequence. Hence $\text{grade}(I_{r_2}(\mathfrak{B}_h^2)) \geq 2$

(3) We take

- $\mathfrak{L}_h^{[31]} := [1(4h) \dots (6h-3) \dots (8h-3)(8h-1)(8h)(8h+3)(8h+4)|12 \dots (4h+1)(4h+3)] = x_1^{4h+2},$
- $\mathfrak{L}_h^{[32]} := [1 \dots 4h(8h+1)(8h+2)|1 \dots (4h+2)] = x_4^{6h}(-x_2^{2h} + x_3^{2h-1}x_0)$
-

$$\mathfrak{L}_h^{[33]} := [3 \dots (2h+1)(4h+1)(6h) \dots 8h(8h+3)|1 \dots (4h+2)]$$

$$\begin{aligned} &= x_1x_2^{6h-3}x_4 - x_0 \left(\sum_{i=0}^{2h-1} x_1^i x_2^{4h-2-i} x_3^{2h-i} x_4^i \right) - x_0 x_1^{2h-1} x_2^{2h-1} x_3 x_4^{2h-1} \\ &+ x_0 x_1^{2h-1} x_2^{2h-2} x_3 x_4^{2h} + x_0^2 x_1^{2h-2} x_3^{2h+1} x_4^{2h-2} \end{aligned}$$

Let us consider the ideal $\langle \mathfrak{L}_h^{[31]}, \mathfrak{L}_h^{[32]} \rangle$. Then primary decomposition of this ideal is $\langle x_1^{4h+2}, x_4^{6h} \rangle \cap \langle x_1^{4h+2}, (-x_2^{2h} + x_3^{2h-1}x_0) \rangle$. Hence associated primes are $\langle x_1, x_4 \rangle, \langle x_1, (-x_2^{2h} + x_3^{2h-1}x_0) \rangle$. We observe that $\mathfrak{L}_h^{[33]} \notin \langle x_1, x_4 \rangle$ and $\mathfrak{L}_h^{[33]} \notin \langle x_1, (-x_2^{2h} + x_3^{2h-1}x_0) \rangle$. If $\mathfrak{L}_h^{[33]} \in \langle x_1, x_4 \rangle$ then $x_0 x_2^{4h-2} x_3^{2h} \in \langle x_1, x_4 \rangle$ which is a contradiction, if $\mathfrak{L}_h^{[33]} \in \langle x_1, (-x_2^{2h} + x_3^{2h-1}x_0) \rangle$ then again $x_0 x_2^{4h-2} x_3^{2h} \in \langle x_1, (-x_2^{2h} + x_3^{2h-1}x_0) \rangle$ a contradiction. Therefore $\{\mathfrak{L}_h^{[31]}, \mathfrak{L}_h^{[32]}, \mathfrak{L}_h^{[33]}\}$ forms a regular sequence hence $\text{grade}(I_{r_3}(\mathfrak{B}_h^3)) \geq 3$.

(4) We take

- $\Gamma_{(4h,1)} = -x_4^{2h}$,
- $\Gamma_{(4h+1,1)} = -x_0$,
- $\Gamma_{(4h+2,1)} = -x_1$,
- $\Gamma_{(4h+3,1)} = x_2$.

Hence $\text{grade}(I_{r_4}(\mathfrak{B}_h^4)) \geq 4$.

Therefore the complex \mathfrak{B}_h is exact, minimality follows from the fact, all entries of matrices \mathfrak{B}_h^i , $1 \leq i \leq 4$ are lies in homogenous maximal ideal $\langle x_0, \dots, x_4 \rangle$. \square

5. SYZYGIES OF PROJECTIVE CLOSURE OF ARSLAN CURVES

In [1] Arslan gave a necessary and sufficient criterion for Cohen-macaulayness of tangent cone of a monomial curve at origin using Gröbner basis. He introduced the following curves, for $h \geq 1$, $\mathfrak{A}_h := C(h(h+1), h(h+1)+1, (h+1)^2, (h+1)^2+1)$ and showed that tangent cone of this curves at origin are Cohen-Macaulay. It has been shown in [4] that Arslan curves are arithmetically Cohen-Macaulay using Gröbner basis. Here we will find syzygies of projective closure of Arslan curves, i.e $\overline{\mathfrak{A}_h}$ and will show that all betti numbers are unbounded function of h .

Lemma 5.1. *Let us consider the polynomials, for $h \geq 2$,*

- $w = x_2x_3 - x_1x_4$
- $g_i = x_1^i x_3^{h-i+1} - x_2^{i+1} x_4^{h-i}$, for $0 \leq i \leq h-1$
- $g_h = x_2^{h+1} - x_1^h x_3$
- $q_j = x_1^{j+1} x_2^{h-j} - x_3^j x_4^{h-j} x_0$, for $0 \leq j \leq h$.

Suppose $\mathfrak{A}_h = \{w, g_i, q_j \mid 1 \leq i, j \leq h\}$ Then \mathfrak{A}_h is a Gröbner basis of defining ideal of projective closure of the Arslan's curve \mathfrak{A}_h w.r.t. degree reverse lexicographic monomial order on the polynomial ring $k[x_0, x_1, x_2, x_3, x_4]$ induced by $x_1 > x_2 > x_3 > x_4 > x_0$.

Proof. We use proposition 5.2 in [4] and theorem 2.6.

Notations 5.1. For $h \geq 2$, let $\mathfrak{I}_h := \mathfrak{p}(h(h+1), h(h+1)+1, (h+1)^2, (h+1)^2+1)$ and $\overline{\mathfrak{I}_h} := \overline{\mathfrak{p}(h(h+1), h(h+1)+1, (h+1)^2, (h+1)^2+1)}$. Also for $h \geq 2$, we define the following matrices.

$$(1) \mathfrak{A}_h^1 := (w \quad g_0 \quad \dots \quad g_h \quad q_0 \quad \dots \quad q_h)$$

$$(2) \mathfrak{A}_h^2 := (\Phi_{ij})_{(2h+3) \times (4h+1)} \text{ where } \Phi_{ij} \text{ are defined as follows,}$$

- $\Phi_{(h+2,1)} = x_1, \Phi_{(h+3,1)} = -x_2, \Phi_{(2h+3,1)} = x_3, \Phi_{(2,1)} = x_0, \Phi_{(i,1)} = 0$, for $i \in \{1, \dots, 4h+1\} \setminus \{2, h+2, h+3, 2h+3\}$.
- $\Phi_{(1,2)} = x_3^i x_4^{h-i-1} x_0, \Phi_{(h+3+i,2)} = -x_1, \Phi_{(h+4+i,2)} = x_2, \Phi_{(i,2)} = 0$, for $i \in \{1, \dots, 4h+1\} \setminus \{2, h+3+i, h+4+i\}, 0 \leq i \leq h-1$.

- $\Phi_{(1,h+2+i)} = x_2^{i+1}x_4^{h-i-1}$, $\Phi_{(2+i,h+2+i)} = -x_1$, $\Phi_{(3+i,h+2+i)} = x_3$, $\Phi_{(j,h+2+i)} = 0$, for $j \in \{1, \dots, 4h+1\} \setminus \{1, 2+i, 3+i\}$, $0 \leq i \leq h-2$
 - $\Phi_{(1,2h+1)} = x_2^h$, $\Phi_{(h+1,2h+1)} = -x_1$, $\Phi_{(h+2,2h+1)} = -x_3$, $\Phi_{(j,2h+1)} = 0$, for $j \in \{1, \dots, 4h+1\} \setminus \{1, h+1, h+2\}$
 - $\Phi_{(1,2h+2+i)} = x_1^i x_3^{h-i}$, $\Phi_{(2+i,2h+2+i)} = -x_2$, $\Phi_{(3+i,2h+2+i)} = x_4$, $\Phi_{(j,2h+2+i)} = 0$, for $j \in \{1, \dots, 4h+1\} \setminus \{1, 2+i, 3+i\}$, $0 \leq i \leq h-2$
 - $\Phi_{(1,3h+1)} = x_1^{h-1}x_3$, $\Phi_{(h+1,3h+1)} = -x_2$, $\Phi_{(h+2,3h+1)} = -x_4$, $\Phi_{(j,2h+2+i)} = 0$, for $j \in \{1, \dots, 4h+1\} \setminus \{1, h+1, h+2\}$
 - $\Phi_{(1,3h+2+i)} = x_1^{i+1}x_2^{h-i-1}$, $\Phi_{(h+3+i,3h+2+i)} = -x_3$, $\Phi_{(h+4+i,3h+2+i)} = x_4$, $\Phi_{(j,3h+2+i)} = 0$, for $j \in \{1, \dots, 4h+1\} \setminus \{1, h+3+i, h+4+i\}$, $0 \leq i \leq h-1$
- (3) $\mathfrak{A}_h^3 := (\Psi_{ij})_{(4h+1) \times (2h-1)}$ where Ψ_{ij} are defined as follows,
- $\Psi_{(2h+2+i,1+i)} = x_1$, $\Psi_{(h+2+i,1+i)} = -x_2$, $\Psi_{(2h+3+i,1+i)} = -x_3$, $\Psi_{(h+3+i,1+i)} = x_4$, $\Psi_{(j,1+i)} = 0$, for $j \in \{1, \dots, 2h-1\} \setminus \{2h+2+i, h+2+i, 2h+3+i, h+3+i\}$, $0 \leq i \leq h-2$
 - $\Psi_{(3h+2+i,h+i)} = x_1$, $\Psi_{(3h+3+i,h+i)} = -x_2$, $\Psi_{(2+i,h+i)} = -x_3$, $\Psi_{(3+i,h+i)} = x_4$, $\Psi_{(j,h+i)} = 0$, for $j \in \{1, \dots, 2h-1\} \setminus \{3h+2+i, 3h+3+i, 2+i, 3+i\}$, $0 \leq i \leq h-2$
 - $\Psi_{(3h+1,2h-1)} = x_1^2$, $\Psi_{(2h+1,2h-1)} = -x_1x_2$, $\Psi_{(3h+2,2h-1)} = x_2^2$, $\Psi_{(4h+1,2h-1)} = -x_1x_3$, $\Psi_{(1,2h-1)} = -x_2x_3$, $\Psi_{(h+1,2h-1)} = x_3^2$, $\Psi_{(1,2h-1)} = x_1x_4$, $\Psi_{(2,2h-1)} = -x_2x_4$, $\Psi_{(2h+2,2h-1)} = -x_3x_0$, $\Psi_{(h+2,2h-1)} = x_0x_4$

Theorem 5.2. For $h \geq 2$, a graded free minimal resolution of $\overline{\mathfrak{J}_h}$ (the defining ideal of the projective closure of Arslan's curve \mathfrak{A}_h) is

$$\mathfrak{M}_h : 0 \longrightarrow R^{2h-1} \xrightarrow{\mathfrak{A}_h^3} R^{4h+1} \xrightarrow{\mathfrak{A}_h^2} R^{2h+3} \xrightarrow{\mathfrak{A}_h^1} R \longrightarrow R/\overline{\mathfrak{J}_h} \longrightarrow 0.$$

Where the matrices \mathfrak{A}_h^i , $1 \leq i \leq 3$ has defined in 5.1.

Proof. We proceed exactly same way as in the proof of theorem 4.4 that is we use Buchsbaum-Eisenbud acyclicity criterion. Let r_i be the i th expected rank of \mathfrak{A}_h . Then $r_1 = (2h+3) - (4h+1) + (2h-1) - 1 = 1$, $r_2 = 4h+1 - (2h-1) = 2h+2$, $r_3 = 2h-1$. We need to show that $\text{grade}(I_{r_i}(\mathfrak{A}_h^i)) \geq i$, $1 \leq i \leq 3$.

(1) We take following minors from \mathfrak{A}_h^2

- $\mathfrak{D}_h^{[21]} := [13 \cdots (2h+3) | 1 \cdots (2h+2)] = (x_2^{h+1} - x_1^h x_3)(-x_3^{h+1} + x_2 x_4^h) x_3^{h-1}$
- $\mathfrak{D}_h^{[22]} := [2 \cdots (2h+3) | 12(2h+2) \cdots (4h+1)] = (x_2 x_3 - x_1 x_4)(-x_1 x_2^h + x_4^h x_0) x_4^{h-1}$.

It is clear from factorization that $\mathfrak{D}_h^{[21]}, \mathfrak{D}_h^{[22]}$ forms a regular sequence. Hence $\text{grade}(I_{r_2}(\mathfrak{A}_h^2)) \geq 2$.

(2) We take following minors from \mathfrak{A}_h^3

- $\mathfrak{D}_h^{[31]} := [2 \cdots h(2h+2) \cdots (3h+1)|1 \cdots (2h-1)] = x_1^{h+1}x_3^{h-1} - x_3^{2h-1}x_0 = x_3^{h-1}(x_1^{h+1} - x_3^h x_0)$
- $\mathfrak{D}_h^{[32]} := [2 \cdots 2h|1 \cdots (2h-1)] = x_2^{h-1}x_3^{h+1} - x_2^{2h}x_4^{2h} = x_2^{h-1}(x_3^{h+1} - x_2^{h+1}x_4^{2h})$
- $\mathfrak{D}_h^{[33]} := [(h+2) \cdots (2h+1)(3h+2) \cdots 4h|1 \cdots (2h-1)] = x_1^h x_2^h - x_1^{h-1}x_4^h x_0 = x_1^{h-1}(x_1 x_2^h - x_4^h x_0)$

From factorization of polynomials, primary decomposition of the ideal $\langle \mathfrak{D}_h^{[31]}, \mathfrak{D}_h^{[32]} \rangle$ is

$$\langle x_3^{h-1}, x_2^{h-1} \rangle \cap \langle x_3^{h-1}, x_3^{h+1} - x_2^{h+1}x_4^{2h} \rangle \cap \langle x_1^{h+1} - x_3^h x_0, x_2^{h-1} \rangle \cap \langle x_1^{h+1} - x_3^h x_0, x_3^{h+1} - x_2^{h+1}x_4^{2h} \rangle.$$

Therefore associated primes of the ideal $\langle \mathfrak{D}_h^{[31]}, \mathfrak{D}_h^{[32]} \rangle$ are,

- $\langle x_3, x_2 \rangle$,
- $\langle x_3, x_3^{h+1} - x_2^{h+1}x_4^{2h} \rangle$,
- $\langle x_1^{h+1} - x_3^h x_0, x_2 \rangle$,
- $\langle x_1^{h+1} - x_3^h x_0, x_3^{h+1} - x_2^{h+1}x_4^{2h} \rangle$.

One can easily check that the polynomial $\mathfrak{D}_h^{[33]}$ does not lie in any associated primes of the ideal $\langle \mathfrak{D}_h^{[31]}, \mathfrak{D}_h^{[32]} \rangle$. Therefore $\{\mathfrak{D}_h^{[31]}, \mathfrak{D}_h^{[32]}, \mathfrak{D}_h^{[33]}\}$ forms a regular sequence. Hence $\text{grade}(I_{r_3}(\mathfrak{A}_h^3)) \geq 3$. Minimality follows from the fact, all entries of matrices $\mathfrak{A}_h^i, 1 \leq i \leq 3$ are lies in homogenous maximal ideal $\langle x_0, \dots, x_4 \rangle$.

□

REFERENCES

- [1] F.Arslan, *Cohen-Macaulayness of Tangent Cones*, Proceedings of the American Mathematical Society, Aug., 2000, Vol. 128, No. 8
- [2] R. Fröberg, C. Gottlieb, R. Häggkvist, *On Numerical Semigroups*, Semigroup Forum Vol. 35 (1987) 63-83
- [3] W. Gastinger, *Über die Verschwindungsideale monomialer Kurven*, PhD thesis, Univ. Regensburg, Landshut (1989).
- [4] J. Herzog, D. Stamate, *Cohen-Macaulay Criteria for projective monomial curves via Gröbner bases*, Acta Mathematica Vietnamica(2019)44: 51-64. <https://doi.org/10.1007/s40306-018-00302-5>
- [5] H. Bresinsky, *On Prime Ideals with Generic Zero $x_i = t^{n_i}$* , Proceedings of the American Mathematical Society, Vol.47, No.2, february 1975.
- [6] H. Bresinsky and L. T. Hoa, *Minimal generating sets for a family of monomial curves in \mathbb{A}^4* , Commutative algebra and algebraic geometry (Ferrara), Lecture Notes in Pure and Appl. Math. 206 (Dekker, New York, 1999) 5–14.

- [7] Decker, W.; Greuel, G.-M.; Pfister, G.; Schönemann, H.: SINGULAR 4.0.1 — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2014).
- [8] Decker, W., Lossen, C., *Computing in Algebraic Geometry A Quick Start using SINGULAR*, Hindustan Book Agency, October 2005.
- [9] J. Herzog, *Generators and relations of abelian semigroups and semigroup rings*, Manuscripta Mathematica, Vol. 2, No. 3, (1970), 175-193.
- [10] R.Mehta, J.Saha, I.Sengupta, *Betti numbers of Bresinsky's curves in \mathbb{A}^4* , Journal of Algebra and Its Applications, DOI: 10.1142/S0219498819501433
- [11] Bayer, D., Stillman, M., 1992. Computation of Hilbert functions. J. Symbolic Comput. 14, 31–50.

Stat Math Unit, Indian Statistical Institute, Kolkata, West-Bengal,700108, INDIA.

Email address: saha.joydip56@gmail.com

Discipline of Mathematics, IIT Gandhinagar, Palaj, Gandhinagar, Gujarat 382355, INDIA.

Email address: indranathsg@iitgn.ac.in

Discipline of Mathematics, IIT Gandhinagar, Palaj, Gandhinagar, Gujarat 382355, INDIA.

Email address: pranjal.srivastava@iitgn.ac.in