

CIRCUIT COMPLEXITY FROM SUPERSYMMETRIC QUANTUM FIELD THEORY WITH MORSE FUNCTION

Sayantana Choudhury^{1,2‡, §}, Sachin Panneer Selvam³, K. Shirish⁴,

¹*National Institute of Science Education and Research, Bhubaneswar, Odisha - 752050, India*

²*Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai - 400085, India*

³*Department of Physics, Birla Institute of Technology and Science, Pilani, Hyderabad Campus, Hyderabad - 500078, India*

⁴*Visvesvaraya National Institute of Technology, Nagpur, Maharashtra, 440010, India*

Abstract

Computation of circuit complexity has gained much attention in the Theoretical Physics community in recent times to gain insights about the chaotic features and random fluctuations of fields in the quantum regime. Recent studies of circuit complexity take inspiration from the geometric approach of Nielsen, which itself is based on the idea of optimal quantum control in which a cost function is introduced for the various possible path to determine the optimum circuit. In this paper, we study the relationship between the circuit complexity and Morse theory within the framework of algebraic topology using which we study circuit complexity in supersymmetric quantum field theory describing both simple and inverted harmonic oscillators up to higher orders of quantum corrections. The expression of circuit complexity in quantum regime would then be given by the Hessian of the Morse function in supersymmetric quantum field theory, and try to draw conclusion from their graphical behavior. We also provide a technical proof of the well known universal connecting relation between quantum chaos and circuit complexity of the supersymmetric quantum field theories, using the general description of Morse theory.

Keywords: Circuit Complexity, Supersymmetric QFT, Morse Theory, AdS/CFT.

[‡] *Corresponding author, E-mail : sayantan.choudhury@niser.ac.in, sayanphysicsisi@gmail.com*

[§] *NOTE: This project is the part of the non-profit virtual international research consortium "Quantum Aspects of Space-Time & Matter" (QASTM) .*

Contents

1	Introduction	1
2	Circuit Complexity for dummies	4
3	Lie algebra formulation of the Inverted Harmonic Oscillator (IHO)	5
4	Brief review of Morse Theory	7
4.1	Definition of Morse function	7
4.2	Gradient flow of Morse function	7
5	Circuit complexity in SUSY QFT via Morse function	8
6	Effect on circuit complexity from IHO perturbation theory	11
7	Effect on circuit complexity from SHO perturbation theory	13
7.1	Circuit complexity for ϕ^3 term	15
7.2	Circuit complexity for ϕ^4 term	15
8	Comparative Analysis	16
8.1	SHO	16
8.2	IHO	18
9	Graphical Analysis	19
9.1	SHO	19
9.2	IHO	22
10	Quantum Chaos from Morse function	26
11	Conclusions	28

1 Introduction

AdS/CFT correspondence has helped in providing great insights on the geometry of the bulk from information in the boundary CFT [1–6]. However, probing the physics of behind the horizon is still a major challenge. It has been inspected that even if the entangled entropy of an eternal AdS black hole saturates after reaching the equilibrium, the size of the Einstein-Rosen(ER) bridge continues to grow with time, posing a problem to the dual description of boundary CFT. Considering this Susskind has introduced new observables in bulk geometry [7–13]. One is the volume of a maximal co-dimension-one bulk surface extending to the boundary of AdS space-time and the second is the action defined on the Wheeler-De-Witt patch. These new observables, according to the conjecture is a dual description of complexity of the boundary field theory.

One of these crucial observables is *volume*, which according to “*Complexity = Volume*” conjecture, states that volume $\mathcal{V}(\mathcal{B})$ of a maximal co-dimension-one bulk surface \mathcal{B} that extends to the AdS boundary and asymptotic to the time slice Σ is proportional to the complexity of the boundary state, $C_{\mathcal{V}}(\Sigma)$, which is given by:

$$C_{\mathcal{V}}(\Sigma) = \max \left[\frac{\mathcal{V}(\mathcal{B})}{G_N l} \right]. \quad (1.1)$$

The second observable is the *gravitational action* estimated in the Wheeler-De-Witt (WDW) patch in the bulk region \mathcal{I}_{WDW} , which according to the another important conjecture, namely the “*Complexity = Action*” conjecture; is proportional to complexity of the boundary field theory [14], given by:

$$C_A = \frac{\mathcal{I}_{WDW}}{\pi \hbar}. \quad (1.2)$$

Complexity in quantum field theory and various other quantum system has been the attraction of theoretical physics community in recent times, because it not only provides insights about what happens inside the horizon in a gravity dual, but also helps in the study the sensitive dependence of initial conditions of boundary fields. In other words, it helps in understanding the out-of-time-order-correlation (OTOC) function of observables in a quantum system which depicts the chaotic and random of quantum fields[15]. OTOC within the framework of supersymmetric quantum mechanics has been studied in the simple harmonic oscillator, one dimensional infinite potential well and various other models in [16] and also the computation of complexity for non-supersymmetric quantum field theories has been done in earlier works [16]. The main idea was to put the free scalar field theory on a lattice, which reduces it to the family of coupled oscillators, then identifying

the circuit as a path ordered exponential of Hamiltonian which forms the representation of $GL(2, R)$, and then construct the euclidean metric. Minimizing the length of this metric would finally give the expression for circuit complexity. These results were then obtained in the framework of the inverted harmonic oscillator and interacting field theories to look for chaotic behavior in quantum field theories(QFT). In this paper, our main objective is to calculate circuit complexity for supersymmetric quantum field theories for simple and inverted harmonic oscillators for various higher-order quantum corrections. To do this we will take an unusual approach and first try to bring out the connection between complexity in quantum field theories namely, supersymmetric field theories with the Morse function. In general Morse theory acts as an essential tool to study the topology of manifolds by studying the differentiable functions through which we could identify the critical points on that manifold namely the minima, maxima, and saddle points. To study the connection between circuit complexity in supersymmetric quantum field theories with the Morse function, we will first identify the so-called “cost function”, an important parameter in computing the circuit complexity which counts the number of gates operating at a particular time t , to construct the optimal circuit with the Morse function on a given manifold. By computing the *Hessian* of the Morse function we obtain the expression for circuit complexity in the present context of discussion. Our motivation for doing so lies around the fact that the eigenvalues of supersymmetric Hamiltonian are closely concentrated near the critical points of the Morse function and the number of zero eigenvalues of the ground state is exactly equal to the Betti number of the manifold. In this paper, we will show how the increase in the number of critical points of the manifold captures the amount of chaos present in supersymmetric quantum field theories for the inverted harmonic oscillator (IHO) and how the mathematical structure of supersymmetry (SUSY) in the regime inverted harmonic oscillators whose potentials forms the generators of $SL(2, R)$ group underlies chaos, and makes the number of critical points increase by a factor of exponential with respect to the superpotential appearing in supersymmetric quantum field theory. For the scalar field theories in the regime where it can be described as a simple harmonic oscillator (SHO), we will identify the non-dynamical auxiliary field namely the F - term in the lagrangian of SUSY theory involving scalar fields, with the gradient of the Morse function, which passes through every critical point on the manifold. In turn we will show that the complexity of supersymmetric quantum field theories for simple harmonic oscillator only depends on the absolute value of the non-dynamical auxiliary field, which also acts as an order parameter for supersymmetry breaking. This way identifying the F - term with the gradient of the Morse function provides another way to check when the supersymmetry is broken. If the gradient flow passes through the critical points, then it could be said that there exists no ground states with zero energy or SUSY is spontaneously broken. Following these lines we will then compute the Hessian in the space of super-coordinates and will derive the expression for complexity. In this connection, it is important to know that Edward Witten has shown that supersymmetric quantum field theory is basically a

Hodge Derham cohomology and has derived the Morse inequality using the formalism of supersymmetry, however, in this paper, we will not work along the same lines but our main objective would be to give quantum chaos a topological flavor in supersymmetric theories. At present, Out-of-time ordered correlation (OTOC) function has been a very important theoretical tool to capture the effect of chaos in any quantum system at late time scales. By following this very important notion, in this paper, we will explicitly give a technical proof of the *universal relation* relating circuit complexity with the quantum chaos in terms of the previously mentioned Out-of-Time Ordered Correlation Function (OTOC) within the framework of supersymmetric quantum field by using the general description of Morse theory.

Organization of the paper is as follows:

- In [section 2](#), we provide a brief review of the concept of circuit complexity in the general context of quantum information theory.
- In [section 3](#), we will explain the Lie algebra formulation and how the potential of Inverted Harmonic Oscillator (IHO) could be embedded in the structure of the manifold.
- In [section 4](#), we will give a brief review of Morse theory namely the Morse function and the gradient flow of the Morse function for completeness.
- In [section 5](#), we will comment on the relationship between circuit complexity and Morse function over a manifold in supersymmetric quantum mechanics.
- In [section 6](#), we will explicitly compute the expression of circuit complexity for the Inverted Harmonic Oscillator (IHO) upto higher order of quantum corrections.
- In [section 7](#), we will compute the complexity for supersymmetric field theories for simple harmonic oscillators up to higher orders of quantum corrections in terms of non-dynamical auxiliary field.
- In [section 8](#), we will compare the results of complexity between supersymmetric and non-supersymmetric quantum field theory for SHO and IHO in [table 8.1](#) and [8.3](#).
- In [section 9](#), we will provide the detailed numerical and graphical analysis of the obtained theoretical results for both SHO and IHO using the prescription of supersymmetric Morse quantum field theory.
- In [section 10](#), we will derive the universal relation between the circuit complexity and quantum chaos expressed in terms of OTOC function for supersymmetric quantum field theory prescription using Morse function.

- Finally [section 11](#), we will conclude our result and comment on how the behavior of complexity for the supersymmetric model is already implemented in the structure of a manifold.

2 Circuit Complexity for dummies

The notion of circuit complexity was first introduced in information theory to find the minimum number of gates to get the desired state from an initial state. It involves acting the initial state with a unitary operator or a set of quantum gates to obtain the desired target state.

$$|\psi_T\rangle = U |\psi_i\rangle \quad (2.1)$$

There exists many such set of unitary operators to get the desired final state but to find a minimum number of such operations to execute the task is what constitutes an optimal quantum circuit. Taking inspiration from this Nielsen and collaborators have developed this idea further and have taken a geometric approach to finding the most optimal quantum circuit to get the desired target state via unitary operations in physics [17–22],

$$U(t) = \mathcal{P} \exp \left(-i \int_0^t H(t) dt \right) \text{ where } H(t) = \sum Y^I(t) M_I, \quad (2.2)$$

where M_I are the Pauli matrices and $Y^I(t)$ are referred to as the control function that decides the nature of gate that will act at a certain value of parameter t . This approach of identifying the action of quantum gates via control function could be morphed in terms of finding the extremal curves, i.e. geodesics. Hence we define a cost function $F(U(t), \dot{U}(t))$ which is a function of the unitary operator U and a vector at a point in the tangent space formed by unitaries. The idea is to minimize this cost function for various possible paths, which is described by the following expression:

$$\mathcal{D}(U(t)) = \int_0^1 \left(F(U(t), \dot{U}(t)) \right) dt \quad (2.3)$$

The task now is to determine the cost function which as described above counts the number of gates to construct the optimal quantum circuit. A general class of cost functions are:

$$F_p(U, Y) = \sum p_I |Y^I|, \quad (2.4)$$

$$F_q(U, Y) = \sqrt{\sum q_I(Y^I)^2}, \quad (2.5)$$

where minimizing F_2 would give us the expression for the length of geodesic of a Riemann surface. This technique uses the concepts of differential geometry to construct the optimal quantum circuit and comment on the complexity of various models in theoretical physics.

3 Lie algebra formulation of the Inverted Harmonic Oscillator (IHO)

To work out the circuit complexity in various models researchers have considered a simple, exactly solvable system known as the Inverted Harmonic Oscillator (IHO), which is described by the Hamiltonian function:

$$H(p, x) = \frac{p^2}{2} - \frac{1}{2}\omega^2 x^2. \quad (3.1)$$

This encapsulates the sensitivity to initial conditions exhibited by the chaotic systems [23–25]. The Inverted Harmonic Oscillator (IHO) differs from the Simple Harmonic Oscillator (SHO) in a lot of ways. For instance, the energy spectrum of IHO has a continuous energy spectrum, whereas the regular SHO has a discrete energy spectrum $(n + \frac{1}{2})\hbar\omega$. Unlike SHO, the IHO is exactly solvable and while the SHO provides a good description of the deviations from the stable equilibrium, the IHO models the decay from an unstable equilibrium. The IHO in recent times has been proved to be incredibly useful to show the equivalence among various diverse fields. The IHO appears in the quantum hall effect and in the mechanism of Rindler Hamiltonian, whose time evolution would give rise to the Hawking-Unruh effect [26–28]. The equivalence between the two phenomena can be shown in terms of the isomorphism of the underlying lie algebra [29, 30].

In this paper, our main objective will not be to formally describe the lie algebra isomorphism but instead using that to calculate the circuit complexity over a manifold via Morse function for supersymmetric quantum field theories. To do this, first we have to show how the potential of IHO forms the generators of Lie algebra. We start with the settings of the quantum hall effect (QHE), the three quadratic potentials generate the Hamiltonian dynamics in the lowest Landau level (LLL):

$$P_{LLL}V_1P_{LLL} = \lambda(R_x^2 + R_y^2), \quad (3.2)$$

$$P_{LLL}V_2P_{LLL} = \lambda(R_X R_Y + R_Y R_X), \quad (3.3)$$

$$P_{LLL}V_3P_{LLL} = \lambda(R_x^2 - R_y^2). \quad (3.4)$$

Furthermore, one can write:

$$P_{LLL}V_1P_{LLL} = \frac{-4\lambda_B^2}{\hbar P_{LLL}LP_{LLL}}, \quad (3.5)$$

$$P_{LLL}V_2P_{LLL} = 2\lambda_B^2 P_{LLL}j_i P_{LLL}, \quad (3.6)$$

$$P_{LLL}V_3P_{LLL} = \frac{-4\lambda_B^2}{\hbar P_{LLL}j_j P_{LLL}}. \quad (3.7)$$

where P_{LLL} is the projection operator to the lowest Landau level, R_X, R_Y are bilinears and j_a, j_b are strain generators. Therefore by projecting the potentials to the LLL both bilinears and the strain generators lead to the quadratic Hamiltonian which is similar to the electrostatic potential and IHO appears in quantum hall effect as might expected.

We now rename the three quadratic potentials in a quantum hall effect V_1, V_2, V_3 as K_1, K_2, K_3 and we identify:

$$P = \frac{R_x}{l_B}, \quad X = \frac{R_y}{l_B}. \quad (3.8)$$

Consequently, we get the Hamiltonians in the LLL to be of the following form:

$$K_1 = (P^2 + X^2), \quad (3.9)$$

$$K_2 = (PX + XP), \quad (3.10)$$

$$K_3 = (P^2 - X^2). \quad (3.11)$$

On the basis of LLL wave functions the potentials could be written the form of differential operators namely:

$$K_1 = \frac{1}{4} \left(-\frac{\partial^2}{\partial X^2} + X^2 \right), \quad (3.12)$$

$$K_2 = \frac{i}{2} \left(X \frac{\partial}{\partial x} + \frac{1}{2} \right), \quad (3.13)$$

$$K_3 = \frac{1}{4} \left(-\frac{\partial^2}{\partial X^2} - X^2 \right). \quad (3.14)$$

These are precisely the generators of $SL(2, R)$ Lie-algebra[31], which act as an area preserving deformations of a two dimensional manifold and satisfy the following non-commuting relations:

$$[K_2, K_3] = -iK_1, \quad (3.15)$$

$$[K_1, K_2] = iK_3, \quad (3.16)$$

$$[K_3, K_1] = iK_2. \quad (3.17)$$

With these techniques at hand, our main objective will now be to describe a Morse function on a manifold formed by the Lie-algebra of $SL(2, R)$ namely the potential of the IHO, and show, how the critical points of Morse function on a manifold actually play the role of the cost function, which decides the gate in action at a particular parameter t to form the optimal circuit.

4 Brief review of Morse Theory

In this section, we are going to give a very brief review of Morse theory. The Morse function helps in the classification of surfaces upto homeomorphism when it passes critical point of index 0, 1, 2. The index 2, 1, and 0 represents the maximum, minimum and saddle points of the manifold.

4.1 Definition of Morse function

To define Morse function, we consider a manifold \mathcal{M} and a function f such that:

Theorem 1:

A smooth map $f : X^n \rightarrow R$ is a Morse function if, for every critical point $p \in X$, \exists coordinates x_1, \dots, x_n , and a coordinate y around $f(p)$ w.r.t. which,

$$f(x_1, \dots, x_n) = -x_1^2 - x_2^2 - \dots - x_k^2 + x_{k+1}^2 + \dots + x_n^2. \quad (4.1)$$

such that the value of the function at the critical points vanishes.

Theorem 2:

A map $f \rightarrow R$ is a morse function if its critical points vanishes and the Hessian of f at each critical point is non-singular.

A Morse function on a compact manifold X helps to determine it's topology, by mapping it's critical points to an axis on a one-dimensional plane, which helps to encode a lot of information about \mathcal{M} [32]. The goal of the Morse theory is to find the invariant of the manifold by counting the critical points of chosen Morse function.

4.2 Gradient flow of Morse function

The gradient flow of the Morse function creates a vector field on the surface of the manifold, which helps to define a notion of transport from one point to another. Let us

suppose we have an integral curve $\gamma_x: R \rightarrow M$ such that:

$$\gamma_x(t) = \phi_t(x), \quad \phi_0(x) = 0. \quad (4.2)$$

Here ϕ is a smooth one-parameter group of diffeomorphism on \mathcal{M} . Then, we get:

$$\begin{aligned} \frac{d}{dt}f(\gamma_x(t)) &= \frac{d}{dt}(f \circ \phi_t(x)) \\ &= df_{\phi_t(x)} \circ \frac{d}{dt}\phi_t(x) \\ &= df_{\phi_t(x)}(-\nabla f)\phi_t(x) \\ &= -\|(\nabla f)\phi_t(x)\|^2 \leq 0. \end{aligned} \quad (4.3)$$

Thus the gradient flow of f is found to be decreasing down the lines of γ_x . One of the reasons why we have defined the gradient flow of the Morse function will become clear in later part of this paper, where it is going to play a very crucial part in deriving the complexity of supersymmetric field theories for the SHO.

5 Circuit complexity in SUSY QFT via Morse function

To compute circuit complexity within the framework of supersymmetric quantum field theory using Morse function, we will consider $SL(2, R)$ modular curve which encodes the potential of IHO as generators of Lie algebra and identify it with a Riemann manifold \mathcal{M} , and the Morse function to the cost function as described in the previous section, and thereby associate the critical points to the action of the control function which decides, which quantum gate will be active at a particular time t .

The supersymmetric operators in terms of an exterior derivative and its adjoint can be described as:

$$Q_1 = d + d^*, \quad (5.1)$$

$$Q_2 = i(d - d^*), \quad (5.2)$$

$$H = dd^* + d^*d. \quad (5.3)$$

The connection between supersymmetric quantum field theories and Derham operators could found in the ref. [33]. Then further by taking into account the following crucial fact:

$$d^2 = 0 = d^{*2}, \quad (5.4)$$

we get the following subsequent supersymmetric relations in the present context, which are given by:

$$Q_1^2 = Q_2^2 = H, \quad (5.5)$$

$$Q_1 Q_2 + Q_2 Q_1 = 0. \quad (5.6)$$

Now let us consider a Morse function f on the surface of the manifold M , and t be a real number. Then,

$$d_t = \exp(-ht) d \exp(ht), \quad (5.7)$$

$$d_t^* = \exp(-ht) d^* \exp(ht). \quad (5.8)$$

Here we can show that:

$$d_t^2 = 0 = d_t^{*2}, \quad (5.9)$$

using which we get the following expressions:

$$Q_{1t} = d + d^*, \quad (5.10)$$

$$Q_{2t} = i(d_t - d_t^*), \quad (5.11)$$

$$H_t = d_t d_t^* + d_t^* d_t. \quad (5.12)$$

We will now explicitly calculate the formula for H_t in terms f , to understand how critical points come into the picture. Let, $v^k(p)$ and v^{k*} be an orthonormal basis of tangent vectors and the corresponding dual vectors at each point p in \mathcal{M} . The a^k and a^{k*} could be regarded as creation and annihilation operators in the present context. We could calculate the covariant second derivative of Morse function f in the dual basis of v^k as $\left(\frac{D^2}{Dx^i Dx^j}\right) f$, with these accords, one could then calculate the Laplacian operator H_t acting on p forms on manifold \mathcal{M} :

$$H_t = dd^* + d^*d + t^2(df)^2 + \sum t \left(\frac{D^2}{Dx^i Dx^j}\right) [v^{*i}, v^j]. \quad (5.13)$$

Here we define $(df)^2$ by the following expression:

$$(df)^2 = g^{ij} \left(\frac{df}{dx^i}\right) \left(\frac{df}{dx^j}\right). \quad (5.14)$$

which is the square of the gradient of the Morse function f , measured with respect to the Riemannian metric g^{ij} of M . Here the term $t^2(df)^2$ plays the role of IHO potential, and also the critical points for the Morse function f lies, where we have

$$df = \sqrt{g^{ij} \left(\frac{df}{dx^i}\right) \left(\frac{df}{dx^j}\right)} = 0. \quad (5.15)$$

Let us now calculate the circuit complexity in supersymmetric quantum field theory by explicitly identifying the cost function with Morse function,

$$F(Q(t), \dot{Q}(t)) \longrightarrow f. \quad (5.16)$$

The role of $F(Q(t), \dot{Q}(t))$ is to count the number of gates required to construct the optimal quantum circuit, but in the present contest it is to find the minimum of supersymmetric charges to get:

$$Q_1 |E^n\rangle_{bosonic} = E_{bosonic}^{n-i}. \quad (5.17)$$

One could see in the above mentioned argument, that eigenfunction of H_t for large t , are concentrated near the critical points of f therefore for the identification, we could say that the eigenvalues of the cost function are actually the critical values of $SL(2, R)$ Riemann surface.

The eigenvalues of H_t acting on p forms for large t can be expressed in the following way:

$$\lambda_p^n(t) = t \left(A_p^n + \frac{B_t^n}{t} + \frac{C_p^n}{t^2} + \dots \right). \quad (5.18)$$

Thus we could see that for large t the above equation, with in few orders agrees with the computations of out-of-time-order-correlations (OTOC) from supersymmetric quantum mechanics computed in [16].

Now to calculate the general expression for circuit complexity, we will make use of the Hessian matrix. Remember a function f is a morse function if and only if the Hessian of f at each critical point is non-singular. This can be described as:

$$H(f(x)) = j(\nabla f(x))(Hf)_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}. \quad (5.19)$$

we will now identify Hessian of f in the tangent space of unitary operator as:

$$T_p(M) \times T_p(M) \longrightarrow R. \quad (5.20)$$

such that we have:

$$\frac{dU(s)}{ds} = -iY_s^I = (Hf)_{ij}, \quad (5.21)$$

where Y_s^I is the control function, which counts the number of operators acting on the reference state to make the optimal circuit.

Consequently, we compute:

$$\mathcal{D}(Q)(s) = \sqrt{(Y^{11})^2 + (Y^{12})^2 + (Y^{21})^2 + (Y^{22})^2}, \quad (5.22)$$

and the corresponding circuit complexity can be computed as:

$$\boxed{\mathcal{C}(Q) = |H(f(x))| = |j(\nabla f(x))(Hf)_{ij}| = \left| \frac{\partial^2 f}{\partial x_i \partial x_j} \right|}. \quad (5.23)$$

6 Effect on circuit complexity from IHO perturbation theory

To explicitly calculate circuit complexity for IHO in a supersymmetric case, we will make use of the concept of Witten index and show how critical point near which the eigenvalues of $H(t)$ are concentrated grows exponentially with respect to the superpotential.

The excited states of supersymmetric quantum field theory always come with pair of states, this could be seen in the algebra of supersymmetry which has no one-dimensional representation [34]:

$$\{Q, Q^*\} = 2H \quad \text{such that} \quad Q^2 = Q^{*2} = 0. \quad (6.1)$$

The Witten index $\text{Tr}[(-1)^F e^{-\beta H}]$, where $(-1)^F$ is the well known fermion number operator, carries interesting information about the ground state of supersymmetric quantum system especially when it is non-zero, i.e. the system has at least one ground state when the Witten index is non-zero, however, it doesn't make any comment on the number of ground states of the system when it is zero [35]. We will take the Hamiltonian of the supersymmetric quantum system in IHO regime by replacing $W(x)$ by $iW(x)$ and show how the critical points of Morse function or complexity grow exponentially with respect to the superpotential, which is given by:

$$H(P, W) = \frac{P^2}{2} - \frac{W^2}{2}. \quad (6.2)$$

have a ground state wave function, which is defined as:

$$\psi = \exp\left(-i\sigma_2 \int_{-\infty}^{\infty} W dx\right). \quad (6.3)$$

Now by doing supersymmetry on the two dimensional manifold the Witten index can be expressed by the following expression [36]:

$$\text{Tr}[(-1)^F \exp(-\beta H)] = \chi(M). \quad (6.4)$$

where $\chi(M)$ is the Euler characteristic of the manifold, thereby using the strong Morse

inequality, we further get:

$$\chi(M) = \text{Tr}[(-1)^F \exp(-\beta H)] = \sum (-1)^F \exp(-\beta H) = \sum (-1)^\gamma C^\gamma . \quad (6.5)$$

where C^γ is the number of critical points of index γ , using which one can find out the following simplified expression:

$$C^\gamma = \left| (-1)^{F-\gamma} \exp \left(-\beta \left(\frac{P^2}{2} - \frac{W^2}{2} \right) \right) \right| . \quad (6.6)$$

Circuit complexity for higher-order interacting term could be calculated in terms of super-field formalism Φ , in which all super-partners related by SUSY transformations could be treated as a single field. In this paper we will restrict ourselves to $\mathcal{N} = 1$ supersymmetry, such that we will have only one fermionic annihilation and creation operator and thereby only one supercharge[37–39]. Scalars and fermions related by supersymmetry correspond to different components of super-field. The most general form of super-fields in terms of super-space variables can be expressed as follows:

$$\Phi(z) = \phi(x) + \theta\psi + \bar{\theta}\bar{\chi}(x) + \theta\sigma^\mu\bar{\theta}A_\mu(x) + \theta^2 F(x) . \quad (6.7)$$

Now, to write the super-field in terms of any single field component, we will apply the SUSY transformations by applying the operator $i(\xi Q + \bar{\xi}\bar{Q})$ under the projection:

$$\theta = \bar{\theta} = 0, \quad (6.8)$$

Expressing the result in terms of other components:

$$\delta_\xi \phi(x) = i(\xi D + \bar{\xi}\bar{D})\Phi|_{\theta=\bar{\theta}=0} = -\xi\psi(x) - \bar{\xi}\bar{\chi}(x), \quad (6.9)$$

where D is a super-covariant derivative of the super-field which anti-commutes with the supersymmetric charges Q and \bar{Q} and under transformations maps super-field to super-field, given by:

$$\delta_\xi D_\alpha \Phi(z) = D_\alpha \delta_\xi \Phi(z) = i(\xi Q + \bar{\xi}\bar{Q})D_\alpha \Phi(z) \quad (6.10)$$

They satisfy the same algebra as supersymmetric charges, thus the scalar component $\phi(x)$, the super-field can be recovered by exponentiating supersymmetry transformations with θ as the parameter:

$$\Phi = \exp(-\delta_\theta)\phi(x), \quad (6.11)$$

Thus, the super-fields can be constructed by applying the operator $\exp(-\delta_\theta)$ to any com-

ponent field. Hence under the previously mentioned projection $\theta = \bar{\theta} = 0$ the complexity for higher order interactions in the super-potential could be written is as follow:

$$\boxed{\text{Complexity for } \phi^3 \text{ term : } C^\gamma = (-1)^{F-\gamma} \exp \left(-\beta \left(\frac{P^2}{2} - \frac{m}{2} \phi^2 - \frac{\lambda}{3} \phi^3 \right) \right)} \quad (6.12)$$

$$\boxed{\text{Complexity for } \phi^2 + \phi^4 \text{ term : } C^\gamma = (-1)^{F-\gamma} \exp \left(-\beta \left(\frac{P^2}{2} - \frac{m}{2} \phi^2 - \frac{\lambda}{4} \phi^4 \right) \right)} \quad (6.13)$$

7 Effect on circuit complexity from SHO perturbation theory

To calculate the circuit complexity for supersymmetric quantum field theory, it is convenient to work in the super-space formalism [40, 41], i.e. we extend the 4 commuting space-time coordinates x_ν to 4 commuting and 4 ant-commuting coordinates $x_\nu, \theta^\alpha, \bar{\theta}^\alpha$. These new coordinates satisfy the following anti-commuting relations:

$$\{\theta_\alpha, \theta_{\dot{\beta}}\} = \{\theta_\alpha, \theta_\beta\} = \{\theta_{\dot{\alpha}}, \theta_{\dot{\beta}}\} = 0. \quad (7.1)$$

Now, any super-multiplet in super-space coordinates could be communicated in terms of super-fields and can be expressed as:

$$\Phi = \phi(x) - i\theta\sigma^\mu\bar{\theta}\partial_\mu\phi(x) - \frac{1}{4}\theta^2\bar{\theta}^2\partial^2\phi(x) + \sqrt{2}\theta\eta + \frac{i}{\sqrt{2}}\theta^2\partial_\mu\eta\sigma^\mu\bar{\theta} + \sqrt{2}\theta^2F(x). \quad (7.2)$$

where η is a Weyl fermion having 4 off-shell degrees of freedom and σ are the Pauli matrices and ϕ is a complex scalar having two degrees of freedom. The supersymmetric Lagrangian remains invariant even after the addition of the term, i.e.

$$\delta\mathcal{L} = \mu^2F + h.c. \quad (7.3)$$

The F term is an auxiliary complex bosonic field which has two off-shell degrees of freedom to match the 4 off-shell degrees of freedom of a weyl fermion, also the F term is an order parameter for SUSY breaking, substituting $F = -\mu^2$ the ground state will not be invariant and supersymmetry will be spontaneously broken.

The Lagrangian in terms of components fields up to second order is:

$$\mathcal{L} = |\partial_\mu \phi|^2 + i\eta^\dagger \partial_\mu \bar{\sigma}^\mu \eta + |F|^2 + \left(mF\phi - \frac{m}{2}\eta\eta + h.c \right). \quad (7.4)$$

We will now identify the auxiliary field F with the gradient of the Morse function which passes through every critical point on the surface, such that these points act as an order parameter for SUSY breaking. By doing the coordinate transformation, $U(u, v) = (x(u, v), \theta(u, v))$ we could define:

$$h(u, v) = g \circ U(u, v) \quad (7.5)$$

such that:

$$\frac{\partial}{\partial x} g = F. \quad (7.6)$$

The eigenvalues of $H(t)$ are concentrated at the critical points of g , hence precisely at the critical point where:

$$dg = F = 0 \quad (7.7)$$

represents the scalar potential of the theory with no supersymmetric ground state. Then Hessian of h can be calculated as:

$$\frac{\partial^2 h}{\partial u^2} = Fx_u^2 + 2Fx_u\theta_u, \quad (7.8)$$

$$\frac{\partial^2 h}{\partial v^2} = Fx_v^2 + 2Fx_v\theta_v, \quad (7.9)$$

$$\frac{\partial^2 h}{\partial u \partial v} = Fx_u x_v + Fx_v x_u + F\theta_u \theta_v + Fx_{uv} + Fx_v \theta_u. \quad (7.10)$$

the above dependence on the derivative of super-space variable with respect to u and v is actually the Jacobian due to the change of the variables as mentioned above. Now solving the equation of motion for F , the circuit complexity can be evaluated as:

$$\boxed{\mathcal{C}(\mathcal{Q}) = \mathcal{F}_1(x_u, x_v, \theta_u, \theta_v, \phi) - \mathcal{I}_1(x_u, x_v, x_{uv}, \theta_u, \phi)}, \quad (7.11)$$

where the newly introduced function $\mathcal{F}_1(x_u, x_v, \theta_u, \theta_v, \phi)$ and $\mathcal{I}_1(x_u, x_v, x_{uv}, \theta_u, \phi)$ are defined by the following expressions:

$$\mathcal{F}_1(x_u, x_v, \theta_u, \theta_v, \phi) := \left(\frac{m\phi}{2}x_u^2 + 2\frac{m\phi}{2}x_u\theta_u \right) \left(\frac{m\phi}{2}x_v^2 + 2\frac{m\phi}{2}x_v\theta_v \right), \quad (7.12)$$

$$\mathcal{I}_1(x_u, x_v, x_{uv}, \theta_u, \phi) := \left(\frac{m\phi}{2} x_u x_v + \frac{m\phi}{2} x_v x_u + \frac{m\phi}{2} x_{uv} + \frac{m\phi}{2} x_v \theta_u \right)^2. \quad (7.13)$$

7.1 Circuit complexity for ϕ^3 term

To calculate the complexity for higher order interacting terms we will follow the same procedure as above, notice that circuit complexity for SUSY field theories only depends on the value of auxiliary field which also act an order parameter for soft SUSY breaking. The Lagrangian for cubic interactions could be defined as:

$$\mathcal{L} = |\partial_\mu \phi|^2 + i\eta^\dagger \partial_\mu \bar{\sigma}^\mu \eta + |F|^2 + \left(mF\phi + \lambda F\phi^2 - \frac{m}{2}\eta\eta - \lambda\phi\eta\eta + h.c \right) \quad (7.14)$$

The kinetic term is a Khähler potential corresponding to the $\theta\bar{\theta}$ term which is invariant under the SUSY transformations. A general Khähler potential could give rise to complicated terms in the Lagrangian, but for simplicity we will consider the most canonical kinetic terms. λ here is the coupling constant, and the corresponding F term is given by the following expression:

$$F(x) = -\frac{\phi(m+2\lambda)}{2}. \quad (7.15)$$

As described above we will now identify gradient of the Morse function with absolute value of the auxiliary field F which is invariant under the SUSY transformations, and then compute the hessian of the Morse function and thereby the complexity as described above:

$$\boxed{\mathcal{C}(\mathcal{Q}) = \mathcal{F}_2(x_u, x_v, \theta_u, \theta_v, \phi) - \mathcal{I}_2(x_u, x_v, x_{uv}, \theta_u, \phi)}, \quad (7.16)$$

where the newly introduced function $\mathcal{F}_2(x_u, x_v, \theta_u, \theta_v, \phi)$ and $\mathcal{I}_2(x_u, x_v, x_{uv}, \theta_u, \phi)$ are defined by the following expressions:

$$\mathcal{F}_2(x_u, x_v, \theta_u, \theta_v, \phi) := \left(\frac{\phi(m+2\lambda)}{2} x_u^2 + 2\frac{\phi(m+2\lambda)}{2} x_u \theta_u \right) \left(\frac{\phi(m+2\lambda)}{2} x_v^2 + 2\frac{\phi(m+2\lambda)}{2} x_v \theta_v \right), \quad (7.17)$$

$$\mathcal{I}_2(x_u, x_v, x_{uv}, \theta_u, \phi) := \left(\frac{\phi(m+2\lambda)}{2} x_u x_v + \frac{\phi(m+2\lambda)}{2} x_v x_u + \frac{\phi(m+2\lambda)}{2} x_{uv} + \frac{\phi(m+2\lambda)}{2} x_v \theta_u \right)^2. \quad (7.18)$$

7.2 Circuit complexity for ϕ^4 term

Super-potential allows to introduce a variety of supersymmetric interactions, here we will study complexity for quartic interaction terms namely $\phi^2 + \phi^4$. The representative Lagrangian involving quartic interaction in terms of components of super-fields is given

by:

$$\mathcal{L} = |\partial_\mu \phi|^2 + i\eta^\dagger \partial_\mu \bar{\sigma}^\mu \eta + |F|^2 + \left(mF\phi + \lambda\phi^3 - \frac{m}{2}\eta\eta - \lambda\phi^2\eta\eta + h.c \right). \quad (7.19)$$

Here F is again representing the auxiliary field as described above, responsible for soft SUSY breaking. By solving the equation of motion for F term we get:

$$F = -\frac{m\phi + \lambda\phi^3}{2}. \quad (7.20)$$

Again identifying the auxiliary field by gradient of the Morse function as described in previous section, we could compute the circuit complexity by computing the Hessian of the Morse function for the corresponding perturbed term which then gives:

$$\boxed{\mathcal{C}(\mathcal{Q}) = \mathcal{F}_3(x_u, x_v, \theta_u, \theta_v, \phi) - \mathcal{I}_3(x_u, x_v, x_{uv}, \theta_u, \phi)}, \quad (7.21)$$

where the newly introduced function $\mathcal{F}_3(x_u, x_v, \theta_u, \theta_v, \phi)$ and $\mathcal{I}_3(x_u, x_v, x_{uv}, \theta_u, \phi)$ are defined by the following expressions:

$$\mathcal{F}_3(x_u, x_v, \theta_u, \theta_v, \phi) := \left(\frac{m\phi + \lambda\phi^3}{2} x_u^2 + 2\frac{m\phi + \lambda\phi^3}{2} x_u \theta_u \right) \left(\frac{m\phi + \lambda\phi^3}{2} x_v^2 + 2\frac{m\phi + \lambda\phi^3}{2} v\theta_v \right), \quad (7.22)$$

$$\mathcal{I}_3(x_u, x_v, x_{uv}, \theta_u, \phi) := \left(\frac{m\phi + \lambda\phi^3}{2} x_u x_v + \frac{m\phi + \lambda\phi^3}{2} x_v x_u + \frac{m\phi + \lambda\phi^3}{2} x_{uv} + \frac{m\phi + \lambda\phi^3}{2} x_v \theta_u \right)^2. \quad (7.23)$$

Hence we see that circuit complexity for supersymmetric field theories only depends on the absolute value of the auxiliary field coming into light from the linear term in the super-potential. The above method for calculating the complexity for SHO is similar to what we have used for IHO. The number of zero eigenvalues of supersymmetric H are precisely equal to the Euler number of the manifold, therefore by identifying the F term with the gradient of the Morse function, the critical points are exactly where the F term becomes zero and determines the ground state of the system.

8 Comparative Analysis

8.1 SHO

Parameter	Non-SUSY QFT		SUSY QFT	
	$\phi^4 + \phi^2$	ϕ^2	$\phi^4 + \phi^2$	ϕ^2
Mass	Complexity for Non-supersymmetric field theories has polynomial as well as logarithmic dependence on the mass parameter as $\log(m\delta)$	Circuit complexity for Non-SUSY QFT has logarithmic dependence on the square of the mass parameter $\log(m^2\delta)$, and in the infrared (IR) region it takes the form $-\log^k(mk)$.	Complexity for Supersymmetric field theories has only polynomial dependence namely the quadratic exponent of mass parameter. For large masses we observe a decrease in rate and reaches a saturation value.	Complexity for Supersymmetric field theories in case of quadratic perturbations also has polynomial namely the quadratic dependence on mass parameter.
Topological dependence	Complexity for Non-supersymmetric field theories has a fractional dependence on the volume of lattice for interacting terms, such as $V^{\frac{1}{2}}$ for dimension $d = 3$	Complexity due to just quadratic perturbations in Non-SUSY QFT doesn't have any topological dependence but depends on the dimension of lattice used for computations	Complexity for supersymmetric field theories due to quadratic interactions depends on the Hessian of the Morse function whose gradient has been identified with the auxiliary field and also on the critical points of the manifold and hence has topological dependence	SUSY complexity due to quadratic term in the superpotential also depends on topological parameters (Hessian of the Morse function) defined on the manifold.
Dependence on field	Complexity for Non-supersymmetric field theories depends on various parameters of quantized field in theories and the strength of interaction with each other and normal frequency modes	Due to quadratic perturbations complexity for Non-SUSY field theories on a lattice depends on components of the momentum vectors and the number of oscillators in the lattice formalism and doesn't depend on the coupling parameter.	Complexity for SUSY QFT only depends on the absolute value of non-dynamical auxiliary field namely the F -term as $F = \frac{m\phi + \lambda\phi^3}{2}$ which is identified as the gradient of the Morse function which passes through every critical points on the surface	In case of quadratic perturbation complexity changes due to the change of F -term which by solving equation of motion is given by $\frac{m\phi}{2}$ and doesn't have any dependence on coupling constant
Growth of complexity	Complexity for perturbing term ϕ^4 for dimension $d > 0$ breaks down and in the limit $\lambda \rightarrow 0$ the circuit complexity have a continuous limit	In the infrared scale i.e $\omega_0 \ll \frac{1}{\delta}$ there is an additional logarithmic factor in the complexity and lead to divergences in the limit $\delta \rightarrow 0$.	When we change coupling λ from 1 to -1 we observe that complexity first rises and then have a sudden dip. As we go to more negative values of lambda we observe the rate of saturation to be faster.	In supersymmetric field theories we have strangely observed that complexity first rapidly grows and then saturates doesn't change much when we change the coupling λ to -1

Table 8.1: Comparison in circuit complexity between SUSY & NON-SUSY QFT for SHO

Parameters of SHO	Features of graph and Lyapunov
General features	The graphs rise fast initially and then slowly saturate. The rate of saturation and complexity at saturation point depends on the order for perturbation with ϕ^4 term being the slowest to saturate. Hence the slope is significant and the Lyapunov exponent is expected to be larger for ϕ^4 theory. The ϕ^3 theory saturates very quickly giving a smaller slope than the rest and a smaller Lyapunov exponent.
Mass	For large masses we observe a decrease in rate and saturation value of ϕ^4 theory. It becomes indifferentiable to the ϕ^2 term as we approach massive fields (hence a smaller Lyapunov exponent). For smaller masses the ϕ^3 graph approaches ϕ^2 , decreasing slightly in rate. We can expect a slight increment in the Lyapunov exponent.
λ	When we make λ negative we observe the saturation is slower in ϕ^2 and ϕ^3 perturbations, and in the case of ϕ^4 perturbation we encounter a zero thereby right-shifting the point of initial rise which will increase the value of the Lyapunov exponent. As we go to more negative values of lambda we observe the rate of saturation to be faster, and we expect the Lyapunov exponent to be smaller.

Table 8.2: Discussions on Lyapunov exponent for SHO complexity

8.2 IHO

Parameter	Non-supersymmetric QFT (ϕ^2)	Supersymmetric QFT (ϕ^2)
Mass	Circuit complexity for Non-SUSY QFT in the regime of inverted harmonic oscillators has quadratic dependence on a mass parameter in the exponential type function namely the inverse cosine hyperbolic function. Also it has been observed that complexity starts to increase before the critical value $\lambda = m^2$.	Circuit complexity for supersymmetric field theories for the inverted harmonic oscillator also has quadratic dependence on the mass parameter in the exponential function. We also observe that as mass increases the rate of change of complexity increases.
Topological dependence	Circuit complexity of Non-supersymmetric quantum field theories doesn't have any topological dependence, however it depends on the number of oscillators and dimension of the lattice	Circuit complexity for supersymmetric field theories depends on the critical points of the manifold, also for even values of $F - \gamma$ we see that complexity increases exponentially.
λ	The complexity starts to increase for $\lambda < \lambda_c$. At the critical point the complexity sharply increases and beyond the critical value $\lambda = m^2$ the model becomes unstable and we expect the complexity to grow rapidly with decreasing pick up time	The change in value of λ contributes to the rate with increasing values resulting in faster rates of increase and hence higher slopes. For negative values of λ we observe that complexity decreases exponentially and model becomes irrelevant.
Growth in complexity	For inverted harmonic oscillator, complexity for non-supersymmetric field theories for initial time is nearly zero, after which it exhibit linear growth.	On the contrary for the inverted harmonic oscillator complexity doesn't exhibit any exponential or linear growth which could be seen in figure (9.5).

Table 8.3: Comparison in complexity between NON-SUSY & SUSY QFT for IHO

Parameters of IHO	Features of graph and Lyapunov
General features	We do not observe any saturation behavior, and the complexity values rise quickly to very high values. Although we can not associate a Lyapunov exponent we make general statements about the rate of increase (hence the slope) of the graphs. We observe increased slopes upon adding perturbation terms.
$(F - \gamma)$	For even values the graphs are increasing exponentially whereas for odd values we see negative complexity values and hence have not included them in our graphical analysis.
Momentum p	The constant momentum factor acts as a scale multiplying the overall complexity value and is hence redundant.
Mass	For ϕ^3 perturbation, we observe as mass increases the rate also increases resulting in larger slopes. This is also true in the case of ϕ^4 theory, with the only difference being the symmetry along vertical-axis.
Temperature	The dependence on temperature can be evaluated by varying β (the inverse temperature). By varying this we can observe that for high temperatures, the rate of increase is much lesser as compared to lower temperatures in the case of IHO. This is true for both ϕ^3 and ϕ^4 perturbations. One can note that this behavior is contradictory to the upper bound that we can set for Lyapunov exponents giving us more incentive to not associate the slope of IHO with the Lyapunov exponent.
λ	For negative λ value we observe a mirror inversion of the graph about vertical-axis and hence the complexity are exponentially decreasing. One can simply interpret the opposite behavior of the graphs with mass and temperature variation in the case of negative λ . The change in value of λ contributes to the rate with increasing values resulting in faster rates of increase and hence higher slopes.

Table 8.4: Discussions on Lyapunov exponent for IHO complexity

9 Graphical Analysis

With the computed formulae for complexity for IHO and SHO in the previous sections we plot the graphs using different values of parameters.

9.1 SHO

For SHO, the parameters involved are the mass m and coupling coefficient λ for higher order theories. We have chosen a simple linear co-ordinate transformation between the field variables x and θ and u, v . This ensures that the Hessian is a constant and our computations become much simpler. The particular coefficients of the transformation have been chosen so that complexity is positive and rising for all cases.

In Fig. 9.1 we have plotted the three different complexities for a intermediate value of mass, keeping the coupling coefficient fixed. We observe that the complexity rises and saturates as expected.

It is important to know how the complexity value behaves for different masses. For lighter particles as we see in Fig. 9.2. We see a slight overall decrease in the complexity




Figure 9.1: Complexity against ϕ for $m = 1, \lambda = 1$




Figure 9.2: Complexity against ϕ for $m = 0.01, \lambda = 1$

values but more importantly we observe that the graph of ϕ^3 term tends very close to the ϕ^2 term. We conclude that as mass grows smaller and smaller, the ϕ^3 graph will inch closer and be in-differentiable to the ϕ^2 graph.




Figure 9.3: Complexity against ϕ for $m = 100$, $\lambda = 1$

For larger masses we see a stark difference. Here it is the graph of the ϕ^4 theory that inches closer to the ϕ^2 graph as seen in Fig. 9.3 and we can once again expect that it will become in-differentiable for much larger masses.

The other important parameter that we need to vary is the value of λ and we start by seeing what will happen when it is made negative as seen in Fig. 9.4. We see the behavior of the complexity remains same i.e. it grows and saturates as we go right. An interesting behavior occurs in the ϕ^4 graph - We see an initial rise and dip, before the rise and saturation. This indicates that we have a zero in the ϕ^4 theory.

We observe no behavioral changes when we vary the specific value of λ apart from the fact that for higher values it saturates much quickly (pointing to a larger Lyapunov exponent) and saturates slower for smaller values (smaller Lyapunov exponent).




Figure 9.4: Complexity against ϕ for $m = 1$, $\lambda = -1$

9.2 IHO

For the plots of IHO we need to keep in mind that there are many more parametric values. We will do a graphical analysis mainly by varying mass and temperature. We will merely state the redundancy involved in other parameters and therefore explain our choice of fixing it.

- If one looks closer to the formulae given for IHO in Sec. 6 we see the value of the momentum - p just adds an overall factor that multiplies the function thereby acting as a scaling factor. Hence we can set this to $p = 1$ without any issues.
- Another important parameter that might affect our plots is whether the value of $F - \gamma$ is Odd or Even. When it is even, we have positive complexity and when it is odd we find that we are dealing with negative complexity values which we safely ignore in the present context of discussion.

Before we proceed to varying the inverse temperature (β) and the mass (m), we can take a look at the general feature of the graphs of different perturbations in the field for both positive and negative λ as we have done in Fig. 9.5 and Fig. 9.6.

We see that for positive coupling coefficient the complexity values in all perturbations rise, but in different rates, whereas in the case of a negative coupling coefficient we see that they decrease rapidly after an initial period of rising (almost an inverse behavior of the positive λ case). We also observe symmetry for negative values of field in the $\phi^2 + \phi^4$ graph. It is important to note that although we can calculate the values of slope, one




Figure 9.5: Complexity against ϕ for different perturbations of field with $\lambda = 1$






Figure 9.6: Complexity against ϕ for different perturbations of field with $\lambda = -1$

cannot associate a *Lyapunov exponent* for the same because saturation does not exist. Maybe we can regard the small rise and saturation before the dip in the case of negative Lambda values, and associate a *Lyapunov exponent* to it.



(a) Variation of inverse temperature



(b) Variation of mass

Figure 9.7: behavior of the complexity of $\phi^2 + \phi^3$ perturbation with varying β and m

In Fig. 9.7, we have plotted the how the complexity values vary when we vary the inverse temperature and the mass. A very similar behavior is observed in the case of





Figure 9.8: behavior of the complexity of $\phi^2 + \phi^4$ perturbation with varying β and m

$\phi^2 + \phi^4$ perturbation too, which is shown in Fig. 9.8.

Since we have already established the behavior of the complexity in the case of negative λ

we take the liberty of not explicitly plotting the above variations in the inverse temperature and mass for that case. One simply has to imagine a decreasing (cannot associate a Lyapunov exponent) mirror inversion in the negative valued domain of the field. This is true for the cases of both perturbations that we have discussed in the present context.

10 Quantum Chaos from Morse function

Emergence of chaos in quantum phenomena can be estimated using an out-of-time-order correlation function which is firmly associated to the commutator of operator, split up in time. However, the universal relation $\mathcal{C} = -\ln(\text{OTOC})$ relating complexity with OTOC [42–47] has been studied greatly in recent times to diagnose chaos in various physical models. In this section we will prove this universal relation for supersymmetric case, relating the complexity with OTOC using the frame of Morse theory, then we will comment on the upper bound of chaos namely the Lyapunov exponent. In the above sections, by identifying the Morse function on a manifold with the cost function, we have calculated the complexity for supersymmetric field theories in various regimes in terms of the Hessian $H(f)$ and also have made use of the fact that the eigenvalues of supersymmetric Hamiltonian H_t are concentrated near the critical points of the Morse function defined on the manifold. In this section we will make use of the same facts and derive the universality relation.

Theorem - *Let $\phi(t)$ be an integral curve which represents the state of the particle at various time, then if x_1, \dots, x_m be a local coordinate chart around a critical point $p \in M$ such that $\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^m}$ is an orthonormal basis for $T_p M$ with respect to the metric g , then for any $t \in \mathbb{R}$ the out-of-time-correlator at p is equal to the exponential of the minus the matrix of the Hessian at p , i. e.*

$$\frac{\partial}{\partial x} \phi_t|_p = \exp(-H_p(f)t). \quad (10.1)$$

By identifying ϕ_t as a smooth function on the surface of the manifold we have

$$\frac{d}{dt} \phi(t, x) = -(\nabla f) \left(\frac{\partial}{\partial x} \phi(t, x) \right). \quad (10.2)$$

By changing the order of differentiation for any $x \in M$ we get,

$$\frac{d}{dt} \left(\frac{\partial}{\partial x} \phi(t, x) \right) = - \left(\frac{\partial}{\partial x} \nabla f \right) \left(\frac{\partial}{\partial x} \phi(t, x) \right). \quad (10.3)$$

Therefore by defining:

$$\Phi(t, x) = \frac{\partial}{\partial x} \phi(t, x) \quad (10.4)$$

is a solution of the linear system of ODE's:

$$\frac{d}{dt} \Phi(t, x) = - \left(\frac{\partial}{\partial x} \nabla f \right) \Phi(t, x). \quad (10.5)$$

Because $\exp \left(- \left(\frac{\partial}{\partial x} \nabla f \right) t \right)$ is also a solution to the linear ODE's, we finally get:

$$\Phi(t, x) = \frac{\partial}{\partial x} \phi(t, x) = \exp \left(- \left(\frac{\partial}{\partial x} \nabla f \right) t \right). \quad (10.6)$$

Since the solution is unique, thereby, at the location of critical point p we have:

$$\frac{\partial}{\partial x} \phi_t|_p = \exp(-H_p(f)t) \quad (10.7)$$

Hence by identifying complexity of SUSY field theories with the Hessian as described in this paper, the above equation under good approximations could be written as:

$$\mathcal{C} = - \ln(OTOC). \quad (10.8)$$

From the above equation, we could see that the Hessian of the Morse function which is a great tool to compute the complexity of supersymmetric theories, under good approximations is perfectly consistent with *universal relation* and is of great interest to capture the effect of chaos in supersymmetric field theories.

we will now comment on the behavior of Lyapunov exponent in SUSY field theories especially in the framework of an inverted harmonic oscillator under which it is expected to have chaotic features, depending upon the increase in the number of critical points as described in section 6. Thus we could write the expression for complexity in the IHO regime as:

$$\mathcal{C}_i(t) \approx c \exp(\lambda_i t) \quad \forall \quad i = 1, 2, \dots, n. \quad (10.9)$$

It is to be noted that the above equation is valid only for the IHO. The index i indicates the higher order quantum corrections in the Lagrangian for which the complexity has been measured, then mathematically Lyapunov exponent could be written as:

$$\lambda_i = \left(\frac{d \ln \mathcal{C}_i(t)}{dt} \right) \quad \forall \quad i = 1, 2, \dots, n. \quad (10.10)$$

Now using the universal relation relating complexity with the Out-of-Time Ordered Correlation (OTOC) function, one could write:

$$OTOC = \exp(-c \exp(\lambda t)) \quad (10.11)$$

where λ is identified as Lyapunov exponent as in with reference [48–50] which captures the effect of chaos in the quantum regime and relates different measures of complexity with OTOC through the universal relation. The above universal relation between complexities in different order of perturbations in the super-potential can be translated to the Lyapunov exponent through the MSS bound as:

$$\lambda_i \leq \lambda \leq \frac{2\pi}{\beta} \quad \forall \quad i = 1, 2, \dots, n. \quad (10.12)$$

where β is the inverse temperature. In this we have shown that the Morse function which could be used to classifies the topology of surfaces also captures the effect of chaos in the quantum regime of supersymmetric field theories.

11 Conclusions

Out-of-order-correlation-function (OTOC) in the framework of supersymmetry has been studied before and as a result didn't show any chaotic behavior in the regime of SHO. Our main focus in this paper was to bring out the relationship between circuit complexity and Morse function and comment on the complexity of supersymmetric quantum field theory, in the regime of the simple and inverted harmonic oscillator (IHO), by formulating the potential of IHO as the generators of $SL(2, Z)$ group. Circuit complexity has been calculated for various models starting from quantum field theory (QFT) to cosmology, which involves working out the cost functions for the particular case and then minimizing it, however, in this paper, we haven't done the same, by pointing out the relationship between the cost and Morse function on a manifold we have explicitly shown how the critical points on the surface encapsulate the action of the supersymmetric charge on the given reference state. We have explicitly made use of the fact that the eigenvalues of the supersymmetric hamiltonian are concentrated near the critical points of the Morse function in the manifold and then using the Witten index which comments on the symmetry breaking of the theory we commented on the complexity of the supersymmetric field theories for the IHO, which increases by a factor of exponential. For computations of complexity in the regime of simple harmonic oscillator, we have identified the auxiliary field in the SUSY lagrangian with the gradient of the Morse function and then computed the hessian, and found out that circuit complexity didn't show any dependence on initial conditions or exponential behavior. Next, we have the well known universal relation relating complexity and out-of-time ordered correlation function $\mathcal{C} = -\ln(OTOC)$ using the general description of Morse

theory.

- **Remark I:**

The circuit complexity for supersymmetric field theories has very deep connections with the Morse function defined on the surface of the manifold. We have obtained the expression for complexity for SUSY field theories in terms of the Hessian of the Morse function, in doing so, we have made use of the fact the eigenvalues of the supersymmetric hamiltonian are concentrated near the critical points of the Morse function.

- **Remark II:**

The behavior of complexity for supersymmetric field theories for the inverted harmonic oscillator is of prime importance, we have found that the growth of complexity in the regime of IHO is directly related to the growth of number of critical points on the manifold, which in turn grows exponentially with respect to the superpotential, we observed similar behavior for higher order quantum corrections namely ϕ^3 and ϕ^4 theories.

- **Remark III:**

We have also computed complexity for the simple harmonic oscillator, by identifying the auxiliary field in the SUSY lagrangian with the gradient of the Morse function and then computed the Hessian in the superspace coordinates, and found out that circuit complexity didn't show any dependence on initial conditions or exponential behavior. It is also worth mentioning that complexity for supersymmetric scalar field theories only depends on the absolute value of the non-dynamical auxiliary field. The F - term which we have identified with gradient of the Morse function determines whether the supersymmetry is spontaneously broken or not depending upon whether the gradient has passed through the critical points. On passing through the critical points we get $F = 0$ which tells that there exists no zero energy supersymmetric ground states.

- **Remark IV:**

We have proved the well known universal relation $\mathcal{C} = -\ln(OTOC)$ which relates complexity with the out-of-time ordered correlation function for supersymmetric field theories using Morse theory. The out-of-time ordered correlation function is an excellent gadget to capture the effect of chaos in the quantum regime. In this paper, we have obtained an upper bound on the Lyapunov exponent and also commented on its various features for supersymmetric field theories purely for SHO and IHO in table ?? and 8.3 which we have obtained purely on the basis of general description of Morse theory which gives quantum chaos a mathematical structure. The main point of Witten's paper on supersymmetry and Morse theory was to provide super-

symmetry a mathematical structure, various results like Witten index which tells if the supersymmetry is broken or not wouldn't have been possible by the normal description of particle physics.

- **Remark V:**

We have found that complexity for supersymmetric field theories differ significantly from ordinary quantum field theories in the sense that for non-SUSY QFT complexity, slowly starts to increase at the critical point, however for SUSY field theories the graph rises fast initially and then slowly tends to saturate, and the rate of saturation depends on the order of quantum corrections. The ϕ^3 theory saturates very rapidly and have the smallest Lyapunov exponent among the other studied perturbations, while the complexity for theory involving ϕ^4 term saturates slowly. For IHO we observed that complexity increases exponentially and quickly rose to very high values unlike ordinary QFT where it has a linear growth.

- **Remark VI:**

We have explicitly studied the dependence of mass on the behavior of circuit complexity and have observed that for massive fields there is a decrease in rate of change of complexity for ϕ^4 theory, and it is interesting to note that the graph becomes nearly indistinguishable from that of free field theory. However for smaller masses the complexity for ϕ^3 graph approaches to ϕ^2 slowly with a slight decrease in the rate of complexity and hence we expect an increase in the value of Lyapunov exponent. In the case of IHO, we observe that as mass increases the rate of change of complexity w.r.t ϕ also increases.

- **Remark VII:**

We have also commented on the behavior of complexity w.r.t the coupling constant λ and have observed that for negative values of λ in the regime of SHO the saturation is slower for ϕ^2 and ϕ^3 perturbations and in the case of ϕ^4 theory there is a sudden dip at the initial stage to zero thereby right shifting the point of initial rise of complexity. In the case of IHO, increase in the value of λ results in increase in the rate of change of complexity, however for negative values of λ we found that complexity decreases exponentially for SUSY field theories.

The future prospects of the present work are appended below:

- **Prospect I:**

In this paper we have restricted ourselves to supersymmetric scalar field theories,

however similar computations could be done for supersymmetric gauge and non-abelian gauge theories by taking in consideration the dynamical D - term[51? , 52], which would give further understanding about complexity and effect of chaos in supersymmetric field theories.

- **Prospect II:** circuit complexity for interacting quantum field theories and its relation with renormalization group has been studied by Arpan Bhattacharyya and collaborator and thus it will be interesting to see what new mathematical structure does renormalization group flow brings out and how it is related to complexity[53–55].
- **Prospect III:** In this paper, we have computed the complexity for SUSY scalar field theories by making use of the properties of the Hessian matrix, however we hope that this is not all[56]. Use of other remarkable properties of the Morse function could help in gaining much broader perspective in supersymmetric field theories and its matter content and interactions and the effect it has on the expansion of universe[57].
- **Prospect IV:** The study of supersymmetry and its complexity in terms of Morse theory has given it a geometrical structure, however for theories of supergravity it is still not quit clear what the right mathematical structure is[58, 59], we suppose that the work on this direction would bring interesting connections between quantum chaos and various other mathematical theories.

Acknowledgements

The research fellowship of SC is supported by the J. C. Bose National Fellowship of Sudhakar Panda. Also SC take this opportunity to thank sincerely to Sudhakar Panda for his constant support and providing huge inspiration. SC also would like to thank School of Physical Sciences, National Institute for Science Education and Research (NISER), Bhubaneswar for providing the work friendly environment. SC also thank all the members of our newly formed virtual international non-profit consortium “Quantum Structures of the Space-Time & Matter” (QASTM) for elaborative discussions. Satyaki Choudhury, Sachin Panner Selvam and K. Shirish would like to thank NISER Bhubaneswar, BITS Hyderabad, VNIT Nagpur respectively for providing fellowships. Last but not the least, we would like to acknowledge our debt to the people belonging to the various part of the world for their generous and steady support for research in natural sciences.

References

- [1] N. Lashkari, M. B. McDermott, and M. Van Raamsdonk, “Gravitational dynamics from entanglement ‘thermodynamics’,” *JHEP* **04** (2014) 195, [arXiv:1308.3716 \[hep-th\]](#).
- [2] M. Van Raamsdonk, “Building up spacetime with quantum entanglement,” *Gen. Rel. Grav.* **42** (2010) 2323–2329, [arXiv:1005.3035 \[hep-th\]](#).
- [3] J. Maldacena and L. Susskind, “Cool horizons for entangled black holes,” *Fortsch. Phys.* **61** (2013) 781–811, [arXiv:1306.0533 \[hep-th\]](#).
- [4] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” *Int. J. Theor. Phys.* **38** (1999) 1113–1133, [arXiv:hep-th/9711200](#).
- [5] T. Li, J. Liu, Y. Xin, and Y. Zhou, “Supersymmetric SYK model and random matrix theory,” *JHEP* **06** (2017) 111, [arXiv:1702.01738 \[hep-th\]](#).
- [6] S. Choudhury, A. Dey, I. Halder, L. Janagal, S. Minwalla, and R. Poojary, “Notes on melonic $O(N)^{q-1}$ tensor models,” *JHEP* **06** (2018) 094, [arXiv:1707.09352 \[hep-th\]](#).
- [7] L. Susskind, “Entanglement is not enough,” *Fortsch. Phys.* **64** (2016) 49–71, [arXiv:1411.0690 \[hep-th\]](#).
- [8] L. Susskind and Y. Zhao, “Switchbacks and the Bridge to Nowhere,” [arXiv:1408.2823 \[hep-th\]](#).
- [9] S. H. Shenker and D. Stanford, “Black holes and the butterfly effect,” *JHEP* **03** (2014) 067, [arXiv:1306.0622 \[hep-th\]](#).
- [10] B. Swingle and M. Van Raamsdonk, “Universality of Gravity from Entanglement,” [arXiv:1405.2933 \[hep-th\]](#).
- [11] S. Choudhury and S. Panda, “Entangled de Sitter from stringy axionic Bell pair I: an analysis using Bunch–Davies vacuum,” *Eur. Phys. J.* **C78** no. 1, (2018) 52, [arXiv:1708.02265 \[hep-th\]](#).
- [12] S. Choudhury and S. Panda, “Quantum entanglement in de Sitter space from stringy axion: An analysis using α vacua,” *Nucl. Phys.* **B943** (2019) 114606, [arXiv:1712.08299 \[hep-th\]](#).
- [13] S. Choudhury, S. Panda, and R. Singh, “Bell violation in the Sky,” *Eur. Phys. J.* **C77** no. 2, (2017) 60, [arXiv:1607.00237 \[hep-th\]](#).
- [14] D. A. Roberts, D. Stanford, and L. Susskind, “Localized shocks,” *JHEP* **03** (2015) 051, [arXiv:1409.8180 \[hep-th\]](#).
- [15] S. Akhtar, S. Choudhury, S. Chowdhury, D. Goswami, S. Panda, and A. Swain, “Open Quantum Entanglement: A study of two atomic system in static patch of de Sitter space,” *Eur. Phys. J.* **C80** no. 8, (2020) 748, [arXiv:1908.09929 \[hep-th\]](#).
- [16] K. Y. Bhagat, B. Bose, S. Choudhury, S. Chowdhury, R. N. Das, S. G. Dastider, N. Gupta, A. Maji, G. D. Pasquino, and S. Paul, “The Generalized OTOC from Supersymmetric Quantum Mechanics: Study of Random Fluctuations from Eigenstate Representation of

- Correlation Functions,” *Symmetry* **13** no. 1, (2020) 44, [arXiv:2008.03280 \[hep-th\]](#).
- [17] M. R. Dowling and M. A. Nielsen, “The geometry of quantum computation,” *arXiv preprint quant-ph/0701004* (2006) .
- [18] R. Jefferson and R. C. Myers, “Circuit complexity in quantum field theory,” *JHEP* **10** (2017) 107, [arXiv:1707.08570 \[hep-th\]](#).
- [19] M. A. Nielsen, “A geometric approach to quantum circuit lower bounds,” *arXiv preprint quant-ph/0502070* (2005) .
- [20] R. Auzzi, S. Baiguera, G. B. De Luca, A. Legramandi, G. Nardelli, and N. Zenoni, “On the geometry of quantum complexity,” [arXiv:2011.07601 \[hep-th\]](#).
- [21] D. A. Roberts and B. Yoshida, “Chaos and complexity by design,” *JHEP* **04** (2017) 121, [arXiv:1610.04903 \[quant-ph\]](#).
- [22] S. Choudhury, A. Dutta, and D. Ray, “Chaos and Complexity from Quantum Neural Network: A study with Diffusion Metric in Machine Learning,” [arXiv:2011.07145 \[hep-th\]](#).
- [23] A. Bhattacharyya, W. Chemissany, S. S. Haque, J. Murugan, and B. Yan, “The Multi-faceted Inverted Harmonic Oscillator: Chaos and Complexity,” [arXiv:2007.01232 \[hep-th\]](#).
- [24] V. Subramanyan, S. S. Hegde, S. Vishveshwara, and B. Bradlyn, “Physics of the Inverted Harmonic Oscillator: From the lowest Landau level to event horizons,” [arXiv:2012.09875 \[cond-mat.mes-hall\]](#).
- [25] G. Barton, “Quantum Mechanics of the Inverted Oscillator Potential,” *Annals Phys.* **166** (1986) 322.
- [26] S. Hawking, “Black hole explosions,” *Nature* **248** (1974) 30–31.
- [27] N. Birrell and P. Davies, *Quantum Fields in Curved Space*. Cambridge Monographs on Mathematical Physics. Cambridge Univ. Press, Cambridge, UK, 2, 1984.
- [28] L. C. Crispino, A. Higuchi, and G. E. Matsas, “The Unruh effect and its applications,” *Rev. Mod. Phys.* **80** (2008) 787–838, [arXiv:0710.5373 \[gr-qc\]](#).
- [29] N. Read and E. Rezayi, “Hall viscosity, orbital spin, and geometry: paired superfluids and quantum Hall systems,” *Phys. Rev. B* **84** (2011) 085316, [arXiv:1008.0210 \[cond-mat.mes-hall\]](#).
- [30] M. Fecko, *Differential geometry and Lie groups for physicists*. Cambridge University Press, 3, 2011.
- [31] M. Kerr, “Notes on the representation theory of $sl_2(\mathbb{r})$,” *Hodge Theory, Complex Geometry, and Representation Theory* **608** (2014) 173.
- [32] J. Milnor, *Morse theory.(AM-51)*, vol. 51. Princeton university press, 2016.
- [33] E. Witten, “Constraints on Supersymmetry Breaking,” *Nucl. Phys. B* **202** (1982) 253.

- [34] M. Drees, “An Introduction to supersymmetry,” in *Inauguration Conference of the Asia Pacific Center for Theoretical Physics (APCTP)*. 11, 1996. [arXiv:hep-ph/9611409](#).
- [35] E. Witten, “Dynamical Breaking of Supersymmetry,” *Nucl. Phys. B* **188** (1981) 513.
- [36] E. Witten, “Supersymmetry and Morse theory,” *J. Diff. Geom.* **17** no. 4, (1982) 661–692.
- [37] S. Weinberg, *The quantum theory of fields. Vol. 3: Supersymmetry*. Cambridge University Press, 6, 2013.
- [38] S. J. Gates, M. T. Grisaru, M. Rocek, and W. Siegel, *Superspace Or One Thousand and One Lessons in Supersymmetry*, vol. 58 of *Frontiers in Physics*. 1983. [arXiv:hep-th/0108200](#).
- [39] F. Quevedo, S. Krippendorf, and O. Schlotterer, “Cambridge Lectures on Supersymmetry and Extra Dimensions,” [arXiv:1011.1491 \[hep-th\]](#).
- [40] Y. Shirman, “Introduction to Supersymmetry and Supersymmetry Breaking,” in *Theoretical Advanced Study Institute in Elementary Particle Physics: The Dawn of the LHC Era*, pp. 359–422. 2010. [arXiv:0907.0039 \[hep-ph\]](#).
- [41] J. Wess and B. Zumino, “Supergauge Invariant Extension of Quantum Electrodynamics,” *Nucl. Phys. B* **78** (1974) 1.
- [42] S. Choudhury, S. Chowdhury, N. Gupta, A. Mishara, S. P. Selvam, S. Panda, G. D. Pasquino, C. Singha, and A. Swain, “Magical Chaotic Cosmological Islands: Generating Page Curve to solve Black Hole Information loss problem from Cosmological Chaos – Complexity connection,” [arXiv:2012.10234 \[hep-th\]](#).
- [43] P. Bhargava, S. Choudhury, S. Chowdhury, A. Mishara, S. P. Selvam, S. Panda, and G. D. Pasquino, “Quantum aspects of chaos and complexity from bouncing cosmology: A study with two-mode single field squeezed state formalism,” [arXiv:2009.03893 \[hep-th\]](#).
- [44] K. Hashimoto, K.-B. Huh, K.-Y. Kim, and R. Watanabe, “Exponential growth of out-of-time-order correlator without chaos: inverted harmonic oscillator,” *Journal of High Energy Physics* **2020** no. 11, (2020) 1–25.
- [45] S. Choudhury, “The Cosmological OTOC: Formulating new cosmological micro-canonical correlation functions for random chaotic fluctuations in Out-of-Equilibrium Quantum Statistical Field Theory,” *Symmetry* **12** no. 9, (2020) 1527, [arXiv:2005.11750 \[hep-th\]](#).
- [46] S. Choudhury and S. Panda, “Cosmological Spectrum of Two-Point Correlation Function from Vacuum Fluctuation of Stringy Axion Field in De Sitter Space: A Study of the Role of Quantum Entanglement,” *Universe* **6** no. 6, (2020) 79, [arXiv:1809.02905 \[hep-th\]](#).
- [47] S. Choudhury, A. Mukherjee, P. Chauhan, and S. Bhattacharjee, “Quantum Out-of-Equilibrium Cosmology,” *Eur. Phys. J.* **C79** no. 4, (2019) 320, [arXiv:1809.02732 \[hep-th\]](#).
- [48] J. Maldacena, S. H. Shenker, and D. Stanford, “A bound on chaos,” *JHEP* **08** (2016) 106, [arXiv:1503.01409 \[hep-th\]](#).

- [49] V. Bunakov, “Quantum signatures of chaos or quantum chaos?,” *Phys. Atom. Nucl.* **79** no. 6, (2016) 995–1009.
- [50] X. Han and S. A. Hartnoll, “Quantum Scrambling and State Dependence of the Butterfly Velocity,” *SciPost Phys.* **7** no. 4, (2019) 045, [arXiv:1812.07598 \[hep-th\]](#).
- [51] H. Itoyama and N. Maru, “D-term Triggered Dynamical Supersymmetry Breaking,” *Phys. Rev. D* **88** no. 2, (2013) 025012, [arXiv:1301.7548 \[hep-ph\]](#).
- [52] O. Piguet, “Introduction to supersymmetric gauge theories,” in *1st School on Field Theory and Gravitation*. 4, 1997. [arXiv:hep-th/9710095](#).
- [53] M. Bonini and F. Vian, “Wilson renormalization group for supersymmetric gauge theories and gauge anomalies,” *Nucl. Phys. B* **532** (1998) 473–497, [arXiv:hep-th/9802196](#).
- [54] D. Kazakov, “Supersymmetry in particle physics: the renormalization group viewpoint,” *Phys. Rept.* **344** (2001) 309–353, [arXiv:hep-ph/0001257](#).
- [55] K. Maruyoshi and J. Song, “Enhancement of Supersymmetry via Renormalization Group Flow and the Superconformal Index,” *Phys. Rev. Lett.* **118** no. 15, (2017) 151602, [arXiv:1606.05632 \[hep-th\]](#).
- [56] J. Murugan, D. Stanford, and E. Witten, “More on supersymmetric and 2d analogs of the syk model,” *Journal of High Energy Physics* **2017** no. 8, (2017) 146.
- [57] N. Bilic, “Supersymmetric dark energy,” *Rom. J. Phys.* **57** (2012) 793–802, [arXiv:1004.4984 \[hep-th\]](#).
- [58] A. Coimbra, C. Strickland-Constable, and D. Waldram, “Supergravity as Generalised Geometry I: Type II Theories,” *JHEP* **11** (2011) 091, [arXiv:1107.1733 \[hep-th\]](#).
- [59] C. Lazaroiu and C. Shahbazi, “Four-dimensional geometric supergravity and electromagnetic duality: a brief guide for mathematicians,” [arXiv:2006.16157 \[math.DG\]](#).