

**PATH-DEPENDENT HAMILTON-JACOBI EQUATIONS
WITH SUPER-QUADRATIC GROWTH IN THE GRADIENT
AND THE VANISHING VISCOSITY METHOD**

ERHAN BAYRAKTAR AND CHRISTIAN KELLER

ABSTRACT. The non-exponential Schilder-type theorem in Backhoff-Veraguas, Lacker and Tangpi [Ann. Appl. Probab., 30 (2020), pp. 1321–1367] is expressed as a convergence result for path-dependent partial differential equations with appropriate notions of generalized solutions. This entails a non-Markovian counterpart to the vanishing viscosity method.

We show uniqueness of maximal subsolutions for path-dependent viscous Hamilton-Jacobi equations related to convex super-quadratic backward stochastic differential equations.

We establish well-posedness for the Hamilton-Jacobi-Bellman equation associated to a Bolza problem of the calculus of variations with path-dependent terminal cost. In particular, uniqueness among lower semi-continuous solutions holds and state constraints are admitted.

1. INTRODUCTION

Backhoff-Veraguas, Lacker and Tangpi [2] derived a non-exponential Schilder-type theorem, which they used to obtain new limit theorems for backward stochastic differential equations (BSDEs) and, in the Markovian case, for the corresponding partial differential equations (PDEs). They posed the question whether it is possible to have a corresponding PDE result in the non-Markovian case as well. Purpose of our work is to provide an answer to this question.

We establish well-posedness for (second order) path-dependent viscous Hamilton-Jacobi equations and for (first-order) path-dependent Hamilton-Jacobi-Bellman (HJB) equations with possibly super-quadratic growth in the gradient. Together with a modification of the Schilder type theorem in [2], we obtain a non-Markovian vanishing viscosity result for path-dependent PDEs and thereby address the mentioned open problem in [2].

The notions of solutions for our path-dependent PDEs are in the spirit of contingent solutions for PDEs (see, e.g., [23]), also known as Dini solutions (see, e.g., [3] and [45]) or minimax solutions (see, e.g., [42]).

In the context of contingent or Dini solutions for first-order standard PDEs related to Bolza problems, [13] and also [35] are very close to our approach. More recent works in this direction are [33], [5] and [6]. Regarding the possible use of viscosity solution techniques, we refer the reader to the remarks on p. 1202 in [7].

2010 *Mathematics Subject Classification.* 60H30; 60F10, 35F21, 35K10, 49J22.

Key words and phrases. Large deviations, vanishing viscosity method, path-dependent partial differential equations, minimax solutions, Dini solutions, calculus of variations, optimal control, state constraints, backward stochastic differential equations, nonsmooth analysis.

The research of the first author was supported in part by the National Science Foundation.

In particular, fast growth in the gradient, discontinuity of the Lagrangian, and extended real-valued lower semi-continuous terminal data (to allow right-end point constraints in optimal control problems) cause non-trivial issues. For example, solutions of HJB equations can be expected then to be only lower semi-continuous.

In the second-order case, the only works we are aware of that use contingent-type solutions are [44] and [43], where Isaacs equations corresponding to Markovian stochastic differential games with drift control and bounded control spaces are studied. However, similar constructions (stochastic or Gaussian derivatives) are also used in [25] and [26].

In the context of first-order path-dependent PDEs, [28] is most relevant. It deals with a calculus of variations problem involving a path-dependent terminal cost and the related path-dependent HJB equation. The setting is very close to our problem (DOC) below. The main difference is that in [28] the terminal cost is required to be Lipschitz continuous, which leads to Lipschitz continuity of the corresponding value function and also makes it possible in [28] to develop a viscosity solution theory. In our work, we require only continuity resp. lower semi-continuity of the terminal cost, which is one of the reasons why we establish a Dini resp. minimax solution theory. For the current state of the art for first-order path-dependent PDEs and for further relevant works, see [24] and the references therein.

In the literature [11, 18–22, 29, 34, 36–41] on viscosity solutions of second-order path-dependent PDEs, the Hamiltonian is required to grow at most linearly in the gradient (the same condition is also needed in [12], where a notion of strong-viscosity solutions is used). Overcoming this restriction for any notion of generalized solutions has been a longstanding open problem. By proving wellposedness of maximal (Dini) subsolutions for a class of second-order path-dependent PDEs with quadratic and even super-quadratic growth in the gradient, we establish first results related to this problem.

Non-Markovian large deviation problems and their connections to path-dependent PDEs are also studied in [32]. In contrast to our work, in [32] only the (limiting) rate function is characterized as a solution of a (first-order) path-dependent PDE. Moreover, the terminal condition is required to be Lipschitz continuous whereas we need only continuity.

2. SETUP

2.1. Notation. Let $\Omega = C([0, T], \mathbb{R}^d)$. The canonical process on Ω is denoted by X , i.e., $X(t, \omega) = \omega(t)$ for each $(t, \omega) \in [0, T] \times \Omega$. Let $\mathbb{F} = \{\mathcal{F}_t\}_{t \in [0, T]}$ be the (raw) filtration generated by X . Given a probability measure P on (Ω, \mathcal{F}_T) , denote by $\mathbb{F}^P = \{\mathcal{F}_t^P\}_{t \in [0, T]}$ the P -completion of the right-limit of \mathbb{F} .

We equip Ω with the supremum norm $\|\cdot\|_\infty$ and $[0, T] \times \Omega$ with the pseudo-metric \mathbf{d}_∞ defined by

$$\mathbf{d}_\infty((t_1, \omega_1), (t_2, \omega_2)) := |t_1 - t_2| + \sup_{s \in [0, T]} |\omega_1(s \wedge t_1) - \omega_2(s \wedge t_2)|.$$

Continuity and semi-continuity of functions defined on Ω (resp. $[0, T] \times \Omega$) are to be understood with respect to $\|\cdot\|_\infty$ (resp. \mathbf{d}_∞). Note that semi-continuous functions on $[0, T] \times \Omega$ are \mathbb{F} -progressive. From now on, we write l.s.c. (resp. u.s.c.) instead of lower semi-continuous (resp. upper semi-continuous).

Given $(t_0, x_0, n) \in [0, T] \times \Omega \times \mathbb{N}$, denote by $P_{t_0, x_0, n}$ be the unique probability measure on (Ω, \mathcal{F}_T) such that $\sqrt{n}X|_{[t_0, T]}$ is a d -dimensional standard $(P_{t_0, x_0, n}, \mathbb{F})$ -Wiener process starting at $x_0(t_0)$ and that $P_{t_0, x_0, n}(X|_{[0, t_0]} = x_0|_{[[0, t_0]}) = 1$. We write $\mathbb{E}_{t_0, x_0, n}$ for the corresponding expected value. Moreover, $P_{t_0, x_0} := P_{t_0, x_0, 1}$ and $\mathbb{F}^{t_0, x_0, n} := \mathbb{F}^{P_{t_0, x_0, n}}$.

As space of controls, the set \mathcal{L}_b of all bounded \mathbb{F} -progressive processes from $[0, T] \times \Omega$ to \mathbb{R}^d is used (whereas in [2] the controls are $\mathbb{F}^{P_{0,0}}$ -progressive).

We denote by dom the effective domain of an extended real-valued function.

2.2. Data. Let $h : \Omega \rightarrow \mathbb{R} \cup \{\infty\}$ and $\ell : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\infty\}$ be measurable functions. We use the following hypotheses for ℓ .

(H1) The function $\ell = \ell(t, a)$ satisfies the Tonelli-Nagumo condition, i.e., there is a function $\phi : [0, \infty) \rightarrow \mathbb{R}$ bounded from below with $\phi(r)/r \rightarrow \infty$ as $r \rightarrow \infty$ such that $\ell(t, a) \geq \phi(|a|)$ on $[0, T] \times \mathbb{R}^d$. Moreover, $\ell(t, \cdot)$ is l.s.c., proper, and convex for every $t \in [0, T]$. (H2) $\int_0^T \ell(t, 0) dt < \infty$.

These hypotheses are nearly identical with the corresponding condition (TI) for the Lagrangian in [2] (where it is denoted by g). In some of our main results, we will, in addition to (H1) and (H2), also assume that ℓ is continuous and finite-valued. In those cases, (TI) is satisfied as pointed out in [2].

2.3. The optimal control problems and HJB equations. Let $n \in \mathbb{N}$. The value for our stochastic optimal control problem (SOC $_n$) with initial data $(t_0, x_0) \in [0, T] \times \Omega$ is given by

$$v_n(t_0, x_0) := \inf_{a \in \mathcal{L}_b^{t_0}} \mathbb{E}_{t_0, x_0, n} \left[\int_{t_0}^T \ell(t, a(t)) dt + h(X + A^a) \right],$$

where $\mathcal{L}_b^{t_0} = \{a \in \mathcal{L}_b : a|_{[0, t_0]} = 0\}$ and A^a is a continuous process on $[0, T] \times \Omega$ defined by $A^a(t) := \int_0^t a(s) ds$. The terminal value problem involving the corresponding HJB equation is

$$\begin{aligned} \text{(TVP } n) \quad \left(\partial_t + \frac{1}{2n} \partial_{xx} \right) u(t, x) + \inf_{a \in \mathbb{R}^d} [a \cdot \partial_x u(t, x) + \ell(t, a)] &= 0 \quad \text{in } [0, T] \times \Omega, \\ u(T, x) &= h(x) \quad \text{on } \Omega. \end{aligned}$$

Remark 2.1. If (H1) and (H2) hold and h is bounded, then $\sup_{n \geq 1} \|v_n\|_\infty < \infty$.

The value for our deterministic optimal control problem (DOC) with initial data $(t_0, x_0) \in [0, T] \times \Omega$ is given by

$$v_0(t_0, x_0) := \inf_{x \in \mathcal{X}^{1,1}(t_0, x_0)} \left[\int_{t_0}^T \ell(t, x'(t)) dt + h(x) \right],$$

where

$$\mathcal{X}^{1,1}(t_0, x_0) := \{x \in \Omega : x|_{[0, t_0]} = x_0|_{[0, t_0]} \text{ and } x|_{[t_0, T]} \in W^{1,1}(t_0, T; \mathbb{R}^d)\}.$$

The terminal value problem involving the corresponding HJB equation is

$$\begin{aligned} \text{(TVP)} \quad \partial_t u(t, x) + \inf_{a \in \mathbb{R}^d} [a \cdot \partial_x u(t, x) + \ell(t, a)] &= 0 \quad \text{in } [0, T] \times \Omega, \\ u(T, x) &= h(x) \quad \text{on } \Omega. \end{aligned}$$

3. NOTIONS OF SOLUTIONS OF PATH-DEPENDENT HJB EQUATIONS

3.1. Dini solutions. Given a function $u : [0, T] \times \Omega \rightarrow \mathbb{R} \cup \{\infty\}$, we define the *lower* and *upper Dini derivative*

$$d_-u(t_0, x_0)(1, a) := \liminf_{\delta \downarrow 0} \frac{u(t_0 + \delta, x_0(\cdot \wedge t_0) + A^a - A^a(t_0)) - u(t_0, x_0)}{\delta} \quad \text{and}$$

$$d_+u(t_0, x_0)(1, a) := \limsup_{\delta \downarrow 0} \frac{u(t_0 + \delta, x_0(\cdot \wedge t_0) + A^a - A^a(t_0)) - u(t_0, x_0)}{\delta}$$

at points $(t_0, x_0) \in [0, T] \times \Omega$ (resp. $(0, T] \times \Omega$) in direction $(1, a) \in \mathbb{R} \times \mathbb{R}^d$. Here, in unison with the process A^a in Subsection 2.3, $A^a(t) = at$.

The following (path-dependent) notion of Dini semi-solutions is motivated by the notion of (contingent) solutions used in Theorem 4.1 of [13] for HJB equations related to Bolza problems. Our notion is also related to the infinitesimal version of minimax solutions for path-dependent Isaacs equations used in [31].

Definition 3.1. Let $u : [0, T] \times \Omega \rightarrow \mathbb{R} \cup \{\infty\}$.

(i) We call u a *Dini supersolution* of (TVP) if u is l.s.c., $u(T, \cdot) \geq h$, and, for every $(t_0, x_0) \in \text{dom}(u)$ with $t_0 < T$,

$$(3.1) \quad \inf_{a \in \mathbb{R}^d} [d_-u(t_0, x_0)(1, a) + \ell(t_0, a)] \leq 0.$$

(ii) We call u a *Dini subsolution* of (TVP) if u is u.s.c., $u(T, \cdot) \leq h$, and, for every $(t_0, x_0) \in \text{dom}(u)$ with $t_0 < T$,

$$(3.2) \quad \inf_{a \in \mathbb{R}^d} [d_+u(t_0, x_0)(1, a) + \ell(t_0, a)] \geq 0.$$

We call u a *maximal Dini subsolution* of (TVP) if u is a Dini subsolution of (TVP) and, for every Dini subsolution v of (TVP), we have $u \geq v$.

Example 3.2. Let $d = 1$, $h = \infty \cdot \mathbf{1}_{K^c}$, where $K := \{t \mapsto t^{1/2}\} \subset \Omega$, and ℓ be defined by $\ell(t, a) = |a|^{3/2}$. Then v_0 satisfies

$$v_0(t_0, x_0) = \begin{cases} \int_{t_0}^T 2^{-3/2} t^{-3/4} dt = 2^{1/2} (T^{1/4} - t_0^{1/4}) & \text{if } x_0|_{[0, t_0]}(t) = t^{1/2}, \\ \infty & \text{otherwise.} \end{cases}$$

Note that, for each $(t_0, x_0) \in \text{dom}(v_0)$ with $t_0 < T$ and each $a \in \mathbb{R}$, we have $d_-v(t_0, x_0)(1, a) = \infty$ for which the constant perturbation involving a is responsible. (To obtain a finite value for our Dini derivative, we would need to permit the non-constant perturbation $t \mapsto t^{1/2}$.) Thus $\inf_{a \in \mathbb{R}} [d_-v(t_0, x_0)(1, a) + |a|^{3/2}] \leq 0$ is never satisfied. Hence, in our example there does not exist a Dini supersolution of (TVP). This justifies the need for an appropriate weaker notion of solution (see Subsection 3.2).

Given $u : [0, T] \times \Omega \rightarrow \mathbb{R}$, we define the *upper stochastic Dini derivative*

$$d_+^{1,2}u(t_0, x_0)(1, a, n^{-1}I_d) := \limsup_{\delta \downarrow 0} \frac{\mathbb{E}_{t_0, x_0, n} [u(t_0 + \delta, X + A^a - A^a(t_0)) - u(t_0, x_0)]}{\delta}.$$

at points $(t_0, x_0) \in [0, T] \times \Omega$ in direction $(1, a, n^{-1}I_d) \in \mathbb{R} \times \mathbb{R}^d \times \mathbb{R}^{d \times d}$ (cf. [25, 26, 43, 44]).

The following notion of subsolutions for second order path-dependent PDEs is motivated by the minimax solutions used in [43, 44] in a Markovian framework.

Definition 3.3. Let $u : [0, T] \times \Omega \rightarrow \mathbb{R}$. We call u a *Dini subsolution* of (TVP n) if u is u.s.c., $u(T, \cdot) \leq h$, and, for every $(t_0, x_0) \in [0, T] \times \Omega$,

$$(3.3) \quad \inf_{a \in \mathbb{R}^d} \left[d_+^{1,2} u(t_0, x_0)(1, a, n^{-1} I_d) + \ell(t_0, a) \right] \geq 0.$$

We call u a *maximal Dini subsolution* of (TVP n) if u is a Dini subsolution of (TVP n) and, for every Dini subsolution v of (TVP n), we have $u \geq v$.

Remark 3.4. In our specific setting, the use of Dini type semiderivatives such as those introduced in this section suffices. This motivates us to call our generalized solutions Dini solutions. For more general data, path-dependent counterparts of contingent derivatives such as the Clio derivatives in [1] need to be utilized. Corresponding generalized solutions would be called contingent solutions.

3.2. Minimax solutions. Here, we introduce a weaker notion of solution, which is an adjustment of the infinitesimal notion of minimax solutions in [4]. It is also motivated by the notion of (l.s.c. contingent) solutions used in Theorem 5.1 of [13]. The problem in Example 3.2, which partially motivated this weaker notion, is overcome by allowing non-constant perturbations (see also the notion of contingent solutions in [9] that are defined via contingent derivatives with function-valued directions).

Definition 3.5. Let $u : [0, T] \times \Omega \rightarrow \mathbb{R} \cup \{\infty\}$.

(i) We call u a *minimax supersolution* of (TVP) if u is l.s.c., $u(T, \cdot) \geq h$, and, for every $(t_0, x_0) \in \text{dom}(u)$ with $t_0 < T$,

$$(3.4) \quad \inf_{x \in \mathcal{X}^{1,1}(t_0, x_0)} \lim_{\delta \downarrow 0} \left[u(t_0 + \delta, x) - u(t_0, x_0) + \int_{t_0}^{t_0 + \delta} \ell(s, x'(s)) ds \right] \delta^{-1} \leq 0.$$

(ii) We call u an *l.s.c. minimax subsolution* of (TVP) if u is l.s.c., $u(T, \cdot) \leq h$, and, for every $(t_0, x_0) \in [0, T] \times \Omega$ and every $(t, x) \in ((t_0, T] \times \mathcal{X}^{1,1}(t_0, x_0)) \cap \text{dom}(u)$ with $\int_{t_0}^t \ell(s, x'(s)) ds < \infty$,

$$(3.5) \quad \lim_{\delta \downarrow 0} \left[u(t - \delta, x) - u(t, x) - \int_{t - \delta}^t \ell(s, x'(s)) ds \right] \delta^{-1} \leq 0.$$

(iii) We call u an *l.s.c. minimax solution* of (TVP) if u is a minimax super- and an l.s.c. minimax subsolution of (TVP).

4. MAIN RESULTS

Theorem 4.1. *Assume (H1) and (H2). Let h be continuous and bounded. Then (v_n) converges to v_0 uniformly on compacta and v_0 is continuous. Moreover, v_0 is the unique l.s.c. minimax solution of (TVP) that is bounded from below. If, in addition, ℓ is continuous and finite-valued, then we have the following:*

(i) *For each $n \in \mathbb{N}$, the function v_n is the unique bounded maximal Dini subsolution of (TVP n)*

(ii) *The function v_0 is the unique bounded maximal Dini subsolution of (TVP).*

Proof. The uniform convergence of (v_n) to v_0 on compacta and the continuity of v_0 are proven in Section 5. Theorem 4.3 characterizes v_0 as the unique minimax solution resp. as the unique bounded maximal Dini subsolution of (TVP). Theorem 7.2 addresses the remaining part, i.e., wellposedness of (TVP n). \square

Remark 4.2. Well-posedness of (TVP n) requires h to be only u.s.c. and bounded (see Theorem 7.2). The corresponding result in the Markovian case treated in [2] is of similar strength (well-posedness holds for maximal viscosity supersolutions of the corresponding viscous Hamilton-Jacobi equations, which is due to [17]).

Theorem 4.3. *Assume (H1).*

(a) *Let h be l.s.c., proper, and bounded from below. Then the value function v_0 is the unique l.s.c. minimax solution of (TVP) that is bounded from below.*

(b) *Assume (H2). Let ℓ be continuous and finite-valued. Let h be u.s.c. and bounded. Then v_0 is the unique maximal bounded Dini subsolution of (TVP).*

Proof. See Section 8. □

5. PROOF OF THE CONVERGENCE RESULT

Consider the semicontinuous envelopes v_* and v^* defined by

$$\begin{aligned} v_*(t_0, x_0) &:= \sup_{\substack{\delta > 0, \\ n \in \mathbb{N}}} \inf_{\substack{(t,x) \in O_\delta(t_0, x_0), \\ m \geq n}} v_m(t, x), \\ v^*(t_0, x_0) &:= \inf_{\substack{\delta > 0, \\ n \in \mathbb{N}}} \sup_{\substack{(t,x) \in O_\delta(t_0, x_0), \\ m \geq n}} v_m(t, x) \end{aligned}$$

for every $(t_0, x_0) \in [0, T] \times \Omega$. Here, $O_\delta(t_0, x_0)$ is the open δ -neighborhood of (t_0, x_0) in $([0, T] \times \Omega, \mathbf{d}_\infty)$.

First, we establish an auxiliary result.

Lemma 5.1. *Assume (H1). Let $t_n \rightarrow t_0$ in $[0, T]$ as $n \rightarrow \infty$. Consider a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$. Let (a_n) be a sequence in $L^1 = L^1([0, T] \times \tilde{\Omega}, dt \otimes d\tilde{P}; \mathbb{R}^d)$ that converges weakly to some $a \in L^1$. Then $\mathbb{E}^{\tilde{P}} \int_{t_0}^T \ell(t, a(t)) dt \leq \underline{\lim}_n \mathbb{E}^{\tilde{P}} \int_{t_n}^T \ell(t, a_n(t)) dt$.*

Proof. We follow the lines of the proof of the corresponding deterministic closure theorem 10.8.ii in [10]. First, note that, by lower semi-continuity and convexity of ℓ , for every $\delta > 0$, we have $\mathbb{E}^{\tilde{P}} \int_{t_0+\delta}^T \ell(t, a(t)) dt \leq \underline{\lim}_n \mathbb{E}^{\tilde{P}} \int_{t_0+\delta}^T \ell(t, a_n(t)) dt$ (more details can be found the proof of Lemma A.1 of [2]). Next, fix $\varepsilon > 0$. Since ℓ is bounded from below, there is a $\delta_0 > 0$ independent from n such that, for all $\delta \in (0, \delta_0)$, we have $\mathbb{E}^{\tilde{P}} \int_{t_n}^{t_0+\delta} \ell(t, a_n(t)) dt > -\varepsilon$ and thus $\underline{\lim}_n \mathbb{E}^{\tilde{P}} \int_{t_n}^T \ell(t, a_n(t)) dt + \varepsilon \geq \mathbb{E}^{\tilde{P}} \int_{t_0+\delta}^T \ell(t, a(t)) dt$. Again, as ℓ is bounded from below, either the right-hand side of the previous inequality converges to $\mathbb{E}^{\tilde{P}} \int_{t_0}^T \ell(t, a(t)) dt$ as $\delta \rightarrow 0$ or otherwise the left-hand side equals ∞ . This concludes the proof as ε was chosen arbitrarily. □

The following two statements adapt Theorem 2.2 in [2] to our slightly more general setting.

Lemma 5.2. *Assume (H1) and (H2). Let h be l.s.c. and bounded from below. Then $v_0 \leq v_*$.*

Proof. Let $(t_0, x_0) \in [0, T] \times \Omega$. It suffices to consider the case $v_*(t_0, x_0) < \infty$. Let $(t_n, x_n)_n$ be a sequence in $[0, T] \times \Omega$ that converges to (t_0, x_0) in \mathbf{d}_∞ and that satisfies $v_*(t_0, x_0) = \underline{\lim}_n v_n(t_n, x_n)$. Let (a_n) be a sequence in \mathcal{L}_b such that each a_n belongs to $\mathcal{L}_b^{t_n}$ and is an n^{-1} -minimizer of (SOC $_n$) with initial data (t_n, x_n) .

Then there exists a subsequence $(\theta_k)_k := (t_{n_k}, x_{n_k}, n_k)_k$ of $(t_n, x_n, n)_n$ with

$$v_*(t_0, x_0) = \lim_k \mathbb{E}_{\theta_k} \left[\int_{t_{n_k}}^T \ell(t, a_{n_k}(t)) dt + h(X + A^{a_{n_k}}) \right]$$

and, for all $k \in \mathbb{N}$,

$$v_*(t_0, x_0) - 1 \leq \mathbb{E}_{\theta_k} \int_{t_{n_k}}^T \ell(t, a_{n_k}(t)) dt + \mathbb{E}_{\theta_k} h(X + A^{a_{n_k}}) \leq v_*(t_0, x_0) + 1.$$

Since ℓ and h are bounded from below, we can assume that

$$v^1 = \lim_k \mathbb{E}_{\theta_k} \int_{t_{n_k}}^T \ell(t, a_{n_k}(t)) dt \text{ and } v^2 = \lim_k \mathbb{E}_{\theta_k} h(X + A^{a_{n_k}})$$

for some $v^1, v^2 \in \mathbb{R}$ with $v_*(t_0, x_0) = v^1 + v^2$ (cf. Theorem 11.1.i and its proof in [10]). As $\sup_k \mathbb{E}_{\theta_k} \int_{t_{n_k}}^T \ell(t, a_{n_k}(t)) dt < \infty$ and $a_{n_k}|_{[0, t_{n_k}]} = 0$ for all $k \in \mathbb{N}$, one can deduce nearly exactly as in the proof of Lemma A.1 in [2] that the continuous processes $A^{a_{n_k}}$ are tight. Let us also note that the sequence (P_{θ_k}) is weakly convergent because for each $\eta \in C_b(\Omega)$,

$$\mathbb{E}_{\theta_k} \eta = \mathbb{E}_{0,0} \eta \left(x_{n_k}(\cdot \wedge t_{n_k}) + \frac{1}{\sqrt{n_k}} (X(t_{n_k} + \cdot)|_{[t_{n_k}, T]} - X(t_{n_k})) \right) \rightarrow \eta(x_0(\cdot \wedge t_0))$$

as $k \rightarrow \infty$, i.e., a sequence of copies of X converges in distribution to the constant $x_0(\cdot \wedge t_0)$. Thus, by Skorohod's representation theorem, there exists a probability space $(\bar{\Omega}, \bar{\mathcal{F}}, \bar{P})$ with a sequence of $\Omega \times \Omega$ -valued random variables $(\bar{A}_{n_k}, \bar{X}_{n_k})_k$ that satisfies $\bar{P} \circ (\bar{A}_{n_k}, \bar{X}_{n_k})^{-1} = P_{\theta_k} \circ (A^{a_{n_k}}, X)^{-1}$ for each $n \in \mathbb{N}$ and that converges (after passing to a subsequence), \bar{P} -a.s., to some random variable (\bar{A}_0, \bar{X}_0) . Next, define a sequence (\bar{a}_k) of \mathbb{R}^d -valued processes on $[0, T] \times \bar{\Omega}$ by $\bar{a}_k(t) := a_{n_k}(t, \bar{X}_{n_k})$. Again as in the proof of Lemma A.1 in [2], one can deduce that (\bar{a}_k) is equiabsolutely integrable and thus has a subsequential weak limit in $L^1([0, T] \times \bar{\Omega}, dt \otimes d\bar{P}; \mathbb{R}^d)$ that we denote by \bar{a}_0 and that satisfies, by Lemma 5.1,

$$\mathbb{E}^{\bar{P}} \int_{t_0}^T \ell(t, \bar{a}_0(t)) dt \leq \varliminf_i \mathbb{E}^{\bar{P}} \int_{t_{n_{k_i}}}^T \ell(t, \bar{a}_{k_i}(t)) dt = \varliminf_i \mathbb{E}_{\theta_{k_i}} \int_{t_{n_{k_i}}}^T \ell(t, a_{n_{k_i}}(t)) dt$$

as well as $\bar{A}_0(t) = \int_0^t \bar{a}_0(s) ds$, \bar{P} -a.s., for every $t \in [0, T]$. Moreover, $\bar{A}_0|_{[0, t_0]} = 0$, \bar{P} -a.s. Hence, together with h being l.s.c. and $\bar{X}_0 = x_0(\cdot \wedge t_0)$, $\bar{\mathbb{P}}$ -a.s., we have

$$(5.1) \quad v_*(t_0, x_0) = v^1 + v^2 \geq \mathbb{E}^{\bar{P}} \left[\int_{t_0}^T \ell(t, \bar{a}_0(t)) dt + h(x_0(\cdot \wedge t_0) + \bar{A}_0) \right].$$

To conclude the proof, it suffices to note that there exists some $x \in \mathcal{X}^{1,1}(t_0, x_0)$ such that the right-hand side (5.1) is greater than or equal to $\int_{t_0}^T \ell(t, x'(t)) dt + h(x)$ (cf. also Remark 2.6 of [27]). \square

Lemma 5.3. *Assume (H1) and (H2). Let h be u.s.c. and bounded. Then $v^* \leq v_0$.*

Proof. It suffices to follow the arguments of the first paragraph of the proof of Theorem 2.2 in [2] and make the obvious adjustments. For the convenience of the reader, we quickly go over it. Fix $(t_0, x_0) \in [0, T] \times \Omega$ and $a \in L^\infty(0, T; \mathbb{R}^d)$ with $\int_0^T \ell(t, a(t)) dt < \infty$. Given $N \in \mathbb{N}$, define $a^N(t) := a(t)$ for $|a(t)| \leq N$ and

$a^N(t) := (N/|a(t)|)a(t)$ for $|a(t)| > N$. Next, consider a sequence (t_n, x_n) that converges to (t_0, x_0) in $[0, T] \times \Omega$ and satisfies $v^*(t_0, x_0) = \lim_n v_n(t_n, x_n)$. Then

$$\begin{aligned} v^*(t_0, x_0) &\leq \overline{\lim}_n \mathbb{E}_{t_n, x_n, n} \left[\int_{t_n}^T \ell(t, a^N(t)) dt + h(X + A^{a^N} - A^{a^N}(t_n)) \right] \\ &\leq \overline{\lim}_n \int_{t_n}^T \ell(t, a^N(t)) dt \\ &\quad + \overline{\lim}_n \mathbb{E}_{0,0} h \left(x_{n_k}(\cdot \wedge t_{n_k}) + \frac{1}{\sqrt{n_k}} (X(t_{n_k} + \cdot)|_{[t_{n_k}, T]} - X(t_{n_k})) + A^{a^N} - A^{a^N}(t_n) \right) \\ &\leq \int_{t_0}^T \ell(t, a^N(t)) dt + h(x_0(\cdot \wedge t_0) + A^{a^N} - A^{a^N}(t_0)) \end{aligned}$$

as h is u.s.c. as well as bounded and $\int_0^T \ell(t, a^N(t)) dt < \infty$, which follows from $\int_0^T \ell(t, a(t)) dt < \infty$, (H2), and the convexity of $\ell(t, \cdot)$ (cf. the first displayed equation after (38) in [2]). Finally, letting $N \rightarrow \infty$ concludes the proof as in [2]. \square

Remark 5.4 (No Lavrentiev phenomenon). Assume (H1) and (H2). Let h be continuous and bounded. Then

$$v_0(t_0, x_0) = \inf_{x \in \mathcal{X}^{1, \infty}(t_0, x_0)} \left[\int_{t_0}^T \ell(t, x'(t)) dt + h(x) \right],$$

where

$$\mathcal{X}^{1, \infty}(t_0, x_0) := \{x \in \Omega : x|_{[0, t_0]} = x_0|_{[0, t_0]} \text{ and } x|_{[t_0, T]} \in W^{1, \infty}(t_0, T; \mathbb{R}^d)\}.$$

This result follows from the proofs of Lemmata 5.2 and 5.3 but with each $P_{t_0, x_0, n}$, $n \in \mathbb{N}$, replaced by the unique probability measure under which $X = x(\cdot \wedge t_0)$ a.s. For a more direct proof, it suffices to slightly modify the proof of Proposition 4.1 in [8] (this result is due to [14]), where the case $h \equiv 0$ is treated.

Proof of the first conclusion of Theorem 4.1. By Lemmata 5.2 and 5.3, $v_0 \leq v_* \leq v^* \leq v_0$. Thus (v_n) converges to v_0 uniformly on compacta and v_0 is continuous (cf. Lemmata V.1.5 and V.1.9 of [3] and keep Remark 2.1 in mind). \square

6. CONNECTIONS TO BSDEs

We present parts of the theory of (convex) superquadratic BSDEs from [15] that are relevant for our work. Fix $(t_0, x_0) \in [0, T] \times \Omega$, $t_1 \in [t_0, T]$, $n \in \mathbb{N}$, and a \mathcal{F}_{t_1} -measurable random variable $\xi : \Omega \rightarrow \mathbb{R} \cup \{\infty\}$. Consider the BSDE

$$(6.1) \quad \begin{aligned} dY(t) &= - \sup_{a \in \mathbb{R}^d} [a \cdot Z(t) - \ell(t, a)] dt + Z(t) dX(t), \text{ on } [t_0, t_1], P_{t_0, x_0, n}\text{-a.s.}, \\ Y(t_1) &= \xi. \end{aligned}$$

Definition 6.1. A pair (Y, Z) is a *supersolution* of (6.1) if

- $Y : [t_0, t_1] \times \Omega \rightarrow \mathbb{R}$ is a càdlàg and $\mathbb{F}^{t_0, x_0, n}$ -adapted process,
- $Z : [t_0, t_1] \times \Omega \rightarrow \mathbb{R}^d$ is an $\mathbb{F}^{t_0, x_0, n}$ -predictable process with

$$\mathbb{E}_{t_0, x_0, n} \int_{t_0}^{t_1} |Z(t)|^2 dt < \infty,$$

- $(t, \omega) \mapsto \left[\int_{t_0}^t Z(r) dX(r) \right] (\omega)$, $[t_0, t_1] \times \Omega \rightarrow \mathbb{R}$, is an $(\mathbb{F}^{t_0, x_0, n}, \mathbb{P}_{t_0, x_0, n})$ -supermartingale, and,
- for every $t, s \in [t_0, t_1]$ with $t \leq s$, we have, $P_{t_0, x_0, n}$ -a.s.,

$$Y(s) \geq Y(t) + \int_t^s \left\{ - \sup_{a \in \mathbb{R}^d} [a \cdot Z(r) - \ell(r, a)] \right\} dr + \int_t^s Z(r) dX(r),$$

$$Y(t_1) \geq \xi.$$

A pair (Y, Z) is a *minimal supersolution* of (6.1) if it is a supersolution of (6.1) and, for every supersolution (\tilde{Y}, \tilde{Z}) of (6.1), we have $Y \leq \tilde{Y}$, $dt \otimes dP_{t_0, x_0, n}$ -a.e.

Proposition 6.2. *Assume (H1) and (H2). Let ξ be bounded. Then (6.1) has a unique minimal supersolution (Y, Z) and we write*

$$(6.2) \quad \mathcal{E}_{t, t_1}^{t_0, x_0, n}(\xi) := Y_t, \quad t \in [t_0, t_1].$$

Proof. By the proof of Theorem 3.4 in [16] and by Theorem A.1 in [16], the set of supersolutions of (6.1) is non-empty. Thus we can apply Theorem 4.17 in [15], which concludes the proof. \square

From now on, let $T = 1$ in this section for the sake of simplicity. Fix $(t_0, x_0) \in [0, T] \times \Omega$. Recall that $\mathcal{L}_b^{t_0} = \{a \in \mathcal{L}_b : a|_{[0, t_0]} = 0\}$. Given $a \in \mathcal{L}_b$, define $A^a : [0, T] \times \Omega \rightarrow \mathbb{R}^d$ by $A^a(t, \omega) := \int_0^t a(s, \omega) ds$. We also identify a control $a \in \mathcal{L}_b$ with the function $\Omega \rightarrow L^0(\Omega, \mathcal{F}_T)$, $\omega \mapsto a(\omega)$, where $[a(\omega)](t) = a(t, \omega)$. Recall that $X(t, \omega) = \omega(t)$, $(t, \omega) \in [0, T] \times \Omega$, and that $\mathbb{E}_{t_0, x_0} = \mathbb{E}^{P_{t_0, x_0}}$.

For the next statement, we borrow the following constructions from [2]. Given $t \in [0, 1]$, $\omega_1, \omega_2 \in \Omega$ with $\omega_2(0) = 0$, a path $\omega_1 \odot_t \omega_2 \in \Omega$ is defined by

$$(6.3) \quad (\omega_1 \odot_t \omega_2)(s) := \omega_1(s \wedge t) + \sqrt{1-t} \omega_2 \left(\frac{s-t}{1-t} \right) \mathbf{1}_{[t, 1]}(s)$$

(note that in [2] \otimes is used instead of \odot) and a path $\omega_1^{(t)} \in \Omega$ is defined by

$$\omega_1^{(t)}(s) := \frac{1}{\sqrt{1-t}} [\omega_1(t + s(1-t)) - \omega_1(t)].$$

Moreover, we use the function $\ell^{(t_0)} : [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\infty\}$ defined by

$$(6.4) \quad \ell^{(t_0)}(t, a) := (1-t_0) \ell \left(t_0 + t(1-t_0), \frac{a}{\sqrt{1-t_0}} \right).$$

Also set $(\omega_1 \odot_1 \omega_2) := \omega_1$, $\omega_1^{(1)} \equiv 0$, and $\ell^{(1)} \equiv 0$.

Lemma 6.3. *Assume (H1) and (H2). Let h be bounded. Then*

$$(6.5) \quad \sup_{a \in \mathcal{L}_b^{t_0}} \mathbb{E}_{t_0, x_0} \left[(-h)(X + A^{a(X)}) - \int_{t_0}^1 \ell(t, a(t, X)) dt \right]$$

$$= \sup_{\check{a} \in \mathcal{L}_b} \mathbb{E}_{0, 0} \left[(-h)(x_0 \odot_{t_0} [X + A^{\check{a}(X)}]) - \int_0^1 \ell^{(t_0)}(t, \check{a}(t, X)) dt \right].$$

The calculations in our proof are essentially the same as those in the proof of Lemma 5.1 in [2]. Although of similar nature, the corresponding statements are different. This permits a more elementary proof in our case.

Proof of Lemma 6.3. One should keep in mind, that $X = x_0 \odot_{t_0} X^{(t_0)}$, P_{t_0, x_0} -a.s. We only consider the case $t_0 < 1$ as otherwise (6.5) is clearly satisfied.

Step 1. Fix $a \in \mathcal{L}_b^{t_0}$. Note that

$$\begin{aligned} \mathbb{E}_{t_0, x_0} \left[\int_{t_0}^1 \ell(t, a(t_0, x_0 \odot_{t_0} X^{(t_0)})) dt \right] &= \mathbb{E}_{0,0} \left[\int_{t_0}^1 \ell(t, a(t_0, x_0 \odot_{t_0} X)) dt \right] \\ &= \mathbb{E}_{0,0} \left[(1-t_0) \int_0^1 \ell(t_0 + t(1-t_0), a(t_0 + t(1-t_0), x_0 \odot_{t_0} X)) dt \right] \\ &= \mathbb{E}_{0,0} \left[\int_0^1 \ell^{(t_0)}(t, \check{a}(t, X)) dt \right], \end{aligned}$$

where $\check{a} \in \mathcal{L}_b$ is defined by

$$\check{a}(t, \omega) := \sqrt{1-t_0} a(t_0 + t(1-t_0), x_0 \odot_{t_0} \omega).$$

Also note that

$$\begin{aligned} \mathbb{E}_{t_0, x_0} \left[(-h)(x_0 \odot_{t_0} [X + A^{a(x_0 \odot_{t_0} X^{(t_0)})}]^{(t_0)}) \right] \\ &= \mathbb{E}_{0,0} \left[(-h)(x_0 \odot_{t_0} [X + (A^{a(x_0 \odot_{t_0} X)})]^{(t_0)}) \right] \\ &= \mathbb{E}_{0,0} \left[(-h)(x_0 \odot_{t_0} [X + A^{\check{a}}]) \right] \end{aligned}$$

because, for every $t \in [0, 1]$,

$$(A^{a(x_0 \odot_{t_0} X)})^{(t_0)} = \frac{1}{\sqrt{1-t_0}} \int_{t_0}^{t_0+t(1-t_0)} a(s, x_0 \odot_{t_0} X) ds = \int_0^t \check{a}(s, X) ds.$$

Consequently, the left-hand side of (6.5) is less than or equal to the right-hand side of (6.5).

Step 2. Fix $\check{a} \in \mathcal{L}_b$. Define $a \in \mathcal{L}_b^{t_0}$ by

$$a(t, \omega) := \frac{1}{\sqrt{1-t_0}} \check{a} \left(\frac{t-t_0}{1-t_0}, \omega^{(t_0)} \right) \mathbf{1}_{[t_0, 1] \times \{x_0|_{[0, t_0]}\}}(t, \omega|_{[0, t_0]}).$$

Going over the calculations in Step 1 backward, one can deduce that the right-hand side of (6.5) is less than or equal to the left-hand side of (6.5). \square

The following statement together with Theorem 4.1 provide a non-Markovian non-linear Feynman-Kac formula that connects maximal Dini subsolutions of path-dependent PDEs with minimal supersolutions of convex superquadratic BSDEs, for which we use the notation (6.2).

Theorem 6.4. *Assume (H1) and (H2). Let ℓ be continuous and finite-valued. Let h be bounded. Fix $(t_0, x_0) \in [0, T] \times \Omega$ and $n \in \mathbb{N}$. Then $v_n(t_0, x_0) = -\mathbb{E}_{t_0, x_0, n}[\mathcal{E}_{t_0, T}^{t_0, x_0, n}(-h)]$. Moreover, for $P_{t_0, x_0, n}$ -a.e. $\omega \in \Omega$ and every $t \in [t_0, T]$, we have $v_n(t, \omega) = -\mathcal{E}_{t, T}^{t_0, x_0, n}(-h)(\omega)$.*

Proof. We prove the theorem only for the case $n = 1$ and $T = 1$. Note that, on the one hand, $-v_1(t_0, x_0)$ equals, by definition, the left-hand side of (6.5). On the other hand, by (BBD) in [2], the right-hand side of (6.5) equals $\rho^{\ell^{(t_0)}}[(-h)(x_0 \odot_{t_0} X)]$, where ρ is defined in Section 2 of [2]. Moreover, by a straight-forward adjustment of the proof of Lemma 5.1 in [2], $\rho^{\ell^{(t)}}[(-h)(\omega \odot_t X)] = Y^{t_0, x_0, 1}(t, \omega)$ for every $t \in [t_0, T]$ and P_{t_0, x_0} -a.e. $\omega \in \Omega$. In this context, note that Lemma 5.1 and Lemma A.2, both in [2], formally require continuity of h , which, however, is not necessary (cf the

corresponding material in [16]). Using Lemma 6.3, we can deduce that $v_1(t, \cdot) = -Y^{t_0, x_0, 1}(t)$, P_{t_0, x_0} -a.s., for every $t \in [t_0, T]$. \square

7. THE SECOND ORDER HJB EQUATIONS

Lemma 7.1. *Let $n \in \mathbb{N}$. Assume (H1). Let $\ell = \ell(t, a)$ be continuous in t . A bounded u.s.c. function $u : [0, T] \times \Omega \rightarrow \mathbb{R}$ is a Dini subsolution of (TVP n) if and only if $u(T, \cdot) \leq h$ and, for every $(t_0, x_0) \in [0, T] \times \Omega$, $t \in (t_0, T]$, and $a \in \mathbb{R}^d$,*

$$(7.1) \quad u(t_0, x_0) \leq \mathbb{E}_{t_0, x_0, n} \left[\int_{t_0}^t \ell(s, a) ds + u(t, X + A^a - A^a(t_0)) \right].$$

Proof. We adapt the proof of Theorem V.3 in [43], which is situated in a Markovian context with bounded control spaces, to our non-Markovian setting with the unbounded control space \mathbb{R}^d .

(i) Let u be a Dini subsolution of (TVP n). Fix $(t_0, x_0) \in [0, T] \times \Omega$. Put $\mathbb{E} = \mathbb{E}_{t_0, x_0, n}$ and $P = P_{t_0, x_0, n}$. Assume that there are $t_1 \in (t_0, T]$ and $a \in \mathbb{R}^d$, and $\varepsilon > 0$ such that

$$(7.2) \quad \mathbb{E} \left[u(t_1, X + A^a - A^a(t_0)) - u(t_0, x_0) + \int_{t_0}^{t_1} \ell(s, a) ds \right] < -\varepsilon.$$

Consider the set S of all \mathbb{F} -stopping times τ that satisfy $t_0 \leq \tau \leq t_1$, P -a.s., and

$$(7.3) \quad \mathbb{E} \left[u(\tau, X + A^a - A^a(t_0)) - u(t_0, x_0) + \int_{t_0}^{\tau} \ell(s, a) ds \right] \geq \frac{\mathbb{E}[\tau] - t_0}{t_1 - t_0} \cdot (-\varepsilon).$$

We equip S with an order \preceq defined by $\tau_1 \preceq \tau_2$ if and only if $\tau_1 \leq \tau_2$, P -a.s. Now, let \tilde{S} be a totally ordered non-empty subset of S . Note that there exists a sequence (τ_k) in \tilde{S} such that $\mathbb{E}[\tau_k] \uparrow \sup_{\tau \in \tilde{S}} \mathbb{E}[\tau]$ as $k \rightarrow \infty$. Since \tilde{S} is totally ordered and \mathbb{E} is linear, (τ_k) is increasing and converges to $\tilde{\tau} := \sup_k \tau_k$. We have $\tilde{\tau} \in S$ because

$$\begin{aligned} & \mathbb{E} \left[u(\tilde{\tau}, X + A^a - A^a(t_0)) + \int_{t_0}^{\tilde{\tau}} \ell(s, a) ds \right] \\ & \geq \overline{\lim}_k \mathbb{E} \left[u(\tau_k, X + A^a - A^a(t_0)) + \int_{t_0}^{\tau_k} \ell(s, a) ds \right] \geq u(t_0, x_0) + \frac{\mathbb{E}[\tilde{\tau}] - t_0}{t_1 - t_0} \cdot (-\varepsilon) \end{aligned}$$

thanks to (7.3) as well as to u being u.s.c and bounded. Hence, by Zorn's lemma, S has a maximal element τ_0 . We show that $\tau_0 = t_1$, P -a.s. To this end, assume that $P(\tau_0 < t_1) > 0$. Define a set-valued function M on $[t_0, t_1] \times \Omega$ by

$$M(t, \omega) := \left\{ \delta \in [0, T - t_0] : t + \delta \leq T \text{ and } \mathbb{E}_{t, \omega, n} \left[u(t + \delta, X + A^a - A^a(t)) - u(t, \omega) + \int_t^{t+\delta} \ell(s, a) ds \right] \geq \frac{\delta}{t_1 - t_0} \cdot (-\varepsilon) \right\}.$$

The sets $M(t, \omega)$ are non-empty because they contain 0 and they are compact because, for any sequence (δ_k) in $[0, T - t]$ that converges to some δ , we have

$$\begin{aligned} & \mathbb{E}_{t, \omega, n} \left[u(t + \delta, X + A^a - A^a(t)) + \int_t^{t+\delta} \ell(s, a) ds \right] \\ & \geq \overline{\lim}_k \mathbb{E}_{t, \omega, n} \left[u(t + \delta_k, X + A^a - A^a(t)) + \int_t^{t+\delta_k} \ell(s, a) ds \right] \end{aligned}$$

as u is u.s.c. and bounded. Moreover, for every sequence $(s_k, \omega_k)_k$ in $[t_0, t_1] \times \Omega$ that converges to some (t, ω) with respect to the metric $((\tilde{s}_1, \tilde{\omega}_1), (\tilde{s}_2, \tilde{\omega}_2)) \mapsto |\tilde{s}_1 - \tilde{s}_2| + \|\tilde{\omega}_1 - \tilde{\omega}_2\|_\infty$ and hence also with respect to \mathbf{d}_∞ , every sequence (δ_k) in $[0, T - t_0]$ with $\delta_k \in M(s_k, \omega_k)$ has a subsequential limit that belongs to $M(t, \omega)$ because, for every subsequential limit δ of (δ_k) (there is at least one), we have, by possibly passing to a subsequence,

$$\begin{aligned}
& \mathbb{E}_{t, \omega, n} \left[u(t + \delta, X + A^a - A^a(t)) + \int_t^{t+\delta} \ell(s, a) ds \right] \\
&= \mathbb{E}_{0, 0, n} \left[u \left((t + \delta, \omega(\cdot \wedge t)) + \frac{1}{\sqrt{n}} (X(t + \cdot)|_{[t, T]} - X(t) + A^a - A^a(t)) \right) \right. \\
&\quad \left. + \int_t^{t+\delta} \ell(s, a) ds \right] \\
&\geq \overline{\lim}_k \mathbb{E}_{0, 0, n} \left[u \left((s_k + \delta_k, \omega_k(\cdot \wedge s_k)) + \frac{1}{\sqrt{n}} (X(s_k + \cdot)|_{[s_k, T]} - X(s_k) \right. \right. \\
&\quad \left. \left. + A^a - A^a(s_k)) \right) + \int_{s_k}^{s_k + \delta_k} \ell(s, a) ds \right] \\
&= \overline{\lim}_k \mathbb{E}_{s_k, \omega_k, n} \left[u(s_k + \delta_k, X + A^a - A^a(s_k)) + \int_{s_k}^{s_k + \delta_k} \ell(s, a) ds \right].
\end{aligned}$$

Hence, M considered as a map from $[t_0, t_1] \times \Omega$ equipped here with the metric $((s_1, \omega_1), (s_2, \omega_2)) \mapsto |s_1 - s_2| + \|\omega_1 - \omega_2\|_\infty$ into the set of all compact non-empty subsets of $[0, T - t_0]$ is u.s.c. (Theorem 2.2 on p. 31 in [30]) and thus measurable (Theorem 2.1 on p. 29 in [30]; see p. 41 in [30] for the definition of measurability). Consequently, by Theorem 3.13 on p. 49 in [30], there exists a $\mathcal{B}([t_0, t_1]) \otimes \mathcal{F}_T$ -measurable selector $\delta(\cdot, \cdot)$ of M that satisfies $|T - t_0 - \delta(t, \omega)| = \text{dist}(T - t_0, M(t, \omega))$ for every $(t, \omega) \in [t_0, t_1] \times \Omega$. Therefore, the map $\omega \mapsto \delta[\omega] := \delta(\tau_0(\omega), \omega(\cdot \wedge \tau_0(\omega)) + A^a(\tau_0(\omega)) - A^a(t_0))$ is \mathcal{F}_{τ_0} -measurable. Since, by (3.3) and the continuity of $\ell(\cdot, a)$, every set $M(t, \omega)$ contains a strictly positive element, we have $|T - t_0 - \delta(t, \omega)| < |T - t_0|$ and hence $\delta(t, \omega) > 0$. Thus, by Lemma V.3 in [43], $\tilde{\tau}_0 := (\tau_0 + \delta[\cdot]) \wedge t_1$ is an \mathbb{F} -stopping time with $\tilde{\tau}_0 > \tau_0$ on $\{\tau_0 < t_1\}$. Finally,

$$\begin{aligned}
& \mathbb{E} \left[u(\tilde{\tau}_0, X + A^a - A^a(t_0)) + \int_{t_0}^{\tilde{\tau}_0} \ell(s, a) ds \right] \\
&= \int_\Omega \left(\mathbb{E} \left[u(\tilde{\tau}_0, X + A^a - A^a(t_0)) + \int_{\tau_0}^{\tilde{\tau}_0} \ell(s, a) ds \middle| \mathcal{F}_{\tau_0} \right] (\omega) \right. \\
&\quad \left. + \int_{t_0}^{\tau_0(\omega)} \ell(s, a) ds \right) P(d\omega) \\
&= \int_\Omega \left(\mathbb{E}_{\tau_0(\omega), \omega, n} \left[u(\tilde{\tau}_0(\omega), X + A^a - A^a(t_0)) + \int_{\tau_0(\omega)}^{\tilde{\tau}_0(\omega)} \ell(s, a) ds \right] \right. \\
&\quad \left. + \int_{t_0}^{\tau_0(\omega)} \ell(s, a) ds \right) P(d\omega)
\end{aligned}$$

$$\begin{aligned}
&= \int_{\Omega} \left(\mathbb{E}_{\tau_0(\omega), \omega + A^a(\tau_0(\omega)) - A^a(t_0), n} \left[u(\tilde{\tau}_0(\omega), X + A^a - A^a(\tau_0(\omega))) \right. \right. \\
&\quad \left. \left. + \int_{\tau_0(\omega)}^{\tilde{\tau}_0(\omega)} \ell(s, a) ds \right] + \int_{t_0}^{\tau_0(\omega)} \ell(s, a) ds \right) P(d\omega) \\
&\geq \int_{\Omega} \left(u(\tau_0(\omega), \omega + A^a(\tau_0(\omega)) - A^a(t_0)) + \frac{\delta[\omega]}{t_1 - t_0} \cdot (-\varepsilon) + \int_{t_0}^{\tau_0(\omega)} \ell(s, a) \right) P(d\omega) \\
&\geq u(t_0, x_0) + \frac{\mathbb{E}[\delta[\cdot] + \tau_0] - t_0}{t_1 - t_0} \cdot (-\varepsilon),
\end{aligned}$$

i.e., $\tilde{\tau}_0 \in S$. Since $\tilde{\tau}_0 \geq \tau_0$ and $P(\{\tilde{\tau}_0 > \tau_0\} \cap \{\tau_0 < t_1\}) > 0$, we have $\tau_0 \preceq \tilde{\tau}_0$ but not $\tau_0 = \tilde{\tau}_0$, P -a.s., i.e., we have a contradiction to τ_0 being the maximal element in S . Hence, $\tau_0 = t_1$, P -a.s. However, this in turn contradicts (7.2), which concludes the proof of this direction.

(ii) Showing the remaining direction is straight-forward. \square

Theorem 7.2. *Assume (H1) and (H2). Fix $n \in \mathbb{N}$. Let h be u.s.c. and bounded. Let ℓ be continuous and finite-valued. Then v_n is the unique bounded maximal Dini subsolution of (TVP n). Moreover, each bounded Dini subsolution of (TVP n) is dominated from above by v_n even if we dispense with the assumption that h is u.s.c.*

Proof. (i) Existence and regularity: First, we show that v_n is bounded and u.s.c. Boundedness of h , (H1), and (H2) yield boundedness of v_n . Next, fix $(t_0, x_0) \in [0, T] \times \Omega$. Consider a sequence $(t_k, x_k)_{k \geq 1}$ in $[0, T] \times \Omega$ that converges to (t_0, x_0) and satisfies $\lim_k v_n(t_k, x_k) = \overline{\lim}_{(t,x) \rightarrow (t_0, x_0)} v_n(t, x)$. We will use \odot defined by (6.3) and the notation (6.4). For this reason, we assume that $T = 1$ and $n = 1$ but this assumption is not restrictive. Note that $(t, x) \mapsto (x \odot_t \omega)$, $[0, T] \times \Omega \rightarrow \Omega$, is continuous for every $\omega \in \Omega$ with $\omega(0) = 0$. Thus Lemma 6.3 yields

$$\begin{aligned}
\lim_k v_n(t_k, x_k) &\leq \overline{\lim}_k \mathbb{E}_{0,0,n} \left[h(x_k \odot_{t_k} [X + A^a]) + \int_0^T \ell^{(t_k)}(t, a(t)) \wedge (C + 1) dt \right] \\
&\leq \mathbb{E}_{0,0,n} \left[h(x_0 \odot_{t_0} [X + A^a]) + \int_0^T \ell^{(t_0)}(t, a(t)) \wedge (C + 1) dt \right] \quad \text{for every } a \in \mathcal{L}_b
\end{aligned}$$

with $\sup_{k \geq 0} \mathbb{E}_{0,0,n} \int_0^T \ell^{(t_k)}(t, a(t)) dt < C + 1$, where C is an upper bound of v_n . Again, by Lemma 6.3, taking the infimum of the right-hand side over all such $a \in \mathcal{L}_b$ yields that v_n is u.s.c.

Next, we establish the subsolution property via the BSDE connection in Theorem 6.4. To this end, we use the notation (6.2) and fix $(t_0, x_0) \in [0, T] \times \Omega$, $t \in [t_0, T]$, and $a \in \mathbb{R}^d$. Note first that, by Proposition 3.6 (1) in [15],

$$(7.4) \quad \mathcal{E}_{t_0, T}^{t_0, x_0, n}(-h) = \mathcal{E}_{t_0, t}^{t_0, x_0, n} \left(\mathcal{E}_{t, T}^{t_0, x_0, n}(-h) \right).$$

Since v_n is bounded and, by Theorem 6.4, $v_n(t, \cdot) = -\mathcal{E}_{t, T}^{t_0, x_0, n}(-h)$, $P_{t_0, x_0, n}$ -a.s., we can apply Theorem 3.4 in [16] (see also Lemma A.2 in [2]) together with (3) in [2] to deduce that, with $Q_{t_0, x_0, n}^a := \mathbb{P}_{t_0, x_0, n} \circ (X + A^a - A^a(t_0))^{-1}$,

$$\mathbb{E}_{t_0, x_0, n} \left[\mathcal{E}_{t_0, t}^{t_0, x_0, n} \left(\mathcal{E}_{t, T}^{t_0, x_0, n}(-h) \right) \right] \geq \mathbb{E}^{Q_{t_0, x_0, n}^a} \left[\mathcal{E}_{t, T}^{t_0, x_0, n}(-h) - \int_{t_0}^t \ell(s, a) ds \right]$$

$$= \mathbb{E}_{t_0, x_0, n} \left[\mathcal{E}_{t, T}^{t_0, x_0, n}(-h)(X + A^a - A^a(t_0)) - \int_{t_0}^t \ell(s, a) ds \right].$$

Thus, by (7.4) and Theorem 6.4, v_n satisfies (7.1) in place of u . Hence, by Lemma 7.1, v_n is a Dini subsolution of (TVP n).

(ii) Uniqueness: Let u be a bounded Dini subsolution of (TVP n). Without loss of generality, let $n = 1$. Fix $(t_0, x_0) \in [0, T] \times \Omega$ and $\varepsilon > 0$. It suffices to show that $u(t_0, x_0) \leq v(t_0, x_0) + \varepsilon$. By (a slight modification) of (36) and (BBD), both in [2], there exists an $a \in \mathcal{L}_b^{t_0}$ with $a(t, \omega) = \sum_{i=1}^m a_i(\omega) \cdot \mathbf{1}_{[s_{i-1}, s_i)}(t)$, where $t_0 = s_0 < s_1 < \dots < s_m = T$ and each $a_i : \Omega \rightarrow \mathbb{R}^d$ is $\mathcal{F}_{s_{i-1}}$ -measurable, such that $\tilde{\mathbb{E}}_{t_0, x_0} \left[\int_{t_0}^T \ell(t, a(t)) dt + h(X) \right] \leq v_n(t_0, x_0) + \varepsilon$. Here, for any $(s, x) \in [t_0, T] \times \Omega$, $\tilde{P}_{s, x} := P_{s, x} \circ (\tilde{X}^{s, x})^{-1}$, $\tilde{\mathbb{E}} := \mathbb{E}^{\tilde{P}_{s, x}}$, and $\tilde{X}^{s, x} : [0, T] \times \Omega \rightarrow \mathbb{R}^d$ is the unique solution of $\tilde{X}^{s, x}(t) = \tilde{X}^{s, x}(s) + \int_s^t a(r, \tilde{X}^{s, x}) dr + X(t) - X(s)$, $t \in [s, T]$, with initial condition $\tilde{X}^{s, x}|_{[0, s]} = x|_{[0, s]}$. Next, note that $a_1 \circ \tilde{X}^{t_0, x_0} = a_1 \circ x_0 =: \tilde{a}_1 \in \mathbb{R}^d$ and $X + A^{\tilde{a}_1} - A^{\tilde{a}_1}(t_0) = \tilde{X}^{t_0, x_0}$ on $[0, s_1]$, P_{t_0, x_0} -a.s. Thus, by Lemma 7.1,

$$\begin{aligned} u(t_0, x_0) &\leq \mathbb{E}_{t_0, x_0} \left[\int_{t_0}^{s_1} \ell(t, \tilde{a}_1) dt + u(s_1, \tilde{X}^{t_0, x_0}) \right] \\ &= \tilde{\mathbb{E}}_{t_0, x_0} \left[\int_{t_0}^{s_1} \ell(t, a(t)) dt + u(s_1, X) \right]. \end{aligned}$$

Also note that, for P_{t_0, x_0} -a.e. $\omega \in \Omega$, we have, with $x_1 = \tilde{X}^{t_0, x_0}(\omega)$, $a_2 \circ \tilde{X}^{s_1, x_1} = a_2 \circ x_1 \in \mathbb{R}^d$ and $X + A^{a_2 \circ x_1} - A^{a_2 \circ x_1}(s_1) = \tilde{X}^{s_1, x_1}$ on $[0, s_2]$, P_{s_1, x_1} -a.s., and thus, by Lemma 7.1,

$$\begin{aligned} u(s_1, x_1) &\leq \mathbb{E}_{s_1, x_1} \left[\int_{s_1}^{s_2} \ell(t, \tilde{a}_2) dt + u(s_2, \tilde{X}^{s_1, x_1}) \right] \\ &= \tilde{\mathbb{E}}_{s_1, x_1} \left[\int_{s_1}^{s_2} \ell(t, a(t)) dt + u(s_2, X) \right]. \end{aligned}$$

Therefore

$$\begin{aligned} \tilde{\mathbb{E}}_{t_0, x_0} [u(s_1, X)] &\leq \int_{\Omega} \tilde{\mathbb{E}}_{s_1, x_1} \left[\int_{s_1}^{s_2} \ell(t, a(t)) dt + u(s_2, X) \right] \Big|_{x_1 = \omega} \tilde{P}_{t_0, x_0}(d\omega) \\ &= \tilde{\mathbb{E}}_{t_0, x_0} \left(\tilde{\mathbb{E}}_{t_0, x_0} \left[\int_{s_1}^{s_2} \ell(t, a(t)) dt + u(s_2, X) \right] \Big| \mathcal{F}_{s_1} \right) \\ &= \tilde{\mathbb{E}}_{t_0, x_0} \left[\int_{s_1}^{s_2} \ell(t, a(t)) dt + u(s_2, X) \right]. \end{aligned}$$

Repeating this procedure and noting that $u(T, \cdot) \leq h$ yields

$$u(t_0, x_0) \leq \tilde{\mathbb{E}}_{t_0, x_0} \left[\int_{t_0}^T \ell(t, a(t)) dt + h(X) \right] \leq v_n(t_0, x_0) + \varepsilon,$$

which concludes the proof. \square

8. THE FIRST ORDER HJB EQUATION

Lemma 8.1. *Assume (H1). An l.s.c. function $u : [0, T] \times \Omega \rightarrow \mathbb{R} \cup \{\infty\}$ bounded from below is a minimax supersolution of (TVP) if and only if $u(T, \cdot) \geq h$ and, for*

every $(t_0, x_0) \in [0, T] \times \Omega$ and $t \in (t_0, T]$, there exists an $x \in \mathcal{X}^{1,1}(t_0, x_0)$ such that

$$(8.1) \quad u(t_0, x_0) \geq \int_{t_0}^t \ell(s, x'(s)) ds + u(t, x).$$

Proof. (i) Let u be a minimax supersolution of (TVP). Assume that there exist a $(t_0, x_0) \in \text{dom}(u)$ with $t_0 < T$ and a $t_1 \in (t_0, T]$ such that, for all $x \in \mathcal{X}^{1,1}(t_0, x_0)$, $u(t_1, x) - u(t_0, x_0) + \int_{t_0}^{t_1} \ell(s, x'(s)) ds > 0$. Then there is an $\varepsilon > 0$ such that

$$(8.2) \quad u(t_1, x) - u(t_0, x_0) + \int_{t_0}^{t_1} \ell(s, x'(s)) ds > \varepsilon$$

for all $x \in \mathcal{X}^{1,1}(t_0, x_0)$ because $\inf_{x \in \mathcal{X}^{1,1}(t_0, x_0)} \left[\int_{t_0}^{t_1} \ell(s, x'(s)) ds + u(t_1, x) \right]$ has a minimizer (cf. [10]). Next, consider the (non-void) set S of all $(t, x) \in [t_0, t_1] \times \mathcal{X}^{1,1}(t_0, x_0)$ for which $u(t, x) - u(t_0, x_0) + \int_{t_0}^t \ell(s, x'(s)) ds \leq \frac{t-t_0}{t_1-t_0} \cdot \varepsilon$ holds. Let $\tilde{s} := \sup\{t : (t, x) \in S \text{ for some } x\}$. This supremum is attained. To see this, consider a sequence $(s_n, x_n)_n$ in S with $s_n \rightarrow \tilde{s}$. Since $\sup_n \int_{t_0}^{s_n} \ell(s, x'_n(s)) ds < \infty$ and u is bounded from below, one can show as in Sections 11.1 and 11.2 of [10] that there is a subsequence $(s_{n_k}, x_{n_k})_k$ such that (x_{n_k}) converges to some \tilde{x} in $\mathcal{X}^{1,1}(t_0, x_0)$, i.e., $\|x_{n_k} - \tilde{x}\|_\infty + \|x'_{n_k} - \tilde{x}'\|_{L^1(t_0, T; \mathbb{R}^d)} \rightarrow 0$. Lower semi-continuity of u together with Theorem 10.8.ii in [10] yield

$$\begin{aligned} u(\tilde{s}, \tilde{x}) + \int_{t_0}^{\tilde{s}} \ell(s, \tilde{x}'(s)) ds &\leq \liminf_k u(s_{n_k}, x_{n_k}) + \liminf_k \int_{t_0}^{s_{n_k}} \ell(s, x'_{n_k}(s)) ds \\ &\leq \frac{\tilde{s} - t_0}{t_1 - t_0} \cdot \varepsilon + u(t_0, x_0). \end{aligned}$$

Thus $(\tilde{s}, \tilde{x}) \in \text{dom}(u)$ and, by (8.2), $\tilde{s} < t_1$. Hence, by (3.4), there is a $\delta > 0$ and an $x \in \mathcal{X}^{1,1}(\tilde{s}, \tilde{x})$ such that $u(\tilde{s} + \delta, x) - u(\tilde{s}, \tilde{x}) + \int_{\tilde{s}}^{\tilde{s} + \delta} \ell(s, x'(s)) ds \leq \frac{\delta}{t_1 - t_0} \cdot \varepsilon$ and $\tilde{s} + \delta < t_1$. Thus $(\tilde{s} + \delta, x) \in S$, which is a contradiction to the maximality of \tilde{s} .

(ii) Fix $(t_0, x_0) \in \text{dom}(u)$ with $t_0 < T$. Suppose that, for every $(\tilde{t}_0, \tilde{x}_0) \in [t_0, T] \times \mathcal{X}^{1,1}(t_0, x_0)$ for every $t \in (\tilde{t}_0, T]$, there exists an $x \in \mathcal{X}^{1,1}(\tilde{t}_0, \tilde{x}_0)$ such that (8.1) holds with (t_0, x_0) replaced by $(\tilde{t}_0, \tilde{x}_0)$. Then one can proceed similarly as in the proof of Lemma 3.6 in [4] to show that there exists a sequence (x_n) in $\mathcal{X}^{1,1}(t_0, x_0)$ and an increasing sequence (A_n) of finite subsets of $[t_0, T]$ whose union A is dense in $[t_0, T]$ such that, for each $n \in \mathbb{N}$ and every $t \in A_n$, (8.1) holds with x replaced by x_n . Thus $\sup_{t \in A, n \in \mathbb{N}} \int_{t_0}^t \ell(s, x'_n(s)) ds \leq u(t_0, x_0) + c < \infty$, where $-c$ is a lower bound of u . Note that we are in a similar situation as in part (i) of this proof and thus it can be shown in the same way that there is a subsequence (x_{n_k}) of (x_n) that converges to some \tilde{x} in $\mathcal{X}^{1,1}(t_0, x_0)$. Now, fix $t \in (t_0, T]$ and a sequence (s_k) in A with $s_k \in A_{n_k}$ for each k and with $s_k \rightarrow t$ as $k \rightarrow \infty$. By Theorem 10.8.ii in [10] and lower semi-continuity of u ,

$$u(t_0, x_0) \geq \liminf_k \left[\int_{t_0}^{s_k} \ell(s, x'_{n_k}(s)) ds + u(s_k, x_{n_k}) \right] \geq \int_{t_0}^t \ell(s, \tilde{x}'(s)) ds + u(t, \tilde{x}),$$

i.e., there is an $x \in \mathcal{X}^{1,1}(t_0, x_0)$ such that, for every $t \in [t_0, T]$, (8.1) holds. From this point, (3.4) follows easily. \square

Lemma 8.2. *Assume (H1). An l.s.c. function $u : [0, T] \times \Omega \rightarrow \mathbb{R} \cup \{\infty\}$ is an l.s.c. minimax subsolution of (TVP) if and only if $u(T, \cdot) \leq h$ and, for every*

$(t_0, x_0) \in [0, T] \times \Omega$, $t \in (t_0, T]$, and $x \in \mathcal{X}^{1,1}(t_0, x_0)$, we have

$$(8.3) \quad u(t_0, x_0) \leq \int_{t_0}^t \ell(s, x'(s)) ds + u(t, x).$$

Proof. (i) Let u be an l.s.c. minimax subsolution of (TVP). For the sake of a contradiction, assume that there exist a $(t_0, x_0) \in [0, T] \times \Omega$, a $t_1 \in (t_0, T]$, an $x \in \mathcal{X}^{1,1}(t_0, x_0)$, and an $\varepsilon > 0$ such that

$$(8.4) \quad u(t_0, x_0) - u(t_1, x) - \int_{t_0}^{t_1} \ell(s, x'(s)) ds > \varepsilon$$

as well as $(t_1, x) \in \text{dom}(u)$ and $\int_t^{t_1} \ell(s, x'(s)) ds < \infty$ for all $t \in [t_0, t_1]$. Put

$$\tilde{s} := \inf \left\{ t \in (t_0, t_1] : u(t, x) - u(t_1, x) - \int_t^{t_1} \ell(s, x'(s)) ds \leq \frac{t_1 - t}{t_1 - t_0} \cdot \varepsilon \right\}.$$

We show that this infimum is attained. To this end, consider a sequence (s_n) in $(t_0, t_1]$ with $s_n \rightarrow \tilde{s}$ and $u(s_n, x) - u(t_1, x) - \int_{s_n}^{t_1} \ell(s, x'(s)) ds \leq \frac{t_1 - s_n}{t_1 - t_0} \cdot \varepsilon$. Then

$$\begin{aligned} u(\tilde{s}, x) - \int_{\tilde{s}}^{t_1} \ell(s, x'(s)) ds &\leq \liminf_n u(s_n, x) - \lim_n \int_{s_n}^{t_1} \ell(s, x'(s)) ds \\ &\leq \liminf_n \left[u(s_n, x) - \int_{s_n}^{t_1} \ell(s, x'(s)) ds \right] \leq \frac{t_1 - \tilde{s}}{t_1 - t_0} \cdot \varepsilon + u(t_1, x), \end{aligned}$$

i.e., \tilde{s} is a minimum and $(\tilde{s}, x) \in \text{dom}(u)$. Moreover, by (8.4), $t_0 < \tilde{s} \leq t_1$. Finally, by (3.5), there is a $\delta \in (0, \tilde{s} - t_0]$ such that $u(\tilde{s} - \delta, x) - u(\tilde{s}, x) - \int_{\tilde{s} - \delta}^{\tilde{s}} \ell(s, x'(s)) ds \leq \frac{\delta}{t_1 - t_0} \cdot \varepsilon$. Hence, $u(\tilde{s} - \delta, x) - u(t_1, x) - \int_{\tilde{s} - \delta}^{t_1} \ell(s, x'(s)) ds \leq \frac{t_1 - (\tilde{s} - \delta)}{t_1 - t_0} \cdot \varepsilon$, which is a contradiction to the minimality of \tilde{s} .

(ii) Showing the remaining direction is straight-forward. \square

Proof of Theorem 4.3. (a) Establishing the lower semi-continuity of v_0 is quite standard. It is very similar to the proof of Lemma 5.2 and actually slightly easier as no probability is involved (cf. also Proposition 3.1 in [13] for the non-path-dependent case). To deduce that v_0 is an l.s.c. minimax solution, it suffices to apply the dynamic programming principle with the existence of a minimizer for (DOC) (cf. [10]) together with Lemmata 8.1 and 8.2. Finally, we can apply Lemmata 8.1 and 8.2 again to obtain a comparison principle between l.s.c. minimax subsolutions and minimax supersolutions, from which uniqueness follows.

(b) Taking Remark 5.4 into account, one can see that the proof follows essentially from the content of Section 7. The measures $\mathbb{P}_{t_0, x_0, n}$ need to be replaced by the Dirac measures under which $X = x(\cdot \wedge t_0)$ a.s. for each $(t_0, x_0) \in [0, T] \times \Omega$ and one should note that the domains of the controls are different ($[0, T]$ here vs. $[0, T] \times \Omega$ in Section 7). Moreover, the BSDE argument in part (i) of the proof of Theorem 7.2 needs to be replaced by the deterministic dynamic programming principle. \square

REFERENCES

- [1] J.-P. Aubin and G. Haddad, *History path dependent optimal control and portfolio valuation and management*, Positivity **6** (2002), no. 3, 331–358. Special issue of the mathematical economics.
- [2] J. Backhoff-Veraguas, D. Lacker, and L. Tangpi, *Nonexponential Sanov and Schilder theorems on Wiener space: BSDEs, Schrödinger problems and control*, Annals of Applied Probability **30** (2020), no. 3, 1321–1367.

- [3] M. Bardi and I. Capuzzo-Dolcetta, *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*, Systems & Control: Foundations & Applications, Birkhäuser Boston, Inc., Boston, MA, 1997. With appendices by M. Falcone and P. Soravia.
- [4] E. Bayraktar and C. Keller, *Path-dependent Hamilton-Jacobi equations in infinite dimensions*, J. Funct. Anal. **275** (2018), no. 8, 2096–2161.
- [5] J. Bernis and P. Bettiol, *Solutions to the Hamilton-Jacobi equation for Bolza problems with discontinuous time dependent data*, ESAIM Control Optim. Calc. Var. **26** (2020), Paper No. 66, 35 pp.
- [6] ———, *Solutions to the Hamilton-Jacobi equation for Bolza problems with state constraints and discontinuous time dependent data*, Large-Scale Scientific Computing. LSSC 2019. Lecture Notes in Computer Science, vol 11958. Springer, Cham, 2020.
- [7] P. Bettiol and R. Vinter, *The Hamilton Jacobi equation for optimal control problems with discontinuous time dependence*, SIAM J. Control Optim. **55** (2017), no. 2, 1199–1225.
- [8] G. Buttazzo and M. Belloni, *A survey on old and recent results about the gap phenomenon in the calculus of variations*, Recent developments in well-posed variational problems, Math. Appl., 331, Kluwer Acad. Publ., Dordrecht, 1995, pp. 1–27.
- [9] O. Carja, *The minimum time function for semilinear evolutions*, SIAM J. Control Optim. **50** (2012), no. 3, 1265–1282.
- [10] L. Cesari, *Optimization—theory and applications*, Applications of Mathematics (New York), vol. 17, Springer-Verlag, New York, 1983. Problems with ordinary differential equations.
- [11] A. Cosso, S. Federico, F. Gozzi, M. Rosestolato, and N. Touzi, *Path-dependent equations and viscosity solutions in infinite dimension*, Ann. Probab. **46** (2018), no. 1, 126–174.
- [12] A. Cosso and F. Russo, *Strong-viscosity solutions: classical and path-dependent PDEs*, Osaka J. Math. **56** (2019), no. 2, 323–373.
- [13] G. Dal Maso and H. Frankowska, *Value functions for Bolza problems with discontinuous Lagrangians and Hamilton-Jacobi inequalities*, ESAIM Control Optim. Calc. Var. **5** (2000), 369–393.
- [14] R. De Arcangelis, *Some remarks on the identity between a variational integral and its relaxed functional*, Ann. Univ. Ferrara Sez. VII (N.S.) **35** (1989), 135–145 (1990).
- [15] S. Drapeau, G. Heyne, and M. Kupper, *Minimal supersolutions of convex BSDEs*, Ann. Probab. **41** (2013), no. 6, 3973–4001.
- [16] S. Drapeau, M. Kupper, E. Rosazza Gianin, and L. Tangpi, *Dual representation of minimal supersolutions of convex BSDEs*, Ann. Inst. Henri Poincaré Probab. Stat. **52** (2016), no. 2, 868–887.
- [17] S. Drapeau and C. Mainberger, *Stability and Markov property of forward backward minimal supersolutions*, Electron. J. Probab. **21** (2016), Paper No. 41, 15.
- [18] I. Ekren, C. Keller, N. Touzi, and J. Zhang, *On viscosity solutions of path dependent PDEs*, Ann. Probab. **42** (2014), no. 1, 204–236.
- [19] I. Ekren, *Viscosity solutions of obstacle problems for fully nonlinear path-dependent PDEs*, Stochastic Process. Appl. **127** (2017), no. 12, 3966–3996.
- [20] I. Ekren, N. Touzi, and J. Zhang, *Viscosity solutions of fully nonlinear parabolic path dependent PDEs: Part I*, Ann. Probab. **44** (2016), no. 2, 1212–1253.
- [21] ———, *Viscosity solutions of fully nonlinear parabolic path dependent PDEs: Part II*, Ann. Probab. **44** (2016), no. 4, 2507–2553.
- [22] I. Ekren and J. Zhang, *Pseudo-Markovian viscosity solutions of fully nonlinear degenerate PPDEs*, Probab. Uncertain. Quant. Risk **1** (2016), Paper No. 6, 34 pp.
- [23] H. Frankowska, *Optimal trajectories associated with a solution of the contingent Hamilton-Jacobi equation*, Appl. Math. Optim. **19** (1989), no. 3, 291–311.
- [24] M. I. Gomoyunov, N. Y. Lukoyanov, and A. R. Plaksin, *Path-Dependent Hamilton-Jacobi Equations: The Minimax Solutions Revised*, arXiv preprint arXiv:2010.06190 (2020).
- [25] U. G. Haussmann, *A probabilistic approach to the generalized Hessian*, Math. Oper. Res. **17** (1992), no. 2, 411–443.
- [26] ———, *Generalized solutions of the Hamilton-Jacobi equation of stochastic control*, SIAM J. Control Optim. **32** (1994), no. 3, 728–743.
- [27] U. G. Haussmann and J.-P. Lepeltier, *On the existence of optimal controls*, SIAM J. Control Optim. **28** (1990), no. 4, 851–902.

- [28] H. Kaise, T. Kato, and Y. Takahashi, *Hamilton–Jacobi partial differential equations with path-dependent terminal costs under superlinear Lagrangians*, Proceedings of the 23-rd International Symposium on Mathematical Theory of Networks and Systems (MTNS2018), Hong Kong University of Science and Technology, 2018, pp. 692–699.
- [29] C. Keller, *Viscosity solutions of path-dependent integro-differential equations*, Stochastic Process. Appl. **126** (2016), no. 9, 2665–2718.
- [30] M. Kisielewicz, *Differential inclusions and optimal control*, Mathematics and its Applications (East European Series), vol. 44, Kluwer Academic Publishers Group, Dordrecht; PWN—Polish Scientific Publishers, Warsaw, 1991.
- [31] N. Yu. Lukoyanov, *On the properties of the value functional of a differential game with hereditary information*, J. Appl. Math. Mech. **65** (2001), no. 3, 361–370.
- [32] J. Ma, Z. Ren, N. Touzi, and J. Zhang, *Large deviations for non-Markovian diffusions and a path-dependent Eikonal equation*, Ann. Inst. Henri Poincaré Probab. Stat. **52** (2016), no. 3, 1196–1216.
- [33] A. Misztela, *The value function representing Hamilton–Jacobi equation with Hamiltonian depending on value of solution*, ESAIM Control Optim. Calc. Var. **20** (2014), no. 3, 771–802.
- [34] T. Pham and J. Zhang, *Two person zero-sum game in weak formulation and path dependent Bellman–Isaacs equation*, SIAM J. Control Optim. **52** (2014), no. 4, 2090–2121.
- [35] S. Plaskacz and M. Quincampoix, *On representation formulas for Hamilton–Jacobi’s equations related to calculus of variations problems*, Topol. Methods Nonlinear Anal. **20** (2002), no. 1, 85–118.
- [36] Z. Ren, *Viscosity solutions of fully nonlinear elliptic path dependent partial differential equations*, Ann. Appl. Probab. **26** (2016), no. 6, 3381–3414.
- [37] ———, *Perron’s method for viscosity solutions of semilinear path dependent PDEs*, Stochastics **89** (2017), no. 6–7, 843–867.
- [38] Z. Ren and M. Rosestolato, *Viscosity solutions of path-dependent PDEs with randomized time*, SIAM J. Math. Anal. **52** (2020), no. 2, 1943–1979.
- [39] Z. Ren, N. Touzi, and J. Zhang, *An overview of viscosity solutions of path-dependent PDEs*, Stochastic analysis and applications 2014, Springer Proc. Math. Stat., 100, Springer, Cham, 2014, pp. 397–453.
- [40] ———, *Comparison of viscosity solutions of fully nonlinear degenerate parabolic path-dependent PDEs*, SIAM J. Math. Anal. **49** (2017), no. 5, 4093–4116.
- [41] ———, *Comparison of viscosity solutions of semilinear path-dependent PDEs*, SIAM J. Control Optim. **58** (2020), no. 1, 277–302.
- [42] A. I. Subbotin, *Generalized solutions of first-order PDEs*, Systems & Control: Foundations & Applications, Birkhäuser Boston Inc., Boston, MA, 1995. The dynamical optimization perspective, Translated from the Russian.
- [43] N. N. Subbotina, *The method of characteristics for Hamilton–Jacobi equations and applications to dynamical optimization*, J. Math. Sci. (N.Y.) **135** (2006), no. 3, 2955–3091.
- [44] N. N. Subbotina, A. I. Subbotin, and V. E. Tret’jakov, *Stochastic and deterministic control. Differential inequalities*, Problems Control Inform. Theory/Problemy Upravlen. Teor. Inform. **14** (1985), no. 6, 405–419, P1–P15. With the Russian original.
- [45] R. Vinter, *Optimal control*, Modern Birkhäuser Classics, Birkhäuser Boston, Ltd., Boston, MA, 2010. Paperback reprint of the 2000 edition.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, UNITED STATES

Email address: `erhan@umich.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CENTRAL FLORIDA, ORLANDO, FL 32816, UNITED STATES

Email address: `christian.keller@ucf.edu`