

Remarks on pseudo-vertex-transitive graphs with small diameter

Jack H. Koolen^{a,b}, Jae-Ho Lee^{*c}, and Ying-Ying Tan^d

^aSchool of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui, 230026, PR China.

^bCAS Wu Wen-Tsun Key Laboratory of Mathematics, University of Science and Technology of China, 96 Jinzhai Road, Hefei, Anhui, 230026, PR China.

^cDepartment of Mathematics and Statistics, University of North Florida, Jacksonville, FL 32224, USA.

^dSchool of Mathematics & Physics, Anhui Jianzhu University, Hefei, Anhui, 230201, PR China.

Abstract

Let Γ denote a Q -polynomial distance-regular graph with vertex set X and diameter D . Let A denote the adjacency matrix of Γ . For a vertex $x \in X$ and for $0 \leq i \leq D$, let $E_i^*(x)$ denote the projection matrix to the i th subconstituent space of Γ with respect to x . The Terwilliger algebra $T(x)$ of Γ with respect to x is the semisimple subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by $A, E_0^*(x), \dots, E_D^*(x)$. Let $V = \mathbb{C}^X$ denote a \mathbb{C} -vector space consisting of complex column vectors with rows indexed by X . We say Γ is pseudo-vertex-transitive whenever for any vertices $x, y \in X$, both (i) the Terwilliger algebras $T(x)$ and $T(y)$ of Γ are isomorphic; and (ii) there exists a \mathbb{C} -vector space isomorphism $\rho : V \rightarrow V$ such that $(\rho A - A\rho)V = 0$ and $(\rho E_i^*(x) - E_i^*(y)\rho)V = 0$ for all $0 \leq i \leq D$. In this paper we discuss pseudo-vertex transitivity for distance-regular graphs with diameter $D \in \{2, 3, 4\}$. In the case of diameter two, a strongly regular graph Γ is thin, and Γ is pseudo-vertex-transitive if and only if every local graph of Γ has the same spectrum. In the case of diameter three, Taylor graphs are thin and pseudo-vertex-transitive. In the case of diameter four, antipodal tight graphs are thin and pseudo-vertex-transitive.

Keywords: Distance-regular graph; Terwilliger algebra; Pseudo-vertex-transitive; Strongly regular graph; Taylor graph; Antipodal tight graph

2020 Mathematics Subject Classification: 05E30

1 Introduction

Let Γ denote a Q -polynomial distance-regular graph with vertex set X and diameter D . We note that Γ can be regarded as a combinatorial analogue of a compact rank one symmetric space [3, p. 311]. We recall the Terwilliger algebra of Γ . Let A denote the adjacency matrix of Γ . Fix a *base vertex* $x \in X$. For $0 \leq i \leq D$, let $E_i^* = E_i^*(x)$ denote the projection matrix onto the i -th

*Corresponding author.

E-mail: koolen@ustc.edu.cn (J.H. Koolen), jaeho.lee@unf.edu (J.-H. Lee), tansusan1@ahjzu.edu.cn (Y.-Y. Tan)

subconstituent space of Γ with respect to x . Let $T(x)$ denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by $A, E_0^*, E_1^*, \dots, E_D^*$. The algebra $T(x)$ is semisimple and called the *Terwilliger algebra* (or *subconstituent algebra*) of Γ with respect to x [26]. Let $T(y)$ denote the Terwilliger algebra of Γ with respect to another base vertex $y \in X$. We say the Terwilliger algebras $T(x)$ and $T(y)$ are *isomorphic* whenever there exists a \mathbb{C} -algebra isomorphism from $T(x)$ to $T(y)$ that sends A to A and E_i^* to $E_i^*(y)$ for $0 \leq i \leq D$. We remark that the isomorphism class of $T(x)$ may depend on the choice of the base vertex x . For instance, the twisted Grassmann graph, introduced by van Dam and Koolen [8], has two orbits of the automorphism group on its vertex set, say X_1 and X_2 . For vertices $x \in X_1$ and $y \in X_2$, the Terwilliger algebras $T(x)$ and $T(y)$ of the twisted Grassmann graph are not isomorphic; cf [2]. Let V denote a \mathbb{C} -vector space consisting of complex column vectors with rows indexed by X . Observe that $\text{Mat}_X(\mathbb{C})$ acts on V by left multiplication. View V as a left module for $T(x)$ and call this the *standard module of $T(x)$* (or *standard $T(x)$ -module*). Since $T(x)$ is semisimple, V decomposes into a direct sum of irreducible $T(x)$ -modules. We say Γ is *pseudo-vertex-transitive* whenever for every pair of vertices $x, y \in X$ both (i) the Terwilliger algebras $T(x)$ and $T(y)$ of Γ are isomorphic; and (ii) there exists a \mathbb{C} -vector space isomorphism $\rho : V \rightarrow V$ such that $(\rho A - A\rho)V = 0$ and $(\rho E_i^*(x) - E_i^*(y)\rho)V = 0$ for all $0 \leq i \leq D$. In other words, when Γ is pseudo-vertex-transitive, it means that the isomorphism class of the standard $T(x)$ -module of Γ does not depend on the base vertex x ; consequently, the multiplicities of irreducible $T(x)$ -modules do not depend on the base vertex x .

We recall the notion of the thinness of Γ . Let W denote an irreducible $T(x)$ -module. Then W is a direct sum of nonzero spaces among $E_0^*W, E_1^*W, \dots, E_D^*W$, and also a direct sum of nonzero spaces among E_0W, E_1W, \dots, E_DW , where E_i is the primitive idempotent of Γ . We note that the dimension of E_i^*W is at most 1 for $0 \leq i \leq D$ if and only if the dimension of E_iW is at most 1 for $0 \leq i \leq D$ [26, Lemma 3.9]; in this case, W is called *thin*. The graph Γ is called *thin* whenever every irreducible $T(x)$ -module is thin for every vertex $x \in X$.

In the present paper, we discuss the thinness and pseudo-vertex transitivity of a Q -polynomial distance-regular graph with small diameter $D \in \{2, 3, 4\}$. In the case of $D = 2$, a distance-regular graph is strongly regular with parameters, say (n, k, a, c) . We show that a strongly regular graph Γ is thin and its pseudo-vertex transitivity is determined by the spectrum of the local graph of Γ ; cf. Proposition 5.8 and Theorem 5.11. In the case of $D \in \{3, 4\}$, we discuss only the antipodal double covers. We will discuss the general case of Q -polynomial distance-regular graphs with diameter $D \geq 3$ in a future paper. Let Γ denote a distance-regular antipodal double cover with diameter $D \in \{3, 4\}$. If $D = 3$, then Γ is a Taylor graph and thin; cf. Proposition 7.4. We show that the isomorphism class of the standard module of the Terwilliger algebra of Γ is determined by its intersection numbers, that is, Γ is pseudo-vertex-transitive; cf. Theorem 7.6. If $D = 4$, then Γ is tight, denoted by $\text{AT4}(p, q, 2)$, where p and $-q$ ($p > -q$) are nontrivial eigenvalues of the first subconstituent of Γ . We show $\text{AT4}(p, q, 2)$ is thin and pseudo-vertex-transitive; cf. Proposition 8.6 and Theorem 8.7.

We remark that the thin property of a Q -polynomial distance-regular antipodal double cover with $D \in \{3, 4\}$ plays a role in determining pseudo-vertex transitivity; cf. Sections 7 and 8. In this context, we could ask the following question: for $D \geq 3$ does the thinness of a Q -polynomial

distance-regular graph guarantee its pseudo-vertex transitivity? Recently, Ito and Koolen suggested the following problem:

Problem 1.1 (The Ito-Koolen Problem [23]). Classify all thin pseudo-vertex-transitive Q -polynomial distance-regular graphs with large enough diameter.

The ultimate goal of our study is to classify thin pseudo-vertex-transitive Q -polynomial distance-regular graphs. The present paper shows that such graphs with small diameter are difficult to classify.

This paper is organized as follows. Sections 2 and 3 review some preliminaries concerning distance-regular graphs and the Terwilliger algebra. Section 4 introduces the notion of pseudo-vertex-transitive distance-regular graphs. It also contains what it means for two thin Q -polynomial distance-regular graphs to be T -isomorphic. Section 5 deals with the thinness of a strongly regular graph with parameters (n, k, a, c) and contains a characterization of pseudo-vertex transitivity for a strongly regular graph. It also contains several examples of cospectral strongly regular graphs and discusses their pseudo-vertex transitivity. Section 6 recalls the notion of antipodal double covers and tight distance-regular graphs and their properties. Section 7 deals with Taylor graphs and their thinness and pseudo-vertex transitivity. Section 8 discusses the case when Γ is an antipodal tight graph. The paper ends with a brief summary and directions for future work in Section 9.

Throughout the paper, we use the following notation. Let X denote a nonempty finite set. Let $\text{Mat}_X(\mathbb{C})$ denote the \mathbb{C} -algebra consisting of all complex matrices whose rows and columns are indexed by X . Let $V = \mathbb{C}^X$ denote the \mathbb{C} -vector space consisting of all complex column vectors with rows indexed by X . We endow V with the standard Hermitian inner product $\langle u, v \rangle = u^t \bar{v}$ for $u, v \in V$. We view V as a left module for $\text{Mat}_X(\mathbb{C})$, called the *standard module*.

2 Preliminaries: distance-regular graphs

In this section, we recall some definitions and notation. Let Γ denote a finite, simple, undirected, connected graph with vertex set X and diameter $D = \max\{\partial(x, y) \mid x, y \in X\}$, where ∂ denotes the path-length distance function for Γ . For a vertex $x \in X$, define $\Gamma_i(x) = \{y \in X \mid \partial(x, y) = i\}$ for $0 \leq i \leq D$. The subgraph $\Delta_i(x)$, $0 \leq i \leq D$, of Γ induced by $\Gamma_i(x)$ is called the *i th subconstituent* of Γ with respect to x . The number $|\Gamma_1(x)|$ is called the *valency* of x in Γ . A graph Γ is said to be *regular with valency k* (or *k -regular*) if each vertex of Γ has valency k . We abbreviate $\Delta(x) = \Delta_1(x)$, the first subconstituent of Γ with respect to x , and call this the *local graph* of Γ at x . Let λ denote an eigenvalue of $\Delta(x)$. We call λ a *local eigenvalue* of Γ with respect to x . We say a graph Γ is *locally Δ* whenever all local graphs of Γ are isomorphic to Δ .

We say Γ is *distance-regular* whenever for all integers $0 \leq h, i, j \leq D$ and for all vertices $x, y \in X$ with $\partial(x, y) = h$, the number $p_{ij}^h = |\Gamma_i(x) \cap \Gamma_j(y)|$ is independent of x and y . The constants p_{ij}^h are called the *intersection numbers* of Γ . We abbreviate $a_i = p_{1,i}^i$ ($0 \leq i \leq D$), $b_i = p_{1,i+1}^i$ ($0 \leq i \leq D-1$) and $c_i = p_{1,i-1}^i$ ($1 \leq i \leq D$). Observe Γ is regular with valency $k = b_0$, and $c_i + a_i + b_i = k$ for $0 \leq i \leq D$, where we define $c_0 = 0$ and $b_D = 0$. The sequence $\{b_0, b_1, \dots, b_{D-1}; c_1, c_2, \dots, c_D\}$ is called the *intersection array* of Γ .

We assume that Γ is distance-regular with diameter D . For $0 \leq i \leq D$, let A_i denote the matrix in $\text{Mat}_X(\mathbb{C})$ defined by

$$(A_i)_{xy} = \begin{cases} 1 & \text{if } \partial(x, y) = i, \\ 0 & \text{if } \partial(x, y) \neq i, \end{cases} \quad (x, y \in X).$$

We call A_i the i th *distance matrix* of Γ . We abbreviate $A := A_1$, called the *adjacency matrix* of Γ . Observe (i) each A_i is real symmetric; (ii) $A_0 = I$; (iii) $\sum_{i=0}^D A_i = J$, the all-ones matrix; (iv) $A_i A_j = \sum_{h=0}^D p_{ij}^h A_h$ ($0 \leq i, j \leq D$). By these facts, we find that A_0, A_1, \dots, A_D is a basis for a commutative subalgebra M of $\text{Mat}_X(\mathbb{C})$, which we call the *Bose-Mesner algebra* of Γ . It is known that A generates M . The algebra M has a second basis E_0, E_1, \dots, E_D such that (i) $E_0 = |X|^{-1} J$; (ii) $\sum_{i=0}^D E_i = I$; (iii) $E_i E_j = \delta_{ij} E_i$; cf. [1, p.45]. We call E_i the i th *primitive idempotent* of Γ . Since $\{E_i\}_{i=0}^D$ is a basis for M , there exist complex scalars $\{\theta_i\}_{i=0}^D$ such that $A = \sum_{i=0}^D \theta_i E_i$. Observe $A E_i = E_i A = \theta_i E_i$ for $0 \leq i \leq D$. The scalars $\{\theta_i\}_{i=0}^D$ are real [3, p.197], and mutually distinct as A generates M . We call θ_i the *eigenvalue* of Γ associated with E_i for $0 \leq i \leq D$. Observe $V = E_0 V + E_1 V + \dots + E_D V$, an orthogonal direct sum. For $0 \leq i \leq D$, $E_i V$ is the eigenspace of A associated with θ_i . For $0 \leq i \leq D$, we denote by m_i the rank of E_i and observe $m_i = \dim(E_i V)$. We call m_i the *multiplicity* of θ_i . By the *spectrum* of Γ , we mean the multiset containing its eigenvalues, each with its multiplicity, denoted by $\text{Spec}(\Gamma) = \{\theta_0^{m_0}, \theta_1^{m_1}, \dots, \theta_D^{m_D}\}$.

We recall the notion of the Q -polynomial property of Γ . Let \circ denote the entrywise product in $\text{Mat}_X(\mathbb{C})$. Since $A_i \circ A_j = \delta_{ij} A_i$ ($0 \leq i, j \leq D$), the Bose-Mesner algebra M is closed under \circ . Since $\{E_i\}_{i=0}^D$ is a basis for M , there exist complex scalars q_{ij}^h such that

$$E_i \circ E_j = |X|^{-1} \sum_{h=0}^D q_{ij}^h E_h, \quad (0 \leq i, j \leq D).$$

By [1, p.48, 49] the scalars q_{ij}^h are real and nonnegative. The q_{ij}^h are called the *Krein parameters* of Γ . We say Γ is *Q-polynomial* (with respect to the ordering E_0, E_1, \dots, E_D) whenever for all integers $0 \leq h, i, j \leq D$, $q_{ij}^h = 0$ (resp. $q_{ij}^h \neq 0$) if one of h, i, j is greater than (resp. equal to) the sum of the other two [1, p.235]. From now on, unless otherwise stated, we assume that a Q -polynomial distance-regular graph discussed in this paper has the ordering E_0, E_1, \dots, E_D . For more background information about distance-regular graphs, we refer the reader to [1, 3, 9].

3 Preliminaries: the Terwilliger algebra

In this section, we recall the Terwilliger algebra of a Q -polynomial distance-regular graph. Let Γ denote a Q -polynomial distance-regular graph with vertex set X and diameter D . Fix a vertex $x \in X$. We refer to x as a “base” vertex. For $0 \leq i \leq D$, we define the diagonal matrix $E_i^* = E_i^*(x) \in \text{Mat}_X(\mathbb{C})$ with diagonal entry

$$(E_i^*)_{yy} = \begin{cases} 1 & \text{if } \partial(x, y) = i, \\ 0 & \text{if } \partial(x, y) \neq i, \end{cases} \quad (y \in X).$$

We call E_i^* the i th *dual primitive idempotent* of Γ with respect to x . Observe $\sum_{i=0}^D E_i^* = I$ and $E_i^* E_j^* = \delta_{ij} E_i^*$. By these facts, $E_0^*, E_1^*, \dots, E_D^*$ is a basis for a commutative subalgebra $M^* = M^*(x)$ of $\text{Mat}_X(\mathbb{C})$ with respect to x , which we call the *dual Bose-Mesner algebra* of Γ . Recall the primitive idempotent E_i ($0 \leq i \leq D$) of Γ . For $0 \leq i \leq D$, define the diagonal matrix $A_i^* = A_i^*(x) \in \text{Mat}_X(\mathbb{C})$ with diagonal entry $(A_i^*)_{yy} = |X|(E_i)_{xy}$ for $y \in X$. By [26, p.379], $A_0^*, A_1^*, \dots, A_D^*$ is a basis for M^* . Moreover $A_0^* = I$ and $A_i^* A_j^* = \sum_{h=0}^D q_{ij}^h A_h^*$. We call A_i^* the i th *dual distance matrix* of Γ with respect to x . We abbreviate $A^* = A_1^*$ and call this the *dual adjacency matrix* of Γ with respect to x . The matrix A^* generates M^* [26, Lemma 3.11]. Since $\{E_i^*\}_{i=0}^D$ is a basis for M^* , there exist complex scalars $\{\theta_i^*\}_{i=0}^D$ such that $A^* = \sum_{i=0}^D \theta_i^* E_i^*$. Observe $A^* E_i^* = E_i^* A^* = \theta_i^* E_i^*$ for $0 \leq i \leq D$. The scalars $\{\theta_i^*\}_{i=0}^D$ are real [26, Lemma 3.11] and mutually distinct. We call θ_i^* the *dual eigenvalue* of Γ associated with E_i^* . Observe $V = E_0^* V + E_1^* V + \dots + E_D^* V$, an orthogonal direct sum. For $0 \leq i \leq D$, $E_i^* V$ is the eigenspace of A^* associated with θ_i^* , which we call the i th *subconstituent space* of Γ with respect to x .

Recall the Bose-Mesner algebra M of Γ and the dual Bose-Mesner algebra of $M^*(x)$ of Γ with respect to x . Let $T = T(x)$ denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by M and M^* . We call T the *Terwilliger algebra* (or *subconstituent algebra*) of Γ with respect to x [26]. Note that A and A^* generate T . The algebra T is finite-dimensional and noncommutative. The algebra T is semisimple since it is closed under the conjugate-transpose map. By [26, Lemma 3.2], we have the following relations in T . For $0 \leq i, j \leq D$,

$$\begin{aligned} E_i^* A E_j^* &= 0 & \text{if } |i - j| > 1, \\ E_i A^* E_j &= 0 & \text{if } |i - j| > 1. \end{aligned}$$

By a T -module, we mean a subspace W of V such that $BW \subseteq W$ for all $B \in T$. Observe that V is a T -module, called the *standard module of T* (or *standard T -module*). Two T -modules W, W' are *isomorphic* if there exists a \mathbb{C} -vector space isomorphism $\sigma : W \rightarrow W'$ such that

$$(\sigma B - B \sigma)W = 0,$$

for all $B \in T$. Let W be a T -module and let U be a T -submodule of W . The orthogonal complement of U in W is a T -module since T is closed under the conjugate transpose map. It follows that W decomposes into an orthogonal direct sum of irreducible T -modules. We observe that V decomposes an orthogonal direct sum of irreducible T -modules; choose one of the irreducible T -modules in this decomposition of V , denoted by W . By the *multiplicity* of W , we mean the number of irreducible T -modules in this decomposition which are isomorphic to W as T -modules.

Let W be an irreducible T -module. Then W decomposes into a direct sum of nonzero spaces among $E_0^* W, E_1^* W, \dots, E_D^* W$, and also a direct sum of nonzero spaces among $E_0 W, E_1 W, \dots, E_D W$. By the *endpoint* of W , we mean $\min\{i \mid 0 \leq i \leq D, E_i^* W \neq 0\}$. By the *dual endpoint* of W , we mean $\min\{i \mid 0 \leq i \leq D, E_i W \neq 0\}$. By the *diameter* of W , we mean $|\{i \mid 0 \leq i \leq D, E_i^* W \neq 0\}| - 1$. Let r denote the endpoint of W and d the diameter of W . By [26, Lemma 3.9], $E_i^* W \neq 0$ if and only if $r \leq i \leq r + d$, and $W = \sum_{h=0}^d E_{r+h}^* W$, an orthogonal direct sum; in addition, $\dim(E_i^* W) \leq 1$ for $0 \leq i \leq D$ if and only if $\dim(E_i W) \leq 1$ for $0 \leq i \leq D$. An irreducible T -module W is said to be *thin* whenever $\dim(E_i^* W) \leq 1$ for $0 \leq i \leq D$. There exists a unique thin irreducible T -module

with endpoint 0 and diameter D , which we call the *primary* T -module. Let W denote the primary T -module. Let v_0, v_1, \dots, v_D be a sequence of vectors of W , not all zero. This sequence is said to be a *standard basis* for W whenever both (i) $v_i \in E_i^*W$ for $0 \leq i \leq D$; and (ii) $\sum_{i=0}^D v_i \in E_0W$. For instance, the sequence $E_0^*\mathbb{1}, E_1^*\mathbb{1}, \dots, E_D^*\mathbb{1}$ is a standard basis for W [26, Lemma 3.6], where $\mathbb{1}$ is the all-ones vector. We note that the action of A on a standard basis for W is given by

$$Av_i = b_{i-1}v_{i-1} + a_iv_i + c_{i+1}v_{i+1} \quad (0 \leq i \leq D), \quad (3.1)$$

where a_i, b_i, c_i are the intersection numbers of Γ and $b_{-1}v_{-1} := 0$ and $c_{D+1}v_{D+1} := 0$.

Let W be a thin irreducible T -module with endpoint r and diameter d . We recall the actions of A and A^* on W . Take a nonzero vector $v_0 \in E_r^*W$. For $1 \leq i \leq d$, define $v_i = E_{r+i}^*Av_{i-1}$ in E_{r+i}^*W . Observe that the vector v_i is a basis for E_{r+i}^*W for each $0 \leq i \leq d$, and thus v_0, v_1, \dots, v_d is an orthogonal basis for W . Define the scalars $a_i(W), 0 \leq i \leq d$, and $x_i(W), 1 \leq i \leq d$, by

$$a_i(W) = \text{trace}(E_{r+i}^*A|_W), \quad x_i(W) = \text{trace}(E_{r+i}^*AE_{r+i-1}^*A|_W), \quad (3.2)$$

where $B|_W$ denotes the restriction of B to W . Let θ_i be the eigenvalue of Γ associated with E_i for $0 \leq i \leq D$. Note that (cf. [6, Lemma 5.10])

$$\sum_{i=0}^d a_i(W) = \sum_{i=0}^d \theta_{t+i}, \quad (3.3)$$

where t is the dual endpoint of W . For $0 \leq i \leq d$, the action of A on v_i is given as

$$Av_i = v_{i+1} + a_i(W)v_i + x_i(W)v_{i-1}, \quad (3.4)$$

where $v_{d+1} := 0$ and $x_0(W)v_{-1} := 0$; cf. [6, Theorem 5.7]. The action of A^* on v_i is given as $A^*v_i = \theta_{r+i}^*v_i$, where θ_j^* is the dual eigenvalue of Γ associated with E_j^* .

The graph Γ is said to be *thin with respect to x* whenever every irreducible $T(x)$ -module is thin. The graph Γ is said to be *thin* whenever Γ is thin with respect to every vertex x of Γ . See [27, Section 6] for examples of thin Q -polynomial distance-regular graphs.

Recall the local graph $\Delta(x)$ of Γ at $x \in X$. Observe that $\Delta(x)$ has $b_0 = k$ vertices and is a_1 -regular. Let $a_1 = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$ denote the local eigenvalues of Γ with respect to x . The spectrum of $\Delta(x)$ is called the *local spectrum* of Γ at x . Let W denote a thin irreducible T -module with endpoint 1. We observe that E_1^*W is a one-dimensional eigenspace for $E_1^*AE_1^*$ with corresponding eigenvalue, say λ . Note that λ is one of $\lambda_2, \dots, \lambda_k$. We call λ the *local eigenvalue* of W .

We finish this section with a comment, which will be useful later when calculating the dimension of the Terwilliger algebra.

Proposition 3.1. *Let $T = T(x)$ be the Terwilliger algebra of Γ with respect to x and V the standard T -module. Suppose that V has exactly r non-isomorphic irreducible T -modules $W_{11}, W_{21}, \dots, W_{r1}$ with $\dim(W_{i1}) = n_i$. For $1 \leq i \leq r$, let m_i be the multiplicity of W_{i1} . Then V decomposes into a direct sum of irreducible T -modules:*

$$V = \bigoplus_{i=1}^r \left(\bigoplus_{j=1}^{m_i} W_{ij} \right),$$

where W_{ij} and $W_{i'j'}$ are isomorphic as T -modules if and only if $i = i'$. The algebra T is isomorphic to the semisimple algebra

$$\text{Mat}_{n_1}(\mathbb{C}) \oplus \cdots \oplus \text{Mat}_{n_r}(\mathbb{C})$$

and

$$\dim(T) = \sum_{i=1}^r n_i^2. \quad (3.5)$$

Proof. Follows from Wedderburn's theory [7]. ■

4 Pseudo-vertex transitivity

In this section, we recall the notion of pseudo-vertex-transitive graphs. Throughout this section, we denote by Γ a Q -polynomial distance-regular graph with vertex set X and diameter D . Fix a base vertex $x \in X$. Let A denote the adjacency matrix of Γ and $A^*(x)$ the dual adjacency matrix of Γ with respect to x . Let $T(x)$ denote the Terwilliger algebra of Γ with respect to x .

We consider another Q -polynomial distance-regular graph Γ' with vertex set X' and diameter D . Fix a vertex $x' \in X'$. Denote by A' the adjacency matrix of Γ' and $A^{*'}(x')$ the dual adjacency matrix of Γ' with respect to x' . Let $T'(x')$ be the Terwilliger algebra of Γ' with respect to x' . We say the Terwilliger algebras $T(x)$ and $T'(x')$ are *isomorphic* whenever $|X| = |X'|$ and there exists a \mathbb{C} -algebra isomorphism from $T(x)$ to $T'(x')$ that sends $A \mapsto A'$ and $A^*(x) \mapsto A^{*'}(x')$. In this case, such an algebra isomorphism is called a *T -algebra isomorphism*.

Recall $V = \mathbb{C}^X$ the standard $T(x)$ -module of Γ . Let $V' = \mathbb{C}^{X'}$ denote the standard $T'(x')$ -module of Γ' . The standard modules V and V' are said to be *T -isomorphic* whenever (i) the Terwilliger algebras $T(x)$ and $T'(x')$ are isomorphic; and (ii) there exists a \mathbb{C} -vector space isomorphism $\rho : V \rightarrow V'$ such that

$$(\rho B - B' \rho)V = 0,$$

where $B \in T$ and $B' = f(B)$ with a T -algebra isomorphism f from $T(x)$ to $T'(x')$. That is, the diagram

$$\begin{array}{ccc} V & \xrightarrow{\rho} & V' \\ B \downarrow & & \downarrow B' \\ V & \xrightarrow{\rho} & V' \end{array}$$

commutes for all $B \in T(x)$. From this definition, we have the following lemma.

Lemma 4.1. *With the above notation, the standard modules V and V' are T -isomorphic if and only if there are ordered bases \mathcal{B} for V and \mathcal{B}' for V' such that the matrix representing A (resp. $A^*(x)$) with respect to \mathcal{B} is equal to the matrix representing A' (resp. $A^{*'}(x')$) with respect to \mathcal{B}' .*

Proof. Routine. ■

Definition 4.2. Two Q -polynomial distance-regular graphs Γ and Γ' are said to be *T -isomorphic* whenever for every pair of vertices $x \in X$ and $x' \in X'$, the standard $T(x)$ -module and the standard $T'(x')$ -module are T -isomorphic.

Lemma 4.3. *If Γ and Γ' are T -isomorphic, then the intersection array of Γ is equal to the intersection array of Γ' .*

Proof. Fix vertices $x \in X$ and $x' \in X'$. Let V denote the standard $T(x)$ -module of Γ and V' the standard $T'(x')$ -module of Γ' . Write $T = T(x)$, $T' = T'(x')$, $E_i^* = E_i^*(x)$, $E_i^{*'} = E_i^{*'}(x')$ for $0 \leq i \leq D$. Since Γ and Γ' are T -isomorphic, there exists a \mathbb{C} -vector space isomorphism $\rho : V \rightarrow V'$ such that $(\rho B - f(B)\rho)V = 0$ for all $B \in T$, where f is a T -algebra isomorphism from T to T' . Let W denote the primary T -module. Set $W' = \rho(W)$. Then W' is a T' -module since for all $B' \in T'$ we have $B'W' = B'\rho(W) = \rho(BW) \in \rho(W) = W'$. Clearly, the T' -module W' is irreducible. Moreover, W' has endpoint 0 since $E_0^*W \neq 0$ and $\rho(E_0^*W) = E_0^{*'}\rho(W) = E_0^{*'}W' \neq 0$. By these comments, W' is the primary T' -module; cf. [12, Proposition 8.4].

Let $\bar{\rho}$ denote the restriction of ρ to W . Then W and W' are T -isomorphic because for all $w \in W$ and for all $B \in T$ we have $\bar{\rho}(Bw) = \rho(Bw) = f(B)\rho(w) = f(B)\bar{\rho}(w)$. Recall the standard basis $\{E_i^*\mathbf{1}\}_{i=0}^D$ for W . We observe that $\bar{\rho}(E_i^*\mathbf{1}) = E_i^{*'}\bar{\rho}(\mathbf{1})$ for $0 \leq i \leq D$ and the sequence $\{E_i^{*'}\bar{\rho}(\mathbf{1})\}_{i=0}^D$ is a standard basis for W' . Since $\bar{\rho}(AE_i^*\mathbf{1}) = A'E_i^{*'}\bar{\rho}(\mathbf{1})$ for $0 \leq i \leq D$, the matrix representing A with respect to $\{E_i^*\mathbf{1}\}_{i=0}^D$ is equal to the matrix representing A' with respect to $\{E_i^{*'}\bar{\rho}(\mathbf{1})\}_{i=0}^D$. By (3.1), these matrices are tridiagonal whose entries are the intersection numbers of Γ and Γ' , respectively. The result follows. \blacksquare

Recall a Q -polynomial distance-regular graph Γ and the standard module $V = \mathbb{C}^X$. Now we define the pseudo-vertex transitivity of Γ .

Definition 4.4. The graph Γ is said to be *pseudo-vertex-transitive* whenever for every pair of vertices $x, y \in X$, both (i) the Terwilliger algebras $T(x)$ and $T(y)$ of Γ are isomorphic; and (ii) there exists a \mathbb{C} -vector space isomorphism $\rho : V \rightarrow V$ such that

$$(\rho A - A\rho)V = 0, \quad (\rho A^*(x) - A^*(y)\rho)V = 0.$$

That is, Γ is pseudo-vertex-transitive whenever for all vertices $x, y \in X$, the standard $T(x)$ -module and the standard $T(y)$ -module of Γ are T -isomorphic. By definition, we note that the isomorphism class of the Terwilliger algebra of a pseudo-vertex-transitive Γ does not depend on the base vertex.

Remark 4.5. (i) Let $\text{Aut}(\Gamma)$ denote the automorphism group of Γ . If $\text{Aut}(\Gamma)$ has a single group orbit on X , then Γ is pseudo-vertex-transitive. From this, it follows that every vertex-transitive graph is pseudo-vertex-transitive. The converse of this statement is not true; see Example 5.15 and Remark 7.7.

(ii) The twisted Grassmann graph $\tilde{J}_q(2D+1, D)$ is not vertex-transitive; it has two orbits of the automorphism group on its vertex set, say X_1 and X_2 . In [2], Bang *et al.* showed that all irreducible modules for the Terwilliger algebra $T(x)$ of $\tilde{J}_q(2D+1, D)$ with respect to $x \in X_2$ are thin, and there are non-thin irreducible modules for the Terwilliger algebra $T(y)$ of $\tilde{J}_q(2D+1, D)$ with respect to $y \in X_1$. It implies that $\tilde{J}_q(2D+1, D)$ is not pseudo-vertex-transitive. We remark that Tanaka and Wang determined all irreducible $T(x)$ -modules of $\tilde{J}_q(2D+1, D)$ for the thin case; cf [25].

(iii) With reference to (ii), let $x \in X_2$ be a base vertex of $\tilde{J}_q(2D+1, D)$ and let $T(x)$ be the Terwilliger algebra of $\tilde{J}_q(2D+1, D)$. We recall the Grassmann graph $J_q(2D+1, D)$. We note

that $J_q(2D + 1, D)$ is thin [27]. For a base vertex y of $J_q(2D + 1, D)$, let $T(y)$ be the Terwilliger algebra of $J_q(2D + 1, D)$. The local eigenvalues of $\tilde{J}_q(2D + 1, D)$ at x equal to the local eigenvalues of $J_q(2D + 1, D)$ at y . From this, it follows that the isomorphism classes of the irreducible $T(x)$ -modules with endpoint 1 are the same as the isomorphism classes of the irreducible $T(y)$ -modules with endpoint 1. However, their multiplicities are different; cf. [2]. In addition, it turns out that the Terwilliger algebras $T(x)$ and $T(y)$ are not isomorphic to each other; cf. [25].

Lemma 4.6. *Suppose that Γ is pseudo-vertex-transitive. Then for every pair of vertices $x, y \in X$ there exist ordered bases \mathcal{B}_x and \mathcal{B}_y for V such that the matrix representing A (resp. $A^*(x)$) with respect to \mathcal{B}_x is equal to the matrix representing A (resp. $A^*(y)$) with respect to \mathcal{B}_y .*

Proof. Immediate from Definition 4.4 and Lemma 4.1. ■

We finish this section with a comment. Suppose that Γ is pseudo-vertex-transitive. For $x \in X$ and for $0 \leq i \leq D$, let $\Delta_i(x)$ denote the i th subconstituents of Γ with respect to x . Then the spectrum of $\Delta_i(x)$ is determined by the characteristic polynomial for $E_i^*(x)AE_i^*(x)$. Let y be another vertex in X . Since Γ is pseudo-vertex-transitive, by Lemma 4.6 there are bases \mathcal{B}_1 and \mathcal{B}_2 for V such that the matrix representing $E_i^*(x)AE_i^*(x)$ with respect to \mathcal{B}_1 is equal to the matrix representing $E_i^*(y)AE_i^*(y)$ with respect to \mathcal{B}_2 . It follows that the characteristic polynomial for $E_i^*(x)AE_i^*(x)$ is equal to the characteristic polynomial for $E_i^*(y)AE_i^*(y)$. Therefore we obtain the following lemma.

Lemma 4.7. *Suppose that Γ is pseudo-vertex-transitive. Then for all integers $0 \leq i \leq D$ and for all vertices $x, y \in X$, the spectrum of $\Delta_i(x)$ and the spectrum of $\Delta_i(y)$ are equal to each other.*

Lemma 4.8. *Suppose that Γ and Γ' are T -isomorphic. Then for all integers $0 \leq i \leq D$ and for all vertices $x \in X$, $x' \in X'$, the spectrum of $\Delta_i(x)$ and the spectrum of $\Delta_i(x')$ are equal to each other.*

Proof. Similar to Lemma 4.7. Use Lemma 4.1. ■

5 Strongly regular graphs

In this section, we summarize the results of [28] and describe a characterization of strongly regular graphs that are pseudo-vertex-transitive. In addition, we give several examples of cospectral strongly regular graphs and discuss their pseudo-vertex transitivity.

5.1 Preliminaries

We begin by recalling the notion of strongly regular graphs. Let Γ be a k -regular graph with n vertices. We say Γ is *strongly regular* with parameters (n, k, a, c) whenever each pair of adjacent vertices has the same number a of common neighbors, and each pair of distinct non-adjacent vertices has the same number c of common neighbors. Note that a connected strongly regular graph with parameters (n, k, a, c) is distance-regular with diameter two and intersection array $\{k, k - a - 1; 1, c\}$.

For the rest of this section, we denote by Γ a connected strongly regular graph with parameters (n, k, a, c) . The graph Γ has exactly three eigenvalues k, σ, τ with

$$\sigma = \frac{a - c + \sqrt{D}}{2}, \quad \tau = \frac{a - c - \sqrt{D}}{2}, \quad (5.1)$$

where $D = (a - c)^2 + 4(k - c)$. The multiplicities m_σ and m_τ of σ and τ , respectively, are given by

$$m_\sigma = \frac{(n - 1)\tau + k}{\tau - \sigma}, \quad m_\tau = \frac{(n - 1)\sigma + k}{\sigma - \tau}. \quad (5.2)$$

Let X denote the vertex set of Γ . Fix a vertex $x \in X$. Consider the distance partition $\{\Gamma_i(x)\}_{i=0}^2$ of X . With respect to this partition, we write the adjacency matrix A of Γ in the partitioned matrix form:

$$A = \begin{pmatrix} 0 & \mathbf{1}^t & \mathbf{0} \\ \mathbf{1} & B_1 & N \\ \mathbf{0} & N^t & B_2 \end{pmatrix}, \quad (5.3)$$

where B_i ($i = 1, 2$) is the adjacency matrix of the i th subconstituent $\Delta_i(x)$ of Γ . For notational convenience, we denote by

$$\tilde{A}_1 := E_1^* A E_1^* = \begin{pmatrix} 0 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & B_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \end{pmatrix}, \quad \tilde{A}_2 := E_2^* A E_2^* = \begin{pmatrix} 0 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & B_2 \end{pmatrix},$$

where $E_i^* = E_i^*(x)$ ($i = 1, 2$) is the i th dual primitive idempotent of Γ . Note that $\Delta_1(x)$ is a -regular with k vertices and that $\Delta_2(x)$ is $(k - c)$ -regular with $n - k - 1$ vertices. It follows that $\Delta_1(x)$ and $\Delta_2(x)$ have trivial eigenvalues a and $k - c$, respectively. We say that an eigenvalue of $\Delta_i(x)$ ($i = 1, 2$) is *local* if it is not equal to an eigenvalue of Γ and has an eigenvector orthogonal to $\mathbf{1}$. The following is a characterization of a local eigenvalue of $\Delta_i(x)$ ($i = 1, 2$).

Lemma 5.1 (cf. [14, Theorem 10.6.3]). *Let $(i, j) = (1, 2)$ or $(i, j) = (2, 1)$. Then λ is a local eigenvalue of $\Delta_i(x)$ if and only if $a - c - \lambda$ is a local eigenvalue of $\Delta_j(x)$, with equal multiplicities.*

Lemma 5.2 (cf. [28, Proposition 3.1, Lemma 3.2]). *Let σ, τ be eigenvalues of Γ as in (5.1). Fix a vertex x of Γ and write $E_i^* = E_i^*(x)$ for $0 \leq i \leq 2$ and $T = T(x)$. Let A be the adjacency matrix of Γ as in (5.3). Let V be the standard T -module. Then the following (I) and (II) hold.*

- (I) *Let $v \in E_1^* V$ be an eigenvector of \tilde{A}_1 with an eigenvalue λ such that $\langle \mathbf{1}, v \rangle = 0$. Then $E_2^* A E_1^* v = 0$ if and only if $\lambda \in \{\sigma, \tau\}$. Let W be the subspace of V spanned by $v, E_2^* A E_1^* v$. Then W is a thin irreducible T -module. Moreover, the following (i) and (ii) hold.*
- (i) *Suppose $\lambda \notin \{\sigma, \tau\}$. Then $E_2^* A E_1^* v$ is an eigenvector of \tilde{A}_2 with an eigenvalue $\sigma + \tau - \lambda$. The T -module W has endpoint 1 and diameter 1. The vector v is a basis for $E_1^* W$ and $E_2^* A E_1^* v$ is a basis for $E_2^* W$. The matrix representing A with respect to a basis $\{v, E_2^* A E_1^* v\}$ for W is*

$$\begin{pmatrix} \lambda & -(\lambda - \sigma)(\lambda - \tau) \\ 1 & \sigma + \tau - \lambda \end{pmatrix}. \quad (5.4)$$

(ii) If $\lambda \in \{\sigma, \tau\}$, then $W = E_1^*W$. The T -module W has endpoint 1 and diameter 0. The action of A on W is given by $Av = \lambda v$.

(II) Let $u \in E_2^*V$ be an eigenvector of \tilde{A}_2 with an eigenvalue λ' such that $\langle \mathbf{1}, u \rangle = 0$. Then $E_1^*AE_2^*u = 0$ if and only if $\lambda' \in \{\sigma, \tau\}$. Let W be the subspace of V spanned by $u, E_1^*AE_2^*u$. Then W is a thin irreducible T -module. Moreover, the following (i) and (ii) hold.

(i) Suppose $\lambda' \notin \{\sigma, \tau\}$. Then $E_1^*AE_2^*u$ is an eigenvector of \tilde{A}_1 with an eigenvalue $\sigma + \tau - \lambda'$. The T -module W has endpoint 1 and diameter 1. The vector u is a basis for E_2^*W and $E_1^*AE_2^*u$ is a basis for E_1^*W . The matrix representing A with respect to a basis $\{E_1^*AE_2^*u, u\}$ for W is

$$\begin{pmatrix} \lambda' & -(\lambda' - \sigma)(\lambda' - \tau) \\ 1 & \sigma + \tau - \lambda' \end{pmatrix}.$$

(ii) If $\lambda' \in \{\sigma, \tau\}$, then $W = E_2^*W$. The T -module W has endpoint 2 and diameter 0. The action of A on W is given by $Au = \lambda' u$.

Recall the first subconstituent space E_1^*V of Γ . Note that $\dim(E_1^*V) = k$. Let

$$v_1, v_2, \dots, v_r, v_{r+1}, \dots, v_k \tag{5.5}$$

denote a sequence of eigenvectors of \tilde{A}_1 in E_1^*V such that $v_1 := E_1^*\mathbf{1}$ and $\langle v_i, v_j \rangle = 0$ for $i \neq j$. Observe that $\{v_i\}_{i=1}^k$ is an orthogonal basis for E_1^*V . Let

$$a = \lambda_1, \lambda_2, \dots, \lambda_r, \lambda_{r+1}, \dots, \lambda_k$$

denote the eigenvalues of \tilde{A}_1 corresponding to the eigenvectors in (5.5), respectively. Assume that $\lambda_i \notin \{\sigma, \tau\}$ for $1 \leq i \leq r$ and $\lambda_j \in \{\sigma, \tau\}$ for $r+1 \leq j \leq k$. We consider the second subconstituent space E_2^*V of Γ . Note that $\dim(E_2^*V) = n - k - 1$. For each vector v_i ($1 \leq i \leq r$) in (5.5), abbreviate $u_i = E_2^*AE_1^*v_i$. Observe that u_1 is an eigenvector of \tilde{A}_2 with eigenvalue $k - c$. By Lemma 5.2(I)(i), u_i ($2 \leq i \leq r$) is an eigenvector of \tilde{A}_2 associated with the eigenvalue $\sigma + \tau - \lambda_i \notin \{\sigma, \tau\}$. We choose the rest of the eigenvectors of \tilde{A}_2 in E_2^*V , denoted by

$$u_{r+1}, u_{r+2}, \dots, u_s,$$

where $s := n - k - 1$, such that $\{u_i\}_{i=1}^s$ spans E_2^*V and $\langle u_i, u_j \rangle = 0$ for $1 \leq i \neq j \leq s$. Observe that $\{u_i\}_{i=1}^s$ is a basis for E_2^*V . Let λ'_h denote the eigenvalue of \tilde{A}_2 associated with u_h for $r+1 \leq h \leq s$. By Lemma 5.2(II), we find $\lambda'_h \in \{\sigma, \tau\}$. Let $W_i = Tv_i$ for $1 \leq i \leq k$ and let $W'_h = Tu_h$ for $r+1 \leq h \leq s$. We say W_i (resp. W'_h) is associated with λ_i (resp. λ'_h). We find that W_i and W'_h are thin irreducible T -modules. In the following table, we display all thin irreducible T -modules of Γ :

thin irreducible T -module	basis	dimension	endpoint	diameter
W_1	$\{E_n^*AE_1^*v_1\}_{n=0}^2$	3	0	2
W_i ($2 \leq i \leq r$)	$v_i, E_2^*AE_1^*v_i$	2	1	1
W_j ($r+1 \leq j \leq k$)	v_j	1	1	0
W'_h ($r+1 \leq h \leq s$)	u_h	1	2	0

Since T is semisimple, the standard T -module V decomposes into an orthogonal direct sum of thin irreducible T -submodules:

$$V = W_1 \oplus \left(\bigoplus_{i=2}^r W_i \right) \oplus \left(\bigoplus_{j=r+1}^k W_j \right) \oplus \left(\bigoplus_{h=r+1}^s W'_h \right). \quad (5.7)$$

The following proposition classifies thin irreducible T -modules of Γ up to isomorphism.

Proposition 5.3 (cf. [28, Lemma 3.4]). *Let W_i (resp. W'_h) denote an irreducible T -module associated with an eigenvalue λ_i (resp. λ'_h) as in (5.6). Then the following (i) and (ii) hold.*

(i) *For $2 \leq i, j \leq k$, W_i and W_j are isomorphic as T -modules if and only if $\lambda_i = \lambda_j$.*

(ii) *For $r+1 \leq h, l \leq s$, W'_h and W'_l are isomorphic as T -modules if and only if $\lambda'_h = \lambda'_l$.*

Let ℓ_1 (resp. ℓ_2) denote the number of distinct eigenvalues of $\Delta(x)$ (resp. $\Delta_2(x)$) contained in $\{\sigma, \tau\}$ with respect to the eigenvectors orthogonal to $E_1^* \mathbf{1}$ (resp. $E_2^* \mathbf{1}$). Let ℓ'_1 (resp. ℓ'_2) denote the number of distinct eigenvalues of $\Delta(x)$ (resp. $\Delta_2(x)$) not contained in $\{\sigma, \tau\}$ with respect to the eigenvectors orthogonal to $E_1^* \mathbf{1}$ (resp. $E_2^* \mathbf{1}$). By Proposition 5.3, the isomorphism classes of irreducible T -modules of Γ are determined by $\text{Spec}(\Gamma)$, $\text{Spec}(\Delta(x))$, and $\text{Spec}(\Delta_2(x))$. It follows that there are $\ell_1 + \ell_2$ pairwise non-isomorphic irreducible $T(x)$ -modules of dimension one, and there are ℓ'_1 ($= \ell'_2$) pairwise non-isomorphic thin irreducible $T(x)$ -modules of dimension two. There is only one thin irreducible $T(x)$ -module of dimension three, namely the primary $T(x)$ -module. By Proposition 3.1 and (5.7), the algebra $T(x)$ is isomorphic to the semisimple algebra

$$(\text{Mat}_1(\mathbb{C})^{\oplus \ell_1}) \oplus (\text{Mat}_1(\mathbb{C})^{\oplus \ell_2}) \oplus (\text{Mat}_2(\mathbb{C})^{\oplus \ell'_1}) \oplus \text{Mat}_3(\mathbb{C}). \quad (5.8)$$

We call $(\ell_1, \ell'_1, \ell_2, \ell'_2)$ the *dimension sequence* of $T(x)$. Note that the dimension sequence of $T(x)$ depends on the base vertex x of Γ .

Lemma 5.4 (cf. [28, Theorem 1.1]). *Let x be a base vertex of Γ and let $T(x)$ be the Terwilliger algebra of Γ with respect to x . Then*

$$\begin{aligned} \dim T(x) &= \ell_1 + \ell_2 + 4\ell'_1 + 9 \\ &= \ell_1 + \ell_2 + 4\ell'_2 + 9, \end{aligned}$$

where $(\ell_1, \ell'_1, \ell_2, \ell'_2)$ is the dimension sequence of $T(x)$.

Proof. By (5.8) and Proposition 3.1. ■

5.2 A characterization of pseudo-vertex-transitive strongly regular graphs

In this subsection, we characterize pseudo-vertex-transitive strongly regular graphs. First, we discuss a relationship between $\text{Spec}(\Delta(x))$ and $\text{Spec}(\Delta_2(x))$ of Γ .

Lemma 5.5. *Let k, σ, τ be eigenvalues of Γ as in (5.1) with multiplicities $1, m_\sigma, m_\tau$, respectively. Fix a vertex x of Γ . The spectrum of the second subconstituent $\Delta_2(x)$ of Γ is determined by the spectrum of the local graph $\Delta(x)$ of Γ and the parameters (n, k, a, c) .*

Proof. Let us denote the spectrum of $\Delta(x)$ by

$$\{a^1, \sigma^{f_\sigma}, \tau^{f_\tau}, \lambda_1^{m_1}, \dots, \lambda_s^{m_s}\},$$

for some s . Here, it is possible that $f_\sigma = 0$ and/or $f_\tau = 0$. By Lemma 5.1, we can write the spectrum of $\Delta_2(x)$ as

$$\{(k-c)^1, \sigma^{g_\sigma}, \tau^{g_\tau}, (a-c-\lambda_1)^{m_1}, \dots, (a-c-\lambda_s)^{m_s}\},$$

where it is possible that $g_\sigma = 0$ and/or $g_\tau = 0$. Since $|\Gamma_1(x)| = k$ and $|\Gamma_2(x)| = n - k - 1$, we have

$$1 + f_\sigma + f_\tau + m_1 + \dots + m_s = k, \quad 1 + g_\sigma + g_\tau + m_1 + \dots + m_s = n - k - 1. \quad (5.9)$$

Using the equations in (5.9), we obtain

$$g_\sigma + g_\tau = n - 2k - 1 + f_\sigma + f_\tau. \quad (5.10)$$

Since the sum of all the eigenvalues of $\Delta_2(x)$ is the trace of \tilde{A}_2 , which is zero, it follows

$$\sigma g_\sigma + \tau g_\tau = -(k-c) - m_1(a-c-\lambda_1) - \dots - m_s(a-c-\lambda_s). \quad (5.11)$$

Solve the system of equations (5.10) and (5.11) for g_σ and g_τ to obtain

$$g_\sigma = -k + m_\sigma + f_\tau, \quad g_\tau = -k + m_\tau + f_\sigma. \quad (5.12)$$

The result follows. ■

Remark 5.6. By (5.12) we have

$$f_\sigma = k - m_\sigma + g_\tau, \quad f_\tau = k - m_\tau + g_\sigma.$$

Thus, the spectrum of $\Delta(x)$ is also determined by the spectrum of $\Delta_2(x)$.

Example 5.7. Consider the Johnson graph $J(8, 2)$ with vertex set $X = \binom{\Omega}{2}$, the collection of all 2-subsets of $\Omega = \{1, 2, \dots, 8\}$. The graph $J(8, 2)$ is strongly regular with parameters $(28, 12, 6, 4)$. Fix a vertex $x \in X$. The spectrum of $J(8, 2)$ is $\{12^1, 4^7, -2^{20}\}$ and the spectrum of the local graph $\Delta(x)$ is given by $\{6^1, 4^1, -2^5, 0^5\}$. Using Lemma 5.5, we obtain the spectrum of $\Delta_2(x)$ of $J(8, 2)$:

$$\{8^1, 4^0, -2^9, 2^5\}.$$

Proposition 5.8. *A connected strongly regular graph Γ is thin.*

Proof. By (5.6), Γ is thin with respect to x . By Proposition 5.3 and Lemma 5.5, the result follows. ■

Recall $E_i^* = E_i^*(x)$ the i th dual primitive idempotent of Γ ($i = 0, 1, 2$). Consider another connected strongly regular graph Γ' with parameters (n', k', a', c') and vertex set X' . Fix $x' \in X'$. Let A' be the adjacency matrix of Γ' and $A^{*'} = A^{*'}(x')$ the dual adjacency matrix of Γ' with respect to x' . Write $E_i^{*'} = E_i^{*'}(x')$ for $0 \leq i \leq 2$ and $T' = T'(x')$. With these notations, we state the following two lemmas.

Lemma 5.9. *Suppose that the parameters of Γ and Γ' are same, that is, $(n, k, a, c) = (n', k', a', c')$. Set $v = E_1^* \mathbf{1}$ and $v' = E_1'^* \mathbf{1}$. Then the T -modules Tv and $T'v'$ are T -isomorphic.*

Proof. Observe that Tv is the primary T -module with a basis $\{E_i^* A E_1^* v\}_{i=0}^2$; see (5.6). Define the vectors $\{v_i\}_{i=0}^2$ by

$$v_0 = E_0^* A E_1^* v, \quad v_1 = a^{-1} E_1^* A E_1^* v, \quad v_2 = c^{-1} E_2^* A E_1^* v.$$

Observe that $v_i = E_i^* \mathbf{1}$ ($0 \leq i \leq 2$), and thus $\{v_i\}_{i=0}^2$ is a standard basis for Tv . Similarly, we define the vectors $\{v'_i\}_{i=0}^2$ by

$$v'_0 = E_0'^* A' E_1'^* v', \quad v'_1 = a^{-1} E_1'^* A' E_1'^* v', \quad v'_2 = c^{-1} E_2'^* A' E_1'^* v'.$$

Then $\{v'_i\}_{i=0}^2$ is a standard basis for the primary T' -module $T'v'$. Define a vector space isomorphism $\rho : Tv \rightarrow T'v'$ that sends v_i to v'_i . Since Γ and Γ' have same parameters (n, k, a, c) , from (3.1) the matrix representing A with respect to $\{v_i\}_{i=0}^2$ and the matrix representing A' with respect to $\{v'_i\}_{i=0}^2$ are the same as the tridiagonal matrix with entries of the intersection numbers of Γ :

$$\begin{pmatrix} 0 & k & 0 \\ 1 & a & k - a - 1 \\ 0 & c & k - c \end{pmatrix}.$$

Therefore, we have $(\rho A - A' \rho)v_i = 0$ for $0 \leq i \leq 2$. Moreover, we readily check that $(\rho E_j^* - E_j'^* \rho)v_i = 0$ for $0 \leq i, j \leq 2$. The result follows. \blacksquare

Lemma 5.10. *Suppose that the parameters of Γ and Γ' are same, that is, $(n, k, a, c) = (n', k', a', c')$. Let $i \in \{1, 2\}$. Let $v \in E_i^* V$ denote an eigenvector of $E_i^* A E_i^*$ with an eigenvalue λ such that $\langle \mathbf{1}, v \rangle = 0$. Let $v' \in E_i'^* V$ denote an eigenvector of $E_i'^* A' E_i'^*$ with an eigenvalue λ' such that $\langle \mathbf{1}, v' \rangle = 0$. Then $\lambda = \lambda'$ if and only if the T -module Tv and the T' -module $T'v'$ are T -isomorphic.*

Proof. Set $i = 1$. Suppose that $\lambda = \lambda'$. If $\lambda \notin \{\sigma, \tau\}$, then by Lemma 5.2(I)(i) Tv and $T'v'$ have a basis $v, E_2^* A E_1^* v$ and a basis $v', E_2'^* A' E_1'^* v'$, respectively. Abbreviate $w = E_2^* A E_1^* v$ and $w' = E_2'^* A' E_1'^* v'$. Define a vector space isomorphism $\rho : Tv \rightarrow T'v'$ by $\rho(v) = v'$ and $\rho(w) = w'$. By Lemma 5.2(I)(i) and since $\lambda = \lambda'$, the matrix representing A with respect to a basis v, w and the matrix representing A' with respect to a basis v', w' are both equal to the matrix (5.4). From this, it follows that $(\rho A - A' \rho)Tv = 0$. It is clear that $(\rho E_i^* - E_i'^* \rho)Tv = 0$ for $0 \leq i \leq 2$. Thus, Tv and $T'v'$ are T -isomorphic. If $\lambda \in \{\sigma, \tau\}$, use Lemma 5.2(I)(ii) in a similar manner as demonstrated above to get the result. Conversely, if Tv and $T'v'$ are T -isomorphic, the the matrix representing A with respect to a basis v, w must be the same as the matrix representing A' with respect to a basis v', w' . By this and using Lemma 5.2(I), we have $\lambda = \lambda'$. For $i = 2$, use Lemma 5.2(II). The result follows. \blacksquare

We now characterize pseudo-vertex-transitive strongly regular graphs.

Theorem 5.11. *A connected strongly regular graph Γ is pseudo-vertex-transitive if and only if for every pair of vertices $x, y \in X$, the spectrum of the local graph $\Delta(x)$ of Γ at x is equal to the spectrum of the local graph $\Delta(y)$ of Γ at y .*

Proof. If Γ is pseudo-vertex-transitive, by Lemma 4.7 it immediately follows that $\text{Spec}(\Delta(x)) = \text{Spec}(\Delta(y))$. Conversely, for $x, y \in X$ suppose that $\text{Spec}(\Delta(x)) = \text{Spec}(\Delta(y))$. By Lemma 5.5, it follows that $\text{Spec}(\Delta_2(x)) = \text{Spec}_2(\Delta_2(y))$. Using these, we will show that the standard $T(x)$ -module and the standard $T(y)$ -module are T -isomorphic. First, choose an eigenvalue $\lambda \in \Delta(x) = \Delta(y)$. Let $v \in E_1^*(x)V$ denote an eigenvector of $E_1^*(x)AE_1^*(x)$ corresponding to λ , and let $v' \in E_1^*(y)V$ denote an eigenvector of $E_1^*(y)AE_1^*(y)$ corresponding to λ . If $\lambda = a$, then Tv and $T'v'$ are the primary modules of $T(x)$ and $T(y)$, respectively. By Lemma 5.9, $T(x)v$ and $T(y)v'$ are T -isomorphic. If $\lambda \neq a$, we observe that T -modules $T(x)v$ and $T(y)v'$ have both endpoint 1. By Lemma 5.10, $T(x)v$ and $T(y)v'$ are T -isomorphic. Next, choose an eigenvalue $\mu \in \Delta_2(x) = \Delta_2(y)$ such that $\mu \in \{\sigma, \tau\}$. Let $u \in E_2^*(x)V$ denote an eigenvector of $E_2^*(x)AE_2^*(x)$ corresponding to μ , and let $u' \in E_2^*(y)V$ denote an eigenvector of $E_2^*(y)AE_2^*(y)$ corresponding to μ . Observe that both T -modules $T(x)u$ and $T(y)u'$ have endpoint 2 and diameter 0. By Lemma 5.10, $T(x)u$ and $T(y)u'$ are T -isomorphic. By these comments and (5.7), the standard modules for $T(x)$ and $T(y)$ are T -isomorphic. Since x, y is an arbitrary pair of vertices in X , Γ is pseudo-vertex-transitive. ■

In the following, we characterize connected strongly regular graphs which are T -isomorphic.

Theorem 5.12. *Let Γ and Γ' be connected strongly regular graphs with parameters (n, k, a, c) and (n', k', a', c') , respectively. Then Γ and Γ' are T -isomorphic if and only if both (i) $(n, k, a, c) = (n', k', a', c')$ and (ii) for every pair of vertices $x \in X$ and $x' \in X'$ the spectrum of the local graph $\Delta(x)$ of Γ is equal to the spectrum of the local graph $\Delta'(x')$ of Γ' .*

Proof. Suppose that Γ and Γ' are T -isomorphic. By Lemma 4.3, the intersection arrays of Γ and Γ' are the same. From this, it follows $(n, k, a, c) = (n', k', a', c')$. Also, by Lemma 4.8, it follows $\text{Spec}(\Delta(x)) = \text{Spec}(\Delta'(x'))$. Conversely, suppose that both (i) and (ii) hold. We claim that $\text{Spec}(\Delta_2(x)) = \text{Spec}(\Delta'_2(x'))$. We denote by

$$\begin{aligned}\text{Spec}(\Delta_2(x)) &= \{(n-k)^1, \sigma^{\alpha_\sigma}, \tau^{\alpha_\tau}, \lambda_1^{m_1}, \dots, \lambda_s^{m_s}\}, \\ \text{Spec}(\Delta'_2(x')) &= \{(n-k)^1, \sigma^{\beta_\sigma}, \tau^{\beta_\tau}, \lambda_1^{n_1}, \dots, \lambda_t^{n_t}\},\end{aligned}$$

where σ, τ are eigenvalues of Γ from (5.1). Since $\text{Spec}(\Delta(x)) = \text{Spec}(\Delta'(y))$ and by Lemma 5.1, it follows that $s = t$ and $m_i = n_i$ and $\lambda_i = \lambda'_i$ for $1 \leq i \leq s$. The sum of multiplicities of all eigenvalues of $\Delta_2(x)$ is $n-1-k$, and the sum of all eigenvalues of $\Delta_2(x)$ is zero. Moreover, the sum of multiplicities of all eigenvalues of $\Delta'_2(y)$ is $n'-1-k' (= n-1-k)$ and the sum of all eigenvalues of $\Delta'_2(y)$ is zero. From these comments, we have

$$\begin{aligned}\alpha_\sigma + \alpha_\tau &= \beta_\sigma + \beta_\tau, \\ \sigma\alpha_\sigma + \tau\alpha_\tau &= \sigma\beta_\sigma + \tau\beta_\tau.\end{aligned}$$

Solving these equations using the fact $\sigma \neq \tau$, we find that $\alpha_\sigma = \beta_\sigma$ and $\alpha_\tau = \beta_\tau$. The desired claim follows. Now, we show that the standard $T(x)$ -module and the standard $T'(x')$ -module are T -isomorphic. It is similar to the proof of Theorem 5.11; use Lemma 5.9 and Lemma 5.10. Since $x \in X$ and $x' \in X'$ are arbitrary, it follows that Γ and Γ' are T -isomorphic. ■

5.3 Examples

In this subsection, we give some concrete examples of cospectral strongly regular graphs and discuss their pseudo-vertex transitivity.

Example 5.13. Let Γ denote the *Shrikhande graph*, a Cayley graph of $\mathbb{Z}_4 \times \mathbb{Z}_4$ relative to the generating set $\{\pm(1,0), \pm(0,1), \pm(1,1)\}$. Let Γ' denote the *4 × 4-grid graph*, the line graph of $K_{4,4}$. The graphs Γ and Γ' are cospectral strongly regular with parameters $(16, 6, 2, 2)$ and the eigenvalues $k = 6, \sigma = 2, \tau = -2$. Moreover, Γ and Γ' are both vertex-transitive and thus pseudo-vertex-transitive. The following table shows the spectrum of the local graph Δ_G of $G \in \{\Gamma, \Gamma'\}$ and the dimension of the Terwilliger algebra T_G of G .

Strongly regular graph with $(16, 6, 2, 2)$		
graph G	$\text{Spec}(\Delta_G)$	$\dim T_G$
Γ	$\{2^1, 1^2, (-1)^2, (-2)^1\}$	20
Γ'	$\{2^2, (-1)^4\}$	15

By Theorem 5.12, it follows that Γ and Γ' are not T -isomorphic.

Example 5.14 (cf. [28, Example 4.2]). Recall the Johnson graph $\Gamma = J(8, 2)$ with vertex set $X = \binom{\Omega}{2}$, where $\Omega = \{1, 2, \dots, 8\}$. The automorphism group $\text{Aut}(\Gamma)^1$ acts on X transitively, and hence Γ is pseudo-vertex-transitive. We recall the *Chang graphs*. The three Chang graphs $\Gamma', \Gamma'', \Gamma'''$ can be obtained from Γ by Seidel switching with respect to one of the sets (cf. [1, pp.105])

- (i) $\{\{1, 5\}, \{2, 6\}, \{3, 7\}, \{4, 8\}\}$ for Γ' ,
- (ii) $\{\{1, 2\}, \{2, 3\}, \{3, 4\}, \{4, 5\}, \{5, 6\}, \{6, 7\}, \{7, 8\}, \{8, 1\}\}$ for Γ'' ,
- (iii) $\{\{1, 2\}, \{2, 3\}, \{3, 1\}, \{4, 5\}, \{5, 6\}, \{6, 7\}, \{7, 8\}, \{8, 4\}\}$ for Γ''' .

Note that none of three Chang graphs (i)–(iii) is vertex-transitive. For case (i), the action of $\text{Aut}(\Gamma')$ on X has two orbits:

$$U'_1 = \{\{1, 5\}, \{2, 6\}, \{3, 7\}, \{4, 8\}\}, \quad U'_2 = X \setminus U'_1.$$

For case (ii), the action of $\text{Aut}(\Gamma'')$ on X has two orbits:

$$U''_1 = \{\{1, 5\}, \{2, 6\}, \{3, 7\}, \{4, 8\}\}, \quad U''_2 = X \setminus U''_1.$$

For case (iii), the action of $\text{Aut}(\Gamma''')$ on X has three orbits²

$$U'''_1 = \{\{1, 2\}, \{2, 3\}, \{3, 1\}\}, \quad U'''_2 = \{(i, j) \mid 4 \leq i < j \leq 8\}, \quad U'''_3 = X \setminus \{U'''_1 \cup U'''_2\}.$$

We remark that the Johnson graph $J(8, 2)$ and three Chang graphs (i)–(iii) are cospectral strongly regular with parameters $(28, 12, 6, 4)$ and the eigenvalues $k = 12, \sigma = 4, \tau = -2$. In

¹Note that $\text{Aut}(\Gamma) = S_8$.

²In [28, Example 4.2], the orbit U'''_1 was missed.

the following table we display the spectrum of the local graph Δ_G of $G \in \{\Gamma, \Gamma', \Gamma'', \Gamma'''\}$ and the dimension of the Terwilliger algebra T_G of G .

Strongly regular graph with (28, 12, 6, 4)			
G	base vertex x	$\text{Spec}(\Delta_G(x))$	$\dim T_G(x)$
Γ	$x \in X$	$\{6^1, 0^5, 4^1, (-2)^5\}$	16
Γ'	$x \in U'_1$	$\{6^1, 2^3, 0^2, (-2)^6\}$	20
	$x \in U'_2$	$\{6^1, (1 + \sqrt{5})^1, 2^1, 0^3, (1 - \sqrt{5})^1, (-2)^5\}$	27
Γ''	$x \in U''_1$	$\{6^1, (1 + \sqrt{3})^2, 0^2, (1 - \sqrt{3})^2, (-2)^5\}$	23
	$x \in U''_2$	$\{6^1, (1 + \sqrt{3})^1, 2^1, \sqrt{2}^1, 0^1, (1 - \sqrt{3})^1, (-\sqrt{2})^1, (-2)^5\}$	35
Γ'''	$x \in U'''_1$	$\{6^1, 3^1, (\frac{1+\sqrt{5}}{2})^2, (\frac{1-\sqrt{5}}{2})^2, (-1)^1, (-2)^5\}$	27
	$x \in U'''_2$	$\{6^1, (\frac{1+\sqrt{13}}{2})^2, 1^2, (\frac{1-\sqrt{13}}{2})^2, (-2)^5\}$	23
	$x \in U'''_3$	$\{6^1, (1 + \sqrt{3})^1, 2^1, \sqrt{2}^1, 0^1, (1 - \sqrt{3})^1, (-\sqrt{2})^1, (-2)^5\}$	35

From this table and by Proposition 5.11, we find that the three Chang graphs are not pseudo-vertex-transitive. We finish this example with comments.

- (i) For $x \in U'_2$ and $y \in U'''_1$, the Terwilliger algebras $T_{\Gamma'}(x)$ and $T_{\Gamma'''}(y)$ have the same dimension 27, but they are not T -algebra isomorphic since their local spectra are not equal to each other. Note that the dimension sequences of $T_{\Gamma'}(x)$ and $T_{\Gamma'''}(y)$ are equal to $(1, 4, 1, 4)$, and thus they are semisimple algebra isomorphic to each other.
- (ii) In a similar way to (i), for $x \in U''_1$ and $y \in U'''_2$, the Terwilliger algebras $T_{\Gamma''}(x)$ and $T_{\Gamma'''}(y)$ are not T -algebra isomorphic, but their dimension sequences are equal to $(1, 3, 1, 3)$. Thus they are semisimple algebra isomorphic to each other.
- (iii) For $x \in U''_2$ and $y \in U'''_3$, the local graphs $\Delta_{\Gamma''}(x)$ and $\Delta_{\Gamma'''}(y)$ have the same spectrum. Therefore, the Terwilliger algebras $T_{\Gamma''}(x)$ and $T_{\Gamma'''}(y)$ are T -algebra isomorphic.

Example 5.15. A *generalized quadrangle* is an incidence structure such that: (i) any two points are on at most one line, and hence any two lines meet in at most one point, (ii) If p is a point not on a line L , then there is a unique point p' on L such that p and p' are collinear. If every line contains $s + 1$ points, and every point lies on $t + 1$ lines, we say that the generalized quadrangle has order (s, t) , denoted by $GQ(s, t)$.

The *point graph* of a generalized quadrangle is the graph with the points of the quadrangle as its vertices, with two points adjacent if and only if they are collinear. The point graph Γ of a $GQ(s, t)$ is strongly regular with parameters $((s + 1)(t + 1), s(t + 1), s - 1, t + 1)$ and eigenvalues $k = s(t + 1), \sigma = s - 1, \tau = -t - 1$, with respective multiplicities

$$1, \quad \frac{st(s + 1)(t + 1)}{s + t}, \quad \frac{s^2(st + 1)}{s + t}.$$

By construction, Γ is locally a disjoint union of the $(t + 1)$ cliques of size s . Thus the spectrum of a local graph Δ of Γ is determined by s, t :

$$\text{Spec}(\Delta) = \{(s - 1)^{t+1}, -1^{(s-1)(t+1)}\}.$$

From these comments and by Theorem 5.11, it follows that Γ is pseudo-vertex-transitive. Moreover, the point graph Γ' of a $GQ(s', t')$ is T -isomorphic to Γ if and only if $(s, t) = (s', t')$. The dimension of the Terwilliger algebra $T(x)$ of Γ with respect to a vertex x is given as follows.

$$\dim T(x) = \begin{cases} 10, & \text{if } t = s = 1; \\ 15, & \text{if } t = 1, s \neq 1, \text{ or } t \neq 1, s \neq 1, s^2 = t; \\ 11, & \text{if } t \neq 1, s = 1; \\ 16, & \text{if } t \neq 1, s \neq 1, s^2 \neq t. \end{cases}$$

We finish this example with a comment. The isomorphism class of the Terwilliger algebra for the point graph of a $GQ(s, t)$ only depends on s and t , and thus it is pseudo-vertex-transitive. However, there exist generalized quadrangles whose point graphs are not vertex-transitive. For example, there are at least three pairwise nonisomorphic $GQ(q, q^2)$ of which at least one is not vertex-transitive, provided that $q = 2^{2h+1}$, $h \geq 1$; see [21, Page 45].

6 Antipodal double covers

In this section, we recall some preliminaries concerning antipodal distance-regular graphs and tight distance-regular graphs, which will be used in the next sections. Let Γ denote a distance-regular graph with vertex set X and diameter D . The graph Γ is called *antipodal* whenever the vertices at distance D from a given vertex are all at distance D from each other. Note that being at distance D or zero induces an equivalence relation on X . With respect to this equivalence relation, the equivalence classes are called *antipodal classes*. We say Γ is an *antipodal double cover* if the equivalence class has size 2. That is, for each vertex $x \in X$ there is a unique vertex $y \in X$ with $\partial(x, y) = D$. Such a unique vertex is called the *antipode* of x , denoted by \hat{x} . An antipodal double cover with diameter $D \geq 3$ has the intersection array $\{b_0, b_1, \dots, b_{D-1}; c_1, c_2, \dots, c_D\}$, where $b_i = c_{D-i}$ for $0 \leq i \leq D - 1$, and has a dual bipartite Q -polynomial structure (i.e., $q_{1i}^i = 0$ for $0 \leq i \leq D$), provided that Γ is not bipartite; cf. [10, Theorem 1.1].

Lemma 6.1. *Let Γ be a distance-regular antipodal double cover with diameter $D \geq 3$. Let x be a vertex of Γ and \hat{x} the antipode of x . Then W is an irreducible $T(x)$ -module if and only if W is an irreducible $T(\hat{x})$ -module.*

Proof. Let W denote an irreducible $T(x)$ -module. Then W is invariant under the actions of A and $E_j^*(x)$ for $0 \leq j \leq D$. Since Γ is an antipodal double cover, it follows that $E_j^*(x) = E_{D-j}^*(\hat{x})$ for $0 \leq j \leq D$. Thus W is invariant under the actions of A and $E_h^*(\hat{x})$ for $0 \leq h \leq D$. The result follows. ■

We recall the tight distance-regular graphs. Let Γ denote a Q -polynomial distance-regular graph with diameter $D \geq 3$ and eigenvalues $k = \theta_0 > \theta_1 > \dots > \theta_D$. In [17], Jurišić *et al.* proved that the intersection numbers a_1, b_1 satisfy

$$\left(\theta_1 + \frac{k}{a_1 + 1}\right) \left(\theta_D + \frac{k}{a_1 + 1}\right) \geq -\frac{ka_1b_1}{(a_1 + 1)^2}. \quad (6.1)$$

We say Γ is *tight* whenever Γ is not bipartite and equality holds in (6.1). In [13], Go and Terwilliger characterized tight distance-regular graphs using the Terwilliger algebra. Define

$$b^+ := -1 - \frac{b_1}{1 + \theta_D}, \quad b^- := -1 - \frac{b_1}{1 + \theta_1}. \quad (6.2)$$

We say Γ is *tight with respect to x* whenever every irreducible $T(x)$ -module with endpoint 1 is thin with local eigenvalue b^+ or b^- . We state the following lemma.

Lemma 6.2 ([13, Theorem 13.7], [17, Theorem 12.6]). *Let Γ denote a distance-regular graph with diameter $D \geq 3$. The following are equivalent.*

- (i) Γ is tight.
- (ii) For $x \in X$ the local graph of Γ at x is connected strongly regular with eigenvalues a_1, b^+, b^- .
- (iii) Γ is non-bipartite and tight with respect to each vertex.

In [20], Pascasio showed that the non-bipartite antipodal Q -polynomial distance-regular graphs are tight. For instance, the Johnson graph $J(2D, 2)$, the halved $2D$ -cube, the non-bipartite Taylor graphs and the Meixner1 graph are non-bipartite antipodal Q -polynomial distance-regular graphs, and therefore they are tight; see [11, 17].

We recall the following lemmas, which will be used later.

Lemma 6.3 (cf. [13, Theorem 10.1]). *Let Γ be a Q -polynomial distance-regular graph with $D \geq 3$. Fix $x \in X$ and write $T = T(x)$ the Terwilliger algebra of Γ . Let W be a thin irreducible T -module with endpoint 1. Then W has the local eigenvalue b^+ (resp. b^-) if and only if W has dual endpoint 1 (resp. 2).*

Lemma 6.4 (cf. [13, Theorem 10.7]). *Let Γ be a Q -polynomial distance-regular graph with $D \geq 3$. Fix $x \in X$ and write $T = T(x)$ and $E_i^* = E_i^*(x)$ for $0 \leq i \leq D$. Let W be a thin irreducible T -module with endpoint 1 and local eigenvalue $\lambda \in \{b^+, b^-\}$. Then W has dimension $D - 1$. For $0 \leq i \leq D$, E_i^*W is zero if $i \in \{0, D\}$ and has dimension 1 if $i \notin \{0, D\}$. Moreover, $E_iW = 0$ if $i \in \{0, n\}$ and has dimension 1 if $i \notin \{0, n\}$, where $n = 1$ if $\lambda = b^-$; and $n = D$ if $\lambda = b^+$.*

Lemma 6.5 (cf. [13, Theorem 11.1]). *Let Γ be a Q -polynomial distance-regular graph with $D \geq 3$. Fix a vertex x of Γ and write $T = T(x)$. Let W be a thin irreducible T -module with endpoint 1 and local eigenvalue $\lambda \in \{b^+, b^-\}$. Let W' be an irreducible T -module. Then W and W' are isomorphic as T -modules if and only if W' is thin with endpoint 1 and local eigenvalue λ .*

Lemma 6.6 (cf. [5, Corollary 11.5]). *Let Γ be a Q -polynomial antipodal distance-regular graph with $D \geq 3$. Fix a vertex x of Γ and write $T = T(x)$. Let W be an irreducible T -module with diameter d . Recall the scalars $\{a_i(W)\}_{i=0}^d$ from (3.2). Then*

$$a_i(W) = a_{d-i}(W) \quad (0 \leq i \leq d). \quad (6.3)$$

Proof. Apply the result of [5, Corollary 11.5] to the dual bipartite Q -polynomial distance-regular graphs. ■

Remark 6.7. In [5] Caughman showed that every bipartite Q -polynomial distance-regular graph with $D \geq 3$ is thin and dual thin; moreover, for a given irreducible T -module W the isomorphism class of W as T -module is determined by the endpoint and diameter of W . Let Γ denote a dual bipartite (or antipodal) Q -polynomial distance-regular graph with $D \geq 3$. Applying the results of [5] to Γ , we find that every dual bipartite Q -polynomial distance-regular graph with $D \geq 3$ is thin. Furthermore, the isomorphism class of the Terwilliger algebra of Γ does not depend on the base vertex. However, it does not guarantee that Γ is pseudo-vertex-transitive in general because the isomorphism class of the standard T -module of Γ could depend on the base vertex. We take note of [5, Conjecture 15.8] that the multiplicities of all irreducible T -modules of Γ are determined by the parameters s , q , D , and d , not by the base vertex. Therefore, if the conjecture is true, then we can conclude that Γ is pseudo-vertex-transitive.

The following sections discuss pseudo-vertex transitivity for distance-regular antipodal double covers with diameter three and four in detail.

7 Taylor graphs

In this section, we discuss the pseudo-vertex transitivity of Taylor graphs. A *Taylor graph* is a distance-regular antipodal double cover of a complete graph with diameter three, that is, a distance-regular graph with intersection array $\{k, b, 1; 1, b, k\}$, where $b < k - 1$. By the comment below Lemma 6.2, non-bipartite Taylor graphs are tight. We note that tight distance-regular graphs with diameter three are exactly the non-bipartite Taylor graphs [19]. The spectrum of a Taylor graph is

$$\{k^1, \theta_1^f, \theta_2^k, \theta_3^g\},$$

where

$$\theta_1 = \frac{k - 2b - 1 + \sqrt{\mathcal{D}}}{2}, \quad \theta_2 = -1, \quad \theta_3 = \frac{k - 2b - 1 - \sqrt{\mathcal{D}}}{2}, \quad (7.1)$$

and

$$f = \left(\frac{1}{2} - \frac{k - 2b - 1}{2\sqrt{\mathcal{D}}} \right) (k + 1), \quad g = \left(\frac{1}{2} + \frac{k - 2b - 1}{2\sqrt{\mathcal{D}}} \right) (k + 1),$$

where $\mathcal{D} = (k - 2b - 1)^2 + 4k$.

For the rest of this section, we denote by Γ a Taylor graph with vertex set X and the eigenvalues $k = \theta_0 > \theta_1 > \theta_2 > \theta_3$. By the comment below Lemma 6.2, Γ is tight. By Lemma 6.2, for each $x \in$

X the local graph $\Delta = \Delta(x)$ of Γ is connected strongly regular with parameters $(n_\Delta, k_\Delta, a_\Delta, c_\Delta)$, where

$$n_\Delta = k, \quad k_\Delta = k - b - 1, \quad a_\Delta = \frac{2k - 3b - 4}{2}, \quad c_\Delta = \frac{k - b - 1}{2}.$$

From (5.1), the three eigenvalues of Δ are given by

$$k_\Delta, \quad \sigma = \frac{k - 2b - 3 + \sqrt{\mathcal{D}}}{4}, \quad \tau = \frac{k - 2b - 3 - \sqrt{\mathcal{D}}}{4}, \quad (7.2)$$

where $\mathcal{D} = (k - 2b - 1)^2 + 4k$. One can verify that

$$\sigma = -1 - \frac{b}{1 + \theta_3}, \quad \tau = -1 - \frac{b}{1 + \theta_1}. \quad (7.3)$$

Thus, $\sigma = b^+$ and $\tau = b^-$, where b^+ and b^- are from (6.2). By (5.2) the multiplicities m_σ and m_τ of σ and τ , respectively, are given by

$$m_\sigma = \frac{k - 1}{2} - \frac{(k + 1)(k - 2b - 1)}{2\sqrt{\mathcal{D}}}, \quad m_\tau = \frac{k - 1}{2} + \frac{(k + 1)(k - 2b - 1)}{2\sqrt{\mathcal{D}}}. \quad (7.4)$$

By these comments we have the following lemma.

Lemma 7.1. *For each $i \in \{1, 2\}$ and for all $x \in X$, the spectrum of $\Delta_i(x)$ of Γ is determined by the intersection numbers k and b .*

Proof. For the case $i = 1$, it follows from (7.2) and (7.4). For the case $i = 2$, since Γ is an antipodal double cover it follows $\Delta_1(x) = \Delta_1(\hat{x}) = \Delta_2(x)$. The result follows. \blacksquare

By Remark 6.7, Γ is thin. In the following, however, we prove the thinness of Γ in our context.

Lemma 7.2. *Fix a vertex $x \in X$ and let $T = T(x)$ be the Terwilliger algebra of Γ . Let W be a T -module. Then the endpoint of W is either 0 or 1.*

Proof. Suppose that W has the endpoint r and diameter d . Clearly, the possible value for r is 0, 1, 2, or 3. If $r = 2$, then $d = 0$ or $d = 1$. If $d = 0$, then W is a one-dimensional T -module. It implies that an eigenvalue of $E_2^*(x)AE_2^*(x)$ on W is an eigenvalue of Γ . By Lemma 7.1, the eigenvalue of $E_2^*(x)AE_2^*(x)$ is $\lambda \in \{\sigma, \tau\}$ in (7.3). But, $\lambda \notin \{k, \theta_1, \theta_2, \theta_3\}$ in (7.1), the eigenvalues of Γ . If $d = 1$, then $\dim(W) = 2$ and $E_3^*(x)W \neq 0$. Take $0 \neq w \in E_3^*(x)W = E_0^*(\hat{x})W \subseteq W$. Since W is a T -module, it follows $Mw \subseteq W$. Observe that Mw is the primary $T(\hat{x})$ -module, and thus $\dim(Mw) = 4$, a contradiction. If $r = 3$, then diameter of W must be 0. This case is similar to the case of $r = 2$ and $d = 1$, so it is not possible. The result follows. \blacksquare

Lemma 7.3. *Recall $\theta_0 > \theta_1 > \theta_2 > \theta_3$ the eigenvalues of Γ . Fix $x \in X$ and write $T = T(x)$. Let W be an irreducible T -module with endpoint 1 and local eigenvalue $\lambda \in \{\sigma, \tau\}$, where σ, τ are from (7.3). Then the following (i), (ii) and (iii) hold.*

(i) W has diameter 1 and is thin.

(ii) Let $w_0 \in E_1^*W$ be a nonzero vector. Set $w_1 = E_2^*Aw_0$. Then $\{w_0, w_1\}$ is a basis for W . With respect to this basis, the matrix representing A is given by

$$\begin{pmatrix} \lambda & (\lambda - \theta_t)^2 \\ 1 & \lambda \end{pmatrix}, \quad (7.5)$$

where $t = 1$ if $\lambda = \sigma$ and $t = 2$ if $\lambda = \tau$.

(iii) The local eigenvalues σ, τ of W are given by $\sigma = (\theta_1 - \theta_2)/2$ and $\tau = (\theta_2 - \theta_3)/2$.

Proof. (i): For the first assertion, it is similar to the proof of Lemma 7.2. For the second assertion, one knows that Γ is tight. By Lemma 6.2(iii), W is thin.

(ii): Clearly, $\{w_0, w_1\}$ is an orthogonal basis for W . By (3.4), the action of A on w_0 is given by $Aw_0 = a_0(W)w_0 + w_1$. Observe that $a_0(W) = \lambda$. Next, the action of A on w_1 is given by $Aw_1 = x_1(W)w_0 + a_1(W)w_1$, where the scalars $a_1(W)$ and $x_1(W)$ are from (3.2). As Γ is an antipodal double cover, by (6.3) we find $a_1(W) = a_0(W) = \lambda$. Recall the intersection numbers $b_0(W)$ and $c_1(W)$ of W . Referring to [6, Section 9], we find that $a_0(W) + b_0(W) = \theta_t$ and $c_1(W) + a_1(W) = \theta_t$, where t is dual endpoint of W . Using these two equations and the fact that $x_1(W) = b_0(W)c_1(W)$, we find $x_1(W) = (\lambda - \theta_t)^2$. By Lemma 6.3, the dual endpoint of W is $t = 1$ (resp. $t = 2$) if the local eigenvalue λ is σ (resp. τ). By these comments, we obtain the matrix (7.5).

(iii): By (3.3), we have $a_0(W) + a_1(W) = 2\lambda = \theta_t + \theta_{t+1}$, where t is the dual endpoint of W . By Lemma 6.3, the result follows. \blacksquare

Proposition 7.4. *A Taylor graph is thin.*

Proof. By Lemma 7.2 and Lemma 7.3. \blacksquare

We recall the following lemma, which classifies all irreducible T -modules of Γ with endpoint one.

Lemma 7.5 (cf. [13, Theorem 11.1]). *Fix $x \in X$ and write $T = T(x)$. Let W be a thin irreducible T -module with endpoint 1 and local eigenvalue $\lambda \in \{\sigma, \tau\}$, where σ, τ are from (7.2). Let W' denote an irreducible T -module. Then W and W' are isomorphic as T -modules if and only if W' is thin with endpoint 1 and local eigenvalue λ .*

Proof. Apply Lemma 6.5 to Γ . The result immediately follows. \blacksquare

Recall the standard T -module V . Since T is semisimple and using Lemma 7.2, Lemma 7.3, and Lemma 7.5, V decomposes into an orthogonal direct sum of thin irreducible T -modules

$$V = W_0 \oplus \left(\bigoplus_i W_i \right) \oplus \left(\bigoplus_j W'_j \right), \quad (7.6)$$

where we denote by

T -module	dimension	endpoint	diameter	comment
W_0	4	0	3	the primary module
W_i	2	1	1	local eigenvalue σ
W'_j	2	1	1	local eigenvalue τ

where σ and τ are from (7.3).

Theorem 7.6. *A Taylor graph is pseudo-vertex-transitive.*

Proof. For a given vertex $x \in X$, let $T(x)$ be the Terwilliger algebra of Γ . By Lemma 7.1 and Lemma 7.5, the isomorphism class of an irreducible $T(x)$ -module is determined by the parameters k and b . Moreover, by Lemma 7.3 and line (7.4) the multiplicities of irreducible $T(x)$ -modules are determined by k and b . Therefore, the isomorphism class of the standard $T(x)$ -module (7.6) of Γ is completely determined by k and b . The result follows. \blacksquare

Remark 7.7. Taylor graphs are not vertex-transitive in general. For example, there are four different Taylor graphs with intersection array $\{25, 12, 1; 1, 12, 25\}$ of which only one is vertex-transitive; cf. [22].

We finish this section with a comment.

Proposition 7.8. *For each $x \in X$, the dimension of $T(x)$ of Γ is given by*

$$\dim(T(x)) = 24.$$

Proof. Applying Proposition 3.1 to (7.6), the algebra $T(x)$ is isomorphic to

$$\text{Mat}_4(\mathbb{C}) \oplus \text{Mat}_2(\mathbb{C}) \oplus \text{Mat}_2(\mathbb{C}).$$

From (3.5), it follows that $\dim(T(x)) = 4^2 + 2^2 + 2^2 = 24$. \blacksquare

8 Antipodal tight graphs $\text{AT4}(p, q, 2)$

In this section, we consider a distance-regular antipodal double cover with diameter four. We recall some properties of this graph that are needed in this section; for more information, we refer the reader to [16, 18, 19]. Let Γ denote a non-bipartite distance-regular antipodal double cover with diameter four. The intersection array of Γ is parameterized by p, q (cf. [19, Section 5]):

$$\{q(pq + p + q), (q^2 - 1)(p + 1), q(p + q)/2, 1; 1, q(p + q)/2, (q^2 - 1)(p + 1); q(pq + p + q)\}, \quad (8.1)$$

where $p \geq 1, q \geq 2$ are integers. The spectrum of Γ is given by $\{\theta_0^{m_0}, \theta_1^{m_1}, \theta_2^{m_2}, \theta_3^{m_3}, \theta_4^{m_4}\}$, where

$$\theta_0 = q(pq + p + q), \quad \theta_1 = pq + p + q, \quad \theta_2 = p, \quad \theta_3 = -q, \quad \theta_4 = -q^2, \quad (8.2)$$

and $m_0 = 1$,

$$\begin{aligned} m_1 &= \frac{q(pq^2 + q^2 + pq - p)}{p + q}, & m_2 &= \frac{q(pq + p + q)(q^2 - 1)(2q + pq + p)}{(p + q)(p + q^2)}, \\ m_3 &= \frac{(pq^2 + q^2 + pq - p)(pq + p + q)}{p + q}, & m_4 &= \frac{(p + 1)(pq + p + q)(pq^2 + q^2 + pq - p)}{(p + q)(p + q^2)}. \end{aligned}$$

Recall that the graph Γ is Q -polynomial and tight [19]. For each $x \in X$ the local graph $\Delta = \Delta(x)$ of Γ is connected strongly regular with parameters $(n_\Delta, k_\Delta, a_\Delta, c_\Delta)$, where (cf. [19, Section 5])

$$n_\Delta = q(pq + p + q), \quad k_\Delta = p(q + 1), \quad a_\Delta = 2p - q, \quad c_\Delta = p. \quad (8.3)$$

The nontrivial eigenvalues of Δ are $b^+ = p$ and $b^- = -q$ and their multiplicities are

$$m_{b^+} = \frac{(q^2 - 1)(pq + p + q)}{p + q}, \quad m_{b^-} = \frac{pq(q + 1)(p + 1)}{p + q}. \quad (8.4)$$

Let $\text{AT4}(p, q, 2)$ denote an antipodal tight double cover of diameter four with parameters p and q . There are three known examples of an $\text{AT4}(p, q, 2)$, namely the following graphs with array $\{c_1, c_2, c_3, c_4\}$:

- (i) $\text{AT4}(2, 2, 2)$, the Johnson graph $J(8, 4)$ with $\{1, 4, 9, 16\}$;
- (ii) $\text{AT4}(4, 2, 2)$, the half-cube $\frac{1}{2}H(8, 2)$ with $\{1, 6, 15, 28\}$;
- (iii) $\text{AT4}(8, 4, 2)$, the Meixner1 graph with $\{1, 24, 135, 176\}$.

Remark 8.1. In [11, Theorem 1.1], Dickie and Terwilliger gave a family of graphs with array $\{1, \beta\eta, (\beta^2 - 1)(2\eta - \beta + 1), \beta(2\eta + 2\eta\beta - \beta^2)\}$, where $\beta \geq 2$, $\eta \geq 3\beta/4$ are integers and η divides $\beta^2(\beta^2 - 1)/2$. We remark that $\beta\eta$ is an even integer by [18, Corollary 3.2]. With respect to the graph $\text{AT4}(p, q, 2)$, we note that $p = 2\eta - \beta$ and $q = \beta$. Using this and by (8.2), the eigenvalues of $\text{AT4}(p, q, 2)$ are given in terms of β and η as follows.

$$\theta_0 = \beta(2\eta + 2n\beta - \beta^2), \quad \theta_1 = 2\eta + 2n\beta - \beta^2, \quad \theta_2 = 2\eta, \quad \theta_3 = -\beta, \quad \theta_4 = -\beta^2.$$

If $(\beta, \eta) = (2, 2)$ we have $\text{AT4}(2, 2, 2)$, and if $(\beta, \eta) = (2, 3)$ we have $\text{AT4}(4, 2, 2)$; these graphs are all examples known for $\beta = 2$. In addition, if $(\beta, \eta) = (4, 6)$ we have the Meixner1 graph $\text{AT4}(8, 4, 2)$, which is the only known example for $\beta > 2$ so far.

For the rest of this section, we denote by Γ an antipodal tight graph $\text{AT4}(p, q, 2)$ with vertex set X . By Remark 6.7, Γ is thin. In the following, however, we prove the thinness of Γ in our context. Fix $x \in X$ and write $T = T(x)$ the Terwilliger algebra of Γ and $E_i^* = E_i^*(x)$ the i th dual primitive idempotent of Γ for $0 \leq i \leq 4$. We recall a classification of all thin irreducible T -modules of Γ with endpoint 1.

Lemma 8.2. *Let W denote a thin irreducible T -module with endpoint 1 and local eigenvalue $\lambda \in \{p, -q\}$. Then W has diameter 2. If W' is an irreducible T -module, then W and W' are isomorphic as T -modules if and only if W' is thin with endpoint 1 and local eigenvalue λ .*

Proof. Apply Lemma 6.4 and Lemma 6.5. ■

Next, we discuss irreducible T -modules of Γ with endpoint 2.

Lemma 8.3. *If W is an irreducible T -module with endpoint 2, then W has diameter 0 and is thin.*

Proof. Let d denote the diameter of W . Observe that the possible value for d is 0, 1, or 2. If $d = 2$, then $\dim(W) = 3$ and $E_4^*W = E_0^*(\hat{x})W \neq 0$. Take a nonzero vector $w \in E_0^*(\hat{x})W$. Since W is a T -module, we have $Mw \subseteq W$. Observe that Mw is the primary $T(\hat{x})$ -module. It follows $\dim(Mw) = 5$, which is greater than $\dim(W) = 3$, a contradiction. If $d = 1$, then $E_3^*W = E_1^*(\hat{x})W$ is nonzero. It follows that W is an irreducible $T(\hat{x})$ -module with endpoint 1. As Γ is tight, by Lemma 6.2(iii) W is thin and has local eigenvalue $\lambda \in \{p, -q\}$. By Lemma 8.2, we have $d = 2$, a contradiction. By these comments, we have $d = 0$. Since W is an irreducible T -module, the dimension of W is one. The result follows. \blacksquare

Lemma 8.4. *Let W be an irreducible T -module with endpoint 2. Let η be an eigenvalue of $E_2^*AE_2^*$ with an eigenspace W . If W' is another irreducible T -module with endpoint 2, then W and W' are isomorphic as T -modules if and only if W' is an eigenspace associated with the eigenvalue η .*

Proof. Let η' denote an eigenvalue of $E_2^*AE_2^*$ with the eigenspace W' . By Lemma 8.3, W and W' are both one-dimensional. Let ρ denote a vector space isomorphism from W and W' . Apparently, $(E_i^*\rho - \rho E_i^*)W = 0$ for all $0 \leq i \leq 4$. Also, we find that $(A\rho - \rho A)W = 0$ if and only if $\eta = \eta'$. The result follows. \blacksquare

Lemma 8.5. *Recall Γ an antipodal tight graph $AT4(p, q, 2)$. For each vertex $x \in X$ and for each $1 \leq i \leq 4$, the spectrum of i th subconstituent $\Delta_i(x)$ of Γ is determined by the parameters p, q .*

Proof. Fix a vertex x in Γ . Since Γ is tight, $\Delta_1(x) = \Delta_3(x)$ is strongly regular with the parameters $(q(pq+p+q), p(q+1), 2p-q, p)$, and has the nontrivial eigenvalues $p, -q$. It suffices to show that the spectrum of the second subconstituent $\Delta_2 = \Delta_2(x)$ of Γ is determined by p and q . We first assume the Krein parameter $q_{44}^4 = 0$. By [16, Theorem 5.5], Δ_2 is an antipodal distance-regular graph with diameter four, and intersection array and the spectrum of Δ_2 are determined by p and q . Now we assume $q_{44}^4 \neq 0$. As the graph Γ is 1-homogeneous in the sense of Nomura (cf. [17, Theorem 11.7]), Δ_2 is edge-regular. We claim that (i) Δ_2 has at most 7 distinct eigenvalues; and (ii) each eigenvalue and its multiplicity are determined by p and q .

First, we find that a_2 is the eigenvalue of Δ_2 with multiplicity 1. Next, let W denote a thin irreducible T -module with endpoint 1 and local eigenvalue $\lambda \in \{p, -q\}$. By Lemma 8.2, W has diameter 2. Take a nonzero vector $w_0 \in E_1^*W$. Set $w_i = E_i^*Aw_{i-1}$ for $i = 1, 2$. Then $\{w_i\}_{i=0}^2$ is a basis for W , and from (3.4) the action of A on the basis $\{w_i\}_{i=0}^2$ is given by

$$Aw_i = w_{i+1} + a_i(W)w_i + x_i(W)w_{i-1}.$$

Observe that $a_0(W) = \lambda$ and $a_0(W) = a_2(W) = \lambda$ by (6.3). Observe also that $a_1(W)$ is an eigenvalue of $E_2^*AE_2^*$, i.e., an eigenvalue of Δ_2 . By this and using (3.3) and Lemma 6.3, we have

$$a_1(W) = \theta_t + \theta_{t+1} + \theta_{t+2} - 2\lambda,$$

where $t = 1$ if $\lambda = p$ and $t = 2$ if $\lambda = -q$. From this and by (8.2), we find that $a_1(W)$ is determined by p, q . Since each λ yields the scalar $a_1(W)$, multiplicities of λ and $a_1(W)$ are equal to each other. By Lemma 8.2 there are two non-isomorphic T -modules with endpoint 1. It follows that there are two eigenvalues of Δ_2 , which are obtained from irreducible T -modules with endpoint 1.

Finally, let W denote a thin irreducible T -module with endpoint 2. By Lemma 8.3, W is a one-dimensional eigenspace for $E_2^*AE_2^*$. It follows that the corresponding eigenvalue of Δ_2 becomes an eigenvalue of Γ . From this, we find that Δ_2 has four possible eigenvalues $\theta_1, \theta_2, \theta_3, \theta_4$. For each $1 \leq i \leq 4$, let m_i denote the multiplicity of θ_i . Abbreviate $B = E_2^*AE_2^*$. Since Δ_2 is edge-regular and $\text{trace}(B^\ell) = \sum_{i=1}^4 \theta_i^\ell m_i$, $\ell = 0, 1, 2, 3$, we can solve this system of equations for m_i , $1 \leq i \leq 4$, where m_i are determined by p, q . The result follows. ■

We remark that the proof of Lemma 8.5 gives a potential new feasibility condition for an $\text{AT4}(p, q, 2)$.

Proposition 8.6. *An antipodal tight graph $\text{AT4}(p, q, 2)$ is thin.*

Proof. For any vertex x , an irreducible $T(x)$ -module of $\text{AT4}(p, q, 2)$ with endpoint 1 is thin since $\text{AT4}(p, q, 2)$ is tight. In addition, every irreducible $T(x)$ -module of $\text{AT4}(p, q, 2)$ with endpoint 2 is thin by Lemma 8.3. The result follows. ■

Recall the standard T -module V . Since T is semisimple and using Lemma 8.2 and Lemma 8.4, V decomposes into an orthogonal direct sum of thin irreducible T -modules

$$V = W_0 \oplus \left(\bigoplus_i W_i \right) \oplus \left(\bigoplus_j W'_j \right) \oplus \left(\bigoplus_h W''_h \right), \quad (8.5)$$

where we denote by

T -module	dimension	endpoint	diameter	comment
W_0	5	0	4	the primary module
W_i	3	1	2	local eigenvalue p
W'_j	3	1	2	local eigenvalue $-q$
W''_h	1	2	0	eigenspace of $E_2^*AE_2^*$

Theorem 8.7. *An antipodal tight graph $\text{AT4}(p, q, 2)$ is pseudo-vertex-transitive.*

Proof. For a given vertex $x \in X$, let $T(x)$ be the Terwilliger algebra of $\text{AT4}(p, q, 2)$. By Lemma 8.2 and Lemma 8.4, the isomorphism class of an irreducible $T(x)$ -module is determined by the parameters p and q . Moreover, by Lemma 8.5 together with (8.4), the multiplicities of irreducible $T(x)$ -modules are determined by p and q . Therefore, the isomorphism class of the standard $T(x)$ -module (8.5) is completely determined by p and q . The result follows. ■

We finish this section with a comment.

Proposition 8.8. *For each $x \in X$ the dimension of $T(x)$ of $\text{AT4}(p, q, 2)$ is given by*

$$\dim(T(x)) = \ell + 43,$$

where $\ell(\leq 7)$ is the number of distinct eigenvalues of $\Delta_2(x)$ except for the eigenvalue a_2 .

Proof. Applying Proposition 3.1 to (8.5), the algebra $T(x)$ is isomorphic to

$$\text{Mat}_5(\mathbb{C}) \oplus \text{Mat}_3(\mathbb{C}) \oplus \text{Mat}_3(\mathbb{C}) \oplus (\text{Mat}_1(\mathbb{C})^{\oplus \ell}),$$

where ℓ is the number of distinct eigenvalues of $\Delta_2(x)$ except for the eigenvalue a_2 . From (3.5), it follows that

$$\dim(T(x)) = \ell(1)^2 + 3^2 + 3^2 + 5^2 = \ell + 43.$$

■

9 Concluding remarks

In this paper, we discussed pseudo-vertex transitivity of Q -polynomial distance-regular graphs with small diameter $D \in \{2, 3, 4\}$. We briefly summarize the results. For the case of $D = 2$, we showed that the local spectrum of a strongly regular graph characterizes its pseudo-vertex transitivity. In addition, we showed that two strongly regular graphs are T -isomorphic if and only if they have the same parameters and their local spectra are equal to each other. For the case of $D = 3$, we were concerned with a Taylor graph Γ with intersection array $\{k, b, 1; 1, b, k\}$. We showed that Γ is thin and that the isomorphism classes of the standard module of the Terwilliger algebra $T(x)$ of Γ only depend on the parameters k, b , and hence Γ is pseudo-vertex-transitive. For the case of $D = 4$, we were concerned with an antipodal tight graph $\text{AT4}(p, q, 2)$. We proved that $\text{AT4}(p, q, 2)$ is thin and that the isomorphism classes of the standard module of Terwilliger algebra $T(x)$ of $\text{AT4}(p, q, 2)$ only depend on the parameters p, q , and hence $\text{AT4}(p, q, 2)$ is pseudo-vertex-transitive. In both cases, $D = 3$ and $D = 4$, the thin property of distance-regular antipodal double covers played an important role in showing their pseudo-vertex transitivity. We anticipate that the thinness of a distance-regular graph still plays a significant role in determining pseudo-vertex transitivity for $D \geq 3$. We give the following conjecture.

Conjecture 9.1. A thin Q -polynomial distance-regular graph with diameter $D \geq 3$ is pseudo-vertex-transitive.

As we saw in Problem 1.1, the Ito-Koolen problem asks to classify all thin pseudo-vertex-transitive Q -polynomial distance-regular graphs with large enough diameter. As a first step towards solving the Ito-Koolen problem, Tan *et al.* [23] found an upper bound of the intersection number c_2 and parameter α , respectively, for a thin Q -polynomial distance-regular graph with classical parameters (D, b, α, β) . Each of upper bound is given by a function in b , where $b = b_1/(\theta_1 + 1)$ and where θ_1 is the second largest eigenvalue of Γ . We give some suggestions for further research.

Problem 9.2. With reference to the result of [23], improve the upper bound on c_2 .

Problem 9.3. Assume Γ has the same parameters as a Grassmann graph $J_q(2D + 1, D)$ and that Γ is thin. Prove or disprove that Γ and $J_q(2D + 1, D)$ are T -isomorphic. We remark that Liang *et al.* [24] have proved this problem for large enough diameter D .

Acknowledgements

The authors would like to express many thanks to the anonymous referees for their corrections and valuable comments. J.-H. Lee would like to thank Paul Terwilliger for several valuable conversations and comments. J.H. Koolen is partially supported by the National Natural Science Foundation of China (No. 12071454), Anhui Initiative in Quantum Information Technologies (No. AHY150000) and the National Key R and D Program of China (No. 2020YFA0713100). Y.-Y Tan is supported by the National Natural Science Foundation of China (No. 11801007, 12171002) and Natural Science Foundation of Anhui Province (No. 1808085MA17) and the foundation of Anhui Jianzhu University (No. 2018QD22).

References

- [1] A. E. Brouwer, A. M. Cohen and A. Neumaier, Distance-Regular Graphs, Springer-Verlag, Berlin, 1989.
- [2] S. Bang, T. Fujisaki and J. H. Koolen, The spectra of the local graphs of the twisted Grassmann graphs, *European J. Combin.* **30**(3) (2009) 638-654.
- [3] E. Bannai and T. Ito, Algebraic Combinatorics I: Association Schemes, Benjamin-Cummings, Menlo Park, 1984.
- [4] J. S. Caughman IV, Spectra of bipartite P - and Q -polynomial association schemes, *Graphs Comb.*, **14** (1998), 321–343.
- [5] J. S. Caughman IV, The Terwilliger algebra of bipartite P - and Q - polynomial schemes, *Discrete Math.*, **196** (1999), 65–95.
- [6] D. R. Cerzo, Structure of thin irreducible modules of a Q -polynomial distance- regular graph, *Linear Algebra Appl.* **433** (2010), 1573–1613.
- [7] C. W. Curtis and I. Reiner, Representation Theory of Finite Groups and Associative Algebras, Pure and Applied Mathematics, Vol. XI, Interscience Publishers, a division of John Wiley & Sons, New York-London, 1962.
- [8] E. R. van Dam and J. H. Koolen, A new family of distance-regular graphs with unbounded diameter, *Invent. Math.* **162** (2005) 189–193.
- [9] E. R. van Dam, J. H. Koolen and H. Tanaka, Distance-regular graphs, *Electron. J. Combin.* (2016) #DS22.
- [10] G. Dickie, Twice Q -polynomial distance-regular graphs are thin, *European J. Combin.* **16** (1995) 555-560.
- [11] G. Dickie and P. Terwilliger, Dual bipartite Q -polynomial distance-regular graphs, *European J. Combin.* **17** (1996) 613–623.

- [12] E. Egge, A generalization of the Terwilliger algebra, *J. Algebra* **233** (2000), 213–252.
- [13] J. T. Go and P. Terwilliger, Tight distance-regular graphs and the subconstituent algebra, *European J. Combin.* **23** (2002), no. 7, 793–816.
- [14] C. D. Godsil and G. Royle, *Algebraic Graph Theory*, Springer-Verlag, Berlin, 2001.
- [15] A. Jurišić, Antipodal covers, Thesis (Ph.D.), University of Waterloo (Canada). 1995. 163 pp.
- [16] A. Jurišić, AT4 family and 2-homogeneous graphs, *Discrete Math.*, **264** (2003), 127–148.
- [17] A. Jurišić, J. H. Koolen and P. Terwilliger, Tight distance-regular graphs, *J. Algebra Combin.* **12(2)**(2000) 163-197.
- [18] A. Jurišić and J. H. Koolen, Nonexistence of some Antipodal Distance-regular Graphs of Diameter Four, *European J. Combin.* **21** (2000), no. 8, 1039–1046.
- [19] A. Jurišić and J. H. Koolen, Krein parameters and antipodal tight graphs with diameter 3 and 4, *Discrete Math.*, **244**(1–3) (2002) 181–202.
- [20] A. A. Pascasio, Tight distance-regular graphs and the Q -polynomial property, *Graphs Combin.* **17** (2001), 149–169.
- [21] S. E. Payne and J. A. Thas, *Finite generalized quadrangles: Second Edition*, EMS Series of Lectures in Mathematics, European Mathematical Society (EMS), Zürich, 2009.
- [22] E. Spence, Regular Two-graphs, <http://www.maths.gla.ac.uk/~es/twograph/reg2Graph.php>
- [23] Y.-Y Tan, J. H. Koolen, M.-Y Cao and J. Park, Thin Q -polynomial distance-regular graphs have bounded c_2 , *preprint*.
- [24] X. Liang, Y.-Y Tan and J. H. Koolen, Thin distance-regular graphs with classical parameters $(D, q, q, \frac{q^t-1}{q-1})$ with $t > D$ are the Grassmann graphs, *submitted*.
- [25] H. Tanaka and T. Wang, The Terwilliger algebra of the twisted Grassmann graph: the thin case, *Electron. J. Combin.* **28** (2020) #P4.15 (22pp.)
- [26] P. Terwilliger, The subconstituent algebra of an association scheme I, *J. Algebra Combin.* **1(4)**(1992) 363-388.
- [27] P. Terwilliger, The subconstituent algebra of an association scheme III, *J. Algebra Combin.* **2(2)**(1993) 177-210.
- [28] M. Tomiyama and N. Yamazaki, The subconstituent algebra of a strongly regular graph, *Kyushu J. Math.* **48(2)**(1994) 323-334.