

# Unified Bernoulli-Euler polynomials of Apostol type

Hacène Belbachir, Yahia Djemmada, Slimane  
Hadj-Brahim

Received: date / Accepted: date

**Abstract** The object of this paper is to introduce and study properties of unified Apostol-Bernoulli and Apostol-Euler polynomials noted by  $\{\mathfrak{B}_n(x; \lambda; \mu)\}_{n \geq 0}$ . We study some arithmetic properties of  $\{\mathfrak{B}_n(x; \lambda; \mu)\}_{n \geq 0}$  as their connection to Apostol-Euler polynomials and Apostol-Bernoulli polynomials. Also, we give derivation and integration representations of  $\{\mathfrak{B}_n(x; \lambda; \mu)\}_{n \geq 0}$ . Finally, we use the umbral calculus approach to deduce symmetric identities.

**Keywords** Euler polynomials · Bernoulli polynomials · Apostol-Bernoulli and Apostol-Euler polynomials · generating function.

**Mathematics Subject Classification (2010)** 11B68 · 11B83 · 11C08 · 11C20.

## 1 Introduction

The Bernoulli  $\{B_n(x)\}_{n \geq 0}$  and the Euler  $\{E_n(x)\}_{n \geq 0}$  polynomials respectively are generated by the following power series (see [4, 6]):

$$\frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} \quad (|t| < 2\pi)$$

and

$$\frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \quad (|t| < \pi).$$

As a particular case, for  $x = 0$ , we denote  $B_n := B_n(0)$  and  $E_n := E_n(0)$ , which are called the Bernoulli and the Euler numbers respectively. They have numerous important applications in various fields of mathematics, as number theory, analysis and combinatorics.

---

H. BELBACHIR  
USTHB, Faculty of Mathematics, RECITS Laboratory P. Box 32,  
El-Alia, 16111 Bab-Ezzouar, Algiers, Algeria.  
E-mail: hacenebelbachir@gmail.com or hbelbachir@usthb.dz

Y. DJEMMADA (corresponding author)  
USTHB, Faculty of Mathematics, RECITS Laboratory P. Box 32,  
El-Alia, 16111 Bab-Ezzouar, Algiers, Algeria.  
E-mail: ydjemmada@usthb.dz or yahia.djem@gmail.com

S. HADJ-BRAHIM  
USTHB, Faculty of Mathematics, RECITS Laboratory P. Box 32,  
El-Alia, 16111 Bab-Ezzouar, Algiers, Algeria.  
E-mail: bhadj@usthb.dz

Apostol [1] introduced and investigated the extended form of the classical Bernoulli polynomials and numbers, which are known as the Apostol-Bernoulli polynomials and numbers. The Apostol-Euler and the Apostol-Genocchi polynomials were introduced by Srivastava [12]. Belbachir et al. [2, 3] proposed a new family of polynomials called Euler-Genocchi polynomials and studied their properties like linear recurrences and difference equations using a determinantal approach and generating function.

## 2 Determinantal representation of the Bernoulli-Euler polynomials of Apostol type

According to [9], the Apostol-Bernoulli polynomials  $\{\mathfrak{B}_n(x; \lambda)\}_{n \geq 0}$  and the Apostol-Euler polynomials  $\{\mathfrak{E}_n(x; \lambda)\}_{n \geq 0}$  are generated by the following power series:

$$\frac{t}{\lambda e^t - 1} e^{xt} = \sum_{n=0}^{\infty} \mathfrak{B}_n(x; \lambda) \frac{t^n}{n!} \quad (|t + \ln \lambda| < 2\pi, \lambda \in \mathbb{R}_+^*) \quad (1)$$

and

$$\frac{2}{\lambda e^t + 1} e^{xt} = \sum_{n=0}^{\infty} \mathfrak{E}_n(x; \lambda) \frac{t^n}{n!} \quad (|t + \ln \lambda| < \pi, \lambda \in \mathbb{R}_+^*), \quad (2)$$

The Apostol-Bernoulli numbers  $\mathfrak{B}_n(\lambda)$  and the Apostol-Euler numbers  $\mathfrak{E}_n(\lambda)$  are given by  $\mathfrak{B}_n(\lambda) = \mathfrak{B}_n(0; \lambda)$  and  $\mathfrak{E}_n(\lambda) = \mathfrak{E}_n(0; \lambda)$ .

Letting

$$T(x, \lambda, t) = \frac{2}{\lambda e^t + 1} e^{xt} \times \frac{t}{\lambda e^t - 1} e^{xt} = \frac{2t}{\lambda^2 e^{2t} - 1} e^{2xt}.$$

Taking into account the right hand side of (1) and (2), a direct computation gives

$$\lambda^2 T(x+1, \lambda, t) - T(x, \lambda, t) = \sum_{n=0}^{\infty} \left\{ \sum_{k=0}^n \binom{n}{k} [\lambda^2 \mathfrak{B}_{n-k}(x, \lambda) \mathfrak{E}_k(x, \lambda) - \mathfrak{B}_{n-k}(x, \lambda) \mathfrak{E}_k(x, \lambda)] \right\} \frac{t^n}{n!}.$$

On the other hand, we have

$$\lambda^2 T(x+1, \lambda, t) - T(x, \lambda, t) = \lambda^2 \frac{2t}{\lambda^2 e^{2t} - 1} e^{2(x+1)t} - \frac{2t}{\lambda^2 e^{2t} - 1} e^{2xt} = 2te^{2xt} = \sum_{n=0}^{+\infty} n 2^n x^{n-1} \frac{t^n}{n!}.$$

Comparing the two expansions of  $\lambda^2 T(x+1, \lambda, t) - T(x, \lambda, t)$ , we formulate the next result.

**Theorem 1** *Let  $x$  be a real number and  $n$  an integer. Then*

$$x^n = \sum_{k=0}^{n+1} A_{n,k} \times \Delta_{n+1-k,k}(x, \lambda), \quad (3)$$

$$\text{where } A_{n,k} = \frac{1}{2^{n+1}(n+1)} \binom{n+1}{k} \text{ and } \Delta_{n,k}(x, \lambda) = \begin{vmatrix} \lambda \mathfrak{B}_n(x+1, \lambda) & \mathfrak{E}_k(x, \lambda) \\ \mathfrak{B}_n(x, \lambda) & \lambda \mathfrak{E}_k(x+1, \lambda) \end{vmatrix}.$$

In particular, taking  $\lambda = 1$  in (3), we get the following result in terms of the Bernoulli and the Euler polynomials.

**Corollary 1** [3] *Let  $x$  be a real number and an integer  $n \geq 0$ , we have*

$$x^n = \frac{1}{2^{n+1}(n+1)} \sum_{k=0}^{n+1} \binom{n+1}{k} \begin{vmatrix} B_{n-(k-1)}(x+1) & E_k(x) \\ B_{n-(k-1)}(x) & E_k(x+1) \end{vmatrix}. \quad (4)$$

### 3 Unified Bernoulli-Euler polynomials of Apostol type

In this section, we give a definition of the unified Bernoulli-Euler polynomials of Apostol type and study their properties using power series.

**Definition 1** Let  $\lambda \in \mathbb{R}_+^*$  and  $\mu \in \mathbb{R}_+ - \{1\}$ , we define the unified Bernoulli-Euler polynomials of Apostol type  $\mathfrak{V}_n(x; \lambda; \mu)$  by the following power series:

$$\frac{2 - \mu + \frac{\mu}{2}t}{\lambda e^t + (1 - \mu)} e^{xt} = \sum_{n \geq 0} \mathfrak{V}_n(x; \lambda; \mu) \frac{t^n}{n!}, \tag{5}$$

where

$$\begin{cases} \left| \ln \left( \frac{\lambda}{1-\mu} \right) + t \right| < 2\pi, \text{ for } 0 \leq \mu < 1; \\ \left| \ln \left( \frac{\lambda}{\mu-1} \right) + t \right| < \pi, \text{ otherwise.} \end{cases}$$

Furthermore, the unified Bernoulli-Euler numbers of Apostol type, denoted  $\mathfrak{V}_n(\lambda; \mu)$ , are given by

$$\mathfrak{V}_n(\lambda; \mu) := \mathfrak{V}_n(0; \lambda; \mu). \tag{6}$$

We summarize in the following table some special polynomials related to this extension.

| Parameters             | Generating functions   | Polynomials                   |
|------------------------|--|-------------------------------|
| $\mu = 0, \lambda = 1$ | $\frac{2}{e^t + 1} e^{xt} = \sum_{n \geq 0} E_n(x) \frac{t^n}{n!}, \quad  t  < \pi$  | Euler polynomials             |
| $\mu = 2, \lambda = 1$ | $\frac{t}{e^t - 1} e^{xt} = \sum_{n \geq 0} B_n(x) \frac{t^n}{n!}, \quad  t  < 2\pi$   | Bernoulli polynomials         |
| $\mu = 2$              | $\frac{t}{\lambda e^t - 1} e^{xt} = \sum_{n \geq 0} \mathfrak{B}_n(x; \lambda) \frac{t^n}{n!}, \quad  t + \ln \lambda  < 2\pi$ | Apostol-Bernoulli polynomials |
| $\mu = 0$              | $\frac{2}{\lambda e^t + 1} e^{xt} = \sum_{n \geq 0} \mathfrak{E}_n(x; \lambda) \frac{t^n}{n!}, \quad  t + \ln \lambda  < \pi$  | Apostol-Euler polynomials     |

We list some properties of the unified Bernoulli-Euler polynomials of Apostol type using generating function approach.

**Theorem 2** Let  $n$  be nonnegative integer, we have

$$\mathfrak{V}_n(x + y; \lambda; \mu) = \sum_{k=0}^n \binom{n}{k} \mathfrak{V}_k(x; \lambda; \mu) y^{n-k}. \tag{7}$$

In particular, for  $x := 0$  and  $y := x$ , the above relation becomes

$$\mathfrak{V}_n(x; \lambda; \mu) = \sum_{k=0}^n \binom{n}{k} \mathfrak{V}_k(\lambda; \mu) x^{n-k}. \tag{8}$$

*Proof* We establish the power series defined in (5) for  $\mathfrak{V}_n(x + y; \lambda; \mu)$ , we have

$$\sum_{n \geq 0} \mathfrak{V}_n(x + y; \lambda; \mu) \frac{t^n}{n!} = \left( \frac{2 - \mu + \frac{\mu}{2}t}{\lambda e^t + (1 - \mu)} \right) e^{(x+y)t} = \sum_{n \geq 0} \sum_{k \geq 0} \mathfrak{V}_n(x; \lambda; \mu) y^k \frac{t^{n+k}}{n!k!}.$$

Applying the product series and then comparing the coefficients of  $t^n$  on both sides, we obtain Identity (7).

**Remark 1** Expression (7) allows us to obtain  $\mathfrak{V}_n(\lambda; \mu)$  the unified Bernoulli-Euler numbers of Apostol type in terms of the unified Bernoulli-Euler polynomials of Apostol type. Indeed, it suffices to replace  $y$  by  $-x$  in Formula (7), we get the following expression:

$$\mathfrak{V}_n(\lambda; \mu) = \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} \mathfrak{V}_k(x; \lambda; \mu) x^{n-k}.$$

As a first consequence of Theorem 1, we show that the unified Bernoulli-Euler polynomials of Apostol type,  $\{\mathfrak{V}_n(x, \lambda; \mu)\}_{n \geq 0}$  given by the power series in (5), can be expressed in terms of the Apostol-Bernoulli and the Apostol-Euler polynomials. That is, by a straightforward calculation, the substitution of  $x^n$  given by (3) in Expression (8) allows us to obtain the following formula:

**Proposition 1** Let  $n, k$  and  $j$  be three integers, it holds that

$$\mathfrak{V}_n(x; \lambda; \mu) = \sum_{k=0}^n \binom{n}{k} \mathfrak{V}_{n-k}(\lambda; \mu) \sum_{j=0}^{k+1} A_{k,j} \times \Delta_{k+1-j,j}(x, \lambda).$$

#### 4 Generalized Raabe Theorem

In this section, we give an extension of Raab Theorem for the unified Bernoulli-Euler polynomials of Apostol type.

**Theorem 3** Let  $r$  and  $m$  be nonnegative integers with  $m$  odd, for  $\lambda = 1 - \mu$  and  $\mu \neq 1$ , we have

$$\sum_{k=0}^{m-1} (-1)^k \mathfrak{V}_n\left(\frac{x+k}{m}; 1-\mu; \mu\right) = \frac{1-m}{m^n} \left(\frac{\mu-2}{2(\mu-1)}\right) E_n(x) + \frac{1}{m^{n-1}} \mathfrak{V}_n(x; 1-\mu; \mu).$$

*Proof* It follows from (5) that

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{k=0}^{m-1} (-1)^k \mathfrak{V}_n\left(\frac{x+k}{m}; 1-\mu; \mu\right) \frac{t^n}{n!} \\ &= \sum_{k=0}^{m-1} (-1)^k \sum_{n=0}^{\infty} \mathfrak{V}_n\left(\frac{x+k}{m}; 1-\mu; \mu\right) \frac{t^n}{n!} \\ &= \sum_{k=0}^{m-1} (-1)^k \frac{2-\mu+\frac{\mu}{2}t}{(1-\mu)e^t+(1-\mu)} e^{(x+k)t/m} \\ &= \frac{2-\mu+\frac{\mu}{2}t}{(1-\mu)(e^t+1)} e^{xt/m} \frac{[1-(-e^{t/m})^m]}{1+e^{t/m}} \\ &= \frac{1}{(1-\mu)} \left(\frac{2-\mu+\frac{\mu}{2}t}{e^{t/m}+1}\right) e^{xt/m} \\ &= \frac{1-m}{2} \left(\frac{2-\mu}{1-\mu}\right) \left(\frac{2}{e^{t/m}+1}\right) e^{xt/m} + \frac{m}{1-\mu} \left(\frac{2-\mu+\frac{\mu t}{2m}}{e^{t/m}+1}\right) e^{xt/m} \\ &= \frac{1-m}{2} \left(\frac{2-\mu}{1-\mu}\right) \sum_{n=0}^{\infty} E_n(x) \frac{(t/m)^n}{n!} + \sum_{n=0}^{\infty} \frac{1}{m^{n-1}} \mathfrak{V}_n(x; 1-\mu; \mu) \frac{t^n}{n!}. \end{aligned}$$

By simple manipulations and equating the coefficients of  $t^n$  on both sides, we get the result.

As a consequence of Theorem 3, for  $\mu = 2$  and  $\mu = 0$  respectively, we have a multiplication Theorem for Euler and Bernoulli polynomials proved by Raabe in [10], as specified by Kargin and Kurt [8]. They are given as follows:

$$\sum_{k=0}^{m-1} B_n \left( \frac{x+k}{m} \right) = \frac{1}{m^{n-1}} B_n(x)$$

and

$$\sum_{k=0}^{m-1} (-1)^k E_n \left( \frac{x+k}{m} \right) = \frac{1}{m^n} E_n(x).$$

## 5 Some explicit formulas

In this section, we give some explicit formulas of the unified Bernoulli-Euler polynomials of Apostol type.

**Theorem 4** For  $\lambda \in \mathbb{R}_+^*$  and  $\mu \in \mathbb{R}_+ - \{1\}$ , it holds that

$$\mathfrak{V}_n(x; \lambda; \mu) = \frac{1}{2(\mu-1)} \left[ (\mu-2) \mathfrak{E}_n \left( x; \frac{\lambda}{1-\mu} \right) - \frac{\mu n}{2} \mathfrak{E}_{n-1} \left( x; \frac{\lambda}{1-\mu} \right) \right] \quad (n \in \mathbb{N}). \quad (9)$$

*Proof* We can reformulate (5) as follows

$$\begin{aligned} \sum_{n=0}^{\infty} \mathfrak{V}_n(x; \lambda; \mu) \frac{t^n}{n!} &= \left( \frac{1}{2(\mu-1)} \right) \left( (\mu-2) - \frac{\mu}{2} t \right) \left( \frac{2}{1 + \frac{\lambda}{1-\mu} e^t} \right) e^{xt} \\ &= \frac{1}{2(\mu-1)} \left[ (\mu-2) \sum_{n=0}^{\infty} \mathfrak{E}_n \left( x; \frac{\lambda}{1-\mu} \right) \frac{t^n}{n!} - \frac{\mu}{2} \sum_{n=1}^{\infty} \mathfrak{E}_n \left( x; \frac{\lambda}{1-\mu} \right) \frac{t^{n+1}}{n!} \right] \\ &= \frac{1}{2(\mu-1)} \sum_{n=1}^{\infty} \left[ (\mu-2) \mathfrak{E}_n \left( x; \frac{\lambda}{1-\mu} \right) - \frac{\mu n}{2} \mathfrak{E}_{n-1} \left( x; \frac{\lambda}{1-\mu} \right) \right] \frac{t^n}{n!}. \end{aligned}$$

Equating the coefficients of  $\frac{t^n}{n!}$  on both sides, we obtain Identity (9).

Here, we give an explicit formula as a dual convex combination of classical Bernoulli and Euler polynomials of Apostol type.

**Theorem 5** Let  $n$  be nonnegative integer and a real number  $\mu \neq 1$ , we have

$$\mathfrak{V}_n(x; \lambda; \mu) = \frac{1}{1-\mu} \left[ \left( 1 - \frac{\mu}{2} \right) \mathfrak{E}_n \left( x; \frac{\lambda}{1-\mu} \right) - \frac{\mu}{2} \mathfrak{B}_n \left( x; \frac{\lambda}{\mu-1} \right) \right]. \quad (10)$$

*Proof* From (5), we have

$$\sum_{n=0}^{\infty} \mathfrak{V}_n(x; \lambda; \mu) \frac{t^n}{n!} = \frac{2-\mu + \frac{\mu}{2}t}{\lambda e^t + (1-\mu)} e^{xt} = \frac{2-\mu}{2(1-\mu)} \frac{2}{\frac{\lambda}{1-\mu} e^t + 1} e^{xt} + \frac{\mu}{2(\mu-1)} \frac{t}{\frac{\lambda}{\mu-1} e^t - 1} e^{xt}.$$

Using (1) and (2) leads to get (10).

**Theorem 6** For  $\mu \neq 1$  and  $n \geq 1$ , the following formula holds:

$$(\mu-1) \sum_{k=1}^n \binom{n}{k} \Upsilon_{n-k,k} \left( \frac{x}{2}, \lambda, \mu \right) + \left( \frac{\mu}{2} - 1 \right) \Delta_{n,0} \left( \frac{x}{2}, \frac{\lambda}{1-\mu} \right) = n(\mu-2)x^{n-1} - n(n-1)\mu x^{n-2},$$

where  $\Upsilon_{n,k}(x, \lambda, \mu) = \left| \begin{array}{cc} \frac{\lambda}{1-\mu} \mathfrak{B}_n(x+1, \lambda) & \mathfrak{V}_k(x; \lambda; \mu) \\ \mathfrak{B}_n(x, \lambda) & \frac{\lambda}{1-\mu} \mathfrak{V}_k(x+1; \lambda; \mu) \end{array} \right|$ .

*Proof* From Theorem 4, we have

$$\begin{aligned} & \sum_{k=1}^n \binom{n}{k} \Upsilon_{n-k,k}(x, \lambda, \mu) \\ &= \frac{1}{2(\mu-1)} \left\{ (\mu-2) \sum_{k=1}^n \binom{n}{k} \Delta_{n-k,k} \left( x, \frac{\lambda}{1-\mu} \right) - \mu \sum_{k=1}^n k \binom{n}{k} \Delta_{n-k,k-1} \left( x, \frac{\lambda}{1-\mu} \right) \right\} \\ &= \frac{1}{2(\mu-1)} \left\{ (\mu-2) \sum_{k=1}^n \binom{n}{k} \Delta_{n-k,k} \left( x, \frac{\lambda}{1-\mu} \right) - \mu n \sum_{k=1}^n \binom{n-1}{k-1} \Delta_{n-k,k-1} \left( x, \frac{\lambda}{1-\mu} \right) \right\}. \end{aligned}$$

Applying Theorem 1 and a straightforward computation, we obtain

$$\begin{aligned} \sum_{k=1}^n \binom{n}{k} \Upsilon_{n-k,k}(x, \lambda, \mu) &= \frac{1}{\mu-1} \left\{ n(\mu-2)(2x)^{n-1} - n(n-1)\mu(2x)^{n-2} \right\} \\ &\quad - \frac{\mu-2}{2(\mu-1)} \Delta_{n,0} \left( x, \frac{\lambda}{1-\mu} \right). \end{aligned}$$

Multiplying both sides by  $(\mu-1)$ , we get the desired identity.

## 6 Derivation and integration representations of unified Bernoulli-Euler polynomials of Apostol type

In this section, we present derivation and integration representations for the unified Bernoulli-Euler polynomials of Apostol type.

**Theorem 7** *Let  $l, n$  be two nonnegative integers. Then*

$$\frac{d^l}{dx^l} \mathfrak{V}_n(x; \lambda; \mu) = (n)_l \mathfrak{V}_{n-l}(x; \lambda; \mu), \quad (11)$$

$$\int_x^y \mathfrak{V}_n(z; \lambda; \mu) dz = \frac{1}{(n+1)} [\mathfrak{V}_{n+1}(y; \lambda; \mu) - \mathfrak{V}_{n+1}(x; \lambda; \mu)], \quad (12)$$

where  $(x)_n := x(x-1)\cdots(x-n+1)$  with  $(x)_0 = 1$ .

*Proof* The assertion (11) follows from (5) by successive differentiations with respect to  $x$  and then uses the induction principle on  $l$ . Furthermore, taking  $l = 1$  in (11) and integrating both sides of the resulting equation with respect to  $z$  over the interval  $[x, y]$ , ( $y > x$ ), we obtain the Integral Formula (12).

**Remark 2** *Setting  $\lambda = 1, \mu = 2$  in (11) and (12), we obtain known results due to Luo et al. [9].*

**Corollary 2** *Let  $n$  be a nonnegative integer. Then*

$$\int_x^{x+y} \mathfrak{V}_n(z; \lambda; \mu) dz = \frac{1}{(n+1)} \sum_{k=0}^n \binom{n}{k} \mathfrak{V}_k(x; \lambda; \mu) y^{n-(k-1)}. \quad (13)$$

*Proof* Replacing  $y$  by  $x+y$  in the Integral Formula (12) and using Formula (7), by successive calculations, we obtain the Integral Formula (13).

**Theorem 8** *For  $\mu \in \mathbb{R}_+^* - \{1, 2\}$  and  $n$  nonnegative integer, the following formula holds:*

$$\begin{aligned} \mathfrak{V}_{n+1}(x; \lambda; \mu) - x \mathfrak{V}_n(x; \lambda; \mu) &= \frac{1}{2-\mu} \sum_{i=0}^n \binom{n}{i} (n-i)! \\ &\times \left( \frac{\mu}{2(\mu-2)} \right)^{n-i} \left[ \frac{\mu}{2} \mathfrak{V}_i(x; \lambda; \mu) - \lambda \sum_{k=0}^i \binom{i}{k} \mathfrak{V}_k(\lambda; \mu) \mathfrak{V}_{i-k}(x+1; \lambda; \mu) \right]. \end{aligned} \quad (14)$$

*Proof* Differentiating both sides of (5) with respect to  $t$ , we express the factors  $\left(1 + \frac{\mu}{2(2-\mu)}t\right)^{-1}$  in series form for  $|t| < \frac{2}{\mu}|2-\mu|$ , and using Formulas (5) and (6), we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} \mathfrak{V}_{n+1}(x; \lambda; \mu) \frac{t^n}{n!} &= x \sum_{n=0}^{\infty} \mathfrak{V}_n(x; \lambda; \mu) \frac{t^n}{n!} + \frac{1}{(2-\mu)} \left( \sum_{n=0}^{\infty} \frac{\mu^n}{2^n(\mu-2)^n} t^n \right) \\ &\times \left[ \frac{\mu}{2} \sum_{n=0}^{\infty} \mathfrak{V}_n(x; \lambda; \mu) \frac{t^n}{n!} - \lambda \left( \sum_{n=0}^{\infty} \mathfrak{V}_n(\lambda; \mu) \frac{t^n}{n!} \right) \left( \sum_{n=0}^{\infty} \mathfrak{V}_n(x+1; \lambda; \mu) \frac{t^n}{n!} \right) \right]. \end{aligned} \quad (15)$$

Then taking into account the series product in (15) and equating the coefficients of  $t$ , we get Identity (14).

## 7 Identities inspired via umbral calculus

Umbral calculus approach is a useful tool to get and guess arithmetic and combinatorial identities, see for instance Gessel [7] on some applications of the classical umbral calculus, Di Crescenzo et al. [5] on umbral calculus. See also classical references as those of Roman and Rota [11].

Let  $\mathbf{B}^n(\lambda; \mu)$  be the umbra defined by  $\mathbf{B}^n(\lambda; \mu) := \mathfrak{V}_n(\lambda; \mu)$  and  $(\mathfrak{V}_n(x; \lambda; \mu))_{n \geq 0}$  defined by

$$\sum_{n \geq 0} \mathfrak{V}_n(x; \lambda; \mu) \frac{t^n}{n!} = F(t) e^{xt} = \exp((\mathbf{B}(\lambda; \mu) + x)t),$$

where  $F(t) := \sum_{n \geq 0} \mathfrak{V}_n(\lambda; \mu) \frac{t^n}{n!} = \exp(\mathbf{B}(\lambda; \mu)t)$ . So,  $\mathfrak{V}_n(x; \lambda; \mu)$  admits the umbral representation

$$\mathfrak{V}_n(x; \lambda; \mu) = (\mathbf{B}(\lambda; \mu) + x)^n.$$

**Theorem 9** *Let  $n$  be nonnegative integer. Then*

$$\mathfrak{V}_n(x+1; \lambda; \mu) = \sum_{k=0}^n \binom{n}{k} \mathfrak{V}_k(x; \lambda; \mu).$$

*Proof* By the umbral representation  $\mathfrak{V}_n(x; \lambda; \mu) = (\mathbf{B}(\lambda; \mu) + x)^n$ , we have

$$\mathfrak{V}_n(x+1; \lambda; \mu) = (\mathbf{B}(\lambda; \mu) + (x+1))^n = \sum_{k=0}^n \binom{n}{k} (\mathbf{B}(\lambda; \mu) + x)^k = \sum_{k=0}^n \binom{n}{k} \mathfrak{V}_k(x; \lambda; \mu).$$

**Theorem 10** *Let  $n, m$  be nonnegative integers. Then*

$$\sum_{k=0}^n \binom{n}{k} y^{n-k} \mathfrak{V}_{m+k}(x; \lambda; \mu) = \sum_{k=0}^m \binom{m}{k} (-y)^{m-k} \mathfrak{V}_{n+k}(x+y; \lambda; \mu).$$

*Proof* By the umbral representation  $\mathfrak{V}_n(x; \lambda; \mu) = (\mathbf{B}(\lambda; \mu) + x)^n$ , on the one hand, we have

$$\begin{aligned} (\mathbf{B}(\lambda; \mu) + (x+y))^n (\mathbf{B}(\lambda; \mu) + x)^m &= (\mathbf{B}(\lambda; \mu) + (x+y))^n (\mathbf{B}(\lambda; \mu) + (x+y) - y)^m \\ &= \sum_{k=0}^m \binom{m}{k} (-y)^{m-k} (\mathbf{B}(\lambda; \mu) + (x+y))^{n+k} \\ &= \sum_{k=0}^m \binom{m}{k} (-y)^{m-k} \mathfrak{V}_{n+k}(x+y; \lambda; \mu), \end{aligned}$$

and on the other hand, we have

$$\begin{aligned} (\mathbf{B}(\lambda; \mu) + (x+y))^n (\mathbf{B}(\lambda; \mu) + x)^m &= \sum_{k=0}^n \binom{n}{k} y^{n-k} (\mathbf{B}(\lambda; \mu) + x)^{m+k} \\ &= \sum_{k=0}^n \binom{n}{k} y^{n-k} \mathfrak{V}_{m+k}(x; \lambda; \mu). \end{aligned}$$

Hence, the two expressions give the desired identity.

---

## References

1. APOSTOL, T. M., ET AL. On the lerch zeta function. *Pacific Journal of Mathematics* 1, 2 (1951), 161–167.
2. BELBACHIR, H., AND HADJ-BRAHIM, S. Some explicit formulas for euler-genocchi polynomials. *Integers* 19 (2019), A28.
3. BELBACHIR, H., HADJ-BRAHIM, S., AND RACHIDI, M. On another approach for a family of appell polynomials. *Filomat* 32, 12 (2018), 4155–4164.
4. BERNOULLI, J. *Ars conjectandi*. Impensis Thurnisiorum, fratrum, 1713.
5. DI CRESCENZO, A., AND ROTA, G.-C. Sul calcolo umbrale. *Ricerche di Matematica* 43 (1994), 129–162.
6. EULER, L. Methodus generalis summandi progressionis. *Commentarii academiae scientiarum Petropolitanae* (1738), 68–97.
7. GESSEL, I. M. Applications of the classical umbral calculus. *Algebra Universalis* 49, 4 (2003), 397–434.
8. KARGIN, L., AND KURT, V. On the generalization of the euler polynomials with the real parameters. *Applied Mathematics and Computation* 218, 3 (2011), 856–859.
9. LUO, Q.-M., AND SRIVASTAVA, H. Some generalizations of the apostol–bernoulli and apostol–euler polynomials. *Journal of Mathematical Analysis and Applications* 308, 1 (2005), 290–302.
10. RAABE, J. L. Zurückführung einiger summen und bestimmten integrale auf die jacob-bernoullische function. *Journal für die reine und angewandte Mathematik* 1851, 42 (1851), 348–367.
11. ROMAN, S. M., AND ROTA, G.-C. The umbral calculus. *Advances in Mathematics* 27, 2 (1978), 95–188.
12. SRIVASTAVA, H. M. Some generalizations and basic (or  $q$ -) extensions of the bernoulli, euler and genocchi polynomials. *Appl. Math. Inform. Sci* 5, 3 (2011), 390–444.