

**INFINITE-DIMENSIONAL THURSTON THEORY
AND TRANSCENDENTAL DYNAMICS I:
INFINITE-LEGGED SPIDERS**

KONSTANTIN BOGDANOV

ABSTRACT. We develop techniques that lay out a basis for generalizations of the famous Thurston’s Topological Characterization of Rational Functions for an *infinite* set of marked points and branched coverings of infinite degree. Analogously to the classical theorem we consider the Thurston’s σ -map acting on a Teichmüller space which is this time infinite-dimensional — and this leads to a completely different theory comparing to the classical setting.

We demonstrate our techniques by giving an alternative proof of the result by Markus Förster about the classification of exponential functions with the escaping singular value. This is the first out of four articles which are prepared in order to publish results of the author’s PhD thesis.

CONTENTS

1.	Introduction	1
2.	Prerequisites	6
3.	Setup of Thurston iteration	10
4.	Strict contraction of σ	13
5.	Id-type maps	16
6.	Spiders	18
7.	Invariant compact subset	27
8.	Proof of Classification Theorem	31
9.	Acknowledgements	32
	References	32

1. INTRODUCTION

Motivation from transcendental dynamics. A part of complex dynamics that has been studied intensively over the last decades is the dynamics of transcendental entire functions. Good introductory references for the subject are [EL, S].

As in polynomial dynamics, for a transcendental entire function f an important and natural concept is its *escaping set* $I(f) = \{z \in \mathbb{C} : f^n(z) \rightarrow \infty \text{ as } n \rightarrow \infty\}$. It is proved in [SZ] for the exponential family $\{e^z + \kappa : \kappa \in \mathbb{C}\}$ and in [RRRS] for more general families of functions (of bounded type and finite order) that the escaping set of every function in the family is organized in form of *dynamic rays* [DK].

Roughly speaking, this means that every point $z \in I(f)$ can be joined to ∞ by a unique simple arc contained in $I(f)$ (or is mapped to such arc after finitely many iterations), so that the iterates f^n restricted to the arc are injective, continuous at ∞ (here we agree for the sake of convenience that $f(\infty) = \infty$), and converge to ∞ uniformly. About the escaping points that belong to a dynamic ray we say

that they *escape on rays*. This is the general mode of escape for many important families of transcendental entire functions.

In many cases the points escaping on rays can be described by their potential (or “speed of escape”) and external address (or “combinatorics of escape”, i.e. the sequence of dynamic rays containing the escaping orbit). This is analogous to the Böttcher coordinates for polynomials where the points in the complement of the filled Julia set are encoded by their potential and external angle.

One of the most fruitful directions in the study of holomorphic dynamical systems is the investigation of parameter spaces. The most famous example is the study of the Mandelbrot set which is the set of parameters c in the complex plane for which the critical value c of the polynomial $z^2 + c$ does not escape. The simplest analogue in the transcendental world is the space of complex parameters κ , each associated to the function $e^z + \kappa$. When we consider parameters κ for which the singular value κ of $e^z + \kappa$ escapes, we, roughly speaking, investigate a part of the “complement of the Mandelbrot set of the exponential family”. The results in this direction are contained in [FRS, FS, F]

Now, we can ask a question. (\star) *For a transcendental entire function of finite type (i.e. with finitely many singular values), is there in its parameter space an entire function such that its singular values escape on rays and have initially prescribed potentials and external addresses?*

In this article we set up a flexible machinery for proving this kind of results for general families of entire functions and apply it to reprove the well-known classification result for the exponential family [F].

A main tool is what one might call infinite-dimensional Thurston theory in dynamics. It is a generalization of the famous *Thurston’s Topological Characterization of Rational Functions* [DH]. The subject is interesting not only as a tool but also in its own right and requires an independent exposition.

Infinite-dimensional Thurston theory. One of the questions of big interest in dynamics is the interplay between topology and geometry. Or saying differently, when there exists a geometrical object corresponding in a certain sense to the underlying topological structure? And if it exists, is it unique?

A major result in this direction is proven by William Thurston.

Theorem (Topological Characterization of Rational Functions [DH, H2]). *A post-critically finite branched covering $f : S^2 \rightarrow S^2$ with hyperbolic orbifold is realized via a rational map if and only if there is no Thurston obstruction.*

This result gives a complete answer to the question whether there exists a post-critically finite rational map with a prescribed combinatorics of the covering.

So it is natural to hope that if we prove an analogous “characterization of entire functions” for “topological entire functions” representing some properly chosen combinatorial model, then the corresponding entire function will have properties as required in question (\star). It turns out that this can be done (at least for certain explicit families of functions), but the proof, though keeping some similarities such as iterations on a Teichmüller space, goes in quite a different direction than the proof of the classical theorem. The key differences between our and the classical case are that we have to consider iterations on an *infinite-dimensional* Teichmüller space (since the set of marked points is infinite), and that our topological entire function is a branched covering of *infinite* degree. This drastically changes the approach already on the level of the setup. So far the case of infinite set of marked points for different type of functions was explored in [Br, CT, F]. A generalization

of the Characterization Theorem for post-singularly finite topological exponential functions is provided in [HSS].

Before discussing details we present the “grand scheme” of how one can construct entire function demanded in question (\star). This “grand scheme” is partially inspired by the ideas in [F].

- (1) Construct a (non-holomorphic) “model map” for which the singular values escape on rays with desired “speed of escape” and “combinatorics”. It will define (analogously to the classical case) the Thurston’s σ -map acting on the Teichmüller space of the complement of the singular orbits of the “model map”.
- (2) Construct a compact subspace in the Teichmüller space that is invariant under σ .
- (3) Prove that σ is strictly contracting in the Teichmüller metric on the constructed subspace.
- (4) Using the argument of the Banach Fixed Point Theorem prove that σ has the unique fixed point on the constructed subspace, and this point corresponds to the entire function with the desired conditions on its singular orbits.

This scheme was successfully employed to prove classification results for compositions of polynomials with the exponential [B], and we believe that it works in a much greater generality (after adjustments of its internal ingredients). In this article we develop the techniques of how to treat every item (1)-(4) and use the scheme to answer the question (\star) in the “baby case” of the exponential family. Among all, the item (2) is the “weightiest” one, and we elaborate it in the two largest Sections 6 and 7.

Setup of Thurston iteration. This is what happens in the item (1) of the “grand scheme” and Section 3.

In the parameter space that we consider we pick a function f_0 that has escaping points realizing the desired “speeds of escape” and “combinatorics” (from question (\star)). Based on f_0 we construct (by a postcomposition with a “capture”) a special quasiregular function f for which its singular orbits coincide with the corresponding escaping orbits of f_0 , and which are equal to f_0 outside of some compact set. Moreover, its singular values escape on rays (defined analogously as for entire maps). This function f plays the role of the topological branched covering in the classical theorem.

Based on f we can construct the σ -map acting on the Teichmüller space (Definition 2.21) of $\mathbb{C} \setminus P_f$ where P_f is the post-singular set of f (i.e. the set of marked points). The map $\sigma : [\varphi] \rightarrow [\tilde{\varphi}]$ is defined in the usual way via Thurston diagram (Section 3).

$$\begin{array}{ccc} \mathbb{C}, P_f & \xrightarrow{\tilde{\varphi}} & \mathbb{C}, \tilde{\varphi}(P_f) \\ \downarrow f & & \downarrow g = \varphi \circ f \circ \tilde{\varphi}^{-1} \\ \mathbb{C}, P_f & \xrightarrow{\varphi} & \mathbb{C}, \varphi(P_f) \end{array}$$

An attempt to define the Thurston’s σ -map as in the classical theorem for a *topological* (rather than quasiregular) exponential function would necessarily fail: the σ -map still can be defined on the level of isotopy classes of homeomorphisms, but since the post-singular set is infinite, the isotopy class of $\tilde{\varphi} \circ \varphi^{-1}$ relative $\varphi(P_f)$ might not necessarily contain a quasiconformal map (see Subsection 3.3 for

more details). Yet in this case σ can be interpreted as a map between different Teichmüller spaces, but we do not touch such constructions in this article.

In the article we set up the iteration for the case of exponential family, but it can be easily upgraded to *any* family of transcendental entire functions.

Id-type maps and spiders. The goal of item (2) in the “grand scheme” is to construct a compact subset of \mathcal{T}_f which is invariant under σ .

The space \mathcal{T}_f with its isotopies relative to infinitely many points in P_f is quite complicated, so one might want to take a look at simpler subspaces. These considerations lead us to the notion of id-type maps (Definition 5.2). These are quasiconformal maps that tend to identity as their argument tends to ∞ on the post-singular dynamic rays of f . An important property of these maps is that the images of post-singular dynamic rays of f preserve their asymptotics near ∞ , and this property is invariant under σ (under proper normalization). So id-type maps form a σ -invariant subspace of \mathcal{T}_f that respects the “combinatorics” of f .

To every id-type map one can associate an *infinite-legged spider* and run the so-called spider-algorithm [HS, F]. The spider which is really the driving concept of the proofs, is a convenient presentation of a point in the Teichmüller space that provides a more tame description of its homotopy information (every point in the Teichmüller space is described by the positions of images of P_f , which is moduli space information, and homotopy information). One can introduce certain equivalence relation on the set of spiders, called projective equivalence (Definition 6.26), for which the following theorem holds (more precise version is Theorem 6.28).

Theorem. *Projective equivalence of spiders is Teichmüller equivalence, i.e. two spiders are projectively equivalent if and only if the associated id-type maps represent the same point in the Teichmüller space.*

This way we obtain a description of quite complicated isotopy types of homeomorphisms relative countably many marked points via a sequence of homotopy types of simple arcs, each relative just finitely many marked points — this is a serious simplification. Another merit of the construction is that these homotopy types behave well under Thurston iteration (Theorem 6.23).

We demonstrate how the construction works for the exponential family, but it is generalizable for functions admitting escape on rays.

Invariant compact subset. Such subset is described as a list of conditions that stay invariant under σ . We give a very rough list at the moment, for more details we refer directly to Theorem 7.1.

The compact invariant subset is the set of points represented by quasiconformal maps φ such that :

- (1) φ is of id-type;
- (2) for the marked points P_f there exists an integer $N > 0$ not depending on φ such that the images under φ of the first N points of P_f are inside of the disk $\mathbb{D}_\rho(0)$ with $\rho > 0$ not depending on φ , and images of the other points of P_f are outside of $\mathbb{D}_\rho(0)$;
- (3) for very point $z \in P_f$ outside of $\mathbb{D}_\rho(0)$, $\varphi(z)$ is contained in a small disk U_z around z so that the disks U_z are mutually disjoint;
- (4) inside of $\mathbb{D}_\rho(0)$ the distances between marked points are bounded from below;
- (5) the isotopy type φ relative points of P_f outside of $\mathbb{D}_\rho(0)$ is “almost” the isotopy type of identity;

- (6) the isotopy type φ relative points of P_f inside of $\mathbb{D}_\rho(0)$ is not “too complicated”, i.e. there are quantitative bounds on how many times the marked points “twist” around each other.

Conditions (2),(3) separate the complex plane into two subsets: $\mathbb{D}_\rho(0)$ where we have not so much control on the behavior of φ but have finitely many post-singular points, and the complement of $\mathbb{D}_\rho(0)$ where the homotopy information is trivial but we have infinitely many marked points.

Conditions (2)-(4) describe the position in the moduli space, while (5)-(6) encode homotopy information of the point in the Teichmüller space.

It is natural to expect that these conditions define a compact subset, invariance is less obvious though.

We provide the construction for the exponential family. Generalizations for more than one singular value require additional conditions on the orbits.

Strict contraction of the σ -map. Item (3) of the “grand scheme” is where the infinite-dimensionality of the Teichmüller space shows up.

The statement that σ is strictly contracting in the classical theorem by Thurston is deduced from the well-known theorem of Teichmüller claiming that the quasi-conformal representative minimizing the Teichmüller distance on a surface of finite analytic type is associated to a unique normalized integrable quadratic differential. We consider a surface of infinite analytic type, and in our case this statement is false.

We are interested in finding some other way to prove strict contraction of σ on the Teichmüller space. However, this seems to be not an easy task. Fortunately, we can find a way around. If we consider *asymptotically conformal* points in the Teichmüller space, then from the Strebel’s Frame Mapping Theorem follows the existence of a unique associated normalized integrable quadratic differential, and we can prove strict contraction of σ on an asymptotically conformal subspace.

In particular, the following theorem holds (more precise version is Theorem 4.4).

Theorem. *σ is invariant and strictly contracting on the subset of asymptotically conformal points of the Teichmüller space.*

In the article we prove the theorem for compositions of polynomials with the exponential. For more general families exactly this method does not work, but it might be possible to deduce strict contraction directly from the construction of the invariant compact subset.

Classification Theorem. As was mentioned earlier, the existence of a fixed point in the invariant compact subset follows by a simple argument of Banach Fixed Point Theorem (the subset is asymptotically conformal). We also show that the entire map corresponding to the constructed fixed point has singular values that escape on rays with the prescribed “speed of escape” and “combinatorics”.

In particular, we reprove the well-known classification of exponential functions with singular values escaping on rays.

Theorem 1.1 (Classification Theorem for $e^z + \kappa$. [F]). *Let \underline{s} be an exponentially bounded external addresses that is not (pre-)periodic, and $t > t_{\underline{s}}$ be a real number. Then in the family $e^z + \kappa$ there exists the unique entire function $e^z + \kappa_{\underline{s},t}$ such that its singular value $\kappa_{\underline{s},t}$ escapes on rays, has potential t and external address \underline{s} .*

Conversely, every function in the family $e^z + \kappa$ such that its singular value escapes on non (pre-)periodic rays is one of these.

For the definitions of an exponentially bounded external address and $t_{\underline{s}}$ we refer the reader to Subsection 2.1.

2. PREREQUISITES

2.1. Escaping dynamics in the exponential family. The simplest among the entire functions of finite type are functions having exactly one singular value. Every such function is equal to $A_1 \circ \exp \circ A_2$ for some affine maps $A_1, A_2 : \mathbb{C} \rightarrow \mathbb{C}$ and is conformally conjugate to a function of the form $e^z + \kappa$. Hence, to study dynamical properties of entire functions with one singular value, it is enough to consider the exponential family $e^z + \kappa$.

For the sake of brevity we adopt the following notation.

Definition 2.1 (\mathcal{N}). *Denote by \mathcal{N} the family of transcendental entire functions of the form $e^z + \kappa$ for $\kappa \in \mathbb{C}$.*

In this section we assemble some preliminary results and definitions about the structure of the escaping set of functions in \mathcal{N} . We follow the exposition in [SZ] though use a slightly different notation that is more suitable for us.

It is proven in [SZ] for the exponential family, and in [RRRS] for much more general families, that the escaping set of such functions is organized in form of *dynamic rays*.

Definition 2.2 (Ray tails). *Let f be a transcendental entire function. A ray tail of f is a continuous curve $\gamma : [0, \infty) \rightarrow I(f)$ such that for every $n \geq 0$ the restriction $f^n|_\gamma$ is injective with $\lim_{t \rightarrow \infty} f^n(\gamma(t)) = \infty$, and furthermore $f^n(\gamma(t)) \rightarrow \infty$ uniformly in t as $n \rightarrow \infty$.*

Definition 2.3 (Dynamic rays, escape on rays, endpoints). *A dynamic ray of a transcendental entire function f is a maximal injective curve $\gamma : (0, \infty) \rightarrow I(f)$ such that $\gamma|_{[t, \infty)}$ is a ray tail for every $t > 0$.*

If a point $z \in I(f)$ belongs to a dynamic ray, we say that the point escapes on rays (because in this case every iterate of the point belongs to a dynamic ray).

If there exists a limit $z = \lim_{t \rightarrow 0} \gamma(t)$, then we say that z is an endpoint of the dynamic ray γ .

From now we assume that $f \in \mathcal{N}$. Let $r \in \mathbb{R}$ be such that the right half-plane $\mathbb{H}_r = \{z \in \mathbb{C} : \operatorname{Re} z > r\}$ does not contain the singular value of f . Then the preimage of \mathbb{H}_r under f has countably many path-connected components, called *tracts*, having ∞ as a boundary point, so that for every tract T the restriction $f|_T$ is a conformal isomorphism. We denote by T_n the tract contained in the infinite strip between the straight lines $\operatorname{Im} z = 2\pi i(n - 1/2)$ and $\operatorname{Im} z = 2\pi i(n + 1/2)$. This way we enumerate the tracts by integer numbers. Clearly, this numeration does not depend on a particular choice of r .

Let $\gamma : [0, \infty) \rightarrow I(f)$ be a ray tail of f . Then for every $n \geq 0$ there is $t_n \geq 0$ and $s_n \in \mathbb{Z}$ such that $f^n \circ \gamma|_{[t_n, \infty)}$ is contained in the tract T_{s_n} . Moreover, all t_n except finitely many are equal to 0.

Definition 2.4 (External address). *Let z be a point escaping on rays. We say that z has external address $\underline{s} = (s_0 s_1 s_2 \dots)$ where $s_n \in \mathbb{Z}$, if each $f^n(z)$ belongs to a ray tail contained in T_{s_n} near ∞ .*

In this case we also say that the dynamic ray (or ray tail) containing z has external address \underline{s} .

It is clear that the external address does not depend on a particular choice of \mathbb{H}_r in the definition of tracts.

On the set of external addresses we can consider the usual shift-operator $\sigma : (s_0 s_1 s_2 \dots) \mapsto (s_1 s_2 s_3 \dots)$.

Now we define the function which allows to characterize the “speed of escape” of points escaping on rays.

Definition 2.5 ($F(t)$). Denote by $F : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ the function

$$F(t) := e^t - 1.$$

What one should know about F is that its iterates grow very fast, as shown in the next self-evident lemma.

Lemma 2.6 (Super-exponential growth of F -iterates). For every $t > 0$ we have $F^n(t)/2^n \rightarrow \infty$ as $n \rightarrow \infty$.

Next definition relates the last two notions.

Definition 2.7 (Exponentially bounded external address, $t_{\underline{s}}$). We say that the sequence $\underline{s} = (s_0 s_1 s_2 \dots)$ is exponentially bounded if there exists $t > 0$ such that $s_n/F^n(t) \rightarrow 0$ as $n \rightarrow \infty$. The infimum of such t we denote by $t_{\underline{s}}$.

Next statement claims that no other type of external addresses can appear for points escaping on rays.

Lemma 2.8 (Only exponentially bounded \underline{s} [SZ]). If $z \in I(f)$ escapes on rays and has external address $\underline{s} = (s_0 s_1 s_2 \dots)$, then \underline{s} is exponentially bounded.

The key statement about the escaping set of functions in \mathcal{N} is the following theorem.

Theorem 2.9 (Escape on rays and asymptotic formula [SZ]). Let $f \in \mathcal{N}$. Then for every exponentially bounded external address there exists the unique dynamic ray realizing it.

Further, if $\mathcal{R}_{\underline{s}}$ is the dynamic ray having exponentially bounded external address $\underline{s} = (s_0 s_1 s_2 \dots)$, and no strict forward iterate of $\mathcal{R}_{\underline{s}}$ contains the singular value of f , then $\mathcal{R}_{\underline{s}}$ can be parametrized by $t \in (t_{\underline{s}}, \infty)$ so that

$$\mathcal{R}_{\underline{s}}(t) = t + 2\pi i s_0 + O(e^{-t}), \quad (2.1)$$

and

$$f^n \circ \mathcal{R}_{\underline{s}} = \mathcal{R}_{\sigma^n \underline{s}} \circ F.$$

Asymptotic bounds $O(\cdot)$ for $\mathcal{R}_{\sigma^n \underline{s}}(t)$ are uniform in n on every ray tail contained in $\mathcal{R}_{\underline{s}}$.

Every escaping point is mapped after finitely many iterations either on a dynamic ray or to an endpoint of a ray. If the singular value of f does not escape, then all dynamic rays are parametrized as above, and every escaping point escapes either on rays or as endpoints of rays.

The theorem above allows us to define the notion of potential.

Definition 2.10 (Potential). Let $f \in \mathcal{N}$ and assume that z escapes on rays with escaping address \underline{s} . We say that t is the potential of z if $|f^n(z) - F^n(t) - 2\pi i s_n| \rightarrow 0$ as $n \rightarrow \infty$.

From Theorem 2.1 follows that every point escaping on rays (without singular values on forward iterates) has a well-defined potential, and different points on the same ray have different potentials.

2.2. Quasiconformal maps. Standard references are [A, LV]. We start with a few preliminary definitions.

Definition 2.11 (Quadrilateral). *A quadrilateral $Q(z_1, z_2, z_3, z_4)$ is a Jordan domain Q together with a sequence z_1, z_2, z_3, z_4 of boundary points called vertices of the quadrilateral. Order of vertices agrees with the positive orientation with respect to Q . Arcs z_1z_2 and z_3z_4 are called a -sides, arcs z_2z_3 and z_4z_1 are called b -sides.*

Every quadrilateral Q is conformally equivalent to the unique canonical rectangle with the length of b -sides equal to 1. For a quadrilateral Q , the length of the a -sides of the canonical rectangle is called a (conformal) *modulus* of Q and is denoted by $\text{mod } Q$.

Definition 2.12 (Maximal dilatation). *Let U be a plane domain and ψ be an orientation-preserving homeomorphism of U . The maximal dilatation of ψ is called the number*

$$K(\psi) = \sup_{Q \subset U} \frac{\text{mod } \psi(Q)}{\text{mod } Q},$$

where the supremum is taken over all quadrilaterals Q contained in G together with its boundary.

Now we can define quasiconformal maps.

Definition 2.13 (Quasiconformal map). *An orientation-preserving homeomorphism ψ of a plane domain U is called quasiconformal if its maximal dilatation $K(\psi)$ is finite. If $K(\psi) \leq K < \infty$, then ψ is called K -quasiconformal.*

It is easy to show that the inverse of a K -quasiconformal mapping is K -quasiconformal, and the composition of a K_1 -quasiconformal and K_2 -quasiconformal mapping is K_1K_2 -quasiconformal.

We also provide the analytic definition of quasiconformal maps. It is equivalent to the previous one.

Definition 2.14 (Quasiconformal map). *A homeomorphism ψ of a plane domain U is quasiconformal if there exists $k < 1$ such that*

- (1) ψ has locally integrable, distributional derivatives ψ_z and $\psi_{\bar{z}}$ on U , and
- (2) $|\psi_{\bar{z}}| \leq k|\psi_z|$ almost everywhere.

Such ψ is called K -quasiconformal, where $K = \frac{1+k}{1-k}$.

Every quasiconformal map is determined by its Beltrami coefficient.

Definition 2.15 (Beltrami coefficient). *The function $\mu_\psi(z) = \psi_{\bar{z}}(z)/\psi_z(z)$ (defined a.e. on U) is called the Beltrami coefficient of ψ .*

Providing the Beltrami coefficient is almost the same as giving a quasiconformal map. Suppose $\mu(z)$ is a measurable complex-valued function defined on a domain $U \subset \hat{\mathbb{C}}$ for which $\|\mu\|_{L^\infty} = k < 1$. We can ask whether it is possible to find a quasiconformal map ψ satisfying the Beltrami equation

$$\psi_{\bar{z}}(z) = \mu(z)\psi_z(z) \tag{2.2}$$

where the partial derivatives $\psi_z(z)$ and $\psi_{\bar{z}}(z)$ are defined in the sense of distributions and are locally integrable.

The answer for $U = \hat{\mathbb{C}}$ is contained in Measurable Riemann Mapping Theorem.

Theorem 2.16 (Measurable Riemann Mapping Theorem [GL]). *The Beltrami equation 2.2 gives a one-to-one correspondence between the set of quasiconformal homeomorphisms of $\hat{\mathbb{C}}$ that fix the points $0, 1$ and ∞ and the set of measurable complex-valued functions μ on $\hat{\mathbb{C}}$ for which $\|\mu\|_{L^\infty} < 1$. Furthermore, the normalized solution ψ^μ depends holomorphically on μ .*

2.3. Beurling-Ahlfors extension. Next, we formulate two theorems that help us to reconstruct a quasiconformal map on a half-plane from its boundary values given by a *quasisymmetric function*.

Definition 2.17 (Quasisymmetric function). *Let ξ be a continuous strictly increasing self-homeomorphism of the real line. We say that ξ is ρ -quasisymmetric if there exists a positive constant ρ such that for every $t > 0$ and every $x \in \mathbb{R}$ we have*

$$\frac{1}{\rho} \leq \frac{\xi(x+t) - \xi(x)}{\xi(x) - \xi(x-t)} \leq \rho.$$

The first theorem basically says that the restriction of the quasiconformal self-homeomorphism of the upper half plane is quasisymmetric.

Theorem 2.18 (Boundary values of homeomorphisms [LV, Theorem 6.2]). *Let φ be a self-homeomorphism of the closure of the upper half-plane fixing ∞ such that for every quadrilateral Q with boundary on the real line holds*

$$\text{mod } \varphi(Q) \leq K \text{ mod } Q$$

for some $K \geq 1$.

Then for every $t > 0$ and every $x \in \mathbb{R}$ holds

$$\frac{1}{\lambda(K)} \leq \frac{\varphi(x+t) - \varphi(x)}{\varphi(x) - \varphi(x-t)} \leq \lambda(K),$$

where $\lambda : [0, \infty] \rightarrow \mathbb{R}$ is a continuous function not depending on φ so that $\lambda(1) = 1$.

Remark 2.19. *Strictly speaking, the authors in [LV] prove the theorem for K -quasiconformal maps of the upper half plane but exactly the same proof works in the setting as above.*

The second theorem says that quasisymmetric maps can be promoted to quasiconformal maps with a sharp estimate on the maximal dilatation.

Theorem 2.20 (Beurling-Ahlfors extension [BA]). *Let $\xi : \mathbb{R} \rightarrow \mathbb{R}$ be a function that is ρ -quasisymmetric. Then it extends to a K -quasiconformal self-homeomorphism of the upper half-plane with $K \leq \rho^2$.*

2.4. Teichmüller theory and quadratic differentials. Good references are [GL, H1]. To avoid unnecessary technicality we give *ad hoc* definitions that work specifically in our context.

Definition 2.21 (Teichmüller space of $\mathbb{C} \setminus V$). *Let V be a discrete subset of \mathbb{C} without accumulation points in \mathbb{C} . The Teichmüller space of the Riemann surface $\mathbb{C} \setminus V$ is the set of quasiconformal homeomorphisms of $\mathbb{C} \setminus V$ modulo post-composition with an affine map and isotopy relative V .*

The points in the Teichmüller space of $\mathbb{C} \setminus V$ are equivalence classes $[\varphi]$ of quasiconformal homeomorphisms φ of $\mathbb{C} \setminus V$.

Remark 2.22. *A more standard definition of the Teichmüller space on a Riemann surface involves isotopy relative the ideal boundary rather than the topological boundary. For planar domains the two definitions are equivalent [GL].*

Every Teichmüller space can be equipped with the special metric.

Definition 2.23 (Teichmüller distance). *Let $[\varphi_0], [\varphi_1]$ be two points in the Teichmüller space of $\mathbb{C} \setminus V$. The Teichmüller distance $d_T([\varphi_0], [\varphi_1])$ is defined as*

$$\inf_{\psi \in [\varphi_1 \circ (\varphi_0)^{-1}]} \log K(\psi),$$

that is, a lower bound of the logarithm of maximal dilatations of quasiconformal maps that belong to $[\varphi_1 \circ (\varphi_0)^{-1}]$.

The Teichmüller space of $\mathbb{C} \setminus V$ equipped with the Teichmüller distance is complete metric space.

We also need the notion of holomorphic quadratic differential.

Definition 2.24 (Holomorphic quadratic differential). *A holomorphic quadratic differential on $\mathbb{C} \setminus V$ is a meromorphic function on $\mathbb{C} \setminus V$.*

We say that a holomorphic quadratic differential q is *integrable* if $q \in L^1(\mathbb{C})$ (or equivalently $q \in L^1(\mathbb{C} \setminus V)$). An integrable quadratic differential has either a simple pole, or a removable singularity at punctures (i.e. at points of V). Further, it has a pole of order 4 at infinity and at least two finite simple poles.

The next theorem is an important result of Teichmüller theory.

Theorem 2.25 (Teichmüller Uniqueness Theorem [GL]). *Suppose φ_0 is a quasiconformal map of $\mathbb{C} \setminus V$ having Beltrami coefficient $k_0 \frac{|q_0|}{q_0}$ with an integrable quadratic differential q_0 and $0 \leq k_0 < 1$. Let φ be any other quasiconformal map of $\mathbb{C} \setminus V$ such that $\varphi^{-1} \circ \varphi_0 \in [\text{id}]$ in the Teichmüller space of $\mathbb{C} \setminus \varphi_0(V)$, and let μ_φ be the Beltrami coefficient of φ . Then either there exists a set of positive measure in $\mathbb{C} \setminus V$ on which $|\mu_\varphi(z)| > k_0$, or $\mu_\varphi(z) = k_0 \frac{|q_0|}{q_0}$ almost everywhere.*

In this article we need a special *asymptotically conformal* subset of the Teichmüller space.

Definition 2.26 (Asymptotically conformal points [GL]). *A point $[\varphi]$ in the Teichmüller space of $\mathbb{C} \setminus V$ is called asymptotically conformal if for every $\epsilon > 0$ there is a compact set $C \subset \mathbb{C} \setminus V$ and a representative $\psi \in [\varphi]$ such that $|\mu_\psi| < \epsilon$ a.e. on $(\mathbb{C} \setminus V) \setminus C$.*

Every asymptotically conformal point evidently contains a *frame mapping*.

Definition 2.27 (Frame mapping [GL]). *Let φ_0 an extremal representative of $[\varphi_0]$ (i.e. minimizing the Teichmüller distance in $[\varphi_0]$) in the Teichmüller space of $\mathbb{C} \setminus V$, and K_0 be its maximal dilatation. If there exists a quasiconformal map $\varphi \in [\varphi_0]$ such that $\text{ess sup}_{(\mathbb{C} \setminus V) \setminus C} |K_\varphi| < K_0$ where C is a compact subset of $\mathbb{C} \setminus V$, we say that φ is a frame mapping of the equivalence class $[\varphi_0]$.*

Since every asymptotically conformal point contains a frame map, next theorem provides a presentation of the extremal representatives of the asymptotically conformal points.

Theorem 2.28 (The Frame Mapping Theorem [GL]). *If the point $[\varphi]$ in the Teichmüller space of $\mathbb{C} \setminus V$ has a frame mapping, then it has an extremal representative φ_0 which has Beltrami coefficient $\mu_{\varphi_0} = k_0 \frac{|q_0|}{q_0}$ where q_0 is an integrable holomorphic quadratic differential with $\|q_0\|_{L^1} = 1$ and $0 \leq k_0 < 1$. Further, if $0 < k_0 < 1$, then q_0 is uniquely determined.*

3. SETUP OF THURSTON ITERATION

3.1. Thurston iteration. In this section we are going to introduce a procedure called Thurston iteration on a Teichmüller space. It will be seen later that this procedure allows us to find entire functions with the desired combinatorics of escape of the singular values.

Let f_0 be a transcendental entire function of finite type, and $\xi : \mathbb{C} \rightarrow \mathbb{C}$ be a quasiconformal map. Define the quasiregular function $f = \xi \circ f_0$ and assume that every singular point of f (i.e. image of a singular point of f_0 under ξ) either escapes or is (pre-)periodic under iterations of f . Let P_f be the union of all singular orbits of f (including singular values). It is common to call P_f either the *post-singular* set or the set of *marked points*.

As in the proof of Thurston's Topological Characterization of Rational Functions [DH], one is interested in existence of an entire map g which is Thurston equivalent to f .

Definition 3.1 (Thurston equivalence). *We say that f is Thurston equivalent to an entire map g if there exist two homeomorphisms $\varphi, \psi : \mathbb{C} \rightarrow \mathbb{C}$ such that*

- (1) $\varphi = \psi$ on P_f ,
- (2) the following diagram commutes

$$\begin{array}{ccc} \mathbb{C}, P_f & \xrightarrow{\psi} & \mathbb{C}, \psi(P_f) \\ \downarrow f & & \downarrow g \\ \mathbb{C}, P_f & \xrightarrow{\varphi} & \mathbb{C}, \varphi(P_f) \end{array}$$

- (3) φ is isotopic to ψ relative P_f .

Note that unlike in the classical theory we consider Thurston equivalence relative the *infinite* set of marked points P_f .

Definition 3.2 (\mathcal{T}_f). *Denote by \mathcal{T}_f the Teichmüller space of $\mathbb{C} \setminus P_f$.*

Then the Thurston map

$$\sigma : \mathcal{T}_f \rightarrow \mathcal{T}_f$$

is defined as follows. Let $[\varphi]$ be a point in \mathcal{T}_f where φ is quasiconformal. Then $\varphi \circ f$ defines a conformal structure on \mathbb{C} . By the Uniformization theorem there exists a map $\tilde{\varphi}$ that is a conformal homeomorphism from \mathbb{C} with the new complex structure (given by $\varphi \circ f$) to the standard \mathbb{C} (and it will be quasiconformal with respect to the standard \mathbb{C}). Then by definition $\sigma[\varphi] := [\tilde{\varphi}]$. It is easy to check that σ is well-defined on the level of the Teichmüller space. The procedure is encoded in the following diagram (called *Thurston diagram*):

$$\begin{array}{ccc} \mathbb{C}, P_f & \xrightarrow{\tilde{\varphi}} & \mathbb{C}, \tilde{\varphi}(P_f) \\ \downarrow f = \xi \circ f_0 & & \downarrow g \\ \mathbb{C}, P_f & \xrightarrow{\varphi} & \mathbb{C}, \varphi(P_f) \end{array}$$

The map $g = \varphi \circ f \circ \tilde{\varphi}^{-1}$ corresponding to the right vertical arrow is by construction a transcendental entire function. As in the classical case, the σ -map is evidently continuous.

As in the classical paper [DH], we are looking for a fixed point of σ . It is easy to see that the following lemma holds.

Lemma 3.3 (Thurston equivalence and fixed points of σ). *If $[\varphi] = [\tilde{\varphi}]$, then the map $g = \varphi \circ f \circ \tilde{\varphi}^{-1}$ on the right hand side of the Thurston diagram is Thurston equivalent to f .*

3.2. Captured exponential function. Since our primary goal is to prove the Classification Theorem for the exponential family using Thurston iteration, we have to choose the map f properly. More precisely, let $f_0 \in \mathcal{N}$ be a function with the non-escaping (finite) singular value v . Next, let $\{a_n\}_{n=0}^{\infty}$ be an orbit of f_0 that

escapes on rays \mathcal{R}_n , and R_n be the part of the dynamic ray \mathcal{R}_n from a_n to ∞ (i.e. the ray tail with the endpoint a_n).

Now we describe a construction of the *capture*. It is a carefully chosen quasiconformal homeomorphism of \mathbb{C} that is equal to identity outside of a bounded set and mapping the singular value of f_0 to a_0 .

Fix a bounded Jordan domain $U \subset \mathbb{C} \setminus \bigcup_{n>0} R_n$ containing both v and a_0 . Define an isotopy $c_u : \mathbb{C} \rightarrow \mathbb{C}$, $u \in [0, 1]$ through quasiconformal maps, such that $c_0 = \text{id}$ on \mathbb{C} , $c_u = \text{id}$ on $\mathbb{C} \setminus U$ for $u \in [0, 1]$, and $c_1(v) = a_0$. We say that the quasiconformal map $c = c_1$ is a capture.

Remark 3.4. *Note that the choice of the capture is not unique, so we select one of them.*

Define the map $f := c \circ f_0$. It is a *quasiregular* map whose singular orbits coincide with $\{a_n\}_{n=0}^\infty$, the orbit of a_0 under f_0 (also $R_{n+1} = f(R_n)$). In this article we are mainly interested in the σ -map associated to the map f constructed this way.

3.3. Motivation of the “quasiregular setup”. We need to say a few words in order to explain why we do not follow the classical setup of Thurston iteration but use a *quasiregular* rather than *topological* branched covering f as in the Thurston’s Characterization Theorem. The main trouble is that for post-singularly infinite topological branched coverings f the isotopy classes $[\varphi]$ and $\sigma[\varphi]$ generally belong to different Teichmüller spaces (which never happens when P_f is finite). As an example one could simply consider the exponential.

Lemma 3.5 (Classical definition of σ does not work). *Let $f(z) = e^z$, and let $\{a_n\} = \{f^{\circ n}(0)\}_{n=0}^\infty$ be the orbit of its singular value 0. Then there exist orientation-preserving (not quasiconformal) homeomorphisms $\varphi, \tilde{\varphi} : \mathbb{C} \rightarrow \mathbb{C}$ such that $\varphi \circ f \circ \varphi^{-1}$ is entire transcendental function but the isotopy class $[\varphi \circ \tilde{\varphi}^{-1}]$ (relative $\tilde{\varphi}(P_f)$) does not contain any quasiconformal homeomorphisms.*

Proof. Pick an orientation-preserving homeomorphism φ such that $\varphi(\bar{z}) = \overline{\varphi}(z)$, and $\varphi(a_n) = a_{2n}$. In particular, it preserves the real line.

Let $\tilde{\varphi}$ be a homeomorphism such that $g = \varphi \circ f \circ \tilde{\varphi}^{-1}$ is entire — its existence can be proven as earlier by Uniformization theorem — and normalized so that $g(z) = e^z + \kappa$. Since $\varphi(0) = \varphi(a_0) = a_0 = 0$, we have $\kappa = 0$. Further, we may assume that $\tilde{\varphi}$ also preserves the real line. Hence $\tilde{\varphi}(a_n) = a_{2n+1}$.

Now, consider the orientation-preserving homomorphism $\varphi \circ \tilde{\varphi}^{-1}$, which preserves the real line. We want to prove that the isotopy class $[\varphi \circ \tilde{\varphi}^{-1}]$ does not contain any quasiconformal maps.

Let $\psi \in [\varphi \circ \tilde{\varphi}^{-1}]$ and consider the annuli

$$A_k := \{z \in \mathbb{C} : a_k < |z| < a_{k+2}\}.$$

Then the annulus $B_n = \psi(A_{2n+1})$ separates the points $0, a_{2n}$ from the points a_{2n+2}, ∞ , and this holds for all $n > 0$.

From the general theory of quasiconformal maps (see e.g. [LV]) we know that there is a real constant $C > 0$ so that

$$\text{mod}(B_n) \leq C \text{mod}(A_{2n}).$$

Hence

$$\frac{\text{mod}(\psi(A_{2n+1}))}{\text{mod}(A_{2n+1})} = \frac{\text{mod}(B_n)}{\text{mod}(A_{2n+1})} \leq \frac{C \text{mod}(A_{2n})}{\text{mod}(A_{2n+1})} \rightarrow 0$$

as $n \rightarrow \infty$ and ψ cannot be quasiconformal. □

The solution to this issue of “jumping” between different Teichmüller spaces is to take a quasiregular function f which appears with its own complex plane, i.e. exactly what we did in the setup of the Thurston iteration.

4. STRICT CONTRACTION OF σ

In this section we prove that the σ -map associated to the captured exponential function (defined as in Subsection 3.2) is strictly contracting in the Teichmüller metric on the subset of asymptotically conformal points in the Teichmüller space (Definition 2.26). Instead of showing it directly, we prove a row of more general lemmas which cover the case of the composition of a polynomial with the exponential: it will not require any essential additional efforts but we will be able to use this theorem in the subsequent articles which handle the infinite-dimensional Thurston theory in this case.

We work in the more general setting of Subsection 3.1 (not with the captured exponential function). More precisely, let f_0 be a transcendental entire function of finite type, and $\xi : \mathbb{C} \rightarrow \mathbb{C}$ be a quasiconformal map. Consider the quasiregular function $f = \xi \circ f_0$ and assume that every singular point of f either escapes or is (pre-)periodic. In this case we have the map $\sigma : \mathcal{T}_f \rightarrow \mathcal{T}_f$ associated to f . We want to apply it to the asymptotically conformal points in \mathcal{T}_f .

Lemma 4.1 (Strict contraction of σ). *If points $[\varphi], [\psi] \in \mathcal{T}_f$ and their σ -images $\sigma[\varphi], \sigma[\psi] \in \mathcal{T}_f$ are asymptotically conformal, and $[\varphi] \neq [\psi]$, then*

$$d_T(\sigma[\varphi], \sigma[\psi]) < d_T([\varphi], [\psi]).$$

Proof. Assume that $\sigma[\varphi] \neq \sigma[\psi]$, otherwise the statement is trivial. Consider the joint Thurston diagram for maps φ and ψ :

$$\begin{array}{ccccc} \mathbb{C}, \tilde{\psi}(P_f) & \xleftarrow{\tilde{\psi}} & \mathbb{C}, P_f & \xrightarrow{\tilde{\varphi}} & \mathbb{C}, \tilde{\varphi}(P_f) \\ \downarrow g & & \downarrow f & & \downarrow h \\ \mathbb{C}, \psi(P_f) & \xleftarrow{\psi} & \mathbb{C}, P_f & \xrightarrow{\varphi} & \mathbb{C}, \varphi(P_f) \end{array}$$

Here g and h are entire maps. Since we are only interested in the Teichmüller distance between $\sigma[\varphi]$ and $\sigma[\psi]$, we can denote $\eta := \varphi \circ \psi^{-1}$ and consider the following wrapped diagram:

$$\begin{array}{ccc} \mathbb{C}, \tilde{\psi}(P_f) & \xrightarrow{\tilde{\eta}} & \mathbb{C}, \tilde{\varphi}(P_f) \\ \downarrow g & & \downarrow h \\ \mathbb{C}, \psi(P_f) & \xrightarrow{[\eta]} & \mathbb{C}, \varphi(P_f) \end{array}$$

Since g and h are holomorphic, the L^∞ -norm of the Beltrami differential of η is preserved under the pull-back through g . It follows immediately that

$$d_T(\sigma[\varphi], \sigma[\psi]) \leq d_T([\varphi], [\psi]).$$

Next, observe that $[\eta]$ is an asymptotically conformal point (as a composition of such classes). Then due to Frame Mapping Theorem 2.28 and Teichmüller Uniqueness Theorem 2.25 there exists the unique map η_0 with the smallest maximal dilatation in $[\eta]$, and its Beltrami coefficient has form $\mu_{\eta_0} = k_0 \frac{q_0}{\bar{q}_0}$ where q_0 is an

integrable holomorphic quadratic differential on $\mathbb{C} \setminus \psi(P_f)$ such that $\|q_0\|_{L^1} = 1$. Let $\tilde{\eta}_0$ be a solution of the Beltrami equation with the Beltrami coefficient $\mu_{\eta_0 \circ g}$. Then the Beltrami coefficient of $\tilde{\eta}_0$ will have form $\mu_{\tilde{\eta}_0} = k_0 \frac{|q_0 \circ g| |g'|^2}{|q_0 \circ g| |g'|^2}$.

Since $\sigma[\eta]$ is asymptotically conformal (as a composition of asymptotically conformal points) in $\sigma[\eta]$ there is a unique extremal map $\tilde{\eta}_1$ and its Beltrami differential has the form $\mu_{\tilde{\eta}_1} = k_1 \frac{|q_1|}{q_1}$ where q_1 is an integrable quadratic differential on $\mathbb{C} \setminus \tilde{\psi}(P_f)$ such that $\|q_1\|_{L^1} = 1$, and $k_1 \leq k_0$.

If $k_1 = k_0$, then due to Teichmüller Uniqueness Theorem 2.25 we have $\mu_{\tilde{\eta}_1} = \mu_{\tilde{\eta}_0}$ and $q_0 \circ g g'^2 / q_1 \in \mathbb{R}$ in $\mathbb{C} \setminus \tilde{\psi}(P_f)$. This implies that $q_0 \circ g g'^2 / q_1 \equiv \text{const}$. But this means that q_0 cannot have more than one pole. Indeed, by Great Picard Theorem g attains every value with at most one exception infinitely many times. Since $|P_f| > 1$ (f_0 either has more than one singular value, or belongs to the exponential family and its unique singular value cannot be a fixed point), P_f contains at least one such value z . If q_0 has a pole at z , then q_1 has poles at infinitely many preimages of z under g . This is impossible since at most finitely many of them belong to P_f . So q_1 can have at most one pole in \mathbb{C} . For an integrable holomorphic quadratic differential on $\mathbb{C} \setminus \psi(P_f)$ this is possible only if $q_1 = 0$, and this contradicts to the choice of q_1 so that $\|q_1\|_{L^1} = 1$. So $k_1 < k_0$, and this means that σ is strictly contracting. \square

We want to use Lemma 4.1 to show that if f_0 is a composition of a polynomial with the exponential and ξ is asymptotically conformal as a map (i.e. its maximal dilatation tends to 0 as the argument tends to ∞), then the σ -map is strictly contracting on the subset of asymptotically conformal points in \mathcal{T}_f . In order to do so, we need to show that σ maps every asymptotically conformal point to another asymptotically conformal point. We do this in the following two lemmas.

If $\mu(z)$ is the Beltrami coefficient of a quasiconformal map φ , then we denote $K_\varphi(z) := \frac{1+\mu(z)}{1-\mu(z)}$.

Lemma 4.2 (Correction of the dilatation). *Let φ be a K -quasiconformal automorphism of \mathbb{C} such that for some $R > 0$ we have $\text{ess sup}_{\mathbb{C} \setminus \mathbb{D}_R(0)} |K_\varphi(z)| \leq 1 + \epsilon$. Then there exists $\rho \geq R$ and a quasiconformal automorphism φ' such that $\varphi \equiv \varphi'$ on $\mathbb{C} \setminus \mathbb{D}_\rho(0)$, $\text{ess sup}_{\mathbb{C}} |K_{\varphi'}(0)| \leq (1 + 2\epsilon)^3$, and φ and φ' are isotopic relative $(\mathbb{C} \setminus \mathbb{D}_\rho(0)) \cup \{0\}$.*

Proof. Choose $r > 1$ such that $\varphi(\mathbb{D}_R(0))$ is contained in a disk of diameter $\delta < 1$ with respect to the hyperbolic metric on $\varphi(\mathbb{D}_r(0))$. Let Q be a quadrilateral with its sides on the boundary of $\mathbb{D}_r(0)$, and M, M' be moduli of Q and $\varphi(Q)$, respectively. After pre- and post-composition of $\varphi|_{\mathbb{D}_r}$ with the canonical conformal maps of Q and $\varphi(Q)$ onto the rectangles $[0, M] \times [0, 1]$ and $[0, M'] \times [0, 1]$, respectively, we obtain a K -quasiconformal homeomorphism χ of these rectangles (as quadrilaterals) whose maximal dilatation does not exceed $1 + \epsilon$ outside of some set having image contained in a hyperbolic disk D of diameter $\delta < 1$ centered at $z_0 \in [0, M'] \times [0, 1]$.

Let $a := \inf_{z \in D} \text{Re } z$, $b := \sup_{z \in D} \text{Re } z$, and γ be the hyperbolic geodesic connecting the most left and the most right points of D . Consider the partition of $[0, M'] \times [0, 1]$ into three rectangles A, B, C such that

$$A = [0, a] \times [0, 1],$$

$$B = [a, b] \times [0, 1],$$

$$C = [b, M'] \times [0, 1].$$

Note that if $z \in [0, M'] \times [0, 1]$ then its distance to the boundary of Q does not exceed $M'/2$. Hence using the standard estimates for the hyperbolic metric we obtain

$$\text{mod } B = b - a = M' \int_a^b \frac{1}{2 \frac{M'}{2}} dx \leq M' \int_\gamma \lambda |dz| \leq M' \delta$$

where λ is the hyperbolic density on $[0, M'] \times [0, 1]$.

Thus due to the estimates on the maximal dilatation of χ and Grötzsch inequality for rectangles we have:

$$\text{mod } \varphi(Q) = M' = \text{mod } A + \text{mod } B + \text{mod } C \leq$$

$$(1 + \epsilon)(\text{mod } \chi^{-1}(A) + \text{mod } \chi^{-1}(B)) + M' \delta \leq$$

$$(1 + \epsilon) \text{mod } Q + M' \delta = (1 + \epsilon)M + M' \delta$$

Hence

$$\text{mod } \varphi(Q) \leq \frac{1 + \epsilon}{1 - \delta} \text{mod } Q$$

and by making r bigger (or equivalently by making δ smaller) we can choose the quasimetry constant (Definition 2.17) of $\chi|_{\partial \mathbb{D}_r(0)}$ arbitrarily close to $1 + \epsilon$. Choose it to be $1 + 2\epsilon$. Thus by Theorem 2.20 the Beurling-Ahlfors extension \mathcal{B}_φ of $\varphi|_{\partial \mathbb{D}_r(0)}$ would give us a quasiconformal homeomorphism of $\mathbb{D}_r(0)$ onto $\varphi(\mathbb{D}_r(0))$ with the maximal dilatation not exceeding $(1 + 2\epsilon)^2$. Outside of $\varphi|_{\partial \mathbb{D}_r(0)}$ define \mathcal{B}_φ to be equal to φ .

If we choose a “fat” enough annulus $\mathbb{D}_\rho(0) \setminus \overline{\mathbb{D}_r(0)}$ around $\mathbb{D}_r(0)$ for some $\rho > r$, we can construct a quasiconformal map θ' such that $\theta'|_{\mathbb{C} \setminus \varphi(\mathbb{D}_\rho(0))} \equiv \text{id}$, $\theta'(\mathcal{B}_\varphi(0)) = 0$, and its maximal dilatation on $\varphi(\mathbb{D}_\rho(0))$ is not bigger than $1 + 2\epsilon$. Denote $\varphi' := \theta' \circ \mathcal{B}_\varphi$. Then $\varphi'(0) = 0$, $\varphi'|_{\mathbb{C} \setminus \mathbb{D}_\rho} = \varphi$, its maximal dilatation does not exceed $(1 + 2\epsilon)^3$. That it is isotopic to φ relative to the $(\mathbb{C} \setminus \mathbb{D}_\rho(0)) \cup \{0\}$ follows from Alexander trick. \square

Now we are ready to show that in the case of the composition of a polynomial with the exponential the asymptotically conformal subset is invariant under σ .

Lemma 4.3 (Invariance of asymptotically conformal subset). *Let p be a non-constant polynomial, $f_0 = p \circ \exp$ and $f = \xi \circ f_0$ where $\xi : \mathbb{C} \rightarrow \mathbb{C}$ is asymptotically conformal (as a map). Then the σ -image of every asymptotically conformal point is asymptotically conformal.*

Proof. Let $[\psi] \in \mathcal{T}_f$ be asymptotically conformal. We want to prove that $\sigma[\psi] \in \mathcal{T}_f$ is also asymptotically conformal. After solving Beltrami equation for $\mu_{\psi \circ \xi \circ p}$ we obtain $[\varphi] \in \mathcal{T}_{\exp(P_f) \cup \{0\}}$ (note that $p(0) \in P_f$) such that the following expanded Thurston diagram (with g belonging to the exponential family and a polynomial q) is commutative.

$$\begin{array}{ccc}
\mathbb{C}, P_f & \xrightarrow{\tilde{\psi}} & \mathbb{C}, \tilde{\psi}(P_f) \\
\exp \downarrow & & \downarrow g \\
\mathbb{C}, \exp(P_f) \cup \{0\} & \xrightarrow{\varphi} & \mathbb{C}, \varphi(\exp(P_f) \cup \{0\}) \\
\xi \circ p \downarrow & & \downarrow q \\
\mathbb{C}, P_f & \xrightarrow{\psi} & \mathbb{C}, \psi(P_f)
\end{array}$$

Since $[\psi]$ and ξ are asymptotically conformal (the latter as a map) and the preimage of a neighborhood of ∞ under p is a neighborhood of ∞ , the point $[\varphi] \in \mathcal{T}_{\exp(P_f) \cup \{0\}}$ is also asymptotically conformal.

If $\text{ess sup}_{\mathbb{C} \setminus \mathbb{D}_R} |K_\varphi| \leq 1 + \epsilon$, then using Lemma 4.2 we can find a quasiconformal homeomorphism φ' which is isotopic to φ relative 0 and the complement of a disk $\mathbb{D}_\rho(0)$ (basically we forget finitely many marked points except 0 that are contained in $\mathbb{D}_\rho(0)$), and so that $\text{ess sup}_{\mathbb{C}} |K_{\varphi'}| \leq (1 + 2\epsilon)^3$. Consider its lift $\tilde{\varphi}'$. It is isotopic to $\tilde{\psi}$ relative $P_f \setminus X$ where X is a finite subset of P_f .

Next, we can always find a quasiconformal homeomorphism θ , equal to identity outside of some bounded set so that $\theta \circ \tilde{\varphi}'$ is isotopic to $\tilde{\psi}$ relative P_f . Further, we can also isotope $\theta \circ \tilde{\varphi}'$ in small neighborhoods of marked points in this bounded set so that the obtained map is conformal in these neighborhoods.

Since ϵ can be chosen arbitrarily small, lemma is proven. \square

After choosing $f_0 \in \mathcal{N}$ and a capture c like in Subsection 3.2, from Lemmas 4.1, 4.3 immediately follows

Theorem 4.4 (σ is strictly contracting on as. conf. subset). *If f is the captured exponential function (defined as in Subsection 3.2), then the σ -map is invariant and strictly contracting on the subset of asymptotically conformal points in \mathcal{T}_f .*

5. Id-TYPE MAPS

The purpose of this subsection is to introduce the so-called id-type (or identity-type) maps which are quasiconformal homeomorphisms of \mathbb{C} satisfying certain conditions (see Definition 5.2). They have two important properties: they are almost equal to identity near ∞ on certain dynamic rays, and the set of points $[\varphi] \in \mathcal{T}_f$ that contain a map of id-type in its equivalence class is invariant under σ . This allows to reduce Thurston iteration to the so-called Spider Algorithm [F, HS].

We work with the captured exponential function $f = c \circ f_0$ constructed in Subsection 3.2 and start with a few definitions which take central position in the theory that we develop.

Definition 5.1 (Standard spider). $S_0 = \cup_{n \geq 0} R_n$ is called the standard spider.

Definition 5.2 (Id-type maps). *A quasiconformal map $\varphi : \mathbb{C} \rightarrow \mathbb{C}$ is of id-type if there is an isotopy $\varphi_u : \mathbb{C} \rightarrow \mathbb{C}$, $u \in [0, 1]$ such that $\varphi_1 = \varphi$, $\varphi_0 = \text{id}$ and $|\varphi_u(z) - z| \rightarrow 0$ uniformly in u as $S_0 \ni z \rightarrow \infty$.*

Definition 5.3 (Id-type points in \mathcal{T}_f). *We say that $[\varphi] \in \mathcal{T}_f$ is of id-type if $[\varphi]$ contains an id-type map.*

Alternatively, we can replace S_0 in Definition 5.2 by endpoints of S_0 , i.e. P_f . In this case there are more *quasiconformal* maps identified as of id-type. Nevertheless, it is not hard to show that they define the same subset of id-type points in \mathcal{T}_f . Definition 5.2 is simply more suitable for our purposes.

Definition 5.4 (Isotopy of id-type maps). *We say that ψ_u is an isotopy of id-type maps if ψ_u is an isotopy through maps of id-type such that $|\psi_u(z) - z| \rightarrow 0$ uniformly in u as $S_0 \ni z \rightarrow \infty$.*

The identity is obviously of id-type. Since c^{-1} is equal to identity outside of a compact set, it is also of id-type, as well as the composition $\varphi \circ c$ of the capture with any id-type map.

Next theorem claims that the σ -map is invariant on the subset of id-type points in \mathcal{T}_f .

Theorem 5.5 (Invariance of id-type points). *If $[\varphi]$ is of id-type, then $\sigma[\varphi]$ is of id-type as well.*

More precisely, if φ is of id-type, then there is a unique id-type map $\hat{\varphi}$ such that $\varphi \circ f \circ \hat{\varphi}^{-1}$ is entire.

Moreover, if φ_u is an isotopy of id-type maps, then the functions $\varphi_u \circ f \circ \hat{\varphi}_u^{-1}$ have the form $e^z + \kappa_u$ where κ_u depends continuously on u .

Proof. First, since $\hat{\varphi}$ is defined up to a postcomposition with an affine map, and S_0 contains continuous curves joining finite points to ∞ , uniqueness is obvious. We only have to prove existence.

Since c^{-1} is also of id-type, φ can be joined to c^{-1} through an isotopy of id-type maps: we can simply take in the correct order the concatenation of two isotopies to id (concatenation of $\varphi \sim \text{id}$ and $\text{id} \sim c^{-1}$).

Let $\psi_u, u \in [0, 1]$ be this isotopy with $\psi_0 = c^{-1}$ and $\psi_1 = \varphi$ and let $\tilde{\psi}_u$ be the unique isotopy of quasiconformal homeomorphisms fixing 0, 1 and ∞ (or simply the solutions of the Beltrami equation 2.2 for $\mu_{\psi \circ f}$ normalized to fix 0, 1, ∞ .) Then the maps $h_u := \psi_u \circ f \circ \tilde{\psi}_u^{-1}$ will have the form

$$h_u(z) = \beta_u e^{\alpha_u z} + \kappa_u$$

where $\alpha_u, \beta_u, \kappa_u$ depend continuously on u .

Next define $\hat{\psi}_u(z) := \alpha_u \tilde{\psi}_u(z) + \log(\beta_u)$, where the branch of the logarithm is chosen so that $\hat{\psi}_0 = \text{id}$. In this case

$$g_u(z) = \psi_u \circ f \circ \hat{\psi}_u^{-1}(z) = e^z + \kappa_u$$

is the homotopy of entire functions such that $g_u \in \mathcal{N}$, $g_0 = f_0$ and κ_u is continuous in u and bounded on $[0, 1]$.

Now, let $z \in S_0$ and $w = f(z) = f_0(z)$. Since $|\psi_u(w) - w| \rightarrow 0$ as $S_0 \ni w \rightarrow \infty$ uniformly in u , we can write $\psi_u(w) = w + o(1)$.

Then

$$\hat{\psi}_u(z) = \log(\psi_u(w) - \kappa_u) = \log(f_0(z) - \kappa_u) = \log(e^z + \kappa_0 - \kappa_u).$$

Hence, due to continuity and boundedness of $\kappa_0 - \kappa_u$, for $S_0 \ni z \rightarrow \infty$ we have

$$|\hat{\psi}_u(z) - z| \rightarrow 0$$

uniformly in u .

The theorem is proven with the assignment $\hat{\varphi} := \hat{\psi}_1$. □

Theorem 5.5 particularly means that the notion of external address is preserved when we iterate id-type maps, in a sense that under proper normalization, that is, when we choose the map $\hat{\varphi}$ out of its Teichmüller equivalence class, the images

of dynamic rays under $\hat{\varphi}$ preserve their original asymptotics (horizontal straight lines).

Remark 5.6. *In the sequel we will use the hat-notation from Theorem 5.5. That is, $\hat{\varphi}$ denotes the unique id-type map so that $\varphi \circ f \circ \hat{\varphi}^{-1}$ is entire. Due to Theorem 5.5 this notation makes sense whenever φ is of id-type.*

6. SPIDERS

In this section we introduce the notion of a spider and a special tool which is supposed to bound the complexity of “spider legs” — altogether this allows to provide numerical estimates on the homotopy information of id-type points in the Teichmüller space. We start with the latter.

6.1. Curves in the punctured half-plane. We begin with the definition that slightly extends the notion of the homotopy of curves fixing endpoints.

Definition 6.1 (Pinched endpoints homotopy). *Let $U \subset \hat{\mathbb{C}}$ be a path-connected domain, and $\gamma : (0, \infty) \rightarrow U$ be a curve such that $\lim_{t \rightarrow 0} \gamma(t) \in \bar{U}$ and $\lim_{t \rightarrow \infty} \gamma(t) \in \bar{U}$.*

We denote by $[\gamma]_U$ the equivalence class of γ in U under homotopies preserving the limits at $t = 0$ and $t = \infty$.

If the limit of γ at 0 or ∞ belongs to U , this is the usual homotopy fixing endpoint, but when the limit is in the boundary of U , the homotopies are considered through curves in U that are “pinched” at the endpoint (possibly two) in the boundary.

We consider such “pinched” homotopy types for two reasons: first, it describes the homotopy type of the images of a ray tail (from S_0) under id-type maps representing the same point in the Teichmüller space; second, as will be seen later, we obtain estimates on the “complexity” of the homotopy information of id-type points.

Now, we define *r-curves* (“r” means right) which model ray tails (or more precisely, their union with ∞).

Definition 6.2 (r-curve). *We say that a continuous curve $\gamma : [0, \infty] \rightarrow \hat{\mathbb{C}}$ is an r-curve if*

- (1) $\gamma[0, \infty) \subset \mathbb{C}$,
- (2) $\gamma(\infty) = \infty$,
- (3) $\operatorname{Re} \gamma(t) \rightarrow +\infty$ as $t \rightarrow \infty$,

Let $V = \{v_i\} \subset \mathbb{C}$ be a finite or a countably infinite sequence (of punctures) such that $\operatorname{Re} v_i \rightarrow +\infty$, and let γ be an r-curve such that $\gamma(0) \in V$ and $\gamma|_{\mathbb{R}^+} \subset \mathbb{C} \setminus V$.

Remark 6.3. *In the case when V is infinite the surface $\mathbb{C} \setminus V$ is supposed to model the surface $\mathbb{C} \setminus P_f$.*

Choose a point $r \in \mathbb{R}$ such that

$$r < \min\left\{\min_{v \in V} \{\operatorname{Re} v\}, \min_{t \in [0, \infty]} \{\operatorname{Re} \gamma(t)\}\right\}.$$

Both V and $\gamma|_{\mathbb{R}^+}$ are contained in \mathbb{H}_r while $\gamma(\infty) = \infty \in \partial\mathbb{H}_r$.

We want to have an algebraic description of $[\gamma]_{\mathbb{H}_r \setminus V}$ (or more precisely $[\gamma|_{\mathbb{R}^+}]_{\mathbb{H}_r \setminus V}$), for instance, in terms of certain fundamental group. But the problem is that the endpoints of γ are punctures, that is, the homotopy is not allowed to pass through them. We cure this problem as follows.

Let

$$H_r = H_r(V, \gamma) := (\mathbb{H}_r \setminus V) \cup \{\infty\}.$$

For a curve $\delta : (0, \infty) \rightarrow \mathbb{C} \setminus V$ with $\lim_{t \rightarrow 0} \delta(t) \in V$ and $\lim_{t \rightarrow \infty} \delta(t) = \infty$ let $\bar{\delta} : (0, \infty] \rightarrow H_r$ be the extension of this curve to ∞ , that is, $\bar{\delta}(t) = \delta(t)$ for $t \in (0, \infty)$ and $\bar{\delta}(\infty) = \infty$.

Lemma 6.4. *For curves $\delta : (0, \infty) \rightarrow \mathbb{C} \setminus V$ with $\lim_{t \rightarrow 0} \delta(t) \in V$ and $\lim_{t \rightarrow \infty} \delta(t) = \infty$ the correspondence $[\delta]_{\mathbb{H}_r \setminus V} \mapsto [\bar{\delta}]_{H_r}$ is well defined and injective.*

Proof. That the correspondence is well defined follows from Definition 6.1.

Thus, we only need to prove that the curves δ_1 and δ_2 are homotopic in $\mathbb{H}_r \setminus V$ if $\bar{\delta}_1$ and $\bar{\delta}_2$ are homotopic in H_r .

Let $v \in V$ be the puncture at which start both $\bar{\delta}_1$ and $\bar{\delta}_2$. Let also

$$\Gamma : [0, \infty) \times [0, 1] \rightarrow H_r \cup \{v, \infty\}$$

with $\Gamma(t, 0) = \bar{\delta}_1$ and $\Gamma(t, 1) = \bar{\delta}_2$ be a homotopy between $\bar{\delta}_1$ and $\bar{\delta}_2$ in H_r , that is, for $u \in [0, 1]$ we have $\Gamma(0, u) = v$ and $\Gamma(\infty, u) = \infty$, and no inner point of the strip $B := [0, \infty) \times [0, 1]$ is mapped to v (but can be mapped to ∞).

Without loss of generality we might assume that $\Gamma([0, 1] \times [0, 1])$ is bounded, otherwise just rescale Γ in the first parameter. For every integer $n > 0$ let β_n be the restriction of Γ on $n \times [0, 1]$, that is, a trajectory of a point $\bar{\delta}_1(n)$ under the homotopy. Since the image under Γ of the boundary of the rectangle $[0, n] \times [0, 1]$ is contractible, each β_n is homotopic relative endpoints to a curve β'_n contained in $H_r \setminus \{\infty\}$. Moreover, we can make a joint choice of all β'_n so that $\beta'_n \rightarrow \infty$ as $n \rightarrow \infty$ (that is, the whole curve β'_n appears in arbitrarily small neighborhoods of ∞ in H_r when n is big).

Now, let $\alpha_n : \partial\mathbb{D} \rightarrow H_r \setminus \{\infty\}$ be the closed curve in $H_r \setminus \{\infty\}$ that is equal to the concatenation

$$\beta_n^{-1} \cdot \bar{\delta}_1|_{[n, n+1]} \cdot \beta_{n+1} \cdot (\bar{\delta}_2|_{[n, n+1]})^{-1}.$$

Each α_n is contractible in $H_r \setminus \{\infty\}$, so one can continuously extend α_n to the unit disk and obtain a map $\alpha'_n : \mathbb{D} \rightarrow H_r \setminus \{\infty\}$ that coincides with α_n on $\partial\mathbb{D}$. Moreover, since $\beta'_n \rightarrow \infty$, we can choose α'_n so that $\alpha'_n(\mathbb{D}) \rightarrow \infty$. After “gluing” all α'_n and the restriction $\Gamma|_{[0, 1] \times [0, 1]}$ together we obtain the desired homotopy.

This finishes the proof of the lemma. \square

Now we are ready to describe the homotopy type of γ . In the sequel we only consider the homotopy types in H_r and denote $[\gamma] = [\gamma]_{H_r}$.

Join $\gamma(0)$ to ∞ by the straight horizontal ray $\langle \gamma(0) + t, t > 0 \rangle$. For every intersection of this ray with some point $v \in V$, choose a small closed disk with center at v such that all disks are mutually disjoint and contain exactly one point of V (its center), and replace the parts of the straight line which are the diameters of the disks by the upper half-circles of the disk boundary.

The obtained curve γ_{str} defines the homotopy class $[\gamma_{str}]$ in H_r , containing curves that approximate “from above” the straight horizontal ray from $\gamma(0)$ to ∞ . Consider also a loop $\gamma_{loop} \subset H_r$ with the base point at ∞ such that $[\gamma_{str}] \cdot [\gamma_{loop}] = [\gamma]$ in H_r . It is not hard to show that such a loop exists: for example, one can slightly homotope γ in H_r so that for some $\epsilon > 0$ we have $\gamma|_{(0, \epsilon]}(t) = \gamma(0) + t$ — then $[\gamma_{loop}] = [(\gamma_{str}|_{[\epsilon, \infty)})^{-1}] \cdot [\gamma|_{[\epsilon, \infty)}]$. The loop homotopy type $[\gamma_{loop}]$ can be viewed as an element of the fundamental group $\pi_1(H_r, \infty)$. The homotopy class $[\gamma_{loop}]$ and the point $\gamma(0)$ uniquely define $[\gamma]$.

From now on we assume that V is finite. Any such V and an r-curve γ with $\gamma(0) \in V$ and $\gamma|_{(0, \infty)} \in \mathbb{C} \setminus V$ define the map

$$W : (V, \gamma) \mapsto W(V, \gamma) \in F(V),$$




FIGURE 6.1. Example of splitting γ (above) into γ_{str} and γ_{loop}

where $F(V)$ is the free group on V , as follows.

For every $w \in V$ construct a generator g_w of $\pi_1(H_r, \infty)$ in the following way. Surround each $v \in V$ lying on the straight horizontal ray $\langle w+t, t \geq 0 \rangle$ by a disk D_v with center at v and radius r_v , so that the disks are mutually disjoint and contain only one point of V . Define the injective curve $a : [0, \infty) \rightarrow H_r$ so that $a(t)$ is equal to $w + r_w + t$ when $w + r_w + t$ is outside of the disks D_v and coincides with the upper half-circles of ∂D_v whenever $w + r_w + t$ passes through D_v for $v \neq w$, and let $c(t) := w + r_w e^{-2\pi i t}$ for $t \in [0, 1]$. Then $g_w := [a^{-1} \cdot c \cdot a]$, and it does not depend on a particular choice of radii of the disks D_v . In other words, we start at ∞ , move “to the left” along a straight line passing “from above” the points $v \in V$ lying in our way, then make a loop around w in the clockwise direction, and return to ∞ along the same curve, passing occasional points in V “from above”.

With this preferred set of generators the homotopy type $[\gamma_{loop}]$ has the unique representation in $\pi_1(H_r, \infty)$. That is, $[\gamma_{loop}] = g_{v_1} g_{v_2} \dots g_{v_m}$ where $v_i \in V, m \in \mathbb{N}$. Define

$$W(V, \gamma) := v_1 v_2 \dots v_m.$$

It is clear that W does not depend on a particular choice of H_r (i.e. the choice of r).

Remark 6.5. *There is a freedom of choice for the generators and γ_{str} , at least in the sense of passing points of V below or above, so it is just according to our convention that we pass them from above.*

We finish this subsection with a few straightforward but useful lemmas.




FIGURE 6.2. Example of a generator g_w , for a case when $\text{Im } w = \text{Im } v$

Lemma 6.6. *Let $\gamma \subset \mathbb{C} \setminus V$ be an r -curve with $\gamma(0) \in V$. Further, let*

$$B := [x_1, \infty) \times [y_1, y_2]$$

be a (right-)infinite strip so that $\gamma|_{[0, \infty)} \subset B$. Denote $V_1 := V \cap B$. Then $W(V, \gamma) \in F(V_1)$, where $F(V_1)$ is identified with its image in $F(V)$ under the natural embedding.

Proof. Follows from the construction of $W(V, \gamma)$. □

By $|W(V, \gamma)|$ we denote the length of the word $W(V, \gamma)$ in a free group. It is $|W(V, \gamma)|$ that will soon allow us to measure the “complexity” of the homotopy information of id-type points in the Teichmüller space.

Lemma 6.7. *If $V_1 \subset V$ and $\gamma(0) \in V_1$, then $|W(V_1, \gamma)| \leq |W(V, \gamma)|$.*

Proof. We can first find $W(V, \gamma)$ and then “glue in” the punctures from $V \setminus V_1$. Then some generators of the fundamental group in $H_r(V)$ become trivial. So the length of $W(V_1, \gamma)$ cannot be bigger than the length of $W(V, \gamma)$. □

Lemma 6.8. *Let $\gamma \subset \mathbb{C} \setminus V$ be an r -curve with the strictly increasing real part and such that $\gamma(0) \in V$. Then $|W(V, \gamma)| < C_{|V|}$ where $C_{|V|} > 0$ depends only on the size of V .*

Proof. Follows from the construction of $W(V, \gamma)$. □

6.2. H_r and Hawaiian earrings. This is a minor subsection containing a rather standing aside fact that we will need later.

Let H_r be defined as in Subsection 6.1 for a countably infinite V . Clearly we can choose such representatives g'_{v_n} of the “straight” generators g_{v_n} (defined similarly as for finite V) so that each of them is a Jordan curve containing ∞ , they are pairwise disjoint except at ∞ , and each g'_{v_n} is a concatenation $\alpha_n^{-1} \cdot \beta_n$, where α_n, β_n are horizontal paths with the strictly increasing real part, and parametrized by it.

Recall that the *Hawaiian earring* is the topological space homeomorphic to $\mathcal{H} = \cup C_n$, where $C_n = \{z \in \mathbb{C} : |z - 1/n| = 1/n\}$ for $n \in \mathbb{N}$.

Lemma 6.9 (Hawaiian earring skeleton). *Let $H := \cup g'_{v_n}$. Then H is homeomorphic to \mathcal{H} and is a deformation retract of H_r .*

Sketch of the proof. We can construct the homeomorphism $\psi : H \rightarrow \mathcal{H}$ by mapping homeomorphically the consecutive generators g'_{v_n} to the consecutive outer circles C_n so that $\psi(g'_{v_n}) = C_n$, $\psi(\infty) = 0$, and $\psi(\alpha_n(t)), \psi(\beta_n(t)) \rightarrow 0$ as $t \rightarrow \infty$ uniformly in n whenever defined.

Such ψ is a continuous bijection. Since H is compact and \mathcal{H} is Hausdorff, ψ is homeomorphism.

The second statement of the lemma follows easily from the fact that the punctured closed unit disk deformation retracts onto its outer boundary. \square

6.3. Homotopy type of a preimage. We are now interested in how the homotopy type, and in particular $|W(V, \gamma)|$, changes after taking preimage under an entire map. Although to prove Classification Theorem 1.1 we only need to consider the exponential function, we study (without essential additional efforts) the case of compositions of a polynomial with the exponential, which will be in our focus in the subsequent articles devoted to more general families. We begin with two preliminary lemmas and then provide bounds for $|W(V, \tilde{\gamma})|$ of a preimage curve $\tilde{\gamma}$.

Let $V \subset \mathbb{C}$ be a *finite* set, and let γ be an r -curve such that $\gamma(0) \in V$ and $\gamma|_{\mathbb{R}^+} \subset \mathbb{C} \setminus V$. As earlier choose $H_r = H_r(V, \gamma)$. We again consider homotopy types of γ in H_r .

Lemma 6.10 (Piecewise linear representative). *There exists an r -curve $\gamma_{pl} \in [\gamma]$ consisting of at most $6|W(V, \gamma)| + 2$ straight line segments.*

Proof. Let $m = |W(V, \gamma)|$. Take a generator $[g_w]$ of the fundamental group $\pi_1(H_r, \infty)$ for some $w \in V$. There exists a representative of the generator $[g_w]$ consisting of 5 straight line segments: 1 segment to pass from ∞ to a small neighborhood of w , 3 to make a triangular loop around w in this small neighborhood, and 1 to return back to ∞ .

Analogously $[\gamma_{str}]$ has an representative consisting of at most 2 straight line segments.

Using the presentation of $[\gamma] = [\gamma_{str}] \cdot [\gamma_{loop}]$ and a presentation of $[\gamma_{loop}] = g_{v_1} \dots g_{v_m}$ via m generators, we already have a piece-wise linear representative of $[\gamma]$ containing at most $5m + 2$ straight line segments. Nevertheless, this representative is not an r -curve because it has some midpoints at ∞ : the endpoints of the generators g_{v_i} . But this flaw can be easily adjusted by the ‘‘price’’ of at most m straight line segments: if $z_1, z_2 \in \mathbb{C}$ are such that $\text{Re } z_1 = \text{Re } z_2 > \max_{v \in V} \text{Re } v$, then the concatenation

$$\langle z_1 + t, t \in [0, \infty] \rangle \cdot \langle z_2 + t, t \in [0, \infty] \rangle^{-1}$$

of horizontal rays is homotopic to a straight line segment between z_1 and z_2 . This way we replace pairs of neighboring infinite rays on the generators by Π -shaped parts.

Altogether this means that $[\gamma]$ has an r -curve representative consisting of at most $5m + 2 + m = 6m + 2$ straight line segments. \square

Lemma 6.11 (Intersections bound homotopy type). *For every point $w \in V$, consider the ray $L_w = \langle w - t, t > 0 \rangle$. Assume that $\gamma|_{(0, \infty)}$ intersects L_w at $k_w < \infty$ points, and let $k = \sum_V k_w$. Then*

$$|W(V, \gamma)| \leq (k + 1)|V|.$$

Proof. Let $0 < t_1 < \dots < t_m < \infty$ be the values of t for which $\gamma|_{(0, \infty)}$ intersects at least one of the L_w . Then $m \leq k$. It can also happen that $m < k$ since γ might intersect a few L_u at the same time.

Choose $s_0 \in (0, t_1)$, $s_1 \in (t_1, t_2), \dots, s_{m-1} \in (t_{m-1}, t_m)$, $s_m \in (t_m, \infty)$, and define

$$\gamma_0 = \gamma|_{[0, s_0]},$$

$$\gamma_1 = \gamma|_{[s_0, s_1]},$$

...

$$\gamma_m = \gamma|_{[s_{m-1}, s_m]},$$

$$\gamma_{m+1} = \gamma|_{[s_m, \infty]},$$

so we have $\gamma = \gamma_0 \cdot \gamma_1 \cdot \dots \cdot \gamma_{m+1}$.

For $z \in \mathbb{C}$ denote by δ_z the infinite ray $\langle z + t, t \in [0, \infty) \rangle$. Then

$$[\gamma] = [\gamma_0 \cdot \delta_{s_0}] \cdot [\delta_{s_0}^{-1} \cdot \gamma_1 \cdot \delta_{s_1}] \cdot \dots \cdot [\delta_{s_{m-1}}^{-1} \cdot \gamma_m \cdot \delta_{s_m}] \cdot [\delta_{s_m} \cdot \gamma_{m+1}].$$

Each $[\delta_{s_{i-1}}^{-1} \cdot \gamma_i \cdot \delta_{s_i}]$ for $i = \overline{1, m}$ is homotopic either to a consecutive product (possibly with inverses) of a few generators g_v without repetitions, or it is null-homotopic.

Further, $[\gamma_0 \cdot \delta_{s_0}]$ is homotopic either to γ_{str} , or to a concatenation with a consecutive product (possibly with inverses) of generators g_v without repetitions, whereas $[\delta_{s_m} \cdot \gamma_{m+1}]$ is evidently null-homotopic.

Hence

$$|W(V, \gamma)| \leq m|V| + |V| \leq (k+1)|V|.$$

□

Now we are ready to describe how $|W(V, \gamma)|$ behaves after taking preimages under the exponential or a polynomial. The estimates are justified by the following elementary lemma.

Lemma 6.12 (Preimages stay horizontal). *Let f be either the exponential or a monic polynomial, $\gamma : (0, \infty) \rightarrow \hat{\mathbb{C}} \setminus \{\text{singular values of } f\}$ be an r -curve, and $\tilde{\gamma}$ be one of its lifts under f . The following statements hold.*

- (1) *If f is the exponential, then $\tilde{\gamma}$ is an r -curve.*
- (2) *If f is a monic polynomial of degree $d \geq 1$, then there exists a degree d root of unity ξ such that $\xi\tilde{\gamma}$ is an r -curve.*

Thus, it makes sense to speak about how the homotopy type of an r -curve changes under lifts.

Proposition 6.13 (Bounds for lifts by polynomials). *Let p be a monic polynomial of degree $d \geq 1$ with its critical values contained in V , and let the set \tilde{V} be such that $p(\tilde{V}) \subset V$ and $p|_{\tilde{V}}$ is bijective. Denote by $\tilde{\gamma}$ the homeomorphic r -curve preimage of γ under p .*

If $n = |V|$, and $m = |W(V, \gamma)|$, then

$$|W(\tilde{V}, \tilde{\gamma})| < 6dn^2(m+1).$$

Proof. Choose some $H_{r'} = H_{r'}(\tilde{V}, \tilde{\gamma})$ so that $p(\mathbb{H}_{r'}) \supset \mathbb{H}_r$. Then every homotopy of γ in H_r lifts to a homotopy of $\tilde{\gamma}$ in $H_{r'}$.

Next, let $\gamma_{pl} \in [\gamma]$ be the piecewise linear representative consisting of at most $6m+2$ straight line segments as in the Lemma 6.10 and denote by δ its lift under p which is r -curve. Then δ is homotopic to $\tilde{\gamma}$ in $H_{r'}$.

Elementary computation shows that a preimage of a straight line segment of γ_{pl} under polynomial either intersects each straight line containing an L_w in no more than d points, or is a subset of it. If the second case takes place for some preimages of bounded segments of γ_{pl} , we slightly deform γ_{pl} (in its homotopy class) so that the first condition holds. Note that since δ is horizontal, if the preimage of the unique infinite straight segment of γ_{pl} is a subset of a horizontal straight line, then

it does not intersect any L_w . Hence without loss of generality we can assume that $\delta|_{(0,\infty)}$ intersects each $L_w, w \in \tilde{V}$ (see Lemma 6.11) in at most $d(6m+2)$ points.

Then due to Lemma 6.11

$$\begin{aligned} |W(\tilde{V}, \tilde{\gamma})| &= |W(\tilde{V}, \delta)| \leq (nd(6m+2) + 1)n = \\ 6mn^2d + 2n^2d + n &< 6mn^2d + 6n^2d = 6dn^2(m+1). \end{aligned}$$

□

We are interested in an analogous result for the exponential.

Proposition 6.14 (Bounds for lifts by the exponential). *Let $0 \in V$, $\gamma(0) \neq 0$ and let \tilde{V} be such that $\exp(\tilde{V}) \subset V$, and $\exp|_{\tilde{V}}$ is injective. Denote by $\tilde{\gamma}$ the homeomorphic r -curve preimage of γ under the exponential.*

If $n = |V| = |\tilde{V}| + 1$, and $m = |W(V, \gamma)|$, then

$$|W(\tilde{V}, \tilde{\gamma})| < 6n^2(m+1).$$

Remark 6.15. *Note that in this case $|V| = |\tilde{V}| + 1$ since asymptotic value of \exp does not have a preimage.*

Proof. The proof is identical to the case of a polynomial in Proposition 6.13, except that a preimage of a straight line segment under the exponential either intersects L_w in at most *one* point (instead of d), or is a subset of it. That is why the formula for the exponential coincides with the formula for polynomials of degree $d = 1$. □

At the end of this subsection we provide bounds for how $W(V, \gamma)$ changes under lifts by compositions of a polynomial and the exponential.

Theorem 6.16 (Bounds for lifts by $p \circ \exp$). *Let p be a monic polynomial of degree d , and $g = p \circ \exp$.*

Further, let $V \subset \mathbb{C}$ be a finite set containing singular values of g and $\gamma : (0, \infty) \rightarrow \hat{\mathbb{C}}$ be an r -curve such that $\gamma(0) \subset V \setminus \{\text{singular values of } g\}$ and $\gamma|_{\mathbb{R}^+} \subset \mathbb{C} \setminus V$.

If $\tilde{V} \subset \mathbb{C}$ is a finite set such that $g(\tilde{V}) \subset V \setminus \{\text{singular values of } g\}$, and $g|_{\tilde{V}}$ is injective, then for every r -curve preimage $\tilde{\gamma}$ of γ under g holds

$$|W(\tilde{V}, \tilde{\gamma})| < 42dn^4(m+1)$$

where $n = |V|$ and $m = |W(V, \gamma)|$.

Proof. We obtain the estimate simply by using consecutively Proposition 6.13 and Proposition 6.14 with the same $|V| = n$.

$$|W(\tilde{V}, \tilde{\gamma})| < 6n^2(6dn^2(m+1) + 1) \leq 42dn^4(m+1).$$

□

6.4. Spiders. Now return to our usual setting with the captured exponential function $f := c \circ f_0$ defined as in Subsection 3.2. Moreover, we assume that the external address of a_0 is not (pre-)periodic.

Definition 6.17 (Spider). *An image S_φ of the standard spider S_0 under an id-type map φ is called a spider.*

The notion of a spider comes with a few natural follow-up definitions.

Definition 6.18 (Spider legs). *The image of a ray tail R_n under a spider map is called a leg (of a spider).*

Definition 6.19 (Subspider). *For a spider S_φ , a non-empty union of its legs is called a subspider of S_φ .*

We also want to have equality relation on the set of spiders.

Definition 6.20 (Equal spiders). *We say that the spiders S_φ and S_ψ are equal if for all $n \geq 0$ we have $\varphi(R_n) = \psi(R_n)$ (as point sets).*

Lemma 6.21 (Equal spiders define the same point in \mathcal{T}_f). *Let $S_\varphi = S_\psi$. Then φ is isotopic to ψ relative S_0 , and consequently $[\varphi] = [\psi]$.*

Proof. We might assume that $\varphi|_{S_0} = \psi|_{S_0}$, otherwise just isotope φ in disjoint neighborhoods of the legs $\{\varphi(R_n)\}$ so that the legs get reparametrized. Consider the map $\psi \circ \varphi^{-1}$. It is equal to identity on the locally connected boundary of the simply-connected domain $\mathbb{C} \setminus \varphi(S_0)$. Hence $\psi \circ \varphi^{-1}|_{\mathbb{C} \setminus \varphi(S_0)}$ is isotopic to identity relative $\varphi(S_0)$. This implies that $[\varphi] = [\psi]$. \square

Denote $\mathcal{O}_n = \{a_i \in P_f : i \leq n\}$.

Definition 6.22 (Leg homotopy word). *Let S_φ be a spider. Then the leg homotopy word W_n^φ of $\varphi(R_n)$ is*

$$W_n^\varphi := W(\varphi(\mathcal{O}_n), \varphi(R_n)).$$

Next theorem helps to estimate how the leg homotopy words change under Thurston iteration.

Theorem 6.23 (Combinatorics of preimage). *Let φ be of id-type. Then*

$$|W_n^{\hat{\varphi}}| < A(n+2)^4 \max\{1, |W_{n+1}^\varphi|\},$$

where A is a positive real number.

Proof. This is just a restatement of Theorem 6.16 in a particular context. That is, when $V = \varphi(\mathcal{O}_{n+1})$, $\gamma = \varphi(R_{n+1})$ and $W(V, \gamma) = W_{n+1}^\varphi$. Then if we take $\tilde{\gamma} = \hat{\varphi}(R_n)$,

$$|W_n^{\hat{\varphi}}| < 42(n+2)^4(|W_{n+1}^\varphi| + 1) < A(n+2)^4 \max\{|W_{n+1}^\varphi|, 1\},$$

where $A > 0$ is a positive real number. \square

6.5. Spiders and Teichmüller equivalence. In the last subsection we prove the results showing that the homotopy type of id-type maps can be uniquely encoded using spiders. More precisely, for a map φ of id-type one can uniquely recover its equivalence class $[\varphi] \in \mathcal{T}_f$ from the positions of points in $\varphi(P_f)$ and W_n^φ for all $n \geq 0$.

Definition 6.24 (Equivalence of spiders). *We say that two spiders S_φ and S_ψ are equivalent if for all $n \geq 0$ we have $\varphi(a_{ij}) = \psi(a_{ij})$ and legs with the same index are homotopic, i.e. $[\varphi(R_n)] = [\psi(R_n)]$ in $\mathbb{C} \setminus \varphi(P_f)$.*

Proposition 6.25 (Spider equivalence is Teichmüller equivalence). *Two spiders S_φ and S_ψ are equivalent if and only if $[\varphi] = [\psi]$, i.e. φ is isotopic to ψ relative P_f .*

Proof. (\Leftarrow) “If” direction follows directly from the definitions.

(\Rightarrow) Assume that the spiders S_φ and S_ψ are equivalent.

We need to prove that S_φ can be isotoped into S_ψ via an ambient isotopy of $\mathbb{C} \setminus P_f$. The statement of the proposition will then follow due to Lemma 6.21. To simplify the notation we assume that $\varphi = \text{id}$, i.e. $S_\varphi = S_0$ (in the general case the proof is exactly the same).

The proof uses the same idea as Alexander’s trick, where we shrink all “entanglements” into one point.

First, note that one can introduce a natural linear order on the set of the legs of S_0 : for every pair of legs R_i and R_j we can say which one is *higher* or *lower*.

More precisely, choose a right half-plane \mathbb{H}_r containing both R_i and R_j and join a_i , the endpoint of R_i , to a finite point of $\partial\mathbb{H}_r$ via an injective path γ inside of $\mathbb{H}_r \setminus (R_i \cup R_j)$. The union of two curves γ and R_i divides \mathbb{H}_r into two parts. We say that R_j is *higher* (resp. *lower*) than R_i if it is contained inside of the upper (resp. lower) part.

For every $n \geq 0$ there exists a minimal number $k_n \geq 1$ so that $S_0 \setminus \cup_{i=0}^n R_i$ is a disjoint union of k_n subspiders S_n^j , $j \in \{1, \dots, k_n\}$ such that for every $j \in \{1, \dots, k_n\}$ and $i \leq n$ all legs of the subspider S_n^j are simultaneously higher or lower than R_i . Clearly, $k_n \leq n + 2$.

Let U be a right half-plane containing S_0 . Now define inductively a sequence of shrinking Jordan domains contained in U .

- **(Step 0)** Define k_0 mutually disjoint Jordan domains U_0^j so that

$$S_0^j \subset U_0^j \subset U$$

and the diameter of U_0^j in the spherical metric is less than the diameter of S_0^j plus 1.

- **(Step n)** Define k_n mutually disjoint Jordan domains U_n^j so that for some $i \leq k_{n-1}$

$$S_n^j \subset U_n^j \subset U_{n-1}^i$$

and the diameter of U_n^j in the spherical metric is less than the diameter of S_n^j plus $1/(n+1)$.

We are going to construct the isotopy ψ_u^∞ which is the concatenation of countably many “local” isotopies ψ_u^n , $n \geq 0$. Roughly speaking, via ψ_u^0 we isotope $\psi(R_0)$ into R_0 , then via ψ_u^1 we isotope $\psi_u^0(R_1)$ relative P_f into R_1 without moving R_0, \dots , then via ψ_u^n we isotope $\psi_u^{n-1}(R_n)$ into R_n without moving R_0, R_1, \dots, R_{n-1} , and so on.

We construct this sequence of isotopies inductively. Without loss of generality assume that $\psi(S_0) \subset U$ and $\psi|_{\mathbb{C} \setminus U} = \text{id}$. Note that every leg $\psi(R_n)$ is homotopic to R_n inside of $U \setminus P_f$ (consequence of the fact that ψ is of id-type).

- **(Step 0)** Let ψ_u^0 , $u \in [0, 1]$ be an isotopy of $\hat{\mathbb{C}} \setminus P_f$ (or equivalently of $\mathbb{C} \setminus P_f$) so that
 - (1) $\psi_u^0 = \psi$,
 - (2) $\psi_u^0|_1 = \text{id}$ on $\mathbb{C} \setminus \cup_{j=1}^{k_0} U_0^j$,
 - (3) $\psi_u^0 \equiv \text{id}$ on $\mathbb{C} \setminus U$.
- **(Step n)** Let ψ_u^n , $u \in [0, 1]$ be an isotopy of $\hat{\mathbb{C}} \setminus P_f$ so that
 - (1) $\psi_u^n = \psi_u^{n-1}$,
 - (2) $\psi_u^n|_1 = \text{id}$ on $\mathbb{C} \setminus \cup_{j=1}^{k_n} U_n^j$,
 - (3) $\psi_u^n \equiv \text{id}$ on $\mathbb{C} \setminus \cup_{j=1}^{k_{n-1}} U_{n-1}^j$.

Note that every Step n is well-defined since the preceding Step $n-1$ guarantees that the leg $\psi(R_n)$ is contained inside of some U_{n-1}^j and is homotopic to R_n inside of $U_{n-1}^j \setminus P_f$.

In order to define the concatenation of these infinitely many isotopies we reparametrize them so that ψ_u^n is defined on the interval $[1 - 1/2^n, 1 - 1/2^{n+1}]$. As $n \rightarrow \infty$, maps φ_u^n are not equal to identity only in neighborhoods of ∞ that are getting smaller. Hence the infinite concatenation $\psi_u^\infty = \psi_u^0 \cdot \psi_u^1 \cdot \dots$ is well defined. This finishes the proof of the proposition. \square

Definition 6.26 (Projective equivalence of spiders). *We say that two spiders S_φ and S_ψ are projectively equivalent if for all $n \geq 0$ we have $\varphi(a_n) = \psi(a_n)$ and $W_n^\varphi = W_n^\psi$.*

Proposition 6.27 (Projectively equivalent spiders are equivalent). *Two spiders S_φ and S_ψ are projectively equivalent if and only if they are equivalent.*

Proof. (\Leftarrow) If $\varphi(R_n)$ and $\psi(R_n)$ are homotopic relative to $\psi(P_f)$, then they are homotopic relative any subset of $\psi(P_f)$.

(\Rightarrow) Assume now that the spiders S_φ and S_ψ are projectively equivalent. Let $r \in \mathbb{R}$ be such that \mathbb{H}_r contains both S_φ and S_ψ , and consider $H_r = H_r(\varphi(P_f), \varphi(R_0))$ (defined as before Lemma 6.4).

We want to prove that the spider legs $\varphi(R_n)$ and $\psi(R_n)$ belong to the same homotopy class in H_r (the condition $W_n^\varphi = W_n^\psi$ says only that $\varphi(R_n)$ and $\psi(R_n)$ are homotopic in H_r relative to only a finite set of punctures $\varphi(\{a_i\}_{i=0}^n)$).

We claim that the homotopy type of each $\varphi(R_n)$ in H_r is uniquely defined by the knowledge of W_k^φ for all $k \geq n$.

Pick $k \geq n$. Assume that we know the homotopy type of $\varphi(R_n)$ in the “finitely-punctured” $H_r \cup \varphi(P_f \setminus \{a_i\}_{i=0}^k)$ (for $k = n$ we have it via W_n^φ) and know W_{k+1}^φ . Since $R_{k+1} \cap R_n = \emptyset$ and φ is of id-type, the homotopy type of $\varphi(R_n)$ in $H_r \cup \varphi(P_f \setminus \{a_i\}_{i=0}^{k+1})$ is uniquely defined.

Next, from Lemma 6.9 we know that H_r deformation retracts onto a subspace, homeomorphic to the Hawaiian earring $\mathcal{H} = \cup C_n$, so that the “straight” generator g_{v_n} of $\pi_1(H_r, \infty)$ is mapped to an element of $\pi_1(\mathcal{H}, 0)$ represented by the circle C_n . It is well-known fact about Hawaiian earrings (e.g. see [dS]) that $\pi_1(\mathcal{H}, 0)$ is a subgroup of $\varprojlim F_j$, where F_j is the fundamental group of $\cup_1^j C_i$ with the base point at 0, considered as a subgroup of $\pi_1(\mathcal{H}, 0)$ after collapsing all other C_i to 0. Hence, if $\theta_1, \theta_2 \in \pi_1(\mathcal{H}, 0)$ are such that $\theta_1 = \theta_2$ in F_j for all j , then $\theta_1 = \theta_2$ in $\pi_1(\mathcal{H}, 0)$. From this follows that the homotopy type of $\varphi(R_n)$ in H_r is uniquely determined by its homotopy types in every $H_r \cup \varphi(P_f \setminus \{a_i\}_{i=0}^k)$ for $k \geq n$. \square

From Proposition 6.25 and Proposition 6.27 follows the theorem that identifies the projective equivalence of spiders and Teichmüller equivalence.

Theorem 6.28 (Projective equivalence of spiders is Teichmüller equivalence). *Two spiders S_φ and S_ψ are projectively equivalent if and only if $[\varphi] = [\psi]$, i.e. φ is isotopic to ψ relative P_f .*

Remark 6.29. *Note that the only properties of S_0 that we used in the proofs of Lemma 6.21 and Propositions 6.25, 6.27 are that S_0 is contained in some right half-plane and all R_n are mutually disjoint (except at ∞) injective r -curves with the spherical diameters tending to 0 as $n \rightarrow \infty$. Hence the statements are obviously true for every such more general choice of S_0 (and after replacement of P_f by the set of endpoints of S_0).*

7. INVARIANT COMPACT SUBSET

In this section we construct a compact invariant subset \mathcal{C}_f of the Teichmüller space \mathcal{T}_f . In the first theorem we present the construction and prove invariance. A statement that \mathcal{C}_f is compact will be the content of the second theorem.

Let $f = c \circ f_0$ be the captured exponential function constructed in Subsection 3.2, and for $n \geq 0$ let t_n be the potential of the point $a_n \in P_f$ (then $t_{n+1} = F(t_n)$).

Define the set $\mathcal{P}' := \{\frac{t_{n+1} + t_n}{2} : n \geq 0\}$ and if $\rho = \frac{t_{n+1} + t_n}{2} \in \mathcal{P}'$, let $N(\rho) := n$. Further, define $D_\rho := \mathbb{D}_\rho(0)$ and $M_\rho := e^{2t_{N(\rho)}}$.

The subset \mathcal{C}_f is constructed in the form of a list of conditions.

Theorem 7.1 (Invariant subset). *Fix some $\rho \in \mathcal{P}'$ and denote $N := N(\rho)$. Let $\mathcal{C}_f(\rho) \subset \mathcal{T}_f$ be the closure of the set of points in \mathcal{T}_f represented by id-type maps*

φ for which there exists an isotopy $\varphi_u, u \in [0, 1]$ of id-type maps such that $\varphi_0 = \text{id}, \varphi_1 = \varphi$, and the following conditions are simultaneously satisfied.

(1) (Marked points stay inside of D_ρ) If $n \leq N$,

$$\varphi_u(a_n) \in D_\rho.$$

(2) (Precise asymptotics outside of D_ρ) If $n > N$, then

$$|\varphi_u(a_n) - a_n| < 1/n.$$

(3) (Separation inside of D_ρ) If $k < l \leq N$, then

$$|\varphi_u(a_k) - \varphi_u(a_l)| > \frac{1}{(M_\rho)^{N-l+1}}.$$

(4) (Bounded homotopy) If $n \leq N$, then

$$|W_n^{\varphi_u}| < A^{N-n+1} \left(\frac{(N+2)!}{(n+1)!} \right)^4$$

where A is the constant from Theorem 6.23.

Then if $\rho \in \mathcal{P}'$ is big enough, the subset $\mathcal{C}_f(\rho)$ is well-defined, invariant under the σ -map and contains [id].

Let us briefly discuss the content of the theorem before proving it. Our invariant subset (and compact, as will be seen later) is described as a closure of a set of id-type maps φ satisfying conditions (1)-(4).

Conditions (1)-(2) say that the maps φ have to be “uniformly of id-type”, that is, the marked points outside of a disk D_ρ have precise asymptotics, while inside of D_ρ we allow some more freedom.

Condition (3) tells us that the marked points inside of D_ρ cannot come very close to each other — this is necessary to keep our set bounded in the Teichmüller space. Moreover, it is needed to control the distance to the asymptotic value — if a marked point is too close to it, then after Thurston iteration its preimage has the real part close to $-\infty$, and this spoils condition (1).

Condition (4) takes care of the homotopy information and provides bounds for leg homotopy words of points inside of D_ρ . Note that similar bounds for marked points outside of D_ρ are encoded implicitly in condition (2).

Proof of Theorem 7.1. Let $\mathcal{C}_f^\circ(\rho) \subset \mathcal{C}_f(\rho)$ be the set of points in \mathcal{T}_f of which we take the closure in the statement of the theorem, i.e. represented by id-type maps φ for which there exists an isotopy φ_u of id-type maps such that $\varphi_0 = \text{id}, \varphi_1 = \varphi$ and the conditions (1)-(4) are simultaneously satisfied. Since σ -map is continuous, it is enough to prove invariance of $\mathcal{C}_f^\circ(\rho)$ for big ρ .

First, note that for big ρ the set $\mathcal{C}_f^\circ(\rho)$ contains $[c^{-1}]$: c^{-1} can be joined to identity via the isotopy c_u^{-1} where c_u was constructed in Subsection 3.2.

Further, from the asymptotic formula 2.1 and the fact that $\text{Re } a_n / \text{Im } a_n \rightarrow \infty$ follows that for all $\rho \in \mathcal{P}'$ big enough the first $N+1$ point a_0, a_1, \dots, a_N of the orbit P_f are contained in D_ρ , while the other points are outside and mutual distances between them and from them to D_ρ are bigger than one. In other words, conditions (1)-(2) allow the first $N+1$ points to move under isotopy φ_u only inside of D_ρ , while for every $n > N$ the point a_n moves inside of a disk D_n of radius $1/n$, and all these disks $D_\rho, D_{N+1}, D_{N+2}, \dots$ are mutually disjoint with the mutual distance between them bigger than one. Moreover, we can assume that for $n > N$ we have $\text{Re } a_n + 1 < \text{Re } a_{n+1}$.

For all $\varphi \in \mathcal{C}_f^\circ(\rho)$ after concatenation with c_u^{-1} we obtain the isotopy ψ_u of id-type maps with $\psi_0 = c^{-1}$, $\psi_1 = \varphi$ and satisfying conditions (1)-(4). Then $\hat{\psi}_u$ is an isotopy of id-type maps with $\hat{\psi}_1 = \text{id}$. Let $g_u(z) = \psi_u \circ f \circ \hat{\psi}_u^{-1}(z) = e^z + \kappa_u$. Now we want to prove that $\hat{\psi}_u$ satisfies each of the items (1)-(4): from this would follow that $\hat{\varphi} \in \mathcal{C}_f^\circ(\rho)$.

We prove that each of the conditions (1)-(4) for $\hat{\varphi}$ follows from the conditions (1)-(4) for φ .

(4) Note that since for $n > N$ we have $\text{Re } a_n + 1 < \text{Re } a_{n+1}$, for $n > N$ we have

$$|W_N^{\psi_u}| = 0 < 1.$$

Hence from Theorem 6.23 for $n \leq N$ we get

$$\begin{aligned} |W_n^{\hat{\psi}_u}| &< A(n+2)^4 \max\{|W_{n+1}^{\psi_u}|, 1\} < A(n+2)^4 A^{N-(n+1)+1} \left(\frac{(N+2)!}{(n+2)!}\right)^4 = \\ &A^{N-n+1} \left(\frac{(N+2)!}{(n+1)!}\right)^4. \end{aligned}$$

(3) We use the formula $\psi_u(a_{n+1}) = e^{\hat{\psi}_u(a_n)} + \kappa_u$. Since $\rho = \frac{t_N + t_{N-1}}{2}$ and $\kappa_u \in D_\rho$, for $n \leq N$ and big ρ we have

$$\begin{aligned} \text{Re } \hat{\psi}_u(a_n) &= \log|\psi_u(a_{n+1}) - \kappa_u| < \log|t_{N+1} + 1 + \rho| < \\ &\log(2t_{N+1}) = \log(2F(t_N)) < 2t_N. \end{aligned}$$

Thus for $k < l \leq N$ and big enough ρ

$$|\psi_u(a_{k+1}) - \psi_u(a_{l+1})| = \left| \int_{\hat{\psi}_u(a_k)}^{\hat{\psi}_u(a_l)} e^z dz \right| \leq \int_{\hat{\psi}_u(a_k)}^{\hat{\psi}_u(a_l)} |e^z| |dz| < M_\rho |\hat{\psi}_u(a_k) - \hat{\psi}_u(a_l)|,$$

and

$$|\hat{\psi}_u(a_k) - \hat{\psi}_u(a_l)| > \frac{|\psi_u(a_{k+1}) - \psi_u(a_{l+1})|}{M_\rho} > \frac{1}{(M_\rho)^{N-l+1}}.$$

(2) Since $\hat{\psi}_u$ is continuous in u and $\hat{\psi}_0 = \text{id}$, for $n > N$

$$\begin{aligned} \hat{\psi}_u(a_n) &= \log(\psi_u(a_{n+1}) - \kappa_u) = \log(a_{n+1} + O(1) - \kappa_u) = \\ &a_n + \log\left(\frac{a_{n+1} + O(1) - \kappa_u}{e^{a_n}}\right) = a_n + \log\left(1 + \frac{\kappa_0 + O(1) - \kappa_u}{a_{n+1} - \kappa_0}\right). \end{aligned}$$

If ρ (and consequently N) is big enough, we have $|\hat{\psi}_u(a_n) - a_n| < 1/n$ for all $n > N$ (due to the very fast growth of $|a_n|$).

(1) Instead of proving directly that for $n \leq N$ holds $\hat{\psi}_u(a_n) \in D_\rho$, we prove that for big ρ

$$-\frac{\rho}{2} < \text{Re } \hat{\psi}_u(a_n) < \frac{\rho}{2}$$

and

$$-\frac{\rho}{2} < \text{Im } \hat{\psi}_u(a_n) < \frac{\rho}{2}.$$

From this would follow that $|\hat{\psi}_u(a_n)| < \rho/\sqrt{2} < \rho$.

In the proof of (3) we have shown that for big ρ holds $\text{Re } \hat{\psi}_u(a_n) < 2t_N$. Clearly, $2t_N < \rho/2$ for big ρ .

Using condition (3) for ψ_u we get

$$\text{Re } \hat{\psi}_u(a_n) = \log|\psi_u(a_{n+1}) - \kappa_u| > \log \frac{1}{M_\rho^n} = -2nt_N$$

which is bigger than $-\rho/2$ for all ρ big enough.

Since $|W_{n+1}^{\psi_u}| < A^{N-n} \left(\frac{(N+2)!}{(n+2)!} \right)^4$, the spider leg $\hat{\psi}_u(R_{n+1})$ makes no more than $A^{N-n} \left(\frac{(N+2)!}{(n+2)!} \right)^4 + 1$ loops around the singular value. Hence the difference between $2\pi s_n$ and the imaginary part of $\hat{\psi}_u(a_n)$ will be less than $2\pi \left(A^{N-n} \left(\frac{(N+2)!}{(n+2)!} \right)^4 + 1 \right)$. Due to fast growth of the sequence $\{t_n\}_{n=0}^\infty$, by making ρ bigger we may assume that for all $n \leq N$ we have

$$2\pi \left(A^{N-n} \left(\frac{(N+2)!}{(n+2)!} \right)^4 + 1 + |s_n| \right) < \frac{\rho}{2}.$$

Hence $|\operatorname{Im} \hat{\psi}_u(a_n)| < \rho/2$ and $|\hat{\psi}_u(a_n)| < \rho/\sqrt{2} < \rho$. \square

Next, we prove another key statement claiming compactness of $\mathcal{C}_f(\rho)$. Note that it is very natural to expect this compactness because of the conditions (1)-(4) of the Theorem 7.1. Indeed, note that (1)-(4) are formulated in terms of how the marked points a_n are allowed to move under isotopies φ_u of id-type maps: the points inside of D_ρ have to stay inside, points outside of D_ρ move only slightly. At the same time the points a_n inside of D_ρ are allowed to rotate around points in \mathcal{O}_n only some bounded amount of times. Outside of D_ρ no “rotations” are allowed and basically nothing happens. The final remark is that the marked points are not allowed to come too close to each other.

Theorem 7.2 (Compactness). *For all $\rho \in \mathcal{P}'$ big enough $\mathcal{C}_f(\rho)$ is a compact subset of \mathcal{T}_f . Moreover, all its elements are asymptotically conformal.*

Proof. We are going to prove that the set $\mathcal{C}_f^\circ(\rho)$ from the proof of Theorem 7.1 is pre-compact, or equivalently, since \mathcal{T}_f is a metric space in the Teichmüller metric, that every sequence $\{[\varphi^n]\} \subset \mathcal{C}_f^\circ(\rho)$ has a subsequence that converges to some point $[\varphi] \in \mathcal{T}_f$.

Since each $[\varphi^n] \in \mathcal{C}_f^\circ(\rho)$, we can assume that every φ^n is of id-type and φ_u^n is an isotopy with $\varphi_0^n = \operatorname{id}$, $\varphi_1^n = \varphi^n$ satisfying conditions (1)-(4) of Theorem 7.1.

Recall from the proof of Theorem 7.1 that for all $\rho \in \mathcal{P}'$ big enough the first $N+1$ marked points move under isotopy φ_u only inside of D_ρ , while for every $n > N$ the point a_n moves inside of a disk D_n of radius $1/n$, and all these disks $D_\rho, D_{N+1}, D_{N+2}, \dots$ are mutually disjoint with the mutual distance between them bigger than one. Hence we can assume that when $u \in [0, 1/2]$, we have $\varphi_u^n|_{D_\rho} = \operatorname{id}$ for every n , and when $u \in [1/2, 1]$, we have $\varphi_u^n|_{\bigcup_{k=N+1}^\infty D_k} = \operatorname{id}$ for every n , that is, first the marked points inside of D_ρ do not move, and afterwards do not move the marked point outside of D_ρ . This means that we simply need to prove the theorem in two separate cases: when only the marked points outside of D_ρ move, and when only the marked points inside of D_ρ move.

In the former case the sequence $[\varphi^n]$ clearly has a limit point in \mathcal{T}_f , since marked points move inside of mutually disjoint small disks with mutual distance between them bounded from below.

Assume now that the marked points outside of D_ρ do not move. We are looking for a convergent subsequence in $\{\varphi^n\}$. Let $V := P_f \cap D_\rho$.

We say that $\varphi^k(V)$ and $\varphi^l(V)$ have the same configuration if for every pair of points $a_m, a_n \in V$ we have

$$\begin{aligned} \operatorname{Re} \varphi^k(a_m) < \operatorname{Re} \varphi^k(a_n) &\iff \operatorname{Re} \varphi^l(a_m) < \operatorname{Re} \varphi^l(a_n), \\ \operatorname{Re} \varphi^k(a_m) = \operatorname{Re} \varphi^k(a_n) &\iff \operatorname{Re} \varphi^l(a_m) = \operatorname{Re} \varphi^l(a_n), \\ \operatorname{Im} \varphi^k(a_m) < \operatorname{Im} \varphi^k(a_n) &\iff \operatorname{Im} \varphi^l(a_m) < \operatorname{Im} \varphi^l(a_n), \end{aligned}$$

$$\text{Im } \varphi^k(a_m) = \text{Im } \varphi^l(a_m) \iff \text{Im } \varphi^k(a_n) = \text{Im } \varphi^l(a_n).$$

Since V is finite, only finitely many different configurations are possible. After passing to a subsequence we may assume that all $\varphi^k(a_n)$ converge to some points b_n , $W_n^{\varphi^k} = \text{const}$ for all $a_n \in V$, and all $\varphi^k(V)$ have the same configuration.

Note that for k big enough $\varphi^k(a_n)$ will be compactly contained in disjoint small disks D_n contained in $\mathbb{D}_{\rho+1}(0)$ with centers at the limits b_n , so we can assume this holds for all n .

Now let $I_n^k : [0, 1] \rightarrow \mathbb{C}$ such that

$$I_n^k(u) = \varphi^k(a_n) + u(\varphi^{k+1}(a_n) - \varphi^k(a_n)),$$

and let ξ_u^k to be an isotopy of quasiconformal maps such that it is identity outside of $\cup D_n$, $\xi_0^k = \text{id}$ and $\xi_u^k(\varphi^k(a_n)) = I_n^k(u)$ for all $a_n \in V$.

Note that $[\varphi^{k+1}] = [\xi_1^k \circ \varphi^k]$. Indeed, while moving simultaneously along I_n^k , the sets $\xi_1^k \circ \psi^k(V)$ do not change their configuration. It is easy to see that in this case $W_n^{\xi_1^k \circ \varphi^k} = W_n^{\varphi^k}$. But then from Theorem 6.28 follows that $\xi_1^k \circ \varphi^k$ and φ^{k+1} define the same point in \mathcal{T}_f .

Hence $[\varphi^k] = [\xi_u^0 \cdot \xi_u^1 \cdot \dots \cdot \xi_u^k]$ converges to some $[\varphi]$ in the Teichmüller metric. This is an easy consequence of the fact that Teichmüller space of the unit disk with one marked point is homeomorphic to the unit disk.

The fact that the elements of $\mathcal{C}_f(\rho)$ are asymptotically conformal follows immediately from the construction of $\mathcal{C}_f(\rho)$. \square

8. PROOF OF CLASSIFICATION THEOREM

In this section we are going to finally prove the Classification Theorem 1.1. Before the proof we state a variation of the Banach Fixed Point Theorem that will be needed for this.

Theorem 8.1 (Adjusted Banach Fixed Point Theorem). *Let X be a compact complete metric space, and $\sigma : X \rightarrow X$ be a strictly contracting map. Then X contains a fixed point of σ and it is unique.*

Proof. The proof is essentially the same as for the classical Banach Fixed Point Theorem.

Assume that X does not contain a fixed point of σ and define a map $\kappa : X \rightarrow \mathbb{R}$ so that $\kappa(x) = d(\sigma(x), \sigma^2(x))/d(x, \sigma(x))$ where $d(x, y)$ is the metric function of X . Since X is compact, κ attains its maximum on X . Moreover, because σ is strictly contracting, this maximum is less than 1. But this means that for every $x \in X$ the sequence $x, \sigma(x), \sigma^2(x), \dots$ is Cauchy and due to completeness converges to some point in X . This point will be a fixed point of σ . Uniqueness is straightforward from the strict contraction of σ . \square

Now, we prove the main statement of this article.

Proof of Classification Theorem 1.1. We show existence and uniqueness of g separately.

(Existence.) Choose as some $f_0 \in \mathcal{N}$ so that its singular value does not escape. Then due to Theorem 2.9 in $I(f_0)$ there is a point z that escapes on rays with potential t and external address \underline{g} .

Let $f = c \circ f_0$ be the captured exponential function constructed as in Subsection 3.2 and having singular value z escaping as under f_0 .

From Theorems 7.1 and 7.2 we know that there is a non-empty compact invariant under σ subset $\mathcal{C}_f = \mathcal{C}_f(\rho) \subset \mathcal{T}_f$ such that all its elements are asymptotically

conformal. Next, from Theorem 4.4 follows that σ is strictly contracting in the Teichmüller metric on \mathcal{C}_f .

Thus we have a strictly contracting map σ on a compact complete metric space \mathcal{C}_f . Note that it is *not* true that σ is uniformly strictly contracting on \mathcal{C}_f (think of the map x^2 on the interval $[0, 1/2]$), so we cannot apply the Banach Fixed Point Theorem directly. But from Theorem 8.1 follows that \mathcal{C}_f contains a fixed point $[\varphi_0]$. Note (even though we do not use this fact later) that, because σ is strictly contracting on \mathcal{C}_f , the constructed fixed point $[\varphi_0]$ is the unique fixed point in \mathcal{C}_f .

Now let $g = \varphi_0 \circ f \circ \hat{\varphi}_0^{-1}$. This is an entire function that is Thurston equivalent to f . Moreover, since every point in \mathcal{C}_f is of id-type, without loss of generality we can assume that φ_0 is of id-type.

Now we need to prove that $\varphi_0(a_0)$ escapes on rays with potential t and external address \underline{s} .

Note first that since the map φ_0 is of id-type, the singular value $\varphi_0(a_0)$ escapes on rays. Indeed, the dynamic ray of g having external address \underline{s} is defined for potentials $t > t_{\underline{s}}$ (because none of its images contains the singular value), and the point p of this ray having potential $t > t_{\underline{s}}$ has the same asymptotics under iterations of g as $\varphi_0(a_0)$. Since g^{-1} is uniformly contracting on its tracts near ∞ , we see that for some integer $N > 0$ the point $g^N(p)$ coincides with $g^N(\varphi_0(a_0))$. For the same reason we can assume that for $n \geq N$ the legs $\varphi_0(R_n)$ coincide with the ray tails of g towards $g^n(\varphi_0(a_0))$ (just isotope φ_0 : every such $\varphi_0(R_n)$ is in the same homotopy class as the ray tail). Since the preimages of $\varphi_0(R_n)$ recover according to the external address, $\varphi_0(R_0)$ coincides with the ray tail of g towards $\varphi_0(a_0)$.

(Uniqueness.) Assume that there exists another function $h \in \mathcal{N}$ satisfying the same conditions on the combinatorics and speed of escape of the singular value. Let S_0^g and S_0^h be the standard spiders of g and h , respectively (i.e. union of the ray tails towards the singular orbit).

Since S_0^g and S_0^h have the same vertical order of legs, one can map P_g into P_h with a map χ of id-type belonging to an asymptotically conformal point of \mathcal{T}_g so that $\chi(S_0^g)$ is equivalent to S_0^h as spiders. But then we can define the map $\sigma : \mathcal{T}_g \rightarrow \mathcal{T}_g$ for the function g (i.e. with the capture identically equal to identity), and then the map $\hat{\chi}$ will be isotopic to χ relative P_g — this is an immediate consequence of the fact that χ and $\hat{\chi}$ are of id-type and that χ can be lifted by the branched covering maps g and h . On the other hand, since $[\text{id}]$ is invariant under σ (g is entire), and $[\chi]$ is asymptotically conformal, either the Teichmüller distance from $[\chi]$ to $[\text{id}]$ must be strictly bigger than the Teichmüller distance from $[\hat{\chi}]$ to $[\text{id}]$ (Theorem 4.4) or $[\chi] = [\text{id}]$ and $g = h$. Since $[\chi]$ is a fixed point of σ , only the latter option is possible. \square

9. ACKNOWLEDGEMENTS

We would like to express our gratitude to our research team in Aix-Marseille Université including Dierk Schleicher, Kostiantyn Drach, Mikhail Hlushchanka, Bernhard Reinke, Sergey Shemyakov and Roman Chernov for uncountably many enjoyable and enlightening discussions of this project at different stages. We also want to thank Dzmitry Dudko for his multiple suggestions that helped to advance the project, as well as Lasse Rempe for his long list of comments and relevant questions. Finally, we are grateful to DFG and ERC fundings who have financed this project.

REFERENCES

- [A] Lars Ahlfors, *Lectures on Quasiconformal Mappings*. American Mathematical Society (2006)

- [B] Konstantin Bogdanov, *Infinite-dimensional Thurston theory and transcendental dynamics with escaping singular orbits*. PhD Thesis, Aix-Marseille Université, 2020.
- [BA] Arne Beurling and Lars Ahlfors, *The boundary correspondence under quasiconformal mappings*. Acta Mathematica **96** (1956), 125–142.
- [Br] David Brown, *Thurston equivalence without postcritical finiteness for a family of polynomial and exponential mappings*. PhD Thesis, Cornell University.
- [CT] Guizhen Cui and Lei Tan, *A Characterization of Hyperbolic Rational Maps*. Inventiones Mathematicae **183.3** (2011), 451–516.
- [DH] Adrien Douady and John Hubbard, *A proof of Thurston's topological characterization of rational functions*. Acta Mathematica **171** (1993), 263–297.
- [DK] Robert Devaney and Michal Krych, *Dynamics of $\exp(z)$* . Ergod. Theor. Dyn. Syst. **4** (1) (1984), 35–52.
- [dS] Bart de Smit, *The fundamental group of the Hawaiian earring is not free*. International Journal of Algebra and Computation **2** (1) (1992), 33–37.
- [EL] Alexandre Eremenko and Mikhail Lyubich, *Dynamical properties of some classes of entire functions*. Annales de l'institut Fourier, **42** (1992) no. 4, 989–1020
- [F] Markus Förster, *Exponential maps with escaping singular orbits*. PhD Thesis, International University Bremen, 2006.
- [FRS] Markus Förster, Lasse Rempe, and Dierk Schleicher, *Classification of escaping exponential maps*. Proc Amer Mathe Soc. **136** (2008), 651–663.
- [FS] Markus Förster and Dierk Schleicher, *Parameter rays in the space of exponential maps*. Ergod Theory Dynam Systems **29** (2009), 515–544.
- [GL] Frederick P. Gardiner and Nikola Lakic, *Quasiconformal Teichmüller theory*. American Mathematical Society (2000).
- [H1] John Hubbard, *Teichmüller theory and applications to geometry, topology, and dynamics*. Volume 1: Teichmüller theory. Ithaca, NY: Matrix Editions (2006).
- [H2] John Hubbard, *Teichmüller theory and applications to geometry, topology, and dynamics*, Volume 2: Surface homeomorphisms and rational functions. Ithaca, NY: Matrix Editions (2016).
- [HS] John Hubbard and Dierk Schleicher, *The spider algorithm*. In: R. Devaney (ed.), *Complex dynamics: the mathematics behind the Mandelbrot and Julia sets*. Proc Symp Pure Math **49**, Amer Math Soc (1994), 155–180.
- [HSS] John Hubbard, Dierk Schleicher, and Mitsuhiro Shishikura, *Exponential Thurston maps and limits of quadratic differentials*. Journal Amer Math Soc **22** (2009), 77–117.
- [LV] Olli Lehto and K.I. Virtanen, *Quasiconformal mappings in the plane*. Springer-Verlag Berlin Heidelberg (1973).
- [RRRS] Günter Rottenfuß, Johannes Rückert, Lasse Rempe, and Dierk Schleicher, *Dynamic rays of bounded-type entire functions*. Annals of Mathematics **173** (2011), 77–125.
- [S] Dierk Schleicher, *Dynamics of entire functions*. In: G. Gentili et al (eds), *Holomorphic dynamical systems*. LNS **1998** (2010), 295–339.
- [SZ] Johannes Zimmer and Dierk Schleicher, *Escaping points for exponential maps*. J. Lond. Math. Soc. (2) **67** (2003), 380–400.

I2M, Aix-Marseille Université, France
Email: konstantin.bogdanov@univ-amu.fr