

SMOOTH NUMBERS IN BEATTY SEQUENCES

ROGER BAKER

ABSTRACT. Let θ be an irrational number of finite type and let $\psi \geq 0$. We consider numbers in the Beatty sequence of integer parts,

$$\mathcal{B}(x) = \{\lfloor \theta n + \psi \rfloor : 1 \leq n \leq x\}.$$

Let $C > 3$. Writing $P(n)$ for the largest prime factor of n and $|\dots|$ for cardinality, we show that

$$|\{n \in \mathcal{B}(x) : P(n) \leq y\}| = \frac{1}{\theta} \Psi(\theta x, y) (1 + o(1))$$

as $x \rightarrow \infty$, uniformly for $y \geq (\log x)^C$. Here $\Psi(X, y)$ denotes the number of integers up to X with $P(n) \leq y$. The range of y extends that given by Akbal [1]. The work of Harper [8] plays a key role in the proof.

1. INTRODUCTION

A positive integer n is said to be y -smooth if $P(n)$, the largest prime factor of n , is at most y . We write $\mathfrak{S}(y)$ for the set of y -smooth numbers in \mathbb{N} and

$$\Psi(x, y) = |\mathfrak{S}(y) \cap [1, x]|,$$

where $|\dots|$ denotes cardinality.

Let $\theta > 1$ be an irrational number and $\psi \in [0, \infty)$. Arithmetic properties of the Beatty sequence

$$\mathcal{B}(x) = \{\lfloor \theta n + \psi \rfloor : 1 \leq n \leq x\}$$

(where $\lfloor \cdot \rfloor$ denotes integer part), have been studied in [1, 3, 4, 7], for example. One may conjecture that for x large and $y = y(x)$ not too small, say $x \geq y \geq (\log x)^C$ where $C > 1$, we have

$$(1.1) \quad |\{n \in \mathcal{B}(x) : P(n) \leq y\}| = \frac{1}{\theta} \Psi(\theta x, y) (1 + o(1))$$

2020 *Mathematics Subject Classification*. Primary 11N25; secondary 11L03.

Key words and phrases. Beatty sequence, exponential sums over smooth numbers.

where $o(1)$ denotes a quantity tending to 0 as x tends to infinity. Banks and Shparlinski [4] obtained (1.1) (in a slightly different form) uniformly for

$$\exp((\log x)^{2/3+\varepsilon}) \leq y \leq x.$$

(We write ε for an arbitrary positive number.) Under the additional condition that θ is of finite type, Akbal [1] obtained (1.1) uniformly for

$$(1.2) \quad \exp((\log \log x)^{5/3+\varepsilon}) \leq y \leq x.$$

In the present paper we extend the range (1.2).

Theorem 1. *Let $\theta > 1$ be an irrational number of finite type and $\psi \geq 0$. Then (1.1) holds uniformly for*

$$(\log x)^{3+\varepsilon} \leq y \leq x.$$

We recall that an irrational number θ is said to be of finite type if ($\|\dots\|$ denoting distance to the nearest integer) we have

$$\|m\theta\| \geq \frac{c}{m^\kappa} \quad (m \in \mathbb{N})$$

for some $c > 0$ and $\kappa > 0$. We note that if θ is of finite type, then so is θ^{-1} .

Theorem 1 depends on an estimate for the exponential sum

$$S(\theta) = \sum_{\substack{n \leq x \\ n \in \mathfrak{S}(y)}} e(n\theta).$$

Akbal [1] uses the estimate of Foury and Tenenbaum [6]: for $3 \leq y \leq \sqrt{x}$, $q \in \mathbb{N}$, $(a, q) = 1$, and $\delta \in \mathbb{R}$,

$$S\left(\frac{a}{q} + \delta\right) \ll x(1 + |\delta x|) \log^3 x \left(\frac{y^{1/2}}{x^{1/4}} + \frac{1}{q^{1/2}} + \left(\frac{qy}{x}\right)^{1/2} \right).$$

This is unhelpful when, say, $y = (\log x)^C$ ($C > 1$) since the trivial bound for $S(\theta)$ is $x^{1-1/C+o(1)}$ (see e.g. [9]).

We use a procedure of Harper [8, Theorem 1] and to state his result we introduce some notation. For $2 \leq y \leq x$, let $u = \frac{\log y}{\log x}$ and let $\alpha = \alpha(x, y)$ be the solution of

$$\sum_{p \leq y} \frac{\log p}{p^\alpha - 1} = \log x.$$

For convenience, when $\theta = \frac{a}{q} + \delta$ as above, we write

$$L = 2(1 + |\delta x|)$$

and

$$(1.3) \quad M = u^{3/2} \log u \log x (\log L)^{1/2} (\log qy)^{1/2}.$$

In Theorem 1 of [8] it is shown that whenever

$$(1.4) \quad q^2 L^2 y^3 \leq x,$$

we have

$$(1.5) \quad S\left(\frac{a}{q} + \delta\right) \ll \Psi(x, y) (q(1 + |\delta x|))^{-\frac{1}{2} + \frac{3}{2}(1-\alpha(x, y))} M.$$

We cannot use this bound directly since (1.3) is too restrictive. We adapt Harper's argument to obtain

Theorem 2. *Let f be a completely multiplicative function, $|f(n)| \leq 1$ ($n \in \mathbb{N}$). Let*

$$S(f, \theta) = \sum_{\substack{n \leq x \\ n \in \mathfrak{S}(y)}} f(n) e(\theta n).$$

Let $q \in \mathbb{N}$, $(a, q) = 1$, and $\delta \in \mathbb{R}$. Then, with $\alpha = \alpha(x, y)$, we have

$$S\left(f, \frac{a}{q} + \delta\right) \ll \Psi(x, y) \left\{ (q(1 + |\delta x|))^{-\frac{1}{2} + \frac{3}{2}(1-\alpha)} M + x^{\alpha/2} (qLy^3)^{\frac{1}{2}} \sqrt{\log y \log q} \right\}.$$

To save space, we refer frequently to [8] in our proof of Theorem 2 in Section 2. Theorem 1 is deduced in a straightforward manner from Theorem 2 in Section 3.

The factor $f(n)$ in Theorem 2 is not needed elsewhere in the paper, but it requires no significant effort to include it.

I would like to thank Adam Harper for helpful comments concerning his proof of (1.5).

2. PROOF OF THEOREM 2.

Lemma 1. *Let $2 \leq y \leq x$ and $d \geq 1$. Then we have*

$$\Psi\left(\frac{x}{d}, y\right) \ll \frac{1}{d^{\alpha(x, y)}} \Psi(x, y).$$

Proof. See de la Bretèche and Tenenbaum [5, Théorème 2.4 (i)] □

We write $p(n)$ for the smallest prime factor of $n \in \mathbb{N}$. We begin the proof of Theorem 2 by noting that the result is trivial for $qLy^3 \geq x^\alpha$. Suppose now that $qLy^3 < x^\alpha$. Every y -smooth number in $[qLy^2, x]$ can be written uniquely in the form mn , where

$$qLy < m \leq qLy^2, \quad \frac{m}{P(m)} \leq qLy, \quad m \in \mathfrak{S}(y)$$

and

$$\frac{qLy^2}{m} \leq n \leq \frac{x}{m}, \quad p(n) \geq P(m), \quad n \in \mathfrak{S}(y).$$

(We take m to consist of the product of the smallest prime factors of the number.) With $\theta = \frac{\alpha}{q} + \delta$, we have

$$\begin{aligned} (2.1) \quad S(f, \theta) &= \sum_{\substack{qLy^2 \leq n \leq x \\ n \in \mathfrak{S}(y)}} f(n)e(n\theta) + O(\Psi(qLy^2, y)) \\ &= U + O(\Psi(qLy^2, y)) \end{aligned}$$

where

$$U = \sum_{\substack{qLy < m \leq qLy^2 \\ m/P(m) \leq qLy \\ m \in \mathfrak{S}(y)}} \sum_{\substack{\frac{qLy^2}{m} \leq n \leq \frac{x}{m} \\ p(n) \geq P(m) \\ n \in \mathfrak{S}(y)}} f(mn)e(mn\theta).$$

We now decompose U as

$$(2.2) \quad U = \sum_{0 \leq j \leq \frac{\log y}{\log 2}} U_j$$

where

$$U_j = \sum_{p \leq y} \sum_{\substack{2^j qLy < m \leq qLy \min(2^{j+1}, p) \\ P(m)=p}} f(m) \sum_{\substack{\frac{qLy^2}{m} \leq n \leq \frac{x}{m} \\ p(n) \geq p \\ n \in \mathfrak{S}(y)}} f(n)e(mn\theta),$$

noting that if $P(m) = p \leq y$, then m is y -smooth, and the condition $\frac{m}{P(m)} \leq qLy$ can be written as $m \leq qLyp$.

We apply the Cauchy-Schwarz inequality to U_j . Let \sum_m denote

$$\sum_{\substack{2^j qLy < m \leq qLy \\ P(m)=p}} \min(2^{j+1}, p).$$

We obtain

$$\begin{aligned} U_j &\leq \sqrt{\sum_{p \leq y} \sum_m 1} \sqrt{\left| \sum_{2^j \leq p \leq y} \sum_{\substack{2^j qLy < m' \leq \frac{qLy}{p} \\ \min(2^{j+1}, p)}} \sum_{\substack{\frac{qLy^2}{m'p} \leq n \leq \frac{x}{m'p}, p(n) \geq p, n \in \mathfrak{S}(y)}} f(n) e(m'pn\theta) \right|^2} \\ &\ll \sqrt{\Psi(2^{j+1}qLy, y)} \sqrt{\sum_{2^j \leq p \leq y} \sum_{\substack{n_1, n_2 \leq \frac{x}{2^j qLy} \\ p(n_1), p(n_2) \geq p \\ n_1, n_2 \in \mathfrak{S}(y)}} \min \left\{ \frac{2^{j+1}qLy}{p}, \frac{1}{\|(n_1 - n_2)p\theta\|} \right\}} \end{aligned}$$

For the last step, we open the square and sum the geometric progression over m' . We may restrict the sum over primes to $p \geq 2^j$, since otherwise the sum over m is empty. Our final bound here for U_j exactly matches [8].

Let

$$\mathfrak{I}_j(r) := \max_{1 \leq b \leq r-1} \sum_{\substack{n_1, n_2 \leq \frac{x}{2^j qLy} \\ n_1, n_2 \in \mathfrak{S}(y) \\ n_1 - n_2 \equiv b \pmod{r}}} 1.$$

Just as in [8], after distinguishing the cases $p \mid q$ and $p \nmid q$, we arrive at

$$(2.3) \quad U_j \ll \sqrt{\Psi(2^{j+1}qLy, y)} (\sqrt{S_1} + \sqrt{S_2}),$$

with

$$\begin{aligned} S_1 = S_1(j) &:= \sum_{\substack{2^j \leq p \leq y \\ p \nmid q}} \mathfrak{I}_j(q) \sum_{b=1}^{q-1} \min \left\{ \frac{2^{j+1}qLy}{p}, \frac{q}{p} \right\} \\ &\quad + \sum_{\substack{2^j \leq p \leq y \\ p \mid q}} \mathfrak{I}_j\left(\frac{q}{p}\right) \sum_{b=1}^{(q/p)-1} \min \left\{ \frac{2^{j+1}qLy}{p}, \frac{q}{pb} \right\} \end{aligned}$$

and

$$S_2 = S_2(j) := \sum_{\substack{2^j \leq p \leq y \\ p|q}} \frac{1}{p} \sum_{\substack{n_1, n_2 \leq \frac{x}{2^j qLy} \\ n_1 - n_2 \equiv 0 \pmod{q} \\ n_1, n_2 \in \mathfrak{S}(y)}} \min \left\{ 2^{j+1} qLy, \frac{1}{|(n_1 - n_2)\delta|} \right\}$$

$$\sum_{\substack{2^j \leq p \leq y \\ p|q}} \frac{1}{p} \sum_{\substack{n_1, n_2 \leq \frac{x}{2^j qLy} \\ n_1 - n_2 \equiv 0 \pmod{q/p} \\ n_1, n_2 \in \mathfrak{S}(y)}} \min \left\{ 2^{j+1} qLy, \frac{1}{|(n_1 - n_2)\delta|} \right\}.$$

We now depart to an extent from the argument in [8]; we have extra terms in the upper bounds in Lemma 2 below which arise because we do not have the upper bound (1.4).

Lemma 2. *Let $(\log x)^{1.1} \leq y \leq x^{1/3}$, $q \geq 1$, and $L = 2(1 + |\delta x|)$. Then for any j , $0 \leq j \leq \frac{\log y}{\log 2}$, and any prime $p \mid q$, we have*

$$(2.4) \quad \mathfrak{T}_j(q) \ll \frac{\Psi(x/2^j qLy, y)^2}{q} q^{1-\alpha(x,y)} \log x + y \Psi \left(\frac{x}{2^j qLy}, y \right)$$

and

$$(2.5) \quad \mathfrak{T}_j \left(\frac{q}{p} \right) \ll \frac{\Psi(x/2^j qLy, y)^2}{q/p} \left(\frac{q}{p} \right)^{1-\alpha(x,y)} \log x$$

$$+ y \Psi \left(\frac{x}{2^j qLy}, y \right).$$

Under the same hypotheses, we have

$$(2.6) \quad \sum_{\substack{n_1, n_2 \leq \frac{x}{2^j qLy} \\ n_1 - n_2 \equiv 0 \pmod{q} \\ n_1, n_2 \in \mathfrak{S}(y)}} \min \left\{ 2^{j+1} qLy, \frac{1}{|(n_1 - n_2)\delta|} \right\}$$

$$\ll 2^j y \Psi \left(\frac{x}{2^j qLy}, y \right)^2 (qL)^{1-\alpha(x,y)} \log x \log L$$

$$+ 2^j qLy^2 \Psi \left(\frac{x}{2^j qLy}, y \right)$$

and, for any prime $p \mid q$,

$$\begin{aligned}
(2.7) \quad & \sum_{\substack{n_1, n_2 \leq \frac{x}{2^j q L y} \\ n_1 - n_2 \equiv 0 \pmod{q/p} \\ n_1, n_2 \in \mathfrak{S}(y)}} \min \left\{ 2^{j+1} q L y, \frac{1}{|(n_1 - n_2) \delta|} \right\} \\
& \ll p 2^j y \Psi \left(\frac{x}{2^j q L y}, y \right)^2 \left(\frac{q L}{p} \right)^{1-\alpha(x,y)} \log x \log L \\
& \quad + 2^j q L y^2 \Psi \left(\frac{x}{2^j q L y}, y \right).
\end{aligned}$$

Proof. We have

$$\mathfrak{T}_j(q) = \max_{1 \leq b \leq q-1} \sum_{\substack{n_1 \leq x/2^j q L y \\ n_1 \in \mathfrak{S}(y)}} \sum_{\substack{n_2 \leq x/2^j q L y \\ n_2 \equiv n_1 - b \pmod{q} \\ n_2 \in \mathfrak{S}(y)}} 1,$$

and just as in [8] the inner sum is

$$\ll y + \frac{\Psi(x/2^j q L y)}{q} q^{1-\alpha(x,y)} \log x.$$

This leads to the bound (2.4) on summing over n_1 . The bound (2.5) follows in exactly the same way.

To prove (2.6), (2.7) we distinguish two cases. If $|\delta| \leq 1/x$, then $L = 2(1 + |\delta x|) \asymp 1$, and the bounds can be proved exactly as above on bounding $\min \left\{ 2^{j+1} q L y, \frac{1}{|(n_1 - n_2) \delta|} \right\}$ by $2^{j+1} q L y \ll 2^j q y$.

Now suppose $|\delta| > 1/x$, so that $L \asymp |\delta x|$. We partition the sum in (2.6) dyadically. Let us use \sum^\dagger to denote a sum over pairs of integers $n_1, n_2 \leq \frac{x}{2^j q L y}$ that are y -smooth and satisfy $n_1 - n_2 \equiv 0 \pmod{q}$. Then we have

$$\begin{aligned}
& \sum_{|n_1 - n_2| \leq \frac{x}{2^j q L^2 y}}^\dagger \min \left\{ 2^{j+1} q L y, \frac{1}{|(n_1 - n_2) \delta|} \right\} \\
& \ll 2^j q L y \sum_{\substack{n_1 \leq \frac{x}{2^j q L y} \\ n_1 \in \mathfrak{S}(y)}} \sum_{\substack{|n_2 - n_1| \leq \frac{x}{2^j q L^2 y} \\ n_2 \in \mathfrak{S}(y) \\ n_2 \equiv n_1 \pmod{q}}} 1.
\end{aligned}$$

Following [8], but as above *not* absorbing a term y , the last expression is

$$\begin{aligned} &\ll 2^j qLy \sum_{\substack{n_1 \leq \frac{x}{2^j qLy} \\ n_1 \in \mathfrak{S}(y)}} \left\{ \frac{\Psi(x/2^j qLy, y)}{qL} (qL)^{1-\alpha} \log x + y \right\} \\ &\ll 2^j y \Psi(x/2^j qLy, y)^2 (qL)^{1-\alpha} \log x + 2^j qLy^2 \Psi(x/2^j qLy, y). \end{aligned}$$

Similarly, for any r , $0 \leq r \leq (\log L)/\log 2$, we have

$$\begin{aligned} &\sum_{\substack{\dagger \\ \frac{2^r}{2^j qL^2 y} < |n_1 - n_2| \leq \frac{2^{r+1}x}{2^j qL^2 y}}} \min \left\{ 2^{j+1} qLy, \frac{1}{|(n_1 - n_2)\delta|} \right\} \\ &\ll \frac{2^j qLy}{2^r} \sum_{\substack{n_1 \leq \frac{x}{2^j qLy} \\ n_1 \in \mathfrak{S}(y)}} \sum_{\substack{|n_2 - n_1| \leq \frac{2^{r+1}x}{2^j qL^2 y} \\ n_2 \in \mathfrak{S}(y) \\ n_2 \equiv n_1 \pmod{q}}} 1 \\ &\ll \frac{2^j qLy}{2^r} \sum_{\substack{n_1 \leq \frac{x}{2^j qLy} \\ n_1 \in \mathfrak{S}(y)}} \left\{ \Psi \left(\frac{x}{2^j qLy}, y \right) \left(\frac{2^r}{qL} \right)^\alpha \log x + y \right\} \\ &\ll 2^j y \Psi \left(\frac{x}{2^j qLy}, y \right)^2 \left(\frac{2^r}{qL} \right)^\alpha \log x + \frac{2^j qLy^2}{2^r} \Psi \left(\frac{x}{2^j qLy}, y \right). \end{aligned}$$

The bound (2.6) follows on summing over r .

The bound (2.7) follows in exactly the same way; we lose a factor p^α in the first term on the right-hand side in (2.7) because of the weaker congruence condition. This completes the proof of Lemma 2. \square

We now assemble our bounds to prove Theorem 1. As noted in [8], for any $p \leq y$ we have

$$(2.8) \quad \sum_{b=1}^{q-1} \min \left\{ \frac{2^{j+1} qLy}{p}, \frac{q}{b} \right\} \ll q \log q$$

and, for $p \mid q$,

$$(2.9) \quad \sum_{b=1}^{(q/p)-1} \min \left\{ \frac{2^{j+1} qLy}{p}, \frac{q}{pb} \right\} \ll \frac{q}{p} \log q.$$

Let

$$A_j = \left(\sum_{\substack{2^j \leq p \leq y \\ p \nmid q}} \frac{1}{p} + \sum_{\substack{2^j \leq p \leq y \\ p \mid q}} 1 \right) 2^j y \Psi \left(\frac{x}{2^j q L y}, y \right) (qL)^{1-\alpha} \log x \log L.$$

We deduce from (2.1)–(2.3), Lemma 2, and (2.8)–(2.9) that

$$(2.10) \quad S \left(f, \frac{a}{q} + \delta \right) \ll \Psi(qLy^2, y) + A + B_1 + B_2$$

where

$$\begin{aligned} A &= \sum_{0 \leq j \leq \frac{\log y}{\log 2}} \sqrt{\Psi(2^{j+1}qLy, y)} \sqrt{\frac{y}{\log y} \log q \Psi \left(\frac{x}{2^j q Ly}, y \right)^2 q^{1-\alpha} \log x} \\ &\quad + \sum_{0 \leq j \leq \frac{\log y}{\log 2}} \sqrt{\Psi(2^{j+1}qLy, y)} \sqrt{A_j}, \end{aligned}$$

$$B_1 = \sum_{0 \leq j \leq \frac{\log y}{\log 2}} \sqrt{\Psi(2^{j+1}qLy, y)} \sqrt{\sum_{2^j \leq p \leq y} q \log q \cdot y \Psi \left(\frac{x}{2^j q Ly}, y \right)},$$

and

$$B_2 = \sum_{0 \leq j \leq \frac{\log y}{\log 2}} \sqrt{\Psi(2^{j+1}qLy, y)} \sqrt{\frac{qLy^3}{\log y} \Psi \left(\frac{x}{2^j q y}, y \right)}.$$

For the estimation of A we can appeal to [8]:

$$(2.11) \quad A \ll \Psi(x, y) (qL)^{-\frac{1}{2} + \frac{3}{2}(1-\alpha(x, y))} M.$$

The Ψ functions in B_1 and B_2 are estimated using Lemma 1. Thus

$$\begin{aligned} \sqrt{\Psi \left(\frac{x}{2^j q Ly}, y \right) \Psi(2^{j+1}qLy, y)} &\ll \Psi(x, y) (2^j q Ly)^{-\alpha/2} \left(\frac{x}{2^{j+1} q Ly} \right)^{-\alpha/2} \\ &\ll \Psi(x, y) x^{-\alpha/2}, \end{aligned}$$

leading to (a slightly wasteful) bound

$$(2.12) \quad B_1 + B_2 \ll \Psi(x, y) \sqrt{y^3 q L \log q \log y} x^{-\alpha/2}.$$

The remaining term to be estimated is $\Psi(qLy^2, y)$, and Lemma 1 gives

$$(2.13) \quad \Psi(qLy^2, y) \ll \left(\frac{x}{qLy^2}\right)^{-\alpha} \Psi(x, y).$$

This term can be absorbed into the right-hand side of (2.12) since

$$(2.14) \quad (qLy^3)^{\alpha-\frac{1}{2}} \ll (qLy^3)^{\alpha/2} \ll x^{\alpha/2}.$$

Theorem 2 follows on combining (2.10)–(2.14).

3. PROOF OF THEOREM 2

Lemma 3. *Let $u_1, \dots, u_N \in \mathbb{R}$. Then for any $J \in \mathbb{N}$ and any $\rho \leq \sigma \leq \rho + 1$, we have*

$$\begin{aligned} & \left| |\{1 \leq n \leq N : u_n \in [\rho, \sigma] \bmod 1\}| - (\sigma - \rho)N \right| \\ & \leq \frac{N}{J+1} + 3 \sum_{j=1}^J \frac{1}{j} \left| \sum_{n=1}^N e(ju_n) \right|. \end{aligned}$$

Proof. See e.g. [2], Theorem 2.1. □

Proof of Theorem 1. In view of the result of [1] cited above, we may suppose that

$$(3.1) \quad (\log x)^{3+\varepsilon} \leq y \leq \exp((\log \log x)^2).$$

We note that $\lfloor \theta n + \psi \rfloor = m$ for some m if and only if

$$(3.2) \quad 0 < \left\{ \frac{m+1-\psi}{\theta} \right\} \leq \frac{1}{\theta},$$

so that, applying Lemma 3,

$$\begin{aligned} \sum_{n \in \mathcal{B}(x, y)} 1 &= \sum_{\substack{m \leq \theta x \\ m \in \mathfrak{S}(y) \\ (3.2) \text{ holds}}} 1 + O(1) \\ &= \theta^{-1} \Psi(\theta x, y) + O\left(\frac{\Psi(\theta x, y)}{\log x} + \sum_{j=1}^{\lfloor \log x \rfloor} \frac{1}{j} \left| \sum_{\substack{m \leq \theta x \\ m \in \mathfrak{S}(y)}} e\left(\frac{jm}{\theta}\right) \right| \right) \end{aligned}$$

For our purposes, then, it suffices to show that

$$(3.3) \quad \sum_{\substack{m \leq \theta x \\ m \in \mathfrak{S}(y)}} e(j\theta^{-1}m) \ll \Psi(\theta x, y)(\log x)^{-2}$$

uniformly for $1 \leq j \leq \log x$. We deduce this from Theorem 2.

By Dirichlet's theorem there is $q \in \mathbb{N}$, $1 \leq q \leq x^{1/2}$, and $a \in \mathbb{N}$ with $(a, q) = 1$ such that

$$\delta := j\theta^{-1} - \frac{a}{q} \in \left[-\frac{1}{qx^{1/2}}, \frac{1}{qx^{1/2}} \right].$$

Now

$$|qj\theta^{-1} - a| \geq \frac{c_1}{(qj)^\kappa}$$

for positive constants c_1 and κ , hence

$$x^{-1/2} \geq q|\delta| \geq \frac{c_1}{(qj)^\kappa}$$

giving

$$q \gg x^{1/2\kappa}(\log x)^{-1}.$$

We apply Theorem 2 with $j\theta^{-1}$ in place of θ , and θx in place of x . Note that

$$qL = 2q(1 + |\delta\theta x|) \ll x^{1/2}$$

so that, in view of (3.1),

$$(3.4) \quad qLy^3 \ll x^{5/9}.$$

Now, since $y \geq (\log x)^{3+\varepsilon}$, we have

$$\alpha(\theta x, y) \geq \frac{2}{3} + \eta \quad (\eta = \eta(\varepsilon) > 0);$$

see [9], for example. Thus, abbreviating $\alpha(\theta x, y)$ to α ,

$$q^{-\frac{1}{2} + \frac{3}{2}(1-\alpha)} \ll q^{-\eta} \ll x^{-\eta/3\kappa}.$$

Now, with M as in (1.3) with θx in place of x ,

$$(3.5) \quad q^{-\frac{1}{2} + \frac{3}{2}(1-\alpha)} M \ll x^{-\eta/4\kappa}.$$

Next, recalling (3.4),

$$(3.6) \quad (\theta x)^{-\alpha/2} (qLy^3)^{1/2} \sqrt{\log y \log q} \ll x^{-1/20}.$$

Combining (3.5), (3.6) we obtain (3.3). This completes the proof of Theorem 1. \square

REFERENCES

- [1] Y. Akbal, A note on values of Beatty sequences that are free of large prime factors, *Colloq. Math.* **160** (2020), 53–63.
- [2] R.C. Baker, *Diophantine Inequalities*, Clarendon Press, Oxford, 1986.
- [3] R.C. Baker and L. Zhao, Gaps between primes in Beatty sequences, *Acta Arith.* **172** (2016), 207–242.
- [4] W.D. Banks and I.E. Shparlinski, Prime divisors in Beatty sequences, *J. Number Theory* **123** (2007), 413–425.
- [5] R. de la Brétèche and G. Tenenbaum, Propriétés statistiques des entiers friables, *Ramanujan J.* **9** (2005), 139–202.
- [6] E. Fouvry and G. Tenenbaum, Entiers sans grand facteur premier en progressions arithmétiques, *Proc. London Math. Soc.* **63** (1991), 449–494.
- [7] G. Harman, Primes in Beatty sequences in short intervals, *Mathematika* **62** (2016), 572–586.
- [8] A. Harper, Minor arcs, mean values and restriction theory for exponential sums over smooth numbers, *Compos. Math.* **152** (2016), 1121–1158.
- [9] A. Hildebrand and G. Tenenbaum, Integers without large prime factors, *J. Théor. Nombres Bordeaux* **5** (1993), 411–484.

DEPARTMENT OF MATHEMATICS
BRIGHAM YOUNG UNIVERSITY
PROVO, UT 84602, U.S.A
Email address: baker@math.byu.edu