

# Metropolis–Hastings transition kernel couplings

John O’Leary\* and Guanyang Wang†

January 5, 2022

## Abstract

Couplings play a central role in the analysis of Markov chain convergence and in the construction of new Markov chain Monte Carlo estimators, diagnostics, and variance reduction techniques. Tight bounds and efficient methods require an appropriate choice of coupling, which can be hard to find when the set of possible couplings is intractable. To address this challenge for the Metropolis–Hastings (MH) family of algorithms, we establish a simple characterization of the set of all MH transition kernel couplings. We then extend this result to describe the set of maximal couplings of the MH kernel, resolving an open question of O’Leary et al. [2021]. We conclude with a series of examples to build intuition. Our results represent an advance in understanding the MH kernel and a step forward for coupling this popular class of algorithms.

## 1 Introduction

From early results on Markov chain ergodicity [Doebelin, 1938, Harris, 1955] to the present day, couplings have played an important role in the analysis of convergence to stationarity. In recent years couplings have also gained prominence as a basis for sampling [Propp and Wilson, 1996, Fill, 1997, Neal, 1999, Flegal and Herbei, 2012], convergence diagnosis [Johnson, 1996, 1998, Biswas et al., 2019], variance reduction [Neal and Pinto, 2001, Goodman and Lin, 2009, Piponi et al., 2020], and unbiased estimation [Glynn and Rhee, 2014, Jacob et al., 2020, Heng and Jacob, 2019, Middleton et al., 2019, 2020]. Across these applications, couplings that deliver smaller meeting times are associated with better results.

Given a pair of Markov chains, the distribution of meeting times that a coupling can produce is constrained by the total variation distance between the marginal distributions of the chains at each iteration [Aldous, 1983, Lindvall, 1992]. Couplings that achieve the fastest meeting times allowed by this bound are said to be efficient or maximal and exist in some generality [Griffeath, 1975, Pitman, 1976, Goldstein, 1979]. Unfortunately these couplings are rarely Markovian, while this is a requirement in many of the applications above. Efficient Markovian couplings are known to exist – or not to exist – in special cases, but they remain elusive for discrete-time MCMC algorithms [Burdzy and Kendall, 2000, Connor and Jacka, 2008, Kuwada, 2009, Hsu and Sturm, 2013, Kendall, 2015, Böttcher, 2017, Banerjee and Kendall, 2017]. For techniques like the Metropolis–Hastings (MH) algorithm [Metropolis

---

\*Department of Statistics, Harvard University, Cambridge, USA. Email: joleary@g.harvard.edu

†Department of Statistics, Rutgers University, New Brunswick, USA. Email: guanyang.wang@rutgers.edu

et al., 1953, Hastings, 1970], it is difficult to even characterize the set of possible Markovian couplings, much less to differentiate among them by their implications for meeting times.

In this paper, we consider transition kernel couplings for the MH family of algorithms. These couplings depend only on the current state of each chain and form a natural starting point for inquiry into more general Markovian couplings. We fully characterize the set of kernel couplings for such chains and prove that every such coupling can be expressed as a coupled proposal step followed by a coupled acceptance step. This two-step representation is easy to implement computationally and amenable to theoretical analysis, following e.g. Bou-Rabee et al. [2020]. It should clarify the task of coupling design for applied work and support the search for more efficient couplings for this important class of algorithms.

So far, two approaches to coupling the MH kernel appear in the literature. In one method a coupling is defined directly in terms of the probability density or mass function of the marginal transition kernel [Rosenthal, 1996, 2002, Qin and Hobert, 2021, O’Leary et al., 2021]. This approach recalls the  $\gamma$ -coupling of Lindvall [1992, chap 1.5] and relies on few properties of the underlying transition kernel. The other method is to define one coupling at the MH proposal step followed by a second coupling at the acceptance step, subject to certain conditions [Johnson, 1998, Bou-Rabee et al., 2020, Jacob et al., 2020]. It has been unclear if one strategy is preferable to the other, or if there exist kernel couplings beyond the scope of either approach. In this paper we resolve this question, showing that every MH transition kernel coupling has a two-step representation.

With these results in hand, we turn to the maximal couplings of the MH transition kernel, which are sometimes called greedy couplings [Aldous and Fill, 1995, Hayes and Sinclair, 2005], one-step maximal couplings [Reutter and Johnson, 1995, Hayes and Vigoda, 2003, Kartashov and Golomozyi, 2013], or step-by-step maximal couplings [Pillai and Smith, 2019] of the associated Markov chains. These couplings do not usually achieve the fastest meeting times allowed by the coupling inequality, but they provide a valuable reference point for further analysis. Extending the results described above, we fully characterize such couplings in terms of the properties of the associated proposal and acceptance couplings. We also resolve an open question of O’Leary et al. [2021] on the structure of maximal transition kernel couplings and their relationship with maximal couplings of the proposal distributions.

We note that all of our results hold for simple MH algorithms as well as refinements such as Hamiltonian Monte Carlo [Duane et al., 1987, Neal, 1993, 2011], the Metropolis-adjusted Langevin algorithm [Roberts and Tweedie, 1996], and particle MCMC [Andrieu et al., 2010]. They also hold on both continuous and discrete state spaces, ‘lazy’ implementations where the proposal distribution is not absolutely continuous with respect to the base measure, and methods like Barker’s algorithm [Barker, 1965] where the acceptance rate function differs from the usual MH form. Thus, our characterization of kernel couplings and maximal kernel couplings applies to most Markov chains which involve a sequence of proposal and acceptance steps.

The rest of this paper is organized as follows. In Section 2, we describe our notation and setting, state our main result on transition kernel couplings, and set out a series of definitions and lemmas to prove it. We introduce a running example on a two-point state space and return to it to illustrate concepts and questions along the way. In Section 3, we highlight a few important properties of maximal couplings, consider the relationship between maximal proposal and transition kernel couplings, and prove our main result on the structure of maximal couplings of the MH transition kernel. In Section 4, we analyze the kernel couplings of several widely used MH algorithms such as the random-walk Metropolis-Hastings algorithm and the Metropolis-adjusted Langevin algorithm. Finally, in Section 5 we discuss our results, highlighting open questions and paths forward.

## 2 Metropolis–Hastings kernel couplings

In this section we show that every MH-like transition kernel coupling can be represented as a proposal coupling followed by an acceptance indicator coupling (Theorem 1). After establishing our setting and notation, we build up to this proof through several auxiliary results (Lemmas 1–7). The key step is Lemma 5, in which we show how to map from an arbitrary coupling of MH-like transition kernels to a proposal coupling and a structure we call a ‘coupled acceptance mechanism.’ The representation of transition kernel couplings in terms of coupled proposal and acceptance steps follows from this result.

### 2.1 Notation and setting

Let  $(\mathcal{Z}, \mathcal{G})$  be any measurable space. We say  $\Theta : \mathcal{Z} \times \mathcal{G} \rightarrow [0, 1]$  is a Markov kernel if  $\Theta(z, \cdot)$  is a probability measure on  $(\mathcal{Z}, \mathcal{G})$  for each  $z \in \mathcal{Z}$  and if  $\Theta(\cdot, A) : \mathcal{Z} \rightarrow [0, 1]$  is measurable for each  $A \in \mathcal{G}$ . We say that  $\Theta$  is a sub-probability kernel if it satisfies the same conditions but where  $\Theta(z, \cdot)$  is a sub-probability measure for each  $z \in \mathcal{Z}$  rather than a probability measure. Let  $2^S$  denote the power set of a set  $S$ , let  $\text{Bern}(\alpha)$  denote the Bernoulli distribution on  $\{0, 1\}$  with  $\mathbb{P}(\text{Bern}(\alpha) = 1) = \alpha$  for  $\alpha \in [0, 1]$ , and let  $a \wedge b := \min(a, b)$  for any  $a, b \in \mathbb{R}$ .

Suppose that  $(\mathcal{X}, \mathcal{F})$  is a Polish space with base measure  $\lambda$ . We take  $\mathcal{X}$  to be our state space. Fix a Markov kernel  $Q : \mathcal{X} \times \mathcal{F} \rightarrow [0, 1]$ , which we call the proposal kernel, and a function  $a : \mathcal{X} \times \mathcal{X} \rightarrow [0, 1]$ , which we call the acceptance rate function. To be an acceptance rate function, we require that  $a(x, \cdot)$  is  $Q(x, \cdot)$ -measurable and  $a(x, x) = 1$  for all  $x \in \mathcal{X}$ . The second condition simplifies many proofs and involves no loss of generality, as we discuss below. Given a current state  $x \in \mathcal{X}$  and a measurable set  $A \in \mathcal{F}$ , we think of  $Q(x, A)$  as the probability of proposing a move from  $x$  to  $x' \in A$  and  $a(x, x')$  as the probability of accepting this proposal. We call  $B(x, x') := \text{Bern}(a(x, x'))$  the acceptance indicator distribution associated with the acceptance rate function  $a$ .

In this paper we consider transition kernels which generalize the Metropolis–Hastings algorithm according to the following property:

**Definition 1.** (MH-like transition kernel) We say a Markov kernel  $P : \mathcal{X} \times \mathcal{F} \rightarrow [0, 1]$  is *generated by  $Q$  and  $a$*  if for all  $x \in \mathcal{X}$ ,  $x' \sim Q(x, \cdot)$  and  $b_x \sim B(x, x') = \text{Bern}(a(x, x'))$  imply  $X := b_x x' + (1 - b_x)x \sim P(x, \cdot)$ . We say  $P$  is *generated by  $Q$  and  $B$*  if  $P$  is generated by  $Q$  and  $a$  and if  $B$  is the acceptance indicator distribution associated with  $a$ . Finally, we say that a Markov transition kernel  $P$  is *MH-like* if there exists a Markov kernel  $Q : \mathcal{X} \times \mathcal{F} \rightarrow [0, 1]$  and an acceptance rate function  $a : \mathcal{X} \times \mathcal{X} \rightarrow [0, 1]$  such that  $P$  is generated by  $Q$  and  $a$ .

Most MCMC algorithms that involve alternating proposal and acceptance steps are MH-like. The MH case arises when we fix a target distribution  $\pi \ll \lambda$ , assume  $Q(x, \cdot) \ll \lambda$  for all  $x \in \mathcal{X}$ , and set  $a(x, x') := 1 \wedge \frac{\pi(x')q(x', x)}{\pi(x)q(x, x')}$  with  $\pi(\cdot) := d\pi/d\lambda$  and  $q(x, \cdot) := dQ(x, \cdot)/d\lambda$ . Our analysis also holds for alternative forms of  $a(x, x')$  and for proposal kernels which are not absolutely continuous with respect to  $\lambda$ . The requirement that  $a(x, x) = 1$  for all  $x \in \mathcal{X}$  involves no loss of generality because any transition kernel  $P$  that can be expressed in terms of some  $\tilde{a}$  without this property can also be expressed in terms of another acceptance function  $a$  which has it. Thus our analysis applies to continuous and discrete state spaces, lazy algorithms in which  $Q(x, \{x\}) > 0$ , and the transition kernel of Barker’s algorithm [Barker, 1965]. We refer the reader to Andrieu et al. [2020] for a general framework for understanding MH-like transition kernels.

Next, let  $\mu$  and  $\nu$  be any finite measures on  $(\mathcal{X}, \mathcal{F})$ . A measure  $\gamma$  on  $(\mathcal{X} \times \mathcal{X}, \mathcal{F} \otimes \mathcal{F})$  is called a coupling of  $\mu$  and  $\nu$  if  $\gamma(A \times \mathcal{X}) = \mu(A)$  and  $\gamma(\mathcal{X} \times A) = \nu(A)$  for all  $A \in \mathcal{F}$ . We write  $\Gamma(\mu, \nu)$  for the set of couplings of  $\mu$  and  $\nu$ , sometimes called the Fréchet class of these measures [Kendall, 2017]. When  $\mu$  and  $\nu$  are probability measures, the coupling inequality states that  $\mathbb{P}(X = Y) \leq 1 - \|\mu - \nu\|_{\text{TV}}$  for  $(X, Y) \sim \gamma \in \Gamma(\mu, \nu)$ . Here  $\|\mu - \nu\|_{\text{TV}} = \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)|$  is the total variation distance. See e.g. Lindvall [1992, chap. 1.2] or Levin et al. [2017, chap. 4] for a further discussion of this important inequality. A coupling  $\gamma$  that achieves the coupling inequality bound is said to be maximal, and we write  $\Gamma^{\max}(\mu, \nu)$  for the set of these.

Suppose that  $\Theta$  is a Markov kernel on  $(\mathcal{Z}, \mathcal{G})$ . We follow Douc et al. [2018, chap.19] in calling  $\bar{\Theta} : (\mathcal{Z} \times \mathcal{Z}) \times (\mathcal{G} \otimes \mathcal{G}) \rightarrow [0, 1]$  a kernel coupling based on  $\Theta$  if  $\bar{\Theta}$  is a Markov kernel on  $(\mathcal{Z} \times \mathcal{Z}, \mathcal{G} \otimes \mathcal{G})$  and if  $\bar{\Theta}((z_1, z_2), \cdot) \in \Gamma(\Theta(z_1, \cdot), \Theta(z_2, \cdot))$  for all  $z_1, z_2 \in \mathcal{Z}$ . We write  $\Gamma(\Theta, \Theta)$  for the set of all such kernel couplings. Likewise we say that  $\bar{\Theta}$  is a maximal kernel coupling if each  $\bar{\Theta}((z_1, z_2), \cdot)$  is a maximal coupling of  $\Theta(z_1, \cdot)$  and  $\Theta(z_2, \cdot)$ , and we write  $\Gamma^{\max}(\Theta, \Theta)$  for the set of these. It is important to remember that a maximal kernel coupling  $\bar{P} \in \Gamma^{\max}(P, P)$  does not usually correspond to a maximal coupling of Markov chains in the sense of Aldous [1983]; the former produces the highest probability of meeting at each individual step, while the latter produces the smallest meeting times allowed by the coupling inequality.

Suppose that an MH-like kernel  $P$  is generated by a proposal kernel  $Q$  and an acceptance rate function  $a$ . Then we call any  $\bar{P} \in \Gamma(P, P)$  a transition kernel coupling and any  $\bar{Q} \in \Gamma(Q, Q)$  a proposal kernel coupling. Let  $\mathcal{B}$  be the power set of  $\{0, 1\}^2$ . We call  $\bar{B} : \mathcal{X}^2 \times \mathcal{X}^2 \times \mathcal{B} \rightarrow [0, 1]$  an acceptance indicator coupling if  $\bar{B}((x, y), (x', y'), \cdot)$  is a measure on  $\{0, 1\}^2$  for all  $x, y, x', y' \in \mathcal{X}$  and if  $\bar{B}(\cdot, \cdot, S) : \mathcal{X}^2 \times \mathcal{X}^2 \rightarrow [0, 1]$  is measurable for all  $S \in \mathcal{B}$ . Abusing notation, we usually drop the third argument of  $\bar{B}$  and write  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$  whenever a pair of random variables  $(b_x, b_y) \in \{0, 1\}^2$  follows the acceptance indicator coupling  $\bar{B}$ . Thus we say that a proposal coupling  $\bar{Q}$  and an acceptance indicator coupling  $\bar{B}$  generate a kernel coupling  $\bar{P}$  if for all  $x, y \in \mathcal{X}$ ,  $(x', y') \sim \bar{Q}((x, y), \cdot)$  and  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$  imply that  $(X, Y) := (b_x x' + (1 - b_x)x, b_y y' + (1 - b_y)y) \sim \bar{P}((x, y), \cdot)$ . The key challenges before us are to show that every  $\bar{P} \in \Gamma(P, P)$  arises in this way and to determine conditions on  $\bar{B}$  such that pairs  $(X, Y)$  defined as above follow a kernel coupling  $\bar{P} \in \Gamma(P, P)$ .

We now introduce a simple example that we will return to over the course of this section.

**Example 1.** Let  $\mathcal{X} = \{1, 2\}$  and  $\mathcal{F} = 2^{\mathcal{X}}$ , and assume a current state pair of  $(x, y) = (1, 2)$ . For a finite state-space like this, it is convenient to represent kernels  $\Theta : \mathcal{X} \times \mathcal{F} \rightarrow [0, 1]$  as  $\Theta(z, \cdot) = (\Theta(z, \{1\}), \Theta(z, \{2\}))$  and couplings  $\bar{\Theta} : (\mathcal{X} \times \mathcal{X}) \times (\mathcal{F} \otimes \mathcal{F}) \rightarrow [0, 1]$  by a matrix like the following:

$$\bar{\Theta}((x, y), \cdot) = \begin{array}{c} \begin{array}{c} x \\ \downarrow \end{array} \\ \begin{array}{cc|cc} \hline \Theta((x, y), (1, 1)) & \Theta((x, y), (2, 1)) & & \\ \hline \Theta((x, y), (1, 2)) & \Theta((x, y), (2, 2)) & \leftarrow y & \\ \hline \end{array} \\ \begin{array}{cc} 1 & 2 \end{array} \end{array}$$

The columns of this matrix represent the destination values for the  $x$  chain (e.g. values of  $X$  or  $x'$ ) while the rows represent the equivalent for the  $y$  chain. The small ‘ $x$ ’ and ‘ $y$ ’ serve as a reminder of the current state pair.

For this example, we assume a uniform proposal distribution and a MH acceptance rate function  $a$  based on a target distribution  $\pi = (1/3, 2/3)$ , so that  $a(1, \cdot) = (1, 1)$  and  $a(2, \cdot) = (1/2, 1)$ . Together these imply a transition kernel  $P$  with  $P(1, \cdot) = (1/2, 1/2)$  and  $P(2, \cdot) = (1/4, 3/4)$ . Simple algebra shows that any proposal and transition coupling based on the kernels above must take the following forms, for some  $\rho \in [0, 1/2]$  and  $\lambda \in [0, 1/4]$ :

$$\bar{Q}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{|c|c|} \hline \rho & 1/2 - \rho \\ \hline 1/2 - \rho & \rho \\ \hline \end{array} \leftarrow y \\ \begin{array}{cc} 1 & 2 \end{array} \end{array} \quad \bar{P}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{|c|c|} \hline \lambda & 1/4 - \lambda \\ \hline 1/2 - \lambda & \lambda + 1/4 \\ \hline \end{array} \leftarrow y \\ \begin{array}{cc} 1 & 2 \end{array} \end{array}$$

We return to this example to illustrate the concepts below.

## 2.2 Main result

Johnson [1998] showed that one can start with a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and construct an acceptance indicator coupling  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  generate a valid transition kernel coupling  $\bar{P} \in \Gamma(P, P)$ . There and in subsequent work such as Jacob et al. [2020],  $\bar{B}$  is taken to be a coupling of  $\text{Bern}(a(x, x'))$  and  $\text{Bern}(a(y, y'))$ , so that proposals from  $x$  to  $x'$  and from  $y$  to  $y'$  are each accepted at the MH rate. O’Leary et al. [2021] used a conditional form of  $\bar{B}$  to produce a maximal transition kernel coupling  $\bar{P}$  from any maximal proposal kernel coupling  $\bar{Q}$ . In this paper we have a more general goal, which is to characterize the set of all possible transition kernel couplings  $\Gamma(P, P)$  in terms of proposal and acceptance indicator couplings. In particular, we prove the following result:

**Theorem 1.** *Let  $P$  be the MH-like transition kernel on  $(\mathcal{X}, \mathcal{F})$  generated by a proposal kernel  $Q$  and an acceptance rate function  $a$ . A joint kernel  $\bar{P} \in \Gamma(P, P)$  if and only if it is generated by  $\bar{Q} \in \Gamma(Q, Q)$  and an acceptance indicator coupling  $\bar{B}$  such that for any  $x, y \in \mathcal{X}$  and  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , we have*

1.  $\mathbb{P}(b_x = 1 \mid x, y, x') = a(x, x')$  for  $Q(x, \cdot)$ -almost all  $x'$ , and
2.  $\mathbb{P}(b_y = 1 \mid x, y, y') = a(y, y')$  for  $Q(y, \cdot)$ -almost all  $y'$ .

This result shows that all MH-like transition kernel couplings  $\bar{P}$  are ‘natural’ in the sense that they arise from coupled proposals  $(x', y') \sim \bar{Q}((x, y), \cdot)$  that are accepted or rejected according to coupled acceptance indicators  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ . The  $\bar{B}$  conditions say that from any starting point  $(x, y)$ , the  $x'$  and  $y'$  proposals are marginally accepted at the rate given by  $a$ , potentially with complicated joint behavior. Theorem 1 also confirms that simple conditions on the joint proposal and acceptance steps are enough to yield a kernel coupling  $\bar{P}$  in  $\Gamma(P, P)$ .

It is not hard to show that  $\bar{P} \in \Gamma(P, P)$  when  $\bar{P}$  is generated by a proposal coupling  $\bar{Q}$  and an acceptance indicator coupling  $\bar{B}$  with the properties given above. However, the converse requires a detailed understanding of the structure of transition kernel couplings. One challenge is as follows: a transition from a state pair  $(x, y)$  to a measurable rectangle  $A_x \times A_y$  with  $x \notin A_x$  and  $y \notin A_y$  can only occur if points  $x' \neq x$  and  $y' \neq y$  are proposed and accepted. This makes it simple to work out the behavior of a hypothetical  $\bar{Q}$  and  $\bar{B}$  on these sets. However, transitions to sets like  $A_x \times \{y\}$ ,  $\{x\} \times A_y$ , and  $\{x\} \times \{y\}$  can arise from the partial or full rejection of proposed moves to sets like  $A_x \times \mathcal{X}$ ,  $\mathcal{X} \times A_y$ , and  $\mathcal{X} \times \mathcal{X}$ . It is more challenging to associate these transition probabilities with joint proposal and acceptance probabilities in a consistent way.

Nevertheless, we prove our Theorem 1 by constructing a mapping from an arbitrary kernel  $\bar{P} \in \Gamma(P, P)$  to a joint distribution  $\bar{Q}$  on proposals  $(x', y')$  and a coupling  $\bar{B}$  on acceptance indicators  $(b_x, b_y)$ , such these reproduce  $\bar{P}$  and agree with the original proposal kernel  $Q$  and

acceptance rate function  $a$ . The first step of this procedure involves defining a collection of joint measures  $\Phi$  on  $(x', y')$  and  $(b_x, b_y)$  with certain adding-up properties. A proposal coupling  $\bar{Q}$  follows directly from  $\Phi$ , and a Radon–Nikodym argument shows that we can also extract an acceptance indicator coupling  $\bar{B}$  with the desired properties.

### 2.3 Coupled acceptance mechanisms

To begin, we define a relationship that can exist between an MH-like transition kernel coupling  $\bar{P}$  and an arbitrary proposal coupling  $\bar{Q}$ :

**Definition 2.** We say that a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and a transition kernel coupling  $\bar{P} \in \Gamma(P, P)$  are related by a *coupled acceptance mechanism*  $\Phi = (\Phi_{11}, \Phi_{10}, \Phi_{01}, \Phi_{00})$  if each  $\Phi_{ij}((x, y), \cdot)$  is a sub-probability on  $(\mathcal{X} \times \mathcal{X}, \mathcal{F} \otimes \mathcal{F})$ , and for all  $x, y \in \mathcal{X}$ ,  $A_x, A_y \in \mathcal{F}$  we have

1.  $\bar{Q}((x, y), A_x \times A_y) = (\Phi_{11} + \Phi_{10} + \Phi_{01} + \Phi_{00})((x, y), A_x \times A_y)$
2.  $\bar{P}((x, y), A_x \times A_y) = \Phi_{11}((x, y), A_x \times A_y) + \Phi_{10}((x, y), A_x \times \mathcal{X})1(y \in A_y) + \Phi_{01}((x, y), \mathcal{X} \times A_y)1(x \in A_x) + \Phi_{00}((x, y), \mathcal{X} \times \mathcal{X})1(x \in A_x)1(y \in A_y)$
3.  $Q(x, \{x\}) = (\Phi_{11} + \Phi_{10})((x, y), \{x\} \times \mathcal{X})$  and  $Q(y, \{y\}) = (\Phi_{11} + \Phi_{01})((x, y), \mathcal{X} \times \{y\})$ .

As we will see, the existence of a coupled acceptance mechanism  $\Phi$  relating  $\bar{Q}$  and  $\bar{P}$  is equivalent to the existence of an acceptance indicator coupling  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ . In Lemma 1 we show that if  $(X, Y) \sim \bar{P}((x, y), \cdot)$  is generated by  $(x', y') \sim \bar{Q}((x, y), \cdot)$  and  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , then a coupled acceptance mechanism  $\Phi$  relating  $\bar{Q}$  and  $\bar{P}$  exists and can be defined by  $\Phi_{ij}((x, y), \cdot) := \mathbb{P}((x', y') \in \cdot, b_x = i, b_y = j)$  for  $i, j \in \{0, 1\}$ . Conversely, in Lemma 6, we show that if a coupled acceptance mechanism  $\Phi$  relates  $\bar{Q}$  and  $\bar{P}$ , then there exists an acceptance indicator coupling  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ .

Thus Definition 2 captures a set of relationships that must exist between a proposal coupling  $\bar{Q}$  and a transition kernel coupling  $\bar{P}$  for the former to generate the latter. We think of  $\Phi$  as subdividing the probability in  $\bar{Q}((x, y), \cdot)$  into four acceptance scenarios, depending on whether one, both, or neither proposal is accepted. See Figure 1 for an illustration of this. Condition 1 requires that the probability assigned to these scenarios must add up to  $\bar{Q}$  on all measurable subsets of  $\mathcal{X} \times \mathcal{X}$ . Condition 2 says that the resulting distribution over transitions must agree with  $\bar{P}$ . Condition 3 concerns proposals with  $x' = x$  or  $y' = y$ , and says that  $\Phi$  must agree with the marginal assumption that such proposals are always accepted.

The crux of Theorem 1 is to prove that every coupling of MH-like transition kernels  $\bar{P}$  arises from a proposal and acceptance coupling. Thus we begin by looking for coupled acceptance mechanisms  $\Phi$  that distribute the probability  $\bar{Q}((x, y), A_x \times A_y)$  of a proposal from  $(x, y)$  to  $(x', y') \in A_x \times A_y$  into contributions to the probability of a transition from  $(x, y)$  to  $(X, Y) \in A_x \times A_y$ ,  $A_x \times \{y\}$ ,  $\{x\} \times A_y$ , and  $\{x\} \times \{y\}$ , in a consistent way. The following result establishes the necessity of the conditions in Definition 2 for this task.

**Lemma 1.** *Let  $\bar{P} \in \Gamma(P, P)$  be a coupling of MH-like transition kernels. If a proposal coupling  $\bar{Q}$  and an acceptance indicator coupling  $\bar{B}$  generate  $\bar{P}$ , then there exists a coupled acceptance mechanism  $\Phi$  relating  $\bar{Q}$  and  $\bar{P}$ .*

*Proof.* Fix  $(x, y)$ , let  $(x', y') \sim \bar{Q}((x, y), \cdot)$ , and let  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ . For  $A \in \mathcal{F} \otimes \mathcal{F}$  and  $i, j \in \{0, 1\}$ , define  $\Phi_{ij}((x, y), A) := \mathbb{P}((x', y') \in A, b_x = i, b_y = j \mid x, y)$ . This  $\Phi$  satisfies




Figure 1: Schematic diagram of a coupled acceptance mechanism  $\Phi$  relating a proposal coupling  $\bar{Q}$  and a transition kernel coupling  $\bar{P}$ . Here  $(x, y)$  represents the current state and  $A_x \times A_y$  is a measurable rectangle in  $\mathcal{X} \times \mathcal{X}$ .  $\bar{Q}((x, y), A_x \times A_y)$  gives the probability of a proposal  $(x', y') \in A_x \times A_y$ . The coupled acceptance mechanism  $\Phi = (\Phi_{11}, \Phi_{10}, \Phi_{01}, \Phi_{00})$  distributes this probability into contributions to the probability  $\bar{P}((x, y), \cdot)$  of a transition from  $(x, y)$  to the sets  $A_x \times A_y$ ,  $A_x \times \{y\}$ ,  $\{x\} \times A_y$ , and  $\{x\} \times \{y\}$ . The conditions of Definition 2 ensure that  $\Phi$  agrees with both  $\bar{Q}$  and  $\bar{P}$ .

Condition 1, since  $\bar{Q}((x, y), A_x \times A_y) = \mathbb{P}((x', y') \in A_x \times A_y | x, y)$ . It also satisfies Condition 2, since for all  $A_x, A_y \in \mathcal{F}$  we have

$$\begin{aligned}
\bar{P}((x, y), A_x \times A_y) &= \mathbb{P}(X \in A_x, Y \in A_y | x, y) \\
&= \mathbb{P}(x' \in A_x, y' \in A_y, b_x = 1, b_y = 1 | x, y) + \mathbb{P}(x' \in A_x, y \in A_y, b_x = 1, b_y = 0 | x, y) \\
&\quad + \mathbb{P}(x \in A_x, y' \in A_y, b_x = 0, b_y = 1 | x, y) + \mathbb{P}(x \in A_x, y \in A_y, b_x = 0, b_y = 0 | x, y) \\
&= \Phi_{11}((x, y), A_x \times A_y) + \Phi_{10}((x, y), A_x \times \mathcal{X})1(y \in A_y) \\
&\quad + \Phi_{01}((x, y), \mathcal{X} \times A_y)1(x \in A_x) + \Phi_{00}((x, y), \mathcal{X} \times \mathcal{X})1(x \in A_x)1(y \in A_y).
\end{aligned}$$

For Condition 3, we have

$$\begin{aligned}
(\Phi_{11} + \Phi_{10})((x, y), \{x\} \times \mathcal{X}) &= \mathbb{P}(x' = x, b_x = 1) = a(x, x)Q(x, \{x\}) = Q(x, \{x\}) \\
(\Phi_{11} + \Phi_{01})((x, y), \mathcal{X} \times \{y\}) &= \mathbb{P}(y' = y, b_y = 1) = a(y, y)Q(y, \{y\}) = Q(y, \{y\}).
\end{aligned}$$

Thus  $\Phi$  is a coupled acceptance mechanism relating  $\bar{Q}$  and  $\bar{P}$ .  $\square$

The  $\bar{P}$  condition of Definition 2 takes a more intuitive (if less compact) form when we express it in terms of measurable sets  $A_x, A_y \in \mathcal{F}$  where  $x \notin A_x$  and  $y \notin A_y$ :

**Lemma 2.** *Let  $\bar{P} \in \Gamma(P, P)$  and suppose  $\Phi = (\Phi_{11}, \Phi_{10}, \Phi_{01}, \Phi_{00})$  are a collection of measures on  $(\mathcal{X} \times \mathcal{X}, \mathcal{F} \otimes \mathcal{F})$ . Then Condition 2 of Definition 2 holds if and only if for all  $A_x, A_y \in \mathcal{F}$*

with  $x \notin A_x$  and  $y \notin A_y$ , we have

1.  $\bar{P}((x, y), A_x \times A_y) = \Phi_{11}((x, y), A_x \times A_y)$
2.  $\bar{P}((x, y), A_x \times \{y\}) = \Phi_{11}((x, y), A_x \times \{y\}) + \Phi_{10}((x, y), A_x \times \mathcal{X})$
3.  $\bar{P}((x, y), \{x\} \times A_y) = \Phi_{11}((x, y), \{x\} \times A_y) + \Phi_{01}((x, y), \mathcal{X} \times A_y)$
4.  $\bar{P}((x, y), \{x\} \times \{y\}) = \Phi_{11}((x, y), \{x\} \times \{y\}) + \Phi_{10}((x, y), \{x\} \times \mathcal{X})$   
 $+ \Phi_{01}((x, y), \mathcal{X} \times \{y\}) + \Phi_{00}((x, y), \mathcal{X} \times \mathcal{X}).$

*Proof.* The conditions above follow from Condition 2 of Definition 2 by evaluating that condition at  $\{x\}, A_x \setminus \{x\}$  and  $\{y\}, A_y \setminus \{y\}$ . For the converse, for any  $A_x, A_y \in \mathcal{F}$  we have

$$\begin{aligned} \bar{P}((x, y), A_x \times A_y) &= \bar{P}((x, y), (A_x \setminus \{x\}) \times (A_y \setminus \{y\})) + \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\})1(y \in A_y) \\ &+ \bar{P}((x, y), \{x\} \times (A_y \setminus \{y\}))1(x \in A_x) + \bar{P}((x, y), \{x\} \times \{y\})1(x \in A_x)1(y \in A_y). \end{aligned}$$

Replacing these terms with Conditions 1-4 above yields Condition 2 of Definition 2.  $\square$

Condition 1 of Lemma 2 corresponds to the intuition that the only way to transition from  $(x, y)$  to a point  $(X, Y) \in A_x \times A_y$  with  $x \notin A_x$  and  $y \notin A_y$  is to propose and accept  $(x', y') = (X, Y)$ . Similarly, Condition 2 says that the probability of transitioning to a point  $(X, Y) \in A_x \times \{y\}$  must come from the combined probability of accepting a proposal to such a point or by proposing a move  $(x', y') \in A_x \times \mathcal{X}$ , with  $x'$  accepted and  $y'$  rejected. The other two conditions follow a similar intuition.

Definition 2 raises two questions. First, we may ask if Condition 3 is actually independent of Conditions 1 and 2. Second, we may ask whether a given  $\bar{P} \in \Gamma(P, P)$  can be related to more than one  $\bar{Q} \in \Gamma(Q, Q)$  by coupled acceptance mechanisms. The next example answers both of these questions.

**Example 2.** We return to the setup of Example 1, in which  $\mathcal{X} = \{1, 2\}$ ,  $\mathcal{F} = 2^{\mathcal{X}}$ ,  $(x, y) = (1, 2)$ ,  $Q(1, \cdot) = Q(2, \cdot) = (1/2, 1/2)$ ,  $P(1, \cdot) = (1/2, 1/2)$ , and  $P(2, \cdot) = (1/4, 3/4)$ . Suppose that we have the following proposal and transition kernel couplings:

$$\bar{Q}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{cc|c} 1 & \begin{array}{cc} 1/4 & 1/4 \end{array} & \\ \hline 2 & \begin{array}{cc} 1/4 & 1/4 \end{array} & \leftarrow y \\ \hline & 1 & 2 \end{array} \end{array} \quad \bar{P}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{cc|c} 1 & \begin{array}{cc} 0 & 1/4 \end{array} & \\ \hline 2 & \begin{array}{cc} 1/2 & 1/4 \end{array} & \leftarrow y \\ \hline & 1 & 2 \end{array} \end{array}$$

Using Lemma 2, the first two conditions of Definition 2 imply that any coupled acceptance mechanism  $\Phi$  relating  $\bar{Q}$  and  $\bar{P}$  must take the following form:

$$\begin{aligned} \Phi_{11}((1, 2), \cdot) &= \begin{array}{c} \begin{array}{cc|c} 1 & \begin{array}{cc} 0 & 1/4 \end{array} & \\ \hline 2 & \begin{array}{cc} a & b \end{array} & \\ \hline & 1 & 2 \end{array} \\ \Phi_{10}((1, 2), \cdot) &= \begin{array}{c} \begin{array}{cc|c} 1 & \begin{array}{cc} d & 0 \end{array} & \\ \hline 2 & \begin{array}{cc} c & 1/4 - b \end{array} & \\ \hline & 1 & 2 \end{array} \\ \Phi_{01}((1, 2), \cdot) &= \begin{array}{c} \begin{array}{cc|c} 1 & \begin{array}{cc} 0 & 0 \end{array} & \\ \hline 2 & \begin{array}{cc} e & 0 \end{array} & \\ \hline & 1 & 2 \end{array} \\ \Phi_{00}((1, 2), \cdot) &= \begin{array}{c} \begin{array}{cc|c} 1 & \begin{array}{cc} 1/4 - d & 0 \end{array} & \\ \hline 2 & \begin{array}{cc} 1/4 - a - c - e & 0 \end{array} & \\ \hline & 1 & 2 \end{array} \end{array} \end{aligned}$$



Thus  $(\Phi_{11} + \Phi_{10})((x, y), \{x\} \times \mathcal{X}) = a + c + d$ , and  $(\Phi_{11} + \Phi_{01})((x, y), \mathcal{X} \times \{y\}) = a + b + e$ . Note however that  $a + c + d = Q(x, \{x\}) = 1/2$  and  $a + b + e = Q(y, \{y\}) = 1/2$  do not automatically hold without imposing the third condition of Definition 2. Also observe that even with that condition, any choice of  $a, b, c$  with  $0 \leq c \leq b \leq 1/4$  and  $0 \leq a \leq 1/4 - b \vee c$  will produce a valid  $\Phi$  relating  $Q$  and  $P$ . Thus we see that coupled acceptance mechanisms do not have to be unique.

## 2.4 Existence of a coupled acceptance mechanism

To prove Theorem 1 we must show that any MH-like transition kernel coupling  $\bar{P} \in \Gamma(P, P)$  arises from a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and an acceptance indicator coupling  $\bar{B}$  with certain properties. In this subsection we show that for any  $\bar{P} \in \Gamma(P, P)$ , there exists a kernel coupling  $\bar{Q} \in \Gamma(Q, Q)$  and a coupled acceptance mechanism  $\Phi$  relating  $\bar{Q}$  and  $\bar{P}$ . In the next subsection we show that we can transform this  $\Phi$  into an acceptance coupling  $\bar{B}$ . Finally, we use these results to prove our main theorem. Along the way we use the following objects, which are defined for  $x \in \mathcal{X}$  and  $A \in \mathcal{F}$ :

$$\begin{aligned} \alpha_0(x, A) &:= Q(x, A \setminus \{x\}) - P(x, A \setminus \{x\}) & \alpha_1(x, A) &:= Q(x, A \cap \{x\}) + P(x, A \setminus \{x\}) \\ \beta(x) &:= \begin{cases} \frac{Q(x, \{x\})}{P(x, \{x\})} & \text{if } P(x, \{x\}) > 0 \\ 1 & \text{otherwise} \end{cases} & \mu(x, A) &:= \begin{cases} \frac{\alpha_0(x, A)}{\alpha_0(x, \mathcal{X})} & \text{if } \alpha_0(x, \mathcal{X}) > 0 \\ 1(x \in A) & \text{otherwise.} \end{cases} \end{aligned}$$

Note that these definitions depend only on the marginal kernels  $Q$  and  $P$ . These objects have a simple interpretation in terms of the underlying chains:

**Lemma 3.** *Let  $x \in \mathcal{X}$ ,  $A \in \mathcal{F}$ ,  $x' \sim Q(x, \cdot)$ ,  $b_x \sim B(x, x') = \text{Bern}(a(x, x'))$ , and so  $X := b_x x' + (1 - b_x)x \sim P(x, \cdot)$ . Then  $\alpha_i(x, A) = \mathbb{P}(x' \in A, b_x = i | x)$  for  $i \in \{0, 1\}$ ;  $\beta(x) = \mathbb{P}(b_x = 1 | x, X = x)$  whenever  $P(x, \{x\}) > 0$ ; and  $\mu(x, A) = \mathbb{P}(x' \in A | b_x = 0, x)$  whenever  $\alpha_0(x, \mathcal{X}) > 0$ .*

*Proof.* We have assumed that the acceptance rate function  $a$  has  $a(x, x) = 1$  for all  $x$ . Thus  $\mathbb{P}(x' \in A \cap \{x\}, b_x = 0 | x) = 0$  and so

$$\begin{aligned} \alpha_0(x, A) &= Q(x, A \setminus \{x\}) - P(x, A \setminus \{x\}) = \mathbb{P}(x' \in A \setminus \{x\} | x) - \mathbb{P}(X \in A \setminus \{x\} | x) \\ &= \mathbb{P}(x' \in A | x) - \mathbb{P}(x' \in A, b_x = 1 | x) = \mathbb{P}(x' \in A, b_x = 0 | x). \end{aligned}$$

Similarly,

$$\begin{aligned} \alpha_1(x, A) &= Q(x, A \cap \{x\}) + P(x, A \setminus \{x\}) = \mathbb{P}(x' \in A \cap \{x\} | x) + \mathbb{P}(X \in A \setminus \{x\} | x) \\ &= \mathbb{P}(x' \in A \cap \{x\}, b_x = 1 | x) + \mathbb{P}(x' \in A \setminus \{x\}, b_x = 1 | x) = \mathbb{P}(x' \in A, b_x = 1 | x). \end{aligned}$$

Suppose  $P(x, \{x\}) > 0$ . Then since  $Q(x, \{x\}) = \alpha_x(x, \{x\})$ , we have

$$\beta(x) = \frac{Q(x, \{x\})}{P(x, \{x\})} = \frac{\mathbb{P}(b_x = 1, x' = x | x)}{\mathbb{P}(X = x | x)} = \frac{\mathbb{P}(b_x = 1, X = x | x)}{\mathbb{P}(X = x | x)} = \mathbb{P}(b_x = 1 | x, X = x).$$

Finally, say  $\alpha_0(x, \mathcal{X}) = \mathbb{P}(b_x = 0 | x) > 0$ . Then

$$\mu(x, A) = \frac{\alpha_0(x, A)}{\alpha_0(x, \mathcal{X})} = \frac{\mathbb{P}(x' \in A, b_x = 0 | x)}{\mathbb{P}(b_x = 0 | x)} = \mathbb{P}(x' \in A | b_x = 0, x).$$

Thus each of  $\alpha_0, \alpha_1, \beta$ , and  $\mu$  has a simple interpretation in terms of the joint probability of  $x'$  and  $b_x$  induced by  $Q$  and  $B$ .  $\square$

We will also need the following properties:

**Lemma 4.** *Let  $x \in \mathcal{X}$  and  $A \in \mathcal{F}$ . Then  $\beta(x) \in [0, 1]$ ,  $\mu(x, \cdot)$  is a measure on  $(\mathcal{X}, \mathcal{F})$ ,  $Q(x, A \cap \{x\}) = P(x, A \cap \{x\})\beta(x)$ , and  $Q(x, A) = \alpha_1(x, A) + \mu(x, A)\alpha_0(x, \mathcal{X})$ .*

*Proof.* When  $P(x, \{x\}) > 0$ ,  $\beta(x) \in [0, 1]$  by Lemma 3. Otherwise  $\beta(x) = 1$ , so in either case  $\beta(x) \in [0, 1]$ . Also by Lemma 3, we observe that  $\mu(x, \cdot)$  is the indicator of a measurable set or a well-defined conditional probability, and so in either case it defines a measure on  $(\mathcal{X}, \mathcal{F})$ .

For the first  $Q$  equality, if  $x \notin A$ , then the  $Q(x, A \cap \{x\}) = 0 = P(x, A \cap \{x\})\beta(x)$ . If  $x \in A$  then we want to show  $Q(x, \{x\}) = P(x, \{x\})\beta(x)$ . This holds by the definition of  $\beta(x)$  when  $P(x, \{x\}) > 0$ . Since  $a(x, x) = 1$  for all  $x \in \mathcal{X}$ ,  $P(x, \{x\}) = 0$  implies  $Q(x, \{x\}) = 0$ , so the result also holds in this case.

The second  $Q$  equality holds by the definitions of  $\mu$ ,  $\alpha_0$ , and  $\alpha_1$  when  $\alpha_0(x, \mathcal{X}) > 0$ . This leaves the  $\alpha_0(x, \mathcal{X}) = 0$  case. In general,  $\alpha_0(x, \mathcal{X}) = Q(x, \{x\}^c) - P(x, \{x\}^c) = \int (1 - a(x, x'))Q(x, dx')$ , since  $a(x, x) = 1$  for all  $x \in \mathcal{X}$ . Thus  $\alpha_0(x, \mathcal{X}) = 0$  implies  $a(x, x') = 1$  for  $Q(x, \cdot)$ -almost all  $x'$ , and so  $\alpha_1(x, A) = Q(x, A \cap \{x\}) + \int_{A \setminus \{x\}} a(x, x')Q(x, dx') = Q(x, A)$ . Thus the second equality also holds when  $\alpha_0(x, \mathcal{X}) = 0$ .  $\square$

We now describe the main lemma used in our proof of Theorem 1.

**Lemma 5.** *For any coupling  $\bar{P} \in \Gamma(P, P)$  of MH-like transition kernels, there exists a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and a coupled acceptance mechanism  $\bar{\Phi}$  relating  $\bar{Q}$  and  $\bar{P}$ .*

*Proof.* The proof proceeds in three steps. First we use  $\bar{P}$  to define a collection of sub-probability kernels  $\bar{\Phi} = (\bar{\Phi}_{11}, \bar{\Phi}_{10}, \bar{\Phi}_{01}, \bar{\Phi}_{00})$ . Then we use  $\bar{\Phi}$  to define a kernel coupling  $\bar{Q} \in \Gamma(Q, Q)$ . Finally we show that  $\bar{\Phi}$  satisfies the conditions of Definition 2, making it a coupled acceptance mechanism relating  $\bar{Q}$  and  $\bar{P}$ .

We begin with an explicit formula for  $\bar{\Phi}_{11}$  in terms of  $\bar{P}$ . For  $x, y \in \mathcal{X}$  and  $A \in \mathcal{F} \otimes \mathcal{F}$ , let

$$\begin{aligned} \bar{\Phi}_{11}((x, y), A) &:= \bar{P}\left((x, y), A \cap (\{x\}^c \times \{y\}^c)\right) + \bar{P}\left((x, y), A \cap (\{x\} \times \{y\}^c)\right) \beta(x) \\ &\quad + \bar{P}\left((x, y), A \cap (\{x\}^c \times \{y\})\right) \beta(y) + \bar{P}\left((x, y), A \cap (\{x\} \times \{y\})\right) \beta(x) \beta(y). \end{aligned}$$

Here  $\bar{\Phi}_{11}$  is a sub-probability kernel, since  $\beta(\cdot) \in [0, 1]$  by Lemma 4. We define the other three components of  $\bar{\Phi}$  by a product measure construction. For  $x, y \in \mathcal{X}$  and  $A_x, A_y \in \mathcal{F}$ , let

$$\begin{aligned} \bar{\Psi}_{10}((x, y), A_x) &:= \alpha_1(x, A_x) - \bar{\Phi}_{11}((x, y), A_x \times \mathcal{X}) \\ \bar{\Psi}_{01}((x, y), A_y) &:= \alpha_1(y, A_y) - \bar{\Phi}_{11}((x, y), \mathcal{X} \times A_y). \end{aligned}$$

We claim that  $\bar{\Psi}_{10}((x, y), \cdot)$  and  $\bar{\Psi}_{01}((x, y), \cdot)$  are sub-probabilities on  $(\mathcal{X}, \mathcal{F})$ . Indeed, by Lemma 4 and the fact that  $\bar{P} \in \Gamma(P, P)$ , we have

$$\begin{aligned} \alpha_1(x, A_x) &= P(x, A_x \setminus \{x\}) + P(x, A_x \cap \{x\})\beta(x) \\ &= \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\}^c) + \bar{P}((x, y), (A_x \cap \{x\}) \times \{y\}^c)\beta(x) \\ &\quad + \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\}) + \bar{P}((x, y), (A_x \cap \{x\}) \times \{y\})\beta(x). \end{aligned}$$

Plugging  $A_x \times \mathcal{X}$  into the definition of  $\bar{\Phi}_{11}$  yields analogous terms:

$$\begin{aligned} \bar{\Phi}_{11}((x, y), A_x \times \mathcal{X}) &= \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\}^c) + \bar{P}((x, y), (A_x \cap \{x\}) \times \{y\}^c)\beta(x) \\ &\quad + \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\})\beta(y) + \bar{P}((x, y), (A_x \cap \{x\}) \times \{y\})\beta(x)\beta(y). \end{aligned}$$

Thus  $\Psi_{10}((x, y), A_x) = (\bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\}) + \bar{P}((x, y), (A_x \cap \{x\}) \times \{y\})\beta(x))(1 - \beta(y))$ , and similarly for  $\Psi_{01}((x, y), A_y)$ . It follows that  $\Psi_{10}((x, y), \cdot)$  and  $\Psi_{01}((x, y), \cdot)$  are sub-probabilities. We also set  $\Psi_{00}(x, y) := 1 - \alpha_1(x, \mathcal{X}) - \alpha_1(y, \mathcal{Y}) + \Phi_{11}((x, y), \mathcal{X} \times \mathcal{X})$ . Algebraic manipulations show that  $\Psi_{00}(x, y) = \bar{P}((x, y), \{x\} \times \{y\})(1 - \beta(x))(1 - \beta(y)) \in [0, 1]$ .

With these results in hand, we define  $\Phi_{10}((x, y), \cdot)$ ,  $\Phi_{01}((x, y), \cdot)$ , and  $\Phi_{00}((x, y), \cdot)$  as the product measures extending the following equalities to  $(\mathcal{X} \times \mathcal{X}, \mathcal{F} \otimes \mathcal{F})$ :

$$\begin{aligned}\Phi_{10}((x, y), A_x \times A_y) &= \Psi_{10}((x, y), A_x)\mu(y, A_y) \\ \Phi_{01}((x, y), A_x \times A_y) &= \mu(x, A_x)\Psi_{01}((x, y), A_y) \\ \Phi_{00}((x, y), A_x \times A_y) &= \mu(x, A_x)\mu(y, A_y)\Psi_{00}(x, y).\end{aligned}$$

Thus each component of  $\Phi = (\Phi_{11}, \Phi_{10}, \Phi_{01}, \Phi_{00})$  is a sub-probability kernel.

Next, we define  $\bar{Q}((x, y), A) := (\Phi_{11} + \Phi_{10} + \Phi_{01} + \Phi_{00})((x, y), A)$  for all  $A \in \mathcal{F} \otimes \mathcal{F}$ . To see that  $\bar{Q} \in \Gamma(Q, Q)$ , observe that for any  $x, y \in \mathcal{X}$  and  $A_x \in \mathcal{F}$  we have

$$\begin{aligned}\bar{Q}((x, y), A_x \times \mathcal{X}) &= (\Phi_{11} + \Phi_{10} + \Phi_{01} + \Phi_{00})((x, y), A_x \times \mathcal{X}) \\ &= (\Phi_{11} + \Phi_{10})(A_x \times \mathcal{X}) + \mu(x, A_x)(\Phi_{01} + \Phi_{00})(\mathcal{X} \times \mathcal{X}) \\ &= \alpha_1(x, A_x) + \mu(x, A_x)\alpha_0(x, \mathcal{X}) = Q(x, A_x).\end{aligned}$$

The second-to-last equality follows from the definitions of  $\Psi_{10}$ ,  $\Psi_{01}$  and  $\Psi_{00}$ , while the last equality is due to Lemma 4. A similar argument shows  $\bar{Q}((x, y), \mathcal{X} \times A_y) = Q(y, A_y)$  for any  $x, y \in \mathcal{X}$  and  $A_y \in \mathcal{F}$ . Thus  $\bar{Q} \in \Gamma(Q, Q)$ .

Finally, we show that  $\Phi$  satisfies the conditions of Definition 2. The  $\bar{Q}$  condition is automatically satisfied given the definitions above. For the  $\bar{P}$  condition, we check the four cases described in Lemma 2. For the  $(A_x \setminus \{x\}) \times (A_y \setminus \{y\})$  case, we have

$$\Phi_{11}((x, y), (A_x \setminus \{x\}) \times (A_y \setminus \{y\})) = \bar{P}((x, y), (A_x \setminus \{x\}) \times (A_y \setminus \{y\})).$$

For the  $(A_x \setminus \{x\}) \times \{y\}$  case,

$$\begin{aligned}\Phi_{11}((x, y), (A_x \setminus \{x\}) \times \{y\}) + \Phi_{10}((x, y), (A_x \setminus \{x\}) \times \mathcal{X}) \\ = P(x, A_x \setminus \{x\}) - \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\}^c) = \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\}).\end{aligned}$$

We use the definitions of  $\Phi_{11}$  and  $\Phi_{10}$  and the fact that  $\alpha_1(x, A \setminus \{x\}) = P(x, A \setminus \{x\})$  for the first equality. The second equality follows from the definition of  $\bar{P} \in \Gamma(P, P)$ . By a similar argument, we have  $\Phi_{11}((x, y), \{x\} \times (A_y \setminus \{y\})) + \Phi_{10}((x, y), \mathcal{X} \times (A_y \setminus \{y\})) = \bar{P}((x, y), \{x\} \times (A_y \setminus \{y\}))$ . For the  $\{x\} \times \{y\}$  case we have

$$\begin{aligned}\Phi_{11}((x, y), \{x\} \times \{y\}) + \Phi_{10}((x, y), \{x\} \times \mathcal{X}) + \Phi_{01}((x, y), \mathcal{X} \times \{y\}) + \Phi_{00}((x, y), \mathcal{X} \times \mathcal{X}) \\ = \Phi_{11}((x, y), \{x\} \times \{y\}) + \Psi_{10}((x, y), \{x\}) + \Psi_{01}((x, y), \{y\}) + \Psi_{00}(x, y) \\ = 1 + \Phi_{11}((x, y), \{x\}^c \times \{y\}^c) - \alpha_1(x, \{x\}^c) - \alpha_1(y, \{y\}^c) = \bar{P}((x, y), \{x\} \times \{y\}).\end{aligned}$$

For the last condition of Definition 2, note that by the definitions of  $\Phi$  and  $\alpha_1$  we have

$$\begin{aligned}(\Phi_{11} + \Phi_{10})((x, y), \{x\} \times \mathcal{X}) &= \alpha_1(x, \{x\}) = Q(x, \{x\}) \\ (\Phi_{11} + \Phi_{01})((x, y), \mathcal{X} \times \{y\}) &= \alpha_1(y, \{y\}) = Q(y, \{y\}).\end{aligned}$$

We conclude that  $\Phi$  is a coupled acceptance mechanism relating  $\bar{P}$  and  $\bar{Q}$ .  $\square$

**Example 3.** We return to the setup of Example 1 to construct  $\Phi$  and  $\bar{Q}$  as in the proof of Lemma 5. We continue to assume  $\mathcal{X} = \{1, 2\}$ ,  $\mathcal{F} = 2^{\mathcal{X}}$ ,  $Q(1, \cdot) = Q(2, \cdot) = (1/2, 1/2)$ ,  $a(1, \cdot) = (1, 1)$ ,  $a(2, \cdot) = (1/2, 1)$ ,  $P(1, \cdot) = (1/2, 1/2)$ , and  $P(2, \cdot) = (1/4, 3/4)$ . Thus  $\alpha_0$ ,  $\alpha_1$ ,  $\beta$ ,  $\mu$  work out to

$$\begin{aligned} \alpha_0(1, \cdot) &= (0, 0) & \alpha_1(1, \cdot) &= (1/2, 1/2) & \mu(1, \cdot) &= (1, 0) \\ \alpha_0(2, \cdot) &= (1/4, 0) & \alpha_1(2, \cdot) &= (1/4, 1/2) & \mu(2, \cdot) &= (1, 0) \\ & & \beta(\cdot) &= (1, 2/3). \end{aligned}$$

As in Example 2, we consider a kernel coupling  $\bar{P}$  with the following transition probabilities:

$$\bar{P}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{cc|c} 1 & \begin{array}{|c|c|} \hline 0 & 1/4 \\ \hline \end{array} & \\ 2 & \begin{array}{|c|c|} \hline 1/2 & 1/4 \\ \hline \end{array} & \leftarrow y \\ \hline & 1 & 2 \end{array} \end{array}$$

Following the construction in the proof of Lemma 5 and the values of  $\alpha_0$ ,  $\alpha_1$ ,  $\mu$ , and  $\beta$  given above, we have

$$\Phi_{11}((1, 2), \cdot) = \begin{array}{c} \begin{array}{cc|c} 1 & \begin{array}{|c|c|} \hline 0 & 1/4 \\ \hline \end{array} & \\ 2 & \begin{array}{|c|c|} \hline 1/3 & 1/6 \\ \hline \end{array} & \\ \hline & 1 & 2 \end{array} \end{array} \quad \Phi_{10}((1, 2), \cdot) = \begin{array}{c} \begin{array}{cc|c} 1 & \begin{array}{|c|c|} \hline 1/6 & 1/12 \\ \hline \end{array} & \\ 2 & \begin{array}{|c|c|} \hline 0 & 0 \\ \hline \end{array} & \\ \hline & 1 & 2 \end{array} \end{array}$$

Similar calculations show that both  $\Phi_{01}((1, 2), \cdot)$  and  $\Phi_{00}((1, 2), \cdot)$  consist entirely of zeros. Finally, since  $\bar{Q}((x, y), \cdot) = (\Phi_{11} + \Phi_{10} + \Phi_{01} + \Phi_{00})((x, y), \cdot)$ , we have

$$\bar{Q}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{cc|c} 1 & \begin{array}{|c|c|} \hline 1/6 & 1/3 \\ \hline \end{array} & \\ 2 & \begin{array}{|c|c|} \hline 1/3 & 1/6 \\ \hline \end{array} & \leftarrow y \\ \hline & 1 & 2 \end{array} \end{array}$$

We see that this proposal coupling has the marginal distributions  $Q(1, \cdot) = Q(2, \cdot) = (1/2, 1/2)$ .

## 2.5 Existence of an acceptance indicator coupling

Next, we show that if we have a coupled acceptance mechanism  $\Phi$  relating  $\bar{Q}$  and  $\bar{P}$ , then there exists an acceptance indicator coupling  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ . In the following, we write  $\Delta^{n-1}$  for the set of multinomial distributions on a set with  $n$  elements.

**Lemma 6.** *Suppose  $\bar{P} \in \Gamma(P, P)$ ,  $\bar{Q} \in \Gamma(Q, Q)$ , and  $\Phi$  is a coupled acceptance mechanism relating  $\bar{P}$  and  $\bar{Q}$ . Then there exist  $\bar{Q}((x, y), \cdot)$ -measurable functions  $\phi_{ij}((x, y), \cdot)$  for  $i, j \in \{0, 1\}$  such that  $\phi = (\phi_{11}, \phi_{10}, \phi_{01}, \phi_{00})((x, y), (x', y')) \in \Delta^3$  for  $\bar{Q}((x, y), \cdot)$ -almost all  $(x', y')$ . If we define  $\bar{B}$  such that  $\mathbb{P}(b_x = i, b_y = j \mid x, y, x', y') = \phi_{ij}((x, y), (x', y'))$  when  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , then  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ .*

*Proof.* For each  $i, j \in \{0, 1\}$ , the  $\bar{Q}$  condition in Definition 2 implies  $\Phi_{ij}((x, y), \cdot) \ll \bar{Q}((x, y), \cdot)$ . Thus we can form the Radon–Nikodym derivative  $\phi_{ij}((x, y), \cdot) := d\Phi_{ij}((x, y), \cdot)/d\bar{Q}((x, y), \cdot)$ . The Radon–Nikodym derivative is linear and  $d\bar{Q}((x, y), \cdot)/d\bar{Q}((x, y), \cdot) = 1$ , so for  $\bar{Q}((x, y), \cdot)$ -almost all  $(x', y')$  we have  $\sum_{i,j \in \{0,1\}} \phi_{ij}((x, y), (x', y')) = 1$ . Since  $\phi_{ij} \geq 0$ , this implies that each  $\phi_{ij} \leq 1$ . Thus  $\phi((x, y), (x', y')) \in \Delta^3$  for  $\bar{Q}((x, y), \cdot)$ -almost all  $(x', y')$ .

For points  $(x', y')$  where the above holds, let  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$  be the multinomial random variable on  $\{0, 1\}^2$  with  $\mathbb{P}(b_x = i, b_y = j \mid x, y, x', y') = \phi_{ij}((x, y), (x', y'))$  for  $i, j \in \{0, 1\}$ . Suppose that  $(x', y') \sim \bar{Q}((x, y), \cdot)$  and let  $(X, Y) := (x'b_x + x(1 - b_x), y'b_y + y(1 - b_y))$ . Then for any  $A_x, A_y \in \mathcal{F}$  we have

$$\begin{aligned}\mathbb{P}((X, Y) \in A_x \times A_y, b_x = 1, b_y = 1 \mid x, y) &= \Phi_{11}((x, y), A_x \times A_y) \\ \mathbb{P}((X, Y) \in A_x \times A_y, b_x = 1, b_y = 0 \mid x, y) &= \Phi_{10}((x, y), A_x \times \mathcal{X})1(y \in A_y) \\ \mathbb{P}((X, Y) \in A_x \times A_y, b_x = 0, b_y = 1 \mid x, y) &= \Phi_{01}((x, y), \mathcal{X} \times A_y)1(x \in A_x) \\ \mathbb{P}((X, Y) \in A_x \times A_y, b_x = 0, b_y = 0 \mid x, y) &= \Phi_{00}((x, y), \mathcal{X} \times \mathcal{X})1(x \in A_x)1(y \in A_y).\end{aligned}$$

It follows from these expressions and the definition of a coupled acceptance mechanism  $\Phi$  that  $\mathbb{P}((X, Y) \in A_x \times A_y) = \bar{P}((x, y), A_x \times A_y)$  on all measurable rectangles  $A_x \times A_y$ , and hence that  $\mathbb{P}((X, Y) \in A) = \bar{P}((x, y), A)$  for all  $A \in \mathcal{F} \otimes \mathcal{F}$ . We conclude that  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ .  $\square$

**Lemma 7.** *Let  $\Phi$  be a coupled acceptance mechanism relating  $\bar{P} \in \Gamma(P, P)$  and  $\bar{Q} \in \Gamma(Q, Q)$ , and define  $\phi$  and  $\bar{B}$  as in the proof of Lemma 6. If  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , then*

1.  $\mathbb{P}(b_x = 1 \mid x, y, x') = a(x, x')$  for  $Q(x, \cdot)$ -almost all  $x'$ , and
2.  $\mathbb{P}(b_y = 1 \mid x, y, y') = a(y, y')$  for  $Q(y, \cdot)$ -almost all  $y'$ .

*Proof.* Let  $a_x := \phi_{11} + \phi_{10}$ . Then  $\mathbb{P}(b_x = 1 \mid x, y, x') = \mathbb{E}_{\bar{Q}((x, y), \cdot)}[a_x((x, y), (x', y')) \mid x, y, x']$ , and  $a_x((x, y), \cdot)$  is  $\bar{Q}((x, y), \cdot)$ -measurable. For all  $A \in \mathcal{F}$ , we have

$$\begin{aligned}& \int 1(x' \in A) \mathbb{P}(b_x = 1 \mid x, y, x') \bar{Q}((x, y), (dx', dy')) \\ &= \int 1(x' \in A) a_x((x, y), (x', y')) \bar{Q}((x, y), (dx', dy')) = (\Phi_{11} + \Phi_{10})((x, y), A \times \mathcal{X}) \\ &= Q(x, \{x\} \cap A) + P(x, A \setminus \{x\}) = \int_A a(x, x') Q(x, dx') \\ &= \int 1(x' \in A) a(x, x') \bar{Q}((x, y), (dx', dy')).\end{aligned}$$

The first equality follows from the defining property of conditional expectations. Definition 2, Condition 2 implies  $(\Phi_{11} + \Phi_{10})((x, y), (A \setminus \{x\}) \times \mathcal{X}) = \bar{P}((x, y), (A \setminus \{x\}) \times \mathcal{X})$ , while Condition 3 implies  $(\Phi_{11} + \Phi_{10})((x, y), (A \cap \{x\}) \times \mathcal{X}) = Q(x, A \cap \{x\})$ . These combine to yield the third equality, above.

By the essential uniqueness of the Radon–Nikodym derivative,  $\mathbb{P}(b_x = 1 \mid x, y, x') = a(x, x')$  for all  $x'$  in a measurable set  $\tilde{A} \in \mathcal{F}$  with  $\bar{Q}((x, y), \tilde{A} \times \mathcal{X}) = Q(x, \tilde{A}) = 1$ . Thus this equality holds for  $Q(x, \cdot)$ -almost all  $x' \in \mathcal{X}$ . A similar argument shows that  $\mathbb{P}(b_y = 1 \mid x, y, y') = a(y, y')$  for  $Q(y, \cdot)$ -almost all  $y' \in \mathcal{X}$ .  $\square$

## 2.6 Main result

Having established the lemmas above, we can now prove our main result.

*Proof of Theorem 1.* For the ‘only if’ case, assume  $\bar{Q} \in \Gamma(Q, Q)$  and  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , an acceptance indicator coupling with  $\mathbb{P}(b_x = 1 \mid x, y, x') = a(x, x')$  for  $Q(x, \cdot)$ -almost all  $x'$  and  $\mathbb{P}(b_y = 1 \mid x, y, y') = a(y, y')$  for  $Q(y, \cdot)$ -almost all  $y'$ . Let  $\bar{P}((x, y), \cdot)$  be the law of

$(X, Y) := (b_x x' + (1 - b_x)x, b_y y' + (1 - b_y)y)$ . Then for any  $A \in \mathcal{F}$ ,

$$\begin{aligned} \bar{P}((x, y), A \times \mathcal{X}) &= \mathbb{P}(X \in A | x, y) = \mathbb{P}(x' \in A, b_x = 1 | x, y) + \mathbb{P}(x \in A, b_x = 0 | x, y) \\ &= \int 1(x' \in A) a(x, x') \bar{Q}((x, y), (dx', dy')) + 1(x \in A) \int (1 - a(x, x')) \bar{Q}((x, y), (dx', dy')) \\ &= \int 1(x' \in A) a(x, x') Q(x, dx') + 1(x \in A) \int (1 - a(x, x')) Q(x, dx') = P(x, A). \end{aligned}$$

A similar argument shows that  $\bar{P}((x, y), \mathcal{X} \times A) = P(y, A)$  for any  $A \in \mathcal{F}$ . Thus  $\bar{P} \in \Gamma(P, P)$ .

For the ‘if’ case, take any  $\bar{P} \in \Gamma(P, P)$ . By Lemma 5, there exists a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and a coupled acceptance mechanism  $\bar{\Phi}$  relating  $\bar{Q}$  and  $\bar{P}$ . Then by Lemma 6, there exists an acceptance indicator coupling  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ . Finally by Lemma 7, this  $\bar{B}$  will be such that if  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , then  $\mathbb{P}(b_x = 1 | x, y, x') = a(x, x')$  for  $Q(x, \cdot)$ -almost all  $x'$  and  $\mathbb{P}(b_y = 1 | x, y, y') = a(y, y')$  for  $Q(y, \cdot)$ -almost all  $y'$ . Thus we conclude that  $\bar{P}$  is generated by a proposal coupling  $\bar{Q}$  and an acceptance indicator coupling  $\bar{B}$  with the desired properties.  $\square$

**Example 4.** Returning once more to our running example, we most recently showed that the transition kernel coupling  $\bar{P}$  was related to the proposal coupling  $\bar{Q}$  by a coupled acceptance mechanism  $\bar{\Phi}$  with the following properties:

$$\begin{aligned} \bar{P}((1, 2), \cdot) &= \begin{array}{c} 1 \\ 2 \end{array} \begin{array}{|c|c|} \hline 0 & 1/4 \\ \hline 1/2 & 1/4 \\ \hline \end{array} \begin{array}{c} 1 \\ 2 \end{array} & \bar{Q}((1, 2), \cdot) = \begin{array}{c} 1 \\ 2 \end{array} \begin{array}{|c|c|} \hline 1/6 & 1/3 \\ \hline 1/3 & 1/6 \\ \hline \end{array} \begin{array}{c} 1 \\ 2 \end{array} \\ \Phi_{11}((1, 2), \cdot) &= \begin{array}{c} 1 \\ 2 \end{array} \begin{array}{|c|c|} \hline 0 & 1/4 \\ \hline 1/3 & 1/6 \\ \hline \end{array} \begin{array}{c} 1 \\ 2 \end{array} & \Phi_{10}((1, 2), \cdot) = \begin{array}{c} 1 \\ 2 \end{array} \begin{array}{|c|c|} \hline 1/6 & 1/12 \\ \hline 0 & 0 \\ \hline \end{array} \begin{array}{c} 1 \\ 2 \end{array} \\ \Phi_{01}((1, 2), \cdot) &= \Phi_{00}((1, 2), \cdot) = \begin{array}{c} 1 \\ 2 \end{array} \begin{array}{|c|c|} \hline 0 & 0 \\ \hline 0 & 0 \\ \hline \end{array} \begin{array}{c} 1 \\ 2 \end{array} \end{aligned}$$

Applying the same construction used in the proof of Theorem 1, we find that  $\bar{P}((1, 2), \cdot)$  can be generated from  $\bar{Q}((1, 2), \cdot)$  together with an acceptance indicator coupling  $\bar{B}$  such that if  $(b_x, b_y) \sim \bar{B}((x, y), \cdot)$ , then

$$\mathbb{P}(b_x = b_y = 1 | x, y, \cdot) = \begin{array}{c} 1 \\ 2 \end{array} \begin{array}{|c|c|} \hline 0 & 3/4 \\ \hline 1 & 1 \\ \hline \end{array} \begin{array}{c} 1 \\ 2 \end{array} \quad \mathbb{P}(b_x = 1, b_y = 0 | x, y, \cdot) = \begin{array}{c} 1 \\ 2 \end{array} \begin{array}{|c|c|} \hline 1 & 1/4 \\ \hline 0 & 0 \\ \hline \end{array} \begin{array}{c} 1 \\ 2 \end{array}$$

$$\mathbb{P}(b_x = 0, b_y = 1 | x, y, \cdot) = \mathbb{P}(b_x = b_y = 0 | x, y, \cdot) = \begin{array}{c} 1 \\ 2 \end{array} \begin{array}{|c|c|} \hline 0 & 0 \\ \hline 0 & 0 \\ \hline \end{array} \begin{array}{c} 1 \\ 2 \end{array}$$

We conclude that the given procedure indeed provides a method to find a proposal and acceptance coupling to reproduce the given MH transition coupling at the point  $(x, y) = (1, 2)$ .

### 3 Maximal kernel couplings

In an application of the results above, we now characterize the maximal couplings of MH-like transition kernels. Given such a kernel  $P$ , the identification of couplings  $\bar{P} \in \Gamma(P, P)$  that induce rapid meeting between chains is a challenging but important question for theoretical analysis and especially in applications, where couplings are often required to be Markovian. Maximal couplings  $\bar{P} \in \Gamma^{\max}(P, P)$  represent myopically optimal solutions to the problem of minimizing the meeting time between chains, in that they achieve the largest possible one-step meeting probability  $\mathbb{P}(X = Y | x, y)$  from each state pair  $(x, y)$ . A clear understanding of the structure of  $\Gamma^{\max}(P, P)$  is valuable for designing practical couplings and provides a reference point in the search for more efficient Markovian couplings of MH-like chains.

We write  $\Delta := \{(z, z) : z \in \mathcal{X}\}$  for the diagonal of  $\mathcal{X} \times \mathcal{X}$ ,  $\delta : \mathcal{X} \rightarrow \Delta$  for the map  $z \mapsto (z, z)$ , and  $A_\Delta := \delta(A) = \{(z, z) : z \in A\}$ . We have assumed that  $(\mathcal{X}, \mathcal{F})$  is a Polish space, so  $\Delta \in \mathcal{F} \otimes \mathcal{F}$  and  $\delta$  is a measurable function. As noted above, the coupling inequality states that if  $\mu$  and  $\nu$  are probability measures on  $(\mathcal{X}, \mathcal{F})$  and  $\gamma \in \Gamma(\mu, \nu)$ , then  $\mathbb{P}_{(X, Y) \sim \gamma}(X = Y) \leq 1 - \|\mu - \nu\|_{\text{TV}}$ . A coupling that achieves this upper bound is said to be maximal. In the next section we see that this is equivalent to a measure theoretical condition, which we use in our characterization of maximal couplings of MH-like kernels.

#### 3.1 The Hahn maximality condition

Given probability measures  $\mu$  and  $\nu$  on  $(\mathcal{X}, \mathcal{F})$ , the Hahn-Jordan theorem [e.g. [Dudley, 2002](#), chapter 5.6] states that there exists a measurable set  $S \in \mathcal{F}$  and sub-probability measures  $\mu^r$  and  $\nu^r$  such that  $\mu - \nu = \mu^r - \nu^r$  and  $\mu^r(S^c) = \nu^r(S) = 0$ . The pair  $(S, S^c)$  is called the Hahn decomposition for  $\mu - \nu$ , and it is essentially unique in the sense that if  $R \in \mathcal{F}$  is another set with  $\mu^r(R^c) = \nu^r(R) = 0$ , then  $(\mu - \nu)(S \Delta R) = 0$ . Here  $A \Delta B = (A \setminus B) \cup (B \setminus A)$  denotes the symmetric difference of measurable sets. The pair  $(\mu^r, \nu^r)$  is called the Jordan decomposition of  $\mu - \nu$ , and it is unique.

In the Jordan decomposition of  $\mu - \nu$ ,  $\mu^r$  and  $\nu^r$  are called the upper and lower variation of  $\mu - \nu$ , and  $\mu \wedge \nu := \mu - \mu^r = \nu - \nu^r$  is called the meet or infimum measure of  $\mu$  and  $\nu$ .  $\mu \wedge \nu$  is non-negative and has the defining property that if  $\eta$  is another measure on  $(\mathcal{X}, \mathcal{F})$  with  $\eta(A) \leq \mu(A) \wedge \nu(A)$  for all  $A \in \mathcal{F}$ , then  $\eta(A) \leq (\mu \wedge \nu)(A)$  for all  $A \in \mathcal{F}$ . Note that by the definition of total variation,  $\|\mu - \nu\|_{\text{TV}} = \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)| = \mu^r(\mathcal{X}) = \nu^r(\mathcal{X}) = 1 - (\mu \wedge \nu)(\mathcal{X})$ . See e.g. [Dshalalow \[2012, chap. 5\]](#) or [Aliprantis and Burkinshaw \[1998, sec. 36\]](#) for more on the lattice-theoretic properties of the set of measures on  $(\mathcal{X}, \mathcal{F})$ .

For any measure  $\mu$  on  $(\mathcal{X}, \mathcal{F})$  let  $\delta_\star \mu$  be the pushforward of  $\mu$  by the diagonal map  $\delta$ , so that  $\delta_\star \mu(A) = \mu(\delta^{-1}(A))$  for  $A \in \mathcal{F} \otimes \mathcal{F}$ . This makes  $\delta_\star \mu$  a measure on  $(\mathcal{X} \times \mathcal{X}, \mathcal{F} \otimes \mathcal{F})$ , with  $\delta_\star \mu(A) = \delta_\star \mu(A \cap \Delta)$  for  $A \in \mathcal{F} \otimes \mathcal{F}$ , and  $\delta_\star \mu(B_\Delta) = \mu(B)$  for any  $B \in \mathcal{F}$ . With this notation, we have the following characterization of maximal couplings:

**Lemma 8** ([Douc et al. \[2018\]](#), Theorem 19.1.6). *Let  $\mu - \nu = \mu^r - \nu^r$  be the Jordan decomposition of a pair of probability measures  $\mu$  and  $\nu$  on  $(\mathcal{X}, \mathcal{F})$ . A coupling  $\gamma \in \Gamma(\mu, \nu)$  is maximal if and only if there exists a  $\gamma^r \in \Gamma(\mu^r, \nu^r)$  such that  $\gamma(A) = \gamma^r(A) + \delta_\star(\mu \wedge \nu)(A)$  for all  $A \in \mathcal{F} \otimes \mathcal{F}$ .*

Note that we must have  $\mu^r(\mathcal{X}) = \nu^r(\mathcal{X})$  for  $\Gamma(\mu^r, \nu^r)$  to be nonempty. This follows from the Jordan decomposition, since  $0 = (\mu - \nu)(\mathcal{X}) = (\mu^r - \nu^r)(\mathcal{X})$ . We also have  $\gamma^r(\Delta) = 0$ , since

$$\begin{aligned} 1 - \|\mu - \nu\|_{\text{TV}} &= \gamma(\Delta) = \gamma^r(\Delta) + \delta_\star(\mu \wedge \nu)(\Delta) \\ &= \gamma^r(\Delta) + (\mu \wedge \nu)(\mathcal{X}) = \gamma^r(\Delta) + 1 - \|\mu - \nu\|_{\text{TV}}. \end{aligned}$$

Finally, we observe that Lemma 8 implies the maximal coupling recognition result of Ernst et al. [2019, Lemma 20]. In particular, we have the following characterization of maximal couplings based on the Hahn decomposition:

**Corollary 1** (Hahn Maximality Condition). *Let  $\mu$  and  $\nu$  be measures on  $(\mathcal{X}, \mathcal{F})$ . A coupling  $\gamma \in \Gamma(\mu, \nu)$  is maximal if and only if there is an  $S \in \mathcal{F}$  such that  $\gamma((S^c \times \mathcal{X}) \setminus \Delta) = \gamma((\mathcal{X} \times S) \setminus \Delta) = 0$ . Any  $(S, S^c)$  with this property will be a Hahn decomposition for  $\mu - \nu$ .*

*Proof.* Let  $\mu - \nu = \mu^r - \nu^r$  be a Jordan decomposition, so that for some  $S \in \mathcal{F}$  we have  $\mu^r(S^c) = \nu^r(S) = 0$ . If  $\gamma \in \Gamma(\mu, \nu)$  is maximal, Lemma 8 implies that  $\gamma(A) = \gamma^r(A) + \delta_*(\mu \wedge \nu)(A)$  for all  $A \in \mathcal{F} \otimes \mathcal{F}$ . Thus  $\gamma((S^c \times \mathcal{X}) \setminus \Delta) = \gamma^r(S^c \times \mathcal{X}) = \mu^r(S^c) = 0$ . Similarly,  $\gamma((\mathcal{X} \times S) \setminus \Delta) = 0$ .

For the converse, let  $\gamma \in \Gamma(\mu, \nu)$  and  $\gamma((S^c \times \mathcal{X}) \setminus \Delta) = \gamma((\mathcal{X} \times S) \setminus \Delta) = 0$ . For any  $B \in \mathcal{F}$ ,

$$\begin{aligned}\mu(B) &= \gamma(B \times \mathcal{X}) = \gamma((B \times \mathcal{X}) \setminus \Delta) + \gamma(B_\Delta) \\ \nu(B) &= \gamma(\mathcal{X} \times B) = \gamma((\mathcal{X} \times B) \setminus \Delta) + \gamma(B_\Delta).\end{aligned}$$

By assumption,  $S$  contains the support of  $\gamma((\cdot \times \mathcal{X}) \setminus \Delta)$  and  $S^c$  contains the support of  $\gamma((\mathcal{X} \times \cdot) \setminus \Delta)$ . Thus  $\mu(\cdot) - \nu(\cdot) = \gamma((\cdot \times \mathcal{X}) \setminus \Delta) - \gamma((\mathcal{X} \times \cdot) \setminus \Delta)$  is the Jordan decomposition of  $\mu - \nu$  and  $(S, S^c)$  is a Hahn decomposition. The uniqueness of the Jordan decomposition implies  $\gamma((B \times \mathcal{X}) \setminus \Delta) = \mu^r(B)$  and  $\gamma((\mathcal{X} \times B) \setminus \Delta) = \nu^r(B)$ , which in turn yields  $\gamma^r(\cdot) := \gamma(\cdot \setminus \Delta) \in \Gamma(\mu^r, \nu^r)$ . We also have  $(\mu \wedge \nu)(B) = \mu(B) - \mu^r(B) = \nu(B) - \nu^r(B)$ , so the above implies  $\gamma(B_\Delta) = (\mu \wedge \nu)(B)$  for all  $B \in \mathcal{F}$ . Thus we conclude that for any  $A \in \mathcal{F} \otimes \mathcal{F}$ ,  $\gamma(A) = \gamma(A \setminus \Delta) + \gamma(A \cap \Delta) = \gamma^r(A) + \delta_*(\mu \wedge \nu)(A)$ .  $\square$

We will use the result above to establish conditions in the form of any  $\bar{P} \in \Gamma^{\max}(P, P)$  in terms of proposal and acceptance couplings. First, we consider the relationship of the maximality of transition kernel coupling  $\bar{P}$  and the maximality of a proposal coupling  $\bar{Q}$  that generates it.

### 3.2 Maximal proposal kernel couplings

It may seem that if  $\bar{P} \in \Gamma^{\max}(P, P)$  is generated by a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and some acceptance coupling  $\bar{B}$  then  $\bar{Q}$  might have to be a maximal coupling. The proposal-based maximal coupling of O’Leary et al. [2021] has this property, and it seems plausible that to maximize the probability of  $X = Y$  one might need to start by maximizing the probability of  $x' = y'$ . However, the following shows that no special relationship exists between maximal proposal and transition couplings.

**Lemma 9.** *Suppose the transition kernel coupling  $\bar{P} \in \Gamma^{\max}(P, P)$  is generated by a proposal coupling  $\bar{Q}_m \in \Gamma^{\max}(Q, Q)$  and an acceptance coupling  $\bar{B}_m$ , that  $\bar{Q}_m((x, y), \Delta^c) > 0$  for some  $(x, y)$ , and at that  $(x, y)$ ,  $\mathbb{P}(b_x = b_y = 1 \mid x, y) < 1$  where  $(x'_m, y'_m) \sim \bar{Q}_m((x, y), \cdot)$  and  $(b_x, b_y) \sim \bar{B}_m((x, y), (x'_m, y'_m))$ . Then there exists a non-maximal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and an acceptance indicator coupling  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  also generate  $\bar{P}$ .*

In the following proof, we use  $\bar{Q}_m$  to construct a  $\bar{Q}$  that agrees with  $\bar{Q}_m$  on accepted proposals and independently redraws rejected ones. The hypotheses on the support of  $\bar{Q}_m$  and  $(b_x, b_y)$  are very general but are needed to ensure that this procedure results in a coupling  $\bar{Q}$  that is not maximal.



---

**Algorithm 1** Construction of  $\bar{Q}$  for Lemma 9
 

---

1. Draw  $(x'_m, y'_m) \sim \bar{Q}_m((x, y), \cdot)$  and  $(b_x, b_y) \sim \bar{B}_m((x, y), (x'_m, y'_m))$
  2. For  $z \in \{x, y\}$ :
    - (a) If  $b_z = 1$ , set  $z' = z'_m$ . Else:
    - (b) Draw  $(\tilde{x}, \tilde{y}) \sim \bar{Q}_m((x, y), \cdot)$  and  $(\tilde{b}_x, \tilde{b}_y) \sim \bar{B}_m((x, y), (\tilde{x}, \tilde{y}))$
    - (c) If  $(\tilde{b}_x, \tilde{b}_y) = (b_x, b_y)$ , set  $z' = \tilde{z}$
    - (d) Else: go to 2(b)
  3. Return  $(x', y')$  and  $(b_x, b_y)$
- 

*Proof.* Let  $\bar{Q}((x, y), \cdot)$  be the distribution of the  $(x', y')$  output of Algorithm 1. We claim that  $\bar{Q} \in \Gamma(Q, Q)$ . For  $A \in \mathcal{F}$ ,

$$\begin{aligned}
 & \bar{Q}((x, y), A \times \mathcal{X}) \\
 &= \mathbb{P}(x'_m \in A, b_x = 1 \mid x, y) + \sum_{j \in \{0,1\}} \mathbb{P}(b_x = 0, b_y = j \mid x, y) \mathbb{P}(\tilde{x} \in A \mid \tilde{b}_x = 0, \tilde{b}_y = j, x, y) \\
 &= \mathbb{P}(x'_m \in A, b_x = 1 \mid x, y) + \sum_{j \in \{0,1\}} \mathbb{P}(x'_m \in A, b_x = 0, b_y = j \mid x, y) = \mathbb{P}(x'_m \in A \mid x) = Q(x, A).
 \end{aligned}$$

A similar argument shows  $\bar{Q}((x, y), \mathcal{X} \times A) = Q(y, A)$ . Let  $(x, y)$  be such that  $\bar{Q}_m((x, y), \Delta^c) > 0$ . We have assumed that there is a positive probability of rejecting either  $x'_m$  or  $y'_m$  at  $(x, y)$ . Thus  $\mathbb{P}(x' = y' \mid x, y) < \mathbb{P}(x'_m = y'_m \mid x, y)$ , and so we conclude that  $\bar{Q}$  is not a maximal coupling.

Next, for  $i, j \in \{0, 1\}$  and  $A \in \mathcal{F} \otimes \mathcal{F}$ , define  $\Phi_{ij}((x, y), A) := \mathbb{P}((x', y') \in A, b_x = i, b_y = j)$  using the full output of Algorithm 1. We observe that this is a coupled acceptance mechanism relating  $\bar{Q}$  and  $\bar{P}$ . The first condition of Definition 2 is satisfied by construction. For the second condition, define  $X_m = b_x x'_m + (1 - b_x)x$  and  $Y_m = b_y y'_m + (1 - b_y)y$ . Since  $\bar{Q}_m$  and  $\bar{B}_m$  generate  $\bar{P}$ , we must have  $(X_m, Y_m) \sim \bar{P}((x, y), \cdot)$ . Thus for any  $A_x \times A_y \in \mathcal{F} \otimes \mathcal{F}$ ,

$$\begin{aligned}
 & \Phi_{11}((x, y), A_x \times A_y) + \Phi_{10}((x, y), A_x \times \mathcal{X})1(y \in A_y) \\
 & \quad + \Phi_{01}((x, y), \mathcal{X} \times A_y)1(x \in A_x) + \Phi_{00}((x, y), \mathcal{X} \times \mathcal{X})1(x \in A_x)1(y \in A_y) \\
 &= \mathbb{P}(X_0 \in A_x, Y_0 \in A_y, b_x = 1, b_y = 1 \mid x, y) + \mathbb{P}(X_0 \in A_x, Y_0 = y, b_x = 1, b_y = 0 \mid x, y) \\
 & \quad + \mathbb{P}(X_0 = x, Y_0 \in A_y, b_x = 0, b_y = 1 \mid x, y) + \mathbb{P}(X_0 = x, Y_0 = y, b_x = 0, b_y = 0 \mid x, y) \\
 &= \bar{P}((x, y), A_x \times A_y).
 \end{aligned}$$

The third condition on  $\Phi$  follows from the fact that  $b_x = 1$  if  $x'_m = x$  and  $b_y = 1$  if  $y'_m = y$ . Since  $\Phi$  is a coupled acceptance mechanism relating  $\bar{Q}$  and  $\bar{P}$ , Lemma 6 ensures the existence of a  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ .  $\square$

The result above shows that if a maximal transition coupling  $\bar{P} \in \Gamma^{\max}(P, P)$  is generated by a proposal coupling  $\bar{Q}$  and an acceptance coupling  $\bar{B}$ , then  $\bar{Q}$  does not have to be maximal. The following example shows that some maximal couplings  $\bar{P}$  cannot be generated from any maximal coupling  $\bar{Q} \in \Gamma^{\max}(Q, Q)$ .

**Example 5.** Assume  $\mathcal{X} = \{1, 2, 3\}$ ,  $\mathcal{F} = 2^{\mathcal{X}}$ , and  $(x, y) = (1, 2)$ . From these states, we assume the following proposal and transition kernel distributions:

$$\begin{aligned}
 Q(x = 1, \cdot) &= (0, 1/2, 1/2) & P(x = 1, \cdot) &= (1/2, 1/2, 0) \\
 Q(y = 2, \cdot) &= (1/2, 0, 1/2) & P(y = 2, \cdot) &= (1/2, 0, 1/2).
 \end{aligned}$$

It is straightforward to verify that these distributions correspond to an MH transition kernel obtained when  $Q(3, \cdot) = (0, 1, 0)$  and  $\pi = (2/5, 2/5, 1/5)$ . Simple algebra shows that any coupling in  $\Gamma^{\max}(Q, Q)$  and any coupling in  $\Gamma^{\max}(P, P)$  must have the following properties:

$$\bar{Q}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{ccc|ccc} 1 & 0 & 1/2 & 0 & & \\ 2 & 0 & 0 & 0 & & \\ 3 & 0 & 0 & 1/2 & & \\ \hline & 1 & 2 & 3 & & \end{array} \leftarrow y \end{array} \quad \bar{P}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{ccc|ccc} 1 & 1/2 & 0 & 0 & & \\ 2 & 0 & 0 & 0 & & \\ 3 & 0 & 1/2 & 0 & & \\ \hline & 1 & 2 & 3 & & \end{array} \leftarrow y \end{array}$$

There exists no acceptance indicator coupling  $\bar{B}$  such that it and the unique maximal  $\bar{Q}$  given above generate  $\bar{P}$ , since  $(x', y') \sim \bar{Q}((1, 2), \cdot)$ ,  $(b_x, b_y) \sim \bar{B}((1, 2), (x', y'))$ ,  $X = b_x x' + (1 - b_x)x$ ,  $Y = b_y y' + (1 - b_y)y$ , and  $(X, Y) \sim \bar{P}((1, 2), \cdot)$  yield a contradiction:

$$1/2 = \mathbb{P}(X = 2, Y = 3 \mid x = 1, y = 2) \leq \mathbb{P}(x' = 2, y' = 3 \mid x = 1, y = 2) = 0.$$

Note that in line with Theorem 1, we can still generate  $\bar{P}((1, 2), \cdot)$  using the following non-maximal proposal coupling and the coupled acceptance indicators  $b_x, b_y$ :

$$\tilde{Q}((1, 2), \cdot) = \begin{array}{c} x \\ \downarrow \\ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1/2 & & \\ 2 & 0 & 0 & 0 & & \\ 3 & 0 & 1/2 & 0 & & \\ \hline & 1 & 2 & 3 & & \end{array} \leftarrow y \end{array}$$

$$\begin{aligned} \mathbb{P}(b_x = 1, b_y = 1 \mid (x, y) = (1, 2), (x', y') = (2, 3)) &= 1 \\ \mathbb{P}(b_x = 0, b_y = 1 \mid (x, y) = (1, 2), (x', y') = (3, 1)) &= 1. \end{aligned}$$

One upshot of Lemma 9 and Example 5 is that a coupling  $\bar{P} \in \Gamma^{\max}(P, P)$  requires a certain amount of proposal probability on the diagonal, but the maximality of  $\bar{Q}$  is neither necessary nor sufficient for  $\bar{Q}$  to be able to generate  $\bar{P}$ .

### 3.3 Characterization of maximal kernel couplings

Next we turn to the main result of this section, which extends Theorem 1 to characterize the maximal couplings of an MH-like transition kernel in terms of proposal and acceptance indicator couplings. For each  $x, y \in \mathcal{X}$ , let  $(S_{xy}, S_{xy}^c)$  be any Hahn decomposition for  $P(x, \cdot) - P(y, \cdot)$ . Thus  $S_{xy} \in \mathcal{F}$  and  $P(x, A) \geq P(y, A)$  for any  $A \in \mathcal{F}$  with  $A \subset S_{xy}$ . Note that if  $P(x, \cdot)$  and  $P(y, \cdot)$  have Radon–Nikodym derivatives  $p(x, \cdot)$  and  $p(y, \cdot)$  with respect to a common dominating measure, then we can use  $S_{xy} = \{z : p(x, z) \geq p(y, z)\}$  for these sets.

**Theorem 2.** *Let  $P$  be the MH-like transition kernel on  $(\mathcal{X}, \mathcal{F})$  generated by a proposal kernel  $Q$  and an acceptance rate function  $a$ . Then  $\bar{P} \in \Gamma^{\max}(P, P)$  if and only if  $\bar{P}$  is generated by  $\bar{Q} \in \Gamma(Q, Q)$  and an acceptance indicator coupling  $\bar{B}$  with the following properties: if  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , then for all  $x, y \in \mathcal{X}$ :*

1.  $\mathbb{P}(b_x = 1 \mid x, y, x') = a(x, x')$  for  $Q(x, \cdot)$ -almost all  $x'$
2.  $\mathbb{P}(b_y = 1 \mid x, y, y') = a(y, y')$  for  $Q(y, \cdot)$ -almost all  $y'$

and for  $\bar{Q}((x, y), \cdot)$ -almost all  $(x', y')$ ,




Figure 2: Diagram of acceptance scenarios considered in Theorem 2. The support of a maximal coupling  $\bar{P}((x, y), \cdot)$  is contained in the union of  $S_{xy} \times S_{xy}^c$  (gray box) and  $\Delta$  (the diagonal). Arrows illustrate the relationship of proposals  $(x', y')$  to transitions  $(X, Y)$  under different accept/reject combinations, with transitions outside the support of  $\bar{P}((x, y), \cdot)$  forbidden almost surely. Case 1: the maximality of  $\bar{P}$  does not constrain the acceptance pattern of proposals  $(x', y') \in S_{xy} \times S_{xy}^c$ . Case 2: proposals in  $\Delta$  can be fully accepted ( $b_x = b_y = 1$ ) or fully rejected ( $b_x = b_y = 0$ ), but  $y'$  must be accepted if  $x' \in S_{xy}^c$ , and  $x'$  must be accepted if  $y' \in S_{xy}$ . Case 3: proposals in  $(S_{xy} \times S_{xy}^c)^c \cap \Delta^c$  must be fully rejected unless  $y' = x$  or  $x' = y$ . Case 4: a proposal  $(x', y')$  outside the support of  $\bar{P}((x, y), \cdot)$  may be partially accepted ( $b_x \neq b_y$ ) if it yields a transition to  $(x, x)$  or  $(y, y)$ .

3.  $\mathbb{P}(b_x = b_y = 1 \mid x, y, x', y') = 0$  if  $x' \neq y'$  and either  $x' \in S_{xy}^c$  or  $y' \in S_{xy}$
4.  $\mathbb{P}(b_x = 1, b_y = 0 \mid x, y, x', y') = 0$  if  $x' \neq y$  and either  $x' \in S_{xy}^c$  or  $y \in S_{xy}$
5.  $\mathbb{P}(b_x = 0, b_y = 1 \mid x, y, x', y') = 0$  if  $y' \neq x$  and either  $x \in S_{xy}^c$  or  $y' \in S_{xy}$
6.  $\mathbb{P}(b_x = 0, b_y = 0 \mid x, y, x', y') = 0$  if  $x \neq y$  and either  $x \in S_{xy}^c$  or  $y \in S_{xy}$ .

Recall that by Corollary 1, the maximality of a coupling  $\bar{P} \in \Gamma(P, P)$  is equivalent to a condition on the support of each  $\bar{P}((x, y), \cdot)$ . Conditions 3-6 relate these support constraints to the behavior of a proposal coupling  $\bar{Q}$  and an acceptance indicator coupling  $\bar{B}$ . See Figure 2 for an illustration of the acceptance scenarios considered in these conditions and a visual intuition for why certain ones must be ruled out for  $\bar{Q}$  and  $\bar{B}$  to generate a maximal  $\bar{P}$ .

*Proof of Theorem 2.* Suppose  $\bar{P} \in \Gamma^{\max}(P, P)$ . By Theorem 1, there exists  $\bar{Q} \in \Gamma(Q, Q)$  and an acceptance indicator coupling  $\bar{B}$  such that for any  $x, y \in \mathcal{X}$  and  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , we have

1.  $\mathbb{P}(b_x = 1 \mid x, y, x') = a(x, x')$  for  $Q(x, \cdot)$ -almost all  $x'$ , and
2.  $\mathbb{P}(b_y = 1 \mid x, y, y') = a(y, y')$  for  $Q(y, \cdot)$ -almost all  $y'$ .

Thus Conditions 1 and 2 directly follow from Theorem 1.

Since  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ , we have  $(X, Y) \sim \bar{P}((x, y), \cdot)$  where  $X = b_x x' + (1 - b_x)x$  and  $Y = b_y y' + (1 - b_y)y$ . Since  $\bar{P}$  is maximal, Corollary 1 implies

$$0 = \bar{P}((x, y), (S_{xy}^c \times \mathcal{X}) \setminus \Delta) = \mathbb{P}(X \in S_{xy}^c, X \neq Y \mid x, y).$$

Breaking this up into the four possible acceptance scenarios  $(b_x, b_y) = (1, 1), (1, 0), (0, 1),$  and  $(0, 0)$  yields

$$\begin{aligned} 0 &= \mathbb{E}[1(x' \neq y', x' \in S_{xy}^c) \mathbb{P}(b_x = b_y = 1 \mid x, y, x', y')] \\ 0 &= \mathbb{E}[1(x' \neq y, x' \in S_{xy}^c) \mathbb{P}(b_x = 1, b_y = 0 \mid x, y, x', y')] \\ 0 &= \mathbb{E}[1(y' \neq x, x \in S_{xy}^c) \mathbb{P}(b_x = 0, b_y = 1 \mid x, y, x', y')] \\ 0 &= \mathbb{E}[1(x \in S_{xy}^c \setminus \{y\}) \mathbb{P}(b_x = b_y = 0 \mid x, y, x', y')]. \end{aligned}$$

In turn, these equations imply that for  $\bar{Q}((x, y), \cdot)$ -almost all  $(x', y')$ ,

$$\begin{aligned} \mathbb{P}(b_x = b_y = 1 \mid x, y, x', y') &= 0 \text{ if } x' \neq y' \text{ and } x' \in S_{xy}^c \\ \mathbb{P}(b_x = 1, b_y = 0 \mid x, y, x', y') &= 0 \text{ if } x' \neq y \text{ and } x' \in S_{xy}^c \\ \mathbb{P}(b_x = 0, b_y = 1 \mid x, y, x', y') &= 0 \text{ if } y' \neq x \text{ and } x \in S_{xy}^c \\ \mathbb{P}(b_x = b_y = 0 \mid x, y, x', y') &= 0 \text{ if } x \neq y \text{ and } x \in S_{xy}^c. \end{aligned}$$

This shows that the first either/or case of each of Conditions 3-6 are satisfied. Since  $\bar{P}$  is maximal, Corollary 1 also implies  $0 = \bar{P}((x, y), (\mathcal{X} \times S_{xy}) \setminus \Delta)$ . Proceeding as above shows that the second either/or cases are also satisfied. So we conclude that if  $\bar{P} \in \Gamma^{\max}(P, P)$ , then  $\bar{P}$  is generated by a  $\bar{Q}$  and  $\bar{B}$  satisfying the six conditions stated above.

For the converse, suppose that  $\bar{Q} \in \Gamma(Q, Q)$  and a proposal coupling  $\bar{B}$  generate  $\bar{P}$  and satisfy the given hypotheses. Since Conditions 1 and 2 are equivalent to the conditions of Theorem 1, we have  $\bar{P} \in \Gamma(P, P)$ . Now let  $(x', y') \sim \bar{Q}((x, y), \cdot)$ ,  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ ,  $X = b_x x' + (1 - b_x)x$ , and  $Y = b_y y' + (1 - b_y)y$ .  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ , so

$$\begin{aligned} \bar{P}((x, y), (S_{xy}^c \times \mathcal{X}) \setminus \Delta) &= \mathbb{P}(X \in S_{xy}^c, X \neq Y \mid x, y) \\ &= \mathbb{P}(x' \in A, x' \neq y', b_x = 1, b_y = 1 \mid x, y) + \mathbb{P}(x' \in A, x' \neq y, b_x = 1, b_y = 0 \mid x, y) \\ &\quad + \mathbb{P}(y' \neq x, b_x = 0, b_y = 1 \mid x, y)1(x \in A) + \mathbb{P}(b_x = 0, b_y = 0 \mid x, y)1(x \in A)1(x \neq y) = 0. \end{aligned}$$

The last equality follows directly from Conditions 3-6 of the Theorem, with Condition 3 ensuring that the first term equals zero, Condition 4 ensuring that the second term equals zero, and so on. A similar argument yields  $\bar{P}((x, y), (\mathcal{X} \times S_{xy}) \setminus \Delta) = 0$ . By Corollary 1,  $\bar{P}$  is maximal if and only if there is a measurable set  $S \in \mathcal{F}$  such that  $\bar{P}((x, y), (S^c \times \mathcal{X}) \setminus \Delta) = \bar{P}((x, y), (\mathcal{X} \times S) \setminus \Delta) = 0$ . The argument above shows that  $S_{xy}$  has these properties, so we conclude that  $\bar{P} \in \Gamma^{\max}(P, P)$ .  $\square$

Now we conclude this section. Theorem 2 completely characterizes the maximal couplings of MH-like transition kernels. We show every maximal kernel coupling can be represented by a proposal coupling and an acceptance coupling that satisfies additional constraints on the Hahn decomposition. Meanwhile, perhaps surprisingly, we show in Example 5 that there is no special relationship between maximal proposal couplings and maximal kernel couplings.

## 4 Examples

In Section 2 we showed that any coupling of MH-like transition kernels can be generated by a proposal coupling followed by an acceptance indicator coupling. Here we apply this result to some popular algorithms. In Section 4.1 we give a more explicit form of Theorem 1 that holds under mild assumptions on the proposal distribution  $Q$ . Sections 4.2 and 4.3 concern the random

walk Metropolis algorithm, the Metropolis-adjusted Langevin algorithm, and the Metropolis–Hastings algorithm on a discrete state space. These examples illustrate the range of behavior possible for simple, gradient-based, and discrete forms of the MH algorithm. We note that our methods also apply to algorithms like Hamiltonian Monte Carlo and Metropolis-within-Gibbs and yield qualitatively similar conclusions.

Finally, in Section 4.4 we consider the two-step representation of an MH transition kernel satisfying minorization conditions. Such conditions play an important role in the ‘drift and minorization’ method of bounding convergence to stationarity, as in Rosenthal [1995] and Jones and Hobert [2001]. The splitting argument at the heart of this approach is usually described in terms of the overall transition kernel, but here we show that it takes a natural form in terms of proposal and acceptance indicator couplings.

## 4.1 Simplified characterization of couplings

To prove the ‘only if’ part of Theorem 1 we constructed a map from an arbitrary coupling  $\bar{P} \in \Gamma(P, P)$  to a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and an acceptance indicator coupling  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ . Lemmas 3–7 suggest a complicated relationship between  $\bar{P}$ ,  $\bar{Q}$ , and  $\bar{B}$ , at least when  $P$  is allowed to be any MH-like transition kernel. Corollary 2 shows that this relationship takes a more intuitive form when  $Q(x, \cdot)$  is absolutely continuous with respect to the base measure and when  $Q(x, \{x\}) = 0$  for all  $x \in \mathcal{X}$ . These assumptions often hold, e.g. on a continuous state space when  $\lambda$  is non-atomic, on a discrete state space when  $Q$  represents a (non-lazy) random walk, and in many other cases.

We need to define a few new objects before stating the main result of this section. First, for all  $A_x \in \mathcal{F}$  let  $\bar{P}_y(x, A_x) := \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\})$ , so that  $\bar{P}_y(x, \cdot)$  is a sub-probability on  $(\mathcal{X}, \mathcal{F})$ . For all  $x, y \in \mathcal{X}$ , the measure  $\bar{P}_y(x, \cdot) \ll \lambda$  if  $Q(x, \cdot) \ll \lambda$  since

$$\begin{aligned} \bar{P}_y(x, A_x) &= \bar{P}((x, y), (A_x \setminus \{x\}) \times \{y\}) \leq \bar{P}((x, y), (A_x \setminus \{x\}) \times \mathcal{X}) \\ &= P(x, A_x \setminus \{x\}) = \int_{A_x \setminus \{x\}} a(x, x') q(x, x') \lambda(dx'). \end{aligned}$$

Here  $q(x, \cdot) = dQ(x, \cdot)/d\lambda$ . In this case  $\bar{P}_y(x, \cdot)$  has density  $\bar{p}_y(x, \cdot)$  with respect to  $\lambda$  by the Radon–Nikodym theorem. A similar argument shows that  $\bar{P}_x(y, A_y) := \bar{P}((x, y), \{x\} \times (A_y \setminus \{y\}))$  has density  $\bar{p}_x(y, \cdot)$  with respect to  $\lambda$ .

Let  $r(x) := P(x, \{x\})$ , the probability of a transition from  $x$  to itself under  $P$ . From the definitions in Section 2.4, we have  $\beta(x) = 1(r(x) = 0)$  and

$$\mu(x, A_x) := \begin{cases} \frac{Q(x, A_x) - P(x, A_x \setminus \{x\})}{r(x)} & \text{if } r(x) > 0 \\ 1(x \in A_x) & \text{if } r(x) = 0. \end{cases}$$

It follows that  $\mu(x, A_x)(1 - \beta(x)) = (Q(x, A_x) - P(x, A_x \setminus \{x\}))/r(x)$  if  $r(x) > 0$  and 0 otherwise. Thus  $\mu(x, \cdot)(1 - \beta(x))$  has density  $m(x, x') := q(x, x')(1 - a(x, x'))/r(x)$  when  $r(x) > 0$ , and 0 otherwise.

Next, we separate  $\bar{P}$  into one part that is absolutely continuous with respect to  $\lambda \times \lambda$  and another part that is singular to it. By the Lebesgue decomposition theorem [e.g. Dudley, 2002, chapter 5.5] for each  $x, y \in \mathcal{X}$  we have  $\bar{P}((x, y), \cdot) = \bar{P}_{\ll}((x, y), \cdot) + \bar{P}_{\perp}((x, y), \cdot)$  with  $\bar{P}_{\ll}((x, y), \cdot) \ll \lambda \times \lambda$  and  $\bar{P}_{\perp}((x, y), \cdot) \perp \lambda \times \lambda$ . We then define  $p((x, y), \cdot)$  to be the density of  $\bar{P}_{\ll}((x, y), \cdot)$  with respect to  $\lambda \times \lambda$ . Finally let  $\bar{r}(x, y) := \bar{P}((x, y), \{x\} \times \{y\})$ , the probability of a transition from  $(x, y)$  to itself under  $\bar{P}$ . With these definitions, we can now give an explicit characterization of the couplings  $\bar{P} \in \Gamma(P, P)$  under mild conditions on the proposal distribution.

**Corollary 2.** Let  $P$  be the MH-like transition kernel on  $(\mathcal{X}, \mathcal{F})$  generated by a proposal kernel  $Q$  and an acceptance rate function  $a$ , and suppose that  $Q(x, \cdot) \ll \lambda$  with  $Q(x, \{x\}) = 0$  for all  $x \in \mathcal{X}$ . If a joint kernel  $\bar{P} \in \Gamma(P, P)$ , then it is generated by  $\bar{Q} = \bar{Q}_{\ll} + \bar{Q}_{\perp} \in \Gamma(Q, Q)$  and an acceptance indicator coupling  $\bar{B}$  with the following properties for all  $x, y \in \mathcal{X}$  and  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ :

1.  $\bar{Q}_{\perp}((x, y), A) = \bar{P}_{\perp}((x, y), A \cap (\{x\}^c \times \{y\}^c))$  for  $A \in \mathcal{F}$ ,
2.  $\bar{Q}_{\ll}((x, y), \cdot)$  has the following density with respect to  $\lambda \times \lambda$ :

$$\begin{aligned} q((x, y), (x', y')) \\ = p((x, y), (x', y')) + \bar{p}_y(x, x')m(y, y') + m(x, x')\bar{p}_x(y, y') + m(x, x')m(y, y')\bar{r}(x, y), \end{aligned}$$

3.  $b_x = b_y = 1$  almost surely when  $(x', y')$  is in the support of  $\bar{Q}_{\perp}((x, y), \cdot)$ ,

and for  $\bar{Q}_{\ll}((x, y), \cdot)$ -almost all  $(x', y')$ ,

4.  $\mathbb{P}(b_x = 1 | x, y, x', y') q((x, y), (x', y')) = p((x, y), (x', y')) + \bar{p}_y(x, x')m(y, y')$ ,
5.  $\mathbb{P}(b_y = 1 | x, y, x', y') q((x, y), (x', y')) = p((x, y), (x', y')) + \bar{p}_x(y, y')m(x, x')$ .

*Proof.* Let  $B_{xy} := \{x\}^c \times \{y\}^c$ , and note that for all  $x, y \in \mathcal{X}$  and  $\bar{Q} \in \Gamma(Q, Q)$ , we have  $\bar{Q}((x, y), B_{xy}) = 1$ , since  $B_{xy}^c = (\{x\} \times \mathcal{X}) \cup (\mathcal{X} \times \{y\})$ ,  $\bar{Q}((x, y), \{x\} \times \mathcal{X}) = Q(x, \{x\}) = 0$ , and  $\bar{Q}((x, y), \mathcal{X} \times \{y\}) = Q(y, \{y\}) = 0$ . Thus we may restrict attention to the behavior of  $\bar{Q}((x, y), \cdot)$  on  $B_{xy}$ .

When  $\bar{P} \in \Gamma(P, P)$ , Lemma 5 tells us that there exists a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  and a coupled acceptance mechanism  $\Phi$  relating  $\bar{P}$  and  $\bar{Q}$ . In particular, for  $x, y \in \mathcal{X}$  and  $A_x, A_y \in \mathcal{F}$  with  $x \notin A_x$  and  $y \notin A_y$ , the proof of that Lemma shows that we can use the following measures:

$$\begin{aligned} \bar{Q}((x, y), A_x \times A_y) &= (\Phi_{11} + \Phi_{10} + \Phi_{01} + \Phi_{00})((x, y), A_x \times A_y) \\ \Phi_{11}((x, y), A_x \times A_y) &= \bar{P}((x, y), A_x \times A_y) \\ \Phi_{10}((x, y), A_x \times A_y) &= \bar{P}((x, y), A_x \times \{y\})(1 - \beta(y))\mu(y, A_y) \\ \Phi_{01}((x, y), A_x \times A_y) &= (1 - \beta(x))\mu(x, A_x)\bar{P}((x, y), \{x\} \times A_y) \\ \Phi_{00}((x, y), A_x \times A_y) &= (1 - \beta(x))\mu(x, A_x)(1 - \beta(y))\mu(y, A_y)\bar{P}((x, y), \{x\} \times \{y\}). \end{aligned}$$

From the Lebesgue decomposition of  $\bar{P}$ , we have

$$\bar{P}((x, y), A_x \times A_y) = \bar{P}_{\perp}((x, y), A_x \times A_y) + \int_{A_x \times A_y} p((x, y), (x', y'))d(x', y').$$

The definitions above Corollary 2 also imply that  $\Phi_{10}((x, y), \cdot)$  has density  $p_y(x, x')m(y, y')$ ,  $\Phi_{01}((x, y), \cdot)$  has density  $p_x(y, y')m(x, x')$ , and  $\Phi_{00}((x, y), \cdot)$  has density  $\bar{r}(x, y)m(x, x')m(y, y')$ . Thus  $\bar{Q}((x, y), \cdot)$  is the sum of a singular part  $\bar{Q}_{\perp}((x, y), A) = \bar{P}((x, y), A \cap B_{xy})$  and an absolutely continuous part  $\bar{Q}_{\ll}((x, y), \cdot)$  with the density function specified in condition 2, above.

Next by Lemma 6, we know that there exists an acceptance indicator coupling  $\bar{B}$  such that  $\bar{Q}$  and  $\bar{B}$  generate  $\bar{P}$ . This  $\bar{B}$  has the property that  $\bar{Q}((x, y), \cdot)$ -almost all values of  $(x', y')$ , if  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$ , then

$$\begin{aligned} \mathbb{P}(b_x = 1 | x, y, x', y') &= d(\Phi_{11} + \Phi_{10})((x, y), \cdot) / d\bar{Q}((x, y), \cdot) \\ \mathbb{P}(b_y = 1 | x, y, x', y') &= d(\Phi_{11} + \Phi_{01})((x, y), \cdot) / d\bar{Q}((x, y), \cdot). \end{aligned}$$

For  $A \in \mathcal{F}$  contained in the support of  $\bar{Q}_\perp((x, y), \cdot)$ , we have  $\Phi_{11}((x, y), A) = \bar{Q}((x, y), A)$  and  $\Phi_{10}((x, y), A) = \Phi_{01}((x, y), A) = 0$ , so the Radon-Nikodym derivatives above equal 1 almost surely. This proves condition 3. Otherwise, for  $\bar{Q}((x, y), \cdot)$ -almost all points  $(x', y')$  with  $q((x, y), (x', y')) > 0$  we have

$$\mathbb{P}(b_x = 1 | x, y, x', y') = \frac{d(\Phi_{11} + \Phi_{10})((x, y), \cdot)/d(\lambda \times \lambda)}{d\bar{Q}((x, y), \cdot)/d(\lambda \times \lambda)} = \frac{p((x, y), (x', y')) + \bar{p}_y(x, x')m(y, y')}{q((x, y), (x', y'))}$$

and similarly for  $\mathbb{P}(b_y | x, y, x', y')$ . Finally, we have  $p((x, y), (x', y')) = \bar{p}_y(x, x') = \bar{p}_x(y, y') = 0$  whenever  $q((x, y), (x', y')) = 0$ , thus proving conditions 4 and 5.  $\square$

Corollary 2 applies to MH-like algorithms on both continuous and discrete spaces, as long as  $Q(x, \cdot)$  has a density or mass function and as long as the proposal  $x'$  almost surely differs from the current point  $x$ . It provides an explicit way to represent any kernel coupling in terms of proposal and acceptance indicator couplings.

## 4.2 Metropolis–Hastings algorithms on a continuous state space

We now turn to the implications of our characterization results for some widely-used algorithms. To begin, let  $\mathcal{X} = \mathbb{R}^n$  equipped with Lebesgue measure  $\lambda$ , and let  $Q$  be a proposal kernel on  $\mathcal{X}$  with density  $q$  such that  $q(x, y) = g(\|x - y\|)$  for some  $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  and any  $x, y \in \mathcal{X}$ . Let  $\pi$  be a target distribution with density  $\pi(\cdot)$  and let  $a$  be the associated MH acceptance rate function. Given these hypotheses on  $Q$ , we refer to the MH transition kernel  $P$  generated by  $Q$  and  $a$  as a Random Walk Metropolis (RWM) kernel.

Corollary 2 implies the existence of a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  which can be written as  $\bar{Q} = \bar{Q}_{\ll} + \bar{Q}_\perp$ . Here  $\bar{Q}_\perp((x, y), A) = \bar{P}_\perp((x, y), A \cap (\{x\}^c \times \{y\}^c))$  for  $A \in \mathcal{F}$  is the measure  $\bar{P}$  after removing the mass at its current state  $(x, y)$ , and  $\bar{Q}_{\ll}((x, y), \cdot)$  has density

$$\begin{aligned} q((x, y), (x', y')) \\ = p((x, y), (x', y')) + \bar{p}_y(x, x')m(y, y') + m(x, x')\bar{p}_x(y, y') + m(x, x')m(y, y')\bar{r}(x, y), \end{aligned}$$

with respect to  $\lambda \times \lambda$ , where all the notations are defined at the beginning of Section 4.1. For RWM algorithms, the function  $m$  takes the following form for every  $x, x'$ :

$$m(x, x') := \frac{q(x, x')(1 - a(x, x'))}{r(x)} = \frac{q(x, x')(1 - \frac{\pi(x')}{\pi(x)})}{1 - \int_{\mathcal{X}} q(x, y)(\frac{\pi(y)}{\pi(x)} \wedge 1)\lambda(dy)}.$$

The Metropolis-adjusted Langevin algorithm (MALA) [Roberts and Tweedie, 1996] uses a discretized approximation to the Langevin diffusion  $dX_t = \frac{1}{2}\nabla \log \pi(X_t)dt + dW_t$ , where  $W_t$  represents a Brownian motion. After each update the MH acceptance-rejection mechanism is applied to determine whether to accept the proposed move. Thus MALA may be viewed as a case of the MH algorithm using a proposal distribution  $Q(x, \cdot) \sim N(x + \tau\nabla \log \pi(x), 2\tau\mathbb{I})$ . This distribution has density  $q(x, y) \propto \exp(-\frac{1}{4\tau}\|y - x - \tau\nabla \log \pi(x)\|_2^2)$ , where  $\tau$  is a tuning parameter. Since MALA makes use of gradient information, it typically outperforms RWM in high dimensions [Roberts and Rosenthal, 1998]. For MALA algorithms, the measure  $\mu(x, \cdot)$  has a slightly different integral representation relative to RWM:

$$\mu(x, A) := \begin{cases} \frac{\int_A q(x, y)(1 - (\frac{\pi(y)q(y, x)}{\pi(x)q(x, y)} \wedge 1))\lambda(dy)}{r(x)} & \text{if } r(x) > 0 \\ 1(x \in A) & \text{otherwise.} \end{cases}$$

Here  $r(x) = 1 - \int_{\mathcal{X}} q(x, y) \left( \frac{\pi(y)q(y, x)}{\pi(x)q(x, y)} \wedge 1 \right) \lambda(dy)$  is the rejection probability at  $x$ .

Fix  $x, y$  with  $r(x), r(y) > 0$  and let  $p(x, y) = q(x, y) \left( \frac{\pi(y)q(y, x)}{\pi(x)q(x, y)} \wedge 1 \right)$  be the transition density from  $x$  to  $y$ . Then every kernel coupling  $\bar{P}$  of the MALA algorithm is very similar to Example 4.2. Again by Corollary 2, there exists a proposal coupling  $\bar{Q} \in \Gamma(Q, Q)$  which can be written as  $\bar{Q} = \bar{Q}_{\ll} + \bar{Q}_{\perp}$ . Here  $\bar{Q}_{\perp}((x, y), A) = \bar{P}_{\perp}((x, y), A \cap (\{x\}^c \times \{y\}^c))$  for  $A \in \mathcal{F}$  is the measure  $\bar{P}$  after removing the mass at its current state  $(x, y)$ , and  $\bar{Q}_{\ll}((x, y), \cdot)$  has density

$$\begin{aligned} q((x, y), (x', y')) \\ = p((x, y), (x', y')) + \bar{p}_y(x, x')m(y, y') + m(x, x')\bar{p}_x(y, y') + m(x, x')m(y, y')\bar{r}(x, y), \end{aligned}$$

with respect to  $\lambda \times \lambda$ , where all the notations are defined at the beginning of Section 4.1. For MALA algorithms, the function  $m$  takes the following slightly different form from the RWH algorithms:

$$m(x, x') := \frac{q(x, x')(1 - a(x, x'))}{r(x)} = \frac{q(x, x')(1 - \frac{\pi(x')q(x', x)}{\pi(x)q(x, x')})}{1 - \int_{\mathcal{X}} q(x, y) \left( \frac{\pi(y)q(y, x)}{\pi(x)q(x, y)} \wedge 1 \right) \lambda(dy)}.$$

Comparing with Example 4.2, we find that the two-step representations for both algorithms take the same general form, while the details (such as  $p(\cdot, \cdot)$ ,  $r(\cdot)$ ,  $q(\cdot, \cdot)$ , and  $m(\cdot, \cdot)$ ) differ by algorithm.

### 4.3 The Metropolis–Hastings algorithm on a finite state space

Consider an arbitrary Metropolis–Hastings algorithm on the finite state space  $\mathcal{X} = \{x_1, \dots, x_n\}$  with target distribution  $\pi$  and proposal distribution  $Q$  with the property that  $Q(x, \{x\}) = 0$  for all  $x \in \mathcal{X}$ . Let  $\bar{P}$  be a transition kernel coupling. For each fixed  $x, y \in \mathcal{X}$ ,  $\bar{P}((x, y), \cdot)$  can be represented by a  $n \times n$  matrix  $M_{\bar{P}}$  where the  $(i, j)$ -th entry corresponds to the probability of moving  $(x, y)$  to  $(x_i, x_j)$ . Let  $(r_1, \dots, r_n), (c_1, \dots, c_n)$  be the row and column sums of  $M_{\bar{P}}$ . The definition of a coupling guarantees that  $r_i = P(x, x_i) = Q(x, x_i) \left( \frac{\pi(x_i)Q(x_i, x)}{\pi(x)Q(x, x_i)} \wedge 1 \right)$ ,  $c_j = P(y, x_j) = Q(y, x_j) \left( \frac{\pi(x_j)Q(x_j, y)}{\pi(y)Q(y, x_j)} \wedge 1 \right)$  for every  $i, j$ . The desired transition kernel coupling  $\bar{Q}$  can also be represented as an  $n \times n$  matrix  $M_{\bar{Q}}$  with row sums  $[Q(x, x_1), \dots, Q(x, x_n)]$  and column sums  $[Q(y, x_1), \dots, Q(y, x_n)]$ .

Writing  $x = x_{i_0}$  and  $y = x_{j_0}$ , then Lemma 5 and Corollary 2 implies that  $M_{\bar{Q}}$  can be expressed as the sum of  $n \times n$  matrices  $M_{11}, M_{10}, M_{01}, M_{00}$ , where

$$\begin{aligned} M_{11}(i, j) &= M_{\bar{P}}(i, j) \mathbf{1}_{(i, j) \neq (i_0, j_0)} & M_{10}(i, j) &= M_{\bar{P}}(i, j_0) \mu_y(j) \mathbf{1}_{i \neq i_0} \\ M_{01}(i, j) &= \mu_x(i) M_{\bar{P}}(i_0, j) \mathbf{1}_{j \neq j_0} & M_{00}(i, j) &= M_{\bar{P}}(i_0, j_0) \mu_x(i) \mu_y(j). \end{aligned}$$

Here

$$\mu_x(i) = \begin{cases} \frac{Q(x, x_i) - P(x, x_i)}{1 - r_{i_0}} & \text{if } r_{i_0} < 1 \\ 1_{i=i_0} & \text{otherwise} \end{cases} \quad \text{and} \quad \mu_y(j) = \begin{cases} \frac{Q(y, x_j) - P(y, x_j)}{1 - c_{j_0}} & \text{if } c_{j_0} < 1 \\ 1_{j=j_0} & \text{otherwise.} \end{cases}$$

It is straightforward to verify that the row and column sums of  $\bar{Q}$  recover  $Q(x, \cdot)$  and  $Q(y, \cdot)$ .

Corollary 2 also specifies the necessary acceptance step coupling. After proposing a move from  $(x, y) = (x_{i_0}, x_{j_0})$  to  $(x_{i_1}, x_{j_1})$ , the probability accepting both  $x$  and  $y$  proposals is



$M_{11}(i_1, j_1)/M_{\bar{Q}}(i_1, j_1)$ , the probability of accepting the  $x$  proposal and rejecting the  $y$  proposal is  $M_{10}(i_1, j_1)/M_{\bar{Q}}(i_1, j_1)$ , the probability of rejecting the  $x$  proposal and accepting the  $y$  proposal is  $M_{01}(i_1, j_1)/M_{\bar{Q}}(i_1, j_1)$ , and the probability of rejecting both proposals and remaining at  $(x, y)$  is  $M_{00}(i_1, j_1)/M_{\bar{Q}}(i_1, j_1)$ . Again it is straightforward to confirm that these probabilities yield the desired transition behavior.

#### 4.4 Coupling with a minorization condition

Let  $P$  be an MH-like transition kernel generated by a proposal kernel  $Q$  and an acceptance rate function  $a$ , and assume that  $P$  has stationary distribution  $\pi$ . The kernel  $P$  satisfies a minorization condition corresponding to a set  $\mathcal{C} \subseteq \mathcal{X}$  if there exists  $\epsilon > 0$  and a probability measure  $\nu$  on  $(\mathcal{X}, \mathcal{F})$  such that  $P(x, A) \geq \epsilon \nu(A)$  for all  $x \in \mathcal{C}$  and  $A \in \mathcal{F}$ . If  $\mathcal{C} = \mathcal{X}$ , then a simple argument shows that  $\|P^n(x, \cdot) - \pi(\cdot)\|_{\text{TV}} \leq (1-\epsilon)^n$  for every  $n$  [Meyn and Tweedie, 2012]. When  $\mathcal{C}$  is a strict subset of  $\mathcal{X}$ , convergence rate results require additional assumptions to control the distribution of return times of the chain to the set  $\mathcal{C}$ .

Minorization arguments often employ the following coupling construction, sometimes called a Nummelin splitting [Nummelin, 1978, Athreya and Ney, 1978, Jones and Hobert, 2001, Gelman et al., 2010], to bound the convergence rate of a chain to its stationary distribution. Suppose that we run two identically distributed Markov chains  $(X_n)$  and  $(Y_n)$  in tandem, with one initialized arbitrarily and the other initialized from its stationary distribution  $\pi$ . If  $X_n, Y_n \in \mathcal{C}$  at some step  $n$ , then we flip a coin with probability  $\epsilon$  of heads. If the coin comes up heads we then draw  $X_{n+1} = Y_{n+1} \sim \nu$ , and otherwise we draw  $X_{n+1}$  and  $Y_{n+1}$  independently from their residual distributions so that  $X_{n+1} \sim \nu_x^r := (P(X_n, \cdot) - \epsilon \nu(\cdot))/(1-\epsilon)$  and  $Y_{n+1} \sim \nu_y^r := (P(Y_n, \cdot) - \epsilon \nu(\cdot))/(1-\epsilon)$ . Once the chains have met we update both chains according to the same draws from  $P$ , so that they remain together at all subsequent iterations. Otherwise we repeat the above whenever both chains occupy  $\mathcal{C}$ .

The procedure above yields a coupling  $\bar{P}$  of transition kernels  $P$ , defined without direct reference to the proposal or acceptance distributions. Alternatively, we can use Theorem 1 to represent this coupling in terms of a proposal coupling  $\bar{Q}$  and an acceptance indicator coupling  $\bar{B}$ . For every  $x, y \in \mathcal{C}$ , the kernel coupling  $\bar{P}((x, y), \cdot)$  described above is a mixture of a measure on the diagonal  $\Delta = \{(x, x) : x \in \mathcal{X}\} \subset \mathcal{X} \times \mathcal{X}$  and a product measure on the complement of  $\Delta$ . In particular, for any  $S \in \mathcal{F} \otimes \mathcal{F}$  we have  $\bar{P}((x, y), S) = \epsilon \nu(S \cap \Delta) + (1-\epsilon)(\nu_x^r \times \nu_y^r)(S \setminus \Delta)$ .

To represent  $\bar{P}$  by a two-step coupling it suffices to construct the coupled acceptance measure  $\Phi = (\Phi_{11}, \Phi_{10}, \Phi_{01}, \Phi_{00})$ . We continue to assume that  $Q(x, \cdot) \ll \lambda$  and  $Q(x, \{x\}) = 0$  for all  $x \in \mathcal{X}$ , and as noted above we have  $\beta(x) = 0$  if  $r(x) > 0$  and 1 otherwise, and

$$\mu(x, A_x) = \begin{cases} \frac{Q(x, A_x) - P(x, A_x \setminus \{x\})}{r(x)} & \text{if } r(x) > 0 \\ 1(x \in A_x) & \text{otherwise} \end{cases}$$

and similarly for  $\mu(y, A_y)$ . From the proof of Lemma 5, we can write the components of  $\Phi$  as follows:

$$\begin{aligned} \Phi_{11}((x, y), A_x \times A_y) &= \bar{P}((x, y), A_x \times A_y \cap (\{x\}^c \times \{y\}^c)) \\ \Phi_{10}((x, y), A_x \times A_y) &= \bar{P}((x, y), (A_x \cap \{x\}^c) \times \{y\}) \times \mu(y, A_y) \\ \Phi_{01}((x, y), A_x \times A_y) &= \mu(x, A_x) \times \bar{P}((x, y), \{x\} \times (A_y \cap \{y\}^c)) \\ \Phi_{00}((x, y), A_x \times A_y) &= r(x, y) \mu(x, A_x) \times \mu(y, A_y) \end{aligned}$$

where  $r(x, y) = \bar{P}((x, y), \{x\} \times \{y\})$  the probability that the kernel stays at  $(x, y)$ . In this case, the two chains are independent when meeting does not occur. Thus we have  $r(x, y) = r(x)r(y)$ . After setting all the  $\Phi_{ij}$ , we define  $\bar{Q} = \Phi_{00} + \Phi_{01} + \Phi_{10} + \Phi_{11}$  as our proposal coupling.

Next we consider the acceptance indicator coupling  $\bar{B}$ . Lemma 6 implies the existence of acceptance indicators  $(b_x, b_y) \sim \bar{B}((x, y), (x', y'))$  such that

$$\mathbb{P}(b_x = i, b_y = j \mid x, y, x', y') = \phi_{ij}((x, y), (x', y'))$$

where  $\phi_{i,j}((x, y), \cdot) = d\Phi_{i,j}((x, y), \cdot)/dQ((x, y), \cdot)$ . If we write every term of  $\Phi_{ij}$  in its integration form, then  $\phi_{i,j}$  can also be evaluated at every point as a ratio of densities. For example, let  $x' \neq x, y' \neq y$  be two different points in  $\mathcal{X}$ . Then the probability density of  $\bar{Q}$  for a move from  $(x, y)$  to  $(x', y')$  will be

$$\begin{aligned} q((x, y), (x', y')) &= (p(x, x') - \epsilon\nu(x'))(p(y, y') - \epsilon\nu(y')) + (p(x, x') - \epsilon\nu(x'))(q(y, y') - p(y, y')) \\ &\quad + (p(x, x') - q(x, x'))(q(y, y') - \epsilon\nu(y')) + (q(x, x') - p(x, x'))(q(y, y') - p(y, y')). \end{aligned}$$

Here  $p(x, x') = q(x, x')(\frac{\pi(x')q(x',x)}{\pi(x)q(x,x')} \wedge 1)$  is the transition kernel density from  $x$  to  $x'$ .

Thus, when  $x' \neq x, y' \neq y$ , and  $x' \neq y'$ , we have the following expressions for the distribution of the acceptance indicator pair  $(b_x, b_y)$  conditional on the current state  $(x, y)$  and the proposal state  $(x', y')$ :

$$\begin{aligned} p_{11} &:= \mathbb{P}(b_x = 1, b_y = 1 \mid x, y, x', y') = \frac{(p(x, x') - \epsilon\nu(x'))(p(y, y') - \epsilon\nu(y'))}{q((x, y), (x', y'))} \\ p_{10} &:= \mathbb{P}(b_x = 1, b_y = 0 \mid x, y, x', y') = \frac{(p(x, x') - \epsilon\nu(x'))(q(y, y') - p(y, y'))}{q((x, y), (x', y'))} \\ p_{01} &:= \mathbb{P}(b_x = 0, b_y = 1 \mid x, y, x', y') = \frac{(p(x, x') - q(x, x'))(q(y, y') - \epsilon\nu(y'))}{q((x, y), (x', y'))} \\ p_{00} &:= \mathbb{P}(b_x = 0, b_y = 0 \mid x, y, x', y') = \frac{(q(x, x') - p(x, x'))(q(y, y') - p(y, y'))}{q((x, y), (x', y'))}. \end{aligned}$$

The marginal probability of accepting the  $x$  move is  $p_{10} + p_{11}$  and the probability of accepting the  $y$  move is  $p_{01} + p_{11}$ , consistent with Lemma 7. As a sanity check, the probability density of moving from  $(x, y)$  to  $(x', y')$  according to our coupling and acceptance mechanism is  $q((x, y), (x', y')) \times p_{11} = (p(x, x') - \epsilon\nu(x')) \cdot (p(y, y') - \epsilon\nu(y'))$ , which is indeed the probability density of the minorization coupling  $\bar{P}((x, y), \cdot)$  at point  $(x', y')$ . The other cases, such as  $y' = x'$ , or  $x' = x$ , or  $y' = y$  can be calculated in the same way.

## 5 Discussion

The analysis of couplings is an appealing topic from both a theoretical and an applied point of view. In theoretical work, couplings arise as a proof technique in the study of Markov chains and other stochastic processes. They are also closely related to assignment problems under marginal constraints, a topic of interest since at least Birkhoff [1946]. In applications, couplings play a leading role in new techniques for convergence diagnosis, variance reduction, and unbiased estimation, as detailed in Section 1.

In this study, we have considered kernel and maximal kernel couplings for the Metropolis–Hastings algorithm and its relatives. In Theorem 1, we showed that every transition kernel coupling arises from a proposal coupling and an acceptance indicator with certain properties.

We also showed the converse: any kernel  $\bar{P}$  on the joint space generated this way must be a kernel coupling. In Theorem 2 we took this a step further, demonstrating that maximal kernel couplings correspond to pairs of proposal and acceptance couplings subject to a short list of additional conditions. These characterization results provide a unifying and descriptive framework to specify possible couplings of the MH transition kernel.

Throughout this paper, we have taken the perspective that the transition kernel of interest  $P$  on a state space  $(\mathcal{X}, \mathcal{F})$  arises from a given proposal kernel  $Q$  and acceptance rate function  $a$ . Alternatively, we could have begun with any kernel  $P$  such that there exists another ‘weakly dominating’ kernel  $Q$  such that  $P(x, A) \leq Q(x, A)$  for all  $x \in \mathcal{X}$  and  $A \in \mathcal{F}$  with  $x \notin A$ . With this perspective, our Theorem 1 reads roughly as follows: ‘suppose a kernel  $P$  is weakly dominated by  $Q$ , and let  $a(x, \cdot)$  be the Radon–Nikodym derivative of  $P(x, \cdot)$  with respect to  $Q(x, \cdot)$  away from the current state  $x$ . Then every kernel coupling  $\bar{P} \in \Gamma(P, P)$  is generated by a coupling  $\bar{Q} \in \Gamma(Q, Q)$  followed by an acceptance indicator coupling  $\bar{B}$  based on  $a$ .’ From this perspective, our result appears as a joint probability existence and characterization theorem which includes MH-like kernel couplings as a special case.

Looking forward, we believe that our results can aid in the design of new and more efficient methods based on couplings of the MH algorithm. We also expect our results to support the theoretical analysis of kernel couplings for this broad class of discrete-time Markov chains. For example, it may be possible to derive meeting time bounds or information on the spectrum of  $\bar{P}$  based on underlying data on  $\bar{Q}$  and  $\bar{B}$ , using a version of the drift and minorization arguments of Rosenthal [1995, 2002] or the techniques of Atchadé and Perron [2007]. A better understanding of the set of kernel couplings may also support optimization arguments like those of Boyd et al. [2004, 2006, 2009].

This work represents a step toward relating the properties of  $\Gamma(P, P)$  and  $\Gamma^{\max}(P, P)$  to those of  $\Gamma(Q, Q)$  and the associated acceptance couplings. As noted above, Markovian couplings which are efficient in the sense of Aldous [1983] are well understood in only a few special cases. Beyond that, only a few additional bounds exist on the efficiency of Markovian couplings [Burdzy and Kendall, 2000]. These questions remain largely open for MH-like chains. A clear understanding of the set of Markovian couplings will make it possible to identify better couplings for use in practice and settle the question of how efficient couplings can be for this important case.

## Acknowledgments

The authors would like to thank Pierre E. Jacob, Persi Diaconis, and Qian Qin for helpful comments. John O’Leary gratefully acknowledges support by the National Science Foundation through grant DMS-1844695.

## References

- D. Aldous. Random walks on finite groups and rapidly mixing Markov chains. In *Séminaire de Probabilités XVII 1981/82*, pages 243–297. Springer, 1983. 1, 4, 27
- D. Aldous and J. Fill. *Reversible Markov Chains and Random Walks on Graphs*. Berkeley, 1995. 2
- C. D. Aliprantis and O. Burkinshaw. *Principles of Real Analysis*. Gulf Professional Publishing, 1998. 15

- C. Andrieu, A. Doucet, and R. Holenstein. Particle Markov chain Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 72(3):269–342, 2010. [2](#)
- C. Andrieu, A. Lee, and S. Livingstone. A general perspective on the Metropolis-Hastings kernel. *arXiv preprint arXiv:2012.14881*, 2020. [3](#)
- Y. F. Atchadé and F. Perron. On the geometric ergodicity of Metropolis-Hastings algorithms. *Statistics*, 41(1):77–84, 2007. [27](#)
- K. B. Athreya and P. Ney. A new approach to the limit theory of recurrent Markov chains. *Transactions of the American Mathematical Society*, 245:493–501, 1978. [25](#)
- S. Banerjee and W. S. Kendall. Rigidity for Markovian maximal couplings of elliptic diffusions. *Probability Theory and Related Fields*, 168(1-2):55–112, 2017. [1](#)
- A. A. Barker. Monte Carlo calculations of the radial distribution functions for a proton–electron plasma. *Australian Journal of Physics*, 18(2):119–134, 1965. [2](#), [3](#)
- G. Birkhoff. Three observations on linear algebra. *Univ. Nac. Tucuman, Rev. Ser. A*, 5:147–151, 1946. [26](#)
- N. Biswas, P. E. Jacob, and P. Vanetti. Estimating convergence of Markov chains with L-lag couplings. In *Advances in Neural Information Processing Systems*, pages 7391–7401, 2019. [1](#)
- B. Böttcher. Markovian maximal coupling of Markov processes. *arXiv preprint arXiv:1710.09654*, 2017. [1](#)
- N. Bou-Rabee, A. Eberle, and R. Zimmer. Coupling and convergence for Hamiltonian Monte Carlo. *Annals of Applied Probability*, 30(3):1209–1250, 2020. [2](#)
- S. Boyd, P. Diaconis, and L. Xiao. Fastest mixing Markov chain on a graph. *SIAM Review*, 46(4):667–689, 2004. [27](#)
- S. Boyd, P. Diaconis, J. Sun, and L. Xiao. Fastest mixing Markov chain on a path. *The American Mathematical Monthly*, 113(1):70–74, 2006. [27](#)
- S. Boyd, P. Diaconis, P. Parrilo, and L. Xiao. Fastest mixing Markov chain on graphs with symmetries. *SIAM Journal on Optimization*, 20(2):792–819, 2009. [27](#)
- K. Burdzy and W. S. Kendall. Efficient Markovian couplings: examples and counterexamples. *Annals of Applied Probability*, pages 362–409, 2000. [1](#), [27](#)
- S. Connor and S. Jacka. Optimal co-adapted coupling for the symmetric random walk on the hypercube. *Journal of Applied Probability*, 45(3):703–713, 2008. [1](#)
- W. Doeblin. Exposé de la théorie des chaînes simples constantes de Markov à un nombre fini d'états. *Mathématique de l'Union Interbalkanique*, 2(77-105):78–80, 1938. [1](#)
- R. Douc, E. Moulines, P. Priouret, and P. Soulier. *Markov Chains*. Springer, 2018. [4](#), [15](#)
- J. H. Dshalalow. *Foundations of Abstract Analysis*. Springer Science & Business Media, 2012. [15](#)
- S. Duane, A. D. Kennedy, B. J. Pendleton, and D. Roweth. Hybrid Monte Carlo. *Physics Letters B*, 195(2):216–222, 1987. [2](#)
- R. Dudley. *Real Analysis and Probability*. Cambridge University Press, New York, 2002. [15](#), [21](#)

- P. A. Ernst, W. S. Kendall, G. O. Roberts, and J. S. Rosenthal. MEXIT: Maximal un-coupling times for stochastic processes. *Stochastic Processes and their Applications*, 129(2):355–380, 2019. [16](#)
- J. A. Fill. An interruptible algorithm for perfect sampling via Markov chains. In *Proceedings of the Twenty-Ninth Annual ACM Symposium on Theory of Computing*, pages 688–695, 1997. [1](#)
- J. M. Flegal and R. Herbei. Exact sampling for intractable probability distributions via a Bernoulli factory. *Electronic Journal of Statistics*, 6:10–37, 2012. [1](#)
- A. Gelman, S. Brooks, G. Jones, and X. Meng. *Handbook of Markov Chain Monte Carlo: Methods and Applications*. Chapman & Hall/CRC handbooks of modern statistical methods. CRC Press, 2010. ISBN 9781420079418. [25](#)
- P. W. Glynn and C.-H. Rhee. Exact estimation for Markov chain equilibrium expectations. *Journal of Applied Probability*, 51(A):377–389, 2014. [1](#)
- S. Goldstein. Maximal coupling. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 46(2):193–204, 1979. [1](#)
- J. B. Goodman and K. K. Lin. Coupling control variates for Markov chain Monte Carlo. *Journal of Computational Physics*, 228(19):7127–7136, 2009. [1](#)
- D. Griffeath. A maximal coupling for Markov chains. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 31(2):95–106, 1975. [1](#)
- T. E. Harris. On chains of infinite order. *Pacific Journal of Mathematics*, 5(Suppl. 1):707–724, 1955. [1](#)
- W. K. Hastings. Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, 57(1):97–109, 1970. [2](#)
- T. P. Hayes and A. Sinclair. A general lower bound for mixing of single-site dynamics on graphs. In *46th Annual IEEE Symposium on Foundations of Computer Science (FOCS'05)*, pages 511–520. IEEE, 2005. [2](#)
- T. P. Hayes and E. Vigoda. A non-markovian coupling for randomly sampling colorings. In *44th Annual IEEE Symposium on Foundations of Computer Science, 2003. Proceedings.*, pages 618–627. IEEE, 2003. [2](#)
- J. Heng and P. E. Jacob. Unbiased Hamiltonian Monte Carlo with couplings. *Biometrika*, 106(2):287–302, 2019. [1](#)
- E. P. Hsu and K.-T. Sturm. Maximal coupling of Euclidean Brownian motions. *Communications in Mathematics and Statistics*, 1(1):93–104, 2013. [1](#)
- P. E. Jacob, J. O’Leary, and Y. F. Atchadé. Unbiased Markov chain Monte Carlo methods with couplings. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82(3):543–600, 2020. [1](#), [2](#), [5](#)
- V. E. Johnson. Studying convergence of Markov chain Monte Carlo algorithms using coupled sample paths. *Journal of the American Statistical Association*, 91(433):154–166, 1996. [1](#)
- V. E. Johnson. A coupling-regeneration scheme for diagnosing convergence in Markov chain Monte Carlo algorithms. *Journal of the American Statistical Association*, 93(441):238–248, 1998. [1](#), [2](#), [5](#)

- G. L. Jones and J. P. Hobert. Honest exploration of intractable probability distributions via Markov chain Monte Carlo. *Statistical Science*, pages 312–334, 2001. [21](#), [25](#)
- M. Kartashov and V. Golomozyi. Maximal coupling procedure and stability of discrete Markov chains. ii. *Theory of Probability and Mathematical Statistics*, 87:65–78, 2013. [2](#)
- W. S. Kendall. Coupling, local times, immersions. *Bernoulli*, 21(2):1014–1046, 2015. [1](#)
- W. S. Kendall. Lectures on probabilistic coupling, 2017. [4](#)
- K. Kuwada. Characterization of maximal Markovian couplings for diffusion processes. *Electronic Journal of Probability*, 14:633–662, 2009. [1](#)
- D. A. Levin, Y. Peres, and E. L. Wilmer. *Markov Chains and Mixing Times*, volume 107. American Mathematical Soc., 2017. ISBN 1470429624. [4](#)
- T. Lindvall. *Lectures on the Coupling Method*. Dover Books on Mathematics, 1992. ISBN 0-486-42145-7. [1](#), [2](#), [4](#)
- N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller. Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21(6):1087–1092, 1953. [1](#)
- S. P. Meyn and R. L. Tweedie. *Markov chains and stochastic stability*. Springer Science & Business Media, 2012. [25](#)
- L. Middleton, G. Deligiannidis, A. Doucet, and P. E. Jacob. Unbiased smoothing using particle independent Metropolis-Hastings. In K. Chaudhuri and M. Sugiyama, editors, *Proceedings of Machine Learning Research*, volume 89, pages 2378–2387. PMLR, 16–18 Apr 2019. [1](#)
- L. Middleton, G. Deligiannidis, A. Doucet, and P. E. Jacob. Unbiased Markov chain Monte Carlo for intractable target distributions. *Electronic Journal of Statistics*, 14(2):2842–2891, 2020. ISSN 1935-7524. [1](#)
- R. Neal and R. Pinto. Improving Markov chain Monte Carlo estimators by coupling to an approximating chain. Technical report, Department of Statistics, University of Toronto, 2001. [1](#)
- R. M. Neal. Bayesian learning via stochastic dynamics. In *Advances in Neural Information Processing Systems*, pages 475–482, 1993. [2](#)
- R. M. Neal. Circularly-coupled Markov chain sampling. Technical report, Department of Statistics, University of Toronto, 1999. [1](#)
- R. M. Neal. MCMC using Hamiltonian dynamics. *Handbook of Markov Chain Monte Carlo*, 2(11):2, 2011. [2](#)
- E. Nummelin. Uniform and ratio limit theorems for Markov renewal and semi-regenerative processes on a general state space. *Annales de l’IHP Probabilités et statistiques*, 14(2):119–143, 1978. [25](#)
- J. O’Leary, G. Wang, and P. E. Jacob. Maximal couplings of the metropolis-hastings algorithm. In *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics*, volume 130 of *Proceedings of Machine Learning Research*, pages 1225–1233. PMLR, 2021. [1](#), [2](#), [5](#), [16](#)

- N. S. Pillai and A. Smith. Mixing times for a constrained Ising process on the two-dimensional torus at low density. In *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, volume 55, pages 1649–1678. Institut Henri Poincaré, 2019. [2](#)
- D. Piponi, M. Hoffman, and P. Sountsov. Hamiltonian Monte Carlo swindles. *Proceedings of Machine Learning Research*, 108:3774–3783, 26–28 Aug 2020. [1](#)
- J. Pitman. On coupling of Markov chains. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 35(4):315–322, 1976. [1](#)
- J. G. Propp and D. B. Wilson. Exact sampling with coupled Markov chains and applications to statistical mechanics. *Random Structures and Algorithms*, 9(1-2):223–252, 1996. [1](#)
- Q. Qin and J. P. Hobert. Wasserstein-based methods for convergence complexity analysis of MCMC with applications. *The Annals of Applied Probability*, 2021. [2](#)
- A. Reutter and V. E. Johnson. General strategies for assessing convergence of MCMC algorithms using coupled sample paths. Technical report, Institute of Statistics and Decision Sciences, Duke University, 1995. [2](#)
- G. O. Roberts and J. S. Rosenthal. Optimal scaling of discrete approximations to Langevin diffusions. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 60(1):255–268, 1998. [23](#)
- G. O. Roberts and R. L. Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2(4):341–363, 1996. [2](#), [23](#)
- J. S. Rosenthal. Minorization conditions and convergence rates for Markov chain Monte Carlo. *Journal of the American Statistical Association*, 90(430):558–566, 1995. [21](#), [27](#)
- J. S. Rosenthal. Analysis of the Gibbs sampler for a model related to James-Stein estimators. *Statistics and Computing*, 6(3):269–275, 1996. [2](#)
- J. S. Rosenthal. Quantitative convergence rates of Markov chains: A simple account. *Electronic Communications in Probability*, 7:123–128, 2002. [2](#), [27](#)