

# NONISOTHERMAL RICHARDS FLOW IN POROUS MEDIA WITH CROSS DIFFUSION

ESTHER S. DAUS, JOSIPA PINA MILIŠIĆ, AND NICOLA ZAMPONI

ABSTRACT. The existence of large-data weak entropy solutions to a nonisothermal immiscible compressible two-phase unsaturated flow model in porous media is proved. The model is thermodynamically consistent and includes temperature gradients and cross-diffusion effects. Due to the fact that some terms from the total energy balance are non-integrable in the classical weak sense, we consider so-called variational entropy solutions. A priori estimates are derived from the entropy balance and the total energy balance. The compactness is achieved by using the Div-Curl lemma.

## 1. INTRODUCTION

The modeling of multiphase flow through porous media is an important task in many engineering applications like for example geothermal systems, oil reservoir engineering, ground-water hydrology, and thermal energy storage. In unsaturated soils the flow can be essentially described by a two-phase flow of two immiscible fluids: air and water. A prominent example for the physical processes involved in this kind of flow is the infiltration of contaminants into upper layers of unsaturated soil where the air present in the soil is taken to be at a constant pressure. This situation can be modeled by the nonisothermal Richards equation where the water-phase consists of a mixture of  $N$  chemical components undergoing cross-diffusion effects. The model is based on equations representing mass, momentum and energy balances, and are coupled to constitutive relations derived in a thermodynamically consistent way starting from the Helmholtz free energy. Compared to the main difficulties related to the mathematical analysis of the similar model considered by Amaziane, Jurak, Pankratov and Piatnitski in [1] (coupling, strong nonlinearity, degeneracy of the diffusion term in the saturation equation), in this article we additionally deal with cross-diffusion terms which appear in the diffusion and in the thermal flux due to the presence of chemical components in the water-phase. Note also that in [1] the authors first regularized the phase pressures before they were using the time discretization to get a sequence of elliptic problems on which they could apply the fixed point argument, while in our approach, we apply the techniques of Buliček, Jüngel, Pokorný and Zamponi [8] taking into account the relation between the capillary pressure function and the interfacial boundary energy (3) introduced by Jurak, Koldoba, Konyukhov and Pankratov in [23].

The mathematical analysis of multiphase flows in porous media started due to their importance in applications in geological sciences and reservoir simulations. The first results on the existence of weak solutions of a simplified system describing a nonisothermal two-phase flow in porous media

---

*Date:* December 28, 2021.

*2000 Mathematics Subject Classification.* 35K65, 35K70, 35Q35, 35K55, 76S05.

*Key words and phrases.* Cross diffusion, global pressure, degenerate nonlinear parabolic equation, entropy method, existence of solutions.

All three authors were partially supported by the bilateral projects No. HR 04/2018, HR 19/2020 of the Austrian Exchange Service OeAD and the Ministry of Science and Education of the Republic of Croatia MZO. The first author acknowledges partial support from the Austrian Science Fund (FWF) grants W1245 and P30000. The second author acknowledges the support from the Croatian Science Foundation project IP-2019-04-1140. The third author acknowledges support from the Alexander von Humboldt Foundation (AvH) and from the Austrian Science Fund (FWF) grant P33010.

1 was obtained by Bocharov and Monakhov in [4, 5]. In [28] the authors proved the global existence  
2 of weak solutions for the nonisothermal, one-dimensional multicomponent heat-air-vapor transport  
3 model in porous textile materials. Existence of global weak solutions for Richards' model arising  
4 from the heat and moisture flow through a partially saturated porous medium was obtained by  
5 Beneš and Pažanin in [3]. The existence result for a nonisothermal immiscible incompressible two-  
6 phase flow in heterogeneous porous media was given by Amaziane, Jurak, Pankratov and Piatnitski  
7 in [1] where the authors used the concept of the nonisothermal global pressure introduced by  
8 Bocharov and Monakhov in [4] as the crucial mathematical tool to obtain a priori estimates and  
9 compactness results. Note that all these results were obtained for the single-species case only.

10 The mathematical existence analysis of cross-diffusion systems has been investigated extensively  
11 in the last two decades using entropy methods (see *e.g.* [10, 24, 25]), but the focus on how to  
12 model cross-diffusion effects (for instance in fluid mixtures) in a thermodynamically consistent  
13 way attracted the mathematical community's attention only recently, see *e.g.* [7, 8, 9, 13, 15, 18,  
14 22, 26]. For seminal works on the thermodynamic theory of fluid mixtures we want to mention *e.g.*  
15 [6, 19]. In general, cross diffusion means that the concentration gradient of one species induces a  
16 flux of another species. Mathematically, cross diffusion is usually described by strongly coupled  
17 parabolic systems of reaction-diffusion type, where the diffusion matrix is non-diagonal and in  
18 general neither symmetric nor positive definite, but the system possesses a (formal) gradient-flow  
19 structure which helps to perform the analysis without the use of a maximum principle, which is  
20 in general not available for such strongly coupled systems. Instead, the key tool in these entropy  
21 methods [10, 24, 25] is to perform a change of variables into so called *entropy variables*, which turn  
22 out to have also a physical meaning in terms of the chemical potentials of the system.

23 We want to give a brief overview of some of the works performed so far in the direction of  
24 thermodynamically consistent modelling of physical systems with cross-diffusion terms. In [7],  
25 Buliček and Havrda studied an incompressible fluid mixture, where the diffusion of all species  
26 and the heat flux were driven by the gradient of the chemical potentials and of the temperature,  
27 allowing for cross effects in such a way that the first and second laws of thermodynamics were  
28 still satisfied. In [15], Dreyer, Druet, Gajewski and Guhlke performed the existence analysis  
29 for an improved Nernst-Planck-Poisson model of compressible isothermal electrolytes with cross-  
30 diffusion phenomena due to mass conservation. In [9] an incompressible, multicomponent, heat  
31 conducting, electrically charged fluid model with cross-diffusion effects was studied. In [26] a  
32 single-phase fluid mixture in porous media was described and analyzed by using cross-diffusion  
33 equations. In [13] the authors studied an isothermal, unsaturated two-phase flow mixture with  
34 dynamic capillary pressure in porous media with cross-diffusion effects, where the assumption of  
35 constant temperature simplified the analysis. A recent breakthrough was made by Buliček, Jüngel,  
36 Pokorný and Zamponi in [8] by studying a steady compressible thermodynamically consistent  
37 Navier-Stokes-Fourier system for chemically reacting fluids with cross-diffusion effects. Very recent  
38 results were performed in [22], where the Maxwell-Stefan-Fourier equations in Fick-Onsager form  
39 were investigated and in [18], where a thermodynamically consistent reaction-cross-diffusion system  
40 coupled to an equation for the heat transfer was studied. The aim of this paper, namely to  
41 study in a thermodynamically consistent way the nonisothermal immiscible compressible two-phase  
42 unsaturated flow in porous media with cross diffusion, is completely new, to our best knowledge.

43 This paper is organized as follows. In the next section we start with an introduction of the main  
44 physical quantities, the presentation of the governing equations of the model of interest, followed  
45 by the constitutive relations and main mathematical hypotheses. In subsection 1.3 we present the  
46 concept of weak solutions called variational entropy solution. Section 2 is devoted to the derivation  
47 of the entropy balance equation which is the starting point for obtaining a priori estimates for  
48 smooth solutions. In Section 3, we deal with the weak sequential stability for smooth solutions. In

1 Section 4, we introduce the approximate problem using different regularization parameters. The  
 2 existence of solutions to the approximate system is shown using the Leray-Schauder fixed point  
 3 argument. Finally, using the compactness results established in Section 3, we perform the limit as  
 4 the time-discretization and the regularization parameters go to zero, obtaining finally a solution of  
 5 the degenerate system. We conclude the article with an Appendix where we present some auxiliary  
 6 technical results.

**1.1. The physical quantities and main assumptions.** We consider a nonisothermal Richards  
 flow process with cross-diffusion effects in a porous reservoir  $\Omega \subset \mathbb{R}^3$ , which is a bounded Lipschitz  
 domain, with a nondeformable skeleton. The time interval of interest is  $(0, \mathcal{T})$  and  $\mathcal{Q} = \Omega \times (0, \mathcal{T})$ .  
 The indexes  $w, n, s$  correspond to the wetting (water) phase, the non-wetting (air) phase, and the  
 skeleton.

**The temperature:**  $T = T(x, t)$  stands for the temperature in  $\Omega$ . We assume that the temperature  
 is locally in a heat equilibrium state, i.e.  $T_w = T_n = T_s = T$ .

**The phase pressures:** by  $p_w = p(x, t)$  we denote the water pressure while  $p_n = p_{at}$  represents the  
 constant air-pressure.

**The water mass density:**  $\rho_w = \rho(T, p)$ . We assume that the water phase is a mixture of  $N$  different  
 components. We denote the mass density of the  $i$ -th water component by  $\rho_i$ , for  $i = 1, \dots, N$ . It  
 holds that  $\rho = \sum_{i=1}^N \rho_i$ . Moreover, let be  $\vec{\rho} = (\rho_1, \dots, \rho_N)$ .

**The saturation functions:** By  $S_w = S(x, t)$  we denote the water saturation in  $\Omega$ .

**The energy densities of the water-phase and skeleton energy:** By  $E_w$  we denote the energy density  
 of the water phase. We assume that the skeleton energy density  $E_s$  depends on the temperature  
 only, i.e.  $E_s = E_s(T)$ . Following [23] the total energy density  $E_f$  is given by

$$(1) \quad E_f = E_w S + E_{int}(S).$$

7 By denoting with  $e$  the specific internal energy, we have  $E_w = (\rho e)$ .

8 **The capillary pressure:** The constitutive equation relating the capillary pressure  $P_c$  to the water-  
 9 phase saturation  $S$  is classically given as an algebraic relationship between  $P_c$  and  $S$ :  $P_c(S) =$   
 10  $p_n - p_w = p_{at} - p$ . A detailed discussion of this relationship can be found e.g. in [2]. The  
 11 relationship between  $P_c$  and  $S$  has been generalized on the basis of thermodynamical arguments  
 12 by Gray and Hassanizadeh [21]. They derived an extended relationship, which, in the case of  
 13 Richards flow and after scaling, can be written in the form

$$(2) \quad p = -P_c(S) + \partial_t f(S),$$

14 where  $f$  denotes the dynamic capillary pressure function.

15 **The skeleton and the interfacial boundary energies:** Following [23, 12], by  $E_{int} = E_{int}(S)$  we denote  
 16 the energy of the interphase density. The capillary pressure function is related to the interfacial  
 17 boundary energy by

$$P_c(S) = -\frac{\partial E_{int}}{\partial S}.$$

It follows that, up to a constant, the interface energy density (i.e. the energy density of the  
 boundary between the two phases) reads as

$$(3) \quad E_{int}(S) = \int_S^1 P_c(\xi) d\xi.$$

18 **The viscosity function:** By  $\mu_w = \mu(T)$  we denote the viscosity of water in  $\Omega$ .

19 **The porosity function:** By  $\Phi$  we denote the porosity of the domain  $\Omega$ . We assume that the porosity  
 20  $\Phi : \Omega \rightarrow [0, 1]$  is a Lebesgue-measurable function such that

$$(4) \quad \text{ess inf}_{\Omega} \Phi > 0, \quad \text{ess sup}_{\Omega} \Phi < 1.$$

- 1 The relative permeability function: By  $k_{r,w} = k_r(S)$  we denote the relative permeability of the water-  
 2 phase in  $\Omega$ . We assume that  $k_r$  is a continuous function that satisfies: (i)  $0 \leq k_r \leq 1$  on  $\mathbb{R}$ ; (ii)  
 3  $k_r(S) = 0$  for  $S \leq 0$  and  $k_r(S) = 1$  for  $S \geq 1$ .  
 4 The mobility functions: The water mobility function  $\lambda_w = \lambda(S, T)$  is defined by

$$(5) \quad \lambda(S, T) = \frac{k_r(S)}{\mu(T)},$$

where  $\mu(T)$  is the viscosity of the water phase.

The water-phase velocity: Following [2], the generalized Darcy law gives

$$(6) \quad \mathbf{v} = -K\lambda(S, T)\nabla p,$$

where  $\lambda$  is the mobility function, and  $0 < K < +\infty$  is the constant absolute permeability of the domain  $\Omega$ .

The chemical potentials: By  $\mu_1, \dots, \mu_N$  we denote the chemical potentials of the contaminants in the water-phase.

The diffusion and the thermal flux: Following thermodynamics of irreversible processes (TIP) in linear approximations [20], we have the following expressions for the diffusion and the thermal fluxes (provided that the external forces are neglected):

(i) the diffusion flux of the  $i$ th species is denoted by  $\mathbf{J}_i$  and is given by

$$(7) \quad \mathbf{J}_i = L_{i0}\nabla\left(\frac{1}{T}\right) - \sum_{j=1}^N L_{ij}\nabla\left(\frac{\mu_j}{T}\right), \quad i = 1, \dots, N,$$

where  $L_{ij} = L_{ij}(\vec{\rho}, T)$  are the diffusion coefficients (mobilities) which form the mobility matrix  $(L_{ij})_{i,j=1,\dots,N}$ .

(ii) the heat flux is denoted by  $\mathbf{q}$  and consists of Fourier's law and the molecular diffusion term

$$(8) \quad \mathbf{q} = -\kappa(T)\nabla T + \sum_{j=1}^N L_{0j}\nabla\left(\frac{\mu_j}{T}\right),$$

- 5 where  $\kappa(T) > 0$  is the heat conductivity given by

$$(9) \quad \kappa(T) = L_{00}T^{-2}.$$

- 6 **1.2. Model equations, constitutive relations and hypotheses.** Let  $\mathbf{v}$  be the water velocity  
 7 and  $\mathbf{v}_i$  be the velocity of the  $i$ -th water component, for  $i = 1, \dots, N$ . A water mixture (barycen-  
 8 tric) velocity  $\mathbf{v}$  is defined as

$$\rho\mathbf{v} = \sum_{i=1}^N \rho_i\mathbf{v}_i.$$

- 9 Each component of the water mixture satisfies the mass conservation law, i.e.

$$(10) \quad \Phi \frac{\partial}{\partial t}(S\rho_i) + \operatorname{div}(\rho_i\mathbf{v}_i) = r_i, \quad i = 1, \dots, N,$$

- 10 where  $r_1, \dots, r_N$  are reaction terms modeling e.g. chemical reactions between the components in  
 11 the water mixture. We assume that  $\sum_{i=1}^N r_i = 0$  (total mass conservation).

By summing these equations we get the total mass conservation of the water-phase:

$$\Phi \frac{\partial}{\partial t}(S\rho) + \operatorname{div}(\rho\mathbf{v}) = 0.$$

The transport of each water-component is divided into convective part ( $\rho_i\mathbf{v}$ ) and diffusive part ( $\mathbf{J}_i$ ) by dividing the component flux

$$\rho_i\mathbf{v}_i = \rho_i\mathbf{v} + \mathbf{J}_i,$$

1 where the diffusive flux is defined as  $\mathbf{J}_i = \rho_i(\mathbf{v}_i - \mathbf{v})$ . Obviously,  $\sum_{i=1}^N \mathbf{J}_i = 0$ .

The model consists of the mass conservation equation (10) for each water-component phase  $i$ , the energy conservation equation taken from [27] and the equation for the water saturation (2):

$$(11) \quad \Phi \frac{\partial}{\partial t}(S\rho_i) + \operatorname{div}(\rho_i\mathbf{v} + \mathbf{J}_i) = r_i, \quad i = 1, \dots, N,$$

$$(12) \quad \frac{\partial}{\partial t}(\Phi E_f + (1 - \Phi)E_s) + \operatorname{div}(((\rho e) + p)\mathbf{v} + \mathbf{q}) = 0,$$

$$(13) \quad \partial_t f(S) + P_c(S) + p = 0.$$

We assume the following initial conditions

$$(14) \quad \rho_i(\cdot, 0) = \rho_i^{in}, \quad S(\cdot, 0) = S^{in}, \quad T(\cdot, 0) = T^{in} \quad \text{in } \Omega, \quad i = 1, \dots, N,$$

where the initial data are Lebesgue measurable functions such that

$$(15) \quad \rho_i^{in} > 0, \quad 0 \leq S^{in} \leq 1, \quad T^{in} > 0, \quad \text{a.e. in } \Omega,$$

$$(16) \quad S^{in} \rho^{in}, \quad \Phi E_f^{in} + (1 - \Phi)E_s^{in}, \quad \Phi S^{in}(\rho\eta)^{in} + (1 - \Phi)(\rho\eta)_s^{in} \in L^1(\Omega).$$

Equations (11)-(13) are solved in a bounded domain  $\Omega \subset \mathbb{R}^3$  and are supplemented with complete slip boundary conditions for the velocity, and Robin boundary conditions for the diffusion fluxes and the heat flux on  $\partial\Omega \times (0, \infty)$ ,  $i = 1, \dots, N$ :

$$(17) \quad \mathbf{v} \cdot \nu = 0, \quad \mathbf{J}_i \cdot \nu = \sum_{k=1}^N b_{ik} \left( \frac{\mu_k}{T} - \frac{\mu_{0,k}}{T_0} \right), \quad \mathbf{q} \cdot \nu = \alpha(T - T_0),$$

where  $\alpha > 0$ ,  $T_0 > 0$ ,  $\mu_{0,1}, \dots, \mu_{0,N} \in \mathbb{R}$  are scalar constants, while  $(b_{ij})_{i,j=1,\dots,N} \in \mathbb{R}^{N \times N}$  is a constant, symmetric, positive semidefinite matrix such that

$$(18) \quad \sum_{i=1}^N b_{ij} = 0 \quad j = 1, \dots, N.$$

**Constitutive relations.** We assume that the thermodynamic quantities appearing in (11) and (12) are induced by the Helmholtz free energy density  $\rho\Psi$  in a thermodynamically consistent way. Precisely, we assume that the free energy density  $\rho\Psi$  decomposes into a sum of water- and skeleton-related contributions

$$(19) \quad \begin{aligned} \rho\Psi &= (\rho\Psi)_w + (\rho\Psi)_s, \\ (\rho\Psi)_w &= (\rho\Psi)_w(\vec{\rho}, T), \quad (\rho\Psi)_s = (\rho\Psi)_s(T). \end{aligned}$$

Following [8, Remark 1.2], we take the following expression for the functions  $(\rho\Psi)_w, (\rho\Psi)_s$ , under the simplifying assumption that the molar masses of each water-component phase are all coincident:

$$(20) \quad (\rho\Psi)_w(\vec{\rho}, T) = T \sum_{i=1}^N \rho_i \log \rho_i + \rho^\gamma - c_w \rho T \log T + p_{at},$$

$$(21) \quad (\rho\Psi)_s(T) = T - c_s T \log T.$$

2 Here  $c_w, c_s > 0$  are (scaled) heat capacities (of water and skeleton, respectively),  $p_{at} > 0$  is the  
3 atmospheric pressure, while

$$(22) \quad \gamma > 2.$$

As a consequence, the chemical potentials, pressure, internal energy density and entropy density for each phase are given by the following relations.

Chemical potentials:

$$(23) \quad \mu_i = \frac{\partial(\rho\Psi)_w}{\partial\rho_i} = T(\log\rho_i + 1) + \gamma\rho^{\gamma-1} - c_w T \log T, \quad i = 1, \dots, N.$$

Phase pressure:

$$(24) \quad p = -(\rho\Psi)_w + \sum_{i=1}^N \rho_i \mu_i = T\rho + (\gamma - 1)\rho^\gamma - p_{at}.$$

Phase energy density:

$$(25) \quad (\rho e) = (\rho\Psi)_w - T \frac{\partial(\rho\Psi)_w}{\partial T} = \rho^\gamma + c_w \rho T + p_{at}.$$

Phase entropy densities:

$$(26) \quad (\rho\eta) = -\frac{\partial(\rho\Psi)_w}{\partial T} = -\sum_{i=1}^N \rho_i \log \rho_i + c_w \rho (\log T + 1).$$

We point out that the Gibbs-Duhem relations hold

$$(27) \quad Td(\rho\eta) = d(\rho e) - \sum_{i=1}^N \mu_i d\rho_i,$$

where the differential  $d$  stands here for  $\partial_t$  or  $\partial_{x_i}$ ,  $i = 1, 2, 3$ .

Skeleton entropy and energy:

$$(28) \quad (\rho\eta)_s = -\frac{\partial(\rho\Psi)_s}{\partial T} = c_s - 1 + c_s \log T,$$

$$(29) \quad E_s = (\rho\Psi)_s - T \frac{\partial(\rho\Psi)_s}{\partial T} = c_s T.$$

- 1 **Hypotheses.** In this subsection we collect the main mathematical hypotheses which we imposed throughout this article.
- 2
- 3 **Phase velocity.** We also assume that  $\mu : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is a continuous, uniformly positive and bounded
- 4 function, while  $k_r \in C^0([0, 1])$  is an increasing, nonnegative function such that

$$(30) \quad \exists \alpha_r \in \left( \frac{2}{\gamma}, \frac{14}{3} \right), \quad \exists k_r^* > 0 : \quad \lim_{s \rightarrow 0} s^{-\alpha_r} k_r(s) = k_r^*.$$

**Capillary pressure.** We assume that  $P_c, f : (0, 1) \rightarrow \mathbb{R}$  are  $C^1$  strictly decreasing functions such that

$$(31) \quad \exists c_f > 0, \quad \frac{\gamma}{\gamma - 1} \leq q < 2 : \quad \inf_{0 < s < 1/2} \frac{|P'_c(s)| k_r(s)^{\frac{q}{2(q-1)}}}{|f'(s)|} \geq c_f,$$

$$(32) \quad \lim_{s \rightarrow 1} f(s) = -\infty, \quad \exists c'_f > 0 : \quad \left| \frac{d}{ds} \sqrt{k_r(s)} \right| \leq c'_f |f'(s)|, \quad 0 < s < 1,$$

$$(33) \quad f(0) > 0, \quad P_c(s) > 0, \quad \exists s_0 > 0 : \quad \inf_{s \in (0, s_0)} \frac{P_c(s)}{f(s)} > \frac{p_{at}}{f(0)},$$

$$(34) \quad \exists c_p > 0, \quad k_p \geq 0 : \quad \lim_{s \rightarrow 0^+} s^{k_p} P_c(s) = c_p,$$

$$(35) \quad \inf_{(0,1)} |f'| > 0, \quad \int_0^{1/2} |f'(u)| |\log u| du < \infty.$$

**Heat conductivity.** We assume that the heat conductivity  $\kappa(T)$  satisfies the following assumption

$$(36) \quad \exists \kappa_1, \kappa_2 > 0 : \quad \kappa_1(1 + T^\beta) \leq \kappa(T) \leq \kappa_2(1 + T^\beta),$$

1 where the exponent  $\beta > 0$  satisfies

$$(37) \quad \beta \geq \frac{q}{2-q}, \quad \beta > \frac{\gamma}{\gamma-2}, \quad \beta \geq \frac{4}{3}.$$

2 We point out that assumptions (35) and  $\beta \geq 4/3$  are only needed in the construction of the  
3 approximate scheme. A possible way to consider the case  $\beta < 4/3$  might be to adapt the (more  
4 involved) stationary scheme presented in [8] to the time-dependent system (11)–(13). However,  
5 we prefer here to restrict ourself to the case  $\beta \geq 4/3$  for the sake of simplicity. **Mobilities.** The  
6 Onsager-Casimir reciprocity relations imply the symmetry of the mobilities  $L_{ij} = L_{ji}$ ,  $L_{i0} = L_{0i}$   
7 for  $i, j = 1, \dots, N$ , and the second law of thermodynamics requires the positive semidefiniteness  
8 of the matrix  $(L_{ij}) \in \mathbb{R}^{N \times N}$  as well as  $L_{00} \geq 0$ .

9 Furthermore, we assume that  $L_{ij} = L_{ji} = \tilde{L}_{ij}(\vec{\rho}, T)$ ,  $L_{i0} = L_{0i} = \tilde{L}_{i0}(\vec{\rho}, T)$  for  $i, j = 1, \dots, N$ ,  
10 where  $\tilde{L}_{ij}, \tilde{L}_{i0} \in C^0(\mathbb{R}_+^N \times \mathbb{R}_+)$  such that

$$(38) \quad |\tilde{L}_{ij}(\vec{\rho}, T)| + \frac{1}{T} |\tilde{L}_{i0}(\vec{\rho}, T)| \leq C, \quad i, j = 1, \dots, N.$$

Moreover, we assume that there exists  $C, C' > 0$  such that

$$(39) \quad C |\Pi^N \vec{u}|^2 \leq \sum_{i,j=1}^N \tilde{L}_{ij}(\vec{\rho}, T) u_i u_j \leq C' |\Pi^N \vec{u}|^2, \quad \forall \vec{u} \in \mathbb{R}^N, \vec{\rho} \in \mathbb{R}_+^N, T > 0,$$

where  $\Pi^N = \mathbb{I} - \mathbf{1} \otimes \mathbf{1}/N$ , is the orthogonal projector on  $\text{span}\{\mathbf{1}\}^\perp$ . Thus,

$$(40) \quad (\Pi^N \vec{u})_i = u_i - \frac{1}{N} \sum_{j=1}^N u_j, \quad i = 1, \dots, N.$$

We also point out that, being  $(L_{ij})_{i,j=1,\dots,N}$  symmetric and positive semidefinite, (39) implies that  $\sum_{i=1}^N L_{ij} = 0$  for  $j = 1, \dots, N$ .

**Reaction terms.** For  $i = 1, \dots, N$  we assume  $r_i = \tilde{r}_i(\rho, T, \Pi^N \vec{\mu}/T)$  for some continuous function  $\tilde{r}_i : \mathbb{R}_+ \times (0, \infty) \times \mathbb{R}^N \rightarrow \mathbb{R}$  such that, for every  $\rho \geq 0, T > 0, \vec{\zeta} \in \mathbb{R}^N$ ,

$$(41) \quad \sum_{j=1}^N \tilde{r}_j(\rho, T, \vec{\zeta}) \zeta_j = 0,$$

$$(42) \quad \exists C_0, C_1, C_2 > 0, a > 2 : \quad \begin{cases} \sum_{j=1}^N \tilde{r}_j(\rho, T, \vec{\zeta}) \zeta_j \leq C_0 - C_1 |\Pi^N \vec{\zeta}|^a, \\ \sum_{j=1}^N |\tilde{r}_j(\rho, T, \vec{\zeta})| \leq C_2 \left(1 + |\Pi^N \vec{\zeta}|^{a-1}\right), \end{cases}$$

$$(43) \quad \forall \rho > 0, T > 0, \text{ the mapping } \vec{\zeta} \in \mathbb{R}^N \mapsto - \sum_{j=1}^N \tilde{r}_j(\rho, T, \vec{\zeta}) \zeta_j \in \mathbb{R} \text{ is convex,}$$

$$(44) \quad \exists \omega \in C^0(\mathbb{R}_+), \omega(0) = 0 : \quad \left| \sum_{j=1}^N (\tilde{r}_j(\rho, T, \vec{\zeta}) - \tilde{r}_j(\rho', T', \vec{\zeta})) \zeta_j \right| \leq \omega(|\rho - \rho'| + |T - T'|) |\vec{\zeta}|^a$$

$\forall \rho, \rho', T, T' > 0, \vec{\zeta} \in \mathbb{R}^N.$

1 **1.3. Solution concept, main result and the methodology of the proof.** Here we define the  
 2 concept of a weak solution to (11)–(13), (14), (17) and state the main result of the paper. Due  
 3 to the fact that some terms from the total energy balance are non-integrable in the classical weak  
 4 sense, we consider so-called variational entropy solutions (following the terminology in [31]), i.e.  
 5 solutions which fulfill (11) weakly, the integrated form of the total energy balance (12), and the  
 6 weak formulation of the entropy balance equation (51).

**Definition 1.** (*Variational entropy solution.*) *A variational entropy solution to (11)–(13), (14), (17) is a Lebesgue-measurable function  $(\vec{\rho}, T, S) : \Omega \times \mathbb{R}_+ \rightarrow \mathbb{R}_+^N \times \mathbb{R}_+ \times [0, 1]$  such that, for every  $\mathcal{T} > 0$  and  $i = 1, \dots, N$*

$$\begin{aligned}
 & P_c(S), \quad p, \quad \rho^\gamma \in L^1(Q_{\mathcal{T}}), \\
 & S\rho^\gamma, \quad S(\Phi E_f + (1 - \Phi)E_s) \in L^\infty(0, \mathcal{T}; L^1(\Omega)), \\
 & \log T, \quad T^{\beta/2} \in L^2(0, \mathcal{T}; H^1(\Omega)), \quad T \in L^\infty(0, \mathcal{T}; L^1(\Omega)) \cap L^{\beta + \frac{2}{3}}(Q_{\mathcal{T}}), \\
 & \Pi^N(\vec{\mu}/T) \in L^2(0, \mathcal{T}; H^1(\Omega)) \cap L^a(Q_{\mathcal{T}}), \quad \sqrt{\frac{\lambda(S, T)}{T}} \nabla p \in L^2(Q_{\mathcal{T}}), \\
 & \frac{1}{T}, \quad \sum_{i,j=1}^N b_{ij} \left( \Pi^N \frac{\vec{\mu}}{T} \right)_i \left( \Pi^N \frac{\vec{\mu}}{T} \right)_j \in L^1(\partial\Omega \times (0, \mathcal{T})), \\
 & T^{-1} f'(S) (\partial_t S)^2 \in L^1(Q_{\mathcal{T}}), \quad f(S) \in L^\infty(0, \mathcal{T}; W^{1,q}(\Omega)),
 \end{aligned}$$

for some suitable exponent  $a > 2$  and  $\beta, q$  given by (37), (31) (respectively), the weak formulation of the partial mass balances (11) holds

$$\begin{aligned}
 (45) \quad & \int_0^{\mathcal{T}} \langle \Phi \partial_t (S\rho_i), \varphi \rangle dt - \int_0^{\mathcal{T}} \int_{\Omega} (\rho_i \mathbf{v} + \mathbf{J}_i) \cdot \nabla \varphi dx dt \\
 & + \int_0^{\mathcal{T}} \int_{\partial\Omega} \sum_{k=1}^N b_{ik} \left( \frac{\mu_k}{T} - \frac{\mu_{0,k}}{T_0} \right) \varphi d\sigma dt = \int_0^{\mathcal{T}} \int_{\Omega} r_i \varphi dx dt, \\
 & i = 1, \dots, N, \quad \forall \varphi \in C^1(\overline{Q_{\mathcal{T}}}),
 \end{aligned}$$

as well as the integrated energy balance

$$\begin{aligned}
 (46) \quad & \int_{\Omega} (\Phi E_f(t) + (1 - \Phi)E_s(t)) dx + \alpha \int_0^t \int_{\partial\Omega} (T - T_0) d\sigma dt \\
 & = \int_{\Omega} (\Phi E_f^{in} + (1 - \Phi)E_s^{in}) dx, \quad \forall t \in [0, \mathcal{T}],
 \end{aligned}$$



eq. (13) holds a.e.  $x \in \Omega$ ,  $t > 0$ , the constitutive relations listed in subsection 1.2 are satisfied a.e.  $x \in \Omega$ ,  $t > 0$ , the entropy balance equation holds

$$\begin{aligned}
(47) \quad & \int_0^{\mathcal{T}} \langle \partial_t [\Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s], \varphi \rangle dt \\
& - \int_0^{\mathcal{T}} \int_{\Omega} \left( (\rho\eta)\mathbf{v} - \sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i + \frac{\mathbf{q}}{T} \right) \cdot \nabla \varphi dx dt \\
& = \int_0^{\mathcal{T}} \int_{\Omega} \left( \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) + L_{00} \left| \nabla \left( \frac{1}{T} \right) \right|^2 \right) \varphi dx dt \\
& + \int_0^{\mathcal{T}} \left( \frac{K}{T} \lambda(S, T) |\nabla p|^2 - \Phi \frac{1}{T} f'(S) (\partial_t S)^2 - \sum_{i=1}^N r_i \frac{\mu_i}{T} \right) \varphi dx dt \\
& + \int_0^{\mathcal{T}} \int_{\partial\Omega} \left( \alpha \frac{T_0 - T}{T} + \sum_{i,j=1}^N b_{ij} \frac{\mu_i}{T} \left( \frac{\mu_j}{T} - \frac{\mu_{0,j}}{T_0} \right) \right) \varphi d\sigma dt + \langle \xi, \varphi \rangle,
\end{aligned}$$

for every  $\varphi \in C^1(\overline{Q_{\mathcal{T}}})$ ,  $\varphi \geq 0$  a.e. in  $Q_{\mathcal{T}}$ , for a suitable choice of  $\xi \in \mathcal{M}(\overline{Q_{\mathcal{T}}})$ ,  $\xi \geq 0$  (nonnegative Radon measure), the saturation balance (13) holds a.e. in  $Q_{\mathcal{T}}$ , and the initial conditions (14) hold in the sense

$$(48) \quad \lim_{t \rightarrow 0} \int_{\Omega} \Phi S(t) \rho_i(t) \varphi dx = \int_{\Omega} \Phi S^{in} \rho_i^{in} \varphi dx, \quad i = 1, \dots, N,$$

$$(49) \quad \lim_{t \rightarrow 0} \int_{\Omega} (\Phi E_f(t) + (1 - \Phi) E_s(t)) \varphi dx = \int_{\Omega} (\Phi E_f^{in} + (1 - \Phi) E_s^{in}) \varphi dx,$$

$$(50) \quad \lim_{t \rightarrow 0} S(t) = S^{in} \quad \text{strongly in } L^1(\Omega),$$

1 for all  $\varphi \in C^1(\overline{\Omega})$ .

2 The main result of this paper is the following theorem.

3 **Theorem 1.** (*Large-data existence of solutions.*)

4 Let hypotheses (30)–(42) hold. Then there exists a variational entropy solution triplet  $(\vec{\rho}, T, S)$  to  
5 (11)–(13), (14), (17).

6 **1.4. The methodology of the proof of Theorem 1.** In the proof of Theorem 1 we mostly  
7 follow the approach given in [8]. The main difficulties in the analysis of the system are the  
8 combined presence of cross-diffusion and degenerate terms, as well as the fact that the system  
9 is time dependent. The cross diffusion means a lack of parabolicity (in the standard sense) for  
10 the system, while the presence of degeneracies in some terms prevent us from obtaining useful  
11 gradient estimates. In fact, because of the degeneracy of the relative permeability, we do not have  
12 uniform estimates for the gradient of the pressure or the gradient of the mass densities, which  
13 causes difficulties in the proof of the compactness result. Furthermore, the strong nonlinearities  
14 in the equations imply that a clever use of the Div-Curl Lemma was needed in order to show  
15 compactness of all the involved terms. Moreover, in comparison to [8], we do not have equations  
16 for the linear momentum, which is instead explicitly defined via Darcy's law. As a consequence,  
17 higher integrability estimates *à-la*-Bogovskiy for the mass density [8, Lemma 3.3] are unavailable.  
18 On the other hand, a higher integrability bound for the pressure on domains separated from the  
19 initial time (i.e.  $\Omega \times [\epsilon, \mathcal{T}]$ ,  $\epsilon > 0$  arbitrary) is derived from the equation for the capillary pressure,  
20 and is crucially employed in the compactness argument. Furthermore, while in [8] the stationary  
21 case is considered, the system studied here is time dependent, which clearly makes the analysis

1 quite more complicated. Indeed, a consequence of dealing with the nonstationary case is the  
 2 impossibility of obtaining a weak formulation for the energy balance (we only get an equation for  
 3 the *integrated* energy balance) due to the lack of control of terms in the energy flux. To overcome  
 4 the lack of parabolicity in the equations due to cross diffusion, we start by proving the existence  
 5 of solutions to an approximate system, which is uniformly elliptic with respect to a special set  
 6 of variables (“entropy variables”), i.e. chemical potentials and the logarithm of the temperature.  
 7 Then we derive suitable a priori estimates from the entropy balance equation, and finally pass to the  
 8 limit of vanishing approximation parameters. For the construction of a solution to the approximate  
 9 problem, we employ a time semi-discretization (with timestep  $\tau$ ) as well as lower-order and higher-  
 10 order regularizations with parameters  $\varepsilon > 0$ ,  $\delta > 0$  (respectively) of the mass conservation equation  
 11 for each water-component phase and the conservation of the energy equation. These levels of  
 12 approximation allow us to obtain  $H^2(\Omega)$  regularity for the approximated, time-discretized solutions.  
 13 In particular, regularizations of the cross-diffusion operator and the source terms yield  $H^2(\Omega)$   
 14 bounds for the chemical potentials, which imply that the (approximated) partial mass densities  
 15 are bounded and uniformly positive. A further regularization of the free energy and the capillary  
 16 pressure are required in order to retain higher integrability bounds for the mass densities and  
 17 temperature, as well as uniform positivity of the saturation, after taking the time-continuous limit  
 18  $\tau \rightarrow 0$  and the higher-order regularization limit  $\delta \rightarrow 0$ . We point out that, while making the  
 19 analysis more involved for the aforementioned reasons, the cross-diffusion terms yield in the end  
 20 an  $L^2$  bound for the relative chemical potentials, whose consequence is the following property: at  
 21 any point of the considered space-time domain, either the total density vanishes (i.e. there is a  
 22 vacuum) or each partial mass density is positive. In order to prove the strong convergence of the  
 23 sequence of approximated total density, partial densities and the temperature, we cannot employ  
 24 the Aubin-Lions Lemma because of the high nonlinearities present in the equations; instead, a  
 25 smart application of the Div-Curl lemma (a standard tool in fluid dynamics) is required in order  
 26 to complete the compactness argument. Finally, we point out that the addition of a nonstationary  
 27 term in the expression for the capillary pressure (13) is mainly motivated by the need of a bound for  
 28 the time derivative of (a function of) the saturation, which is crucial in the compactness argument.

## 29 2. ENTROPY BALANCE EQUATION AND A PRIORI ESTIMATES FOR SMOOTH SOLUTIONS

30 In this section we will derive suitable a priori estimates for smooth solutions. The starting point  
 31 is the following result, which shows that the entropy balance equation can be derived from the  
 32 mass and energy balance equations (11) and (12).

**Lemma 2.** (*Entropy balance equation.*)

Let  $(\vec{\rho}, T, S) : \Omega \times \mathbb{R}_+ \rightarrow \mathbb{R}_+^N \times \mathbb{R}_+ \times [0, 1]$  be a smooth solution to (11)–(13), (14), (17). Then the entropy balance equation holds

$$(51) \quad \frac{\partial}{\partial t} [\Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s] + \operatorname{div} \left( (\rho\eta)\mathbf{v} - \sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i + \frac{\mathbf{q}}{T} \right) \\ = \frac{K}{T} \lambda(S, T) |\nabla p|^2 + \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) + L_{00} \left| \nabla \left( \frac{1}{T} \right) \right|^2 - \Phi \frac{1}{T} f'(S) (\partial_t S)^2 - \sum_{i=1}^N r_i \frac{\mu_i}{T}.$$

*Proof.* The Gibbs-Duhem relations (27) allow us to write

$$(52) \quad \partial_t(\rho\eta) = - \sum_{i=1}^N \frac{\mu_i}{T} \partial_t \rho_i + \frac{1}{T} \partial_t(\rho e).$$

Next, from (28), (29) it follows

$$(53) \quad \partial_t(\rho\eta)_s = \frac{1}{T}\partial_t E_s.$$

We also point out that, summing (24) and (25) and employing (23) and (26) leads to

$$(54) \quad p + (\rho e) - \sum_{i=1}^N \rho_i \mu_i = T(\rho\eta).$$

Let us multiply (11) by  $-\mu_i/T$  and sum from  $i = 1, \dots, N$ , multiply (12) by  $1/T$  and then sum the obtained equations. In this way we get

$$(55) \quad \begin{aligned} & \Phi \left( - \sum_{i=1}^N \frac{\mu_i}{T} \partial_t(\rho_i S) + \frac{1}{T} \partial_t E_f \right) + (1 - \Phi) \frac{1}{T} \partial_t E_s \\ & - \sum_{i=1}^N \frac{\mu_i}{T} \operatorname{div}(\rho_i \mathbf{v} + \mathbf{J}_i) + \frac{1}{T} \operatorname{div}((\rho e + p)\mathbf{v} + \mathbf{q}) = - \sum_{i=1}^N r_i \frac{\mu_i}{T}. \end{aligned}$$

Let us now consider the first term on the left-hand side of (55). From (1), (3) we deduce

$$\begin{aligned} & - \sum_{i=1}^N \frac{\mu_i}{T} \partial_t(\rho_i S) + \frac{1}{T} \partial_t E_f \\ & = S \left( - \sum_{i=1}^N \frac{\mu_i}{T} \partial_t \rho_i + \frac{1}{T} \partial_t(\rho e) \right) + \left( - \sum_{i=1}^N \frac{\mu_i}{T} \rho_i + \frac{1}{T}(\rho e) \right) \partial_t S - \frac{1}{T} P_c(S) \partial_t S. \end{aligned}$$

Using (52), (54) and (13) from the equation above we get

$$- \sum_{i=1}^N \frac{\mu_i}{T} \partial_t(\rho_i S) + \frac{1}{T} \partial_t E_f = \partial_t(S(\rho\eta)) + \frac{1}{T} f'(S)(\partial_t S)^2.$$

Thanks to the above-written identity and (53), equation (55) becomes

$$\begin{aligned} & \Phi \partial_t(S(\rho\eta)) + (1 - \Phi) \partial_t(\rho\eta)_s + \frac{1}{T} \Phi f'(S)(\partial_t S)^2 \\ & - \sum_{i=1}^N \frac{\mu_i}{T} \operatorname{div}(\rho_i \mathbf{v} + \mathbf{J}_i) + \frac{1}{T} \operatorname{div}((\rho e + p)\mathbf{v} + \mathbf{q}) = - \sum_{i=1}^N r_i \frac{\mu_i}{T}, \end{aligned}$$

which can be equivalently rewritten as

$$(56) \quad \begin{aligned} & \Phi \partial_t(S(\rho\eta)) + (1 - \Phi) \partial_t(\rho\eta)_s + \frac{1}{T} \Phi f'(S)(\partial_t S)^2 \\ & + \mathbf{v} \cdot \left( - \sum_{i=1}^N \frac{\mu_i}{T} \nabla \rho_i + \frac{1}{T} \nabla(\rho e) + \frac{1}{T} \nabla p \right) \\ & + (\operatorname{div} \mathbf{v}) \left( - \sum_{i=1}^N \frac{\mu_i}{T} \rho_i + \frac{(\rho e)}{T} + \frac{p}{T} \right) + \operatorname{div} \left( - \sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i + \frac{\mathbf{q}}{T} \right) \\ & = - \sum_{i=1}^N \nabla \left( \frac{\mu_i}{T} \right) \cdot \mathbf{J}_i + \nabla \left( \frac{1}{T} \right) \cdot \mathbf{q} - \sum_{i=1}^N r_i \frac{\mu_i}{T}. \end{aligned}$$

Due to (27) one has

$$(57) \quad \nabla(\rho\eta) = - \sum_{i=1}^N \frac{\mu_i}{T} \nabla \rho_i + \frac{1}{T} \nabla(\rho e),$$

while from (7) and (8) it follows

$$(58) \quad \begin{aligned} & - \sum_{i=1}^N \nabla \left( \frac{\mu_i}{T} \right) \cdot \mathbf{J}_i + \nabla \left( \frac{1}{T} \right) \cdot \mathbf{q} \\ & = \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) + L_{00} \left| \nabla \left( \frac{1}{T} \right) \right|^2 \geq 0. \end{aligned}$$

By inserting (57)-(58) into (56) and making use of (54) we conclude

$$\begin{aligned} & \Phi \partial_t (S(\rho\eta)) + (1 - \Phi) \partial_t (\rho\eta)_s + \mathbf{v} \cdot \nabla(\rho\eta) + (\operatorname{div} \mathbf{v})(\rho\eta) \\ & + \frac{1}{T} \mathbf{v} \cdot \nabla p + \operatorname{div} \left( - \sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i + \frac{\mathbf{q}}{T} \right) \\ & = \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) + L_{00} \left| \nabla \left( \frac{1}{T} \right) \right|^2 - \frac{1}{T} \Phi f'(S) (\partial_t S)^2 - \sum_{i=1}^N r_i \frac{\mu_i}{T}, \end{aligned}$$

1 which, thanks to (6), yields the entropy balance equation (51).  $\square$

**Remark 3.** *The right-hand side of (51) is nonnegative thanks to the assumptions that  $(L_{ij}) \in \mathbb{R}^{N \times N}$  is a positive semidefinite matrix,  $L_{00} \geq 0$ , the mobility function  $\lambda$  in (5) is nonnegative,  $f'(S) \leq 0$  and  $-\sum_{i=1}^N r_i \frac{\mu_i}{T} \geq 0$ . Therefore, the second law of thermodynamics holds true*

$$(59) \quad \partial_t [\Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s] + \operatorname{div} \left( (\rho\eta) \mathbf{v} - \sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i + \frac{\mathbf{q}}{T} \right) \geq 0.$$

2 The main a priori estimates are given by the following proposition.

**Lemma 4.** *Any smooth solution to (11)–(13), (14)–(17) satisfies*

$$(60) \quad \|P_c(S)\|_{L^1(Q_T)} + \|p\|_{L^1(Q_T)} + \|\rho\|_{L^\gamma(Q_T)} \leq C,$$

$$(61) \quad \|S^{1/\gamma} \rho\|_{L^\infty(0,T;L^\gamma(\Omega))} + \|Sp\|_{L^\infty(0,T;L^1(\Omega))} + \|\Phi E_f + (1 - \Phi)E_s\|_{L^\infty(0,T;L^1(\Omega))} \leq C,$$

$$(62) \quad \|T\|_{L^\infty(0,T;L^1(\Omega))} + \|\log T\|_{L^2(0,T;H^1(\Omega))} + \|T^{\beta/2}\|_{L^2(0,T;H^1(\Omega))} \leq C,$$

$$(63) \quad \|T\|_{L^{\beta+\frac{2}{3}}(Q_T)} \leq C,$$

$$(64) \quad \|\Pi^N(\bar{\mu}/T)\|_{L^2(0,T;H^1(\Omega))} + \|\Pi^N(\bar{\mu}/T)\|_{L^a(Q_T)} \leq C,$$

$$(65) \quad \left\| \sqrt{\frac{\lambda(S,T)}{T}} \nabla p \right\|_{L^2(Q_T)} \leq C,$$

$$(66) \quad \|T^{-1}\|_{L^1(\partial\Omega \times (0,T))} \leq C,$$

$$(67) \quad \left\| \sum_{i,j=1}^N b_{ij} \frac{\mu_i}{T} \frac{\mu_j}{T} \right\|_{L^1(\partial\Omega \times (0,T))} \leq C,$$

$$(68) \quad \|T^{-1} f'(S) (\partial_t S)^2\|_{L^1(Q_T)} + \|f(S)\|_{L^\infty(0,T;W^{1,q}(\Omega))} \leq C,$$

1 where here and in the following,  $C > 0$  denotes a generic constant depending only on the given  
 2 data and (possibly) on  $\mathcal{T}$ .

*Proof.* We start by integrating the energy balance (12) in space and time:

$$\int_{\Omega} (\Phi E_f + (1 - \Phi) E_s)|_{t=0}^{t=t_1} dx + \int_0^{t_1} \int_{\partial\Omega} (((\rho e) + p)\mathbf{v} + \mathbf{q}) \cdot \nu d\sigma dt = 0, \quad \forall t_1 \in (0, \mathcal{T}).$$

From (17) it follows

$$\int_{\Omega} (\Phi E_f + (1 - \Phi) E_s)|_{t=0}^{t=t_1} dx + \alpha \int_0^{t_1} \int_{\partial\Omega} (T - T_0) d\sigma dt = 0, \quad \forall t_1 \in (0, \mathcal{T}),$$

3 and so

$$(69) \quad \sup_{t \in [0, \mathcal{T}]} \int_{\Omega} (\Phi E_f + (1 - \Phi) E_s) dx + \alpha \int_0^{\mathcal{T}} \int_{\partial\Omega} T d\sigma dt \leq C.$$

The  $L^\infty(0, \mathcal{T}; L^\gamma(\Omega))$  bound for  $S^{1/\gamma}\rho$  and the  $L^\infty(0, \mathcal{T}; L^1(\Omega))$  bound for  $Sp$  are derived from the constitutive relations (1), (24), (25) and estimate (69). Therefore (61) holds. As a byproduct, from (29), (4), (61) it follows

$$(70) \quad \|T\|_{L^\infty(0, \mathcal{T}; L^1(\Omega))} \leq C.$$

Let  $\zeta : [0, 1] \rightarrow [0, 1]$  be a  $C^1([0, 1])$  cutoff function such that  $\zeta = 1$  on  $[0, 1/3]$ ,  $\zeta = 0$  on  $[2/3, 1]$ , and  $\zeta$  is decreasing in  $(1/3, 2/3)$ . Let us multiply (13) by  $\zeta(S)$  and define  $f_1(s) = \int_0^s \zeta(u) f'(u) du$  for  $0 \leq s \leq 1$ . Notice that  $f_1$  is bounded by construction and the assumptions on  $\zeta$ . We obtain

$$\partial_t f_1(S) + \zeta(S) P_c(S) + \zeta(S) p = 0.$$

Integrating the above identity leads to

$$\int_{\Omega} f_1(S(t)) dx + \int_0^t \int_{\Omega} (\zeta(S) P_c(S) + \zeta(S) p) dx dt' = \int_{\Omega} f_1(S^{in}) dx, \quad t > 0.$$

Since  $p \geq -p_{at}$  a.e. in  $Q_{\mathcal{T}}$  and  $P_c$  is a nonnegative function while  $f_1$  is bounded, we obtain that  $\zeta(S) P_c(S)$  is bounded in  $L^1(Q_{\mathcal{T}})$ . Since  $(1 - \zeta) P_c$  is bounded (because  $P_c$  is nonnegative, decreasing, and  $1 - \zeta$  vanishes near 0), we infer

$$(71) \quad \int_0^t \int_{\Omega} P_c(S) dx dt \leq C.$$

On the other hand, multiplying (13) times  $1 - \zeta(S)$  and defining  $f_2(s) = \int_0^s (1 - \zeta(u)) f'(u) du$  yields

$$\partial_t f_2(S) + (1 - \zeta(S)) P_c(S) + (1 - \zeta(S)) p = 0.$$

Integrating the above identity leads to

$$- \int_{\Omega} f_2(S(t)) dx = \int_0^t \int_{\Omega} ((1 - \zeta(S)) P_c(S) + (1 - \zeta(S)) p) dx dt' - \int_{\Omega} f_2(S^{in}) dx, \quad t > 0.$$

We know that  $(1 - \zeta) P_c$  is bounded. Moreover, the properties of  $\zeta$ , the constitutive relations (1), (24), (25) and (61) imply that  $(1 - \zeta(S)) p$  is bounded in  $L^1(Q_{\mathcal{T}})$ . It follows that

$$- \int_{\Omega} f_2(S(t)) dx \leq C, \quad t > 0.$$

Since  $-f_2(s) = -f_2(2/3) + f(2/3) - f(s)$  for  $s > 2/3$  and the fact that  $f$  is bounded in  $[0, 2/3]$  (because it is decreasing, smooth in  $(0, 1)$ , and  $f(0) < \infty$ ), we conclude

$$(72) \quad - \int_{\Omega} f(S(t)) dx \leq C, \quad t > 0.$$

1 Integrating (13) in  $Q_{\mathcal{T}}$  and employing (71), (72) and the fact that  $p \geq -p_{at}$  a.e. in  $Q_{\mathcal{T}}$  yields (60).

Let us now integrate the entropy balance equation (51) with respect to space and time (from  $t = 0$  to  $t = \mathcal{T}$ ) and employ (14)–(17). We get

$$\begin{aligned}
& \int_0^{\mathcal{T}} \int_{\Omega} \left( \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) + \kappa(T) |\nabla \log T|^2 \right) dx dt \\
& + \int_0^{\mathcal{T}} \int_{\Omega} \frac{K}{T} \lambda(S, T) |\nabla p|^2 dx dt - \int_0^{\mathcal{T}} \int_{\Omega} \sum_{i=1}^N r_i \frac{\mu_i}{T} dx dt \\
& + \int_0^{\mathcal{T}} \int_{\partial\Omega} \left( \alpha \frac{T_0 - T}{T} + \sum_{i,j=1}^N b_{ij} \frac{\mu_i}{T} \left( \frac{\mu_j}{T} - \frac{\mu_{0,j}}{T_0} \right) \right) d\sigma dt \\
(73) \quad & = \int_{\Omega} \left( \Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s \right) dx \Big|_{t=0}^{t=\mathcal{T}},
\end{aligned}$$

where  $\kappa(T)$  is given by (9). Let us consider the right-hand side of (73). From (26) and (28) we deduce

$$\int_{\Omega} \left( \Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s \right) dx \Big|_{t=0}^{t=\mathcal{T}} \leq C_1 + C \int_{\Omega} (\Phi S \rho (\log T)_+ + (1 - \Phi)(\log T)_+) dx \Big|_{t=\mathcal{T}},$$

where  $C_1$  depends on the initial datum and  $(u)_+ := \max\{u, 0\}$  denotes the positive part. Since (1), (25), (29) hold, we deduce

$$\int_{\Omega} \left( \Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s \right) dx \Big|_{t=0}^{t=\mathcal{T}} \leq C_1 + C \int_{\Omega} (\Phi E_f + (1 - \Phi) E_s) dx \Big|_{t=\mathcal{T}}.$$

2 From (69) we get

$$(74) \quad \int_{\Omega} \left( \Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s \right) dx \Big|_{t=0}^{t=\mathcal{T}} \leq C.$$

Let us now consider the boundary term in (73). Since the matrix  $(b_{ij})_{i,j=1}^N$  is positive semidefinite and (18) holds, it follows (via Cauchy-Schwartz)

$$\sum_{i,j=1}^N b_{ij} \frac{\mu_i}{T} \left( \frac{\mu_j}{T} - \frac{\mu_{0,j}}{T_0} \right) \geq \frac{1}{2} \sum_{i,j=1}^N b_{ij} \frac{\mu_i}{T} \frac{\mu_j}{T} - \frac{1}{2} \sum_{i,j=1}^N b_{ij} \frac{\mu_{0,i}}{T_0} \frac{\mu_{0,j}}{T_0},$$

which yields (67). Furthermore, (42) implies

$$- \int_0^{\mathcal{T}} \int_{\Omega} \sum_{i=1}^N r_i \frac{\mu_i}{T} dx dt \geq C_1 \|\Pi^N(\vec{\mu}/T)\|_{L^a(Q_{\mathcal{T}})}^a - C_0,$$

which yields the  $L^a(Q_{\mathcal{T}})$  bound for  $\Pi^N \vec{\mu}/\mathcal{T}$ . On the other hand, (39) implies that

$$\begin{aligned}
& \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) = \sum_{i,j=1}^N L_{ij} \sum_{k=1}^N \partial_k \left( \frac{\mu_i}{T} \right) \partial_k \left( \frac{\mu_j}{T} \right) \\
& = \sum_{k=1}^N \left( \sum_{i,j=1}^N L_{ij} \partial_k \left( \frac{\mu_i}{T} \right) \partial_k \left( \frac{\mu_j}{T} \right) \right) \geq C \sum_{k=1}^N |\Pi^N \partial_k(\vec{\mu}/T)|^2 \\
(75) \quad & = C \sum_{k=1}^N \sum_{i=1}^N \left( \Pi^N \partial_k(\vec{\mu}/T) \right)_i^2 = C \sum_{i=1}^N |\nabla \Pi^N(\vec{\mu}/T)_i|^2 = C |\nabla \Pi^N(\vec{\mu}/T)|^2.
\end{aligned}$$

In this way from (73) we get the estimate

$$\|\nabla \Pi^N(\frac{\vec{\mu}}{T})\|_{L^2(Q_T)} \leq C,$$

1 which, together with the  $L^a(Q_T)$  bound for  $\Pi^N \vec{\mu}/\mathcal{T}$ , yields (64).

Clearly (66) follows immediately from (73) and the previous bounds for the boundary integrals and (74). On the other hand, from (36) it follows

$$(76) \quad \kappa(T)|\nabla \log T|^2 \geq \kappa_1(1 + T^\beta)|\nabla \log T|^2 \geq c(|\nabla \log T|^2 + |\nabla T^{\beta/2}|^2).$$

Therefore (66), (70), (73), (76) lead to the uniform bounds for  $T$ ,  $\log T$  in  $L^\infty(0, T; L^1(\Omega))$ ,  $L^2(0, T; H^1(\Omega))$  (respectively) contained in (62). In order to prove the uniform bound for  $T^{\beta/2}$  in  $L^2(0, T; H^1(\Omega))$  (which will yield (62)), it is necessary to distinguish two cases.

*Case  $1 < \beta \leq 2$ .* In this case  $T^{\beta/2}$  is bounded in  $L^\infty(0, T; L^1(\Omega))$  thanks to (70), so Poincaré's Lemma yields the statement.

*Case  $\beta > 2$ .* In this case we observe that

$$\nabla T = \chi_{(0,1)}(T)T\nabla \log T + \chi_{[1,\infty)}(T)\frac{2}{\beta}T^{1-\beta/2}\nabla T^{\beta/2}$$

is bounded in  $L^2(Q_T)$  given the uniform bounds for  $\nabla \log T$  and  $\nabla T^{\beta/2}$  in  $L^2(Q_T)$ . Given that (70) holds, it follows via Poincaré's Lemma that  $T$  is bounded in  $L^2(0, T; H^1(\Omega))$  and therefore (via Sobolev embedding) in  $L^2(0, T; L^6(\Omega))$ . We apply now an iterative interpolation argument to prove that  $T^{\beta/2}$  is bounded in  $L^2(0, T; L^1(\Omega))$ ; the statement will then follow via Poincaré's Lemma. Assume that, for some  $p \geq 2$ ,

$$(77) \quad \|T\|_{L^p(0,T;L^{3p}(\Omega))} \leq C.$$

Since (70) holds, a straightforward interpolation argument yields

$$\|T\|_{L^{2r}(0,T;L^r(\Omega))} \leq C, \quad r = \frac{5}{6} + \frac{p}{2}.$$

2 This means that  $T^r$  is bounded in  $L^2(0, T; L^1(\Omega))$ . If  $\beta \leq 2r$  then we deduce (easy interpolation  
3 with (70)) that  $T^{\beta/2}$  is bounded in  $L^2(0, T; L^1(\Omega))$ , which is what we want. Let us therefore assume  
4 that  $\beta > 2r$ . It follows from the known bounds for  $\nabla \log T$ ,  $\nabla T^{\beta/2}$  in  $L^2(Q_T)$  that  $\nabla T^r$  is bounded  
5 in  $L^2(Q_T)$ . Since  $T^r$  is bounded in  $L^2(0, T; L^1(\Omega))$ , Poincaré's Lemma yields that  $T^r$  is bounded  
6 in  $L^2(0, T; H^1(\Omega))$ , thus Sobolev's embedding implies that  $T^r$  is bounded in  $L^2(0, T; L^6(\Omega))$ , that  
7 is, (77) holds with  $p$  replaced by  $2r = \frac{5}{3} + p$ . Since (77) holds true with  $p = 2$ , an easy induction  
8 argument implies that (77) holds for every  $p$  having the form  $p = 2 + \frac{5}{3}n$ , with  $n \in \mathbb{N}$  arbitrary, as  
9 long as  $2 + \frac{5}{3}n < \frac{\beta}{2}$ . Since  $2 + \frac{5}{3}n \rightarrow \infty$  as  $n \rightarrow \infty$ , iterating this argument allows us to eventually  
10 deduce that  $T^{\beta/2}$  is bounded in  $L^2(0, T; L^1(\Omega))$ . Bound (65) follows immediately from (73).

Let us now show (68). The bound for the first term in (68) follows immediately from (73); let us try to bound the second term. Take the gradient of (13), and multiply the resulting equation times  $|\nabla f(S)|^{q-2}\nabla f(S)$  and integrate in  $\Omega \times [0, t_1]$ , where  $t_1 \in [0, \mathcal{T}]$  is generic. It follows

$$\begin{aligned} & \frac{1}{q} \int_{\Omega} |\nabla f(S)|^q dx \Big|_{t=0}^{t=t_1} + \int_0^{t_1} \int_{\Omega} |f'(S)|^{q-2} f'(S) P'_c(S) |\nabla S|^q dx dt \\ & = - \int_0^{t_1} \int_{\Omega} |\nabla f(S)|^{q-2} \nabla f(S) \cdot \nabla p dx dt. \end{aligned}$$

Young's inequality and (65) lead to

$$\begin{aligned}
& \frac{1}{q} \int_{\Omega} |\nabla f(S)|^q dx \Big|_{t=0}^{t=t_1} + \int_0^{t_1} \int_{\Omega} |f'(S)|^{q-2} f'(S) P'_c(S) |\nabla S|^q dx dt \\
& \leq \frac{\varepsilon}{2} \int_0^{t_1} \int_{\Omega} \frac{T}{\lambda} |\nabla f(S)|^{2(q-1)} dx dt + \frac{1}{2\varepsilon} \int_0^{t_1} \int_{\Omega} \frac{\lambda}{T} |\nabla p|^2 dx dt \\
& \leq \frac{\varepsilon(q-1)}{q} \int_0^{t_1} \int_{\Omega} \lambda^{-q/(2q-2)} |\nabla f(S)|^q dx dt + \frac{\varepsilon(2-q)}{2q} \int_0^{t_1} \int_{\Omega} T^{q/(2-q)} dx dt + C\varepsilon^{-1}.
\end{aligned}$$

Thanks to (5) and the boundedness of  $\mu$ , it follows  $1/\lambda = \mu(T)/k_r(S) \leq C/k_r(S)$ . Furthermore (37), (62) yield that  $\|T^{q/(2-q)}\|_{L^1(Q_{\mathcal{T}})} \leq C$ . From these estimates and the fact that both  $P_c$  and  $f$  are nonincreasing, we deduce

$$\begin{aligned}
& \frac{1}{q} \int_{\Omega} |\nabla f(S)|^q dx \Big|_{t=0}^{t=t_1} + \int_0^{t_1} \int_{\Omega} |f'(S)|^{q-1} |P'_c(S)| |\nabla S|^q dx dt \\
& \leq C\varepsilon \int_0^{t_1} \int_{\Omega} k_r(S)^{-q/(2q-2)} |f'(S)|^q |\nabla S|^q dx dt + C\varepsilon^{-1}.
\end{aligned}$$

Since  $k_r$  is uniformly positive in  $[1/2, 1]$ , it follows

$$\begin{aligned}
& \frac{1}{q} \int_{\Omega} |\nabla f(S)|^q dx \Big|_{t=0}^{t=t_1} + \int_0^{t_1} \int_{\Omega} |f'(S)|^{q-1} |P'_c(S)| |\nabla S|^q dx dt \\
& \leq C\varepsilon \int_{\Omega} |\nabla f(S(t_1))|^q dx \\
& \quad + C\varepsilon \int_{Q_{t_1} \cap \{S < 1/2\}} k_r(S)^{-q/(2q-2)} |f'(S)|^q |\nabla S|^q dx dt + C\varepsilon^{-1}.
\end{aligned}$$

Applying (31) with  $\varepsilon < c_f/qC$  leads to

$$\int_{\Omega} |\nabla f(S)|^q dx \Big|_{t=0}^{t=t_1} \leq C + C \int_{\Omega} |\nabla f(S(t_1))|^q dx, \quad \forall t_1 \in [0, \mathcal{T}],$$

- 1 which via a Gronwall argument yields a bound for  $\nabla f(S)$  in  $L^\infty(0, \mathcal{T}; L^q(\Omega))$ . However, since
- 2 (72) holds and  $f$  is upper bounded, (68) follows via Poincaré's Lemma. This finishes the proof of
- 3 Lemma 4. □

- 4 The following three propositions (Prop. 5, Prop. 6, Prop. 7) give a priori estimates needed for
- 5 the existence result. Their proofs are rather technical and they are given in the Appendix. The
- 6 first one gives estimates for the terms in (11), (12).

**Proposition 5.** *Any smooth solution to (11)–(13), (14)–(17) satisfies*

$$(78) \quad \|\rho \mathbf{v}\|_{L^{\frac{2\beta}{\beta+1}}(0, \mathcal{T}; L^{\frac{6\beta\gamma}{\gamma+6\beta+3\beta\gamma}}(\Omega))} \leq C.$$

$$(79) \quad \|\mathbf{J}_i\|_{L^2(Q_{\mathcal{T}})} \leq C, \quad i = 1, \dots, N.$$

$$(80) \quad \|\rho_i \mathbf{v} + \mathbf{J}_i\|_{L^m(Q_{\mathcal{T}})} \leq C, \quad i = 1, \dots, N.$$

$$(81) \quad \|\partial_t(\Phi S \rho_i)\|_{L^m(0, \mathcal{T}; W^{-1, m}(\Omega))} \leq C,$$

$$(82) \quad \|\partial_t f(S)\|_{L^1(Q_{\mathcal{T}})} \leq C.$$

$$(83) \quad \|\partial_t F(S)\|_{L^\infty(0, \mathcal{T}; L^1(\Omega))} \leq C, \quad F(s) \equiv - \int_s^1 s_1 f'(s_1) ds_1,$$



where  $\gamma$  and  $\beta$  were given by (22), (37) and

$$m \equiv \min \left\{ \frac{2\beta}{\beta+1}, \frac{6\beta\gamma}{\gamma+6\beta+3\beta\gamma} \right\} > 1.$$

The following proposition gives the bound of the total entropy  $\Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s$  and for the entropy flux.

**Proposition 6.** *Any smooth solution to (11)–(13), (14)–(17) satisfies*

$$(84) \quad \|\Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_{\mathcal{T}})} \leq C.$$

4

$$(85) \quad \exists s > 1 : \quad \left\| (\rho\eta)\mathbf{v} - \sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i + \frac{\mathbf{q}}{T} \right\|_{L^s(Q_{\mathcal{T}})} \leq C.$$

Finally, we derive the gradient bound for  $\rho$ , the bound for  $\log \rho_i/\rho$  and the bound for the reaction terms  $r_i$ ,  $i = 1, \dots, N$ .

**Proposition 7.** *Any smooth solution to (11)–(13), (14)–(17) satisfies*

$$(86) \quad \|\nabla[\sqrt{k_r(S)}G(\rho^\gamma)]\|_{L^{a_2}(0,\mathcal{T};L^q(\Omega))} \leq C[G], \quad \forall G \in W^{1,\infty}(\mathbb{R}_+).$$

8

$$(87) \quad \left\| \log \frac{\rho_i}{\rho} \right\|_{L^2(Q_{\mathcal{T}})} \leq C, \quad i = 1, \dots, N.$$

9

$$(88) \quad \|r_i\|_{L^{\frac{a}{a-1}}(Q_{\mathcal{T}})} \leq C, \quad i = 1, \dots, N.$$

### 10 3. WEAK SEQUENTIAL STABILITY FOR SMOOTH SOLUTIONS

The concept of weak sequential stability in the sense of Feireisl [16], together with weak compactness results, constitute an important step in the global existence analysis. Generally speaking, weak sequential stability means that, given a sequence of (smooth) solutions to a PDE system of interest, there exists a subsequence which converges to a (weak) solution to the problem. Weak stability implies that the limiting solution is a solution to the original system.

In our case, the sequence of solutions solves an approximate and regularized system of equations, obtained from the original one by a semidiscretization in time, and the index of the sequence is related to the approximation and regularization parameters. In order to prove weak compactness, the Div-Curl lemma [17, Prop. 3.3], developed by Murat [30] and Tartar [32], has been used, which represents an efficient tool for handling compactness in nonlinear problems, where the classical Rellich–Kondrashev argument is not applicable.

In this section we assume that a sequence  $(\bar{\rho}^{(n)}, T^{(n)}, S^{(n)})$  of variational entropy solutions to (11)–(13), (14), (17) exists, and show that  $(\bar{\rho}^{(n)}, T^{(n)}, S^{(n)})$  converges (up to subsequences) to another variational entropy solution  $(\bar{\rho}, T, S)$  of (11)–(13), (14), (17) as  $n \rightarrow \infty$ .

**Theorem 8 (Weak stability).** *Let  $(\bar{\rho}^{(n)}, T^{(n)}, S^{(n)})$  be a sequence of variational entropy solutions to (11)–(13), (14), (17) according to Def. 1. Assume furthermore that  $\rho_i^{(n)}, T^{(n)}, S^{(n)} > 0$  a.e. in  $Q_{\mathcal{T}}$  for  $i = 1, \dots, N$ ,  $n \in \mathbb{N}$ . Then  $(\bar{\rho}^{(n)}, T^{(n)}, S^{(n)})$  is (up to subsequences) strongly convergent in  $L^1(Q_{\mathcal{T}})$  (as  $n \rightarrow \infty$ ) to some function triplet  $(\bar{\rho}, T, S)$  which is a variational entropy solution to (11)–(13), (14), (17) according to Def. 1.*

We shall prove Theorem 8 in several steps described in the following subsections.

1 **3.1. Convergence results.** In the following lemma we will collect the strong convergence results,  
 2  $\forall \varepsilon > 0$ , needed for passing to the limit in (45)–(46)–(47), when  $n \rightarrow \infty$ .

3 **Lemma 9.** *The following strong convergence results hold:*

$$(89) \quad S^{(n)} \rightarrow S \quad \text{strongly in } L^r(Q_{\mathcal{T}}), \quad \forall r < \infty,$$

4 *where*

$$(90) \quad S > 0 \quad \text{a.e. in } Q_{\mathcal{T}}.$$

5

$$(91) \quad p^{(n)} \rightarrow p \quad \text{strongly in } L^{4/3}(\Omega \times [\varepsilon, \mathcal{T}]) \text{ for every } \varepsilon > 0,$$

6

$$(92) \quad S^{(n)} p^{(n)} \rightarrow Sp \quad \text{strongly in } L^1(Q_{\mathcal{T}}).$$

7

$$(93) \quad \rho^{(n)} \rightarrow \rho \quad \text{strongly in } L^{\gamma-\varepsilon}(Q_{\mathcal{T}}),$$

8

$$(94) \quad \rho_i^{(n)} \rightarrow \rho_i \quad \text{strongly in } L^{\gamma-\varepsilon}(Q_{\mathcal{T}}), \quad i = 1, \dots, N,$$

$$(95) \quad S^{(n)} \rho_i^{(n)} \log \rho_i^{(n)} \rightarrow S \rho_i \log \rho_i, \quad S^{(n)} \rho_i^{(n)} \rightarrow S \rho_i, \quad i = 1, \dots, N, \\ \text{strongly in } L^{\gamma-\varepsilon}(Q_{\mathcal{T}}),$$

$$(96) \quad \log T^{(n)} \rightarrow \log T \quad \text{strongly in } L^{2-\varepsilon}(0, \mathcal{T}; L^{6-\varepsilon}(\Omega)),$$

$$(97) \quad T^{(n)} \rightarrow T \quad \text{strongly in } L^{\beta+\frac{2}{3}-\varepsilon}(Q_{\mathcal{T}}).$$

9 *Proof.* This proof will be divided into 4 steps. In Step 1 we will prove (89), (90) and (92). In Step  
 10 2 we will prove (93). Step 3 will be about proving (94) and (95), and finally Step 4 proves (96)  
 11 and (97). Let us define preliminarily the following vector fields:

$$(98) \quad U_i^{(n)} = (\Phi S^{(n)} \rho_i^{(n)}, \rho_i^{(n)} \mathbf{v}^{(n)} + \mathbf{J}_i^{(n)}), \quad i = 1, \dots, N,$$

12

$$(99) \quad U^{(n)} = \sum_{i=1}^N U_i^{(n)} = (\Phi S^{(n)} \rho^{(n)}, \rho^{(n)} \mathbf{v}^{(n)}),$$

13

$$(100) \quad V^{(n)}[G] = (\sqrt{k_r(S^{(n)})} G((\rho^{(n)})^\gamma), 0, 0, 0), \quad G \in W^{1,\infty}(\mathbb{R}_+),$$

14

$$(101) \quad Z^{(n)}[G] = (G(\Pi^N \bar{\mu}^{(n)}/T^{(n)}), 0, 0, 0), \quad G \in W^{1,\infty}(\mathbb{R}^N),$$

15

$$(102) \quad W^{(n)} = (\Phi S^{(n)} (\rho \eta)^{(n)} + (1 - \Phi) (\rho \eta)_s^{(n)}, (\rho \eta)^{(n)} \mathbf{v}^{(n)} - \sum_{i=1}^N \frac{\mu_i^{(n)}}{T^{(n)}} \mathbf{J}_i^{(n)} + \frac{\mathbf{q}^{(n)}}{T^{(n)}}),$$

16

$$(103) \quad Y^{(n)}[G] = (G(T^{(n)}), 0, 0, 0), \quad G \in W^{1,\infty}(\mathbb{R}_+).$$

**Step 1: Strong convergence, a.e. positivity of saturation and convergence of  $S^{(n)}p^{(n)}$ .**

In this step we show (89)–(92). From (68), (82) it follows

$$\|\partial_t f(S^{(n)})\|_{L^1(0,\mathcal{T};L^1(\Omega))} + \|f(S^{(n)})\|_{L^\infty(0,\mathcal{T};W^{1,q}(\Omega))} \leq C.$$

Since  $W^{1,q}(\Omega) \hookrightarrow L^q(\Omega)$  compactly and  $L^q(\Omega) \hookrightarrow L^1(\Omega)$  continuously, the Aubin-Lions Lemma allows us to deduce that, up to subsequences,

$$f(S^{(n)}) \text{ is strongly convergent in } L^q(Q_{\mathcal{T}}) \text{ as } n \rightarrow \infty.$$

- 1 In particular  $f(S^{(n)})$  is a.e. convergent in  $Q_{\mathcal{T}}$ . Since  $f$  is strictly decreasing (and a fortiori one-
- 2 to-one), it follows that  $S^{(n)}$  is a.e. convergent in  $Q_{\mathcal{T}}$ . Thanks to the uniform  $L^\infty(Q_{\mathcal{T}})$  bounds for
- 3  $S^{(n)}$ , we deduce (89).

We now prove (90). Since  $p^{(n)} \geq -p_{at}$  and (33) holds, it follows

$$\partial_t f(S^{(n)}) + \Gamma f(S^{(n)}) \leq p_{at}, \quad \Gamma \equiv \inf_{0 < s < s_0} \frac{P_c(s)}{f(s)} > \frac{p_{at}}{f(0)}.$$

A Gronwall argument yields

$$f(S^{(n)}(t)) \leq f(S^{in})e^{-\Gamma t} + \frac{p_{at}}{\Gamma}(1 - e^{-\Gamma t}), \quad t > 0, \text{ a.e. in } \Omega.$$

Relation (89) allows us to take the limit  $n \rightarrow \infty$  in the above inequality and obtain

$$(104) \quad f(S(t)) \leq f(S^{in})e^{-\Gamma t} + \frac{p_{at}}{\Gamma}(1 - e^{-\Gamma t}), \quad t > 0, \text{ a.e. in } \Omega.$$

Consider the set  $E := \{x \in \Omega : f(S^{in}(x)) \geq p_{at}/\Gamma\}$ . Clearly  $\{S^{in} = 0\} \subset E$  since  $\Gamma > p_{at}/f(0)$ . Moreover

$$f(S(t)) \leq \frac{p_{at}}{\Gamma}, \quad t > 0, \quad \text{a.e. in } \Omega \setminus E,$$

which means that

$$S(t) \geq f^{-1}\left(\frac{p_{at}}{\lambda}\right) > 0, \quad t > 0, \quad \text{a.e. in } \Omega \setminus E.$$

On the other hand, the right-hand side of (104) is decreasing in time for  $x \in E$ , so

$$\forall \varepsilon > 0, \quad f(S(t)) \leq f(S^{in})e^{-\lambda\varepsilon} + \frac{p_{at}}{\lambda}(1 - e^{-\lambda\varepsilon}), \quad t \geq \varepsilon, \text{ a.e. in } E.$$

But  $f(S^{in}(x)) \leq f(0)$ , so

$$\forall \varepsilon > 0, \quad S(t) \geq \tilde{s}_\varepsilon \equiv f^{-1}\left(f(0)e^{-\lambda\varepsilon} + \frac{p_{at}}{\lambda}(1 - e^{-\lambda\varepsilon})\right), \quad t \geq \varepsilon, \text{ a.e. in } E.$$

- 4 Clearly  $\tilde{s}_\varepsilon > 0$  for every  $\varepsilon > 0$ . This fact, combined with the uniform positivity of  $S$  on  $(\Omega \setminus E) \times$
- 5  $[0, \mathcal{T}]$ , allows us to deduce

$$(105) \quad \forall \varepsilon > 0 \quad \exists s_\varepsilon > 0 : \quad S(t) \geq s_\varepsilon, \quad t \geq \varepsilon, \quad \text{a.e. in } \Omega.$$

In particular  $S > 0$  a.e. in  $Q_{\mathcal{T}}$  and thus (90) holds. Furthermore, it is immediate to see from the argument above that (105) also holds for every term of the sequence  $S^{(n)}$ ,  $n \in \mathbb{N}$ .

The strong convergence of the saturation (89) and its positivity (90) will help us to show (92). Let us define  $Q_{\mathcal{T}}^\varepsilon = \Omega \times [\varepsilon, \mathcal{T}]$ . From (5), (65), (105) as well as the uniform boundedness of  $\mu$  it follows

$$\forall \varepsilon > 0, \quad \exists C_\varepsilon > 0 : \quad \|(T^{(n)})^{-1/2} \nabla p^{(n)}\|_{L^2(Q_{\mathcal{T}}^\varepsilon)} \leq C_\varepsilon.$$

From the above estimate, (63) and the fact that  $\beta > 2$  we obtain via Hölder's inequality

$$\forall \varepsilon > 0, \quad \exists C_\varepsilon > 0 : \quad \|\nabla p^{(n)}\|_{L^{16/11}(Q_{\mathcal{T}}^\varepsilon)} \leq \|(T^{(n)})^{1/2}\|_{L^{16/3}(Q_{\mathcal{T}}^\varepsilon)} \|(T^{(n)})^{-1/2} \nabla p^{(n)}\|_{L^2(Q_{\mathcal{T}}^\varepsilon)} \leq C_\varepsilon.$$

Via Poincaré Lemma we get

$$\forall \varepsilon > 0, \quad \exists C_\varepsilon > 0 : \quad \left\| p^{(n)} - |\Omega|^{-1} \int_{\Omega} p^{(n)} dx \right\|_{L^{16/11}(Q_{\mathcal{T}}^\varepsilon)} \leq C_\varepsilon.$$

On the other hand, (105), (61) yield

$$\left| \int_{\Omega} p^{(n)}(t) dx \right| \leq \frac{1}{s_\varepsilon} \int_{\Omega} S^{(n)}(t) |p^{(n)}(t)| dx \leq \frac{C}{s_\varepsilon}, \quad t \geq \varepsilon,$$

which means

$$(106) \quad \forall \varepsilon > 0, \quad \exists C_\varepsilon > 0 : \quad \|p^{(n)}\|_{L^{16/11}(Q_{\mathcal{T}}^\varepsilon)} \leq C_\varepsilon.$$

Since  $p^{(n)} \rightarrow p$  a.e. in  $Q_{\mathcal{T}}$ , the Vitali-De la Vallée Poussin Theorem [17, Corollary 10.2] implies (91). Given that (89), (91) hold, it follows

$$S^{(n)} p^{(n)} \rightarrow Sp \quad \text{strongly in } L^1(Q_{\mathcal{T}}^\varepsilon), \quad \forall \varepsilon > 0.$$

However,

$$\|S^{(n)} p^{(n)}\|_{L^1(0,\varepsilon;L^1(\Omega))} \leq \varepsilon \|S^{(n)} p^{(n)}\|_{L^\infty(0,\varepsilon;L^1(\Omega))} \leq C\varepsilon,$$

and a similar estimate holds for  $Sp$  (e.g. via Fatou's Lemma). It follows that

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \|S^{(n)} p^{(n)} - Sp\|_{L^1(Q_{\mathcal{T}})} \\ & \leq \limsup_{n \rightarrow \infty} \|S^{(n)} p^{(n)} - Sp\|_{L^1(Q_{\mathcal{T}}^\varepsilon)} + \limsup_{n \rightarrow \infty} \|S^{(n)} p^{(n)} - Sp\|_{L^1(Q_{\mathcal{T}} \setminus Q_{\mathcal{T}}^\varepsilon)} \\ & \leq \sup_{n \in \mathbb{N}} (\|S^{(n)} p^{(n)}\|_{L^1(0,\varepsilon;L^1(\Omega))} + \|Sp\|_{L^1(0,\varepsilon;L^1(\Omega))}) \leq C\varepsilon. \end{aligned}$$

1 This means that (92) holds.

**Step 2: Strong convergence of the total density.** In this step we show (93).

Since (60) holds, it follows that (up to subsequences)

$$\rho^{(n)} \rightharpoonup \rho \quad \text{weakly in } L^\gamma(Q_{\mathcal{T}}).$$

Let us consider the vector fields  $U^{(n)}$ ,  $V^{(n)}[G]$  defined in (99), (100), with  $G \in W^{1,\infty}(\mathbb{R}_+)$  arbitrary. Thanks to (61), (80), we deduce that  $U^{(n)}$  is bounded in  $L^m(Q_{\mathcal{T}})$  with  $m > 1$ , while  $V^{(n)}[G]$  is trivially bounded in  $L^\infty(Q_{\mathcal{T}})$ . On the other hand, summing (11) in  $i = 1, \dots, N$  yields  $\operatorname{div}_{(t,x)} U^{(n)} = 0$ , while the antisymmetric part of the Jacobian of  $V^{(n)}[G]$ , which we denote by  $\operatorname{curl}_{(t,x)} V^{(n)}[G] = \nabla V^{(n)}[G] - \nabla^T V^{(n)}[G]$ , satisfies

$$|\operatorname{curl}_{(t,x)} V^{(n)}[G]| \leq C |\nabla [\sqrt{k_r(S^{(n)})} G ((\rho^{(n)})^\gamma)]|$$

and is therefore bounded in  $L^1(Q_{\mathcal{T}})$  thanks to (86). In particular both  $\operatorname{div}_{(t,x)} U^{(n)}$ ,  $\operatorname{curl}_{(t,x)} V^{(n)}[G]$  are relatively compact in  $W^{-1,z}(Q_{\mathcal{T}})$  for some  $z > 1$ . Therefore we can apply the Div-Curl Lemma [17, Prop. 3.3] and deduce

$$\overline{U^{(n)} \cdot V^{(n)}[G]} = \overline{U^{(n)}} \cdot \overline{V^{(n)}[G]},$$

where  $\overline{F(U^{(n)})}$  denotes a weak  $L^1$ -limit of the sequence  $(F(U^{(n)}))_{n \in \mathbb{N}}$ , meaning that  $\overline{F(U^{(n)})} \in L^1(\Omega)$  is defined by

$$\int_{\Omega} \overline{F(U^{(n)})} \phi dx := \lim_{n \rightarrow \infty} \int_{\Omega} F(U^{(n)}) \phi dx \quad \text{for all } \phi \in L^\infty(\Omega).$$

This is equivalent to

$$\overline{\Phi S^{(n)} \rho^{(n)} \sqrt{k_r(S^{(n)}) G((\rho^{(n)})^\gamma)}} = \overline{\Phi S^{(n)} \rho^{(n)} \sqrt{k_r(S^{(n)}) G((\rho^{(n)})^\gamma)}}.$$

However, since  $\Phi > 0$  and does not depend on  $n$ , while  $S^{(n)} \rightarrow S$  strongly in  $L^p(Q_T)$  for every  $p < \infty$ , it follows

$$S \sqrt{k_r(S)} \overline{\rho^{(n)} G((\rho^{(n)})^\gamma)} = S \sqrt{k_r(S)} \overline{\rho G((\rho^{(n)})^\gamma)}.$$

It holds  $S \sqrt{k_r(S)} > 0$  because  $S > 0$  a.e. in  $Q_T$ . We deduce

$$(107) \quad \overline{\rho^{(n)} G((\rho^{(n)})^\gamma)} = \overline{\rho G((\rho^{(n)})^\gamma)} \quad \text{on } Q_T, \quad \forall G \in W^{1,\infty}(\mathbb{R}_+).$$

Let us now choose  $G(s) = G_k(s) \equiv \min\{(1+s)^{1/\gamma}, (1+k)^{1/\gamma}\}$ , for  $s \geq 0$ ,  $k \in \mathbb{N}$ . Define also  $G_\infty(s) = (1+s)^{1/\gamma}$ . The weak lower semicontinuity of the  $L^1$  norm and Cauchy-Schwartz inequality allow us to estimate

$$\begin{aligned} & \|\overline{\rho^{(n)} [G_k((\rho^{(n)})^\gamma) - G_\infty((\rho^{(n)})^\gamma)]}\|_{L^1(Q_T)} \\ & \leq \liminf_{n \rightarrow \infty} \|\rho^{(n)} [G_k((\rho^{(n)})^\gamma) - G_\infty((\rho^{(n)})^\gamma)]\|_{L^1(Q_T)} \\ & \leq \sup_{n \in \mathbb{N}} \|\rho^{(n)}\|_{L^2(Q_T)} \|G_k((\rho^{(n)})^\gamma) - G_\infty((\rho^{(n)})^\gamma)\|_{L^2(Q_T)}. \end{aligned}$$

Thanks to (60), it follows (remember that  $\gamma > 2$ ):

$$\begin{aligned} & \|\overline{\rho^{(n)} [G_k((\rho^{(n)})^\gamma) - G_\infty((\rho^{(n)})^\gamma)]}\|_{L^1(Q_T)}^2 \\ & \leq C \sup_{n \in \mathbb{N}} \int_{Q_T \cap \{\rho^{(n)} > k^{1/\gamma}\}} (1 + (\rho^{(n)})^\gamma)^{2/\gamma} dx dt \\ & \leq C k^{2/\gamma-1} \sup_{n \in \mathbb{N}} \int_{Q_T \cap \{\rho^{(n)} > k^{1/\gamma}\}} (\rho^{(n)})^{\gamma-2} (1 + (\rho^{(n)})^\gamma)^{2/\gamma} dx dt \\ & \leq C k^{2/\gamma-1} \sup_{n \in \mathbb{N}} \|\rho^{(n)}\|_{L^\gamma(Q_T)}^\gamma \leq C k^{2/\gamma-1}. \end{aligned}$$

In a similar way one shows that

$$\|\overline{\rho G_k((\rho^{(n)})^\gamma) - G_\infty((\rho^{(n)})^\gamma)}\|_{L^1(Q_T)}^2 \leq C k^{2/\gamma-1}.$$

From the above inequalities and (107) with  $G = G_k$  it follows

$$\begin{aligned} & \|\overline{\rho^{(n)} G_\infty((\rho^{(n)})^\gamma)} - \overline{\rho G_\infty((\rho^{(n)})^\gamma)}\|_{L^1(Q_T)} \\ & \leq \|\overline{\rho^{(n)} [G_k((\rho^{(n)})^\gamma) - G_\infty((\rho^{(n)})^\gamma)]}\|_{L^1(Q_T)} \\ & \quad + \|\overline{\rho G_k((\rho^{(n)})^\gamma) - G_\infty((\rho^{(n)})^\gamma)}\|_{L^1(Q_T)} \leq C k^{1/\gamma-1/2} \rightarrow 0 \quad \text{as } k \rightarrow \infty, \end{aligned}$$

implying that (recall that  $G_\infty(s) = (1+s)^{1/\gamma}$ )

$$(108) \quad \overline{\rho^{(n)} (1 + (\rho^{(n)})^\gamma)^{1/\gamma}} = \overline{\rho (1 + (\rho^{(n)})^\gamma)^{1/\gamma}} \quad \text{on } Q_T.$$

- 1 Since  $s \in \mathbb{R}_+ \mapsto (1+s^\gamma)^{1/\gamma} \in \mathbb{R}_+$  is strictly increasing and strictly convex, we conclude [17,
- 2 Thr. 10.19, Thr. 10.20] that, up to subsequences,  $\rho^{(n)}$  is a.e. convergent in  $Q_T$ . Since  $\rho^{(n)}$  is
- 3 bounded in  $L^\gamma(Q_T)$ , Vitali - De la Vallée Poussin theorem implies that (93) holds.

### **Step 3: Strong convergence of partial densities.**

In this step we show (94) and (95). For every  $i = 1, \dots, N$ , the vector field  $U_i^{(n)}$  defined in (98) is bounded in  $L^m(Q_T)$  thanks to (80), while its time-space divergence  $\operatorname{div}_{(t,x)} U_i^{(n)} = r_i^{(n)}$  is bounded in  $L^{a/(a-1)}(Q_T)$  due to (88), and a fortiori relatively compact in  $W^{-1,p}(Q_T)$  for some  $p > 1$ . On the

other hand, for every  $G \in W^{1,\infty}(\mathbb{R}^N)$ , the vector field  $Z^{(n)}[G]$  defined in (101) is trivially bounded in  $L^\infty(Q_{\mathcal{T}})$ , while (64) implies

$$\|\operatorname{curl}_{(t,x)} Z^{(n)}[G]\|_{L^2(Q_{\mathcal{T}})} \leq C[G] \|\nabla \Pi^N \vec{\mu}^{(n)}/T^{(n)}\|_{L^2(Q_{\mathcal{T}})} \leq C[G].$$

In particular  $\operatorname{curl}_{(t,x)} Z^{(n)}[G]$  is relatively compact in  $H^{-1}(Q_{\mathcal{T}})$ . Therefore we can apply the Div-Curl Lemma with the vector fields  $U_i^{(n)}$ ,  $Z^{(n)}$  and obtain

$$\overline{U_i^{(n)} \cdot Z^{(n)}} = \overline{U_i^{(n)}} \cdot \overline{Z^{(n)}}, \quad i = 1, \dots, N,$$

which means

$$\overline{\Phi S^{(n)} \rho_i^{(n)} G(\Pi^N \vec{\mu}^{(n)}/T^{(n)})} = \overline{\Phi S^{(n)} \rho_i^{(n)}} \overline{G(\Pi^N \vec{\mu}^{(n)}/T^{(n)})}, \quad i = 1, \dots, N.$$

Once again, the strong convergence of  $S^{(n)}$  and the weak convergence of  $\rho_i^{(n)}$ ,  $G(\Pi^N \vec{\mu}^{(n)}/T^{(n)})$  (as well as the positivity of  $\Phi$ ) imply

$$(109) \quad \overline{\rho_i^{(n)} G(\Pi^N \vec{\mu}^{(n)}/T^{(n)})} = \overline{\rho_i} \overline{G(\Pi^N \vec{\mu}^{(n)}/T^{(n)})}, \quad i = 1, \dots, N, \quad \text{on } Q_{\mathcal{T}},$$

1 for every  $G \in W^{1,\infty}(\mathbb{R}^N)$ . However, from (23), (40) it follows

$$(110) \quad (\Pi^N \vec{\mu}^{(n)}/T^{(n)})_i = \log \rho_i^{(n)} - \frac{1}{N} \sum_{j=1}^N \log \rho_j^{(n)}, \quad i = 1, \dots, N,$$

so (109) becomes

$$\overline{\rho_i^{(n)} G \left( \log \rho_i^{(n)} - \frac{1}{N} \sum_{j=1}^N \log \rho_j^{(n)} \right)} = \overline{\rho_i} \overline{G \left( \log \rho_i^{(n)} - \frac{1}{N} \sum_{j=1}^N \log \rho_j^{(n)} \right)}, \quad i = 1, \dots, N, \quad \text{on } Q_{\mathcal{T}}.$$

Let  $k \in \mathbb{N}$  arbitrary. Define  $Q_{\mathcal{T},k} \equiv \{(x,t) \in Q_{\mathcal{T}} : \rho(x,t) \geq 1/k\}$ . Since  $\rho^{(n)} \rightarrow \rho$  strongly in  $L^1(Q_{\mathcal{T}})$ , by Egorov-Severini's theorem  $\rho^{(n)} \rightarrow \rho$  almost uniformly in  $Q_{\mathcal{T},k}$ , that is

$$\forall \epsilon > 0 \quad \exists E_{\epsilon,k} \subset Q_{\mathcal{T},k} : \quad \rho^{(n)} \rightarrow \rho \text{ strongly in } L^\infty(E_{\epsilon,k}), \quad |Q_{\mathcal{T},k} \setminus E_{\epsilon,k}| < \epsilon.$$

We can therefore assume w.l.o.g. that  $\rho^{(n)} \geq 1/2k$  a.e. in  $E_{\epsilon,k}$ ,  $n \in \mathbb{N}$ . Defining  $\sigma_i^{(n)} = \rho_i^{(n)}/\rho^{(n)}$  on  $E_{\epsilon,k}$ , for  $i = 1, \dots, N$ , allows us to write

$$\sum_{i=1}^N \overline{\rho_i^{(n)} G \left( \log \sigma_i^{(n)} - \frac{1}{N} \sum_{j=1}^N \log \sigma_j^{(n)} \right)} = \sum_{i=1}^N \overline{\rho_i} \overline{G \left( \log \sigma_i^{(n)} - \frac{1}{N} \sum_{j=1}^N \log \sigma_j^{(n)} \right)} \quad \text{on } E_{\epsilon,k}.$$

Fix  $i \in \{1, \dots, N\}$  generic. By choosing  $G(\mathbf{u}) = G_M(u_i) = \min\{u_i, M\}$  in the above identity, with  $M \in \mathbb{N}$  generic, exploiting (64) and proceeding like in the proof of (108) one finds out that

$$\sum_{i=1}^N \overline{\rho_i^{(n)} \left( \log \sigma_i^{(n)} - \frac{1}{N} \sum_{j=1}^N \log \sigma_j^{(n)} \right)} = \sum_{i=1}^N \overline{\rho_i} \overline{\left( \log \sigma_i^{(n)} - \frac{1}{N} \sum_{j=1}^N \log \sigma_j^{(n)} \right)} \quad \text{on } E_{\epsilon,k},$$

which, thanks to (93) and (87), leads to

$$\sum_{i=1}^N \overline{\rho_i^{(n)} \log \sigma_i^{(n)}} = \sum_{i=1}^N \overline{\rho_i} \overline{\log \sigma_i^{(n)}} \quad \text{on } E_{\epsilon,k}.$$

Once again, (93) and (87) imply

$$\sum_{i=1}^N \overline{\sigma_i^{(n)} \log \sigma_i^{(n)}} = \sum_{i=1}^N \overline{\sigma_i^{(n)}} \overline{\log \sigma_i^{(n)}} \quad \text{on } E_{\epsilon,k}.$$

1 The fact that  $\log$  is strictly monotone and strictly concave allows us to conclude that (up to  
2 subsequences)

$\sigma_i^{(n)} = \rho_i^{(n)}/\rho^{(n)}$  is a.e. convergent on  $E_{\epsilon,k}$ , which, together with (93), implies that  $\rho_i^{(n)} \rightarrow \rho_i$   
a.e. in  $E_{\epsilon,k}$ . Since  $|Q_{\mathcal{T},k} \setminus E_{\epsilon,k}| < \epsilon$  and (60) holds, we easily deduce that  $\rho_i^{(n)} \rightarrow \rho_i$  strongly in  
 $L^1(Q_{\mathcal{T},k})$  for every  $k \in \mathbb{N}$ . Since  $0 \leq \rho \leq 1/k$  on  $Q_{\mathcal{T}} \setminus Q_{\mathcal{T},k}$ , it follows that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \int_{Q_{\mathcal{T}}} |\rho_i^{(n)} - \rho_i| dx dt &\leq \limsup_{n \rightarrow \infty} \int_{Q_{\mathcal{T},k}} |\rho_i^{(n)} - \rho_i| dx dt + \limsup_{n \in \mathbb{N}} \int_{Q_{\mathcal{T}} \setminus Q_{\mathcal{T},k}} |\rho_i^{(n)} - \rho_i| dx dt \\ &\leq 2 \int_{Q_{\mathcal{T}} \setminus Q_{\mathcal{T},k}} \rho dx dt \leq \frac{2}{k} |Q_{\mathcal{T}}|, \end{aligned}$$

3 and therefore  $\rho_i^{(n)} \rightarrow \rho_i$  strongly in  $L^1(Q_{\mathcal{T}})$ , for  $i = 1, \dots, N$ .

4 Together with (60), we conclude (94). Moreover, (95) holds as it is a straightforward consequence  
5 of (89), (94) and Vitali - De la Vallée Poussin theorem.

**Step 4: Strong convergence of the temperature.**

Finally, we show (96) and (97). Let us consider the vector fields  $W^{(n)}, Y^{(n)}$  defined in (102), (103).  
Thanks to (84), (85) we deduce that  $W^{(n)}$  is bounded in  $L^s(Q_{\mathcal{T}})$  for some  $s > 1$ , while  $Y^{(n)}[G]$   
is bounded in  $L^\infty(Q_{\mathcal{T}})$  for every  $G \in W^{1,\infty}(\mathbb{R}_+)$ . On the other hand, the time-space divergence  
 $\operatorname{div}_{(t,x)} W^{(n)}$  is the right-hand side of (51), which is bounded in  $L^1(Q_{\mathcal{T}})$  thanks to Lemma 4, while

$$|\operatorname{curl}_{(t,x)} Y^{(n)}[G]| \leq C[G] |\nabla T|$$

is bounded in  $L^1(Q_{\mathcal{T}})$  due to (62). In order to see this, we distinguish two cases:

*Case*  $1 < \beta \leq 2$ . In this case it holds  $\nabla T^{(n)} = \frac{2}{\beta} (T^{(n)})^{1-\beta/2} \nabla (T^{(n)})^{\beta/2}$ , which, given that  $0 \leq$   
 $1 - \beta/2 < 1/2$ , implies that  $\nabla T^{(n)}$  is bounded in  $L^1(Q_{\mathcal{T}})$  via the uniform bounds for  $T^{(n)}$  and  
 $\nabla (T^{(n)})^{\beta/2}$  in  $L^\infty(0, T; L^1(\Omega))$  and  $L^2(Q_{\mathcal{T}})$ , respectively.

*Case*  $\beta > 2$ . In this case we observe that

$$\nabla T^{(n)} = \chi_{(0,1)}(T^{(n)}) T^{(n)} \nabla \log T^{(n)} + \chi_{[1,\infty)}(T^{(n)}) \frac{2}{\beta} (T^{(n)})^{1-\beta/2} \nabla (T^{(n)})^{\beta/2}$$

is bounded in  $L^2(Q_{\mathcal{T}})$  given the uniform bounds for  $\nabla \log T^{(n)}$  and  $\nabla (T^{(n)})^{\beta/2}$  in  $L^2(Q_{\mathcal{T}})$ . In  
particular both  $\operatorname{div}_{(t,x)} W^{(n)}$ ,  $\operatorname{curl}_{(t,x)} Y^{(n)}[G]$  are relatively compact in  $W^{-1,r}(Q_{\mathcal{T}})$  for some  $r > 1$ .  
From the Div-Curl Lemma it follows

$$\overline{W^{(n)} \cdot Y^{(n)}[G]} = \overline{W^{(n)}} \cdot \overline{Y^{(n)}[G]}$$

which means

$$\begin{aligned} \overline{(\Phi S^{(n)}(\rho\eta)^{(n)} + (1 - \Phi)(\rho\eta)_s^{(n)}) G(T^{(n)})} &= \overline{(\Phi S^{(n)}(\rho\eta)^{(n)} + (1 - \Phi)(\rho\eta)_s^{(n)})} \overline{G(T^{(n)})} \\ &\text{a.e. in } Q_{\mathcal{T}}, \quad \forall G \in W^{1,\infty}(\mathbb{R}_+). \end{aligned}$$

From the definitions (26), (28), bound (62) and strong convergence relations (95) we deduce

$$(c_w \Phi S \rho + c_s(1 - \Phi)) \overline{(\log T^{(n)}) G(T^{(n)})} = (c_w \Phi S \rho + c_s(1 - \Phi)) \overline{(\log T^{(n)})} \overline{G(T^{(n)})},$$

which, thanks to (4), yields

$$\overline{(\log T^{(n)}) G(T^{(n)})} = \overline{(\log T^{(n)})} \overline{G(T^{(n)})} \quad \text{a.e. in } Q_{\mathcal{T}}, \quad \forall G \in W^{1,\infty}(\mathbb{R}_+).$$

6 Since (62) holds, by arguing in a similar way as the derivation of (108) one obtains

$$(111) \quad \overline{(\log T^{(n)}) T^{(n)}} = \overline{(\log T^{(n)})} \overline{T^{(n)}} \quad \text{a.e. in } Q_{\mathcal{T}}.$$

1 Once again, the strict monotonicity and strict convexity of  $\log$  allows us to conclude that  $T^{(n)}$  is  
 2 (up to subsequences) a.e. convergent in  $Q_{\mathcal{T}}$ . From this fact, (62), (63) and Sobolev's embedding  
 3  $H^1(\Omega) \hookrightarrow L^6(\Omega)$  it follows (96) and (97).  $\square$

4 **3.2. Passing to the limit in the equations (45)–(47), when  $n \rightarrow \infty$ .** In this subsection we will  
 5 use the previous convergence results for passing to the limit in the variational entropy formulation.

*Limit in equation (45).* From (78), (79), (81), (95) it follows

$$\begin{aligned} \Phi \partial_t(S^{(n)} \rho_i^{(n)}) \rightharpoonup^* \Phi \partial_t(S \rho_i) \quad \text{weakly}^* \text{ in } L^m(0, \mathcal{T}; W^{-1,m}(\Omega)), \\ \mathbf{J}_i^{(n)} \rightharpoonup \mathbf{J}_i \quad \text{weakly in } L^2(Q_{\mathcal{T}}), \end{aligned}$$

for  $i = 1, \dots, N$ . Thanks to (6), (65),  $(\lambda(S^{(n)}, T^{(n)})T^{(n)})^{-1/2} \mathbf{v}^{(n)}$  is bounded in  $L^2(Q_{\mathcal{T}})$ ; however, eq. (5), the boundedness of  $k_r \in C^0([0, 1])$  and the uniform positivity of  $\mu$  (see assumptions on phase velocity) imply that  $\lambda(S^{(n)}, T^{(n)}) = k_r(S^{(n)})/\mu(T^{(n)})$  is bounded in  $L^\infty(Q_{\mathcal{T}})$ , yielding that  $(T^{(n)})^{-1/2} \mathbf{v}^{(n)}$  is bounded in  $L^2(Q_{\mathcal{T}})$ . So, from (63) we deduce that  $\mathbf{v}^{(n)}$  is bounded in  $L^{\frac{6\beta+4}{3\beta+5}}(Q_{\mathcal{T}})$ . As a consequence,

$$\mathbf{v}^{(n)} \rightharpoonup \mathbf{v} \quad \text{weakly in } L^{\frac{6\beta+4}{3\beta+5}}(Q_{\mathcal{T}}).$$

Since (94) holds and (thanks to (37))  $\frac{1}{\gamma} + \frac{3\beta+5}{6\beta+4} < 1$ , it follows

$$\rho_i^{(n)} \mathbf{v}^{(n)} \rightharpoonup \rho_i \mathbf{v} \quad \text{weakly in } L^s(Q_{\mathcal{T}}) \text{ for some } s > 1.$$

We must now identify  $\mathbf{v}$ . We start by pointing out that (105) easily holds for  $S^{(n)}$ , too (just remake the same computations). Therefore (6), (5) lead to

$$-K \nabla p^{(n)} \rightharpoonup \frac{\mathbf{v}}{\lambda(S, T)} \quad \text{weakly in } L^{\frac{6\beta+4}{3\beta+5}-\varepsilon_1}(\Omega \times [\varepsilon_2, \mathcal{T}]), \quad \forall \varepsilon_1, \varepsilon_2 > 0.$$

From (91) it follows

$$-K \lambda(S, T) \nabla p = \mathbf{v} \quad \text{a.e. in } \Omega \times [\varepsilon_2, \mathcal{T}], \quad \forall \varepsilon_2 > 0.$$

6 Being  $\varepsilon_2 > 0$  arbitrary and  $\mathbf{v} \in L^{\frac{6\beta+4}{3\beta+5}}(Q_{\mathcal{T}})$ , we deduce that (6) holds a.e. in  $Q_{\mathcal{T}}$ . As a consequence  
 7 of this fact and (65), we also deduce

$$(112) \quad \frac{\lambda(S^{(n)}, T^{(n)})^{1/2}}{(T^{(n)})^{1/2}} \nabla p^{(n)} \rightharpoonup \frac{\lambda(S, T)^{1/2}}{T^{1/2}} \nabla p \quad \text{weakly in } L^2(Q_{\mathcal{T}}).$$

We will now identify the limits of the diffusion fluxes  $\mathbf{J}_1, \dots, \mathbf{J}_N$ . From (64) it follows

$$(113) \quad \Pi^N \vec{\mu}^{(n)} / T^{(n)} \rightharpoonup \vec{\zeta} \quad \text{weakly in } L^2(0, \mathcal{T}; H^1(\Omega)),$$

which, thanks to (38), (39), yields, for  $i = 1, \dots, N$ :

$$(114) \quad \sum_{j=1}^N \tilde{L}_{ij}(\vec{\rho}^{(n)}, T^{(n)}) \nabla(\mu_j^{(n)} / T^{(n)}) \rightharpoonup \sum_{j=1}^N \tilde{L}_{ij}(\vec{\rho}, T) \nabla \zeta_j \quad \text{weakly in } L^2(Q_{\mathcal{T}}).$$

8 In order to identify  $\vec{\zeta}$ , we point out that

$$(115) \quad \text{For a.e. } (x, t) \in Q_{\mathcal{T}}, \quad \rho(x, t) > 0 \quad \Rightarrow \quad \min_{i=1, \dots, N} \rho_i(x, t) > 0.$$

This is a consequence of (87). In fact, since  $\rho_i^{(n)}$  is a.e. convergent in  $Q_{\mathcal{T}}$ , Fatou's Lemma implies

$$\int_{Q_{\mathcal{T}} \cap \{\rho > 0\}} \log(\rho / \rho_i) dx \leq \liminf_{n \rightarrow \infty} \int_{Q_{\mathcal{T}} \cap \{\rho > 0\}} \log(\rho^{(n)} / \rho_i^{(n)}) dx \leq C,$$



1 for  $i = 1, \dots, N$ . This means that  $\rho_i > 0$  on  $Q_{\mathcal{T}} \cap \{\rho > 0\}$  for  $i = 1, \dots, N$ .

Thanks to the a.e. convergence of  $\rho_i^{(n)}$  and (115), we deduce

$$(116) \quad \begin{aligned} \Pi^N \frac{\vec{\mu}^{(n)}}{T^{(n)}} &\rightarrow \vec{\zeta} = \Pi^N \frac{\vec{\mu}}{T} \quad \text{a.e. on } Q_{\mathcal{T}} \cap \{\rho > 0\}, \\ \left( \Pi^N \frac{\vec{\mu}}{T} \right)_i &= \log \left( \frac{\rho_i}{\rho} \right) \quad \text{a.e. in } Q_{\mathcal{T}} \cap \{\rho > 0\}, \quad i = 1, \dots, N. \end{aligned}$$

From (38), (62), (94), (96) we deduce that, for  $i = 1, \dots, N$ ,

$$(117) \quad \begin{aligned} \frac{\tilde{L}_{i0}(\vec{\rho}^{(n)}, T^{(n)})}{T^{(n)}} &\rightarrow \frac{\tilde{L}_{i0}(\vec{\rho}, T)}{T} \quad \text{strongly in } L^p(Q_{\mathcal{T}}), \quad \forall p < \infty, \\ \nabla \log T^{(n)} &\rightharpoonup \nabla \log T \quad \text{weakly in } L^2(Q_{\mathcal{T}}). \end{aligned}$$

Since  $\frac{\tilde{L}_{i0}(\vec{\rho}^{(n)}, T^{(n)})}{T^{(n)}} \nabla \log T^{(n)}$  is bounded in  $L^2(Q_{\mathcal{T}})$ , we deduce (up to subsequences)

$$\frac{\tilde{L}_{i0}(\vec{\rho}^{(n)}, T^{(n)})}{T^{(n)}} \nabla \log T^{(n)} \rightharpoonup \frac{\tilde{L}_{i0}(\vec{\rho}, T)}{T} \nabla \log T \quad \text{weakly in } L^2(Q_{\mathcal{T}}),$$

for  $i = 1, \dots, N$ , which means

$$(118) \quad \tilde{L}_{i0}(\vec{\rho}^{(n)}, T^{(n)}) \nabla \frac{1}{T^{(n)}} \rightharpoonup \tilde{L}_{i0}(\vec{\rho}, T) \nabla \frac{1}{T} \quad \text{weakly in } L^2(Q_{\mathcal{T}}).$$

From (114), (116), (118) we obtain

$$(119) \quad \begin{aligned} \mathbf{J}_i &= L_{i0} \nabla \left( \frac{1}{T} \right) - \sum_{k=1}^N L_{ik} \nabla \zeta_k, \quad \text{a.e. on } Q_{\mathcal{T}}, \\ L_{ij} &= \tilde{L}_{ij}(\vec{\rho}, T), \quad L_{i0} = \tilde{L}_{i0}(\vec{\rho}, T) \quad \text{a.e. on } Q_{\mathcal{T}}. \end{aligned}$$

for  $i, j = 1, \dots, N$ . Let us now focus on the reaction terms. From (88) it follows that

$$r_i^{(n)} \rightharpoonup r_i \quad \text{weakly in } L^{a/(a-1)}(Q_{\mathcal{T}}), \quad i = 1, \dots, N.$$

2 However, since  $r_i^{(n)} = \tilde{r}_i(\rho^{(n)}, T^{(n)}, \Pi^N \vec{\mu}^{(n)}/T^{(n)})$ , the continuity of  $\tilde{r}_i$  as well as (93), (96), (116) we deduce

$$(120) \quad r_i = \tilde{r}_i(\rho, T, \Pi^N \vec{\mu}/T) \quad \text{a.e. in } Q_{\mathcal{T}} \cap \{\rho > 0\}.$$

Finally, we point out that the continuous Sobolev embedding  $H^1(\Omega) \hookrightarrow H^{1/2}(\partial\Omega)$  yields the convergence of the boundary integral:

$$\sum_{k=1}^N b_{ik} \left( \frac{\mu_k^{(n)}}{T^{(n)}} - \frac{\mu_{0,k}}{T_0} \right) \rightharpoonup \sum_{k=1}^N b_{ik} \left( \zeta_k - \frac{\mu_{0,k}}{T_0} \right) \quad \text{weakly in } L^2(\partial\Omega \times (0, \mathcal{T})).$$

4 This shows that (45) holds.

*Limit in equation (47).* Let us now turn our attention to  $\mathbf{q}^{(n)}$ . From (222) it follows

$$(121) \quad \frac{\mathbf{q}^{(n)}}{T^{(n)}} \rightharpoonup \tilde{\mathbf{q}} \quad \text{weakly in } L^{\frac{2+3\beta}{1+3\beta}}(Q_{\mathcal{T}}).$$

On the other hand, from (8) it follows

$$\frac{\mathbf{q}^{(n)}}{T^{(n)}} = -\frac{\kappa(T^{(n)})}{T^{(n)}} \nabla T^{(n)} + \sum_{j=1}^N \frac{\tilde{L}_{0j}(\vec{\rho}^{(n)}, T^{(n)})}{T^{(n)}} \nabla \frac{\mu_j^{(n)}}{T^{(n)}}.$$

From (38), (113), (117) and the boundedness of  $\sum_{j=1}^N \frac{\tilde{L}_{0j}(\vec{\rho}^{(n)}, T^{(n)})}{T^{(n)}} \nabla \frac{\mu_j^{(n)}}{T^{(n)}}$  in  $L^2(Q_{\mathcal{T}})$  we deduce

$$\sum_{j=1}^N \frac{\tilde{L}_{0j}(\vec{\rho}^{(n)}, T^{(n)})}{T^{(n)}} \nabla \frac{\mu_j^{(n)}}{T^{(n)}} \rightharpoonup \sum_{j=1}^N \frac{\tilde{L}_{0j}(\vec{\rho}, T)}{T} \nabla \zeta_j \quad \text{weakly in } L^2(Q_{\mathcal{T}}).$$

On the other hand, (221), (96) imply

$$\frac{\kappa(T^{(n)})}{T^{(n)}} \nabla T^{(n)} \rightharpoonup \frac{\kappa(T)}{T} \nabla T \quad \text{weakly in } L^{\frac{2+3\beta}{1+3\beta}}(Q_{\mathcal{T}}).$$

1 We conclude

$$(122) \quad \tilde{\mathbf{q}} = -\frac{\kappa(T)}{T} \nabla T + \sum_{j=1}^N \frac{\tilde{L}_{0j}(\vec{\rho}, T)}{T} \nabla \zeta_j = \frac{\mathbf{q}}{T}, \quad \text{a.e. in } Q_{\mathcal{T}}.$$

From (220), (116) it follows

$$-\sum_{i=1}^N \frac{\mu_i^{(n)}}{T^{(n)}} \mathbf{J}_i^{(n)} \rightharpoonup \Xi \quad \text{weakly in } L^{\frac{2a}{a+2}}(Q_{\mathcal{T}}), \quad \Xi = -\sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i \quad \text{a.e. in } Q_{\mathcal{T}} \cap \{\rho > 0\}.$$

Finally, consider the quantity

$$\begin{aligned} \xi^{(n)} &= \frac{K}{T^{(n)}} \lambda(S^{(n)}, T^{(n)}) |\nabla p^{(n)}|^2 + \sum_{i,j=1}^N L_{ij}^{(n)} \nabla \left( \frac{\mu_i^{(n)}}{T^{(n)}} \right) \cdot \nabla \left( \frac{\mu_j^{(n)}}{T^{(n)}} \right) + L_{00}^{(n)} \left| \nabla \left( \frac{1}{T^{(n)}} \right) \right|^2 \\ &\quad - \Phi \frac{1}{T^{(n)}} f'(S^{(n)}) (\partial_t S^{(n)})^2 - \sum_{i=1}^N r_i^{(n)} \frac{\mu_i^{(n)}}{T^{(n)}} \\ &= \frac{K}{T} \lambda(S, T) |\nabla p|^2 - \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) - L_{00} \left| \nabla \left( \frac{1}{T} \right) \right|^2 \\ &\quad + \Phi \frac{1}{T} f'(S) (\partial_t S)^2 + \sum_{i=1}^N r_i \frac{\mu_i}{T}. \end{aligned}$$

2 It is clear that  $\xi^{(n)}$  is bounded in  $L^1(Q_{\mathcal{T}})$  and therefore also in  $\mathcal{M}(\overline{Q_{\mathcal{T}}})$ . Therefore  $\xi^{(n)} \rightharpoonup^* \xi$   
3 weakly\* in  $\mathcal{M}(\overline{Q_{\mathcal{T}}})$ .

We will prove that  $\xi$  is a nonnegative measure. Let  $\varphi \in C^0(\overline{Q_{\mathcal{T}}})$ ,  $\varphi \geq 0$  in  $Q_{\mathcal{T}}$ . From (112) and the weak lower semicontinuity of the  $L^2(Q_{\mathcal{T}})$  norm we deduce

$$(123) \quad \liminf_{n \rightarrow \infty} \int_{Q_{\mathcal{T}}} \frac{K}{T^{(n)}} \lambda(S^{(n)}, T^{(n)}) |\nabla p^{(n)}|^2 \varphi dx dt \geq \int_{Q_{\mathcal{T}}} \frac{K}{T} \lambda(S, T) |\nabla p|^2 \varphi dx dt.$$

Furthermore, from (68) it follows that

$$\frac{1}{(T^{(n)})^{1/2}} \sqrt{-f'(S^{(n)})} \partial_t S^{(n)} \rightharpoonup \omega \quad \text{weakly in } L^2(Q_{\mathcal{T}}).$$

Since (97) holds we deduce

$$\sqrt{-f'(S^{(n)})} \partial_t S^{(n)} \rightharpoonup \sqrt{T} \omega \quad \text{weakly in } L^1(Q_{\mathcal{T}}).$$

Testing  $\sqrt{-f'(S^{(n)})}\partial_t S^{(n)}$  against an arbitrary test function, integrating by parts in time and taking the limit  $n \rightarrow \infty$  allows us to easily determine the value of  $\omega$ , yielding

$$\frac{1}{(T^{(n)})^{1/2}} \sqrt{-f'(S^{(n)})}\partial_t S^{(n)} \rightharpoonup \frac{1}{T^{1/2}} \sqrt{-f'(S)}\partial_t S \quad \text{weakly in } L^2(Q_{\mathcal{T}}).$$

Once again, the above relation and the weak lower semicontinuity of the  $L^2(Q_{\mathcal{T}})$  norm lead to

$$(124) \quad \liminf_{n \rightarrow \infty} \int_{Q_{\mathcal{T}}} -\Phi \frac{1}{T^{(n)}} f'(S^{(n)}) (\partial_t S^{(n)})^2 \varphi dxdt \geq \int_{Q_{\mathcal{T}}} -\Phi \frac{1}{T} f'(S) (\partial_t S)^2 \varphi dxdt.$$

Finally, Lemma 12 and assumptions (43), (44) imply that

$$(125) \quad \liminf_{n \rightarrow \infty} \int_{Q_{\mathcal{T}}} \sum_{i,j=1}^N L_{ij}^{(n)} \nabla \left( \frac{\mu_i^{(n)}}{T^{(n)}} \right) \cdot \nabla \left( \frac{\mu_j^{(n)}}{T^{(n)}} \right) \varphi dxdt \geq \int_{Q_{\mathcal{T}}} \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) \varphi dxdt,$$

$$(126) \quad \liminf_{n \rightarrow \infty} \int_{Q_{\mathcal{T}}} L_{00}^{(n)} \left| \nabla \left( \frac{1}{T^{(n)}} \right) \right|^2 \varphi dxdt \geq \int_{Q_{\mathcal{T}}} L_{00} \left| \nabla \left( \frac{1}{T} \right) \right|^2 \varphi dxdt,$$

$$(127) \quad \liminf_{n \rightarrow \infty} \int_{Q_{\mathcal{T}}} - \sum_{i=1}^N r_i^{(n)} \frac{\mu_i^{(n)}}{T^{(n)}} \varphi dxdt \geq \int_{Q_{\mathcal{T}}} - \sum_{i=1}^N r_i \frac{\mu_i}{T} \varphi dxdt.$$

From (123)–(127) and the definition of  $\xi$  we conclude

$$\langle \xi, \varphi \rangle = \lim_{n \rightarrow \infty} \langle \xi^{(n)}, \varphi \rangle \geq 0.$$

1 This means that  $\xi$  is nonnegative. Therefore (47) holds.

*Limit in equation (46).* We will now prove that (46) holds in the limit  $n \rightarrow \infty$ . From (62), (96) it follows that  $T^{(n)} \rightharpoonup T$  weakly in  $L^2(0, \mathcal{T}; H^1(\Omega))$ . Thanks to the Sobolev embedding  $H^1(\Omega) \hookrightarrow H^{1/2}(\partial\Omega)$  this implies that  $T^{(n)} \rightharpoonup T$  weakly in  $L^2(0, \mathcal{T}; H^{1/2}(\partial\Omega))$ . In particular

$$\int_0^t \int_{\partial\Omega} (T^{(n)} - T_0) d\sigma dt' \rightarrow \int_0^t \int_{\partial\Omega} (T - T_0) d\sigma dt', \quad n \rightarrow \infty.$$

Furthermore  $E_s(T^{(n)}) = c_s T^{(n)} \rightarrow c_s T = E_s(T)$  strongly in  $L^2(Q_{\mathcal{T}})$  thanks to (29), (97), while the term (recall (3))

$$E_{int}(S^{(n)}) = \int_{S^{(n)}}^1 P_c(s) ds$$

can be estimated from (34) and (60) through a similar argument as in (129), implying that

$$E_{int}(S^{(n)}) \rightarrow E_{int}(S) \quad \text{strongly in } L^1(Q_{\mathcal{T}}).$$

Therefore we must now only show that the term (recall (25))

$$S^{(n)}(\rho e)^{(n)} = S^{(n)}(\rho^{(n)})^\gamma + c_w \rho^{(n)} T^{(n)} + p_{at}$$

is strongly convergent in  $L^1(Q_{\mathcal{T}})$ . However, from (24) we obtain

$$(\gamma - 1)S^{(n)}(\rho e)^{(n)} - S^{(n)}p^{(n)} = [c_w(\gamma - 1) - 1]S^{(n)}\rho^{(n)}T^{(n)} + \gamma p_{at},$$

which, thanks to (89), (93), (96), (92), leads to

$$\begin{aligned} (\gamma - 1)S^{(n)}(\rho e)^{(n)} &\rightarrow Sp + [c_w(\gamma - 1) - 1]S\rho T + \gamma p_{at} = (\gamma - 1)S(\rho e) \\ &\quad \text{strongly in } L^1(Q_{\mathcal{T}}). \end{aligned}$$

2 We conclude that (46) holds.

1 *Limit in equation (13). We will now show that the saturation balance equation (13) holds in the*  
 2 *limit  $n \rightarrow \infty$ .*

We have by assumption

$$\partial_t f(S^{(n)}) + P_c(S^{(n)}) + p^{(n)} = 0, \quad t > 0, \quad \text{a.e. in } \Omega.$$

Multiplying the equation times  $S^{(n)}$  leads to

$$\partial_t F(S^{(n)}) + S^{(n)} P_c(S^{(n)}) + S^{(n)} p^{(n)} = 0, \quad t > 0, \quad \text{a.e. in } \Omega,$$

where  $F$  is like in (83). Integrating the above equation against a test function  $\varphi \in C_c^1(Q_{\mathcal{T}})$  yields

$$(128) \quad - \int_0^{\mathcal{T}} \int_{\Omega} F(S^{(n)}) \partial_t \varphi dx dt + \int_0^{\mathcal{T}} \int_{\Omega} (S^{(n)} P_c(S^{(n)}) + S^{(n)} p^{(n)}) \varphi dx dt = 0.$$

Since  $F(S) = - \int_S^1 s_1 f'(s_1) ds_1 = -f(1) + S f(S) + \int_S^1 f(s_1) ds_1$  and  $f \in C^0([0, 1])$ , then also  $F \in C^0([0, 1])$ . Therefore from (89) we deduce  $F(S^{(n)}) \rightarrow F(S)$  strongly in  $L^r(Q_{\mathcal{T}})$  for every  $r < \infty$ . On the other hand, (34) and (60) imply

$$(129) \quad \begin{cases} \|S^{(n)} P_c(S^{(n)})\|_{L^\infty(Q_{\mathcal{T}})} \leq C & k_p \leq 1, \\ \|S^{(n)} P_c(S^{(n)})\|_{L^{\frac{k_p}{k_p-1}}(Q_{\mathcal{T}})} \leq C \|P_c(S^{(n)})^{1-1/k_p}\|_{L^{\frac{k_p}{k_p-1}}(Q_{\mathcal{T}})} \leq C & k_p > 1. \end{cases}$$

Since  $z_p > 1$  and (89) holds, we deduce that  $S^{(n)} P_c(S^{(n)}) \rightarrow S P_c(S)$  strongly in  $L^1(Q_{\mathcal{T}})$ . Using (89) and (92) we can take the limit  $n \rightarrow \infty$  in (128) and get

$$- \int_0^{\mathcal{T}} \int_{\Omega} F(S) \partial_t \varphi dx dt + \int_0^{\mathcal{T}} \int_{\Omega} (S P_c(S) + S p) \varphi dx dt = 0, \quad \forall \varphi \in C_c^1(Q_{\mathcal{T}}),$$

which is the weak formulation of

$$\partial_t F(S) + S P_c(S) + S p = 0, \quad t > 0, \quad \text{a.e. in } \Omega.$$

3 Since  $F'(S) = S f'(S)$  and  $S > 0$  a.e. in  $Q_{\mathcal{T}}$ , dividing the above equation times  $S$  yields (13).

4 This finishes the proof of Theorem 8.

#### 5 4. THE APPROXIMATE SCHEME

6 In this section we build a sequence of approximated solutions to (11)–(13), (14), (17).

Let  $K_1 > 0$ ,  $K_2 > 0$ ,  $K_3 > 0$  be generic but fixed positive constants satisfying

$$(130) \quad \begin{cases} K_1 > \frac{6}{5}\gamma & \alpha_r \leq \frac{4}{3}, \\ \frac{6}{5}\gamma < K_1 < \frac{3\alpha_r-2}{3\alpha_r-4}\gamma & \alpha_r > \frac{4}{3}, \end{cases} \quad 3 < K_2 < 3\beta, \quad K_3 > \frac{5K_1 + 6\gamma}{5K_1 - 6\gamma}.$$

7 We point out that, for  $\alpha_r > 4/3$ , it holds  $\frac{3\alpha_r-2}{3\alpha_r-4} > \frac{6}{5}$  thanks to (30).

8 **4.1. Discretization and regularization.** We introduce the following parameters of the approx-  
 9 imate scheme:

- 10 (i)  $\varepsilon > 0$  lower-order regularization of (11)–(12), as well as the regularization in the free energy,  
 11 the source term, and the capillary pressure,
- 12 (ii)  $\delta > 0$  higher-order regularization in (11)–(12),
- 13 (iii)  $\tau > 0$  time-step in the implicit Euler discretization.

1 **Regularization.** We introduce the "regularized" free energy for the approximate system:

$$(131) \quad (\rho\Psi)_{w,\varepsilon} = (\rho\Psi)_w + \varepsilon\rho^{K_1}, \quad (\rho\Psi)_{s,\varepsilon} = (\rho\Psi)_s - \varepsilon T^{K_2}.$$

2 The definitions of the thermodynamics quantities  $\mu_i$ ,  $p$ ,  $\rho e$ ,  $(\rho\eta)_s$ ,  $E_s$  change accordingly:

$$(132) \quad \mu_{i,\varepsilon} = \frac{\partial(\rho\Psi)_{w,\varepsilon}}{\partial\rho_i} = \frac{\partial(\rho\Psi)_w}{\partial\rho_i} + \varepsilon K_1 \rho^{K_1-1},$$

3

$$(133) \quad p_\varepsilon = -(\rho\Psi)_{w,\varepsilon} + \sum_{i=1}^N \rho_i \mu_{i,\varepsilon} = -(\rho\Psi)_w + \sum_{i=1}^N \rho_i \mu_i + \varepsilon(K_1 - 1)\rho^{K_1},$$

4

$$(134) \quad (\rho e)_\varepsilon = (\rho\Psi)_{w,\varepsilon} - T \frac{\partial(\rho\Psi)_{w,\varepsilon}}{\partial T} = (\rho\Psi)_w - T \frac{\partial(\rho\Psi)_w}{\partial T} + \varepsilon \rho^{K_1},$$

5

$$(135) \quad (\rho\eta)_{s,\varepsilon} = -\frac{\partial(\rho\Psi)_s}{\partial T} + \varepsilon K_2 T^{K_2-1},$$

6

$$(136) \quad E_{s,\varepsilon} = (\rho\Psi)_{s,\varepsilon} - T \frac{\partial(\rho\Psi)_{s,\varepsilon}}{\partial T} = (\rho\Psi)_s - T \frac{\partial(\rho\Psi)_s}{\partial T} + \varepsilon(K_2 - 1)T^{K_2}.$$

7 However, in what follows, we will denote the regularized quantities  $\mu_{i,\varepsilon}$ ,  $p_\varepsilon$ ,  $(\rho e)_\varepsilon$ ,  $(\rho\eta)_{s,\varepsilon}$ ,  $E_{s,\varepsilon}$  with  
 8  $\mu_i$ ,  $p$ ,  $\rho e$ ,  $(\rho\eta)_s$ ,  $E_s$  to avoid a too cumbersome notation. Also, notice that the phase entropy  $\rho\eta$   
 9 remains unchanged. We also point out that the regularized skeleton entropy  $(\rho\eta)_s$  is still a concave  
 10 function of the regularized skeleton energy  $E_s$  (easy computations).

We regularize the source term  $r_i$ :

$$(137) \quad r_{i,\varepsilon} = r_i - \varepsilon \left| \frac{\mu_i}{T} \right|^{a-2} \frac{\mu_i}{T}, \quad i = 1, \dots, N.$$

We also regularize the dynamic capillary pressure term  $P_c$  in the following way:

$$(138) \quad P_{c,\varepsilon}(s) = \begin{cases} P_c(s) & \text{if } k_p > 0 \\ P_c(s) - \varepsilon \log s & \text{if } k_p = 0 \end{cases}, \quad E_{f,\varepsilon}^k = (\rho e)_\varepsilon^k S^k - \int_{1/2}^{S^k} P_{c,\varepsilon}(\xi) d\xi,$$

11 where the parameter  $k_p \geq 0$  is defined in (34).

Finally, we truncate the initial value for the saturation:

$$(139) \quad S^{in,\varepsilon} = \max\{\varepsilon, \min\{S^{in}, 1 - \varepsilon\}\}.$$

**Discretization.** Fix  $\mathcal{T} > 0$ . For  $L \in \mathbb{N}$  we define  $\tau = \mathcal{T}/L$ ,  $t_k = \tau k$  ( $k = 0, \dots, L$ ). Consider the implicit Euler time discretization in (11)–(13) with regularizing terms:

$$(140) \quad \begin{aligned} & \int_{\Omega} \tau^{-1} (S^k \rho_i^k - S^{k-1} \rho_i^{k-1}) \Phi \varphi dx - \int_{\Omega} (\rho_i^k \mathbf{v}^k + \mathbf{J}_i^k) \cdot \nabla \varphi dx - \int_{\Omega} r_{i,\varepsilon}^k \varphi dx \\ & + \int_{\partial\Omega} \sum_{\ell=1}^N b_{i\ell} \left( \frac{\mu_\ell^k}{T^k} - \frac{\mu_{0,\ell}}{T_0} \right) \varphi ds = -\varepsilon R_{\varepsilon,1}(\mu_i^k/T^k, \varphi) - \delta \tilde{R}_{\delta,1}(\mu_i^k/T^k, \varphi), \end{aligned}$$

$$\begin{aligned}
& \int_{\Omega} \tau^{-1} (\Phi(E_{f,\varepsilon}^k - E_{f,\varepsilon}^{k-1}) + (1 - \Phi)(E_s^k - E_s^{k-1})) \psi dx \\
& - \int_{\Omega} (((\rho e)^k + p^k) \mathbf{v}^k + \mathbf{q}^k) \cdot \nabla \psi dx + \alpha \int_{\partial\Omega} (T^k - T_0) \varphi ds \\
& = -\varepsilon \left( R_{\varepsilon,2}(T^k, \log T^k, \psi) + R_{\varepsilon,3}(T^k, \log T^k, \psi) \right) \\
(141) \quad & - \delta \left( \tilde{R}_{\delta,2}(T^k, \log T^k, \psi) + \tilde{R}_{\delta,3}(T^k, \log T^k, \psi) \right),
\end{aligned}$$

$$(142) \quad \frac{f(S^k) - f(S^{k-1})}{\tau} + P_{c,\varepsilon}(S^k) + p^k = 0,$$

for every  $\varphi, \psi \in C^2(\overline{\Omega})$ , where we defined:

$$(143) \quad R_{\varepsilon,1}(\mu_i^k/T^k, \varphi) = \int_{\Omega} \left( \nabla \frac{\mu_i^k}{T^k} \cdot \nabla \varphi + \frac{\mu_i^k}{T^k} \varphi \right) dx,$$

$$(144) \quad R_{\varepsilon,2}(T^k, \log T^k, \psi) = \int_{\Omega} ((1 + T^k) \nabla \log T^k \cdot \nabla \varphi + (1 + (T^k)^{-K_3}) (\log T^k) \psi) dx,$$

$$(145) \quad R_{\varepsilon,3}(T^k, \log T^k, \psi) = \int_{\Omega} (T^k)^{-K_3} |\nabla \log T^k|^{K_3-1} \nabla \log T^k \cdot \nabla \psi dx,$$

$$(146) \quad \tilde{R}_{\delta,1}(\mu_i^k/T^k, \varphi) = \int_{\Omega} D^2 \frac{\mu_i^k}{T^k} : D^2 \varphi dx,$$

$$(147) \quad \tilde{R}_{\delta,2}(T^k, \log T^k, \psi) = \int_{\Omega} (1 + T^k) D^2 \log T^k : D^2 \psi dx,$$

$$(148) \quad \tilde{R}_{\delta,3}(T^k, \log T^k, \psi) = \int_{\Omega} (1 + T^k) |\nabla \log T^k|^2 \nabla \log T^k \cdot \nabla \psi dx.$$

- 1 We will first prove the well-posedness of the approximate system and then perform the limits  
2  $\delta \rightarrow 0, \tau \rightarrow 0, \varepsilon \rightarrow 0$  in this order.

- 3 **4.2. Existence of solutions to the approximate system.** We formulate (140)-(141)-(142) as  
4 a fixed-point problem for an operator  $F : X \times [0, 1] \rightarrow X$  defined by means of a linearized problem.

- 5 4.2.1. *Reformulation as a fixed-point problem.* Define the spaces

$$V = W^{1,4}(\Omega; \mathbb{R}^N) \times W^{1,4}(\Omega), \quad V_0 = H^2(\Omega; \mathbb{R}^N) \times H^2(\Omega),$$

and variables

$$\vec{z} = \frac{\vec{\mu}}{T}, \quad w = \log T.$$

- 6 For arbitrary given  $(\vec{z}^*, w^*) \in V \hookrightarrow L^\infty(\Omega; \mathbb{R}^N) \times L^\infty(\Omega)$  (remember  $\Omega \subset \mathbb{R}^3$ ), define  $T^* = \exp(w^*)$ ,  
7  $\vec{\mu}^* = T^* \vec{z}^*$ , and the quantities  $\vec{\rho}^*, \mathbf{v}^*$  etc. accordingly to the constitutive relations.

For  $(\vec{z}^*, w^*) \in V$  and  $\sigma \in [0, 1]$ , let us consider the linearized problem

$$(149) \quad \sigma \left\{ \int_{\Omega} \tau^{-1}(S\rho_i^* - S^{k-1}\rho_i^{k-1})\Phi\varphi dx - \int_{\Omega} (\rho_i^*\mathbf{v}^* + \mathbf{J}_i^*) \cdot \nabla\varphi dx \right. \\ \left. - \int_{\Omega} r_{i,\varepsilon}^*\varphi dx + \int_{\partial\Omega} \sum_{\ell=1}^N b_{i\ell} \left( \frac{\mu_{\ell}^*}{T^*} - \frac{\mu_{0,\ell}}{T_0} \right) \varphi ds \right\} \\ = -\varepsilon R_{\varepsilon,1}(z_i, \varphi) - \delta R_{\delta,1}(z_i, \varphi), \quad \varphi \in H^2(\Omega),$$

$$(150) \quad \sigma \left\{ \int_{\Omega} \tau^{-1}(\Phi(E_{f,\varepsilon}^* - E_{f,\varepsilon}^{k-1}) + (1 - \Phi)(E_s^* - E_s^{k-1}))\psi dx \right. \\ \left. - \int_{\Omega} (((\rho e)^* + p^*)\mathbf{v}^* + \mathbf{q}^*) \cdot \nabla\psi dx + \alpha \int_{\partial\Omega} (T^* - T_0)\psi ds \right\} \\ = -\varepsilon \left( R_{\varepsilon,2}(T^*, w, \psi) + R_{\varepsilon,3}(T^*, w, \psi) \right) - \delta \left( R_{\delta,2}(T^*, w, \psi) + R_{\delta,3}(T^*, w, \psi) \right), \\ \psi \in H^2(\Omega),$$

1 coupled with the nonlinear algebraic equation

$$(151) \quad \tau^{-1}(f(S) - f(S^{k-1})) + \sigma(P_{c,\varepsilon}(S) + p^*) = 0.$$

The strict monotonicity of  $f$ ,  $P_{c,\varepsilon}$  as well as the fact that

$$\lim_{s \rightarrow 0^+} P_{c,\varepsilon}(s) = \infty, \quad \lim_{s \rightarrow 1^-} f(s) = -\infty$$

imply that (151) has a unique solution  $S : \Omega \rightarrow (0, 1)$  which is Lebesgue-measurable. Moreover, the Lax-Milgram Lemma allows us to state that system (149), (150) has a unique solution  $(\vec{z}, w) \in V_0 \hookrightarrow V$ . We consider therefore the mapping

$$F : ((\vec{z}^*, w^*), \sigma) \in V \times [0, 1] \mapsto (\vec{z}, w) \in V.$$

It holds that  $F(\cdot, 0) \equiv 0$ . Furthermore  $F$  is continuous (standard argument) and bounded as an operator  $V \times [0, 1] \rightarrow V_0$  (just choose  $\varphi = z_i$ ,  $\psi = w$  in (149), (150), respectively, and estimate the left-hand sides of the resulting equations in a straightforward way), which means (thanks to the compact Sobolev embedding  $V_0 \hookrightarrow V$ ) that  $F$  is compact. Now we wish to prove that the set

$$(152) \quad \mathcal{F}_{\sigma} \equiv \{(\vec{z}, w) \in V : F((\vec{z}, w), \sigma) = (\vec{z}, w)\}$$

2 is bounded in  $V$  uniformly w.r.t.  $\sigma \in [0, 1]$ . This will allow us to apply the Leray-Schauder  
3 fixed point theorem and deduce the existence of a fix point for  $F(\cdot, 1)$ , that is, a solution to the  
4 approximate system (140)–(142).

4.2.2. *Global entropy and energy balance for approximate solutions.* Let  $(\vec{z}, w) \in \mathcal{F}_{\sigma}$ . Recall that  $T = \exp(w)$ ,  $\vec{\mu} = T\vec{z}$ . Choose  $\varphi = z_i = \mu_i/T$ ,  $\psi = -1/T$  in (149), (150), respectively (also keep in mind that  $(\vec{z}, w) = (\vec{z}^*, w^*)$ , so we omit the  $*$  sign everywhere). It follows (the index  $i$  is summed

from 1 to  $N$ )

$$\begin{aligned}
(153) \quad & \sigma \left\{ \tau^{-1} \int_{\Omega} \left( \Phi(S\rho_i - S^{k-1}\rho_i^{k-1}) \frac{\mu_i}{T} - (\Phi(E_{f,\varepsilon} - E_{f,\varepsilon}^{k-1}) + (1 - \Phi)(E_s - E_s^{k-1})) \frac{1}{T} \right) dx \right. \\
& - \int_{\Omega} (\rho_i \mathbf{v} + \mathbf{J}_i) \cdot \nabla \frac{\mu_i}{T} dx + \int_{\Omega} ((\rho e) + p) \mathbf{v} + \mathbf{q} \cdot \nabla \frac{1}{T} dx + \alpha \int_{\partial\Omega} \frac{T_0 - T}{T} ds \\
& \left. - \int_{\Omega} r_{i,\varepsilon} \frac{\mu_i}{T} dx + \int_{\partial\Omega} \sum_{\ell=1}^N b_{i\ell} \left( \frac{\mu_\ell}{T} - \frac{\mu_{0,\ell}}{T_0} \right) \frac{\mu_i}{T} ds \right\} \\
& = - \varepsilon \int_{\Omega} \left( |\nabla \frac{\mu_i}{T}|^2 + \left| \frac{\mu_i}{T} \right|^2 \right) dx - \delta \int_{\Omega} |D^2 \frac{\mu_i}{T}|^2 dx \\
& - \varepsilon \int_{\Omega} \left( (1 + T) \frac{|\nabla \log T|^2}{T} - (1 + T^{-K_3}) \frac{\log T}{T} \right) dx \\
& - \varepsilon \int_{\Omega} |T|^{-1-K_3} |\nabla \log T|^{K_3+1} dx \\
& - \delta \int_{\Omega} \left( (1 + T) D^2 \log T : D^2 \left( -\frac{1}{T} \right) + (T^{-1} + 1) |\nabla \log T|^4 \right) dx.
\end{aligned}$$

- 1 We consider firstly the terms coming from the regularization and from the discrete time derivatives.
- 2 Note that the Gibbs-Duhem relations (27) give:

$$\frac{\partial(-\rho\eta)}{\partial\rho_i} = \frac{\mu_i}{T}, \quad \frac{\partial(-\rho\eta)}{\partial(\rho e)} = -\frac{1}{T}.$$

In this way we get:

$$\begin{aligned}
& \Phi(S\rho_i - S^{k-1}\rho_i^{k-1}) \frac{\mu_i}{T} - (\Phi(E_{f,\varepsilon} - E_{f,\varepsilon}^{k-1}) + (1 - \Phi)(E_s - E_s^{k-1})) \frac{1}{T} \\
& = S^{k-1} \Phi \left\{ (\rho_i - \rho_i^{k-1}) \frac{\partial(-\rho\eta)}{\partial\rho_i} + ((\rho e) - (\rho e)^{k-1}) \frac{\partial(-\rho\eta)}{\partial(\rho e)} \right\} \\
& + \Phi(S - S^{k-1}) \left\{ \rho_i \frac{\mu_i}{T} - \frac{(\rho e)}{T} \right\} \\
& + \frac{\Phi}{T} \int_{S^{k-1}}^S P_{c,\varepsilon}(\xi) d\xi - (1 - \Phi) c_s \frac{T - T^{k-1}}{T} \\
& - (1 - \Phi) \varepsilon (K_2 - 1) \frac{T^{K_2} - (T^{k-1})^{K_2}}{T}.
\end{aligned}$$

The concavity of the functions

$$(\bar{\rho}, \rho e) \mapsto \rho\eta, \quad T \mapsto \log T, \quad T \mapsto T^{1-1/K_2}, \quad S \mapsto \int_{1/2}^S P_{c,\varepsilon}(\xi) d\xi$$

and (54) lead to

$$\begin{aligned}
& \Phi(S\rho_i - S^{k-1}\rho_i^{k-1}) \frac{\mu_i}{T} - (\Phi(E_{f,\varepsilon} - E_{f,\varepsilon}^{k-1}) + (1 - \Phi)(E_s - E_s^{k-1})) \frac{1}{T} \\
& \geq \Phi S^{k-1} (\rho\eta)^{k-1} + (1 - \Phi) [c_s \log T^{k-1} + \varepsilon K_2 (T^{k-1})^{K_2-1}] - \Phi S (\rho\eta) \\
& - (1 - \Phi) [c_s \log T + \varepsilon K_2 T^{K_2-1}] + \Phi(S - S^{k-1}) \frac{P_{c,\varepsilon}(S) + p}{T}.
\end{aligned}$$



Multiplying the above inequality times  $\sigma\tau^{-1}$  and employing (151) and (28) allows us to obtain

$$(154) \quad \begin{aligned} & \frac{\sigma}{\tau} \left\{ \Phi(S\rho_i - S^{k-1}\rho_i^{k-1})\frac{\mu_i}{T} - (\Phi(E_{f,\varepsilon} - E_{f,\varepsilon}^{k-1}) + (1-\Phi)(E_s - E_s^{k-1}))\frac{1}{T} \right\} \\ & \geq -\frac{\sigma}{\tau} \left\{ \Phi\left(S(\rho\eta) - S^{k-1}(\rho\eta)^{k-1}\right) + (1-\Phi)\left((\rho\eta)_s - (\rho\eta)_s^{k-1}\right) \right\} \\ & \quad - \frac{\Phi}{T}\tau^{-2}(S - S^{k-1})(f(S) - f(S^{k-1})) - \frac{\sigma}{\tau}(1-\Phi)\varepsilon K_2[T^{K_2-1} - (T^{k-1})^{K_2-1}]. \end{aligned}$$

Next, using Young's inequality, direct calculations give the following estimate:

$$\begin{aligned} & \delta \int_{\Omega} ((1+T)D^2 \log T : D^2 \left(-\frac{1}{T}\right) + (T^{-1} + 1)|\nabla \log T|^4) dx \\ & \geq c\delta \int_{\Omega} (|D^2 \log T|^2 + (T^{-1} + 1)|\nabla \log T|^4) dx \end{aligned}$$

1 for some constant  $c > 0$ .

Therefore, from (153), using (58) and the discrete entropy balance equation, we obtain the global entropy balance for the approximate solutions:

$$(155) \quad \begin{aligned} & \sigma \int_{\Omega} \left\{ \sum_{i,j=1}^N L_{ij} \nabla \left(\frac{\mu_i}{T}\right) \cdot \nabla \left(\frac{\mu_j}{T}\right) + \frac{\kappa(T)|\nabla T|^2}{T^2} + \frac{K}{T} \lambda(S, T) |\nabla p|^2 - \vec{r} \cdot \frac{\vec{\mu}}{T} \right\} dx \\ & - \int_{\Omega} \Phi \tau^{-2} (S - S^{k-1})(f(S) - f(S^{k-1})) dx \\ & + \sigma \int_{\partial\Omega} \left( \alpha \frac{T_0 - T}{T} + \sum_{i,j=1}^N b_{ij} \frac{\mu_i}{T} \left( \frac{\mu_j}{T} - \frac{\mu_{0,j}}{T_0} \right) \right) ds \\ & + \varepsilon \int_{\Omega} \left( |\nabla \frac{\mu_i}{T}|^2 + |\frac{\mu_i}{T}|^2 \right) dx + \delta \int_{\Omega} |D^2 \frac{\mu_i}{T}|^2 dx \\ & + \varepsilon \int_{\Omega} \left( (1+T) \frac{|\nabla \log T|^2}{T} - (1+T^{-K_3}) \frac{\log T}{T} \right) dx \\ & + \varepsilon \int_{\Omega} |T|^{-1-K_3} |\nabla \log T|^{K_3+1} dx \\ & + c\delta \int_{\Omega} (|D^2 \log T|^2 + (T^{-1} + 1)|\nabla \log T|^4) dx \\ & \leq \sigma\tau^{-1} \int_{\Omega} \left( \Phi(S(\rho\eta) - S^{k-1}(\rho\eta)^{k-1}) + (1-\Phi)((\rho\eta)_{s,\varepsilon} - (\rho\eta)_{s,\varepsilon}^{k-1}) \right) dx. \end{aligned}$$

On the other hand, choosing  $\psi = 1$  in (150) leads to the global energy balance inequality for the approximate solutions:

$$(156) \quad \begin{aligned} & \sigma \left\{ \int_{\Omega} \tau^{-1} (\Phi(E_{f,\varepsilon} - E_{f,\varepsilon}^{k-1}) + (1-\Phi)(E_s - E_s^{k-1})) dx + \alpha \int_{\partial\Omega} (T - T_0) ds \right\} \\ & = -\varepsilon \int_{\Omega} (1 + T^{-K_3}) \log T dx. \end{aligned}$$

Summing (155) and (156) yields the global entropy-energy inequality for the approximate solutions:

$$\begin{aligned}
(157) \quad & \sigma\tau^{-1} \int_{\Omega} [\Phi(E_{f,\varepsilon} - S(\rho\eta)) + (1 - \Phi)(E_s - (\rho\eta)_{s,\varepsilon})] dx \\
& + \sigma \int_{\Omega} \left\{ \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) + \frac{\kappa(T)|\nabla T|^2}{T^2} + \frac{K}{T} \lambda(S, T) |\nabla p|^2 - \vec{r} \cdot \frac{\vec{\mu}}{T} \right\} dx \\
& - \int_{\Omega} \frac{\Phi}{T} \tau^{-2} (S - S^{k-1})(f(S) - f(S^{k-1})) dx \\
& + \sigma \int_{\partial\Omega} \left( \alpha \left( 1 - \frac{1}{T} \right) (T - T_0) + \sum_{i,j=1}^N b_{ij} \frac{\mu_i}{T} \left( \frac{\mu_j}{T} - \frac{\mu_{0,j}}{T_0} \right) \right) ds \\
& + \varepsilon \int_{\Omega} \left( \left| \nabla \frac{\vec{\mu}}{T} \right|^2 + \left| \frac{\vec{\mu}}{T} \right|^2 \right) dx + \delta \int_{\Omega} \left| D^2 \frac{\vec{\mu}}{T} \right|^2 dx \\
& + \varepsilon \int_{\Omega} \left( (1 + T) \frac{|\nabla \log T|^2}{T} + (1 + T^{-K_3}) \left( 1 - \frac{1}{T} \right) \log T \right) dx \\
& + \varepsilon \int_{\Omega} |T|^{-1-K_3} |\nabla \log T|^{K_3+1} dx \\
& + c\delta \int_{\Omega} (|D^2 \log T|^2 + (T^{-1} + 1) |\nabla \log T|^4) dx \\
& \leq \sigma\tau^{-1} \int_{\Omega} [\Phi(E_{f,\varepsilon}^{k-1} - S^{k-1}(\rho\eta)^{k-1}) + (1 - \Phi)(E_s^{k-1} - (\rho\eta)_{s,\varepsilon}^{k-1})] dx.
\end{aligned}$$

Furthermore, since  $\int_{s_\varepsilon^*}^S P_{c,\varepsilon}(\xi) d\xi \leq C$  thanks to the definition on  $P_{c,\varepsilon}$ , we deduce from the definition of  $E_{f,\varepsilon}$  and (25), (26) that

$$\begin{aligned}
E_{f,\varepsilon} - S(\rho\eta) &= S(\rho^\gamma + c_w \rho T + p_{at}) - \int_{s_\varepsilon^*}^S P_{c,\varepsilon}(\xi) d\xi + S\varepsilon\rho^{K_1} \\
&+ S \sum_{i=1}^N \rho_i \log \rho_i - c_w S \rho (\log T + 1) \\
&\geq -C + S \left( \rho^\gamma + c_w \rho (T - \log T) + \sum_{i=1}^N \rho_i (\log \rho_i - c_w) \right) + S\varepsilon\rho^{K_1} \\
&\geq S\rho^\gamma + S\varepsilon\rho^{K_1} - C,
\end{aligned}$$

while (28), (29) imply

$$\begin{aligned}
E_s - (\rho\eta)_s &= c_s(T - \log T) + 1 - c_s + \varepsilon T^{K_2-1} ((K_2 - 1)T - K_2) \\
&\geq C(T + |\log T| + \varepsilon T^{K_2}) - C'.
\end{aligned}$$

It follows

$$\begin{aligned}
(158) \quad & \Phi(E_{f,\varepsilon} - S(\rho\eta)) + (1 - \Phi)(E_s - (\rho\eta)_s) \\
& \geq c_1 \Phi S(\rho^\gamma + \varepsilon\rho^{K_1}) + c_2(1 - \Phi)(T + |\log T| + \varepsilon T^{K_2}) - C.
\end{aligned}$$

We deduce from (157), (158) that (recall  $w = \log T$ ,  $\vec{z} = \vec{\mu}/T$ )

$$\|\vec{z}\|_{H^2(\Omega; \mathbb{R}^N)} + \|D^2 w\|_{L^2(\Omega)} + \|\nabla w\|_{L^2(\Omega)} + \int_{\Omega} w(1 - e^{-w}) dx \leq C(\varepsilon, \delta, \tau).$$

Since  $C > 0$  exists such that  $|w| \leq C(w(1 - e^{-w}) + 1)$  for every  $w \in \mathbb{R}$ , we obtain through the Poincaré Lemma

$$\|\vec{z}\|_{H^2(\Omega; \mathbb{R}^N)} + \|w\|_{H^2(\Omega)} \leq C(\varepsilon, \delta, \tau).$$

4.2.3. *Conclusion: existence for the approximate problem.* We conclude that the set in (152) is bounded in  $V_0$  (and a fortiori in  $V$ ) uniformly w.r.t.  $\sigma \in [0, 1]$ . Leray-Schauder's fixed point theorem allows us to conclude that  $(\vec{z}, w) \in V_0$  exists such that  $F((\vec{z}, w), 1) = (\vec{z}, w)$ , i.e. the approximate system has a solution  $(\vec{\mu}^k, T^k, S^k)$  such that  $\vec{\mu}^k/T^k = \vec{z} \in H^2(\Omega; \mathbb{R}^N)$ ,  $\log T^k = w \in H^2(\Omega)$ ,  $S \in L^\infty(\Omega)$ ,  $0 < S < 1$  a.e. in  $\Omega$ . In particular  $\vec{\mu}^k \in H^2(\Omega; \mathbb{R}^N)$ ,  $T^k \in H^2(\Omega)$ . Furthermore (157) holds with  $\sigma = 1$ . We conclude this part by stating the following technical result, whose proof can be found in the Appendix.

**Lemma 10.** *The following estimates for  $S_k$  hold:*

$$(159) \quad S^k \geq \varepsilon > 0 \text{ a.e. in } \Omega, \quad k \geq 0, \text{ provided that } 0 < \varepsilon < f^{-1}(p_{at}/\lambda_0),$$

$$(160) \quad \left| \int_{\Omega} f(S^k) dx \right| \leq \left| \int_{\Omega} f(S^{k-1}) dx \right| + \tau C(\varepsilon) \int_{\Omega} (1 + |p^k|) dx, \quad k \geq 0,$$

$$(161) \quad (1 - C\tau) \int_{\Omega} (1 + |\nabla f(S^k)|^2)^{2/3} dx + \tau \int_{\Omega} \frac{|f'(S^k) P'_{c,\varepsilon}(S^k)| |\nabla S^k|^2}{(1 + |\nabla f(S^k)|^2)^{1/3}} dx \\ \leq \int_{\Omega} (1 + |\nabla f(S^{k-1})|^2)^{2/3} dx + \tau \int_{\Omega} |\nabla p^k|^{4/3} dx, \quad k \geq 0.$$

4.3. **Limit  $\delta \rightarrow 0$ .** We denote all the variables with a  $\delta$  apex to put their dependence on  $\delta$  in evidence.

From (157) it follows

$$\sqrt{\varepsilon} \left\| \frac{\vec{\mu}^{(\delta)}}{T^{(\delta)}} \right\|_{H^1(\Omega)} + \|\log T^{(\delta)}\|_{H^1(\Omega)} + \|(T^{(\delta)})^{\frac{\beta}{2}}\|_{H^1(\Omega)} \leq C(\tau).$$

The compact Sobolev's embedding  $H^1(\Omega) \hookrightarrow L^q(\Omega)$  for every  $q < 6$  implies that  $\vec{\mu}^{(\delta)}/T^{(\delta)}$ ,  $\log T^{(\delta)}$ ,  $(T^{(\delta)})^{\beta/2}$  are strongly convergent in  $L^q(\Omega)$  for every  $q < 6$ , and in particular also a.e. in  $\Omega$ . From (23) it follows, for  $i = 1, \dots, N$ :

$$(162) \quad \log \rho_i^{(\delta)} + \frac{\gamma(\rho^{(\delta)})^{\gamma-1}}{T^{(\delta)}} + \varepsilon K_1(\rho^{(\delta)})^{K_1-1} = \frac{\mu_i^{(\delta)}}{T^{(\delta)}} + c_w \log T^{(\delta)} - 1.$$

Taking the exponential of both sides in (162) and summing in  $i = 1, \dots, N$  yields

$$\rho^{(\delta)} \exp \left( \frac{\gamma(\rho^{(\delta)})^{\gamma-1}}{T^{(\delta)}} + \varepsilon K_1(\rho^{(\delta)})^{K_1-1} \right) = \sum_{i=1}^N \exp \left( \frac{\mu_i^{(\delta)}}{T^{(\delta)}} + c_w \log T^{(\delta)} - 1 \right).$$

The right-hand side of the above identity is a.e. convergent in  $\Omega$  to an a.e. positive function, while  $T^{(\delta)} \rightarrow T > 0$  a.e. in  $\Omega$ . By the implicit function theorem  $\rho^{(\delta)}$  is a.e. convergent in  $\Omega$  to an a.e. positive function  $\rho$ . From this fact and (162) it follows that  $\rho_i^{(\delta)}$  is a.e. convergent in  $\Omega$  to  $\rho_i > 0$ , for  $i = 1, \dots, N$ .

From (157), (158), (159) we also deduce

$$(163) \quad \|\rho^{(\delta)}\|_{L^{K_1}(\Omega)} \leq C(\varepsilon, \tau).$$

In particular  $\rho_i^{(\delta)} \rightarrow \rho_i$  strongly in  $L^q(\Omega)$ , for every  $q < K_1$ ,  $i = 1, \dots, N$ .

From (157) and (5) (as well as the boundedness of  $\mu$ ) it follows that

$$\left\| \sqrt{\frac{k_r(S^{(\delta)})}{T^{(\delta)}}} \nabla p^{(\delta)} \right\|_{L^2(\Omega)} \leq C(\tau).$$

However, (159) and the  $L^6(\Omega)$  bound for  $(T^{(\delta)})^{\beta/2}$  yield

$$\begin{aligned} \|\nabla p^{(\delta)}\|_{L^{\frac{6\beta}{1+3\beta}}(\Omega)} &\leq \|\sqrt{T^{(\delta)}}\|_{L^{6\beta}(\Omega)} \left\| \frac{1}{\sqrt{T^{(\delta)}}} \nabla p^{(\delta)} \right\|_{L^2(\Omega)} \\ &\leq C(\varepsilon) \|\sqrt{T^{(\delta)}}\|_{L^{6\beta}(\Omega)} \left\| \sqrt{\frac{k_r(S^{(\delta)})}{T^{(\delta)}}} \nabla p^{(\delta)} \right\|_{L^2(\Omega)} \leq C(\varepsilon, \tau). \end{aligned}$$

The above estimate, together with the bound for  $p^{(\delta)}$  in  $L^1(\Omega)$  which comes from the  $L^1(\Omega)$  estimate for  $E_{f,\varepsilon}^{(\delta)}$  and relation  $p^{(\delta)} \leq CE_{f,\varepsilon}^{(\delta)} + C'$  implies (via Poincaré's Lemma)

$$(164) \quad \|p^{(\delta)}\|_{W^{1, \frac{6\beta}{1+3\beta}}(\Omega)} \leq C(\varepsilon, \tau).$$

Therefore via Sobolev's embedding

$$(165) \quad \|p^{(\delta)}\|_{L^{\frac{6\beta}{1+\beta}}(\Omega)} \leq C(\varepsilon, \tau),$$

$$p^{(\delta)} \rightarrow p \quad \text{strongly in } L^q(\Omega), \quad \forall q < \frac{6\beta}{1+\beta}.$$

Furthermore the trivial bound  $(\rho^{(\delta)})^\gamma \leq C(p^{(\delta)} + 1)$  and the a.e. convergence of  $\rho_i^{(\delta)}$  imply

$$\rho_i^{(\delta)} \rightarrow \rho_i \quad \text{strongly in } L^q(\Omega), \quad \forall q < \frac{6\beta\gamma}{1+3\beta}, \quad i = 1, \dots, N.$$

- 1 The constitutive relations between all the variables (see subsection 1.2) hold because  $\rho_i^{(\delta)}, T^{(\delta)}$  are
- 2 a.e. convergent in  $\Omega$  towards a.e. positive limits.
- 3 Finally, (142), the strong convergence of  $p^{(\delta)}$  and the strict monotonicity of  $f, P_{c,\varepsilon}$  imply that
- 4  $S^{(\delta)}$  is a.e. convergent in  $\Omega$  towards a limit  $S$  which satisfies  $0 < \varepsilon \leq S < 1$  a.e. in  $\Omega$  due to
- 5 (159) and  $\lim_{s \rightarrow 1} f(s) = -\infty$ . Furthermore  $S^{(\delta)}$  is also strongly convergent to  $S$  in  $L^q(\Omega)$  for every
- 6  $q < \infty$ .

It is immediate to see that the  $\delta$ -depending regularization terms in (140), (141) easily tend to 0 when  $\delta \rightarrow 0$ , the only exception being the term

$$\delta \int_{\Omega} T^{(\delta)} |\nabla \log T^{(\delta)}|^2 \nabla \log T^{(\delta)} \cdot \nabla \psi dx$$

which appears in (141). Since  $\beta \geq 4/3$  by assumption, it follows

$$\begin{aligned} \delta \int_{\Omega} T^{(\delta)} |\nabla \log T^{(\delta)}|^3 dx &\leq \delta \|T^{(\delta)}\|_{L^4(\Omega)} \|\nabla \log T^{(\delta)}\|_{L^4(\Omega)}^3 \\ &\leq C\delta \|T^{(\delta)}\|_{L^{3\beta}(\Omega)} \|\nabla \log T^{(\delta)}\|_{L^4(\Omega)}^3. \end{aligned}$$

Since  $T^{(\delta)}$  is bounded in  $L^{3\beta}(\Omega)$  while  $\delta^{1/4} \nabla \log T^{(\delta)}$  is bounded in  $L^4(\Omega)$ , we conclude

$$\delta \int_{\Omega} T^{(\delta)} |\nabla \log T^{(\delta)}|^3 dx \leq C\delta^{1/4}.$$

- 1 Let us now check whether the terms in the energy equation are bounded in  $L^q(\Omega)$  for some  $q > 1$ .
- 2 From (164), (6) and the boundedness of  $\lambda(S^{(\delta)}, T^{(\delta)})$  it follows

$$(166) \quad \|\mathbf{v}^{(\delta)}\|_{L^{\frac{6\beta}{1+3\beta}}(\Omega)} \leq C,$$

and so

$$\mathbf{v}^{(\delta)} \rightharpoonup \mathbf{v} \quad \text{weakly in } L^{\frac{6\beta}{1+3\beta}}(\Omega).$$

Furthermore (165) and the strong convergence of  $S^{(\delta)}, T^{(\delta)}$  imply

$$\begin{aligned} \nabla p^{(\delta)} &\rightharpoonup \nabla p \quad \text{weakly in } L^{\frac{6\beta}{1+3\beta}}(\Omega), \\ \lambda(S^{(\delta)}, T^{(\delta)}) &\rightarrow \lambda(S, T) \quad \text{strongly in } L^q(\Omega), \quad \forall q < \infty. \end{aligned}$$

- 3 It follows  $\mathbf{v} = -K\lambda(S, T)\nabla p$ .

From (165), (166) we deduce

$$p^{(\delta)}\mathbf{v}^{(\delta)} \rightharpoonup p\mathbf{v} \quad \text{weakly in } L^{\frac{3\beta}{1+2\beta}}(\Omega).$$

Notice  $\frac{3\beta}{1+2\beta} > 1$  since  $\beta > 1$ . Also, since  $(\rho e)^{(\delta)} \leq Cp^{(\delta)} + C'$ , it follows

$$(\rho e)^{(\delta)}\mathbf{v}^{(\delta)} \rightharpoonup (\rho e)\mathbf{v} \quad \text{weakly in } L^{\frac{3\beta}{1+2\beta}}(\Omega).$$

Let us consider the heat flux  $\mathbf{q}^{(\delta)}$ . Since

$$\mathbf{q}^{(\delta)} = -\kappa(T^{(\delta)})\nabla T^{(\delta)} + \sum_{j=1}^N L_{0j} \nabla \frac{\mu_j^{(\delta)}}{T^{(\delta)}}$$

and  $\mu_j^{(\delta)}/T^{(\delta)} \rightharpoonup \mu_j/T$  weakly in  $L^2(\Omega)$ , for  $j = 1, \dots, N$ , one only needs to make sure that  $\kappa(T^{(\delta)})\nabla T^{(\delta)}$  converges (weakly) to the correct limit. From the  $\delta$ -uniform bound for  $\nabla(T^{(\delta)})^{K_4/2}$  in  $L^2(\Omega)$  it follows

$$\begin{aligned} \|\kappa(T^{(\delta)})\nabla T^{(\delta)}\|_{L^{\frac{3\beta}{2\beta+1}}(\Omega)} &\leq C\|(T^{(\delta)})^{\beta/2+1}\nabla(T^{(\delta)})^{\beta/2}\|_{L^{\frac{3\beta}{2\beta+1}}(\Omega)} \\ &\leq C\|(T^{(\delta)})^{\beta/2+1}\|_{L^{\frac{6\beta}{\beta+2}}(\Omega)}\|\nabla(T^{(\delta)})^{\beta/2}\|_{L^2(\Omega)} \\ &\leq C\|T^{(\delta)}\|_{L^{3\beta}(\Omega)}^{\beta/2+1}\|\nabla(T^{(\delta)})^{\beta/2}\|_{L^2(\Omega)} \leq C(\varepsilon, \tau). \end{aligned}$$

Therefore

$$\kappa(T^{(\delta)})\nabla T^{(\delta)} \rightharpoonup \kappa(T)\nabla T \quad \text{weakly in } L^{\frac{3\beta}{2\beta+1}}(\Omega).$$

- 4 It is immediate to see that  $E_{f,\varepsilon}^{(\delta)}$  is strongly convergent (in  $L^1(\Omega)$ ) towards the correct limit. At
- 5 this point taking the limit  $\delta \rightarrow 0$  in (140)–(142) is straightforward. The resulting limit equations
- 6 are just like (140)–(142) but without the regularizing terms in  $\delta$ .

4.4. **Limit**  $\tau \rightarrow 0$ . Let us introduce the new notation

$$\rho_i^{(\tau)}(x, t) = \rho_i^{in}(x)\mathcal{X}_{\{0\}}(t) + \sum_{k=1}^N \rho_i^k(x)\mathcal{X}_{((k-1)\tau, k\tau]}(t), \quad i = 1, \dots, N,$$

and the same for the other variables. We introduce also the discrete time derivative

$$D_\tau f(t) = \tau^{-1}(f(t) - f(t - \tau)), \quad t \in [\tau, \mathcal{T}],$$

for any function  $f : [0, \mathcal{T}] \rightarrow X$  ( $X$  is any vector space on  $\mathbb{R}$ ). We can rewrite (140)–(142) (after the limit  $\delta \rightarrow 0$ ) as

$$(167) \quad \begin{aligned} & \int_0^{\mathcal{T}} \int_{\Omega} D_{\tau}(S^{(\tau)} \rho_i^{(\tau)}) \Phi \varphi dxdt - \int_0^{\mathcal{T}} \int_{\Omega} (\rho_i^{(\tau)} \mathbf{v}^{(\tau)} + \mathbf{J}_i^{(\tau)}) \cdot \nabla \varphi dxdt \\ & - \int_0^{\mathcal{T}} \int_{\Omega} r_{i,\varepsilon}^{(\tau)} \varphi dxdt + \int_0^{\mathcal{T}} \int_{\partial\Omega} \sum_{\ell=1}^N b_{i\ell} \left( \frac{\mu_{\ell}^{(\tau)}}{T^{(\tau)}} - \frac{\mu_{0,\ell}}{T_0} \right) \varphi dsdt \\ & = -\varepsilon \int_0^{\mathcal{T}} \int_{\Omega} \left( \nabla \frac{\mu_i^{(\tau)}}{T^{(\tau)}} \cdot \nabla \varphi + \frac{\mu_i^{(\tau)}}{T^{(\tau)}} \varphi \right) dxdt, \end{aligned}$$

$$(168) \quad \begin{aligned} & \int_0^{\mathcal{T}} \int_{\Omega} D_{\tau}(\Phi E_{f,\varepsilon}^{(\tau)} + (1 - \Phi) E_s^{(\tau)}) \psi dxdt \\ & - \int_0^{\mathcal{T}} \int_{\Omega} (((\rho e)^{(\tau)} + p^{(\tau)}) \mathbf{v}^{(\tau)} + \mathbf{q}^{(\tau)}) \cdot \nabla \psi dxdt + \alpha \int_0^{\mathcal{T}} \int_{\partial\Omega} (T^{(\tau)} - T_0) \psi dsdt \\ & = -\varepsilon \int_0^{\mathcal{T}} \int_{\Omega} ((1 + T^{(\tau)}) \nabla \log T^{(\tau)} \cdot \nabla \psi + (1 + (T^{(\tau)})^{-K_3}) (\log T^{(\tau)}) \psi) dxdt \\ & - \varepsilon \int_0^{\mathcal{T}} \int_{\Omega} (T^{(\tau)})^{-K_3} |\nabla \log T^{(\tau)}|^{K_3-1} \nabla \log T^{(\tau)} \cdot \nabla \psi dxdt, \end{aligned}$$

1

$$(169) \quad D_{\tau} f(S^{(\tau)}) + P_{c,\varepsilon}(S^{(\tau)}) + p^{(\tau)} = 0.$$

Estimate (157) can be rewritten as

$$(170) \quad \begin{aligned} & \sup_{[0,\mathcal{T}]} \int_{\Omega} \left[ \Phi(E_{f,\varepsilon}^{(\tau)} - S^{(\tau)}(\rho\eta)^{(\tau)}) + (1 - \Phi)(E_s^{(\tau)} - (\rho\eta)_s^{(\tau)}) \right] dx \\ & + \int_0^{\mathcal{T}} \int_{\Omega} \left\{ \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i^{(\tau)}}{T^{(\tau)}} \right) \cdot \nabla \left( \frac{\mu_j^{(\tau)}}{T^{(\tau)}} \right) + \frac{\kappa(T^{(\tau)}) |\nabla T^{(\tau)}|^2}{|T^{(\tau)}|^2} \right. \\ & \left. + \frac{K}{T^{(\tau)}} \lambda(S^{(\tau)}, T^{(\tau)}) |\nabla p^{(\tau)}|^2 - \bar{r}^{(\tau)} \cdot \frac{\bar{\mu}^{(\tau)}}{T^{(\tau)}} - \Phi(D_{\tau} S^{(\tau)})(D_{\tau} f(S^{(\tau)})) \right\} dx \\ & + \int_0^{\mathcal{T}} \int_{\partial\Omega} \left[ \alpha \left( 1 - \frac{1}{T^{(\tau)}} \right) (T^{(\tau)} - T_0) + \sum_{i,j=1}^N b_{ij} \frac{\mu_i^{(\tau)}}{T^{(\tau)}} \left( \frac{\mu_j^{(\tau)}}{T^{(\tau)}} - \frac{\mu_{0,j}}{T_0} \right) \right] dsdt \\ & + \varepsilon \int_0^{\mathcal{T}} \int_{\Omega} \left( \left| \nabla \frac{\bar{\mu}^{(\tau)}}{T^{(\tau)}} \right|^2 + \left| \frac{\bar{\mu}^{(\tau)}}{T^{(\tau)}} \right|^2 + (1 + T^{(\tau)}) \frac{|\nabla \log T^{(\tau)}|^2}{T^{(\tau)}} \right. \\ & \left. + \left| \nabla \frac{1}{T^{(\tau)}} \right|^{K_3+1} + (1 + (T^{(\tau)})^{-K_3}) \left( 1 - \frac{1}{T^{(\tau)}} \right) \log T^{(\tau)} \right) dxdt \\ & \leq \int_{\Omega} [\Phi(E_{f,\varepsilon}^{in} - S^{in,\varepsilon}(\rho\eta)^{in}) + (1 - \Phi)(E_s^{in} - (\rho\eta)_s^{in})] dx. \end{aligned}$$

The a priori estimates derived in Section 2 are satisfied by the approximate solution. Moreover, the latter satisfies also the following bounds, which follow from (170):

$$(171) \quad \sqrt{\varepsilon} \|\mu_i^{(\tau)} / T^{(\tau)}\|_{L^2(0,\mathcal{T}; H^1(\Omega))} + \varepsilon^{1/a} \|\mu_i^{(\tau)} / T^{(\tau)}\|_{L^a(Q_{\mathcal{T}})} \leq C, \quad i = 1, \dots, N.$$

- 1 Let us now find a better estimate for  $T^{(\tau)}$ . The  $L^\infty(0, \mathcal{T}; L^1(\Omega))$  bound for  $E_{s,\varepsilon}$  (see (170)) as well  
 2 as (136) lead to

$$(172) \quad \|T^{(\tau)}\|_{L^\infty(0, \mathcal{T}; L^{K_2}(\Omega))} \leq C\varepsilon^{-1/K_2},$$

while the  $L^2(Q_{\mathcal{T}})$  bound for  $\nabla(T^{(\tau)})^{\beta/2}$  and the Sobolev embedding  $H^1 \hookrightarrow L^6$  imply

$$\|T^{(\tau)}\|_{L^\beta(0, \mathcal{T}; L^{3\beta}(\Omega))} \leq C.$$

The previous two estimates yield by interpolation (we also employed the fact that  $\beta + (2/3)K_2 > \beta + 2$ , which follows from (130))

$$(173) \quad \|T^{(\tau)}\|_{L^{\beta+2}(Q_{\mathcal{T}})} \leq C \|T^{(\tau)}\|_{L^{\beta+(2/3)K_2}(Q_{\mathcal{T}})} \leq C \|T^{(\tau)}\|_{L^\infty(0, \mathcal{T}; L^{K_2}(\Omega))}^{\frac{(2/3)K_2}{\beta+(2/3)K_2}} \|T^{(\tau)}\|_{L^\beta(0, \mathcal{T}; L^{3\beta}(\Omega))}^{\frac{\beta}{\beta+(2/3)K_2}} \\ \leq C\varepsilon^{-\frac{(2/3)}{\beta+(2/3)K_2}}.$$

Furthermore, from (170) and the fact that

$$(T-1) \log T \geq 0 \quad \forall T > 0, \quad \inf_{0 < T < 1/2} (T-1) \log T = \frac{1}{2} \log 2 > 0,$$

- 3 we also deduce

$$(174) \quad \varepsilon \int_{Q_{\mathcal{T}} \cap \{T^{(\tau)} < 1\}} (T^{(\tau)})^{-K_3-1} \log \frac{1}{T^{(\tau)}} dx dt + \varepsilon^{\frac{1}{1+K_3}} \left\| \frac{1}{T^{(\tau)}} \right\|_{L^{1+K_3}(0, \mathcal{T}; W^{1,1+K_3}(\Omega))} \leq C.$$

Let us now find an estimate for the pressure. From (65), (159), (172) it follows

$$\|\nabla p^{(\tau)}\|_{L^2(0, \mathcal{T}; L^{\frac{2K_2}{K_2+1}}(\Omega))} \leq C(\varepsilon) \|\sqrt{T^{(\tau)}}\|_{L^\infty(0, \mathcal{T}; L^{2K_2}(\Omega))} \left\| \sqrt{\frac{k_r(S^{(\tau)})}{T^{(\tau)}}} \nabla p^{(\tau)} \right\|_{L^2(Q_{\mathcal{T}})} \leq C(\varepsilon),$$

- 4 so via the  $L^\infty(0, T; L^1(\Omega))$  bound for  $p^{(\tau)}$  in (60) and Poincaré's Lemma

$$(175) \quad \|p^{(\tau)}\|_{L^2(0, \mathcal{T}; W^{1, \frac{2K_2}{K_2+1}}(\Omega))} \leq C(\varepsilon).$$

Sobolev's embedding  $W^{1, \frac{2K_2}{K_2+1}}(\Omega) \hookrightarrow L^{\frac{6K_2}{K_2+3}}(\Omega)$  implies

$$\|p^{(\tau)}\|_{L^2(0, \mathcal{T}; L^{\frac{6K_2}{K_2+3}}(\Omega))} \leq C(\varepsilon).$$

The estimate above together with the  $L^\infty(0, \mathcal{T}; L^1(\Omega))$  bound for  $p^{(\tau)}$  coming from (170) leads (by interpolation) to

$$(176) \quad \|p^{(\tau)}\|_{L^{2/\theta}(0, \mathcal{T}; L^\xi(\Omega))} \leq C(\varepsilon), \quad \frac{1}{\xi} = 1 - \theta + \theta \frac{K_2 + 3}{6K_2}, \quad \frac{3(K_2 + 1)}{5K_2 - 3} < \theta < 1.$$

- 5 We point out that thanks to (130) it holds that  $K_2 > 3$ , which implies  $\frac{3(K_2+1)}{5K_2-3} < 1$ . Therefore one  
 6 can choose  $\frac{3(K_2+1)}{5K_2-3} < \theta < 1$  in (176).

Another estimate for  $p^{(\tau)}$  can be found by combining (65) and (173). Noticing that  $\beta + 2 \geq 3$  and (159), (30) hold, one finds

$$\|\nabla p^{(\tau)}\|_{L^{3/2}(Q_{\mathcal{T}})} \leq C\varepsilon^{-\alpha_r} \|\sqrt{T^{(\tau)}}\|_{L^6(Q_{\mathcal{T}})} \left\| \sqrt{\frac{k_r(S^{(\tau)})}{T^{(\tau)}}} \nabla p^{(\tau)} \right\|_{L^2(Q_{\mathcal{T}})} \leq C\varepsilon^{-\alpha_r},$$

- 7 so via the  $L^\infty(0, T; L^1(\Omega))$  bound for  $p^{(\tau)}$  in (60) and Poincaré's Lemma

$$(177) \quad \|p^{(\tau)}\|_{L^{\frac{3}{2}}(0, \mathcal{T}; W^{1, \frac{3}{2}}(\Omega))} \leq C\varepsilon^{-\alpha_r}.$$

Sobolev's embedding  $W^{1,3/2}(\Omega) \hookrightarrow L^3(\Omega)$  yields

$$\|p^{(\tau)}\|_{L^{3/2}(0,\mathcal{T};L^3(\Omega))} \leq C\varepsilon^{-\alpha_r},$$

- 1 which, together with the  $L^\infty(0, \mathcal{T}; L^1(\Omega))$  bound for  $p^{(\tau)}$  coming from (60), allows us to deduce by  
2 interpolation that

$$(178) \quad \|p^{(\tau)}\|_{L^2(Q_{\mathcal{T}})} \leq \|p^{(\tau)}\|_{L^\infty(0,\mathcal{T};L^1(\Omega))}^{1/4} \|p^{(\tau)}\|_{L^{3/2}(0,\mathcal{T};L^3(\Omega))}^{3/4} \leq C\varepsilon^{-\frac{3}{4}\alpha_r}.$$

Inequality (161) implies in the new notation (upon summation in  $k$ )

$$\begin{aligned} & \int_{\Omega} (1 + |\nabla f(S^{(\tau)}(t))|^2)^{2/3} dx + \int_0^t \int_{\Omega} \frac{|f'(S^{(\tau)})P'_{c,\varepsilon}(S^{(\tau)})| |\nabla S^{(\tau)}|^2}{(1 + |\nabla f(S^{(\tau)})|^2)^{1/3}} dx dt' \\ & \leq \int_{\Omega} (1 + |\nabla f(S^{in,\varepsilon})|^2)^{2/3} dx + \int_0^t \int_{\Omega} |\nabla p^{(\tau)}|^{4/3} dx dt' \\ & \quad + C \int_0^t \int_{\Omega} (1 + |\nabla f(S^{(\tau)})|^2)^{2/3} dx dt, \quad t \in (0, \mathcal{T}]. \end{aligned}$$

Gronwall's inequality (in its discrete version) and (177) lead to

$$(179) \quad \|\nabla f(S^{(\tau)})\|_{L^\infty(0,\mathcal{T};L^{\frac{4}{3}}(\Omega))}^{4/3} + \int_0^{\mathcal{T}} \int_{\Omega} \frac{|f'(S^{(\tau)})P'_{c,\varepsilon}(S^{(\tau)})| |\nabla S^{(\tau)}|^2}{(1 + |\nabla f(S^{(\tau)})|^2)^{1/3}} dx dt \leq C(\varepsilon).$$

- 3 Thanks to (35), the above estimate implies

$$(180) \quad \|\nabla S^{(\tau)}\|_{L^\infty(0,\mathcal{T};L^{4/3}(\Omega))} \leq C(\varepsilon).$$

From (160), (60) as well as the fact that  $f$  is upper bounded, one easily deduces that

$$(181) \quad \|f(S^{(\tau)})\|_{L^\infty(0,\mathcal{T};L^1(\Omega))} \leq C(\varepsilon).$$

- 4 From (179), (181) and Poincaré inequality we get

$$(182) \quad \|f(S^{(\tau)})\|_{L^\infty(0,\mathcal{T};W^{1,4/3}(\Omega))} \leq C(\varepsilon).$$

- 5 Furthermore, thanks to (159) and the fact that  $P_{c,\varepsilon}$  is nonnegative, decreasing and continuous in  
6  $(0, 1)$ , we immediately deduce that

$$(183) \quad \|P_{c,\varepsilon}(S^{(\tau)})\|_{L^\infty(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

- 7 From (169), (178), (183), it follows

$$(184) \quad \|D_\tau f(S^{(\tau)})\|_{L^2(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

- 8 Bounds (182), (184) allow us to apply Aubin-Lions Lemma in the version of [14, Thr. 1] and obtain  
9 (up to subsequences)

$$(185) \quad f(S^{(\tau)}) \rightarrow f^* \quad \text{strongly in } L^1(Q_{\mathcal{T}}), \text{ as } \tau \rightarrow 0.$$

- 10 Being  $f$  invertible, this implies that  $S^{(\tau)}$  is a.e. convergent in  $Q_{\mathcal{T}}$ . The uniform  $L^\infty$  bounds for  
11  $S^{(\tau)}$  yield

$$(186) \quad S^{(\tau)} \rightarrow S \quad \text{strongly in } L^q(Q_{\mathcal{T}}), \quad \forall q < \infty.$$

- 12 Given (159), (185), (186) and the fact that  $\lim_{s \rightarrow 1} f(s) = -\infty$ , Fatou's Lemma allows us to state

$$(187) \quad \varepsilon \leq S < 1 \quad \text{a.e. in } Q_{\mathcal{T}}.$$

From (133), (178) it follows

$$(188) \quad \|\rho^{(\tau)}\|_{L^{2K_1}(Q_{\mathcal{T}})} \leq C\varepsilon^{-\frac{3\alpha_r}{4K_1}}.$$



Let us now find an estimate for  $\nabla \rho_i$ ,  $i = 1, \dots, N$ . Differentiating (162) in  $x$  leads to

$$(189) \quad \begin{aligned} \nabla \log \rho_i^{(\tau)} + \gamma \frac{\nabla(\rho^{(\tau)})^{\gamma-1}}{T^{(\tau)}} - \gamma \frac{(\rho^{(\tau)})^{\gamma-1}}{T^{(\tau)}} \nabla \log T^{(\tau)} + \varepsilon K_1 \nabla(\rho^{(\tau)})^{K_1-1} \\ = \nabla \frac{\mu_i^{(\tau)}}{T^{(\tau)}} + c_w \nabla \log T^{(\tau)}, \quad i = 1, \dots, N. \end{aligned}$$

Easy computations allow us to equivalently rewrite the above identity as

$$\begin{aligned} \sum_{j=1}^N \mathcal{M}_{ij} \nabla \rho_j^{(\tau)} &= \mathcal{F}_i, \quad i = 1, \dots, N, \\ \mathcal{M}_{ij} &= \frac{\delta_{ij}}{\rho_i^{(\tau)}} + \frac{\gamma(\gamma-1)}{T^{(\tau)}} (\rho^{(\tau)})^{\gamma-2} + \varepsilon K_1 (K_1 - 1) (\rho^{(\tau)})^{K_1-2}, \\ \mathcal{F}_i &= \nabla \frac{\mu_i^{(\tau)}}{T^{(\tau)}} + \left( c_w + \frac{\gamma(\rho^{(\tau)})^{\gamma-1}}{T^{(\tau)}} \right) \nabla \log T^{(\tau)}. \end{aligned}$$

Since  $\sum_{i,j=1}^N \mathcal{M}_{ij} v_i v_j \geq \sum_{i=1}^N \frac{v_i^2}{\rho_i^{(\tau)}}$ , we deduce

$$\begin{aligned} 4 \sum_{i=1}^N |\nabla \sqrt{\rho_i^{(\tau)}}|^2 &\leq \sum_{i,j=1}^N \mathcal{M}_{ij} \nabla \rho_i^{(\tau)} \cdot \nabla \rho_j^{(\tau)} = \sum_{i=1}^N \mathcal{F}_i \cdot \nabla \rho_i^{(\tau)} \\ &\leq 2 \left( \sum_{i=1}^N |\mathcal{F}_i|^2 \rho_i \right)^{1/2} \left( \sum_{i=1}^N |\nabla \sqrt{\rho_i^{(\tau)}}|^2 \right)^{1/2}, \end{aligned}$$

which implies

$$\begin{aligned} \sum_{i=1}^N |\nabla \sqrt{\rho_i^{(\tau)}}| &\leq C \sum_{i=1}^N |\mathcal{F}_i| \sqrt{\rho_i^{(\tau)}} \\ &\leq C \sqrt{\rho^{(\tau)}} \left| \nabla \frac{\vec{\mu}^{(\tau)}}{T^{(\tau)}} \right| + C \left( 1 + \frac{(\rho^{(\tau)})^{\gamma-1/2}}{T^{(\tau)}} \right) |\nabla \log T^{(\tau)}|. \end{aligned}$$

Multiplying the above estimate times  $\sqrt{\rho^{(\tau)}}$  leads to

$$(190) \quad \sum_{i=1}^N |\nabla \rho_i^{(\tau)}| \leq C \rho^{(\tau)} \left| \nabla \frac{\vec{\mu}^{(\tau)}}{T^{(\tau)}} \right| + C \left( 1 + \frac{(\rho^{(\tau)})^\gamma}{T^{(\tau)}} \right) |\nabla \log T^{(\tau)}|$$

From (130), (188), (174) it follows

$$\left\| \rho^{(\tau)} \nabla \frac{\vec{\mu}^{(\tau)}}{T^{(\tau)}} \right\|_{L^1(Q_\tau)} \leq \left\| \rho^{(\tau)} \right\|_{L^2(Q_\tau)} \left\| \nabla \frac{\vec{\mu}^{(\tau)}}{T^{(\tau)}} \right\|_{L^2(Q_\tau)} \leq C(\varepsilon),$$

as well as

$$\begin{aligned} \left\| \frac{(\rho^{(\tau)})^\gamma}{T^{(\tau)}} \nabla \log T^{(\tau)} \right\|_{L^1(Q_\tau)} &\leq C \left\| (\rho^{(\tau)})^\gamma \right\|_{L^{\frac{5K_1}{3\gamma}}(Q_\tau)} \left\| \frac{1}{T^{(\tau)}} \right\|_{L^{1+K_3}(Q_\tau)} \left\| \nabla \log T^{(\tau)} \right\|_{L^2(Q_\tau)} \\ &\leq C(\varepsilon). \end{aligned}$$

1 We deduce

$$(191) \quad \left\| \nabla \rho_i^{(\tau)} \right\|_{L^1(Q_\tau)} \leq C(\varepsilon), \quad i = 1, \dots, N.$$

From (22) and (130), (188) it follows  $\|\rho_i^{(\tau)}\|_{L^4(Q_{\mathcal{T}})} \leq C(\varepsilon)$  for  $i = 1, \dots, N$ . From this fact, the uniform  $L^\infty$  bounds for  $S^{(\tau)}$ , as well as (180), (191), we conclude

$$\|\nabla(S^{(\tau)}\rho_i^{(\tau)})\|_{L^1(Q_{\mathcal{T}})} \leq C(\varepsilon), \quad i = 1, \dots, N,$$

1 which, thanks to the uniform  $L^1$  bound for  $S^{(\tau)}\rho^{(\tau)}$ , yields

$$(192) \quad \|S^{(\tau)}\rho_i^{(\tau)}\|_{L^1(0, \mathcal{T}; W^{1,1}(\Omega))} \leq C(\varepsilon), \quad i = 1, \dots, N.$$

An estimate for  $D_\tau(S^{(\tau)}\rho_i^{(\tau)})$ ,  $i = 1, \dots, N$ , can be derived from (167) in a rather straightforward way, just like we did in the previous section (devoted to the limit  $\delta \rightarrow 0$ ), although now one has to keep in mind the dependency of every quantity on time. The following bounds are easily derived from (38), (42), (62), (64):

$$\|\mathbf{J}_i^{(\tau)}\|_{L^2(Q_{\mathcal{T}})} + \|r_{i,\varepsilon}^{(\tau)}\|_{L^{a/(a-1)}(Q_{\mathcal{T}})} \leq C, \quad i = 1, \dots, N.$$

On the other hand, from (6), (177), (188) and the fact that  $K_1 > \frac{6}{5}\gamma > \frac{12}{5}$  (see (130), (6)) it follows

$$\exists q > 1 : \quad \|\rho_i^{(\tau)}\mathbf{v}^{(\tau)}\|_{L^q(Q_{\mathcal{T}})} \leq C\|\rho^{(\tau)}\|_{L^{\frac{5}{3}K_1}(Q_{\mathcal{T}})}\|\nabla p^{(\tau)}\|_{L^{\frac{4}{3}}(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

We deduce

$$\|D_\tau(S^{(\tau)}\rho_i^{(\tau)}\Phi)\|_{L^q(0, \mathcal{T}; W^{1,q}(\Omega)')} \leq C(\varepsilon), \quad i = 1, \dots, N.$$

2 Since  $\Phi \in L^\infty(0, \mathcal{T}; W^{1,\infty}(\Omega))$ , from the estimate above and (192) we conclude via Aubin-Lions  
 3 Lemma that (up to subsequences)  $S^{(\tau)}\rho_i^{(\tau)}\Phi$  is strongly convergent in  $L^1(Q_{\mathcal{T}})$ , for  $i = 1, \dots, N$ .  
 4 Given the uniform  $L^\infty$  bounds for  $S^{(\tau)}$  and (188), the convergence holds in  $L^z(Q_{\mathcal{T}})$  for every  
 5  $z < \frac{5}{3}K_1$ . Furthermore, since  $\Phi$  does not depend on  $\tau$  and is uniformly positive, while  $S^{(\tau)}$  is  
 6 strongly convergent in  $L^q(Q_{\mathcal{T}})$  for every  $q < \infty$  and satisfies (159), we conclude

$$(193) \quad \rho_i^{(\tau)} \rightarrow \rho_i \quad \text{strongly in } L^z(Q_{\mathcal{T}}), \quad \forall z < \frac{5}{3}K_1, \quad i = 1, \dots, N.$$

7 We wish to point out that  $\rho_i > 0$  a.e. in  $Q_{\mathcal{T}}$  for  $i = 1, \dots, N$  thanks to Fatou's Lemma and (171).

Now we show that the temperature  $T^{(\tau)}$  is strongly convergent. We begin by estimating the energy flux  $\mathcal{J}_E^{(\tau)}$  given by

$$\mathcal{J}_E^{(\tau)} = ((\rho e)^{(\tau)} + p^{(\tau)})\mathbf{v}^{(\tau)} + \mathbf{q}^{(\tau)} - \varepsilon(1 + T^{(\tau)})\nabla \log T^{(\tau)} - \varepsilon \left| \nabla \frac{1}{T^{(\tau)}} \right|^{K_3-1} \nabla \frac{1}{T^{(\tau)}}.$$

Given the definitions of  $p^{(\tau)}$ ,  $(\rho e)^{(\tau)}$ ,  $\mathbf{v}^{(\tau)}$ , it holds

$$|((\rho e)^{(\tau)} + p^{(\tau)})\mathbf{v}^{(\tau)}| \leq C(p^{(\tau)} + 2p_{at})|\nabla p^{(\tau)}|$$

From (175), (176) it follows

$$\begin{aligned} & \|p^{(\tau)}|\nabla p^{(\tau)}|\|_{L^{\frac{2}{1+\theta}}(0, \mathcal{T}; L^{\xi_2}(\Omega))} \leq C(\varepsilon), \\ & \frac{1}{\xi_2} = \frac{1}{\xi} + \frac{K_2 + 1}{2K_2} = 1 - \theta + \theta \frac{K_2 + 3}{6K_2} + \frac{K_2 + 1}{2K_2}. \end{aligned}$$

Since  $\frac{3(K_2+1)}{5K_2-3} < \theta < 1$ , it follows that  $\xi_2 > 1$ ,  $2/(1+\theta) > 1$ . Therefore

$$\exists q > 1 : \quad \|p^{(\tau)}|\nabla p^{(\tau)}|\|_{L^q(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

As a consequence

$$(194) \quad \exists q > 1 : \quad \|((\rho e)^{(\tau)} + p^{(\tau)})\mathbf{v}^{(\tau)}\|_{L^q(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

Let us now consider the heat flux

$$\mathbf{q}^{(\tau)} = -\kappa(T^{(\tau)})\nabla T^{(\tau)} + \sum_{j=1}^N L_{0j}^{(\tau)} \nabla \frac{\mu_j^{(\tau)}}{T^{(\tau)}}.$$

It is clear that

$$\left\| \sum_{j=1}^N L_{0j}^{(\tau)} \nabla \frac{\mu_j^{(\tau)}}{T^{(\tau)}} \right\|_{L^2(Q_{\mathcal{T}})} \leq C.$$

On the other hand, from (36) one finds

$$\begin{aligned} |\kappa(T^{(\tau)})\nabla T^{(\tau)}| &\leq C(1 + (T^{(\tau)})^\beta) |\nabla T^{(\tau)}| \\ &\leq CT^{(\tau)} |\nabla \log T^{(\tau)}| + C(T^{(\tau)})^{1+\beta/2} |\nabla (T^{(\tau)})^{\beta/2}|. \end{aligned}$$

From (130), (173) it follows that  $T^{(\tau)}$ ,  $(T^{(\tau)})^{1+\beta/2}$  are bounded in  $L^{2+\omega}(Q_{\mathcal{T}})$  for some  $\omega > 0$  uniformly in  $\tau$ . Therefore from (62) one finds

$$(195) \quad \exists q > 1 : \quad \|\kappa(T^{(\tau)})\nabla T^{(\tau)}\|_{L^q(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

1 From (174) it follows immediately

$$(196) \quad \varepsilon \left\| \left| \nabla \frac{1}{T^{(\tau)}} \right|^{K_3-1} \nabla \frac{1}{T^{(\tau)}} \right\|_{L^{\frac{K_3+1}{K_3}}(Q_{\mathcal{T}})} \leq C\varepsilon^{\frac{1}{K_3+1}}.$$

2 Since the exponents  $q$  in (194), (195) do not depend on  $K_3$ , we conclude that, if  $K_3$  is large enough,  
3 the energy flux  $\mathcal{J}_E^{(\tau)}$  can be estimated as follows:

$$(197) \quad \|\mathcal{J}_E^{(\tau)}\|_{L^{\frac{K_3+1}{K_3}}(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

An estimate for the total energy  $\mathcal{E}^{(\tau)} = \Phi E_{f,\varepsilon}^{(\tau)} + (1 - \Phi)E_s^{(\tau)}$  is now required. Since

$$(198) \quad \begin{aligned} \mathcal{E}^{(\tau)} &= \Phi \left( S^{(\tau)}((\rho^{(\tau)})^\gamma + c_w \rho^{(\tau)} T^{(\tau)} + p_{at} + \varepsilon(\rho^{(\tau)})^{K_1}) - \int_{1/2}^{S^{(\tau)}} P_{c,\varepsilon}(\xi) d\xi \right) \\ &\quad + (1 - \Phi) (c_s T^{(\tau)} + \varepsilon(K_2 - 1)(T^{(\tau)})^{K_2}), \end{aligned}$$

if follows from (187) (also thanks to (22) and Young's inequality)

$$|\mathcal{E}^{(\tau)}| \leq C(\varepsilon) (1 + (\rho^{(\tau)})^{K_1} + (T^{(\tau)})^{K_2}).$$

From (130), (188), (173), it follows that

$$\exists q > 1 : \quad \|(\rho^{(\tau)})^{K_1} + (T^{(\tau)})^{K_2}\|_{L^q(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

4 It follows that  $\mathcal{E}^{(\tau)}$  is bounded in  $L^q(Q_{\mathcal{T}})$  uniformly in  $\tau$  for some  $q > 1$  which does not depend  
5 on  $K_3$ . Therefore choosing  $K_3$  large enough yields

$$(199) \quad \|\mathcal{E}^{(\tau)}\|_{L^{\frac{K_3+1}{K_3}}(Q_{\mathcal{T}})} \leq C(\varepsilon).$$

We aim to prove strong convergence of the temperature via the Div-Curl Lemma. For this reason we find it more convenient to work with time-continuous functions. With this idea in mind, we

define now a piecewise linear interpolation operator for the time variable. Given any  $u : [0, \mathcal{T}] \rightarrow X$  (where  $X$  is a generic Banach space), define

$$\mathcal{L}_\tau[u](t) = u(0)\chi_{\{0\}}(t) + \sum_{k=1}^N \left( u(t_{k-1})\frac{t_k - t}{\tau} + u(t_k)\frac{t - t_{k-1}}{\tau} \right) \chi_{(t_{k-1}, t_k]}(t),$$

$$t_k = k\tau, \quad k = 0, \dots, N, \quad N = T/\tau.$$

We point out that, for every function  $u : [0, \mathcal{T}] \rightarrow X$ , the following identity holds:

$$(200) \quad \frac{d}{dt} \mathcal{L}_\tau[u] = D_\tau u^{(\tau)},$$

- 1 where  $u^{(\tau)}$  is the piecewise constant approximation of  $u$  and  $D_\tau$  is the usual discrete time derivative  
2 operator.

Let us now define the vector fields

$$\mathcal{U}^{(\tau)} = (\mathcal{L}_\tau[\mathcal{E}^{(\tau)}], \mathcal{J}^{(\tau)}), \quad \mathcal{V}^{(\tau)}[G] = (G(1/T^{(\tau)}), 0, 0, 0),$$

where  $G \in W^{1,\infty}(\mathbb{R}_+)$  is arbitrary. From (197), (199) and the definition of  $\mathcal{L}_\tau$  it follows

$$\|\mathcal{U}^{(\tau)}\|_{L^q(Q_\mathcal{T})} \leq C(\varepsilon).$$

On the other hand, eq. (168) implies the following identity:

$$(201) \quad \int_0^\mathcal{T} \int_\Omega D_\tau(\mathcal{E}^{(\tau)})\psi dxdt - \int_0^\mathcal{T} \int_\Omega \mathcal{J}_E^{(\tau)} \cdot \nabla \psi dxdt$$

$$= -\varepsilon \int_0^\mathcal{T} \int_\Omega (1 + (T^{(\tau)})^{-K_3})(\log T^{(\tau)})\psi dxdt$$

$$\forall \psi \in L^{K_3+1}(0, \mathcal{T}; W_0^{1,K_3+1}(\Omega)),$$

which, thanks to (200), implies

$$\langle \operatorname{div}_{(t,x)} \mathcal{U}^{(\tau)}, \psi \rangle = -\varepsilon \int_0^\mathcal{T} \int_\Omega (1 + (T^{(\tau)})^{-K_3})(\log T^{(\tau)})\psi dxdt \quad \forall \psi \in C_c^1(Q_\mathcal{T}),$$

which, thanks to (174), yields

$$\|\operatorname{div}_{(t,x)} \mathcal{U}^{(\tau)}\|_{L^1(Q_\mathcal{T})} \leq C.$$

In particular,  $\operatorname{div}_{(t,x)} \mathcal{U}^{(\tau)}$  is relatively compact in  $W^{-1,r}(Q_\mathcal{T})$  for some  $r > 1$ . On the other hand, while  $\|\mathcal{V}^{(\tau)}[G]\|_{L^\infty(Q_\mathcal{T})} \leq C[G]$ , it holds

$$|\operatorname{curl}_{(t,x)} \mathcal{V}^{(\tau)}[G]| \leq C[G]|\nabla(1/T^{(\tau)})|,$$

which, together with (174), leads to

$$\|\operatorname{curl}_{(t,x)} \mathcal{V}^{(\tau)}[G]\|_{L^{K_3+1}(Q_\mathcal{T})} \leq C(\varepsilon).$$

- 3 In particular  $\operatorname{curl}_{(t,x)} \mathcal{V}[G]$  is relatively compact in  $W^{-1,r}(Q_\mathcal{T})$  for some  $r > 1$ .

At this point we are in the condition of applying the Div-Curl lemma and deduce

$$\overline{\mathcal{U}^{(\tau)} \cdot \mathcal{V}^{(\tau)}[G]} = \overline{\mathcal{U}^{(\tau)}} \cdot \overline{\mathcal{V}^{(\tau)}[G]}$$

that is

$$\overline{\mathcal{L}_\tau[\mathcal{E}^{(\tau)}]G(1/T^{(\tau)})} = \overline{\mathcal{L}_\tau[\mathcal{E}^{(\tau)}]} \overline{G(1/T^{(\tau)})} \quad \text{a.e. } Q_\mathcal{T}.$$

We now wish to get rid of the piecewise linear interpolation operator  $\mathcal{L}_\tau$ . Let  $\psi \in C_c^1(Q_\mathcal{T})$  arbitrary and define  $\omega^{(\tau)} = \mathcal{E}^{(\tau)} - \mathcal{L}_\tau[\mathcal{E}^{(\tau)}]$ . From the above identity it follows

$$\begin{aligned}
(202) \quad & \int_{Q_\mathcal{T}} (\overline{\mathcal{E}^{(\tau)} G(1/T^{(\tau)})} - \overline{\mathcal{E}^{(\tau)}} \overline{G(1/T^{(\tau)})}) \psi dxdt \\
&= \int_{Q_\mathcal{T}} ((\overline{\mathcal{E}^{(\tau)} - \mathcal{L}_\tau[\mathcal{E}^{(\tau)}]} G(1/T^{(\tau)})) \psi dxdt \\
&\quad - \int_{Q_\mathcal{T}} ((\overline{\mathcal{E}^{(\tau)} - \mathcal{L}_\tau[\mathcal{E}^{(\tau)}]} \overline{G(1/T^{(\tau)})}) \psi dxdt \\
&= \lim_{\tau, \sigma \rightarrow 0} \frac{1}{2} \int_{Q_\mathcal{T}} (\omega^{(\tau)} - \omega^{(\sigma)}) (G(1/T^{(\tau)}) - G(1/T^{(\sigma)})) \psi dxdt \\
&\leq \lim_{\tau, \sigma \rightarrow 0} \frac{1}{2} \|\omega^{(\tau)} - \omega^{(\sigma)}\|_{L^{\frac{K_3+1}{K_3}}(0, \mathcal{T}; W^{-1, \frac{K_3+1}{K_3}}(\Omega))} \\
&\quad \times \|(G(1/T^{(\tau)}) - G(1/T^{(\sigma)})) \psi\|_{L^{K_3+1}(0, \mathcal{T}; W^{1, K_3+1}(\Omega))} \\
&\leq C \lim_{\tau \rightarrow 0} \|\omega^{(\tau)}\|_{L^{\frac{K_3+1}{K_3}}(0, \mathcal{T}; W^{-1, \frac{K_3+1}{K_3}}(\Omega))} \|G(1/T^{(\tau)}) \psi\|_{L^{K_3+1}(0, \mathcal{T}; W^{1, K_3+1}(\Omega))}.
\end{aligned}$$

Since  $G \in W^{1, \infty}(\mathbb{R}_+)$  and (174) holds, it follows

$$(203) \quad \|G(1/T^{(\tau)}) \psi\|_{L^{K_3+1}(0, \mathcal{T}; W^{1, K_3+1}(\Omega))} \leq C(\varepsilon) \|\psi\|_{L^{K_3+1}(0, \mathcal{T}; W^{1, K_3+1}(\Omega)) \cap L^\infty(Q_\mathcal{T})}.$$

On the other hand, since

$$\begin{aligned}
\omega^{(\tau)}(t) &= \mathcal{E}^{(\tau)}(t) - \mathcal{L}_\tau[\mathcal{E}^{(\tau)}](t) \\
&= \sum_{k=1}^N \frac{t_k - t}{\tau} (\mathcal{E}^{(\tau)}(t_{k-1}) - \mathcal{E}^{(\tau)}(t_k)) \chi_{(t_{k-1}, t_k]}(t) = -\tau \sum_{k=1}^N \frac{t_k - t}{\tau} (D_\tau \mathcal{E}^{(\tau)})(t_k) \chi_{(t_{k-1}, t_k]}(t),
\end{aligned}$$

therefore

$$\begin{aligned}
& \|\omega^{(\tau)}\|_{L^{\frac{K_3+1}{K_3}}(0, \mathcal{T}; W^{-1, \frac{K_3+1}{K_3}}(\Omega))} \\
& \leq \tau \sum_{k=1}^N \left\| \frac{t_k - \cdot}{\tau} \chi_{(t_{k-1}, t_k]} \right\|_{L^{\frac{K_3+1}{K_3}}(0, \mathcal{T})} \|D_\tau \mathcal{E}^{(\tau)}(t_k)\|_{W^{-1, \frac{K_3+1}{K_3}}(\Omega)} \\
& \leq \tau^{1 + \frac{K_3}{K_3+1}} \sum_{k=1}^N \|D_\tau \mathcal{E}^{(\tau)}(t_k)\|_{W^{-1, \frac{K_3+1}{K_3}}(\Omega)} = \tau^{\frac{K_3}{K_3+1}} \|D_\tau \mathcal{E}^{(\tau)}\|_{L^1(0, \mathcal{T}; W^{-1, \frac{K_3+1}{K_3}}(\Omega))}.
\end{aligned}$$

From (174), (197), (201) we deduce

$$\|\omega^{(\tau)}\|_{L^{\frac{K_3+1}{K_3}}(0, \mathcal{T}; W^{-1, \frac{K_3+1}{K_3}}(\Omega))} \leq C \tau^{\frac{K_3}{K_3+1}},$$

which, together with (202), (203) leads to

$$\int_{Q_\mathcal{T}} (\overline{\mathcal{E}^{(\tau)} G(1/T^{(\tau)})} - \overline{\mathcal{E}^{(\tau)}} \overline{G(1/T^{(\tau)})}) \psi dxdt \leq 0 \quad \forall \psi \in C_c^1(Q_\mathcal{T}),$$

which easily implies

$$(204) \quad \overline{\mathcal{E}^{(\tau)} G(1/T^{(\tau)})} - \overline{\mathcal{E}^{(\tau)}} \overline{G(1/T^{(\tau)})} = 0 \quad \text{a.e. in } Q_\mathcal{T}.$$

From the (198) and the strong convergence of  $S^{(\tau)}$ ,  $\rho^{(\tau)}$  it follows

$$(205) \quad \begin{aligned} & (c_w \Phi \rho + c_s(1 - \Phi)) \left( \overline{T^{(\tau)} G(1/T^{(\tau)})} - \overline{T^{(\tau)} G(1/T^{(\tau)})} \right) \\ & + \varepsilon(K_2 - 1)(1 - \Phi) \left( \overline{(T^{(\tau)})^{K_2} G(1/T^{(\tau)})} - \overline{(T^{(\tau)})^{K_2} G(1/T^{(\tau)})} \right) = 0 \\ & \text{a.e. in } Q_{\mathcal{T}}. \end{aligned}$$

Choose now  $G(s) = \arctan(s)$ ,  $s \geq 0$ . The resulting function  $s \in \mathbb{R}_+ \mapsto G(1/s) \in \mathbb{R}_+$  is strictly decreasing and strictly convex. From [17, Thr. 10.19] it follows

$$\begin{aligned} \overline{T^{(\tau)} G(1/T^{(\tau)})} - \overline{T^{(\tau)} G(1/T^{(\tau)})} &\leq 0 \quad \text{a.e. in } Q_{\mathcal{T}}, \\ \overline{(T^{(\tau)})^{K_2} G(1/T^{(\tau)})} - \overline{(T^{(\tau)})^{K_2} G(1/T^{(\tau)})} &\leq 0 \quad \text{a.e. in } Q_{\mathcal{T}}, \end{aligned}$$

which, together with (205), yields

$$\begin{aligned} \overline{T^{(\tau)} G(1/T^{(\tau)})} - \overline{T^{(\tau)} G(1/T^{(\tau)})} &= \overline{(T^{(\tau)})^{K_2} G(1/T^{(\tau)})} - \overline{(T^{(\tau)})^{K_2} G(1/T^{(\tau)})} = 0 \\ &\text{a.e. in } Q_{\mathcal{T}}. \end{aligned}$$

From the above identity and from [17, Thr. 10.20], we conclude that  $T^{(\tau)}$  is (up to subsequences) a.e. convergent in  $Q_{\mathcal{T}}$ . The bounds (173), (174) allow us to deduce by dominated convergence that

$$\begin{aligned} T^{(\tau)} &\rightarrow T \quad \text{strongly in } L^{q_1}(Q_{\mathcal{T}}) \quad \forall q_1 < \beta + \frac{2}{3}K_2, \\ \frac{1}{T^{(\tau)}} &\rightarrow \frac{1}{T} \quad \text{strongly in } L^{q_2}(Q_{\mathcal{T}}) \quad \forall q_2 < K_3 + 1, \\ &T > 0 \quad \text{a.e. in } Q_{\mathcal{T}}. \end{aligned}$$

Since we have already estimated the terms appearing in (167)–(169), we can now take the limit  $\tau \rightarrow 0$  in (167)–(169) and obtain

$$(206) \quad \begin{aligned} & \int_0^{\mathcal{T}} \langle \partial_t(S\rho_i), \Phi\varphi \rangle dt - \int_0^{\mathcal{T}} \int_{\Omega} (\rho_i \mathbf{v} + \mathbf{J}_i) \cdot \nabla \varphi dx dt \\ & - \int_0^{\mathcal{T}} \int_{\Omega} r_{i,\varepsilon} \varphi dx dt + \int_0^{\mathcal{T}} \int_{\partial\Omega} \sum_{\ell=1}^N b_{i\ell} \left( \frac{\mu_{\ell}}{T} - \frac{\mu_{0,\ell}}{T_0} \right) \varphi ds dt \\ & = -\varepsilon \int_0^{\mathcal{T}} \int_{\Omega} \left( \nabla \frac{\mu_i}{T} \cdot \nabla \varphi + \frac{\mu_i}{T} \varphi \right) dx dt, \end{aligned}$$

$$(207) \quad \begin{aligned} & \int_0^{\mathcal{T}} \langle \partial_t(\Phi E_{f,\varepsilon} + (1 - \Phi)E_s)\psi \rangle dt - \int_0^{\mathcal{T}} \int_{\Omega} ((\rho e + p)\mathbf{v} + \mathbf{q}) \cdot \nabla \psi dx dt \\ & + \alpha \int_0^{\mathcal{T}} \int_{\partial\Omega} (T - T_0)\psi ds dt \\ & = -\varepsilon \int_0^{\mathcal{T}} \int_{\Omega} ((1 + T)\nabla \log T \cdot \nabla \psi + (1 + T^{-K_3})(\log T)\psi) dx dt \\ & - \varepsilon \int_0^{\mathcal{T}} \int_{\Omega} T^{-K_3} |\nabla \log T|^{K_3-1} \nabla \log T \cdot \nabla \psi dx dt, \end{aligned}$$

$$(208) \quad \partial_t f(S) + P_{c,\varepsilon}(S) + p = 0.$$

- 1 Let  $\zeta \in C^1(\overline{Q_{\mathcal{T}}})$  arbitrary. Let us choose  $\phi = \frac{\mu_i}{T}\zeta$ ,  $\psi = -\frac{1}{T}\zeta$  in (206), (207), respectively. We point
- 2 out that thanks to (174), (197), the choice  $\psi = -\frac{1}{T}\zeta$  as test function in (207) is admissible.

By proceeding like in the proof of Lemma 2 we obtain the approximate entropy balance equation:

$$\begin{aligned}
(209) \quad & \int_0^{\mathcal{T}} \langle \partial_t [\Phi S(\rho\eta) + (1 - \Phi)(\rho\eta)_s], \zeta \rangle dt \\
& - \int_0^{\mathcal{T}} \int_{\Omega} \left( (\rho\eta)\mathbf{v} - \sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i + \frac{\mathbf{q}}{T} - \varepsilon \sum_{i=1}^N \frac{\mu_i}{T} \nabla \frac{\mu_i}{T} \right. \\
& \quad \left. - \varepsilon \frac{1+T}{T} \nabla \log T - \varepsilon T^{-K_3-1} |\nabla \log T|^{K_3-1} \nabla \log T \right) \cdot \nabla \zeta dx dt \\
& = \int_0^{\mathcal{T}} \int_{\Omega} \left( \sum_{i,j=1}^N L_{ij} \nabla \left( \frac{\mu_i}{T} \right) \cdot \nabla \left( \frac{\mu_j}{T} \right) + \kappa(T) |\nabla \log T|^2 \right) \zeta dx dt \\
& \quad + \int_0^{\mathcal{T}} \left( \frac{K}{T} \lambda(S, T) |\nabla p|^2 - \Phi \frac{1}{T} f'(S) (\partial_t S)^2 - \sum_{i=1}^N r_{i,\varepsilon} \frac{\mu_i}{T} \right) \zeta dx dt \\
& \quad + \int_0^{\mathcal{T}} \int_{\partial\Omega} \left( \alpha \frac{T_0 - T}{T} + \sum_{i,j=1}^N b_{ij} \frac{\mu_i}{T} \left( \frac{\mu_j}{T} - \frac{\mu_{0,j}}{T_0} \right) \right) \zeta d\sigma dt \\
& \quad + \varepsilon \int_0^{\mathcal{T}} \int_{\Omega} \left( \left| \nabla \frac{\bar{\mu}}{T} \right|^2 + \left| \frac{\bar{\mu}}{T} \right|^2 + \frac{1+T}{T} |\nabla \log T|^2 + \left| \nabla \frac{1}{T} \right|^{K_3+1} \right. \\
& \quad \left. + (1 + T^{-K_3}) \frac{1}{T} \log \frac{1}{T} \right) \zeta dx dt, \quad \forall \zeta \in C^1(\overline{Q_{\mathcal{T}}}).
\end{aligned}$$

Choosing  $\psi = \psi(t) = \chi_{[0,t]}(t')$  in (207) yields the approximate integrated total energy balance:

$$\begin{aligned}
(210) \quad & \int_{\Omega} (\Phi E_{f,\varepsilon}(t) + (1 - \Phi) E_s(t)) dx + \alpha \int_0^t \int_{\Omega} (T - T_0) dx dt' \\
& = \int_{\Omega} (\Phi E_{f,\varepsilon}^{in} + (1 - \Phi) E_s^{in}) dx - \varepsilon \int_0^t \int_{\Omega} (1 + T^{-K_3}) \log T dx dt', \\
& \quad t \in [0, \mathcal{T}].
\end{aligned}$$

1 **4.5. Limit  $\varepsilon \rightarrow 0$ .** This limit is essentially the content of the weak stability analysis. The final  
2 equations (13), (45)–(47) will result from taking the limit  $\varepsilon \rightarrow 0$  in (206), (208), (209), (210). We  
3 only need to check the following two things: (i) that the regularizing terms in the equations vanish  
4 as  $\varepsilon \rightarrow 0$ , and (ii) that the regularizing terms do not destroy the argument proving the strong  
5 convergence of the approximate solution.

6 *Step 1: error terms vanish.* Let us begin by showing that the regularizing terms in (206), (208),  
7 (209), (210) tend to zero as  $\varepsilon \rightarrow 0$  (with the exception of the fourth integral on the right-hand  
8 side of (209), which will converge towards a nonnegative Radon measure that will be included in  
9 the term  $\xi$  appearing in (47)).

Clearly (171) implies

$$-\varepsilon \int_0^{\mathcal{T}} \int_{\Omega} \left( \nabla \frac{\mu_i}{T} \cdot \nabla \varphi + \frac{\mu_i}{T} \varphi \right) dx dt \rightarrow 0 \quad \forall \varphi \in L^2(0, \mathcal{T}; H^1(\Omega)).$$

Next, let us consider the regularizing term in (208). This term is only present if  $k_p = 0$  in (34)  
(see the definition (138) of  $P_{c,\varepsilon}$ ). Let  $s_0$  like in (33). Since we are assuming  $k_p = 0$ , (34) implies  
that  $s_1 \in (0, s_0)$  exists such that  $\inf_{(0,s_1)} P_c > p_{at}$ . Let  $\zeta : [0, 1] \rightarrow [0, 1]$  be a cutoff such that  $\zeta = 1$

in  $[0, s_1/2]$ ,  $\zeta = 0$  in  $[s_1, 1]$ ,  $\zeta$  is nonincreasing. Let us test (208) against  $-\zeta(S^{(\varepsilon)}) \log S^{(\varepsilon)}$  and recall that we consider a truncated initial datum for the saturation (given by (139)) in place of  $S^{in}$ :

$$\begin{aligned} & \int_{\Omega} \tilde{f}(S^{(\varepsilon)}(\mathcal{T})) dx - \int_0^{\mathcal{T}} \int_{\Omega} (P_c(S^{(\varepsilon)}) - \varepsilon \log S^{(\varepsilon)}) \zeta(S^{(\varepsilon)}) \log S^{(\varepsilon)} dx dt' \\ &= \int_{\Omega} \tilde{f}(S^{in, \varepsilon}) dx + \int_0^{\mathcal{T}} \int_{\Omega} p \zeta(S^{(\varepsilon)}) \log S^{(\varepsilon)} dx dt' \\ &\leq \int_{\Omega} \tilde{f}(S^{in, \varepsilon}) dx - p_{at} \int_0^{\mathcal{T}} \int_{\Omega} \zeta(S^{(\varepsilon)}) \log S^{(\varepsilon)} dx dt', \end{aligned}$$

where  $\tilde{f}(s) = \int_{s_1}^s f'(u) \zeta(u) (-\log u) du$  and  $s_0$  is like in (33). Since  $\tilde{f} \geq 0$ ,  $\inf_{(0, s_1)} P_c > p_{at}$  and (35) holds, it follows

$$\varepsilon \int_0^{\mathcal{T}} \int_{\Omega} (\log S^{(\varepsilon)})^2 \zeta(S^{(\varepsilon)}) dx dt \leq \int_{\Omega} \tilde{f}(S^{in, \varepsilon}) dx \leq C.$$

We deduce

$$(211) \quad \sqrt{\varepsilon} \|\log S^{(\varepsilon)}\|_{L^2(Q_{\mathcal{T}})} \leq C.$$

1 This implies that

$$(212) \quad P_{c, \varepsilon}(S^{(\varepsilon)}) - P_c(S^{(\varepsilon)}) \rightarrow 0 \quad \text{strongly in } L^2(Q_{\mathcal{T}}),$$

2 as well as

$$(213) \quad E_{f, \varepsilon} - E_f = \varepsilon \int_{1/2}^{S^{(\varepsilon)}} \log(\xi) d\xi \rightarrow 0 \quad \text{strongly in } L^\infty(Q_{\mathcal{T}}).$$

Let us now consider the error term in the entropy flux:

$$(214) \quad \begin{aligned} R^{(\varepsilon)} &= -\varepsilon \sum_{i=1}^N \frac{\mu_i^{(\varepsilon)}}{T^{(\varepsilon)}} \nabla \frac{\mu_i^{(\varepsilon)}}{T^{(\varepsilon)}} - \varepsilon \frac{1 + T^{(\varepsilon)}}{T^{(\varepsilon)}} \nabla \log T^{(\varepsilon)} \\ &\quad - \varepsilon (T^{(\varepsilon)})^{-K_3-1} |\nabla \log T^{(\varepsilon)}|^{K_3-1} \nabla \log T^{(\varepsilon)}. \end{aligned}$$

From (171) it follows

$$\varepsilon \sum_{i=1}^N \frac{\mu_i^{(\varepsilon)}}{T^{(\varepsilon)}} \nabla \frac{\mu_i^{(\varepsilon)}}{T^{(\varepsilon)}} \rightarrow 0 \quad \text{strongly in } L^{\frac{2a}{a+2}}(Q_{\mathcal{T}}).$$

From (62), (174) it follows immediately that

$$\varepsilon \frac{1 + T^{(\varepsilon)}}{T^{(\varepsilon)}} \nabla \log T^{(\varepsilon)} \rightarrow 0 \quad \text{strongly in } L^2(Q_{\mathcal{T}}).$$

Hölder inequality yields

$$\begin{aligned} & \varepsilon \int_{Q_{\mathcal{T}}} (T^{(\varepsilon)})^{-K_3-1} |\nabla \log T^{(\varepsilon)}|^{K_3} dx dt = \varepsilon \int_{Q_{\mathcal{T}}} (T^{(\varepsilon)})^{-1} |\nabla (T^{(\varepsilon)})^{-1}|^{K_3} dx dt \\ & \leq C \varepsilon \left( \int_{Q_{\mathcal{T}}} (T^{(\varepsilon)})^{-K_3-1} dx dt \right)^{\frac{1}{K_3+1}} \left( \int_{Q_{\mathcal{T}}} |\nabla (T^{(\varepsilon)})^{-1}|^{K_3+1} dx dt \right)^{\frac{K_3}{K_3+1}}. \end{aligned}$$

From (174) we deduce

$$\varepsilon \int_{Q_{\mathcal{T}}} (T^{(\varepsilon)})^{-K_3-1} |\nabla \log T^{(\varepsilon)}|^{K_3} dx dt \leq C \left( \varepsilon \int_{Q_{\mathcal{T}}} (T^{(\varepsilon)})^{-K_3-1} dx dt \right)^{\frac{1}{K_3+1}}.$$



On the other hand,

$$\begin{aligned} & \varepsilon \int_{Q_{\mathcal{T}}} (T^{(\varepsilon)})^{-K_3-1} dxdt \\ &= \varepsilon \int_{Q_{\mathcal{T}} \cap \{T^{(\tau)} < \varepsilon^{\frac{1}{2(K_3+1)}}\}} (T^{(\varepsilon)})^{-K_3-1} dxdt + \varepsilon \int_{Q_{\mathcal{T}} \cap \{T^{(\tau)} \geq \varepsilon^{\frac{1}{2(K_3+1)}}\}} (T^{(\varepsilon)})^{-K_3-1} dxdt. \end{aligned}$$

From (174) it follows

$$\begin{aligned} & \varepsilon \int_{Q_{\mathcal{T}} \cap \{T^{(\tau)} < \varepsilon^{\frac{1}{2(K_3+1)}}\}} (T^{(\varepsilon)})^{-K_3-1} dxdt \\ & \leq \frac{2(K_3+1)\varepsilon}{-\log \varepsilon} \int_{Q_{\mathcal{T}} \cap \{T^{(\tau)} < \varepsilon^{\frac{1}{2(K_3+1)}}\}} (T^{(\varepsilon)})^{-K_3-1} \log \frac{1}{T^{(\tau)}} dxdt \leq \frac{C}{-\log \varepsilon}, \end{aligned}$$

while

$$\varepsilon \int_{Q_{\mathcal{T}} \cap \{T^{(\tau)} \geq \varepsilon^{\frac{1}{2(K_3+1)}}\}} (T^{(\varepsilon)})^{-K_3-1} dxdt \leq C\sqrt{\varepsilon},$$

so we deduce

$$(215) \quad \varepsilon \int_{Q_{\mathcal{T}}} (T^{(\varepsilon)})^{-K_3-1} dxdt \leq C \left( \sqrt{\varepsilon} + \frac{1}{-\log \varepsilon} \right) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

- 1 Therefore the error term  $R^{(\varepsilon)}$  in the entropy flux tends to 0 strongly in  $L^1(Q_{\mathcal{T}})$  as  $\varepsilon \rightarrow 0$ . The
- 2 error term on the right-hand side of (209), on the other hand, thanks to the estimates of the error
- 3 terms previously derived, converges weakly-\* in  $\mathcal{M}(\overline{Q_{\mathcal{T}}})$  towards an element  $\tilde{\xi} \in \mathcal{M}(\overline{Q_{\mathcal{T}}})$  satisfying
- 4  $\langle \tilde{\xi}, \varphi \rangle \geq 0$  for every  $\varphi \in C(\overline{Q_{\mathcal{T}}})$  such that  $\varphi \geq 0$  in  $\overline{Q_{\mathcal{T}}}$ .

In the integrated energy balance (210) the error term on the right hand side easily tends to zero as  $\varepsilon \rightarrow 0$  thanks to (215):

$$\varepsilon \int_0^t \int_{\Omega} (1 + T^{-K_3}) \log T dxdt' \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

Let us now turn our attention to the error terms in the thermodynamic quantities defined in (132)–(136). Such error terms will tend to zero strongly in  $L^1(Q_{\mathcal{T}})$  as  $\varepsilon \rightarrow 0$  provided that

$$\varepsilon(\rho^{(\varepsilon)})^{K_1} + \varepsilon(T^{(\varepsilon)})^{K_2} \rightarrow 0 \quad \text{strongly in } L^1(Q_{\mathcal{T}}).$$

From (60), (130), (188) it follows by interpolation

$$\|\rho^{(\varepsilon)}\|_{L^{K_1}(Q_{\mathcal{T}})} \leq \|\rho^{(\varepsilon)}\|_{L^{\gamma}(Q_{\mathcal{T}})}^{\frac{\gamma}{2K_1-\gamma}} \|\rho^{(\varepsilon)}\|_{L^{2K_1}(Q_{\mathcal{T}})}^{\frac{2K_1-2\gamma}{2K_1-\gamma}} \leq C\varepsilon^{-\frac{3\alpha_r}{4K_1} \frac{2K_1-2\gamma}{2K_1-\gamma}},$$

which leads to

$$\int_{Q_{\mathcal{T}}} \varepsilon(\rho^{(\varepsilon)})^{K_1} dxdt \leq C\varepsilon^{1-\frac{3\alpha_r}{4} \frac{2K_1-2\gamma}{2K_1-\gamma}}.$$

From (30), (130) it follows that

$$\int_{Q_{\mathcal{T}}} \varepsilon(\rho^{(\varepsilon)})^{K_1} dxdt \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

On the other hand, (63) yields immediately that  $\varepsilon(T^{(\varepsilon)})^{K_2} \rightarrow 0$  strongly in  $L^1(Q_{\mathcal{T}})$  in the case that  $K_2 \leq \beta + \frac{2}{3}$ . Therefore let us consider the case  $K_2 > \beta + \frac{2}{3}$ ,  $K_2 < 3\beta$  (recall (130)). From

(63), (173) we deduce by interpolation

$$\|T^{(\varepsilon)}\|_{L^{K_2}(Q_{\mathcal{T}})} \leq \|T^{(\varepsilon)}\|_{L^{\beta+\frac{2}{3}}(Q_{\mathcal{T}})}^{1-\theta} \|T^{(\varepsilon)}\|_{L^{\beta+\frac{2}{3}K_2}(Q_{\mathcal{T}})}^{\theta} \leq C\varepsilon^{-\frac{(2/3)\theta}{\beta+(2/3)K_2}},$$

$$\theta = \frac{(3\beta + 2K_2)(K_2 - \beta - 2/3)}{2K_2(K_2 - 1)},$$

which implies

$$\varepsilon \int_{Q_{\mathcal{T}}} (T^{(\varepsilon)})^{K_2} dxdt \leq C\varepsilon^{1-\frac{(2/3)K_2\theta}{\beta+(2/3)K_2}}.$$

It is easy to see that  $1 - \frac{(2/3)K_2\theta}{\beta+(2/3)K_2} > 0$  if and only if  $\beta > 1/3$ , which is true by assumption (37).

This means

$$(216) \quad \int_{Q_{\mathcal{T}}} \varepsilon (T^{(\varepsilon)})^{K_2} dxdt \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

- 1 It follows that the error terms in (132)–(136) tend to zero strongly in  $L^1(Q_{\mathcal{T}})$  as  $\varepsilon \rightarrow 0$ . Step 1 is
- 2 therefore complete.
- 3 *Step 2: strong convergence holds.* The strong convergence of the saturation  $S^{(\varepsilon)}$  as well as the
- 4 positive lower bound for  $S^{(\varepsilon)}$  work exactly like in the weak stability in Section 3.

In the proof of convergence of the total densities, it holds that the only additional error term appearing in  $U^{(\varepsilon)}$  can be found in the definition of the pressure  $p^{(\varepsilon)}$ , and only contributes to the velocity field  $\mathbf{v}^{(\varepsilon)}$  via Darcy's Law; however for the regularized velocity field the same bounds hold, that hold for the not-regularized velocity, therefore the additional error term appearing in  $U^{(\varepsilon)}$  does not influence its integrability, which means that the vector field  $U^{(\varepsilon)}$  is still bounded in  $L^r(Q_{\mathcal{T}})$  for some  $r > 1$ . Furthermore

$$\operatorname{div}_{(t,x)} U^{(\varepsilon)} = \varepsilon \Delta \sum_{i=1}^N \frac{\mu_i^{(\varepsilon)}}{T^{(\varepsilon)}} - \varepsilon \sum_{i=1}^N \frac{\mu_i^{(\varepsilon)}}{T^{(\varepsilon)}},$$

- 5 and therefore, thanks to (171),  $\operatorname{div}_{(t,x)} U^{(\varepsilon)} \rightarrow 0$  strongly in  $L^2(0, \mathcal{T}; H^1(\Omega)')$  (and a fortiori strongly
- 6 in  $W^{-1,z}(Q_{\mathcal{T}})$  for some  $z > 1$ ). This means that the Div-Curl Lemma can still be applied to the
- 7 couple  $U^{(\varepsilon)}$ ,  $V^{(\varepsilon)}$  and argue like in the second step of the weak stability argument to deduce the
- 8 strong convergence of the total density  $\rho^{(\varepsilon)}$ .

- 9 In a similar way the proof of the strong convergence of the partial mass densities  $\rho_i^{(\varepsilon)}$ ,  $i =$
- 10  $1, \dots, N$ , can be adapted to the approximate system.

Concerning the strong convergence of the temperature, the vector field  $W^{(\varepsilon)}$  is still bounded in  $L^r(Q_{\mathcal{T}})$  for some  $r > 1$  since the only error term in  $W^{(\varepsilon)}$  (aside from the one in the velocity field, which can be ignored as in the case of  $U^{(\varepsilon)}$ ) appears in the skeleton entropy  $(\rho\eta)_{s,\varepsilon}$  and is proportional to  $\varepsilon(T^{(\varepsilon)})^{K_2-1}$ , which tends to zero (a fortiori, is bounded) strongly in  $L^{K_2/(K_2-1)}(Q_{\mathcal{T}})$  thanks to (216). On the other hand,

$$\operatorname{div}_{(t,x)} W^{(\varepsilon)} = -\operatorname{div}_x R^{(\varepsilon)} + \Xi^{(\varepsilon)},$$

- 11 where  $R^{(\varepsilon)}$  is given by (214) and  $\Xi$  is the right-hand side of (209). We have already proved that  $\Xi^{(\varepsilon)}$
- 12 is bounded in  $L^1(Q_{\mathcal{T}})$ , while  $R^{(\varepsilon)} \rightarrow 0$  strongly in  $L^1(Q_{\mathcal{T}})$ . We can therefore apply the Div-Curl
- 13 Lemma in the version of [11] to the couple  $(W^{(\varepsilon)}, Y^{(\varepsilon)}[G])$  and, by arguing like in the weak stability
- 14 analysis, prove the strong convergence of the temperature  $T^{(\varepsilon)}$ .

- 15 This finishes the proof of the existence theorem.

1

## 5. APPENDIX

2 To improve the readability of the paper, we present here some proofs and results which are  
3 rather technical, but are nevertheless needed for completeness.

**Proof of Proposition 5.** Since (6), (5) hold and  $\mu$  is uniformly positive, we deduce

$$\rho|\mathbf{v}| \leq C\rho\sqrt{\lambda(S,T)T} \sqrt{\frac{\lambda(S,T)}{T}} |\nabla p| \leq C(k_r(S))^{1/2} \rho T^{1/2} \sqrt{\frac{\lambda(S,T)}{T}} |\nabla p|.$$

From (30) and Hölder's inequality (notice that  $\frac{\gamma+6\beta+3\beta\gamma}{6\beta\gamma} = \frac{1}{\gamma} + \frac{1}{6\beta} + \frac{1}{2}$ ) we get

$$\begin{aligned} & \|\rho(t)\mathbf{v}(t)\|_{L^{\frac{6\beta\gamma}{\gamma+6\beta+3\beta\gamma}}(\Omega)}^{\frac{2\beta}{\beta+1}} \\ & \leq C\|S(t)^{1/\gamma}\rho(t)\|_{L^\gamma(\Omega)}^{\frac{2\beta}{\beta+1}} \|\sqrt{T}(t)\|_{L^{6\beta}(\Omega)}^{\frac{2\beta}{\beta+1}} \left\| \sqrt{\frac{\lambda(S(t),T(t))}{T(t)}} \nabla p(t) \right\|_{L^2(\Omega)}^{\frac{2\beta}{\beta+1}} \\ & \leq C \left( \operatorname{ess\,sup}_{t' \in [0, \mathcal{T}]} \|S(t')^{1/\gamma}\rho(t')\|_{L^\gamma(\Omega)}^{\frac{2\beta}{\beta+1}} \right) \|\sqrt{T}(t)\|_{L^{6\beta}(\Omega)}^{\frac{2\beta}{\beta+1}} \left\| \sqrt{\frac{\lambda(S(t),T(t))}{T(t)}} \nabla p(t) \right\|_{L^2(\Omega)}^{\frac{2\beta}{\beta+1}}, \end{aligned}$$

which, by integration and using Hölder's inequality, leads to

$$\begin{aligned} & \|\rho\mathbf{v}\|_{L^{\frac{2\beta}{\beta+1}}(0, \mathcal{T}; L^{\frac{6\beta\gamma}{\gamma+6\beta+3\beta\gamma}}(\Omega))}^{\frac{2\beta}{\beta+1}} \\ & \leq C\|S^{1/\gamma}\rho\|_{L^\infty(0, \mathcal{T}; L^\gamma(\Omega))}^{\frac{2\beta}{\beta+1}} \|\sqrt{T}\|_{L^{2\beta}(0, \mathcal{T}; L^{6\beta}(\Omega))}^{\frac{2\beta}{\beta+1}} \left\| \sqrt{\frac{\lambda}{T}} \nabla p \right\|_{L^2(Q_{\mathcal{T}})}^{\frac{2\beta}{\beta+1}} \\ & = C\|S\rho^\gamma\|_{L^\infty(0, \mathcal{T}; L^1(\Omega))}^{\frac{2\beta}{\gamma(\beta+1)}} \|T^{\frac{\beta}{2}}\|_{L^2(0, \mathcal{T}; L^6(\Omega))}^{\frac{2}{\beta+1}} \left\| \sqrt{\frac{\lambda}{T}} \nabla p \right\|_{L^2(Q_{\mathcal{T}})}^{\frac{2\beta}{\beta+1}}. \end{aligned}$$

We point out that  $\frac{2\beta}{\beta+1} > 1$ ,  $\frac{6\beta\gamma}{\gamma+6\beta+3\beta\gamma} > 1$  thanks to the assumptions (22), (31), (37) and the parameters  $\gamma, q, \beta$ . From the Sobolev embedding  $H^1(\mathbb{R}^d) \hookrightarrow L^6(\mathbb{R}^d)$  (valid for  $d \leq 3$ ) and (62) it follows

$$\|T^{\frac{\beta}{2}}\|_{L^2(0, \mathcal{T}; L^6(\Omega))} \leq C.$$

4 From the previous estimates as well as (61), (65) we obtain (78).

Let us now estimate  $\mathbf{J}_i$ . From (7) it follows

$$|\mathbf{J}_i| \leq C(|\nabla \log T| + |\nabla \Pi^N(\bar{\mu}/T)|), \quad i = 1, \dots, N,$$

5 so from (62), (64) we deduce (79).

6 From (11), (78), (79) we easily conclude that (80) and (81) hold.

7 An estimate (82) for  $\partial_t f(S)$  is immediately found from (13) and (60).

8 On the other hand, from the fact that  $p \leq C(\rho e)$  and the bound (61) we deduce that  $Sp$  is  
9 bounded in  $L^\infty(0, \mathcal{T}; L^1(\Omega))$ . As a consequence, multiplying (13) by  $S$  and taking the  $L^\infty(0, \mathcal{T}; L^1(\Omega))$   
10 norm leads to (83).

**Proof of Proposition 6.** From (26) it follows

$$\|S(\rho\eta)\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_T)} \leq \sum_{i=1}^N \|S\rho_i \log \rho_i\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_T)} + c_w \|S\rho(\log T + 1)\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_T)}.$$

However, for  $\delta \in (0, \gamma/2]$ , since  $S \leq S^{(1+\delta)/\gamma}$  and  $\rho_i \log \rho_i \leq C(1 + \rho^{1+\delta})$ , it holds

$$\begin{aligned} \sum_{i=1}^N \|S\rho_i \log \rho_i\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_T)} &\leq C + C \|S^{\frac{1+\delta}{\gamma}} \rho^{1+\delta}\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_T)} \\ &= C + C \|S^{1/\gamma} \rho\|_{L^{\frac{2\gamma(1+\delta)}{\gamma+2}}(Q_T)}^{1+\delta}. \end{aligned}$$

Since  $\gamma > 2$  and  $0 < \delta \leq \gamma/2$ , from (61) it follows

$$\sum_{i=1}^N \|S\rho_i \log \rho_i\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_T)} \leq C.$$

Moreover, Hölder's inequality yields

$$\|S\rho(\log T + 1)\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_T)} \leq \|S\rho\|_{L^\gamma(Q_T)} \|\log T + 1\|_{L^2(Q_T)} \leq C$$

1 thanks to (61), (62). We conclude

$$(217) \quad \|S(\rho\eta)\|_{L^{\frac{2\gamma}{\gamma+2}}(Q_T)} \leq C.$$

2 From (28), (62) it follows that  $(\rho\eta)_s$  is bounded in  $L^2(Q_T)$ , so we get (84).

We will now find an estimate for the entropy flux. We begin by considering

$$(\rho\eta)\mathbf{v} = K \left( \sum_{i=1}^N \rho_i \log \rho_i - c_w \rho(\log T + 1) \right) \lambda(S, T) \nabla p,$$

3 where the above equality holds thanks to (26), (6). Let  $s \in \mathbb{R}$  be such that

$$(218) \quad s > 1, \quad \frac{1}{\gamma} + \frac{1}{2\beta} + \frac{1}{2} < \frac{1}{s}.$$

The above definition makes sense since (37) holds. It follows via Hölder's inequality

$$\begin{aligned} \|(\rho\eta)\mathbf{v}\|_{L^s(Q_T)} &\leq C \sum_{i=1}^N \|\rho_i(\log \rho_i) \sqrt{\lambda(S, T)T}\|_{L^{\frac{2s}{2-s}}(Q_T)} \left\| \sqrt{\frac{\lambda(S, T)}{T}} \nabla p \right\|_{L^2(Q_T)} \\ &\quad + C \|\rho(\log T + 1) \sqrt{\lambda(S, T)T}\|_{L^{\frac{2s}{2-s}}(Q_T)} \left\| \sqrt{\frac{\lambda(S, T)}{T}} \nabla p \right\|_{L^2(Q_T)}. \end{aligned}$$

However, since  $\mu$  is uniformly positive and (65) holds, we get

$$\begin{aligned} \|(\rho\eta)\mathbf{v}\|_{L^s(Q_T)} &\leq C \sum_{i=1}^N \|\rho_i(\log \rho_i) \sqrt{k_r(S)T}\|_{L^{\frac{2s}{2-s}}(Q_T)} \\ &\quad + C \|\rho(\log T + 1) \sqrt{k_r(S)T}\|_{L^{\frac{2s}{2-s}}(Q_T)}. \end{aligned}$$

Now, since for every  $\delta > 0$  there exists  $C_\delta > 0$  such that  $x|\log x| \leq C_\delta(1+x^{1+\delta})$  for  $x > 0$ , Hölder's inequality and (218) allow us to state

$$\begin{aligned} \|(\rho\eta)\mathbf{v}\|_{L^s(Q_\tau)} &\leq C_\delta \sum_{i=1}^N \|(1+\rho_i^{1+\delta})\sqrt{k_r(S)}T^{1/2}\|_{L^{\frac{2s}{2-s}}(Q_\tau)} \\ &\quad + C_\delta \|\rho\sqrt{k_r(S)}(1+T^{(1+\delta)/2})\|_{L^{\frac{2s}{2-s}}(Q_\tau)} \\ &\leq C_\delta \sum_{i=1}^N \|(1+\rho_i^{1+\delta})\sqrt{k_r(S)}\|_{L^{\gamma/(1+\delta)}(Q_\tau)} \|T^{1/2}\|_{L^{2\beta}(Q_\tau)} \\ &\quad + C_\delta \|\rho\sqrt{k_r(S)}\|_{L^\gamma(Q_\tau)} \|(1+T^{(1+\delta)/2})\|_{L^{2\beta/(1+\delta)}(Q_\tau)}, \end{aligned}$$

- 1 for some  $\delta > 0$  small enough. Assumption (30) and bounds (61), (62) allow us to conclude (219)
- $$\exists s > 1 : \quad \|(\rho\eta)\mathbf{v}\|_{L^s(Q_\tau)} \leq C.$$

Let us then consider

$$-\sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i = -\sum_{i=1}^N \frac{\mu_i}{T} L_{i0} \nabla \frac{1}{T} + \sum_{i,j=1}^N \frac{\mu_i}{T} L_{ij} \nabla \frac{\mu_j}{T},$$

where the above equality comes from (7). Since (38) holds and  $L_{ij}$  is symmetric and positive semidefinite, we obtain via Cauchy-Schwartz and (39) that

$$\begin{aligned} \left| -\sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i \right| &\leq C \left| \Pi^N \frac{\vec{\mu}}{T} \right| |\nabla \log T| + C \left( \sum_{i,j=1}^N L_{ij} \frac{\mu_i \mu_j}{T^2} \right)^{\frac{1}{2}} \left( \sum_{i,j=1}^N L_{ij} \frac{\nabla \mu_i \cdot \nabla \mu_j}{T^2} \right)^{\frac{1}{2}} \\ &\leq C \left| \Pi^N \frac{\vec{\mu}}{T} \right| \left( |\nabla \log T| + \left| \nabla \Pi^N \frac{\vec{\mu}}{T} \right| \right). \end{aligned}$$

- 2 Given that (62), (64) hold, we obtain

$$(220) \quad \left\| -\sum_{i=1}^N \frac{\mu_i}{T} \mathbf{J}_i \right\|_{L^{\frac{2a}{a+2}}(Q_\tau)} \leq C.$$

Finally, let us consider

$$\frac{\mathbf{q}}{T} = -\kappa(T) \nabla \log T + \sum_{j=1}^N \frac{L_{0j}}{T} \nabla \frac{\mu_j}{T},$$

where the above equality comes from (8), (9). It follows from (36), that

$$|\kappa(T) \nabla \log T| \leq C |\nabla \log T| + CT^{\beta/2} |\nabla T^{\beta/2}|,$$

while (38) implies

$$\left| \sum_{j=1}^N \frac{L_{0j}}{T} \nabla \frac{\mu_j}{T} \right| \leq C \left| \nabla \Pi^N \frac{\vec{\mu}}{T} \right|.$$

From (62)–(64) and Hölder's inequality we conclude

$$(221) \quad \|\kappa(T) \nabla \log T\|_{L^{\frac{2+3\beta}{1+3\beta}}(Q_\tau)} + \left\| \sum_{j=1}^N \frac{L_{0j}}{T} \nabla \frac{\mu_j}{T} \right\|_{L^2(Q_\tau)} \leq C,$$

1 which leads to

$$(222) \quad \left\| \frac{\mathbf{q}}{T} \right\|_{L^{\frac{2+3\beta}{1+3\beta}}(Q_T)} \leq C.$$

2 Putting (219), (220), (222) together allows us to obtain the estimate for the entropy flux (85).

**Proof of Proposition 7.** Let us now find an estimate for  $\nabla\rho$ . Eq. (24) yields:

$$\nabla p = (T + \gamma(\gamma - 1)\rho^{\gamma-1})\nabla\rho + \rho\nabla T,$$

which implies

$$|\nabla\rho^\gamma| \leq |(T + \gamma(\gamma - 1)\rho^{\gamma-1})\nabla\rho| \leq \rho|\nabla T| + |\nabla p|.$$

From (30) and the uniform boundedness of  $\mu$  it follows

$$\sqrt{k_r(S)}|\nabla\rho^\gamma| \leq C\rho S^{1/\gamma}|\nabla T| + T^{1/2}\sqrt{\frac{\lambda(S, T)}{T}}|\nabla p|.$$

From (61), (62), (65), (63) (as well as from Hölder's inequality) it follows

$$\begin{aligned} \|\rho S^{1/\gamma}\nabla T\|_{L^{\frac{2\gamma}{2+\gamma}}(Q_T)} &\leq \|\rho S^{1/\gamma}\|_{L^\gamma(Q_T)}\|\nabla T\|_{L^2(Q_T)} \leq C, \\ \left\| T^{1/2}\sqrt{\frac{\lambda(S, T)}{T}}\nabla p \right\|_{L^{\frac{2\beta+4/3}{\beta+5/3}}(Q_T)} &\leq \|T^{1/2}\|_{L^{2\beta+4/3}(Q_T)} \left\| \sqrt{\frac{\lambda(S, T)}{T}}\nabla p \right\|_{L^2(Q_T)} \leq C, \end{aligned}$$

so we deduce

$$\|\sqrt{k_r(S)}\nabla\rho^\gamma\|_{L^{a_2}(Q_T)} \leq C, \quad a_2 \equiv \min\left\{\frac{2\gamma}{2+\gamma}, \frac{2\beta+4/3}{\beta+5/3}\right\} > 1.$$

In particular we deduce

$$(223) \quad \|\sqrt{k_r(S)}\nabla G(\rho^\gamma)\|_{L^{a_2}(Q_T)} \leq C[G],$$

3 where  $G \in W^{1,\infty}(\mathbb{R}_+)$  is arbitrary, and the constant  $C[G] > 0$  depends on  $G$ .

On the other hand, since  $1/\gamma + 1/q \leq 1$  (see (31)), from (32), (61), (68) it follows

$$\|G(\rho^\gamma)\nabla\sqrt{k_r(S)}\|_{L^\infty(0,T;L^q(\Omega))} \leq \|G(\rho^\gamma)\|_{L^\infty(Q_T)}\|\nabla f(S)\|_{L^\infty(0,T;L^q(\Omega))} \leq C[G],$$

4 which, together with (223), leads to (86).

Let us now find another gradient estimate related to  $\rho_i/\rho$ ,  $i = 1, \dots, N$ . From (110) and (64) we get

$$\int_0^T \int_\Omega \left( \log \frac{\rho_i}{\rho} - \frac{1}{N} \sum_{j=1}^N \log \frac{\rho_j}{\rho} \right)^2 dxdt = \int_0^T \int_\Omega \left( \log \rho_i - \frac{1}{N} \sum_{j=1}^N \log \rho_j \right)^2 dxdt \leq C,$$

for  $i = 1, \dots, N$ . Lemma 11 allows us to deduce

$$\left\| \sum_{j=1}^N \log \frac{\rho_j}{\rho} \right\|_{L^2(Q_T)} \leq C,$$

5 which, together with the previous estimate, leads to (87). Bound (88) on the reaction terms

6  $r_1, \dots, r_N$  comes straightforwardly from (42), (64).

**Proof of Lemma 10.** We show (159), i.e. that  $S^k \geq \varepsilon > 0$  a.e. in  $\Omega$ , for  $k \geq 0$ , provided that  $0 < \varepsilon < f^{-1}(p_{at}/\lambda_0)$ . Without loss of generality we can consider (142) on the set  $\{S^k \leq s_0\}$  where  $s_0$  is defined in (33). From the definition of  $P_{c,\varepsilon}$ , the positivity of  $f$  in  $(0, s_0)$  and (33), it follows

$$\inf_{s \in (0, s_0)} \frac{P_{c,\varepsilon}(s)}{f(s)} \geq \inf_{s \in (0, s_0)} \frac{P_c(s)}{f(s)} \equiv \lambda_0 > 0.$$

The positivity of  $f$  in  $(0, s_0)$ , the fact that  $p^k \geq -p_{at}$  and (142) imply

$$\frac{f(S^k) - f(S^{k-1})}{\tau} + \lambda_0 f(S^k) - p_{at} \leq 0,$$

and so

$$f(S^k) \leq (1 + \tau\lambda_0)^{-1} f(S^{k-1}) + p_{at}\tau(1 + \tau\lambda_0)^{-1}.$$

We prove by induction on  $k \geq 0$  that  $f(S^k) \leq f(\varepsilon)$  for every  $k \geq 0$ . Since  $S^0 = S^{in,\varepsilon} \geq \varepsilon$ , the statement is true for  $k = 0$ . Assume now the statement holds true for  $k - 1$ :  $f(S^{k-1}) \leq f(\varepsilon)$ . It follows

$$f(S^k) \leq (1 + \tau\lambda_0)^{-1} f(\varepsilon) + p_{at}\tau(1 + \tau\lambda_0)^{-1}.$$

The right-hand side of the above inequality is  $\leq f(\varepsilon)$  if  $f(\varepsilon) \geq p_{at}/\lambda_0$ , which holds true since  $0 < \varepsilon < f^{-1}(p_{at}/\lambda_0)$  and  $f$  is decreasing. Therefore the statement holds true for every  $k \geq 0$ . Once again, being  $f$  decreasing, it follows that

$$(224) \quad S^k \geq f^{-1}(f(\varepsilon)) = \varepsilon > 0, \quad k \geq 0, \quad \text{a.e. in } \Omega,$$

1 provided that  $\varepsilon < f^{-1}(p_{at}/\lambda_0)$ .

Bound (159) allows us to immediately find a bound for the spatial integral of  $f(S^k)$ . Indeed, integrating (142) in  $\Omega$  yields

$$\int_{\Omega} f(S^k) dx = \int_{\Omega} f(S^{k-1}) dx - \tau \int_{\Omega} (P_{c,\varepsilon}(S^k) + p^k) dx.$$

2 However, since (159) holds, then  $P_{c,\varepsilon}(S^k) \leq C(\varepsilon)$  for every  $k \geq 0$ , so (160) holds.

Finally, we will show that (161) holds. Let us now take the gradient of (142), test the resulting equation times  $\tau \frac{4}{3} (1 + |\nabla f(S^k)|^2)^{-1/3} \nabla f(S^k)$  and use the elementary inequality (a consequence of the convexity of  $x \mapsto (1 + x^2)^{2/3}$ )

$$\frac{4}{3} \frac{x}{(1 + x^2)^{1/3}} (x - y) \geq (1 + x^2)^{2/3} - (1 + y^2)^{2/3}, \quad x, y \geq 0.$$

It holds (thanks to Young's inequality)

$$\begin{aligned} & \int_{\Omega} (1 + |\nabla f(S^k)|^2)^{2/3} dx + \frac{4}{3} \tau \int_{\Omega} \frac{f'(S^k) P'_{c,\varepsilon}(S^k)}{(1 + |\nabla f(S^k)|^2)^{1/3}} |\nabla S^k|^2 dx \\ & \leq \int_{\Omega} (1 + |\nabla f(S^{k-1})|^2)^{2/3} dx - \frac{4}{3} \tau \int_{\Omega} \frac{\nabla p^k \cdot \nabla f(S^k)}{(1 + |\nabla f(S^k)|^2)^{1/3}} dx \\ & \leq \int_{\Omega} (1 + |\nabla f(S^{k-1})|^2)^{2/3} dx + \tau \int_{\Omega} |\nabla p^k|^{4/3} dx + C\tau \int_{\Omega} \frac{|\nabla f(S^k)|^4}{(1 + |\nabla f(S^k)|^2)^{4/3}} dx. \end{aligned}$$

3 Since  $f' P'_{c,\varepsilon} \geq 0$  in  $(0, 1)$  and  $x^4(1 + x^2)^{-4/3} \leq (1 + x^2)^{2/3}$  for  $x \geq 0$ , we conclude that (161) holds.

1 **5.1. Auxiliary results.** We begin by stating a simple but useful algebraic property.

**Lemma 11.** *For every  $\varepsilon > 0$  there exists  $C_\varepsilon > 0$  such that*

$$\sum_{i=1}^N \left( \log u_i - \frac{1}{N} \sum_{j=1}^N \log u_j \right)^2 \geq C_\varepsilon \left( \sum_{i=1}^N \log u_i \right)^2 - \varepsilon$$

2 *for every  $\vec{u} \in (0, \infty)^N$ , such that  $\sum_{i=1}^N u_i = 1$ .*

*Proof.* By contradiction. Assume there exists  $\varepsilon_0 > 0$  such that, for every  $n \in \mathbb{N}$ , there is  $u^{(n)} \in (0, \infty)^N$  such that

$$\sum_{i=1}^N \left( \log u_i^{(n)} - \frac{1}{N} \sum_{j=1}^N \log u_j^{(n)} \right)^2 < \frac{1}{n} \left( \sum_{i=1}^N \log u_i^{(n)} \right)^2 - \varepsilon_0, \quad \sum_{i=1}^N u_i^{(n)} = 1.$$

As a consequence  $\left( \sum_{i=1}^N \log u_i^{(n)} \right)^2 > 0$ , so we can define

$$v_i^{(n)} = \frac{\log u_i^{(n)}}{\sum_{k=1}^N \log u_k^{(n)}}, \quad i = 1, \dots, N,$$

and it follows

$$(225) \quad \sum_{i=1}^N \left( v_i^{(n)} - \frac{1}{N} \right)^2 < \frac{1}{n} - \frac{\varepsilon_0}{\left( \sum_{i=1}^N \log u_i^{(n)} \right)^2}.$$

We point out that clearly  $u_i^{(n)} \leq 1$  for  $i = 1, \dots, N$ , which implies that  $v_i^{(n)} \geq 0$  for  $i = 1, \dots, N$ . Furthermore  $\sum_{i=1}^N v_i^{(n)} = 1$  by construction, so the sequence  $v^{(n)}$  is bounded. Therefore there exists a subsequence (not relabeled) of  $v^{(n)}$  that is convergent:  $v^{(n)} \rightarrow v$  as  $n \rightarrow \infty$ . Taking the limit  $n \rightarrow \infty$  in (225) yields

$$\lim_{n \rightarrow \infty} v_i^{(n)} = \frac{1}{N} \quad i = 1, \dots, N, \quad \lim_{n \rightarrow \infty} \left| \sum_{k=1}^N \log u_k^{(n)} \right| = \infty.$$

As a consequence,

$$-\log u_i^{(n)} = v_i^{(n)} \left| \sum_{k=1}^N \log u_k^{(n)} \right| \rightarrow \infty \quad \text{as } n \rightarrow \infty, \quad i = 1, \dots, N.$$

3 This means that  $u_i^{(n)} \rightarrow 0$  as  $n \rightarrow \infty$  for  $i = 1, \dots, N$ , which is in contradiction with the fact that  
 4  $\sum_{i=1}^N u_i^{(n)} = 1$ . This finishes the proof of the Lemma.  $\square$

5 The following result is a generalized version of Fatou's Lemma.

**Lemma 12** (Generalized Fatou's Lemma). *Let  $Q \subset \mathbb{R}^d$  be a bounded domain,  $(u_n)_{n \in \mathbb{N}}, (v_n)_{n \in \mathbb{N}}$  sequences of Lebesgue-measurable functions  $Q \rightarrow \mathbb{R}^m, Q \rightarrow \mathbb{R}^N$  (respectively) such that  $u_n \rightarrow u$  a.e. in  $Q$  and  $v_n \rightarrow v$  weakly in  $L^p(Q)$ , for some  $p \geq 1$ . Let  $g : \mathbb{R}^m \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a continuous nonnegative function such that*

*for every  $u \in \mathbb{R}^m$ , the mapping  $v \in \mathbb{R}^N \mapsto g(u, v) \in \mathbb{R}$  is convex,*

$$\exists \omega \in C^0(\mathbb{R}_+), \omega(0) = 0 : \quad |g(u, v) - g(u', v)| \leq \omega(|u - u'|) |v|^p \quad \forall u, u' \in \mathbb{R}^m, v \in \mathbb{R}^N,$$

$$\exists C > 0 : \quad \int_Q g(u_n, v_n) \leq C \quad \forall n \in \mathbb{N}.$$



Then

$$(226) \quad \liminf_{n \rightarrow \infty} \int_Q g(u_n, v_n) \geq \int_Q g(u, v).$$

*Proof.* Since  $u_n \rightarrow u$  a.e. in  $Q$ , Egorov-Severini's Theorem implies that, for every  $\varepsilon > 0$ , there exists  $E_\varepsilon \subset Q$  such that  $|Q \setminus E_\varepsilon| < \varepsilon$  and  $u_n \rightarrow u$  in  $L^\infty(E_\varepsilon)$ . Being  $g$  nonnegative it holds

$$\int_Q g(u_n, v_n) \geq \int_{E_\varepsilon} g(u_n, v_n) = \int_{E_\varepsilon} (g(u_n, v_n) - g(u, v_n)) + \int_{E_\varepsilon} g(u, v_n).$$

Thanks to the assumptions on  $g$ ,  $u_n$ ,  $v_n$

$$\left| \int_{E_\varepsilon} (g(u_n, v_n) - g(u, v_n)) \right| \leq \int_{E_\varepsilon} \omega(|u_n - u|)|v_n|^p \leq \|\omega(|u_n - u|)\|_{L^\infty(E_\varepsilon)} \|v_n\|_{L^p(Q)}^p \rightarrow 0$$

as  $n \rightarrow \infty$ , therefore

$$\liminf_{n \rightarrow \infty} \int_Q g(u_n, v_n) \geq \liminf_{n \rightarrow \infty} \int_{E_\varepsilon} g(u, v_n).$$

Being  $v \mapsto g(u, v)$  convex, the mapping  $v \in L^p(E_\varepsilon) \mapsto \int_{E_\varepsilon} g(u, v) \in \mathbb{R}$  is weakly lower semicontinuous and so

$$\liminf_{n \rightarrow \infty} \int_Q g(u_n, v_n) \geq \int_{E_\varepsilon} g(u, v).$$

- 1 Since  $\varepsilon > 0$  is arbitrary and  $|Q \setminus E_\varepsilon| < \varepsilon$  we conclude that (226) holds. This finishes the proof of  
2 the Lemma. □

3

### REFERENCES

- 4 [1] B. AMAZIANE, M. JURAK, L. PANKRATOV AND A. PIATNITSKI. An existence result for nonisothermal immis-  
5 cible incompressible 2-phase flow in heterogeneous porous media. *Math Meth Appl Sci.* 40 (2017), 7510–7539.  
6 [2] J. BEAR. Dynamics of Fluids in Porous Media, *American Elsevier*, 1972. Reissued by *Dover publication*, 1988.  
7 [3] M. BENEŠ, I. PAŽANIN. Homogenization of degenerate coupled fluid flows and heat transport through porous  
8 media. *J. Math. Anal. Appl.* 446 (2017), 165–192.  
9 [4] O.B. BOCHAROV, V.N. MONAKHOV. Boundary value problems of nonisothermic two-phase filtration in porous  
10 media. *Dinamika Sploshn Sredy* 86 (1988), 47–59.  
11 [5] O.B. BOCHAROV, V.N. MONAKHOV. Nonisothermal filtration of immiscible fluids with variable residual satu-  
12 ration. *Dinamika Sploshn Sredy* 88 (1988), 3–12.  
13 [6] D. BOTHE, W. DREYER. Continuum thermodynamics of chemically reacting fluid mixtures. *Acta Mech.* 226  
14 (2015), 1757–1805.  
15 [7] M. BULIČEK, J. HAVRDA. On existence of weak solution to a model describing incompressible mixtures with  
16 thermal diffusion cross effects. *Z. Angew. Math. Mech.* 95 (2015), 589–619.  
17 [8] M. BULIČEK, A. JÜNGEL, M. POKORNÝ, N. ZAMPONI. Existence analysis of a stationary compressible fluid  
18 model for heat-conducting and chemically reacting mixtures. Submitted for publication (arXiv: 2001.06082).  
19 [9] M. BULIČEK, M. POKORNÝ, N. ZAMPONI. Existence analysis for incompressible fluid model of electrically  
20 charged chemically reacting and heat conducting mixtures. *SIAM J. Math. Anal.* 49 (2017), no. 5, 3776–3830.  
21 [10] M. BURGER, B. SCHLAKE, M.-T. WOLFRAM. Nonlinear Poisson-Nernst-Planck equations for ion flux through  
22 confined geometries. *Nonlinearity* 25 (2012), no. 4, 961–990.  
23 [11] S. CONTI, G. DOLZMANN, S. MÜLLER. The div-curl lemma for sequences whose divergence and curl are  
24 compact in  $W^{-1,1}$ . *Comptes Rendus Mathematique* 349.3-4 (2011), 175–178.  
25 [12] O. COUSSY. Poromechanics, *Wiley* 2004.  
26 [13] E. S. DAUS, J.-P. MILIŠIĆ, N. ZAMPONI. Global existence for a two-phase flow model with cross diffusion.  
27 *DCDS-B* 25.3 (2020), 957–979.  
28 [14] M. DREHER, A. JÜNGEL. Compact families of piecewise constant functions in  $L^p(0, T; B)$ . *Nonlinear Analysis:*  
29 *Theory, Methods & Applications* 75.6 (2012): 3072–3077.  
30 [15] W. DREYER, P.-E. DRUET, P. GAJEWSKI, C. GUHLKE. Analysis of improved Nernst-Planck-Poisson models  
31 of compressible isothermal electrolytes. *Z. Angew. Math. Phys.* 71, 119 (2020).

- 1 [16] E. FEIREISL. Dynamics of Viscous Compressible Fluids. *Oxford University Press*, Oxford, 2004.
- 2 [17] E. FEIEREISL, A. NOVOTNÝ. Singular limits in Thermodynamics of Viscous Fluids. *Birkhäuser*, 2009.
- 3 [18] J. FISCHER, K. HOPF, M. KNIELY, A. MIELKE. Global existence analysis of energy-reaction-diffusion systems.  
4 Submitted for publication (arXiv:2012.03792).
- 5 [19] V. GIOVANGIGLI. Multicomponent Flow Modeling. *Birkhäuser*, Boston, 1999.
- 6 [20] S.R. DE GROOT, P. MAZUR. Non-equilibrium thermodynamics. *Dover publications*, 1984.
- 7 [21] S.M. HASSANIZADEH, W.G. GRAY. Thermodynamic basis of capillary-pressure in porous-media, *Water Re-*  
8 *sour. Res.* 29.10 (1993), 3389–3405.
- 9 [22] C. HELMER, A. JÜNGEL. Analysis of Maxwell-Stefan systems for heat conducting fluid mixtures. *Nonlinear*  
10 *Anal. Real World Appl.* 59 (2021), 103263.
- 11 [23] M. JURAK, A. KOLDOBA, A. KONYUKHOV AND L. PANKRATOV. Nonisothermal immiscible compressible  
12 thermodynamically consistent two-phase flow in porous media. *C. R. Mecanique* 347 (2019), 920–929.
- 13 [24] A. JÜNGEL. Entropy Methods for Diffusive Partial Differential Equations. *Springer Briefs in Mathematics*,  
14 Springer, 2016.
- 15 [25] A. JÜNGEL. The boundedness-by-entropy method for cross-diffusion systems. *Nonlinearity* 28 (2015), no. 6,  
16 1963–2001.
- 17 [26] A. JÜNGEL, J. MIKYŠKA, N. ZAMPONI. Existence analysis of a single-phase flow mixture model with van der  
18 Waals pressure. To appear in *SIAM J. Math. Anal.* (2018), no.1 (arXiv: 1612.04161).
- 19 [27] M. KAVIANY. Principles of Heat Transfer in Porous Media. *Springer*, New York, 1995.
- 20 [28] B. LI, W. SUN. Global existence of weak solution for nonisothermal multicomponent flow in porous textile  
21 media. *SIAM J. Math. Anal.* 42 (2010), 3076–3102.
- 22 [29] A. MOUSSA. Some variants of the classical Aubin-Lions Lemma. *J. Evol. Equ.* 16 (2016), 65–93.
- 23 [30] F. MURAT. Compacité par compensation. *Ann. Sc. Norm. Sup. Pisa*, Cl. Sci. Ser. 5, IV (1978), 489–507.
- 24 [31] A. NOVOTNÝ, M. POKORNÝ. Steady compressible Navier-Stokes-Fourier system for monoatomic gas and its  
25 generalizations. *J. Diff. Eqs.* 251 (2011), 270–315.
- 26 [32] L. TARTAR. Compensated compactness and applications to partial dif- ferential equations. *Nonlinear Anal.*  
27 *and Mech.*, Heriot-Watt Sympos., L.J. Knopps editor, Research Notes in Math 39, Pitman, Boston, 136–211, 1975.
- 28 [33] E. ZEIDLER. *Nonlinear Functional Analysis and its Applications*. Volume II/B. Springer, New York, 1990.
- 29 INSTITUTE FOR ANALYSIS AND SCIENTIFIC COMPUTING, VIENNA UNIVERSITY OF TECHNOLOGY, WIEDNER  
30 HAUPTSTRASSE 8–10, 1040 WIEN, AUSTRIA  
31 *Email address:* esther.daus@tuwien.ac.at
- 32 UNIVERSITY OF ZAGREB, FACULTY OF ELECTRICAL ENGINEERING AND COMPUTING, UNSKA 3, 10000 ZA-  
33 GREB, CROATIA  
34 *Email address:* pina.milisic@fer.hr
- 35 UNIVERSITY OF MANNHEIM, SCHOOL OF BUSINESS INFORMATICS AND MATHEMATICS, B6, 28, 68159 MANNHEIM,  
36 GERMANY  
37 *Email address:* nzamponi@mail.uni-mannheim.de