

A connecting theorem for geodesic flows on the 2-torus

Stefan Klempnauer*

Faculty of Mathematics, Ruhr-University Bochum

December 12, 2021

Abstract

We use a result of J. Mather on the existence of connecting orbits for compositions of monotone twist maps of the cylinder to prove the existence of connecting geodesics on the unit tangent bundle ST^2 of the 2-torus in regions without invariant tori.

The author thanks the SFB CRC/TRR 191 *Symplectic Structures in Geometry, Algebra and Dynamics* of the DFG and the Ruhr-University Bochum for the funding of his research.

1 Introduction and main result

Let $T^2 \cong \mathbb{R}^2/\mathbb{Z}^2$ denote the 2-torus with universal covering $\pi : \mathbb{R}^2 \rightarrow T^2$. The tangent bundle is given by $TT^2 \cong T^2 \times \mathbb{R}^2$. In the following we will recall the definition of a Finsler metric. For a general overview of Finsler geometry see [1].

Definition 1.1. A Finsler metric on T^2 is a map

$$F : TT^2 \rightarrow [0, \infty)$$

with the following properties

1. (Regularity) F is C^∞ on $TT^2 - 0$
2. (Positive homogeneity) $F(x, \lambda y) = \lambda \cdot F(x, y)$ for $\lambda > 0$
3. (Strong convexity) The hessian

$$(g_{ij}(x, v)) = \left(\partial_{v_i v_j} \frac{1}{2} F(x, v)^2 \right)$$

is positive-definite for every $(x, v) \in TT^2 - 0$.

*stefan.klempnauer@rub.de

A Finsler metric is called *reversible* if $F(x, v) = F(x, -v)$ for every $(x, v) \in TT^2$. The *unit tangent bundle* $ST^2 \cong T^2 \times S^1$ is given by $ST^2 = F^{-1}(\{1\})$.

We define the *length* $l_F(c)$ of a piecewise differentiable curve $c : [a, b] \rightarrow T^2$ via

$$l_F(c) = \int_a^b F(\dot{c}(t)) dt$$

Lifting the Finsler metric F to \mathbb{R}^2 allows us to define the length of a curve $c : [a, b] \rightarrow \mathbb{R}^2$ in the universal covering in the same manner. The *geodesic flow* $\phi^t : ST^2 \rightarrow ST^2$ is the restriction of the Euler-Lagrange flow of the Lagrangian L_F , with

$$L_F = \frac{1}{2} F^2$$

to the unit tangent bundle. A *geodesic* we call either a trajectory of the Euler-Lagrange flow, or its projection to T^2 , i.e. a geodesic is a curve $t \mapsto c(t) \subset T^2$ satisfying the Euler-Lagrange equation

$$\partial_x L_F(c, \dot{c}) - \partial_t (\partial_v L_F(c, \dot{c})) = 0$$

In general we will assume a geodesic c to be parametrized by arclength, i.e. $\dot{c} \subset ST^2$. Sometimes it can be convenient to consider lifts $\tilde{c} : \mathbb{R} \rightarrow \mathbb{R}^2$ or $\check{c} : \mathbb{R} \rightarrow T\mathbb{R}^2$ of a geodesic to the universal cover. These can be seen as the geodesics of the lifted \mathbb{Z}^2 -periodic Finsler metric \tilde{F} on \mathbb{R}^2 .

Definition 1.2. A subset $\Lambda \subset ST^2$ is called an *invariant torus* if Λ is the graph of a continuous map $X : T^2 \rightarrow S^1$ and ϕ^t -invariant. An invariant torus (or more generally any ϕ^t -invariant set) Λ has *bounded direction* (with respect to $v \in \mathbb{Z}^2 - \{0\}$), if the lifts $\tilde{c} : \mathbb{R} \rightarrow \mathbb{R}^2$ of geodesics in Λ are graphs over the euclidean line $\mathbb{R}v$.

In analogy to a Birkhoff region of instability we define the following notion of an instability region of the geodesic flow.

Definition 1.3. An *instability region* is a compact invariant subset $U \subset ST^2$ with boundary being the disjoint union of two invariant tori Λ_-, Λ_+ , such that every invariant torus $\Lambda \subset U$ is equal to Λ_- or Λ_+ .

The main result is the following.

Theorem 1.4. Let F be a Finsler metric on T^2 and let $U \subset ST^2$ be an instability region with bounded direction (with respect to e_1). Furthermore, we assume that there are no closed geodesics in the boundary of U . Then there exists an F -geodesic $c : \mathbb{R} \rightarrow T^2$ that connects the two boundary components of U (i.e. there exist sequences $t_n \rightarrow \infty$ and $s_n \rightarrow -\infty$ with $\lim_{n \rightarrow \infty} \dot{c}(t_n) \in \Lambda_+$ and $\lim_{n \rightarrow \infty} \dot{c}(s_n) \in \Lambda_-$, where Λ_- and Λ_+ are the two boundary components of U).

2 Return maps of the geodesic flow on T^2 and twist maps

2.1 Return maps as Lagrangian time-1 maps

Let F be a Finsler metric on the 2-torus T^2 with \tilde{F} denoting the lifted metric on the universal cover \mathbb{R}^2 . For a vector $w = (w_1, w_2) \in \mathbb{R}^2$ we will write w^\perp for the vector $w^\perp = (-w_2, w_1)$.

Definition 2.1. For a Finsler metric F and a prime element $v \in \mathbb{Z}^2 \setminus \{0\}$ we define a time-dependent Lagrangian on \mathbb{R} via

$$\tilde{L}(t, x, r) = \tilde{F}(tv + xv^\perp, v + rv^\perp)$$

which is 1-periodic in the time t and 1-periodic in the first component x . The periodicity is a direct consequence of the \mathbb{Z}^2 -periodicity of \tilde{F} . Since \tilde{L} is periodic in x it can be interpreted as a time-periodic Lagrangian L on S^1 . Note that \tilde{L} is strictly convex, i.e. $\partial_{rr}\tilde{L} > 0$ because of the strict convexity of the Finsler metric \tilde{F} .

We have a correspondence of lifted geodesics, which are graphs over $v\mathbb{R}$ and solutions of L in the following theorem by J.P. Schröder. A special case can be found in [5].

Theorem 2.2. Let F and L as above and let $\theta : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function. Let $\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$ be the curve given by

$$\gamma(t) = tv + \theta(t)v^\perp$$

Then γ is a reparametrization of an \tilde{F} -geodesic if and only if θ is an Euler-Lagrange solution of \tilde{L} .

Proof. Observe that we have the following relation between the Lagrangian action $A_{\tilde{L}}$ and the Finsler length $l_{\tilde{F}}$.

$$\begin{aligned} A_{\tilde{L}}(\theta|_{[a,b]}) &= \int_a^b \tilde{L}(t, \theta(t), \theta'(t)) dt \\ &= \int_a^b \tilde{F}(tv + \theta(t)v^\perp, v + \theta'(t)v^\perp) dt \\ &= \int_a^b \tilde{F}(\gamma(t), \dot{\gamma}(t)) dt \\ &= l_{\tilde{F}}(\gamma|_{[a,b]}) \end{aligned}$$

Assume now that $\gamma : [a, b] \rightarrow \mathbb{R}^2$ is the reparametrization of an \tilde{F} -geodesic, i.e. $\partial_{s=0} l_{\tilde{F}}(\gamma_s) = 0$ for any proper variation of γ . Let $\theta_s : [a, b] \rightarrow \mathbb{R}$ be a proper variation of θ . From θ_s we construct a proper variation γ_s of γ via

$$\gamma_s(t) = tv + \theta_s(t)v^\perp$$

Then we have $A_{\tilde{L}}(\theta_s) = l_{\tilde{F}}(\gamma_s)$ for every s , and hence we have

$$\partial_s|_{s=0} A_{\tilde{L}}(\theta_s) = \partial_s|_{s=0} l_{\tilde{F}}(\gamma_s) = 0.$$

This proves one direction. To prove the other direction assume now that $\theta : [a, b] \rightarrow \mathbb{R}$ is critical with respect to the Lagrangian action and let $X : [a, b] \rightarrow \mathbb{R}^2$ be a vector field along γ with $X(a) = X(b) = 0$. Since $\dot{\gamma}(t) = v + \theta'(t)v^\perp$ the pair of vectors $\{\dot{\gamma}(t), v^\perp\}$ always forms a basis of \mathbb{R}^2 . Thus, we can rewrite the vector field X as

$$X(t) = \underbrace{\lambda(t)\dot{\gamma}(t)}_{A(t)} + \underbrace{\mu(t)v^\perp}_{B(t)}$$

for functions λ, μ with $\lambda(a) = \lambda(b) = \mu(a) = \mu(b) = 0$. Let γ_s be a proper variation of γ corresponding to the variational vector field X and let β_s be a proper variation of γ corresponding to B , i.e.

$$\partial_s|_{s=0} \gamma_s(t) = X(t) \quad \text{and} \quad \partial_s|_{s=0} \beta_s(t) = B(t)$$

Observe that for small $|s|$ the curve $\eta_s : t \mapsto \gamma(t + s\lambda(t))$ is a reparametrization of γ and hence has length independent of s . Thus we have

$$\begin{aligned}
0 &= \partial_s|_{s=0} l_{\tilde{F}}(\eta_s) \\
&= \partial_s|_{s=0} \int_a^b \tilde{F}(\eta_s(t), \partial_t \eta_s(t)) dt \\
&= \int_a^b \partial_s|_{s=0} \tilde{F}(\eta_s(t), \partial_t \eta_s(t)) dt \\
&= \int_a^b \partial_1 \tilde{F}(\eta_0(t), \partial_t \eta_0(t)) \partial_s|_{s=0} \eta_s(t) + \partial_2 \tilde{F}(\eta_0(t), \partial_t \eta_0(t)) \partial_s|_{s=0} \partial_t \eta_s(t) dt \\
&= \int_a^b \partial_1 \tilde{F}(\gamma(t), \partial_t \gamma(t)) \partial_s|_{s=0} \eta_s(t) + \partial_2 \tilde{F}(\gamma(t), \partial_t \gamma(t)) \partial_t \partial_s|_{s=0} \eta_s(t) dt \\
&= \int_a^b \partial_1 \tilde{F}(\gamma(t), \partial_t \gamma(t)) A(t) + \partial_2 \tilde{F}(\gamma(t), \partial_t \gamma(t)) \dot{A}(t) dt
\end{aligned}$$

Hence it follows that

$$\begin{aligned}
\partial_s|_{s=0} l_{\tilde{F}}(\gamma_s) &= \partial_s|_{s=0} \int_a^b \tilde{F}(\gamma_s(t), \dot{\gamma}_s(t)) dt \\
&= \int_a^b \partial_1 \tilde{F}(\gamma_0(t), \dot{\gamma}_0(t)) \partial_s|_{s=0} \gamma_s(t) + \partial_2 \tilde{F}(\gamma_0(t), \dot{\gamma}_0(t)) \partial_s|_{s=0} \dot{\gamma}_s(t) dt \\
&= \int_a^b \partial_1 \tilde{F}(\gamma(t), \dot{\gamma}(t)) \partial_s|_{s=0} \gamma_s(t) + \partial_2 \tilde{F}(\gamma(t), \dot{\gamma}(t)) \partial_t \partial_s|_{s=0} \gamma_s(t) dt \\
&= \int_a^b \partial_1 \tilde{F}(\gamma(t), \dot{\gamma}(t)) X(t) + \partial_2 \tilde{F}(\gamma(t), \dot{\gamma}(t)) \dot{X}(t) dt \\
&= \int_a^b \partial_1 \tilde{F}(\gamma(t), \dot{\gamma}(t)) (A(t) + B(t)) + \partial_2 \tilde{F}(\gamma(t), \dot{\gamma}(t)) (\dot{A}(t) + \dot{B}(t)) dt \\
&= \int_a^b \partial_1 \tilde{F}(\gamma(t), \dot{\gamma}(t)) B(t) + \partial_2 \tilde{F}(\gamma(t), \dot{\gamma}(t)) \dot{B}(t) dt \\
&= \partial_s|_{s=0} l_{\tilde{F}}(\beta_s)
\end{aligned}$$

Consequently, the curve γ is critical with respect to the Finsler length if

$$\partial_s|_{s=0} l_{\tilde{F}}(\beta_s) = 0$$

for every proper variation β_s , which varies γ only in v^\perp direction, i.e. β_s is of the form

$$\beta_s(t) = tv + \theta_s(t)v^\perp$$

For those variations we have already computed that if θ_s is critical with respect to the \tilde{L} -action then β_s is critical with respect to the Finsler length. \square

Remember that $T\mathbb{R}^2 \cong \mathbb{R}^2 \times \mathbb{R}^2$. To establish a connection between return maps of the geodesic flow and the time-1 map of the Lagrangian \tilde{L} we define the following sets

$$\tilde{V} = \{(x, w) \in T\mathbb{R}^2 : \langle w, v \rangle > 0, \tilde{F}(w) = 1\}$$

with sections \tilde{V}_i , where $i \in \mathbb{Z}$, given by

$$\tilde{V}_i = \{(iv + bv^\perp, w) \in T\mathbb{R}^2 : \langle w, v \rangle > 0, \tilde{F}(w) = 1, b \in \mathbb{R}\}$$

In general the Euler-Lagrange solutions of \tilde{L} will not exist for all times. However, because of theorem 2.2 we have the following

Proposition 2.3. *If the Euler-Lagrange flow of \tilde{L} is complete then the set \tilde{V} is invariant and every orbit starting in a section \tilde{V}_i will consecutively pass through the sections \tilde{V}_{i+j} for $j \in \mathbb{N}$.*

Proof. Completeness of the Euler-Lagrange flow means that the Euler-Lagrange solutions of \tilde{L} exist for all times. Let $(x, w) \in \tilde{V}$. We write $(x, w) = (av + bv^\perp, rv + hv^\perp)$ for coefficients $a, b, h, r \in \mathbb{R}$ with $r > 0$. Let $\tilde{c} : \mathbb{R} \rightarrow \mathbb{R}^2$ be the geodesic with $\tilde{c}(a) = x$ and $\dot{\tilde{c}}(a) = \frac{1}{r}w$. To show that \tilde{V} is invariant we have to show that $\langle \dot{\tilde{c}}(t), v \rangle > 0$ for every $t \in \mathbb{R}$. Let $\theta : \mathbb{R} \rightarrow \mathbb{R}$ be the Euler-Lagrange solution of \tilde{L} with $\theta(a) = b$ and $\theta'(a) = \frac{h}{r}$. Since θ exists for all times it follows from Theorem 2.2 that the curve $\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$ with $\gamma(t) = tv + \theta(t)v^\perp$ is a reparametrization of a geodesic. We have $\gamma(a) = av + bv^\perp = x$ and $\dot{\gamma}(a) = v + \frac{h}{r}v^\perp = \frac{1}{r}w$ and hence γ is a reparametrization of the geodesic \tilde{c} . Since $\langle \dot{\gamma}(t), v \rangle = \|v\|^2$ for every t , we have $\langle \dot{\tilde{c}}(t), v \rangle > 0$ for every t . With the same construction one sees that if (x, w) was chosen in \tilde{V}_i the constructed reparametrization γ will pass through \tilde{V}_{i+j} at times $i + j$. \square

By $\tilde{R}_0 : \tilde{V}_0 \rightarrow \tilde{V}_1$ we denote the map that maps an element $(x, w) \in \tilde{V}_0$ to the point of intersection of \tilde{V}_1 with the orbit passing through (x, w) . We define the projected section $V \subset ST^2$ via

$$V := \Pi(V_0)$$

Since the geodesic flow on $T\mathbb{R}^2$ is \mathbb{Z}^2 -invariant, the map \tilde{R}_0 descends to a map $R : V \rightarrow V$, which is the n -th return map of the section V , with respect to the geodesic flow on ST^2 . Here, $n - 1$ is equal to $\|v\|^2 - 1$ ($n - 1$ is equal to the number of deck-transformations of $\mathbb{R}v^\perp$, that lie in between $\mathbb{R}v^\perp$ and $v + \mathbb{R}v^\perp$, which corresponds to the number of integer points in the rectangle spanned by v and v^\perp . If v is a prime element in \mathbb{Z}^2 then this is equal to $\|v\|^2 - 1$ after Pick's theorem).

Proposition 2.4. *If the Euler-Lagrange flow of \tilde{L} is complete the return map R is conjugated to the time-1 map $\varphi_L^{0,1} : Z \rightarrow Z$ of the Lagrangian L .*

Proof. Let $(x, w) \in \tilde{V}_0$. We can rewrite (x, w) as

$$(x, w) = \left(\frac{\langle x, v^\perp \rangle}{\|v\|^2} v^\perp, \frac{\langle w, v \rangle}{\|v\|^2} v + \frac{\langle w, v^\perp \rangle}{\|v\|^2} v^\perp \right)$$

Define real numbers $b, h \in \mathbb{R}$ and $r > 0$, such that $(x, w) = (bv^\perp, rv + hv^\perp)$. Since the Euler-Lagrange flow is complete there exists an Euler-Lagrange solution $\theta : \mathbb{R} \rightarrow \mathbb{R}$ of \tilde{L} with

$$\theta(0) = b \quad \text{and} \quad \theta'(0) = \frac{h}{r}$$

From theorem 2.2 it follows that there exists a reparametrization

$$\gamma(t) = tv + \theta(t)v^\perp$$

of an \tilde{F} -geodesic. It follows then that

$$\tilde{R}_0(x, w) = \left(\gamma(1), \frac{\dot{\gamma}(1)}{\tilde{F}(\gamma(1), \dot{\gamma}(1))} \right)$$

We define the scaled sections $\widetilde{W}_i \subset T\mathbb{R}^2$ for $i \in \mathbb{Z}$ via

$$\widetilde{W}_0 = \{(bv^\perp, v + dv^\perp) \in T\mathbb{R}^2 : b, d \in \mathbb{R}\} \quad \text{and} \quad \widetilde{W}_i = iv + \widetilde{W}_0$$

together with a map $\widetilde{g} : T\mathbb{R}^2 - 0 \rightarrow T\mathbb{R}^2 - 0$ with

$$\widetilde{g}(x, w) = \left(x, \frac{w}{\widetilde{F}(x, w)} \right)$$

Observe that the restrictions $\widetilde{g}|_{\widetilde{W}_i} : \widetilde{W}_i \rightarrow \widetilde{V}_i$ are diffeomorphisms (check that $(x, w) \mapsto (x, \frac{\|v\|^2}{\langle v, w \rangle} w)$ is the inverse of $\widetilde{g}|_{\widetilde{W}_i}$ and both are differentiable) and that

$$\left(\widetilde{g}|_{\widetilde{W}_1} \right)^{-1} \circ \widetilde{R}_0 \circ \widetilde{g}|_{\widetilde{W}_0} = \widetilde{P}_0$$

for the map $\widetilde{P}_0 : \widetilde{W}_0 \rightarrow \widetilde{W}_1$ with

$$\widetilde{P}_0(\gamma(0), \dot{\gamma}(0)) = (\gamma(1), \dot{\gamma}(1))$$

Define diffeomorphisms $\widetilde{l}_i : \widetilde{W}_i \rightarrow \mathbb{R}^2$ via

$$\widetilde{l}_i(x, w) = \left(\frac{\langle x, v^\perp \rangle}{\|v\|^2}, \frac{\langle w, v^\perp \rangle}{\|v\|^2} \right)$$

Then we have

$$\widetilde{l}_1 \circ \widetilde{P}_0 \circ \left(\widetilde{l}_0 \right)^{-1} = \varphi_L^{0,1}$$

The conjugacy of the return map R and the time-1 map $\varphi_L^{0,1}$ follows from the \mathbb{Z}^2 -invariance of the geodesic flow of \widetilde{F} . \square

Example 2.5. Let \widetilde{F} be the flat metric, i.e.

$$\widetilde{F}(x, w) = \|w\| = \sqrt{\langle w, w \rangle}$$

and $v \in \mathbb{Z}^2 \setminus \{0\}$ prime. The Lagrangian \widetilde{L} (see definition 2.1) is then given by

$$\begin{aligned} \widetilde{L}(t, x, r) &= \widetilde{F}(tv + xv^\perp, v + rv^\perp) \\ &= \|v\| \sqrt{1 + r^2} \end{aligned}$$

We obtain that a function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ is an Euler-Lagrange solution if and only if

$$\underbrace{\partial_x \widetilde{L}(t, \theta(t), \theta'(t))}_{=0} - \partial_t (\partial_r \widetilde{L}(t, \theta(t), \theta'(t))) = 0$$

which is equivalent to

$$\|v\| \frac{\theta''}{\sqrt{1 + \theta'^2}^3} = 0$$

and thus θ is an Euler-Lagrange solution if and only if $\theta''(t) = 0$ for every $t \in \mathbb{R}$. Consequently the solutions are of the form

$$\theta(t) = at + b \quad \text{for } a, b \in \mathbb{R}$$

Thus, we have for the time-1 map $\varphi_L^{0,1} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ of the Lagrangian \widetilde{L}

$$\varphi_L^{0,1}(x, y) = (x + y, y)$$

To determine the map \tilde{R}_0 let $(x, w) \in \tilde{V}_0$, i.e. $(x, w) = (bv^\perp, rv + hv^\perp)$ for $b, h \in \mathbb{R}$ and $r > 0$, such that $\|rv + hv^\perp\| = 1$. The Euler-Lagrange solution θ with $\theta(0) = b$ and $\theta'(0) = \frac{h}{r}$ is given by

$$\theta(t) = \frac{h}{r}t + b$$

This yields the following reparametrization γ of an \tilde{F} -geodesic

$$\gamma(t) = tv + \left(\frac{h}{r}t + b\right)v^\perp$$

We calculate

$$\begin{aligned}\tilde{R}_0(x, w) &= \left(\gamma(1), \frac{\dot{\gamma}(1)}{\|\dot{\gamma}(1)\|}\right) \\ &= \left(v + \left(\frac{h}{r} + b\right)v^\perp, \frac{v + \frac{h}{r}v^\perp}{\|v + \frac{h}{r}v^\perp\|}\right) \\ &= \left(x + \frac{1}{r}w, \frac{w}{\|w\|}\right) \\ &= \left(x + \frac{\|v\|^2}{\langle w, v \rangle}w, w\right)\end{aligned}$$

To determine the map \tilde{P}_0 recall that

$$\tilde{P}_0 = \left(\tilde{g}|_{\tilde{W}_1}\right)^{-1} \circ \tilde{R}_0 \circ \tilde{g}|_{\tilde{W}_0}$$

with

$$\tilde{g}|_{\tilde{W}_i}(x, w) = \left(x, \frac{w}{\|w\|}\right) \quad \text{and} \quad \left(\tilde{g}|_{\tilde{W}_i}\right)^{-1}(x, w) = \left(x, \frac{\|v\|^2}{\langle w, v \rangle}w\right)$$

For $(x, w) \in \tilde{W}_0$ we obtain

$$\begin{aligned}\tilde{P}_0(x, w) &= \left(\tilde{g}|_{\tilde{W}_1}\right)^{-1} \circ \tilde{R}_0 \left(x, \frac{w}{\|w\|}\right) \\ &= \left(\tilde{g}|_{\tilde{W}_1}\right)^{-1} \left(x + \frac{\|v\|^2}{\langle w, v \rangle}w, \frac{w}{\|w\|}\right) \\ &= \left(x + \frac{\|v\|^2}{\langle w, v \rangle}w, \frac{\|v\|^2}{\langle w, v \rangle}w\right) \\ &= (x + w, w)\end{aligned}$$

with the last equation following from $(x, w) \in \tilde{W}_0$. To conclude the example we calculate further

$$\begin{aligned}\tilde{l}_1 \circ \tilde{P}_0 \circ (\tilde{l}_0)^{-1}(x, y) &= \tilde{l}_1 \circ \tilde{P}_0(xv^\perp, v + yv^\perp) \\ &= \tilde{l}_1(xv^\perp + v + yv^\perp, v + yv^\perp) \\ &= (x + y, y)\end{aligned}$$

with the last line being equal to the time-1 map of \tilde{L} .

In this example the time-1 map of \tilde{L} is a twist map (the so-called *shear map*). We will see later that in general the strict convexity of L guarantees that the time-1 map $\phi_L^{0,1}$ is conjugated to a finite composition of positive twist maps. This will allow us to apply the theory of monotone twist maps to return maps of the geodesic flow.

2.2 Twist maps of the cylinder and Mather's theorem

In this section we will briefly recall some of the theory on finite compositions of monotone twist maps of the cylinder and Mather's theorem about connecting orbits in Birkhoff regions of instability.

Definition 2.6. Let $f : Z \rightarrow Z$ be a map of the cylinder with a lift $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, such that $F(x, y) = (X(x, y), Y(x, y))$ is a diffeomorphism with

1. F is isotopic to the identity
2. Twist condition: The map $(x, y) \mapsto (x, X(x, y))$ is a diffeomorphism for every y
3. Exact symplectic: $F^*(\lambda) - \lambda = dS$ for some function $S : Z \rightarrow \mathbb{R}$.

Then $f : Z \rightarrow Z$ is called a monotone twist map of the cylinder. The map f is called positive if $\partial_y X > 0$ everywhere. Here, $\lambda = ydx$ is the Liouville form on \mathbb{R}^2 .

For further reading on twist maps we refer the reader to [3]. The following definitions go along the lines of [4]. The following theorem (originally due to Birkhoff [2]) can be found in the Appendix of [4] for maps $f \in \mathcal{P}^1$.

Theorem 2.7. Let $f \in \mathcal{P}^1$. Any f -invariant homotopically non-trivial Jordan curve (homeomorphic to S^1) Γ in the infinite cylinder is the graph of a Lipschitz function $u : S^1 \rightarrow \mathbb{R}$.

By \mathcal{P}^1 we denote the set of finite compositions of positive monotone twist maps of the cylinder.

Definition 2.8. Let $f \in \mathcal{P}^1$. A Birkhoff region of instability $B \subset Z$ is a compact f -invariant subset, such that

1. The boundary of B consists of two components Γ_- and Γ_+ , each homeomorphic to a circle and each non-contractible in Z
2. If $\Gamma \subset B$ is any f -invariant subset, homeomorphic to a circle, non-contractible, then $\Gamma = \Gamma_-$ or $\Gamma = \Gamma_+$

For every $f \in \mathcal{P}^1$ there is a variational principle h (cf [4] chapter 1) and consequently one can define the notion of a minimal orbit of f . Much like if f is a twist map one can prove that every minimal orbit has a rotation number. This let's us define the set $\mathcal{M}_{f,\omega}$ of minimal orbits of f with rotation number ω . Let $\Gamma \subset Z$ be an f -invariant circle (and consequently after theorem 2.7 a Lipschitz graph) with rotation number ω . It follows from [4] proposition 2.8 that $\Gamma \subset \mathcal{M}_{f,\omega}$. If ω is irrational it follows from [4] proposition 2.6 that the projection $\pi : Z \rightarrow S^1$ restricted to $\mathcal{M}_{f,\omega}$ is injective, and hence we have $\Gamma = \mathcal{M}_{f,\omega}$.

A special case of theorem 4.1 in [4] is the following connecting theorem by J. Mather

Theorem 2.9. Let $f \in \mathcal{P}^1$. If f has a Birkhoff region of instability B , such that the rotation numbers $\omega_- < \omega_+$ corresponding to the boundary graphs Γ_- (respectively Γ_+) are irrational, then there exists an f -orbit \mathcal{O} that is α -asymptotic to Γ_- and ω -asymptotic to Γ_+ .

The orbit of a point $x \in S^1 \times \mathbb{R}$ is called ω -asymptotic to Γ_+ if $d(f^n(x), \Gamma_+)$ converges to zero as $n \rightarrow \infty$. It is called α -asymptotic to Γ_- if $d(f^{-n}(x), \Gamma_-)$ converges to zero as $n \rightarrow \infty$.

The next theorem can be found as theorem 39.1 in [3] in a more general setting. It allows us to determine when the time- (s, t) maps of a time-periodic Hamiltonian on Z are twist maps.

Theorem 2.10. *Let $H : S^1 \times S^1 \times \mathbb{R} \rightarrow \mathbb{R}$ be a smooth Hamiltonian, such that the maps $h_{t,x} : \mathbb{R} \rightarrow \mathbb{R}$ with*

$$h_{t,x} : p \mapsto \partial_p H(t, x, p)$$

are diffeomorphisms for every $(t, x) \in S^1 \times S^1$ (global Legendre condition). Then, given any compact set $K \subset S^1 \times \mathbb{R}$ and starting time a , there exists an $\epsilon_0 > 0$, such that, for all $0 < \epsilon < \epsilon_0$ the time- $(a, a + \epsilon)$ map of H is a twist map on K .

3 Proof of the main result

To prove the main theorem we will define a time-periodic Lagrangian L on S^1 , such that the Euler-Lagrange solutions of L correspond to reparametrizations of F -geodesics, whose lifts are graphs over the euclidean line $e_1 \cdot \mathbb{R} \subset \mathbb{R}^2$. After a perturbation of the Lagrangian outside a tube $S^1 \times (-R, R)$ we will be able to work with the conjugated Hamiltonian H . Theorem 2.10 will guarantee that the time-1 map ψ of H is in \mathcal{P}^1 , i.e. a composition of finitely many positive monotone twist maps. We will then see that the instability region of the geodesic flow corresponds to a Birkhoff region of instability of ψ and hence Mather's connecting theorem 2.9 will guarantee the existence of an orbit connecting the boundary graphs of the Hamiltonian (respectively Lagrangian) system. Using the correspondence of Euler-Lagrange solutions and reparametrizations of geodesics we obtain a connecting geodesic with the required asymptotic behaviour.

Let $v \in \mathbb{Z} \setminus \{0\}$. Like in the previous section we associate a smooth and strictly convex time-periodic Lagrangian L on S^1 to the Finsler metric F via

$$L(t, x, r) = F\left(tv + xv^\perp, v + rv^\perp\right)$$

We perturb the Lagrangian for large values of $|r|$: For $R > 0$ we can find a constant $D > 0$, such that there is an extension L_R of $L|_{S^1 \times S^1 \times (-R, R)}$ to $S^1 \times S^1 \times \mathbb{R}$ with

1. $L_R(t, x, r) = \frac{D}{2}r^2$ for large values of $|r|$
2. L_R is strictly convex with second partial derivative bounded and bounded away from zero, i.e. $0 < \frac{1}{C} < \partial_{rr}L_R < C$ for a constant $C > 0$

Remark 3.1. *Observe that the Euler-Lagrange solutions of L_R for large values of $|r|$ are just straight lines, and hence the integral curves of the Euler-Lagrange vector field exist for all times. To see this lift L_R to a 1-periodic Lagrangian on \mathbb{R} and observe that for large values of $|r|$ the Euler-Lagrange equation reduces to $\ddot{\gamma} = 0$ for functions $\gamma : \mathbb{R} \rightarrow \mathbb{R}$. Thus, for large $|r|$ the solutions are straight lines. Consequently, for large $C > 0$ we have that the compact sets $S^1 \times S^1 \times [-C, C]$ are invariant under the Euler-Lagrange flow, and hence every solution exists for all times, i.e. the flow is complete.*

The following theorem is a corollary of theorem 2.2.

Theorem 3.2. *Let F be a Finsler metric on T^2 and let L_R be the above Lagrangian for an $R > 0$. Let $\theta : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function with $-R < \theta' < R$ and let $\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$ be the curve given by*

$$\gamma(t) = tv + \theta(t)v^\perp$$

Then γ is a reparametrization of a lift of an \tilde{F} -geodesic if and only if θ is an Euler-Lagrange solution of \tilde{L}_R .

Proof. The theorem follows immediately for a curve $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $-R < \theta' < R$ since the perturbed Lagrangian L_R agrees with the unperturbed Lagrangian L in a neighbourhood of the curve. Consequently, the Euler-Lagrange equations agree on this neighbourhood and hence θ is an Euler-Lagrange solution for L_R if and only if θ is an Euler-Lagrange solution for L . □

To define the associated Hamiltonian $H_R : S^1 \times S^1 \times \mathbb{R} \rightarrow \mathbb{R}$ of the Lagrangian L_R we need to consider the Legendre-transform $\mathcal{L}_t : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}$ with

$$\mathcal{L}_t : (x, r) \mapsto (x, \partial_r L(t, x, r))$$

For a moment we will omit the dependence on R and just write L instead of L_R . Note that the Legendre-transform is a diffeomorphism because $\partial_{rr}L$ is bounded away from zero. For the above Lagrangian L we can define the associated Hamiltonian $H : S^1 \times S^1 \times \mathbb{R} \rightarrow \mathbb{R}$ via

$$H(t, x, \partial_r L(t, x, r)) = \partial_r L(t, x, r)r - L(t, x, r) \quad (1)$$

The time- (s, t) maps of the Lagrangian and associated Hamiltonian for $s < t$ are conjugated via the Legendre transform

$$\psi^{s,t} = \mathcal{L}_t \circ \varphi^{s,t} \circ \mathcal{L}_s^{-1}$$

It follows from (1) and the strict convexity of L that $\frac{1}{C} < \partial_{pp}H < C$ for a constant $C > 0$.

Proposition 3.3. *The time-1 map $\psi : Z \rightarrow Z$ of the associated Hamiltonian $H_R : Z \rightarrow \mathbb{R}$ of the Lagrangian L_R is in \mathcal{P}^1 .*

Proof. Since $\partial_{pp}H(t, x, p) \neq 0$ the maps $h_{t,x}$ are local diffeomorphisms. Injectivity follows from $\partial_{pp}H > 0$ and surjectivity follows from $\partial_{pp}H$ being bounded away from zero. Consequently, for every large tube $S^1 \times [-P, P]$ (those tubes are actually invariant under the time-dependent flow of H if we chose P to be large enough) there is an $\epsilon > 0$, such that the restriction of $\psi_H^{0,\epsilon}$ to the tube is a twist map. Outside those tubes the Lagrangian L reduces to $L(t, x, r) = \frac{D}{2}r^2$ and hence with the Legendre-transform being equal to $\mathcal{L}_t(x, r) = (x, Dr)$ we find that for the Hamiltonian H it holds that

$$H(t, x, y) = \frac{y^2}{2D} \quad \text{for large } |y|$$

Thus, the Hamiltonian vector field is identical to $X_H(t, x, y) = (\frac{y}{D}, 0)$ outside large tubes and hence the time-1 map outside large tubes is identical to a shear-map of twist $\frac{1}{D}$. Since the starting time a in theorem 2.10 is arbitrary we can piece the time-1 map $\psi = \psi_H^{0,1} : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}$ together from finitely many twist maps, i.e. there exist real numbers $0 = \epsilon_0 < \dots < \epsilon_n = 1$, such that

$$\psi = \psi_H^{\epsilon_{n-1}, \epsilon_n} \circ \dots \circ \psi_H^{\epsilon_0, \epsilon_1}.$$

and $\psi_H^{\epsilon_i, \epsilon_{i+1}}$ is a twist map for every i . □

Lemma 3.4. *If $\Lambda \subset ST^2$ is an invariant torus with bounded direction (with respect to v) and $R > 0$ is chosen large enough, then the time-1 map of the Lagrangian L_R has a corresponding invariant circle Γ . If there are no closed geodesics on Λ the invariant circle has irrational rotation number.*

Remark 3.5. *By "Corresponding" in lemma 3.4 we mean that every Euler-Lagrange solution $t \mapsto (\theta_s(t), \partial_t \theta_s(t))$ of \tilde{L}_R with*

$$(\theta_s(0), \partial_t \theta_s(0)) = (s, \partial_t \theta_s(0)) \in \Gamma$$

corresponds to a reparametrization γ_s of the geodesic in Λ with $\gamma_s(0) = sv^\perp$.

Proof. (of lemma 3.4) It is useful to see Λ as a \mathbb{Z}^2 -periodic continuous graph in $S\mathbb{R}^2 \cong \mathbb{R}^2 \times S^1$, which is invariant under the geodesic flow $\phi : S\mathbb{R}^2 \times \mathbb{R} \rightarrow S\mathbb{R}^2$ of the lifted Finsler metric. First we want to define a reparametrization of the restricted geodesic flow $\phi : \Lambda \times \mathbb{R} \rightarrow \Lambda$ on Λ . Since the projections of the geodesics on Λ are graphs over $v\mathbb{R}$ and because of the periodicity of Λ , we have

$$\langle \pi_2 \circ \phi(w, t), v \rangle \geq \text{const} > 0 \quad (2)$$

for every $(w, t) \in \Lambda \times \mathbb{R}$. We define a smooth map $k : S\mathbb{R}^2 \times \mathbb{R} \rightarrow S\mathbb{R}^2 \times \mathbb{R}$ via

$$k(w, t) = \left(w, \int_0^t \langle \pi_2 \circ \phi(w, s), v \rangle ds \right)$$

It follows from (2) that k is a local diffeomorphism in a neighbourhood of every $(w, t) \in \Lambda \times \mathbb{R}$. Furthermore, it follows also from (2) that the restriction of k to $\Lambda \times \mathbb{R}$ is bijective and consequently the map $k : \Lambda \times \mathbb{R} \rightarrow \Lambda \times \mathbb{R}$ is a homeomorphism. Hence, we have the inverse $k^{-1} : \Lambda \times \mathbb{R} \rightarrow \Lambda \times \mathbb{R}$, which is continuous and differentiable in the second component. We define a reparametrization $\hat{\phi} : \Lambda \times \mathbb{R} \rightarrow \Lambda$ of the geodesic flow via

$$\hat{\phi}(w, t) = \phi \circ k^{-1}(w, \|v\|^2 t)$$

Note, that since the graph Λ is \mathbb{Z}^2 -periodic, we have for $z \in \mathbb{Z}^2$

$$\pi_1 \circ \hat{\phi}(w', t) = \pi_1 \circ \hat{\phi}(w, t) + z \quad \text{if } \pi_1(w') = \pi_1(w) + z \quad (3)$$

From $k \circ k^{-1} = id$ we calculate that

$$\langle \partial_t (\pi_1 \circ \hat{\phi})(w, t), v \rangle = \|v\|^2 \quad (4)$$

for every $(w, t) \in \Lambda$. Hence we can write the map $\pi_1 \circ \hat{\phi} : \Lambda \times \mathbb{R} \rightarrow \mathbb{R}^2$ as

$$\pi_1 \circ \hat{\phi}(w, t) = \left(\frac{1}{\|v\|^2} \langle \pi_1(w), v \rangle + t \right) v + \theta(w, t) v^\perp \quad (5)$$

where $\theta : \Lambda \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and differentiable in the second component. For $i \in \mathbb{Z}$ we define subsets $\Lambda_i \subset \Lambda$ via

$$\Lambda_i = \{w \in \Lambda \mid \pi_1(w) \in \mathbb{R}v^\perp + iv\}$$

Note, that for $w \in \Lambda_0$ we get from (5)

$$\pi_1 \circ \hat{\phi}(w, t) = tv + \theta(w, t)v^\perp \quad (6)$$

As an intermediate step we will show that

$$\hat{\phi}(w, t + t') = \hat{\phi}(\hat{\phi}(w, t'), t) \quad \text{for every } t, t' \in \mathbb{R} \quad (7)$$

To see this we define curves $\gamma_1, \gamma_2 : \mathbb{R} \rightarrow \Lambda$ via

$$\gamma_1(t) = \hat{\phi}(w, t' + t)$$

and

$$\gamma_2(t) = \hat{\phi}(\hat{\phi}(w, t'), t)$$

Since the curves are both reparametrizations of a geodesic on Λ with $\gamma_1(0) = \gamma_2(0)$ they are reparametrizations of the same geodesic. From equation (4) it follows for every t that

$$\langle \partial_t(\pi_1 \circ \gamma_1)(t), v \rangle = \langle \partial_t(\pi_1 \circ \gamma_2)(t), v \rangle$$

together with $\gamma_1(0) = \gamma_2(0)$ we obtain

$$\langle \pi_1 \circ \gamma_1(t), v \rangle = \langle \pi_1 \circ \gamma_2(t), v \rangle$$

for every $t \in \mathbb{R}$. Since every projection of a geodesic in Λ is a graph over $\mathbb{R}v$ this implies that

$$\pi_1 \circ \gamma_1(t) = \pi_1 \circ \gamma_2(t)$$

for every $t \in \mathbb{R}$. Finally, since Λ is a graph over \mathbb{R}^2 we have $\gamma_1 = \gamma_2$ and thus we have proven (7). Next we will prove that

$$\pi_1 \circ \hat{\phi}(w, 1) \in \Lambda_{i+1} \text{ for every } w \in \Lambda_i \quad (8)$$

To see this note that from (5) and $\langle \pi_1(w), v \rangle = i\|v\|^2$ it follows that

$$\langle \pi_1 \circ \hat{\phi}(w, 1), v \rangle = (i+1)\|v\|^2$$

We can now define a graph $\Gamma \subset S^1 \times \mathbb{R}$, which we will later prove to be an invariant circle of the time-1 map of L_R . We set

$$\Gamma = \{(\theta(w, 0), \partial_t \theta(w, 0)) \mid w \in \Lambda_0\}$$

To see that Γ is actually a graph in $S^1 \times \mathbb{R}$ note that for every $w \in \Lambda_0$ we have $\pi_1(w) = sv^\perp$ for an $s \in \mathbb{R}$. Consequently it follows from (6) that $\theta(w, 0) = s$ if $\pi_1(w) = sv^\perp$. To see that Γ is periodic (i.e. a subset in $S^1 \times \mathbb{R}$) note that if $w, w' \in \Lambda_0$ with $\pi_1(w) = sv^\perp$ and $\pi_1(w') = (s+1)v^\perp$ we have $\pi_1 \circ \hat{\phi}(w', t) = \pi_1 \circ \hat{\phi}(w, t) + v^\perp$ because of (3). Thus it follows from (6) that $\partial_t \theta(w, 0) = \partial_t \theta(w', 0)$ and hence Γ can be seen as the graph of the 1-periodic function $h : S^1 \rightarrow \mathbb{R}$ with

$$h(s) = \partial_t \theta(\Lambda(sv^\perp), 0)$$

Note, that Γ is a continuous graph since h is a composition of continuous functions. We can thus pick an $R > 0$ large enough, so that $\Gamma \subset S^1 \times (-R, R)$. To see now that Γ is invariant under the time-1 map $\varphi : S^1 \times \mathbb{R} \rightarrow S^1 \times \mathbb{R}$ of the Lagrangian L_R observe that for a $w \in \Lambda_0$ it follows from (5) and (7) that

$$\begin{aligned} \theta(w, 1+t) &= \frac{1}{\|v\|^2} \langle \pi_1 \circ \hat{\phi}(w, 1+t), v^\perp \rangle \\ &= \frac{1}{\|v\|^2} \langle \pi_1 \circ \hat{\phi}(\hat{\phi}(w, 1), t), v^\perp \rangle \end{aligned}$$

Since $\hat{\phi}(w, 1) \in \Lambda_1$ (see (8)) and because of the invariance (3) there exists a $w' \in \Lambda_0$, such that the last line is equal to

$$\begin{aligned}
&= \frac{1}{\|v\|^2} \langle \pi_1 \circ \hat{\phi}(w', t) + v, v^\perp \rangle \\
&= \frac{1}{\|v\|^2} \langle \pi_1 \circ \hat{\phi}(w', t), v^\perp \rangle \\
&= \theta(w', t)
\end{aligned}$$

for every $t \in \mathbb{R}$. This implies that $\theta(w, 1) = \theta(w', 0)$ and $\partial_t \theta(w, 1) = \partial_t \theta(w', 0)$ and hence using theorem 3.2 the time-1 map φ with

$$\varphi(\theta(w, 0), \partial_t \theta(w, 0)) = (\theta(w, 1), \partial_t \theta(w, 1))$$

maps Γ onto itself. Applying the Legendre transform to the graph Γ yields an invariant graph of the time-1 map ψ of the associated Hamiltonian H_R . Assume now, that Γ has rational rotation number. Consequently there exists a periodic point $x = (\theta(w, 0), \partial_t \theta(w, 0))$ in Γ . This implies that the geodesic with initial velocity w is closed, which is a contradiction to Λ having no closed geodesics. \square

Lemma 3.6. *Let $U \subset ST^2$ be an instability region with boundary graphs Λ_- and Λ_+ . Assume that U has bounded direction (with respect to $v = e_1$). If $R > 0$ is chosen large enough, such that the time-1 map of the Lagrangian L_R has two corresponding invariant circles Λ_- and Λ_+ , then the only invariant circles in the subset $B \subset S^1 \times \mathbb{R}$ enclosed by Λ_- and Λ_+ are the boundary circles Λ_- and Λ_+ .*

Proof. To prove this theorem we have to show that if we have an invariant circle Γ of the map φ , which is not equal to Γ_- or Γ_+ then we also have an ϕ -invariant torus Λ not equal to Λ_- or Λ_+ . Now, assume that $\Gamma \subset S^1 \times \mathbb{R}$ is such an invariant circle of φ . It follows from theorem 2.7 that Γ is equal to the graph $\{(s, h(s)) | s \in S^1\}$ of a Lipschitz function $h : S^1 \rightarrow \mathbb{R}$. For every fixed $s \in S^1$ the map $t \mapsto \varphi^{0,t}(s, h(s))$ is a trajectory of the Euler-Lagrange flow on $S^1 \times \mathbb{R}$. We define maps $\theta_s : \mathbb{R} \rightarrow \mathbb{R}$ via

$$(\theta_s(t), \theta'_s(t)) := \varphi^{0,t}(s, h(s))$$

Note that the maps $(s, t) \mapsto \theta(s, t)$ and $(s, t) \mapsto \theta'_s(t)$ are continuous, since θ_s and θ'_s are the compositions of $\varphi^{0,t}(s, h(s))$ with the projections of $S^1 \times \mathbb{R}$ onto S^1 respectively \mathbb{R} . We extend θ to a map $\theta : \mathbb{R}^2 \rightarrow \mathbb{R}$, $(s, t) \mapsto \theta_s(t)$ by seeing Γ as a 1-periodic graph in \mathbb{R}^2 and using the Euler-Lagrange flow φ of the lifted Lagrangian \tilde{L} in the definition of θ . We then have

$$(\theta_{s+1}(t), \theta'_{s+1}(t)) = (\theta_s(t) + 1, \theta'_s(t))$$

Observe that since $\varphi^{0,0} = id$ we have $\theta_s(0) = s$ for every $s \in \mathbb{R}$. Note that for $r \neq p$ we have $\theta_r(t) \neq \theta_p(t)$. To see this assume that we have $\theta_r(t) = \theta_p(t)$. Since the set $\varphi^{0,t}(\Gamma) = \{(\theta_s(t), \theta'_s(t)) | s \in \mathbb{R}\}$ is invariant under the map $\varphi^{t,t+1}$ it is a Lipschitz graph. Hence it also holds that $\theta'_r(t) = \theta'_p(t)$. We have the associated reparametrizations of geodesics $\gamma_r(t) = tv + \theta_r(t)v^\perp$ and $\gamma_p(t) = tv + \theta_p(t)v^\perp$. This would imply that the geodesics meet at the point $\gamma_r(t) = \gamma_p(t)$ with $\dot{\gamma}_r(t) = \dot{\gamma}_p(t)$, which is not possible because geodesics can not become tangent to each other. As mentioned above, since the sets $\Gamma_t := \varphi^{0,t}(\Gamma)$ are Lipschitz graphs there exist periodic Lipschitz functions $h_t : \mathbb{R} \rightarrow \mathbb{R}$ with $\Gamma_t = \{(s, h_t(s)) | s \in \mathbb{R}\}$ with

$$h_t(\theta_s(t)) = \theta'_s(t).$$

We define a function $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ via

$$g : (s, t) \mapsto h_t(s).$$

We will now show that g is continuous. To see this take real sequences $t_n \rightarrow t$ and $s_n \rightarrow s$. Since Γ_t is a Lipschitz graph there exists exactly one $r_n \in \mathbb{R}$ for every $n \in \mathbb{N}$ with

$$\theta_{r_n}(t_n) = s_n.$$

The sequence r_n is bounded. To see this observe that it follows from the continuity of the map $(s, t) \mapsto \theta_s(t)$ and the graph property of the sets Γ_t that the map $s \mapsto \theta_s(t)$ is strictly increasing for every fixed t . Now, assume that the sequence r_n is unbounded. Without loss of generality we assume that r_n is unbounded from above with $r_n \geq n$ for every $n \in \mathbb{N}$. Then we have

$$\theta_{r_n}(t_n) \geq \theta_n(t_n) = \theta_0(t_n) + n$$

and thus the sequence $\theta_{r_n}(t_n)$ is also unbounded. This contradicts the fact that s_n converges to s and consequently the sequence r_n has to be bounded. To see that r_n actually converges assume that there are two limit points $r = \lim_{i \rightarrow \infty} r_{n_i}$ and $p = \lim_{j \rightarrow \infty} r_{n_j}$. Since we have $\theta_{r_{n_i}}(t_{n_i}) = s_{n_i}$ it follows from the continuity of θ that $\theta_r(t) = s$. Analogously we obtain $\theta_p(t) = s$. As above it follows that $r = p$. To see that g is continuous observe that

$$h_{t_n}(s_n) = h_{t_n}(\theta_{r_n}(t_n)) = \theta'_{r_n}(t_n)$$

and hence it follows from the continuity of $(s, t) \mapsto \theta'_s(t)$ that

$$\lim_{n \rightarrow \infty} h_{t_n}(s_n) = \lim_{n \rightarrow \infty} \theta'_{r_n}(t_n) = \theta'_r(t) = h_t(s)$$

Thus we have proven that $g : (s, t) \mapsto h_t(s)$ is continuous. To construct an invariant graph in ST^2 , observe that since the reparametrized geodesics γ_s with $\gamma_s(t) = tv + \theta_s(t)v^\perp$ can not cross and since $\gamma_{s+1} = \gamma_s + v^\perp$ the geodesics γ_s foliate \mathbb{R}^2 . This allows us to define a graph $\Lambda \subset S\mathbb{R}^2$ via

$$\Lambda = \left\{ \left(\gamma_s(t), \frac{\dot{\gamma}_s(t)}{F(\dot{\gamma}_s(t))} \right) : (s, t) \in \mathbb{R}^2 \right\}$$

Furthermore, Λ is ϕ -invariant since it only consists of images of geodesics. We define a map $X : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ via

$$X : x \mapsto v + h_{\left\langle x, \frac{v}{\|v\|^2} \right\rangle} \left(\left\langle x, \frac{v^\perp}{\|v\|^2} \right\rangle \right) v^\perp$$

Observe that X is continuous (because h is) and maps $\gamma_s(t)$ to $\dot{\gamma}_s(t)$. Hence Λ is the graph of the continuous function

$$x \mapsto \frac{X(x)}{F(X(x))}$$

What is left to prove is that X is \mathbb{Z}^2 periodic, i.e. Λ is actually a graph in ST^2 . We will show that if we assume our instability region to have bounded direction with respect to $v \in \mathbb{Z}^2 - \{0\}$ then X is $v\mathbb{Z} \times v^\perp\mathbb{Z}$ -periodic. As a special case we get that X is \mathbb{Z}^2 -periodic if v is chosen to be equal to e_1 . To prove that X (and thus $\frac{X}{F(X)}$) is $v\mathbb{Z} \times v^\perp\mathbb{Z}$ -periodic observe that for every $(s, t) \in \mathbb{R}^2$ we have

$$\begin{aligned} \gamma_s(t) + v^\perp &= tv + \theta_s(t)v^\perp + v^\perp \\ &= tv + (\theta_s(t) + 1)v^\perp \\ &= tv + \theta_{s+1}(t)v^\perp \\ &= \gamma_{s+1}(t) \end{aligned}$$

Since $\theta'_s(t) = \theta'_{s+1}(t)$ we have $\dot{\gamma}_s(t) = \dot{\gamma}_{s+1}(t)$ and thus X is $v\mathbb{Z}$ -periodic. From the invariance of Γ_t under $\varphi^{t,t+1}$ we obtain that for every $s \in \mathbb{R}$ there exists an $r \in \mathbb{R}$ with

$$\varphi^{t,t+1}(\theta_r(t), \theta'_r(t)) = (\theta_s(t), \theta'_s(t))$$

Thus we have

$$\begin{aligned} \gamma_s(t) + v &= (t+1)v + \theta_s(t)v^\perp \\ &= (t+1)v + \theta_r(t+1)v^\perp \\ &= \gamma_r(t+1) \end{aligned}$$

And since $\theta'_r(t+1) = \theta'_s(t)$ we have $\dot{\gamma}_r(t+1) = \dot{\gamma}_s(t)$. Hence we have shown the $v\mathbb{Z}$ -periodicity of X . □

We can now prove the main theorem.

Proof. (theorem 1.4)

If $R > 0$ is large enough lemma 3.4 guarantees that the time-1 map φ of L_R has two invariant circles Γ_- and Γ_+ in $S^1 \times (-R, R)$, which correspond to the invariant boundary graphs. With lemma 3.6 it follows that the region bounded by the corresponding invariant circles of the Hamiltonian time-1 map is a Birkhoff region of instability. Thus, after going back to the Lagrangian setting, we obtain from theorem 2.9 a point x in the area between the boundary circles, and sequences $t'_n \rightarrow \infty$, $s'_n \rightarrow -\infty$ with

$$\lim_{n \rightarrow \infty} \varphi^{0,t'_n}(x) \in \Gamma_+$$

and

$$\lim_{n \rightarrow \infty} \varphi^{0,s'_n}(x) \in \Gamma_-$$

Using theorem 3.2 we obtain after reparametrization a geodesic $c : \mathbb{R} \rightarrow T^2$ and sequences $t_n \rightarrow \infty$, $s_n \rightarrow -\infty$ with

$$\lim_{n \rightarrow \infty} \dot{c}(t_n) \in \Lambda_+$$

and

$$\lim_{n \rightarrow \infty} \dot{c}(s_n) \in \Lambda_-$$

□

References

- [1] D. Bao, S.-S. Chern, Z. Shen, *An Introduction to Riemann-Finsler Geometry*, Springer (2000)
- [2] G. D. Birkhoff, *Surface transformations and their dynamical applications*, Collected Math. Papers, vol. 2, (1920)
- [3] C. Golé, *Symplectic Twist Maps*, World Scientific Publishing, 2001.
- [4] J. Mather, *Variational Construction of Orbits of Twist Diffeomorphisms*, Journal of the American Math. Soc., 4 (1991), 207 - 263
- [5] J. P. Schröder, *Tonelli Lagrangians on the 2-torus: global minimizers, invariant tori and topological entropy*, Ph.D. thesis, Ruhr-Universität Bochum (2013)