

# New Type I Binary $[72, 36, 12]$ Self-Dual Codes from Composite Matrices and $R_1$ Lifts

Adrian Korban

Department of Mathematical and Physical Sciences

University of Chester

Thornton Science Park, Pool Ln, Chester CH2 4NU, England

Serap Şahinkaya

Tarsus University, Faculty of Engineering

Department of Natural and Mathematical Sciences

Mersin, Turkey

Deniz Ustun

Tarsus University, Faculty of Engineering

Department of Computer Engineering

Mersin, Turkey

December 28, 2021

## Abstract

In this work, we define three composite matrices derived from group rings. We employ these composite matrices to create generator matrices of the form  $[I_n \mid \Omega(v)]$ , where  $I_n$  is the identity matrix and  $\Omega(v)$  is a composite matrix and search for binary self-dual codes with parameters  $[36, 18, 6]$  or  $[8]$ . We next lift these codes over the ring  $R_1 = \mathbb{F}_2 + u\mathbb{F}_2$  to obtain codes whose binary images are self-dual codes with parameters  $[72, 36, 12]$ . Many of these codes turn out to have weight enumerators with parameters that were not known in the literature before. In particular, we find 30 new Type I binary self-dual codes with parameters  $[72, 36, 12]$ .

# 1 Introduction

In this work, we employ the idea of composite matrices from group rings that was introduced in [5], to search for binary self-dual codes. In particular, we consider a generator matrix of the form  $[I_n \mid \Omega(v)]$ , where  $I_n$  is the identity matrix and  $\Omega(v)$  is a composite matrix where  $v$  is an element in the group ring  $RG$ , to search for binary self-dual codes. Similar approach for finding extremal binary self-dual codes of length 68 can be found in [6, 7]. We describe three composite matrices from group rings with the use of groups of orders 18, 9 and 6 which we then employ to search for binary self-dual codes with parameters  $[36, 18, 6$  or  $8]$ . We next lift these codes over the ring  $\mathbb{F}_2 + u\mathbb{F}_2$ , to obtain codes whose binary images are self-dual codes of length 72. Many of these codes turn out to be new Type I binary self-dual codes with parameters  $[72, 36, 12]$ .

The rest of the work is organized as follows. In Section 2, we give preliminary definitions and results on codes, the alphabets we use, special matrices, group rings and composite matrices from group rings. In Section 3, we construct three composite matrices of order 18 by using group rings and groups of different cardinalities, in particular, groups with cardinalities 18, 9 and 6. Next, in Section 4, we define three generator matrices in which we use the composite matrices described in Section 3. We show the conditions that each generator matrix needs to meet in order to produce self-dual codes. In Section 5, we present all the computational results obtained from searching for self-dual codes with different parameters and over different alphabets and with the use of the generator matrices presented in Section 4. We finish with concluding remarks and directions for possible future research.

## 2 Preliminaries

### 2.1 Codes

We begin by recalling the standard definitions from coding theory. A code  $C$  of length  $n$  over a Frobenius ring  $R$  is a subset of  $R^n$ . If the code is a submodule of  $R^n$  then we say that the code is linear. Elements of the code  $C$  are called codewords of  $C$ . Let  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  and  $\mathbf{y} = (y_1, y_2, \dots, y_n)$  be two elements of  $R^n$ . The duality is understood in terms of the Euclidean

inner product, namely:

$$\langle \mathbf{x}, \mathbf{y} \rangle_E = \sum x_i y_i.$$

The dual  $C^\perp$  of the code  $C$  is defined as

$$C^\perp = \{ \mathbf{x} \in R^n \mid \langle \mathbf{x}, \mathbf{y} \rangle_E = 0 \text{ for all } \mathbf{y} \in C \}.$$

We say that  $C$  is self-orthogonal if  $C \subseteq C^\perp$  and is self-dual if  $C = C^\perp$ .

An upper bound on the minimum Hamming distance of a binary self-dual code was given in [17]. Specifically, let  $d_I(n)$  and  $d_{II}(n)$  be the minimum distance of a Type I and Type II binary code of length  $n$ , respectively. Then

$$d_{II}(n) \leq 4 \lfloor \frac{n}{24} \rfloor + 4$$

and

$$d_I(n) \leq \begin{cases} 4 \lfloor \frac{n}{24} \rfloor + 4 & \text{if } n \not\equiv 22 \pmod{24} \\ 4 \lfloor \frac{n}{24} \rfloor + 6 & \text{if } n \equiv 22 \pmod{24}. \end{cases}$$

Self-dual codes meeting these bounds are called *extremal*. Throughout the text, we obtain extremal binary codes of different lengths. Self-dual codes which are the best possible for a given set of parameters is said to be optimal. Extremal codes are necessarily optimal but optimal codes are not necessarily extremal.

## 2.2 The ring $R_1 = \mathbb{F}_2 + u\mathbb{F}_2$

In this section, we recall some theory on self-dual codes over  $\mathbb{F}_2 + u\mathbb{F}_2$ . We refer to [4] where Type II, Type IV, self-dual codes and cyclic codes over  $\mathbb{F}_2 + u\mathbb{F}_2$  were studied.

The ring  $\mathbb{F}_2 + u\mathbb{F}_2$  is a ring of characteristic 2 with 4 elements with the restriction  $u^2 = 0$ . It is defined as

$$\mathbb{F}_2 + u\mathbb{F}_2 = \{ a + bu \mid a, b \in \mathbb{F}_2, u^2 = 0 \},$$

and it is easily seen that  $\mathbb{F}_2 + u\mathbb{F}_2 \cong \mathbb{F}_2[x]/(x^2)$ . A linear code  $C$  of length  $n$  over the ring  $\mathbb{F}_2 + u\mathbb{F}_2$  is an  $\mathbb{F}_2 + u\mathbb{F}_2$ -submodule of  $(\mathbb{F}_2 + u\mathbb{F}_2)^n$ . The elements of  $\mathbb{F}_2 + u\mathbb{F}_2$  are  $0, 1, u, 1 + u$  and their Lee weights are defined as  $0, 1, 2, 1$  respectively. The Hamming ( $d_H$ ) and Lee ( $d_L$ ) distance between  $n$  tuples is then defined as the sum of the Hamming and Lee weights of the difference of

the components of these tuples respectively. The smallest positive Hamming and Lee distance of a code  $C$  is denoted by  $d_H(C)$  and  $d_L(C)$  respectively.

A Gray map  $\phi$  is defined as

$$\begin{aligned}\phi : (\mathbb{F}_2 + u\mathbb{F}_2)^n &\rightarrow \mathbb{F}_2^{2n}, \\ \phi(\bar{a} + \bar{b}u) &= (\bar{b}, \bar{a} + \bar{b}),\end{aligned}$$

where  $\bar{a}, \bar{b} \in \mathbb{F}_2^n$ . The map is a distance preserving isometry from  $((\mathbb{F}_2 + u\mathbb{F}_2)^n, d_L)$  to  $(\mathbb{F}_2^{2n}, d_H)$ , where  $d_L$  and  $d_H$  denote the Lee and Hamming distance in  $(\mathbb{F}_2 + u\mathbb{F}_2)^n$  and  $\mathbb{F}_2^{2n}$  respectively. This means that if  $C$  is a linear code over  $\mathbb{F}_2 + u\mathbb{F}_2$  with parameters  $[n, 2^k, d]$  ( $2^k$  is the number of the codewords), then  $\phi(C)$  is a binary linear code of parameters  $[2n, k, d]$ . The following theorem is a natural result of the Gray map.

**Theorem 2.1.** *If  $C$  is a self-dual code over  $\mathbb{F}_2 + u\mathbb{F}_2$  of length  $n$ , then  $\phi(C)$  is a self-dual binary code of length  $2n$ .*

We can also define a natural projection from  $\mathbb{F}_2 + u\mathbb{F}_2$  to  $\mathbb{F}_2$  as follows:

$$\begin{aligned}\mu : \mathbb{F}_2 + u\mathbb{F}_2 &\rightarrow \mathbb{F}_2, \\ \mu(a + bu) &= a.\end{aligned}$$

If  $D = \mu(C)$  for some linear code  $C$  over  $\mathbb{F}_2 + u\mathbb{F}_2$ , we say that  $D$  is a projection of  $C$  into  $\mathbb{F}_2$ , and that  $C$  is a lift of  $D$  into  $\mathbb{F}_2 + u\mathbb{F}_2$ . It is clear that the projection of a self-orthogonal code is self-orthogonal, but the projection of a self-dual code need not be self-dual. We finish this section with two well known results.

**Theorem 2.2.** *Suppose that  $C$  is a self-dual code over  $\mathbb{F}_2 + u\mathbb{F}_2$  of length  $2n$ , generated by the matrix  $[I_n|A]$ , where  $I_n$  is the  $n \times n$  identity matrix. Then  $\mu(C)$  is a self-dual binary code of length  $2n$ .*

**Theorem 2.3.** *Suppose  $C$  is a linear code over  $\mathbb{F}_2 + u\mathbb{F}_2$  and that  $C' = \mu(C)$ , is its projection to  $\mathbb{F}_2$ . With  $d$  and  $d'$  representing the minimum Lee and Hamming distances of  $C$  and  $C'$  respectively, we have that  $d \leq 2d'$ .*

### 2.3 Special Matrices and Group Rings

To understand the composite construction which we define later in this work, we recall some basic definitions of some special matrices and theory on group rings and the map that sends group ring elements to matrices.

A circulant matrix is one where each row is shifted one element to the right relative to the preceding row. We label the circulant matrix as  $A = \text{circ}(\alpha_1, \alpha_2 \dots, \alpha_n)$ , where  $\alpha_i$  are ring elements. The transpose of a matrix  $A$ , denoted by  $A^T$ , is a matrix whose rows are the columns of  $A$ , i.e.,  $A_{ij}^T = A_{ji}$ . A symmetric matrix is a square matrix that is equal to its transpose.

While group rings can be given for infinite rings and infinite groups, we are only concerned with group rings where both the ring and the group are finite. Let  $G$  be a finite group of order  $n$ , then the group ring  $RG$  consists of  $\sum_{i=1}^n \alpha_i g_i$ ,  $\alpha_i \in R$ ,  $g_i \in G$ .

Addition in the group ring is done by coordinate addition, namely

$$\sum_{i=1}^n \alpha_i g_i + \sum_{i=1}^n \beta_i g_i = \sum_{i=1}^n (\alpha_i + \beta_i) g_i. \quad (2.1)$$

The product of two elements in a group ring is given by

$$\left( \sum_{i=1}^n \alpha_i g_i \right) \left( \sum_{j=1}^n \beta_j g_j \right) = \sum_{i,j} \alpha_i \beta_j g_i g_j. \quad (2.2)$$

It follows that the coefficient of  $g_k$  in the product is  $\sum_{g_i g_j = g_k} \alpha_i \beta_j$ .

The following construction of a matrix was first given for codes over fields by Hurley in [13]. It was extended to Frobenius rings in [8]. Let  $R$  be a finite commutative Frobenius ring and let  $G = \{g_1, g_2, \dots, g_n\}$  be a group of order  $n$  and let  $v = \sum_{i=1}^n \alpha_i g_i \in RG$ . Define the matrix  $\sigma(v) \in M_n(R)$  to be

$$\sigma(v) = \begin{pmatrix} \alpha_{g_1^{-1}g_1} & \alpha_{g_1^{-1}g_2} & \alpha_{g_1^{-1}g_3} & \dots & \alpha_{g_1^{-1}g_n} \\ \alpha_{g_2^{-1}g_1} & \alpha_{g_2^{-1}g_2} & \alpha_{g_2^{-1}g_3} & \dots & \alpha_{g_2^{-1}g_n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{g_n^{-1}g_1} & \alpha_{g_n^{-1}g_2} & \alpha_{g_n^{-1}g_3} & \dots & \alpha_{g_n^{-1}g_n} \end{pmatrix}. \quad (2.3)$$

We note that the elements  $g_1^{-1}, g_2^{-1}, \dots, g_n^{-1}$  are the elements of the group  $G$  in a some given order.

## 2.4 Composite Matrices from Group Rings

In this section, we recall a matrix construction which is an extension of the matrix  $\sigma(v)$  in Equation (2.3). This extension was first introduced in [5]. With this extension, one can produce many interesting  $n \times n$  matrices for different choices of not just one group as in the case of the matrix  $\sigma(v)$ , but for different choices of more than one group, please see [5] for more details. Such matrices are referred to as composite constructions or just composite matrices.

Let  $R$  be a finite commutative Frobenius ring. Let  $\{g_1, g_2, \dots, g_n\}$  be a fixed listing of the elements of  $G$ . Let  $\{(h_i)_1, (h_i)_2, \dots, (h_i)_r\}$  be a fixed listing of the elements of  $H_i$ , where  $H_i$  is any group of order  $r$ . Let  $r$  be a factor of  $n$  with  $n > r$  and  $n, r \neq 1$ . Also, let  $G_r$  be a subset of  $G$  containing  $r$  distinct elements of  $G$ . Define the map:

$$\begin{array}{ccc} \phi : H & \mapsto & G_r \\ h_1 & \xrightarrow{\phi} & g_1 \\ h_2 & \xrightarrow{\phi} & g_2 \\ \vdots & \vdots & \vdots \\ h_r & \xrightarrow{\phi} & g_r. \end{array}$$

This map sends  $r$  distinct elements of the group  $H$  to  $r$  distinct elements of the group  $G$ .

Let  $v = \alpha_{g_1}g_1 + \alpha_{g_2}g_2 + \dots, \alpha_{g_n}g_n \in RG$ . Define the matrix  $\Omega(v) \in M_n(R)$  to be

$$\Omega(v) = \begin{pmatrix} A_1 & A_2 & A_3 & \dots & A_{\frac{n}{r}} \\ A_{\frac{n}{r}+1} & A_{\frac{n}{r}+2} & A_{\frac{n}{r}+3} & \dots & A_{\frac{2n}{r}} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ A_{\frac{(r-1)n}{r}+1} & A_{\frac{(r-1)n}{r}+2} & A_{\frac{(r-1)n}{r}+3} & \dots & A_{\frac{n^2}{r^2}} \end{pmatrix}, \quad (2.4)$$

where at least one block has the following form:

$$A_l = \begin{pmatrix} \alpha_{g_j^{-1}g_k} & \alpha_{g_j^{-1}g_{k+1}} & \cdots & \alpha_{g_j^{-1}g_{k+(r-1)}} \\ \alpha_{g_{j+1}^{-1}g_k} & \alpha_{g_{j+1}^{-1}g_{k+1}} & \cdots & \alpha_{g_{j+1}^{-1}g_{k+(r-1)}} \\ \alpha_{g_{j+2}^{-1}g_k} & \alpha_{g_{j+2}^{-1}g_{k+1}} & \cdots & \alpha_{g_{j+2}^{-1}g_{k+(r-1)}} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{g_{j+r-1}^{-1}g_k} & \alpha_{g_{j+r-1}^{-1}g_{k+1}} & \cdots & \alpha_{g_{j+r-1}^{-1}g_{k+(r-1)}} \end{pmatrix},$$

and the other blocks are of the form:

$$A'_l = \begin{pmatrix} \alpha_{g_j^{-1}g_k} & \alpha_{g_j^{-1}g_{k+1}} & \cdots & \alpha_{g_j^{-1}g_{k+(r-1)}} \\ \alpha_{\phi_l((h_i)_2^{-1}(h_i)_1)} & \alpha_{\phi_l((h_i)_2^{-1}(h_i)_2)} & \cdots & \alpha_{\phi_l((h_i)_2^{-1}(h_i)_r)} \\ \alpha_{\phi_l((h_i)_3^{-1}(h_i)_1)} & \alpha_{\phi_l((h_i)_3^{-1}(h_i)_2)} & \cdots & \alpha_{\phi_l((h_i)_3^{-1}(h_i)_r)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_l((h_i)_r^{-1}(h_i)_1)} & \alpha_{\phi_l((h_i)_r^{-1}(h_i)_2)} & \cdots & \alpha_{\phi_l((h_i)_r^{-1}(h_i)_r)} \end{pmatrix},$$

where  $l = \{1, 2, 3, \dots, \frac{n^2}{r^2}\}$  and where:

$$\begin{array}{ccc} \phi_l : H_i & \mapsto & G_r \\ (h_i)_1 & \xrightarrow{\phi_l} & g_j^{-1}g_k \\ (h_i)_2 & \xrightarrow{\phi_l} & g_j^{-1}g_{k+1} \\ \vdots & \vdots & \vdots \\ (h_i)_r & \xrightarrow{\phi_l} & g_j^{-1}g_{k+(r-1)}. \end{array}$$

Here we notice that when  $l = 1$  then  $j = 1, k = 1$ , when  $l = 2$  then  $j = 1, k = r + 1$ , when  $l = 3$  then  $j = 1, k = 2r + 1, \dots$  when  $l = \frac{n}{r}$  then  $j = 1, k = n - r + 1$ . When  $l = \frac{n}{r} + 1$  then  $j = r + 1, k = 1$ , when  $l = \frac{n}{r} + 2$  then  $j = r + 1, k = r + 1$ , when  $l = \frac{n}{r} + 3$  then  $j = r + 1, k = 2r + 1, \dots$  when  $l = \frac{2n}{r}$  then  $j = r + 1, k = n - r + 1, \dots$ , and so on.

### 3 Composite Matrices

In this section, we employ the matrix construction given in Equation (2.4) to build three composite matrices. We particularly consider some groups of orders 18, 9 and 6 to construct composite matrices of order 18 which we later use to search for binary self-dual codes.

1. Let  $G = \langle x, y \mid x^9 = y^2 = x^y = x^{-1} \rangle \cong D_{18}$ . Let  $v_1 = \sum_{i=0}^8 \sum_{j=0}^1 \alpha_{1+i+9j} y^j x^i \in RD_{18}$ . Also, let  $H = \langle a \mid a^9 = 1 \rangle \cong C_{3,3}$ . We now define the composite matrix over  $R$  as

$$\Omega(v_1) = \begin{pmatrix} A'_1 & A'_2 \\ A'_3 & A'_4 \end{pmatrix}$$

where

$$A'_1 = \begin{pmatrix} \alpha_{g_1^{-1}g_1} & \alpha_{g_1^{-1}g_2} & \alpha_{g_1^{-1}g_3} & \cdots & \alpha_{g_1^{-1}g_9} \\ \alpha_{\phi_1(h_2^{-1}h_1)} & \alpha_{\phi_1(h_2^{-1}h_2)} & \alpha_{\phi_1(h_2^{-1}h_3)} & \cdots & \alpha_{\phi_1(h_2^{-1}h_9)} \\ \alpha_{\phi_1(h_3^{-1}h_1)} & \alpha_{\phi_1(h_3^{-1}h_2)} & \alpha_{\phi_1(h_3^{-1}h_3)} & \cdots & \alpha_{\phi_1(h_3^{-1}h_9)} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_1(h_9^{-1}h_1)} & \alpha_{\phi_1(h_9^{-1}h_2)} & \alpha_{\phi_1(h_9^{-1}h_3)} & \cdots & \alpha_{\phi_1(h_9^{-1}h_9)} \end{pmatrix}$$

with

$$\phi_1 : h_j^{-1}h_i \xrightarrow{\phi_1} g_1^{-1}g_i$$

for when

$$j = 2 : i = 1, 2, \dots, 9$$

$$j = 3 : i = 1, 2, \dots, 9$$

$\vdots$

$$j = 9 : i = 1, 2, \dots, 9,$$

$$A'_2 = \begin{pmatrix} \alpha_{g_1^{-1}g_{10}} & \alpha_{g_1^{-1}g_{11}} & \alpha_{g_1^{-1}g_{12}} & \cdots & \alpha_{g_1^{-1}g_{18}} \\ \alpha_{\phi_2(h_2^{-1}h_1)} & \alpha_{\phi_2(h_2^{-1}h_2)} & \alpha_{\phi_2(h_2^{-1}h_3)} & \cdots & \alpha_{\phi_2(h_2^{-1}h_9)} \\ \alpha_{\phi_2(h_3^{-1}h_1)} & \alpha_{\phi_2(h_3^{-1}h_2)} & \alpha_{\phi_2(h_3^{-1}h_3)} & \cdots & \alpha_{\phi_2(h_3^{-1}h_9)} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_2(h_9^{-1}h_1)} & \alpha_{\phi_2(h_9^{-1}h_2)} & \alpha_{\phi_2(h_9^{-1}h_3)} & \cdots & \alpha_{\phi_2(h_9^{-1}h_9)} \end{pmatrix}$$

with

$$\phi_2 : h_j^{-1}h_k \xrightarrow{\phi_2} g_1^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 10; k = 2, i = 11; \dots k = 9, i = 18,$$

$$j = 3 : k = 1, i = 10; k = 2, i = 11; \dots k = 9, i = 18,$$

$\vdots$



$$j = 9 : k = 1, i = 10; k = 2, i = 11; \dots k = 9, i = 18,$$

$$A'_3 = \begin{pmatrix} \alpha_{g_{10}^{-1}g_1} & \alpha_{g_{10}^{-1}g_2} & \alpha_{g_{10}^{-1}g_3} & \dots & \alpha_{g_{10}^{-1}g_9} \\ \alpha_{\phi_3(h_2^{-1}h_1)} & \alpha_{\phi_3(h_2^{-1}h_2)} & \alpha_{\phi_3(h_2^{-1}h_3)} & \dots & \alpha_{\phi_3(h_2^{-1}h_9)} \\ \alpha_{\phi_3(h_3^{-1}h_1)} & \alpha_{\phi_3(h_3^{-1}h_2)} & \alpha_{\phi_3(h_3^{-1}h_3)} & \dots & \alpha_{\phi_3(h_3^{-1}h_9)} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_3(h_9^{-1}h_1)} & \alpha_{\phi_3(h_9^{-1}h_2)} & \alpha_{\phi_3(h_9^{-1}h_3)} & \dots & \alpha_{\phi_3(h_9^{-1}h_9)} \end{pmatrix}$$

with

$$\phi_3 : h_j^{-1}h_i \xrightarrow{\phi_3} g_{10}^{-1}g_i$$

for when

$$j = 2 : i = 1, 2, \dots, 9$$

$$j = 3 : i = 1, 2, \dots, 9$$

$\vdots$

$$j = 9 : i = 1, 2, \dots, 9,$$

and

$$A'_4 = \begin{pmatrix} \alpha_{g_{10}^{-1}g_{10}} & \alpha_{g_{10}^{-1}g_{11}} & \alpha_{g_{10}^{-1}g_{12}} & \dots & \alpha_{g_{10}^{-1}g_{18}} \\ \alpha_{\phi_4(h_2^{-1}h_1)} & \alpha_{\phi_4(h_2^{-1}h_2)} & \alpha_{\phi_4(h_2^{-1}h_3)} & \dots & \alpha_{\phi_4(h_2^{-1}h_9)} \\ \alpha_{\phi_4(h_3^{-1}h_1)} & \alpha_{\phi_4(h_3^{-1}h_2)} & \alpha_{\phi_4(h_3^{-1}h_3)} & \dots & \alpha_{\phi_4(h_3^{-1}h_9)} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_4(h_9^{-1}h_1)} & \alpha_{\phi_4(h_9^{-1}h_2)} & \alpha_{\phi_4(h_9^{-1}h_3)} & \dots & \alpha_{\phi_4(h_9^{-1}h_9)} \end{pmatrix}$$

with

$$\phi_4 : h_j^{-1}h_k \xrightarrow{\phi_4} g_{10}^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 10; k = 2, i = 11; \dots k = 9, i = 18,$$

$$j = 3 : k = 1, i = 10; k = 2, i = 11; \dots k = 9, i = 18,$$

$\vdots$

$$j = 9 : k = 1, i = 10; k = 2, i = 11; \dots k = 9, i = 18.$$

This results in a composite matrix over  $R$  of the following form:

$$\begin{aligned}\Omega(v_1) &= \begin{pmatrix} A'_1 & A'_2 \\ A'_3 & A'_4 \end{pmatrix} = \begin{pmatrix} B & C \\ D & E \end{pmatrix} = \\ &= \begin{pmatrix} B_1 & B_2 & B_3 & C_1 & C_2 & C_3 \\ B'_3 & B_1 & B_2 & C'_3 & C_1 & C_2 \\ B'_2 & B'_3 & B_1 & C'_2 & C'_3 & C_1 \\ D_1 & D_2 & D_3 & E_1 & E_2 & E_3 \\ D'_3 & D_1 & D_2 & E'_3 & E_1 & E_2 \\ D'_2 & D'_3 & D_1 & E'_2 & E'_3 & E_1 \end{pmatrix},\end{aligned}$$

where

$$\begin{aligned}B_1 &= \text{circ}(\alpha_1, \alpha_2, \alpha_3), B_2 = \text{circ}(\alpha_4, \alpha_5, \alpha_6), B_3 = \text{circ}(\alpha_7, \alpha_8, \alpha_9), \\ B'_2 &= \text{circ}(\alpha_6, \alpha_4, \alpha_5), B'_3 = \text{circ}(\alpha_9, \alpha_7, \alpha_8), C_1 = \text{circ}(\alpha_{10}, \alpha_{11}, \alpha_{12}), \\ C_2 &= \text{circ}(\alpha_{13}, \alpha_{14}, \alpha_{15}), C_3 = \text{circ}(\alpha_{16}, \alpha_{17}, \alpha_{18}), C'_2 = \text{circ}(\alpha_{15}, \alpha_{13}, \alpha_{14}), \\ C'_3 &= \text{circ}(\alpha_{18}, \alpha_{16}, \alpha_{17}), D_1 = \text{circ}(\alpha_{10}, \alpha_{18}, \alpha_{17}), D_2 = \text{circ}(\alpha_{16}, \alpha_{15}, \alpha_{14}), \\ D_3 &= \text{circ}(\alpha_{13}, \alpha_{12}, \alpha_{11}), D'_2 = \text{circ}(\alpha_{14}, \alpha_{16}, \alpha_{15}), D'_3 = \text{circ}(\alpha_{11}, \alpha_{13}, \alpha_{12}), \\ E_1 &= \text{circ}(\alpha_1, \alpha_9, \alpha_8), E_2 = \text{circ}(\alpha_7, \alpha_6, \alpha_4), E_3 = \text{circ}(\alpha_3, \alpha_2, \alpha_1), \\ E'_2 &= \text{circ}(\alpha_4, \alpha_7, \alpha_6) \text{ and } E'_3 = \text{circ}(\alpha_1, \alpha_3, \alpha_2).\end{aligned}$$

2. Let  $G = \langle x, y \mid x^6 = y^3 = 1, xy = yx \rangle \cong C_3 \times C_6$ . Let  $v_2 = \sum_{i=0}^5 \sum_{j=0}^2 \alpha_{1+i+6j} x^i y^j \in R(C_3 \times C_6)$ . Also, let  $H = \langle a, b \mid a^3 = b^2 = 1, a^b = a^{-1} \rangle \cong D_6$ . We now define the composite matrix over  $R$  as

$$\Omega(v_2) = \begin{pmatrix} A'_1 & A'_2 & A'_3 \\ A'_4 & A'_5 & A'_6 \\ A'_7 & A'_8 & A'_9 \end{pmatrix}$$

where

$$A'_1 = \begin{pmatrix} \alpha_{g_1^{-1}g_1} & \alpha_{g_1^{-1}g_2} & \cdots & \alpha_{g_1^{-1}g_6} \\ \alpha_{\phi_1(h_2^{-1}h_1)} & \alpha_{\phi_1(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_1(h_2^{-1}h_6)} \\ \alpha_{\phi_1(h_3^{-1}h_1)} & \alpha_{\phi_1(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_1(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_1(h_6^{-1}h_1)} & \alpha_{\phi_1(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_1(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_1 : h_j^{-1} h_i \xrightarrow{\phi_1} g_1^{-1} g_i$$

for when

$$j = 2 : i = 1, 2, \dots, 6$$

$$j = 3 : i = 1, 2, \dots, 6$$

$\vdots$

$$j = 6 : i = 1, 2, \dots, 6,$$

$$A'_2 = \begin{pmatrix} \alpha_{g_1^{-1} g_7} & \alpha_{g_1^{-1} g_8} & \cdots & \alpha_{g_1^{-1} g_{12}} \\ \alpha_{\phi_2(h_2^{-1} h_1)} & \alpha_{\phi_2(h_2^{-1} h_2)} & \cdots & \alpha_{\phi_2(h_2^{-1} h_6)} \\ \alpha_{\phi_2(h_3^{-1} h_1)} & \alpha_{\phi_2(h_3^{-1} h_2)} & \cdots & \alpha_{\phi_2(h_3^{-1} h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_2(h_6^{-1} h_1)} & \alpha_{\phi_2(h_6^{-1} h_2)} & \cdots & \alpha_{\phi_2(h_6^{-1} h_6)} \end{pmatrix}$$

with

$$\phi_2 : h_j^{-1} h_k \xrightarrow{\phi_2} g_1^{-1} g_i$$

for when

$$j = 2 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$j = 3 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$\vdots$

$$j = 6 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$A'_3 = \begin{pmatrix} \alpha_{g_1^{-1} g_{13}} & \alpha_{g_1^{-1} g_{14}} & \cdots & \alpha_{g_1^{-1} g_{18}} \\ \alpha_{\phi_3(h_2^{-1} h_1)} & \alpha_{\phi_3(h_2^{-1} h_2)} & \cdots & \alpha_{\phi_3(h_2^{-1} h_6)} \\ \alpha_{\phi_3(h_3^{-1} h_1)} & \alpha_{\phi_3(h_3^{-1} h_2)} & \cdots & \alpha_{\phi_3(h_3^{-1} h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_3(h_6^{-1} h_1)} & \alpha_{\phi_3(h_6^{-1} h_2)} & \cdots & \alpha_{\phi_3(h_6^{-1} h_6)} \end{pmatrix}$$

with

$$\phi_3 : h_j^{-1} h_k \xrightarrow{\phi_3} g_1^{-1} g_i$$

for when

$$\begin{aligned}
j = 2 : & \quad k = 1, i = 13; \quad k = 2, i = 14; \quad \dots \quad k = 6, i = 18, \\
j = 3 : & \quad k = 1, i = 13; \quad k = 2, i = 14; \quad \dots \quad k = 6, i = 18, \\
& \quad \vdots \\
j = 6 : & \quad k = 1, i = 13; \quad k = 2, i = 14; \quad \dots \quad k = 6, i = 18,
\end{aligned}$$

$$A'_4 = \begin{pmatrix} \alpha_{g_7^{-1}g_1} & \alpha_{g_7^{-1}g_2} & \dots & \alpha_{g_7^{-1}g_6} \\ \alpha_{\phi_4(h_2^{-1}h_1)} & \alpha_{\phi_4(h_2^{-1}h_2)} & \dots & \alpha_{\phi_4(h_2^{-1}h_6)} \\ \alpha_{\phi_4(h_3^{-1}h_1)} & \alpha_{\phi_4(h_3^{-1}h_2)} & \dots & \alpha_{\phi_4(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_4(h_6^{-1}h_1)} & \alpha_{\phi_4(h_6^{-1}h_2)} & \dots & \alpha_{\phi_4(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_4 : h_j^{-1}h_i \xrightarrow{\phi_4} g_7^{-1}g_i$$

for when

$$j = 2 : \quad i = 1, 2, \dots, 6$$

$$j = 3 : \quad i = 1, 2, \dots, 6$$

$\vdots$

$$j = 6 : \quad i = 1, 2, \dots, 6,$$

$$A'_5 = \begin{pmatrix} \alpha_{g_7^{-1}g_7} & \alpha_{g_7^{-1}g_8} & \dots & \alpha_{g_7^{-1}g_{12}} \\ \alpha_{\phi_5(h_2^{-1}h_1)} & \alpha_{\phi_5(h_2^{-1}h_2)} & \dots & \alpha_{\phi_5(h_2^{-1}h_6)} \\ \alpha_{\phi_5(h_3^{-1}h_1)} & \alpha_{\phi_5(h_3^{-1}h_2)} & \dots & \alpha_{\phi_5(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_5(h_6^{-1}h_1)} & \alpha_{\phi_5(h_6^{-1}h_2)} & \dots & \alpha_{\phi_5(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_5 : h_j^{-1}h_k \xrightarrow{\phi_5} g_7^{-1}g_i$$

for when

$$j = 2 : \quad k = 1, i = 7; \quad k = 2, i = 8; \quad \dots \quad k = 6, i = 12,$$

$$j = 3 : \quad k = 1, i = 7; \quad k = 2, i = 8; \quad \dots \quad k = 6, i = 12,$$

⋮

$$j = 6 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$A'_6 = \begin{pmatrix} \alpha_{g_7^{-1}g_{13}} & \alpha_{g_7^{-1}g_{14}} & \dots & \alpha_{g_7^{-1}g_{18}} \\ \alpha_{\phi_6(h_2^{-1}h_1)} & \alpha_{\phi_6(h_2^{-1}h_2)} & \dots & \alpha_{\phi_6(h_2^{-1}h_6)} \\ \alpha_{\phi_6(h_3^{-1}h_1)} & \alpha_{\phi_6(h_3^{-1}h_2)} & \dots & \alpha_{\phi_6(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_6(h_6^{-1}h_1)} & \alpha_{\phi_6(h_6^{-1}h_2)} & \dots & \alpha_{\phi_6(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_6 : h_j^{-1}h_k \xrightarrow{\phi_6} g_7^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$$j = 3 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

⋮

$$j = 6 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$$A'_7 = \begin{pmatrix} \alpha_{g_{13}^{-1}g_1} & \alpha_{g_{13}^{-1}g_2} & \dots & \alpha_{g_{13}^{-1}g_6} \\ \alpha_{\phi_7(h_2^{-1}h_1)} & \alpha_{\phi_7(h_2^{-1}h_2)} & \dots & \alpha_{\phi_7(h_2^{-1}h_6)} \\ \alpha_{\phi_7(h_3^{-1}h_1)} & \alpha_{\phi_7(h_3^{-1}h_2)} & \dots & \alpha_{\phi_7(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_7(h_6^{-1}h_1)} & \alpha_{\phi_7(h_6^{-1}h_2)} & \dots & \alpha_{\phi_7(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_7 : h_j^{-1}h_i \xrightarrow{\phi_7} g_{13}^{-1}g_i$$

for when

$$j = 2 : i = 1, 2, \dots, 6$$

$$j = 3 : i = 1, 2, \dots, 6$$

⋮

$$j = 6 : i = 1, 2, \dots, 6,$$

$$A'_8 = \begin{pmatrix} \alpha_{g_{13}^{-1}g_7} & \alpha_{g_{13}^{-1}g_8} & \cdots & \alpha_{g_{13}^{-1}g_{12}} \\ \alpha_{\phi_8(h_2^{-1}h_1)} & \alpha_{\phi_8(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_8(h_2^{-1}h_6)} \\ \alpha_{\phi_8(h_3^{-1}h_1)} & \alpha_{\phi_8(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_8(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_8(h_6^{-1}h_1)} & \alpha_{\phi_8(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_8(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_8 : h_j^{-1}h_k \xrightarrow{\phi_8} g_{13}^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$j = 3 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$\vdots$

$$j = 6 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

and

$$A'_9 = \begin{pmatrix} \alpha_{g_{13}^{-1}g_{13}} & \alpha_{g_{13}^{-1}g_{14}} & \cdots & \alpha_{g_{13}^{-1}g_{18}} \\ \alpha_{\phi_9(h_2^{-1}h_1)} & \alpha_{\phi_9(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_9(h_2^{-1}h_6)} \\ \alpha_{\phi_9(h_3^{-1}h_1)} & \alpha_{\phi_9(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_9(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_9(h_6^{-1}h_1)} & \alpha_{\phi_9(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_9(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_9 : h_j^{-1}h_k \xrightarrow{\phi_9} g_{13}^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$$j = 3 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$\vdots$

$$j = 6 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18.$$

This results in a composite matrix over  $R$  of the following form:

$$\Omega(v_2) = \begin{pmatrix} A'_1 & A'_2 & A'_3 \\ A'_4 & A'_5 & A'_6 \\ A'_7 & A'_8 & A'_9 \end{pmatrix} = \begin{pmatrix} B & C & D \\ D & B & C \\ C & D & B \end{pmatrix} =$$

$$= \begin{pmatrix} B_1 & B_2 & C_1 & C_2 & D_1 & D_2 \\ B_2^T & B_1^T & C_2^T & C_1^T & D_2^T & D_1^T \\ D_1 & D_2 & B_1 & B_2 & C_1 & C_2 \\ D_2^T & D_1^T & B_2^T & B_1^T & C_2^T & C_1^T \\ C_1 & C_2 & D_1 & D_2 & B_1 & B_2 \\ C_2^T & C_1^T & D_2^T & D_1^T & B_2^T & B_1^T \end{pmatrix},$$

where

$$B_1 = \text{circ}(\alpha_1, \alpha_2, \alpha_3), B_2 = \text{circ}(\alpha_4, \alpha_5, \alpha_6), C_1 = \text{circ}(\alpha_7, \alpha_8, \alpha_9),$$

$$C_2 = \text{circ}(\alpha_{10}, \alpha_{11}, \alpha_{12}), D_1 = \text{circ}(\alpha_{13}, \alpha_{14}, \alpha_{15}), D_2 = \text{circ}(\alpha_{16}, \alpha_{17}, \alpha_{18}).$$

3. Let  $G = \langle x, y \mid x^6 = y^3 = 1, xy = yx \rangle \cong C_3 \times C_6$ . Let  $v_3 = \sum_{i=0}^5 \sum_{j=0}^2 \alpha_{1+i+6j} x^i y^j \in R(C_3 \times C_6)$ . Also, let  $H = \langle a \mid a^6 = 1 \rangle \cong C_{3,2}$ . We now define the composite matrix over  $R$  as

$$\Omega(v_3) = \begin{pmatrix} A'_1 & A'_2 & A'_3 \\ A'_4 & A'_5 & A'_6 \\ A'_7 & A'_8 & A'_9 \end{pmatrix}$$

where

$$A'_1 = \begin{pmatrix} \alpha_{g_1^{-1}g_1} & \alpha_{g_1^{-1}g_2} & \cdots & \alpha_{g_1^{-1}g_6} \\ \alpha_{\phi_1(h_2^{-1}h_1)} & \alpha_{\phi_1(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_1(h_2^{-1}h_6)} \\ \alpha_{\phi_1(h_3^{-1}h_1)} & \alpha_{\phi_1(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_1(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_1(h_6^{-1}h_1)} & \alpha_{\phi_1(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_1(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_1 : h_j^{-1}h_i \xrightarrow{\phi_1} g_1^{-1}g_i$$

for when

$$j = 2 : i = 1, 2, \dots, 6$$

$$j = 3 : i = 1, 2, \dots, 6$$

$\vdots$

$$j = 6 : i = 1, 2, \dots, 6,$$

$$A'_2 = \begin{pmatrix} \alpha_{g_1^{-1}g_7} & \alpha_{g_1^{-1}g_8} & \cdots & \alpha_{g_1^{-1}g_{12}} \\ \alpha_{\phi_2(h_2^{-1}h_1)} & \alpha_{\phi_2(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_2(h_2^{-1}h_6)} \\ \alpha_{\phi_2(h_3^{-1}h_1)} & \alpha_{\phi_2(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_2(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_2(h_6^{-1}h_1)} & \alpha_{\phi_2(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_2(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_2 : h_j^{-1}h_k \xrightarrow{\phi_2} g_1^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$j = 3 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$\vdots$

$$j = 6 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$A'_3 = \begin{pmatrix} \alpha_{g_1^{-1}g_{13}} & \alpha_{g_1^{-1}g_{14}} & \cdots & \alpha_{g_1^{-1}g_{18}} \\ \alpha_{\phi_3(h_2^{-1}h_1)} & \alpha_{\phi_3(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_3(h_2^{-1}h_6)} \\ \alpha_{\phi_3(h_3^{-1}h_1)} & \alpha_{\phi_3(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_3(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_3(h_6^{-1}h_1)} & \alpha_{\phi_3(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_3(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_3 : h_j^{-1}h_k \xrightarrow{\phi_3} g_1^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$$j = 3 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$\vdots$

$$j = 6 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$



$$A'_4 = \begin{pmatrix} \alpha_{g_7^{-1}g_1} & \alpha_{g_7^{-1}g_2} & \cdots & \alpha_{g_7^{-1}g_6} \\ \alpha_{\phi_4(h_2^{-1}h_1)} & \alpha_{\phi_4(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_4(h_2^{-1}h_6)} \\ \alpha_{\phi_4(h_3^{-1}h_1)} & \alpha_{\phi_4(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_4(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_4(h_6^{-1}h_1)} & \alpha_{\phi_4(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_4(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_4 : h_j^{-1}h_i \xrightarrow{\phi_4} g_7^{-1}g_i$$

for when

$$j = 2 : i = 1, 2, \dots, 6$$

$$j = 3 : i = 1, 2, \dots, 6$$

$\vdots$

$$j = 6 : i = 1, 2, \dots, 6,$$

$$A'_5 = \begin{pmatrix} \alpha_{g_7^{-1}g_7} & \alpha_{g_7^{-1}g_8} & \cdots & \alpha_{g_7^{-1}g_{12}} \\ \alpha_{\phi_5(h_2^{-1}h_1)} & \alpha_{\phi_5(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_5(h_2^{-1}h_6)} \\ \alpha_{\phi_5(h_3^{-1}h_1)} & \alpha_{\phi_5(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_5(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_5(h_6^{-1}h_1)} & \alpha_{\phi_5(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_5(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_5 : h_j^{-1}h_k \xrightarrow{\phi_5} g_7^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$j = 3 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$\vdots$

$$j = 6 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$A'_6 = \begin{pmatrix} \alpha_{g_7^{-1}g_{13}} & \alpha_{g_7^{-1}g_{14}} & \cdots & \alpha_{g_7^{-1}g_{18}} \\ \alpha_{\phi_6(h_2^{-1}h_1)} & \alpha_{\phi_6(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_6(h_2^{-1}h_6)} \\ \alpha_{\phi_6(h_3^{-1}h_1)} & \alpha_{\phi_6(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_6(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_6(h_6^{-1}h_1)} & \alpha_{\phi_6(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_6(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_6 : h_j^{-1}h_k \xrightarrow{\phi_6} g_7^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$$j = 3 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$\vdots$

$$j = 6 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$$A'_7 = \begin{pmatrix} \alpha_{g_{13}^{-1}g_1} & \alpha_{g_{13}^{-1}g_2} & \cdots & \alpha_{g_{13}^{-1}g_6} \\ \alpha_{\phi_7(h_2^{-1}h_1)} & \alpha_{\phi_7(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_7(h_2^{-1}h_6)} \\ \alpha_{\phi_7(h_3^{-1}h_1)} & \alpha_{\phi_7(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_7(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_7(h_6^{-1}h_1)} & \alpha_{\phi_7(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_7(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_7 : h_j^{-1}h_i \xrightarrow{\phi_7} g_{13}^{-1}g_i$$

for when

$$j = 2 : i = 1, 2, \dots, 6$$

$$j = 3 : i = 1, 2, \dots, 6$$

$\vdots$

$$j = 6 : i = 1, 2, \dots, 6,$$

$$A'_8 = \begin{pmatrix} \alpha_{g_{13}^{-1}g_7} & \alpha_{g_{13}^{-1}g_8} & \cdots & \alpha_{g_{13}^{-1}g_{12}} \\ \alpha_{\phi_8(h_2^{-1}h_1)} & \alpha_{\phi_8(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_8(h_2^{-1}h_6)} \\ \alpha_{\phi_8(h_3^{-1}h_1)} & \alpha_{\phi_8(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_8(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_8(h_6^{-1}h_1)} & \alpha_{\phi_8(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_8(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_8 : h_j^{-1}h_k \xrightarrow{\phi_8} g_{13}^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$$j = 3 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

$\vdots$

$$j = 6 : k = 1, i = 7; k = 2, i = 8; \dots k = 6, i = 12,$$

and

$$A'_9 = \begin{pmatrix} \alpha_{g_{13}^{-1}g_{13}} & \alpha_{g_{13}^{-1}g_{14}} & \cdots & \alpha_{g_{13}^{-1}g_{18}} \\ \alpha_{\phi_9(h_2^{-1}h_1)} & \alpha_{\phi_9(h_2^{-1}h_2)} & \cdots & \alpha_{\phi_9(h_2^{-1}h_6)} \\ \alpha_{\phi_9(h_3^{-1}h_1)} & \alpha_{\phi_9(h_3^{-1}h_2)} & \cdots & \alpha_{\phi_9(h_3^{-1}h_6)} \\ \vdots & \vdots & \vdots & \vdots \\ \alpha_{\phi_9(h_6^{-1}h_1)} & \alpha_{\phi_9(h_6^{-1}h_2)} & \cdots & \alpha_{\phi_9(h_6^{-1}h_6)} \end{pmatrix}$$

with

$$\phi_9 : h_j^{-1}h_k \xrightarrow{\phi_9} g_{13}^{-1}g_i$$

for when

$$j = 2 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$$j = 3 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18,$$

$\vdots$

$$j = 6 : k = 1, i = 13; k = 2, i = 14; \dots k = 6, i = 18.$$

This results in a composite matrix over  $R$  of the following form:

$$\Omega(v_3) = \begin{pmatrix} A'_1 & A'_2 & A'_3 \\ A'_4 & A'_5 & A'_6 \\ A'_7 & A'_8 & A'_9 \end{pmatrix} = \begin{pmatrix} B & C & D \\ D & B & C \\ C & D & B \end{pmatrix} =$$

$$= \begin{pmatrix} B_1 & B_2 & C_1 & C_2 & D_1 & D_2 \\ B'_2 & B_1 & C'_2 & C_1 & D'_2 & D_1 \\ D_1 & D_2 & B_1 & B_2 & C_1 & C_2 \\ D'_2 & D_1 & B'_2 & B_1 & C'_2 & C_1 \\ C_1 & C_2 & D_1 & D_2 & B_1 & B_2 \\ C'_2 & C_1 & D'_2 & D_1 & B'_2 & B_1 \end{pmatrix},$$

where

$$\begin{aligned} B_1 &= \text{circ}(\alpha_1, \alpha_2, \alpha_3), B_2 = \text{circ}(\alpha_4, \alpha_5, \alpha_6), B'_2 = \text{circ}(\alpha_6, \alpha_4, \alpha_5), \\ C_1 &= \text{circ}(\alpha_7, \alpha_8, \alpha_9), C_2 = \text{circ}(\alpha_{10}, \alpha_{11}, \alpha_{12}), C'_2 = \text{circ}(\alpha_{12}, \alpha_{10}, \alpha_{11}), \\ D_1 &= \text{circ}(\alpha_{13}, \alpha_{14}, \alpha_{15}), D_2 = \text{circ}(\alpha_{16}, \alpha_{17}, \alpha_{18}), D'_2 = \text{circ}(\alpha_{18}, \alpha_{16}, \alpha_{17}). \end{aligned}$$

## 4 Generator Matrices

In this section, we consider generator matrices of the form  $[I \mid \Omega(v_i)]$ , where  $I$  is the identity matrix and  $\Omega(v_i)$  with  $i = \{1, 2, 3\}$  are the composite matrices from the previous section. For each, we show under what conditions such generator matrix produces self-dual codes over the ring  $R$ . We assume that the ring  $R$  has characteristic 2.

**Theorem 4.1.** *The generator matrix*

$$[I \mid \Omega(v_1)] \tag{4.1}$$

where  $\Omega(v_1)$  is the composite matrix defined in previous section, generates a self-dual code over the ring  $R$  if and only if the following hold in  $R$ :

$$BB^T + CC^T = I_9, \tag{4.2}$$

$$BD^T + CE^T = \mathbf{0}, \tag{4.3}$$

$$DB^T + EC^T = \mathbf{0}, \tag{4.4}$$

$$DD^T + EE^T = I_9. \tag{4.5}$$

*Proof.* Follows from the standard proof that  $(I_m \mid A)$  generates a self-dual of length  $2m$  if and only if  $AA^T = I_m$ .  $\square$

**Theorem 4.2.** *The generator matrix*

$$[I \mid \Omega(v_2)] \tag{4.6}$$

where  $\Omega(v_2)$  is the composite matrix defined in previous section, generates a self-dual code over the ring  $R$  if and only if the following hold in  $R$  :

$$BB^T + CC^T + DD^T = I_6, \tag{4.7}$$

$$BD^T + CB^T + DC^T = \mathbf{0}, \tag{4.8}$$

$$BC^T + CD^T + DB^T = \mathbf{0}. \tag{4.9}$$

*Proof.* Follows from the standard proof that  $(I_m \mid A)$  generates a self-dual of length  $2m$  if and only if  $AA^T = I_m$ .  $\square$

**Theorem 4.3.** *The generator matrix*

$$[I \mid \Omega(v_3)] \tag{4.10}$$

where  $\Omega(v_3)$  is the composite matrix defined in previous section, generates a self-dual code over the ring  $R$  if and only if the following hold in  $R$  :

$$BB^T + CC^T + DD^T = I_6, \tag{4.11}$$

$$BD^T + CB^T + DC^T = \mathbf{0}, \tag{4.12}$$

$$BC^T + CD^T + DB^T = \mathbf{0}. \tag{4.13}$$

*Proof.* Follows from the standard proof that  $(I_m \mid A)$  generates a self-dual of length  $2m$  if and only if  $AA^T = I_m$ .  $\square$

## 5 New Type I Binary Self-Dual Codes of length 72

In this section, we search for binary self-dual codes by employing the generator matrices defined in Section 4. In particular, we search for binary self-dual codes with parameters  $[36, 18, 6$  or  $8]$  which we then lift to the ring  $\mathbb{F}_2 + u\mathbb{F}_2$  to obtain self-dual codes of length 36 whose binary images are self-dual codes with parameters  $[72, 36, 12]$ .

The possible weight enumerators for Type I [72, 36, 12] codes are as follows ([9]):

$$W_{72,1} = 1 + 2\beta y^{12} + (8640 - 64\gamma)y^{14} + (124281 - 24\beta + 384\gamma)y^{16} + \dots$$

$$W_{72,2} = 1 + 2\beta y^{12} + (7616 - 64\gamma)y^{14} + (134521 - 24\beta + 384\gamma)y^{16} + \dots$$

where  $\beta$  and  $\gamma$  are parameters.

Many codes for different values of  $\beta$  and  $\gamma$  have been constructed in [2, 3, 9, 10, 11, 12, 14, 16, 18, 19, 20, 21]. For an up-to-date list of all known Type I and Type II binary self-dual codes with parameters [72, 36, 12] please see [15].

All the upcoming computational results were obtained by performing searches in the software package MAGMA ([1]).

1. Here, we employ the generator matrix  $[I \mid \Omega(v_1)]$  to search for binary self-dual codes of length 36. Since the matrix  $\Omega(v_1)$  is fully defined by the first row, we only list the first rows of the matrices  $B$  and  $C$  which we label as  $r_B$  and  $r_C$  respectively. We summarise the results in the table below.

Table 1: Type I [36, 18, 6 – 8] Codes from Theorem 4.1

	Type	$r_B$	$r_C$	$ Aut(C_i) $
$C_1$	[36, 18, 6]	(0, 0, 0, 0, 0, 1, 0, 1, 1)	(1, 0, 1, 1, 1, 0, 1, 0, 1)	$2^2 \cdot 3^2$
$C_2$	[36, 18, 6]	(0, 0, 0, 0, 1, 1, 0, 1, 1)	(1, 0, 0, 1, 0, 1, 1, 1, 0)	$2^2 \cdot 3^2$
$C_3$	[36, 18, 8]	(0, 0, 1, 0, 0, 1, 0, 0, 1)	(1, 0, 0, 1, 1, 0, 1, 1, 1)	$2^2 \cdot 3^2$
$C_4$	[36, 18, 8]	(0, 1, 0, 0, 0, 1, 0, 1, 1)	(1, 0, 0, 1, 0, 0, 1, 1, 1)	$2^2 \cdot 3^2$

We now apply the  $R_1$ -lift to each code from the above table to obtain codes whose binary images are self-dual codes with parameters [72, 36, 12]. We only list codes that have not been known in the literature before.

Table 2: New Type I [72, 36, 12] Codes from  $R_1$ -lift of  $C_1$ 

Type	$r_B$	$r_C$	$\gamma$	$\beta$	$ Aut(\mathcal{C}_i) $	
$\mathcal{C}_1$	$W_{72,1}$	$(u, 0, u, u, u, 1, u, u + 1, 1)$	$(1, 0, 1, 1, u + 1, 0, 1, u, 1)$	0	192	36
$\mathcal{C}_2$	$W_{72,1}$	$(u, 0, 0, u, 0, 1, u, 1, 1)$	$(1, 0, u + 1, 1, u + 1, 0, 1, u, u + 1)$	0	198	36
$\mathcal{C}_3$	$W_{72,1}$	$(u, u, 0, u, u, 1, u, 1, u + 1)$	$(1, u, u + 1, 1, u + 1, 0, 1, 0, u + 1)$	0	336	36
$\mathcal{C}_4$	$W_{72,1}$	$(0, u, 0, 0, 0, 1, 0, 1, u + 1)$	$(1, u, u + 1, 1, u + 1, 0, 1, 0, u + 1)$	18	234	36
$\mathcal{C}_5$	$W_{72,1}$	$(u, u, 0, u, 0, 1, u, u + 1, 1)$	$(1, u, u + 1, 1, 1, u, 1, 0, u + 1)$	18	345	36
$\mathcal{C}_6$	$W_{72,1}$	$(0, u, 0, 0, 0, 1, 0, u + 1, 1)$	$(1, u, 1, 1, u + 1, u, 1, u, 1)$	18	378	36
$\mathcal{C}_7$	$W_{72,1}$	$(u, u, u, u, 0, 1, u, u + 1, u + 1)$	$(1, u, 1, 1, u + 1, 0, 1, 0, 1)$	18	396	36
$\mathcal{C}_8$	$W_{72,1}$	$(0, 0, u, 0, u, 1, 0, 1, u + 1)$	$(1, u, 1, 1, 1, 0, 1, u, 1)$	18	441	36
$\mathcal{C}_9$	$W_{72,1}$	$(u, u, 0, u, 0, 1, u, 1, u + 1)$	$(1, u, 1, 1, 1, 0, 1, u, 1)$	18	453	36

Table 3: New Type I [72, 36, 12] Codes from  $R_1$ -lift of  $C_2$ 

Type	$r_B$	$r_C$	$\gamma$	$\beta$	$ Aut(\mathcal{C}_i) $	
$\mathcal{C}_{10}$	$W_{72,1}$	$(0, u, 0, 0, 1, 1, 0, 1, u + 1)$	$(1, u, u, 1, u, u + 1, 1, 1, 0)$	0	219	36
$\mathcal{C}_{11}$	$W_{72,1}$	$(u, 0, u, u, 1, 1, u, u + 1, u + 1)$	$(1, u, 0, 1, 0, 1, 1, u + 1, u)$	0	345	36
$\mathcal{C}_{12}$	$W_{72,1}$	$(0, 0, u, 0, 1, 1, 0, 1, u + 1)$	$(1, u, 0, 1, 0, 1, 1, 1, 0)$	0	408	36
$\mathcal{C}_{13}$	$W_{72,1}$	$(u, 0, 0, u, 1, u + 1, u, u + 1, u + 1)$	$(1, 0, u, 1, 0, u + 1, 1, u + 1, 0)$	18	261	36
$\mathcal{C}_{14}$	$W_{72,1}$	$(u, u, u, u, 1, 1, u, 1, u + 1)$	$(1, 0, 0, 1, u, 1, 1, u + 1, 0)$	18	270	36
$\mathcal{C}_{15}$	$W_{72,1}$	$(u, 0, u, u, 1, u + 1, u, 1, u + 1)$	$(1, 0, u, 1, u, 1, 1, u + 1, 0)$	18	357	36

Table 4: New Type I [72, 36, 12] Codes from  $R_1$ -lift of  $C_3$ 

Type	$r_B$	$r_C$	$\gamma$	$\beta$	$ Aut(\mathcal{C}_i) $	
$\mathcal{C}_{16}$	$W_{72,1}$	$(u, u, 1, u, 0, 1, u, 0, 1)$	$(1, u, 0, 1, u + 1, u, 1, 1, u + 1)$	0	120	36
$\mathcal{C}_{17}$	$W_{72,1}$	$(u, u, 1, u, 0, 1, u, u, 1)$	$(1, 0, u, 1, 1, 0, 1, 1, u + 1)$	0	282	36
$\mathcal{C}_{18}$	$W_{72,1}$	$(u, u, 1, u, 0, u + 1, u, 0, 1)$	$(1, u, 0, 1, u + 1, u, 1, 1, u + 1)$	0	300	36
$\mathcal{C}_{19}$	$W_{72,1}$	$(u, u, 1, u, 0, u + 1, u, 0, 1)$	$(1, u, u, 1, 1, 0, 1, 1, 1)$	18	336	36
$\mathcal{C}_{20}$	$W_{72,1}$	$(u, 0, 1, u, 0, 1, u, u, 1)$	$(1, 0, 0, 1, 1, 0, 1, u + 1, u + 1)$	36	435	36

Table 5: New Type I [72, 36, 12] Codes from  $R_1$ -lift of  $C_4$ 

Type	$r_B$	$r_C$	$\gamma$	$\beta$	$ Aut(\mathcal{C}_i) $	
$\mathcal{C}_{21}$	$W_{72,1}$	$(0, 1, u, 0, u, 1, 0, u + 1, u + 1)$	$(1, u, u, 1, u, u, 1, u + 1, u + 1)$	0	366	36
$\mathcal{C}_{22}$	$W_{72,1}$	$(u, 1, u, u, u, 1, u, 1, u + 1)$	$(1, u, 0, 1, 0, 0, 1, u + 1, 1)$	0	372	36
$\mathcal{C}_{23}$	$W_{72,1}$	$(0, 1, u, 0, u, 1, 0, u + 1, u + 1)$	$(1, 0, 0, 1, 0, 0, 1, u + 1, u + 1)$	0	384	36
$\mathcal{C}_{24}$	$W_{72,1}$	$(u, 1, u, u, u, 1, u, 1, u + 1)$	$(1, u, u, 1, u, 0, 1, u + 1, 1)$	0	390	36
$\mathcal{C}_{25}$	$W_{72,1}$	$(u, 1, 0, u, 0, 1, u, u + 1, 1)$	$(1, u, 0, 1, 0, 0, 1, 1, u + 1)$	0	399	36
$\mathcal{C}_{26}$	$W_{72,1}$	$(u, 1, u, u, u, 1, u, u + 1, 1)$	$(1, 0, u, 1, 0, 0, 1, u + 1, u + 1)$	18	264	36
$\mathcal{C}_{27}$	$W_{72,1}$	$(u, 1, 0, u, u, u + 1, u, u + 1, 1)$	$(1, 0, u, 1, 0, u, 1, 1, u + 1)$	18	285	36
$\mathcal{C}_{28}$	$W_{72,1}$	$(0, 1, u, 0, 0, u + 1, 0, 1, 1)$	$(1, u, u, 1, u, 0, 1, 1, 1)$	18	300	36

2. Here, we employ the generator matrix  $[I \mid \Omega(v_2)]$  to search for binary self-dual codes of length 36. Since the matrix  $\Omega(v_2)$  is fully defined by the first row, we only list the first rows of the matrices  $B, C$  and  $D$  which we label as  $r_B, r_C$  and  $r_D$  respectively. We summarise the results in the table below

Table 6: Type I [36, 18, 6 – 8] Codes from Theorem 4.2

Type	$r_B$	$r_C$	$r_D$	$ Aut(C_i) $
$C_5$ [36, 18, 6]	(0, 0, 0, 0, 0, 1)	(1, 1, 1, 0, 0, 1)	(1, 1, 1, 0, 1, 0)	$2^5 \cdot 3^4 \cdot 5$
$C_6$ [36, 18, 6]	(0, 0, 0, 0, 1, 1)	(0, 0, 0, 0, 1, 1)	(1, 1, 1, 1, 0, 1)	$2^5 \cdot 3^4 \cdot 5$
$C_7$ [36, 18, 6]	(0, 0, 0, 0, 1, 1)	(0, 1, 1, 0, 1, 1)	(0, 1, 1, 1, 0, 0)	$2^5 \cdot 3^2$
$C_8$ [36, 18, 6]	(0, 0, 1, 0, 0, 1)	(0, 0, 1, 1, 1, 0)	(1, 1, 1, 0, 0, 1)	$2^5 \cdot 3^2$

We now apply the  $R_1$ -lift to each code from the above table to obtain codes whose binary images are self-dual codes with parameters [72, 36, 12]. We only list codes that have not been known in the literature before.

Table 7: New Type I [72, 36, 12] Codes from  $R_1$ -lift of  $C_7$

Type	$r_B$	$r_C$	$r_D$	$\gamma$	$\beta$	$ Aut(C_i) $
$C_{29}$ $W_{72,1}$	(0, 0, 0, $u$ , 1, 1)	( $u$ , 1, $u+1$ , $u$ , 1, 1)	( $u$ , $u+1$ , 1, $u+1$ , 0, 0)	0	471	144

3. Here, we employ the generator matrix  $[I \mid \Omega(v_3)]$  to search for binary self-dual codes of length 36. Since the matrix  $\Omega(v_3)$  is fully defined by the first row, we only list the first rows of the matrices  $B, C$  and  $D$  which we label as  $r_B, r_C$  and  $r_D$  respectively. We summarise the results in the table below

Table 8: Type I [36, 18, 6 – 8] Codes from Theorem 4.2

Type	$r_B$	$r_C$	$r_D$	$ Aut(C_i) $
$C_9$ [36, 18, 6]	(0, 0, 0, 0, 0, 1)	(0, 1, 1, 0, 1, 1)	(1, 0, 1, 1, 0, 1)	$2^5 \cdot 3^2$
$C_{10}$ [36, 18, 6]	(0, 0, 0, 0, 0, 1)	(1, 1, 1, 0, 0, 1)	(1, 1, 1, 0, 1, 0)	$2^5 \cdot 3^4 \cdot 5$
$C_{11}$ [36, 18, 6]	(0, 0, 0, 0, 1, 1)	(0, 0, 0, 0, 1, 1)	(1, 1, 1, 1, 0, 1)	$2^5 \cdot 3^4 \cdot 5$
$C_{12}$ [36, 18, 6]	(0, 0, 0, 0, 1, 1)	(0, 1, 1, 0, 0, 1)	(1, 1, 0, 1, 0, 1)	$2^5 \cdot 3^2$

We now apply the  $R_1$ -lift to each code from the above table to obtain codes whose binary images are self-dual codes with parameters



[72, 36, 12]. We only list codes that have not been known in the literature before.

Table 9: New Type I [72, 36, 12] Codes from  $R_1$ -lift of  $C_9$

	Type	$r_B$	$r_C$	$r_D$	$\gamma$	$\beta$	$ Aut(\mathcal{C}_i) $
$\mathcal{C}_{30}$	$W_{72,1}$	$(0, u, u, u, u, 1)$	$(u, 1, 1, u, u + 1, 1)$	$(u + 1, u, u + 1, 1, u, u + 1)$	0	621	432

## 6 Conclusion

In this paper, we presented three composite matrices of order 18 derived from group rings. We formed three generator matrices which consist of the composite matrices and searched for binary self-dual codes with parameters [36, 18, 6 or 8]. We then lifted the codes over the ring  $R_1 = \mathbb{F}_2 + u\mathbb{F}_2$  to obtain codes whose binary images are self-dual codes with parameters [72, 36, 12]. We were able to construct 30 Type I binary [72, 36, 12] self-dual codes with new weight enumerators in  $W_{72,1}$ :

$$\begin{aligned}
 &(\gamma = 0, \beta = \{120, 192, 198, 219, 282, 300, 336, 345, 366, 372, 384, 390, 399, 408, \\
 &\quad 471, 621\}), \\
 &(\gamma = 18, \beta = \{234, 261, 264, 270, 285, 300, 336, 345, , 357, 378, 396, 441, 453\}), \\
 &(\gamma = 36, \beta = \{435\}).
 \end{aligned}$$

A suggestion for future work is to consider composite constructions of higher orders than 18 and to form generator matrices that can be used to search for optimal binary self-dual codes of different lengths.

## References

- [1] W. Bosma, J. Cannon and C. Playoust, “The Magma algebra system. I. The user language”, J. Symbolic Comput., vol. 24, pp. 235–265, 1997.
- [2] I. Bouyukliev, V. Fack and J. Winna, “Hadamard matrices of order 36”, European Conference on Combinatorics, Graph Theory and Applications, pp. 93–98, 2005.
- [3] R. Dontcheva, “New binary self-dual [70, 35, 12] and binary [72, 36, 12] self-dual doubly-even codes”, Serdica Math. J., vol. 27, pp. 287–302, 2002.

- [4] S.T. Dougherty, P. Gaborit, M. Harada, P. Sole, “Type II codes over  $\mathbb{F}_2 + u\mathbb{F}_2$ ”, IEEE Trans. Inform. Theory, vol. 45, pp. 32–45, 1999.
- [5] S. T. Dougherty, J. Gildea and A. Korban, “Extending an Established Isomorphism between Group Rings and a Subring of the  $n \times n$  Matrices”, **In press.**
- [6] S.T. Dougherty, J. Gildea, A. Korban and A. Kaya, “Composite Constructions of Self-Dual Codes from Group Rings and New Extremal Self-Dual Binary Codes of length 68”, Advances in Mathematics of Communications, vol. 14, pp. 677–702, 2020.
- [7] S.T. Dougherty, J. Gildea, A. Korban and A. Kaya, “New Extremal Self-Dual Binary Codes of length 68 via Composite Construction,  $\mathbb{F}_2 + u\mathbb{F}_2$  Lifts, Extensions and Neighbours”, International Journal of Information and Coding Theory (IJICOT), vol. 5, no. 3/4, 2020.
- [8] S.T. Dougherty, J. Gildea, R. Taylor and A. Tylshchak, “Group Rings, G-Codes and Constructions of Self-Dual and Formally Self-Dual Codes”, Des., Codes and Cryptog., Designs, vol. 86, no. 9, pp. 2115–2138, 2018.
- [9] S.T. Dougherty, T.A. Gulliver, M. Harada, “Extremal binary self dual codes”, IEEE Trans. Inform. Theory, vol. 43, no. 6, pp. 2036–2047, 1997.
- [10] S.T. Dougherty, J-L. Kim and P. Sole, “Double circulant codes from two class association schemes”, Advances in Mathematics of Communications, vol. 1, no. 1, pp. 45–64, 2007.
- [11] T.A. Gulliver, M. Harada, “On double circulant doubly-even self-dual  $[72, 36, 12]$  codes and their neighbors”, Australas. J. Comb., vol. 40, pp. 137-144, 2008.
- [12] M. Gurel, N. Yankov, “Self-dual codes with an automorphism of order 17”, Mathematical Communications, vol. 21, no. 1, pp. 97–101, 2016.
- [13] T. Hurley, “Group Rings and Rings of Matrices”, Int. Jour. Pure and Appl. Math, vol. 31, no. 3, pp. 319–335, 2006.
- [14] A. Kaya, B. Yildiz and I. Siap, “New extremal binary self-dual codes of length 68 from quadratic residue codes over  $\mathbb{F}_2 + u\mathbb{F}_2 + u^2\mathbb{F}_2$ ”, Finite Fields and Their Applications, vol. 29, pp. 160–177, 2014.

- [15] A. Korban, All known Type I and Type II [72, 36, 12] binary self-dual codes, available online at <https://sites.google.com/view/adriankorban/binary-self-dual-codes>.
- [16] A. Korban, S. Sahinkaya, D. Ustun, “A Novel Genetic Search Scheme Based on Nature – Inspired Evolutionary Algorithms for Self-Dual Codes”, arXiv:2012.12248.
- [17] E.M. Rains, “Shadow Bounds for Self-Dual Codes”, IEEE Trans. Inf. Theory, vol. 44, pp. 134–139, 1998.
- [18] N. Tufekci, B. Yildiz, “On codes over  $R_{k,m}$  and constructions for new binary self-dual codes”, Mathematica Slovaca, vol. 66, no. 6, pp. 1511–1526, 2016.
- [19] N. Yankov, M.H. Lee, M. Gurel and M. Ivanova, “Self-dual codes with an automorphism of order 11”, IEEE, Trans. Inform. Theory, vol. 61, pp. 1188–1193, 2015.
- [20] A. Zhdanov, “New self-dual codes of length 72”, arXiv:1705.05779.
- [21] A. Zhdanov, “Convolutional encoding of 60, 64, 68, 72-bit self-dual codes”, arXiv:1702.05153.