

INTERTWINING PERIODS AND DISTINCTION FOR P-ADIC GALOIS SYMMETRIC PAIRS

NADIR MATRINGE AND OMER OFFEN

ABSTRACT. We consider distinction of representations in the context of p -adic Galois symmetric spaces. We provide new sufficient conditions for distinction of parabolically induced representations in terms of similar conditions on the inducing data and deduce a characterization for distinction of representations parabolically induced from cuspidal. We explicate the results further for classical groups and give several applications, in particular, concerning the preservation of distinction via Langlands functoriality. We relate our results with a conjecture of Dipendra Prasad.

CONTENTS

1. Introduction	2
Acknowledgement	5
2. Notation and preliminaries	5
2.1. Parabolic subgroups and Iwasawa decomposition	5
2.2. Roots and Weyl groups	5
2.3. Symmetric spaces	7
2.4. Parabolic induction and standard intertwining operators	7
2.5. Meromorphic families of linear forms	9
2.6. The result of Blanc and Delorme	9
3. Intertwining periods	10
3.1. An informal introduction	10
3.2. The strategy	10
3.3. The set up	11
3.4. A graph of involutions	11
3.5. Minimal vertices	11
3.6. The cone of convergence	12
3.7. Admissible orbits	12
3.8. The intertwining periods modulo convergence	13
3.9. Intertwining periods for minimal vertices	13
3.10. Convergence in the unramified case	13
3.11. The main result	15
3.12. Applications to distinction	16
4. Classical Galois pairs	18

Date: September 12, 2021.

4.1.	Further notation and preliminaries	18
4.2.	Classical Galois pairs-The set up	20
4.3.	A classification of orbits	21
5.	Distinction for classical Galois pairs-statement of the main result	26
5.1.	The standard parabolic subgroups of G°	26
5.2.	Explication of Corollary 2 for classical symmetric pairs	27
5.3.	A necessary condition for distinction	28
6.	Distinction for classical Galois pairs-proof of the main result	29
6.1.	A reduction step	29
6.2.	Admissible orbits	29
6.3.	The signed permutation group	29
6.4.	The split even orthogonal case	30
6.5.	The set $W_G(M)$ and elementary symmetries.	30
6.6.	A choice of representatives in G for $W_G(M)$	31
6.7.	Orbits and stabilizers in $X \cap N_{G^\circ}(M^\circ)$	32
6.8.	The contribution of an admissible orbit	37
6.9.	Completion of the proof of Theorem 2	37
7.	Distinction and conjugate-selfduality in the split odd orthogonal case	38
8.	On distinction and local Langlands correspondence	41
8.1.	On Galois distinction for GL_N	41
8.2.	L-groups and transfer maps	42
8.3.	Transfer of induced generic representations	44
8.4.	A consequence of a conjecture of Prasad	45
8.5.	Transfer of distinction for principal series	48
Appendix A.	Explicit computations of Prasad's quadratic character and opposition group	55
A.1.	Prasad's quadratic character	55
A.2.	Prasad's global character	64
A.3.	The opposition group	64
References		65

1. INTRODUCTION

Let E/F be a quadratic extension of p -adic fields with Galois involution θ , \mathbb{H} a connected reductive group defined over F , and $\mathbb{G} = \mathrm{Res}_{E/F}(\mathbb{H})$. Set $G = \mathbb{G}(F)$ and $H = \mathbb{H}(F)$. In this paper we provide a step in reducing the study of H -distinction of irreducible representations of G to that of cuspidal representations.

Recall that a smooth representation π of G is called H -distinguished if it admits a nonzero H -invariant linear form. Also, any irreducible representation π of G is the quotient of an induced representation of the form $I_P^G(\sigma)$ for σ a cuspidal representation of a Levi part of the parabolic subgroup P of G . In particular if π is H -distinguished then so is $I_P^G(\sigma)$. One

of our main results is a necessary and sufficient condition on σ for $I_P^G(\sigma)$ to be distinguished, it is the content of Corollary 2.

Of course one is far from understanding when π is distinguished from this result, and to go further one needs to understand when a nonzero H -invariant linear form L on $I_P^G(\sigma)$ descends to π .

One of the motivations of the present work is the paper [Pra] that makes precise predictions on the Langlands parameter of an irreducible representation of G that is H -distinguished, and on the multiplicity of H -invariant linear forms on π in terms of such a parameter. In fact if we want to state results in terms of Langlands parameters, one has to go via more steps. More precisely, π is the unique irreducible quotient of a standard module $I_P^G(\tau)$ thanks to the Langlands quotient theorem. Here τ is an essentially tempered representation and therefore a summand of a representation induced from an essentially discrete series representation. Finally, one studies distinction of discrete series representations via distinction of their cuspidal support. At each step one would like a necessary and sufficient condition on the inducing datum of the induced representation considered to be H -distinguished, and an understanding of when an H -invariant linear form on the induced representation descends to a given irreducible quotient.

Necessary conditions for distinction of an induced representation of G in terms of distinction of the inducing datum are usually obtained by means of the Bernstein-Zelevinsky geometric lemma, and this technique is nowadays very standard. In fact in the case of representations induced from a cuspidal representation the necessary condition obtained from the geometric lemma becomes sufficient, as we show in Corollary 2.

To show that the necessary condition given by the geometric lemma is sufficient is not as straightforward and we make use of local intertwining periods which in fact allow us to obtain sufficient conditions for induced representations to be distinguished in a more general context, see Proposition 3. This can be seen as the most basic application of intertwining periods to the theory. More generally to understand when a nonzero H -invariant linear form on an induced representation descends to an irreducible quotient is a notoriously difficult problem in the field. One way to tackle it is to study local intertwining periods on induced representations and their functional equation (see Remark 3). We expect our result to have further applications concerning distinction of generic discrete series of classical groups.

Intertwining periods were introduced in the global setting in [Jac95] and further studied in [JLR99] and [LR03]. Informally, suppose for a moment that F is a number field with ring of adèles \mathbb{A} . The global intertwining periods are meromorphic families of $\mathbb{H}(\mathbb{A})$ -invariant linear forms on spaces obtained by parabolic induction from automorphic representations of Levi subgroups of $\mathbb{G}(\mathbb{A})$. The linear forms are defined by (the meromorphic continuation of) certain integrals. They arise naturally in the study of period integrals of Eisenstein series and consequently appear in the spectral expansion of the relative trace formula. They have also been considered for non-Galois symmetric pairs (see e.g. [Off06] where they were used to study the $\mathrm{Sp}_{2n}(\mathbb{A})$ -period integral on the residual spectrum of $\mathrm{GL}_{2n}(\mathbb{A})$).

Here, building on [Off17], we develop in Section 3 their local theory for Galois pairs, in analogy with the global work [LR03]. Returning to the local setting where F is p -adic, our

main result on local intertwining periods is their convergence in a cone and meromorphic continuation. More precisely, let P be a standard parabolic subgroup of G (with respect to a fixed minimal parabolic subgroup and a fixed θ -stable maximal F -split torus in it) and M the standard Levi subgroup of P . Let $x \in G$ be such that $\theta(x) = x^{-1}$ and assume further that the involution $\theta_x = \text{Ad}(x) \circ \theta$ of G stabilizes M . Denote by S_x the θ_x -fixed points in a subset S of G .

For a finite length (smooth complex valued) representation σ of M with bounded matrix coefficients, $\ell \in \text{Hom}_{M_x}(\sigma, \mathbf{1}_{M_x})$ and χ an unramified character of M anti-invariant under θ_x , we show that for χ in a certain sufficiently positive cone, the integral

$$\int_{P_x \backslash G_x} \ell(\varphi_\chi(h)) dh$$

converges absolutely for all holomorphic sections $\varphi_\chi \in I_P^G(\chi \otimes \sigma)$ and admits a meromorphic continuation with respect to χ . When ℓ is nonzero, by taking a leading term of such an intertwining period we prove in Proposition 3 the existence of a nonzero G_x -invariant linear form on $I_P^G(\sigma)$. The existence of such ℓ is our new sufficient condition for distinction. It immediately leads to our characterization of distinction when σ is cuspidal (Corollary 2).

The Levi subgroup M acts naturally by twisted conjugation on the set of all x as above and the condition on a cuspidal σ in Corollary 2 is expressed in terms of the M -orbits. The formulation is not always convenient to use in view of the motivation, obtaining results in terms of Langlands parametrization. Hence in Section 5, we make our criterion completely explicit when \mathbb{H} is a classical group. This is the content of Theorem 2. Its proof requires a careful analysis of orbits and stabilizers that is carried out in Section 6. In particular, for this task we obtain in §4.3 the classification of the G -orbits in the symmetric space $X = \{x \in G : \theta(x) = x^{-1}\}$ with respect to twisted conjugation $g \cdot x = gx\theta(g)^{-1}$.

Let K/k be a quadratic extension of number fields with Galois involution still denoted by θ . As an application of Theorem 2 we prove in Theorem 3 that an irreducible generic cuspidal automorphic representation of the split odd special orthogonal group $\text{SO}_{2n+1}(\mathbb{A}_K)$ which is $\text{SO}_{2n+1}(\mathbb{A}_k)$ -distinguished (or more generally, distinguished by a certain K/k -inner form) is automatically θ -conjugate self-dual. The rigidity theorem of [JS03] is a key ingredient in our proof. From this and a globalization result of Beuzart-Plessis [BP] we deduce a local analogue of this result in Theorem 4 that slightly generalizes, for p -adic SO_{2n+1} , a Theorem in [BP] for general p -adic quasi-split groups. We also deduce in Proposition 7 similar results for some non-generic representations.

In Section 8 we study a relation between Galois distinction for representations of quasi-split classical groups and their transfer to a general linear group. As explained by Proposition 8 this relation is a consequence of Dipendra Prasad's Conjecture 1. As an application of the main results in this work, we prove this relation in Theorem 7 for a certain class of generic irreducible representations of the classical group, that contains all irreducible generic principal series.

Finally in Appendix A we provide explicit formulas for Prasad's quadratic character for classical groups. Most of these formulas are well-known but we give proofs for convenience of the reader. We also establish further properties of this character.

ACKNOWLEDGEMENT

We warmly thank Raphaël Beuzart-Plessis for his help in the writing of Section A.1 of Appendix A, and Ahmed Moussaoui for useful conversations and explanations. We also thank Colette Moeglin for useful correspondence. The first named author thanks the CNRS for giving him a “délégation” in 2020.

2. NOTATION AND PRELIMINARIES

Let F be a non-archimedean local field of characteristic zero and $|\cdot|$ the normalized absolute value on F . For an affine algebraic variety \mathbb{X} defined over F set $X = \mathbb{X}(F)$. Let \mathbb{G} be a connected reductive group defined over F . We denote by $X^*(G)$ the group of F -rational characters from \mathbb{G} to the multiplicative group \mathbb{G}_m and let $\mathfrak{a}_G^* = X^*(G) \otimes_{\mathbb{Z}} \mathbb{R}$. Let $\mathfrak{a}_G = \text{Hom}(\mathfrak{a}_G^*, \mathbb{R})$ be the dual vector space and let $H_G : G \rightarrow \mathfrak{a}_G$ be the map defined by

$$e^{\langle \chi, H_G(g) \rangle} = |\chi(g)|, \quad \chi \in X^*(G), \quad g \in G.$$

For \mathfrak{a} a real vector space we denote by $\mathfrak{a}_{\mathbb{C}} = \mathfrak{a} \otimes_{\mathbb{R}} \mathbb{C}$ its complexification.

Throughout the paper, we mostly avoid mention of the underlying algebraic groups defined over F and consider only their groups of F -points. When we say that P is a parabolic subgroup of G we mean that $P = \mathbb{P}(F)$ where \mathbb{P} is a parabolic subgroup of \mathbb{G} defined over F . We use a similar convention for other algebraic properties such as, (split) tori, unipotent subgroups etc.

Let A_G be the split component of G , that is, the maximal split torus in the center of G . We recall that restriction to A_G identifies \mathfrak{a}_G^* with $\mathfrak{a}_{A_G}^*$.

For groups $A \subseteq B$ denote by $Z_B(A)$ the centralizer of A in B and by $N_B(A)$ the normalizer of A in B . For a locally compact group Q denote by δ_Q its modulus function.

2.1. Parabolic subgroups and Iwasawa decomposition. Let P_0 be a minimal parabolic subgroup of G and T a maximal F -split torus of G contained in P_0 . The group P_0 admits a Levi decomposition $P_0 = M_0 \ltimes U_0$ where $M_0 = Z_G(T)$ and U_0 is the unipotent radical of P_0 . Let $\mathfrak{a}_0 = \mathfrak{a}_T = \mathfrak{a}_{M_0}$ and $\mathfrak{a}_0^* = \mathfrak{a}_T^* = \mathfrak{a}_{M_0}^*$.

A parabolic subgroup P of G is called semi-standard if it contains T , and standard if it contains P_0 . A semi-standard parabolic subgroup P of G with unipotent radical U admits a unique Levi decomposition $P = M \ltimes U$ with the Levi subgroup M containing T . Such a Levi subgroup is called semi-standard. If in addition P is standard, M is called a standard Levi subgroup. Whenever we write $P = M \ltimes U$ is a (semi-)standard parabolic subgroup we implicitly assume the above (semi-)standard Levi decomposition.

Fix a maximal compact open subgroup K of G in good position with respect to P_0 (see [MW95, I.1.4]). For a standard parabolic subgroup $P = M \ltimes U$ of G this allows one to extend H_M to a function on $G = UMK$ by

$$H_M(umk) = H_M(m), \quad u \in U, \quad m \in M, \quad k \in K.$$

2.2. Roots and Weyl groups.

2.2.1. *Roots.* Denote by $R(T, G)$ the root system of G with respect to T , by $R(T, P_0)$ the set of positive roots of $R(T, G)$ with respect to P_0 , and by Δ_0 the corresponding set of simple roots. Note that $R(T, G)$ lies in \mathfrak{a}_0^* . For every $\alpha \in R(T, G)$ we denote by $\alpha^\vee \in \mathfrak{a}_0$ the corresponding coroot.

There is a unique element $\rho_0 \in \mathfrak{a}_0^*$ (half the sum of positive roots with multiplicities) such that

$$\delta_{P_0}(p) = e^{\langle 2\rho_0, H_{M_0}(p) \rangle}, \quad p \in P_0.$$

For a standard parabolic subgroup $P = M \times U$ of G let $R(A_M, G)$ be the set of non-trivial restrictions to A_M of elements of $R(T, G)$, $R(A_M, P)$ the subset of non-zero restrictions to A_M of elements of $R(T, P_0)$ and Δ_P the subset of non-zero restrictions to A_M of elements in Δ_0 . For $\alpha \in R(A_M, G)$ we write $\alpha > 0$ if $\alpha \in R(A_M, P)$ and $\alpha < 0$ otherwise.

Let $Q = L \times V$ be a parabolic subgroup of G containing P . The restriction map from A_M to A_L defines a projection $\mathfrak{a}_M^* \rightarrow \mathfrak{a}_L^*$ that gives rise to a direct sum decomposition $\mathfrak{a}_M^* = \mathfrak{a}_L^* \oplus (\mathfrak{a}_M^L)^*$ (the second component is the kernel of this projection) and a compatible decomposition $\mathfrak{a}_M = \mathfrak{a}_L \oplus \mathfrak{a}_M^L$ for the dual spaces. For $\lambda \in \mathfrak{a}_M^*$ (respectively $\nu \in \mathfrak{a}_M$) we write $\lambda = \lambda_L + \lambda_M^L$ (respectively, $\nu = \nu_L + \nu_M^L$) for the corresponding decomposition. Set

$$\Delta_P^Q = \{\alpha \in \Delta_P, \alpha_L = 0\}.$$

(Note that Δ_P^Q identifies with $\Delta_{P \cap L}$ defined with respect to $(L, L \cap P_0)$ replacing (G, P_0) .)

The coroot $\alpha^\vee \in \mathfrak{a}_M$ associated to $\alpha \in R(A_M, G)$ is defined as follows: let $\alpha_0 \in R(T, G)$ be such that $\alpha = (\alpha_0)_M$ then one sets $\alpha^\vee = (\alpha_0^\vee)_M$ (it is independent of the choice of α_0).

Let $\rho_P = (\rho_0)_M$. We have the relation

$$\delta_P(p) = e^{\langle 2\rho_P, H_M(p) \rangle}, \quad p \in P.$$

2.2.2. *Weyl groups and Bruhat decomposition.* We denote by $W_G := N_G(T)/M_0$ the Weyl group of G . Note that for a semi-standard parabolic subgroup $P = M \times U$ of G the Levi part M contains M_0 . Therefore expressions such as wP and Pw are well defined for $w \in W_G$. Furthermore, W_M is a subgroup of W_G . Let $P = M \times U$ and $Q = L \times V$ be semi-standard parabolic subgroups of G . The Bruhat decomposition is the bijection $W_M w W_L \mapsto P w Q$ from $W_M \backslash W_G / W_L$ to $P \backslash G / Q$.

2.2.3. *Elementary symmetries.* For standard Levi subgroups $M \subseteq L$ of G let $W_L(M)$ be the set of elements $w \in W_L$ such that w is of minimal length in wW_M and wMw^{-1} is a standard Levi subgroup of G . Note that for $w \in W_L(M)$ and $w' \in W_L(wMw^{-1})$ we have $w'w \in W_L(M)$. By definition $W_L(M)$ is the disjoint union over standard Levi subgroups M' of G of the sets

$$W_L(M, M') = \{w \in W_L(M) : wMw^{-1} = M'\}.$$

Set $W(M) = W_G(M)$. In [MW95, I.1.7, I.1.8] the elementary symmetries $s_\alpha \in W(M)$ attached to each $\alpha \in \Delta_P$ are introduced and used in order to define a length function ℓ_M on $W(M)$. There is a unique element of $W_L(M)$ of maximal length for ℓ_M and we denote it by w_M^L .

2.3. Symmetric spaces. By a symmetric pair (G, θ) we mean that θ is an involution of \mathbb{G} defined over F . We say that the symmetric pair (G, θ) is a Galois pair if $\mathbb{G} = \text{Res}_{E/F} \mathbb{H}$ is the Weil restriction of scalars of a connected reductive group \mathbb{H} defined over F , E/F is a quadratic field extension and θ is the associated Galois action. Thus, for a Galois pair (G, θ) we have $G = \mathbb{H}(E)$ and $G^\theta = \mathbb{H}(F)$.

To a symmetric pair (G, θ) we associate the symmetric space

$$X = \{g \in G, \theta(g) = g^{-1}\}$$

equipped with the twisted conjugation G -action

$$g \cdot x = gx\theta(g)^{-1}, \quad g \in G, x \in X.$$

By [HW93, Proposition 6.15], X consists of a finite number of G -orbits. For $x \in X$ we denote by θ_x the involution of G given by

$$\theta_x(g) = x\theta(g)x^{-1}.$$

For $x \in X$ and Q a subgroup of G , we denote by Q_x the stabilizer of x in Q , so that

$$Q_x = (Q \cap \theta_x(Q))^{\theta_x}.$$

Such an involution θ will induce linear involutions on different vector spaces. Whenever θ is an involution acting linearly on a vector space V over \mathbb{R} or \mathbb{C} , we write

$$V = V_\theta^+ \oplus V_\theta^-$$

for the corresponding decomposition into the θ -eigenspaces V_θ^\pm with corresponding eigenvalues ± 1 .

2.4. Parabolic induction and standard intertwining operators. By a representation of a subgroup of G we always mean a smooth complex valued representation.

2.4.1. Induced representations. Let $P = M \ltimes U$ be a standard parabolic subgroup of G and σ a representation of M . We denote by $I_P^G(\sigma)$ the representation of G by right translations on the space of functions φ on G with values in the space of σ that are right invariant by some open subgroup of G and satisfy

$$\varphi(umg) = \delta_P(m)^{1/2} \sigma(m) \varphi(g), \quad u \in U, m \in M, g \in G.$$

2.4.2. Holomorphic sections. For $\lambda \in \mathfrak{a}_{M, \mathbb{C}}^*$ we denote by $\sigma[\lambda]$ the representation of M on the space of σ given by $\sigma[\lambda](m) = e^{\langle \lambda, H_M(m) \rangle} \sigma(m)$. In order to make sense of meromorphic families of linear forms on induced representations we realize all the representations $I_P^G(\sigma[\lambda])$ for $\lambda \in \mathfrak{a}_{M, \mathbb{C}}^*$ on the space of $I_P^G(\sigma)$.

For $\lambda \in \mathfrak{a}_{M, \mathbb{C}}^*$ and $\varphi \in I_P^G(\sigma)$ write $\varphi_\lambda(g) = e^{\langle \lambda, H_M(g) \rangle} \varphi(g)$ and let $I_P^G(\sigma, \lambda)$ be the representation of G on the space $I_P^G(\sigma)$ defined by

$$(I_P^G(g, \sigma, \lambda) \varphi)_\lambda(x) = \varphi_\lambda(xg).$$

The map $\varphi \mapsto \varphi_\lambda$ is an isomorphism of representations $I_P^G(\sigma, \lambda) \rightarrow I_P^G(\sigma[\lambda])$.

2.4.3. *Intertwining operators.* Let $w \in W(M)$ and select a representative n of w in $N_G(T)$. Set $M_1 = wMw^{-1} = nMn^{-1}$ and let $P_1 = M_1 \times U_1$ be the standard parabolic subgroup of G with Levi subgroup M_1 . For $c \in \mathbb{R}$ let

$$\mathcal{D}^{M,w}(c) = \{\lambda \in \mathfrak{a}_{M,\mathbb{C}}^* : \langle \operatorname{Re}(\lambda), \alpha^\vee \rangle > c, \forall \alpha \in R(A_M, G), \alpha > 0, w\alpha < 0\}.$$

We denote by $n\sigma$ the representation of M_1 on the space of σ defined by $(n\sigma)(nmn^{-1}) = \sigma(m)$, $m \in M$. Note that the isomorphism class of $n\sigma$ is independent of the choice of representative n for w and we denote it by $w\sigma$. For $\lambda \in \mathfrak{a}_{M,\mathbb{C}}^*$ we have $w\lambda \in \mathfrak{a}_{M_1,\mathbb{C}}^*$ and the standard intertwining operator

$$M(n, \sigma, \lambda) : I_P^G(\sigma, \lambda) \rightarrow I_{P'}^G(w\sigma, w\lambda)$$

is the meromorphic continuation of the operator given by the following convergent integral for $\lambda \in \mathcal{D}^{M,w}(c_\sigma)$ for some constant c_σ

$$(M(n, \sigma, \lambda)\varphi)_{w\lambda}(g) = \int_{U_1 \cap wUw^{-1} \setminus U_1} \varphi_\lambda(n^{-1}ug) \, du.$$

By definition, the convergence of the integral in the appropriate domain means that for every element v^\vee in the smooth dual of σ the scalar valued integral

$$\int_{U_1 \cap wUw^{-1} \setminus U_1} v^\vee[\varphi_\lambda(n^{-1}ug)] \, du$$

converges. We record the following simple consequence.

Lemma 1. *With the above notation, let $\lambda \in \mathcal{D}^{M,w}(c_\sigma)$ and ℓ a linear form on the space of σ such that for every $\varphi \in I_P^G(\sigma)$ and $g \in G$ the integral*

$$\int_{U_1 \cap wUw^{-1} \setminus U_1} |\ell(\varphi_\lambda(n^{-1}ug))| \, du$$

converges. Then

$$\ell((M(n, \sigma, \lambda)\varphi)_{w\lambda}(g)) = \int_{U_1 \cap wUw^{-1} \setminus U_1} \ell(\varphi_\lambda(n^{-1}ug)) \, du.$$

Proof. Replacing φ with $I_P^G(g, \sigma, \lambda)\varphi$ we assume without loss of generality that g is the identity e of G . Let \mathcal{M}_1 be a compact open subgroup of M_1 such that φ is right \mathcal{M}_1 -invariant and $\mathcal{M} = n^{-1}\mathcal{M}_1n$ a compact open subgroup of M . Let

$$\ell_0(v) = \int_{\mathcal{M}} \ell(\sigma(m)v) \, dm$$

for every v in the space of σ where integration is over the probability measure of \mathcal{M} . Then ℓ_0 lies in the smooth dual of σ . For $m \in \mathcal{M}$ we have

$$\sigma(m)[(M(n, \sigma, \lambda)\varphi)_{w\lambda}(e)] = (M(n, \sigma, \lambda)\varphi)_\lambda(nmn^{-1}) = (M(n, \sigma, \lambda)\varphi)_{w\lambda}(e)$$

and therefore

$$\ell((M(n, \sigma, \lambda)\varphi)_{w\lambda}(g)) = \ell_0((M(n, \sigma, \lambda)\varphi)_{w\lambda}(g)) = \int_{U_1 \cap wUw^{-1} \setminus U_1} \ell_0(\varphi_\lambda(n^{-1}ug)) \, du.$$

Let $(\mathcal{U}_k)_{k \in \mathbb{N}}$ be an increasing sequence of compact open subgroups of U_1 with union U_1 and such that \mathcal{M}_1 normalizes each \mathcal{U}_k . (It is easy to see that such a sequence exists.) By the dominated convergence theorem we have

$$\int_{U_1 \cap wUw^{-1} \setminus U_1} \ell_0(\varphi_\lambda(n^{-1}u)) \, du = \lim_{k \rightarrow \infty} \int_{\mathcal{U}_k \cap wUw^{-1} \setminus \mathcal{U}_k} \ell_0(\varphi_\lambda(n^{-1}u)) \, du.$$

Note that for every k we have

$$\begin{aligned} \int_{\mathcal{U}_k \cap wUw^{-1} \setminus \mathcal{U}_k} \ell_0(\varphi_\lambda(n^{-1}u)) \, du &= \int_{\mathcal{U}_k \cap wUw^{-1} \setminus \mathcal{U}_k} \int_{\mathcal{M}} \ell(\sigma(m)[\varphi_\lambda(n^{-1}u)]) \, dm \, du \\ &= \int_{\mathcal{U}_k \cap wUw^{-1} \setminus \mathcal{U}_k} \int_{\mathcal{M}} \ell(\varphi_\lambda(mn^{-1}u)) \, dm \, du. \end{aligned}$$

After the change of variables $m \mapsto n^{-1}mn$ this becomes

$$\int_{\mathcal{U}_k \cap wUw^{-1} \setminus \mathcal{U}_k} \int_{\mathcal{M}_1} \ell(\varphi_\lambda(n^{-1}mu)) \, dm \, du.$$

Changing order of integration, making the change of variables $u \mapsto m^{-1}um$ and applying the right \mathcal{M}_1 -invariance of φ this becomes

$$\int_{\mathcal{U}_k \cap wUw^{-1} \setminus \mathcal{U}_k} \ell(\varphi_\lambda(n^{-1}u)) \, du.$$

Applying the dominated convergence theorem again we have that

$$\lim_{k \rightarrow \infty} \int_{\mathcal{U}_k \cap wUw^{-1} \setminus \mathcal{U}_k} \ell(\varphi_\lambda(n^{-1}u)) \, du = \int_{U_1 \cap wUw^{-1} \setminus U_1} \ell(\varphi_\lambda(n^{-1}u)) \, du$$

and the lemma follows. \square

2.5. Meromorphic families of linear forms. For a finite dimensional complex vector space V and a complex vector space \mathcal{V} we say that $L(\lambda)_{\lambda \in V}$ is a meromorphic family of linear forms on \mathcal{V} if there is a non-zero polynomial function f on V such that $f(\lambda)L(\lambda)$ is a linear form on \mathcal{V} for every $\lambda \in V$ and for all $\nu \in \mathcal{V}$ the map

$$\lambda \mapsto f(\lambda)L(\lambda)\nu$$

from V to \mathbb{C} is holomorphic.

2.6. The result of Blanc and Delorme. For (G, θ) a symmetric pair, let $P = M \ltimes U$ be a parabolic subgroup of G such that $P \cap \theta(P) = M$. Note that in this case $P^\theta = M^\theta$ is unimodular. Furthermore, since M is θ -stable, θ acts as a linear involution on \mathfrak{a}_M^* so that

$$\mathfrak{a}_M^* = (\mathfrak{a}_M^*)_\theta^+ \oplus (\mathfrak{a}_M^*)_\theta^-.$$

There is a similar decomposition to the dual space \mathfrak{a}_M so that $(\mathfrak{a}_M)_\theta^\pm$ is dual to $(\mathfrak{a}_M^*)_\theta^\pm$. In particular, we have

$$\langle \lambda, H_M(m) \rangle = 0, \quad \lambda \in (\mathfrak{a}_M^*)_\theta^-, \quad m \in M^\theta.$$

For $c > 0$ let

$$\mathcal{D}_{M, \theta}(c) = \{\lambda \in (\mathfrak{a}_M^*)_\theta^- : \langle \lambda, \alpha^\vee \rangle > c, \forall \alpha \in R(A_M, P)\}.$$

The following is a consequence of [BD08]. Let σ be a representation of M of finite length. There exists $c > 0$ such that for all $\varphi \in I_P^G(\sigma)$, $\ell \in \text{Hom}_{M^\theta}(\sigma, \mathbf{1}_{M^\theta})$ and $\lambda \in \mathcal{D}_{M,\theta}(c)$ the integral

$$J_P^G(\varphi; \theta, \ell, \sigma, \lambda) = \int_{M^\theta \backslash G^\theta} \ell(\varphi_\lambda(g)) dg$$

is absolutely convergent. Furthermore, it admits a meromorphic continuation to a meromorphic family of linear forms $J_P^G(\theta, \ell, \sigma, \lambda) \in \text{Hom}_{G^\theta}(I_P^G(\sigma, \lambda), \mathbf{1}_{G^\theta})$, $\lambda \in (\mathfrak{a}_{M,\mathcal{C}}^*)_\theta^-$.

3. INTERTWINING PERIODS

Our goal in this section is to construct certain meromorphic families of invariant linear forms, local intertwining periods, on induced representations associated to a p-adic Galois pair. In the global setting, intertwining periods were introduced in [Jac95] and studied further in a more general setting in [JLR99] and [LR03].

3.1. An informal introduction. The local intertwining periods emerge as follows. Consider a symmetric pair (G, θ) . Let χ be a character of G^θ . For a representation π of G we denote by $\text{Hom}_{G^\theta}(\pi, \chi)$ the space of (G^θ, χ) -equivariant linear forms on the space of π .

Consider a parabolic subgroup P of G with a θ -stable Levi subgroup M and let U be the unipotent radical of P so that $P^\theta = M^\theta \ltimes U^\theta$. Let σ be a representation of M and $\ell \in \text{Hom}_{M^\theta}(\sigma, \delta_{P^\theta} \delta_P^{-1/2} \chi)$. For $\varphi \in I_P^G(\sigma)$, the map $g \mapsto \chi(g)^{-1} \ell(\varphi(g))$ is left $(P^\theta, \delta_{P^\theta})$ -equivariant and therefore at least formally the integral

$$L(\varphi) = \int_{P^\theta \backslash G^\theta} \chi(g)^{-1} \ell(\varphi(g)) dg$$

makes sense. If it converges it defines a linear form $L \in \text{Hom}_{G^\theta}(I_P^G(\sigma), \chi)$. In general, however, the integral fails to converge. By considering unramified twists of σ one hopes to define the linear form by a convergent integral on some cone of unramified twists and by meromorphic continuation in general.

In this section we achieve this goal when (G, θ) is a Galois pair and χ is the trivial character.

3.2. The strategy. Our proof of convergence in a cone and meromorphic continuation is based on the study of the geometry and combinatorics of the P -orbits on G/G^θ carried out in [LR03, §3-4]. It allows us to study the intertwining periods inductively along a certain directed graph associated to the P -orbits. Roughly speaking, vertices in the graph are associated to certain P -orbits and edges to a relation between them defined via twisted conjugation. For a directed edge one expresses the intertwining period associated with the origin as a double integral that formally corresponds with the composition of the intertwining period associated with the destination and an intertwining operator. These are the local analogues of the simple functional equations obtained in [LR03, Proposition 10.1.1]. They allow reduction of the meromorphic continuation from origin to destination. In order to reduce convergence of the origin intertwining period to the destination one it is required to prove that the above composition expressed as a double integral is absolutely

convergent. It is done in two steps. First, we prove it for unramified representations induced from real exponents. In this case the integrands at hand are positive and it is enough to prove convergence as an iterated integral. This reduction is carried out applying the Gindikin-Karpelevich formula and convergence of the standard intertwining operators. In the second step we use the bounds obtained by Lagier in [Lag08, Theorem 4] in order to reduce to the first case. This inductive process along the graph allows us to reduce convergence in a cone and meromorphic continuation to certain minimal cases. By a local analogue of [LR03, Lemma 5.4.1], the minimal case is reduced to the main results established by Blanc and Delorme in [BD08]. The entire proof is a local analogue of the one carried by Lapid and Rogawski in [LR03]. We remark that Lapid and Rogawski assume that G admits a θ -stable minimal parabolic subgroup. Thanks to the analysis of P -orbits on G/G^θ for a general involution θ of G in [Off17] we do not need to make this assumption.

3.3. The set up. For the rest of this section assume that (G, θ) is a Galois symmetric pair. Fix the minimal parabolic subgroup P_0 of G . According to [HW93, Lemma 2.4] there exists a θ -stable maximal split torus T of G contained in P_0 . Thus, P_0 admits a Levi decomposition $P_0 = M_0 \ltimes U_0$ where $M_0 = Z_G(T)$ is a θ -stable Levi subgroup of P_0 .

3.4. A graph of involutions. For a standard Levi subgroup M of G let

$$X[M] = \{x \in X : \theta_x(M) = M\}.$$

Note for example that $X[M_0] = X \cap N_G(T)$.

We define a directed labeled graph $\mathcal{G} = \mathcal{G}(G, \theta, P_0, T)$ as follows. The vertices of \mathcal{G} are the pairs (M, x) where M is a standard Levi subgroup of G and $x \in X[M]$. The edges of \mathcal{G} are given by

$$(M, x) \xrightarrow{n} (M_1, x_1)$$

if there is $\alpha \in \Delta_P$ with $-\alpha \neq \theta_x(\alpha) < 0$ such that $n \in s_\alpha M$ where $s_\alpha \in W(M)$ is the elementary symmetry associated to α , $M_1 = nMn^{-1}$ and $x_1 = n \cdot x$. Note that $(M_1)_{x_1} = nM_xn^{-1}$.

3.5. Minimal vertices. Recall that the Weyl group W_G acts (by conjugation) simply transitively on the set of minimal semi-standard parabolic subgroups of G . There is therefore a unique $\tau \in W_G$ such that $\theta(P_0) = \tau P_0 \tau^{-1}$.

Let (M, x) be a vertex in \mathcal{G} and P the standard parabolic subgroup of G with Levi subgroup M . Set $P' = \tau^{-1}\theta(P)\tau$. Then $P' = M' \ltimes U'$ is a standard parabolic subgroup of G with standard Levi subgroup $M' = \tau^{-1}\theta(M)\tau = \tau^{-1}x^{-1}Mx\tau$ and unipotent radical $U' = \tau^{-1}\theta(U)\tau$. We say that (M, x) is minimal if there exists a standard parabolic subgroup $Q = L \ltimes V$ of G containing P such that

- $\theta(L) = \tau L \tau^{-1}$;
- $Mx\tau M' = Mw_{M'}^L M'$;
- $\theta_x(\alpha) = -\alpha$, $\alpha \in \Delta_P^Q$.

This definition coincides with [Off17, Definition 6.6]. Note that the conditions $M \subseteq L$ together with $\theta(L) = \tau L \tau^{-1}$ imply that $M' \subseteq L$ and therefore $w_{M'}^L$ makes sense. The following is a straightforward consequence of the definition.

Lemma 2. *Let (M, x) be a minimal vertex in \mathcal{G} . In the above notation Q , L and V are θ_x stable.*

Proof. Let $n \in N_G(T)$ be a representative of τ . It follows from the second point of the definition that $xn \in L$ and from the first that $L = n^{-1}\theta(L)n$. Conjugating with xn we conclude that $L = \theta_x(L)$. Since $n^{-1}\theta(Q)n$ is a standard parabolic subgroup of G with Levi subgroup $n^{-1}\theta(L)n = L$ we conclude that $n^{-1}\theta(Q)n = Q$ and conjugation with xn similarly shows that $\theta_x(Q) = Q$. Since V is the unipotent radical of Q it immediately follows that $\theta_x(V) = V$. \square

As a consequence of [LR03, Lemma 3.2.1 and Proposition 3.3.1] (see [Off17, Corollary 6.7]) we have the following.

Proposition 1. *Let (M, x) be a vertex in the graph \mathcal{G} . There exists a path*

$$(M, x) \searrow^{n_1} (M_1, \theta_1) \searrow^{n_2} \cdots \searrow^{n_k} (M_k, x_k)$$

in \mathcal{G} such that (M_k, x_k) is a minimal vertex. \square

3.6. The cone of convergence. Let (M, x) be a vertex in \mathcal{G} . Recall that θ_x acts as an involution on \mathfrak{a}_M^* and let

$$(\mathfrak{a}_M^*)_x^- = (\mathfrak{a}_M^*)_{\theta_x}^-.$$

For $c > 0$ let

$$\mathcal{D}_{M,x}(c) = \{\lambda \in (\mathfrak{a}_M^*)_x^- : \langle \lambda, \alpha^\vee \rangle > c, \forall \alpha \in R(A_M, G), \alpha > 0, \theta_x(\alpha) < 0\}.$$

For an edge $(M, x) \searrow^n (M_1, x_1)$ in \mathcal{G} let $\alpha \in \Delta_P$ be such that $n \in s_\alpha M$. By [LR03, Lemma 5.2.1] we have

$$\mathcal{D}_{M,x}(c) = s_\alpha^{-1} \mathcal{D}_{M_1, x_1}(c) \cap \{\lambda \in (\mathfrak{a}_M^*)_x^- : \langle \lambda, \alpha^\vee \rangle > c\}.$$

3.7. Admissible orbits. Let $P = M \ltimes U$ be a standard parabolic subgroup of G . For $x \in X$ let $w \in W_G$ be such that $Px\theta(P) = Pw\theta(P)$. (By the Bruhat decomposition, the double coset $W_M w W_{\theta(M)}$ is uniquely determined by x .)

We say that x (or its P -orbit in X) is *P -admissible* if $M = w\theta(M)w^{-1}$. If a P -orbit O in X is M admissible then it contains an element of $X[M]$. In fact $O \mapsto O \cap X[M]$ is a bijection from the admissible P -orbits in X to the M -orbits in $X[M]$. (See [Off17, Lemma 3.2]). It further follows from [Off17, Corollary 6.9] that

$$(1) \quad \delta_P^{1/2}|_{P_x} = \delta_{P_x}, \quad x \in X[M].$$

As a consequence, for a representation σ of M , $x \in X[M]$, $\ell \in \text{Hom}_{M_x}(\sigma, \mathbf{1}_{M_x})$, $\varphi \in I_P^G(\sigma)$ and $\lambda \in (\mathfrak{a}_{M,\mathbb{C}}^*)_x^-$ we have

$$(2) \quad \ell(\varphi_\lambda(pg)) = \delta_{P_x}(p)\ell(\varphi_\lambda(g)), \quad p \in P_x, g \in G.$$

3.8. The intertwining periods modulo convergence. It follows from (2) in its notation that the following integral formally makes sense

$$J_P^G(\varphi; x, \ell, \sigma, \lambda) = \int_{P_x \backslash G_x} \ell(\varphi_\lambda(g)) dg.$$

If λ is such that the integral is convergent for all $\varphi \in I_P^G(\sigma)$ then it defines a linear form $J_P^G(x, \ell, \sigma, \lambda) \in \text{Hom}_{G_x}(I_P^G(\sigma, \lambda), \mathbf{1}_{G_x})$.

Assuming that σ is of finite length and with bounded matrix coefficients we prove that there exists $c > 0$ such that the integrals are absolutely convergent as long as $\text{Re}(\lambda) \in \mathcal{D}_{M,x}(c)$ and admit meromorphic continuation to $\lambda \in (\mathfrak{a}_{M,\mathbb{C}}^*)_x^-$. We will prove this by induction along a path as in Proposition 1 in the graph \mathcal{G} .

3.9. Intertwining periods for minimal vertices. Our first step is the observation that the results of Blanc and Delorme (§2.6) imply the base of the induction.

Corollary 1. *Let (M, x) be a minimal vertex in \mathcal{G} and σ a representation of M of finite length. Then for any $\ell \in \text{Hom}_{M_x}(\sigma, \mathbf{1}_{M_x})$ the linear form $J_P^G(x, \ell, \sigma, \lambda)$ is defined by a convergent integral for $\lambda \in \mathcal{D}_{M,x}(c)$ and admits a meromorphic continuation to $\lambda \in (\mathfrak{a}_{M,\mathbb{C}}^*)_x^-$.*

Proof. We use the notation of §3.5 for a minimal vertex and integrate $J_P^G(\varphi; x, \ell, \sigma, \lambda)$ in stages. We have

$$J_P^G(\varphi; x, \ell, \sigma, \lambda) = \int_{P_x \backslash G_x} \ell(\varphi_\lambda(g)) dg = \int_{Q_x \backslash G_x} \int_{P_x \backslash Q_x} \delta_{Q_x}(m)^{-1} \ell(\varphi_\lambda(mg)) dm dg.$$

Recall that Q, L and V are θ_x -stable (Lemma 2). It follows that $Q_x \backslash G_x$ is compact (see [GO16, Lemma 3.1]). By smoothness, the outer integral may therefore be replaced by a finite sum and it is enough to prove convergence and meromorphic continuation of the inner integral.

It further follows from [Off17, Lemma 6.3] that $Q_x = L_x \rtimes V_x$ and $P_x = M_x \rtimes U_x$. Since $U = (U \cap L) \rtimes V$ we conclude that $U_x = (U_x \cap L) \rtimes V_x$. By the definition of minimality $U_x \cap L = \{e\}$ and therefore $P_x = M_x \rtimes V_x$. The inner integral is therefore

$$\int_{M_x \backslash L_x} \delta_{Q_x}(m)^{-1} \ell(\varphi_\lambda(mg)) dm.$$

Recall that by (1) we have $\delta_Q^{1/2}|_{Q_x} = \delta_{Q_x}$ and that the function $m \mapsto \delta_Q^{-1/2}(m)\varphi(mg)$, $m \in L$ lies in $I_{P \cap L}^L(\sigma)$. Thus θ_x is an involution on L and by minimality of the vertex, the parabolic subgroup $P \cap L$ of L satisfies $\theta_x(P \cap L) \cap (P \cap L) = M$. Note further that since V is θ_x -stable we have $\mathcal{D}_{M,x}(c) = \mathcal{D}_{M, \theta_x|_L}(c)$ where the right hand side is defined in §2.6. The result therefore follows from §2.6. \square

3.10. Convergence in the unramified case. Let $P = M \rtimes U$ be a standard parabolic subgroup of G and $\mathbf{1}_M$ the trivial representation of M . For any $\lambda \in \mathfrak{a}_M^*$ we consider

the induced representation $I_P^G(\mathbf{1}_M, \lambda)$ realized on the space $I_P^G(\mathbf{1}_M)$. Let $x \in X[M]$ and $\lambda \in (\mathfrak{a}_M^*)_x^-$. For $\varphi \in I_P^G(\mathbf{1}_M)$ consider the integral

$$J_P^G(\varphi; x, \lambda) = \int_{P_x \backslash G_x} \varphi_\lambda(g) dg.$$

Note that $\varphi_\lambda(pg) = \delta_P^{1/2}(p)\varphi_\lambda(g)$, $p \in P_x$ and $g \in G$ and by (1) the integral formally makes sense.

Proposition 2. *There exists $c > 0$ such that $J_P^G(\varphi; x, \lambda)$ is defined by a convergent integral for every standard parabolic subgroup $P = M \times U$ of G , $\varphi \in I_P^G(\mathbf{1}_M)$, $x \in X[M]$ and $\lambda \in \mathcal{D}_{M,x}(c)$.*

Proof. Let \mathcal{M} be the finite set of standard Levi subgroups of G . For a parabolic subgroup P of G there are finitely many P -orbits in X , [HW93, Proposition 6.15]. Consequently, there are finitely many M -orbits in $X[M]$ for every $M \in \mathcal{M}$ (see §3.7). Note further that for a standard parabolic subgroup $P = M \times U$, $x \in X[M]$ and $m \in M$ we have

$$J_P^G(\varphi; m \cdot x, \lambda) = e^{\langle \lambda + \rho_P, H_M(m) \rangle} J_P^G(I_P^G(m^{-1}, \mathbf{1}_M, \lambda)\varphi; x, \lambda)$$

where the left hand side is defined by a convergent integral if and only if the right side is.

It therefore follows from Corollary 1 that there exists $c > 0$ such that the integral $J_P^G(\varphi, x, \lambda)$ converges for all standard parabolic subgroups $P = M \times U$ of G , $\varphi \in I_P^G(\mathbf{1}_M)$, $x \in X[M]$ such that (M, x) is a minimal vertex in \mathcal{G} and $\lambda \in \mathcal{D}_{M,x}(c)$. We may further assume that c is large enough so that the intertwining operators $M(n, \mathbf{1}_M, \lambda)$ are defined by a convergent integral for every standard parabolic subgroup $P = M \times U$ of G , $\alpha \in \Delta_P$, $n \in s_\alpha M$ and $\lambda \in \mathcal{D}^{M, s_\alpha}(c)$. Note further that if $x \in X[M]$ is such that $-\alpha \neq \theta_x(\alpha) < 0$ then $\mathcal{D}_{M,x}(c) \subseteq \mathcal{D}^{M, s_\alpha}(c)$ (see §2.4.3 and §3.6).

It follows from Proposition 1 that it is enough to prove the following. Assume that $(M, x) \xrightarrow{n} (M_1, x_1)$ is an edge on the graph \mathcal{G} and $\alpha \in \Delta_P$ is such that $n \in s_\alpha M$. If $J_{P_1}^G(\varphi; x_1, s_\alpha \lambda)$ is defined by a convergent integral for every $\varphi \in I_{P_1}^G(\mathbf{1}_{M_1})$ and $\lambda \in \mathcal{D}_{M_1, x_1}(c)$ then $J_P^G(\varphi; x, \lambda)$ is defined by a convergent integral for every $\varphi \in I_P^G(\mathbf{1}_M)$ and $\lambda \in \mathcal{D}_{M,x}(c)$.

Let ξ be the spherical vector in $I_P^G(\mathbf{1}_M)$ normalized by taking the value one at the identity. Thanks to the Iwasawa decomposition, for every $\varphi \in I_P^G(\mathbf{1}_M)$ there is a positive constant A such that $|\varphi| \leq A\xi$ hence $|\varphi_\lambda| \leq A\xi_\lambda$ for all $\lambda \in \mathfrak{a}_M^*$. It therefore suffices to consider $\varphi = \xi$.

Let $P_1 = M_1 \times U_1$ be the standard parabolic subgroup with Levi subgroup M_1 and $Q = L \times V$ the standard parabolic subgroup containing P such that $\Delta_P^Q = \{\alpha\}$. For $\lambda \in \mathcal{D}_{M,x}(c)$ we have

$$J_P^G(\xi; x, \lambda) = \int_{P_x \backslash G_x} \xi_\lambda(g) dg = \int_{nP_x n^{-1} \backslash G_{x_1}} \xi_\lambda(n^{-1}gn) dg.$$

It follows from [Off17, Lemma 6.4] that $nU_x n^{-1} \subseteq (U_1)_{x_1}$ and from [Off17, Lemma 6.3] that $P_x = M_x \times U_x$ and $(P_1)_{x_1} = (M_1)_{x_1} \times (U_1)_{x_1}$.

Since $(M_1)_{x_1} = nM_x n^{-1}$ we deduce that $nP_x n^{-1} \subseteq (P_1)_{x_1}$ and the inclusion $(U_1)_{x_1} \subseteq (P_1)_{x_1}$ induces an isomorphism $nU_x n^{-1} \backslash (U_1)_{x_1} \simeq nP_x n^{-1} \backslash (P_1)_{x_1}$. Applying [Off17, Lemma

6.4] again the projection from Q to L defines an isomorphism $nU_x n^{-1} \backslash (U_1)_{x_1} \simeq L \cap U_1 \simeq (U_1 \cap nU n^{-1}) \backslash U_1$. The local analogues of [LR03, Lemma 4.3.1 (3) and (4)] easily follow from [Off17, Lemma 6.4]. Proceeding as in the proof of [LR03, Proposition 10.1.1] we get that

$$(3) \quad J_P^G(\xi; x, \lambda) = \int_{(P_1)_{x_1} \backslash G_{x_1}} \int_{(U_1 \cap nU n^{-1}) \backslash U_1} \xi_\lambda(n^{-1} u g n) \, du \, dg.$$

By our assumption on c , the inner integral is absolutely convergent and we have

$$\int_{(U_1 \cap nU n^{-1}) \backslash U_1} \xi_\lambda(n^{-1} u g n) \, du = (M(n, \mathbf{1}_M, \lambda) \xi)_{s_\alpha \lambda}(g n).$$

By the Gindikin-Karpelevich formula we have $(M(n, \mathbf{1}_M, \lambda) \xi)_{s_\alpha \lambda}(g) = c(\alpha, \lambda) \xi_{s_\alpha \lambda}(g)$, $g \in G$ where

$$c(\alpha, \lambda) = \frac{L(\langle \lambda, \alpha^\vee \rangle)}{L(1 + \langle \lambda, \alpha^\vee \rangle)},$$

$L(s) = (1 - q^{-s})^{-1}$ and q is the size of the residual field of F . By assumption the integral

$$\int_{(P_1)_{x_1} \backslash G_{x_1}} \xi_{s_\alpha \lambda}(g n) \, dg$$

converges (see §3.6) and by positivity of the integrand it follows that the double integral (3) converges. The proposition follows. \square

Remark 1. It follows from the above proposition and its proof that there exists $c = c(G, P_0) > 0$ such that the integrals defining

- the linear forms $J_P^G(x, \lambda)$ and
- the intertwining operators $M(n, \mathbf{1}_M, \lambda)$

are absolutely convergent for every standard parabolic subgroup $P = M \rtimes U$ of G , $x \in X[M]$, $\lambda \in \mathcal{D}_{M,x}(c)$, $\alpha \in \Delta_P$ such that $-\alpha \neq \theta_x(\alpha) < 0$ and $n \in s_\alpha M$. For the next section we fix such a c and write $\mathcal{D}_{M,x} = \mathcal{D}_{M,x}(c)$.

3.11. The main result. Next, we show that the intertwining periods can be defined in the Galois setting. The method of proof also provides us with the simple functional equations that they satisfy.

Theorem 1. *Let $P = M \rtimes U$ be a standard parabolic subgroup of G , σ a representation of M of finite length with bounded matrix coefficients, $x \in X[M]$ and $\ell \in \text{Hom}_{M_x}(\sigma, \mathbf{1}_{M_x})$. For every $x \in X$ such that $\theta_x(M) = M$, $\lambda \in (\mathfrak{a}_{M,\mathbb{C}}^*)_{x}^{-}$ such that $\text{Re}(\lambda) \in \mathcal{D}_{M,x}$ and $\varphi \in I_P^G(\sigma)$ the integral*

$$J_P^G(\varphi; x, \ell, \sigma, \lambda) = \int_{P_x \backslash G_x} \ell(\varphi_\lambda(g)) \, dg$$

converges. Furthermore, the linear form $J_P^G(x, \ell, \lambda)$ admits a meromorphic continuation to $\lambda \in (\mathfrak{a}_{M,\mathbb{C}}^)_{x}^{-}$ and satisfies the following functional equations. Whenever $\alpha \in \Delta_P$ and*

$n \in s_\alpha M$ are such that $(M, x) \xrightarrow{n} (M_1, x_1)$ in \mathcal{G} we have

$$J_P^G(\varphi; x, \ell, \sigma, \lambda) = J_{P_1}^G(M(n, \sigma, \lambda) I_P^G(n, \sigma, \lambda) \varphi; x_1, \ell, s_\alpha \sigma, s_\alpha \lambda).$$

Proof. We first prove the convergence for $\lambda \in (\mathfrak{a}_{M,\mathbb{C}}^*)_{x}^{-}$ such that $\operatorname{Re}(\lambda) \in \mathcal{D}_{M,x}$. Taking the absolute integral we may assume that $\lambda = \operatorname{Re}(\lambda)$. For every $g \in G$ write $g = u_g m_g k_g$ with $u_g \in U$, $m_g \in M$ and $k_g \in K$. Then

$$\ell(\varphi_\lambda(g)) = e^{\langle \lambda + \rho_P, H_M(m_g) \rangle} \ell(\sigma(m_g)\varphi(k_g)) = e^{\langle \lambda + \rho_P, H_M(g) \rangle} \ell(\sigma(m_g)\varphi(k_g)).$$

It follows from [Lag08, Théorème 4] that the function $m \mapsto \ell(\sigma(m)v)$ is bounded for every v in the space of σ . Since φ takes finitely many values on K it follows that there exists $C > 0$ such that

$$\int_{P_x \backslash G_x} |\ell(\varphi_\lambda(g))| dg \leq C \int_{P_x \backslash G_x} e^{\langle \lambda + \rho_P, H_M(g) \rangle} dg.$$

The convergence therefore follows from Proposition 2.

If (M, x) is a minimal vertex in \mathcal{G} the meromorphic continuation follows from Corollary 1. Now that convergence is granted, we show that if $\alpha \in \Delta_P$ and $n \in s_\alpha M$ are such that $(M, x) \xrightarrow{n} (M_1, x_1)$ in \mathcal{G} and $P_1 = M_1 \ltimes U_1$ is the standard parabolic subgroup of G with standard Levi subgroup M_1 then the meromorphic continuation of $J_{P_1}^G(x_1, \ell, \lambda)$ implies that of $J_P^G(x, \ell, \lambda)$. In the process we establish the desired functional equation (which already appeared in the proof of the unramified case). In fact the computation hereunder is a completely parallel generalization of that done in Proposition 2. We will use the notation of its proof and will not repeat all the arguments. In particular, $Q = L \ltimes V$ is the standard parabolic subgroup containing P such that $\Delta_P^Q = \{\alpha\}$.

Since σ has bounded matrix coefficients, it is straightforward that the integral defining $M(n, \sigma, \lambda)$ is absolutely convergent for $\lambda \in \mathcal{D}_{M,x}$ (see Remark 1). If $\operatorname{Re}(\lambda) \in \mathcal{D}_{M,x}$ the arguments in the proof of Proposition 2 imply that

$$\begin{aligned} J_P^G(\varphi; x, \ell, \sigma, \lambda) &= \int_{P_x \backslash G_x} \ell(\varphi_\lambda(g)) dg = \int_{nP_x n^{-1} \backslash G_{x_1}} \ell(\varphi_\lambda(g)) dg \\ &= \int_{(P_1)_{x_1} \backslash G_{x_1}} \int_{(U_1 \cap nU n^{-1}) \backslash U_1} \ell(\varphi_\lambda(n^{-1}ugn)) du dg \end{aligned}$$

and by Lemma 1 this equals

$$\int_{(P_1)_{x_1} \backslash G_{x_1}} \ell[(M(n, \sigma, \lambda)\varphi)_{s_\alpha \lambda}(gn)] dg.$$

The theorem now follows by induction based on Proposition 1. \square

3.12. Applications to distinction. In this section we continue to denote by (G, θ) a Galois symmetric pair and by $X = \{x \in G : \theta(x) = x^{-1}\}$ the associated symmetric space. Fix a standard parabolic subgroup $P = M \ltimes U$ of G and a representation σ of M and let $\pi = I_P^G(\sigma)$.

For every P orbit O in X there exists a standard parabolic subgroup $Q = L \ltimes V$ of G contained in P and $x \in O$ such that $M \cap \theta_x(M) = L$. We say that O contributes to σ if the normalized Jacquet module $r_{L,M}(\sigma)$ is L_x -distinguished. This property depends only on O .

Let $x \in X$. As a consequence of [Off17, Theorem 4.2 and Corollary 6.9], applied to the Galois symmetric pair (G, θ_x) , if π is G_x -distinguished then there exists a P -orbit in $G \cdot x \subseteq X$ that contributes to σ . The following is a partial converse for this necessary condition for distinction.

Proposition 3. *With the above notation, if σ is of finite length with bounded matrix coefficients and there exists $x' \in G \cdot x$ such that $\theta_{x'}(M) = M$ and σ is $M_{x'}$ -distinguished then $I_P^G(\sigma, \lambda)$ is G_x -distinguished for every $\lambda \in (\mathfrak{a}_{M, \mathbb{C}}^*)_{x'}^-$.*

Proof. We construct a meromorphic family of linear forms $L_\lambda \in \text{Hom}_{G_x}(I_P^G(\sigma, \lambda), \mathbf{1}_{G_x})$, $\lambda \in (\mathfrak{a}_{M, \mathbb{C}}^*)_{x'}^-$ on $I_P^G(\sigma)$ as follows. Let $\eta \in G$ be such that $x' = \eta \cdot x$ (so that $G_{x'} = \eta G_x \eta^{-1}$), $0 \neq \ell \in \text{Hom}_{M_{x'}}(\sigma, \mathbf{1}_{M_{x'}})$ and $L_\lambda = J_P^G(x', \ell, \sigma, \lambda) \circ I_P^G(\eta, \sigma, \lambda)$. By Theorem 1, indeed, L_λ is a meromorphic family of linear forms as required and it is clearly non zero. The proposition now follows by taking a leading term of L_λ along a generic direction. That is, for $\lambda \in (\mathfrak{a}_{M, \mathbb{C}}^*)_{x'}^-$ there exist $\lambda_0 \in (\mathfrak{a}_{M, \mathbb{C}}^*)_{x'}^-$ (a generic direction) and a non-negative integer m such that $s \mapsto s^m L_{\lambda+s\lambda_0} \in \text{Hom}_{G_x}(I_P^G(\sigma, \lambda+s\lambda_0), \mathbf{1}_{G_x})$ is holomorphic from \mathbb{C} to the linear dual of $I_P^G(\sigma)$ and non-zero at $s = 0$. Its value at $s = 0$ gives a non-zero linear form in $\text{Hom}_{G_x}(I_P^G(\sigma, \lambda), \mathbf{1}_{G_x})$. \square

As a consequence, for a representation of G induced from cuspidal, contribution of a P -orbit characterizes distinction.

Corollary 2. *Let $x \in X$ and σ a cuspidal representation of M of finite length. Then $I_P^G(\sigma)$ is G_x -distinguished if and only if there exists $x' \in G \cdot x$ such that $\theta_{x'}(M) = M$ and σ is $M_{x'}$ -distinguished.*

Proof. If L is a proper Levi subgroup of M then $r_{L, M}(\sigma) = 0$. The only if direction is therefore immediate from the necessary condition obtained by [Off17, Theorem 4.2 and Corollary 6.9].

Assume now that $x' \in G \cdot x \cap X[M]$ and σ is $M_{x'}$ -distinguished. By cuspidality σ is a direct sum of irreducible representations and it is enough to assume that σ is irreducible.

Let ω_σ be the central character of σ and note that ω_σ is trivial on $(Z_M)_{x'}$. Therefore, there exists $\mu \in (\mathfrak{a}_M^*)_{x'}^-$ such that $|\omega_\sigma| = e^{(-\mu, H_M(\cdot))}$ and therefore the unramified twist $\sigma[\mu]$ of σ is unitary and in particular has bounded matrix coefficients. Furthermore, $\text{Hom}_{M_{x'}}(\sigma, \mathbf{1}_{M_{x'}}) = \text{Hom}_{M_{x'}}(\sigma[\mu], \mathbf{1}_{M_{x'}})$. Since $I_P^G(\sigma) \simeq I_P^G(\sigma[\mu], -\mu)$ the corollary follows from Proposition 3 applied to $\sigma[\mu]$. \square

We end this section with a somewhat weaker variant of Proposition 3 that, nevertheless, removes the bounded hypothesis on the inducing representation and avoids the use of intertwining periods. For irreducibly induced representations it provides the same sufficient condition for distinction.

Proposition 4. *Suppose that $x' \in G \cdot x$ is such that $\theta_{x'}(M) = M$ and σ is $M_{x'}$ -distinguished. Then there exists $w \in W(M)$ such that $I_{P_1}^G(w\sigma)$ is G_x -distinguished. (Here P_1 is the standard parabolic subgroup of G with Levi subgroup wMw^{-1} .) In particular, if in addition $I_P^G(\sigma)$ is irreducible then it is G_x -distinguished.*

Proof. Note that for $w \in W(M)$ and a representative n of w in G we have

$$\mathrm{Hom}_{(wMw^{-1})_{n,x'}}(n\sigma, \mathbf{1}_{(wMw^{-1})_{n,x'}}) = \mathrm{Hom}_{M_{x'}}(\sigma, \mathbf{1}_{M_{x'}}).$$

Consequently, applying Proposition 1 it is enough to assume that (M, x') is in fact minimal and prove that $I_P^G(\sigma)$ is G_x -distinguished. But then let L and Q be as in Section 3.5. The representation $\sigma' = I_{P \cap L}^L(\sigma)$ is $L_{x'}$ -distinguished thanks to the proof of [Off17, Proposition 7.2]. Now $I_P^G(\sigma) \simeq I_Q^G(\sigma')$ is $G_{x'}$ -distinguished, hence G_x -distinguished, by the proof of [Off17, Proposition 7.1]. \square

Remark 2. The results of §3 also hold for symmetric pairs of Prasad and Takloo–Bighash type, i.e. those of the form $(\mathrm{GL}_m(D), C_E(\mathrm{GL}_m(D)))$ where D is an F -central division algebra of index d with md even, and $C_E(\mathrm{GL}_m(D))$ is the centralizer of an embedding of E in the space $M_m(D)$ of $m \times m$ matrices with entries in D . This follows from the fact that Equation (1) holds in this setting (see [BM19, Equation (5.3) and Remark 5.4]).

Remark 3. In particular, Proposition 3 and Corollary 2 become more powerful when $I_P^G(\sigma)$ is reducible. Note that any irreducible representation π of G is the quotient of a representation of the form $I_P^G(\sigma)$ for σ as in Proposition 3 or Corollary 2, and if π is G_x -distinguished, certainly $I_P^G(\sigma)$ is as well. Moreover, Proposition 3 provides a particular G_x -invariant linear form L on $I_P^G(\sigma)$. In the situation where $\mathrm{Hom}_{G_x}(I_P^G(\sigma), \mathbf{1}_{G_x})$ is one dimensional, this means that π is distinguished if and only if L descends to π . Such an observation is in practice very useful and has been used to study distinction of Langlands quotients in a variety of situations, see [FLO12], [Gur15], [Mat17a] or [Mat17b].

If $I_P^G(\sigma, \lambda)$ satisfies that $\mathrm{Hom}_{G_x}(I_P^G(\sigma, \lambda), \mathbf{1}_{G_x})$ is one dimensional for a generic choice of λ then the corresponding intertwining periods should satisfy functional equations more general than those obtained in this work. Namely, for any $w \in W(M)$ (represented by $n \in G$) and P_1 the standard parabolic subgroup of G with Levi subgroup wMw^{-1} , the intertwining period $J_{P_1}^G(x', \ell, \sigma, \lambda)$ satisfies an identity of the form

$$J_{P_1}^G(n \cdot x', \ell, w\sigma, w\lambda) \circ M(n, \sigma, \lambda) \circ I_P^G(n, \sigma, \lambda) = \gamma_w(\sigma, \lambda) J_P^G(x', \ell, \sigma, \lambda)$$

for some proportionality constant γ_w that is meromorphic in λ . In some cases γ_w can be explicitly expressed in terms of local invariants associated to σ , and again this has turned out very helpful to understand when the linear form L descends to π , see [FLO12], [Mat17b] or [SX].

4. CLASSICAL GALOIS PAIRS

In the next section we explicate Corollary 2 for a Galois symmetric pair (G, θ) where G is a classical group. In this section we introduce the setting and classify the G -orbits in X . This classification is new and of interest for its own sake.

4.1. Further notation and preliminaries. Let k be a non Archimedean local field of characteristic zero, K/k a field extension of degree one or two and ρ a generator of the Galois group $\mathrm{Gal}(K/k)$. Let

$$(K/k)_1 = \{a \in K^* : aa^\rho = 1\} \quad \text{and} \quad N(K/k) = \{aa^\rho : a \in K^*\}.$$

That is, $(K/k)_1$ is the subgroup $\ker(N_{K/k})$ if K/k is a quadratic extension and $N_{K/k} : K^* \rightarrow k^*$ is the norm map associated with K/k and the group $\{\pm 1\}$ otherwise while $N(K/k)$ is the subgroup $N_{K/k}(K^*)$ of k^* if K/k is a quadratic extension and the group k^{*2} of squares in k^* otherwise.

For $\varepsilon \in \{\pm 1\}$ we denote by

$$Y_n(K/k, \varepsilon) = \{y \in \mathrm{GL}_n(K) : {}^t y^\rho = \varepsilon y\}$$

the space of ε -hermitian invertible matrices with respect to the field extension K/k . It comes naturally equipped with the following right $\mathrm{GL}_n(K)$ -action

$$y \star g = {}^t g^\rho y g, \quad y \in Y_n(K/k, \varepsilon), \quad g \in \mathrm{GL}_n(K).$$

We denote by $G(y, K/k)$ the stabilizer of $y \in Y_n(K/k, \varepsilon)$ in $\mathrm{GL}_n(K)$.

This set up is a standard unified notation for considering symplectic, orthogonal and unitary groups. We recall the classification of $\mathrm{GL}_n(K)$ -orbits in $Y_n(K/k, \varepsilon)$ and explicate the stabilizers in each case.

- (1) **The symplectic case:** $K = k$, $\varepsilon = -1$ and n is even. The space $Y_n(k/k, -1)$ is not empty if and only if n is even and in that case it is the space of alternating matrices in $\mathrm{GL}_n(k)$. For $m \in \mathbb{N}$ we have that $Y_{2m}(k/k, -1) = \begin{pmatrix} & I_m \\ -I_m & \end{pmatrix} \star \mathrm{GL}_{2m}(k)$ is a unique $\mathrm{GL}_{2m}(k)$ -orbit.

The stabilizer $G(y, k/k)$ is the symplectic group $\mathrm{Sp}(y, k)$ associated to an element $y = -{}^t y$ in $\mathrm{GL}_n(k)$.

- (2) **The orthogonal case:** $K = k$ and $\varepsilon = 1$. The space $Y_n(k/k, 1)$ is the space of symmetric matrices in $\mathrm{GL}_n(k)$. The $\mathrm{GL}_n(k)$ -orbits in $Y_n(k/k, 1)$ are determined by the discriminant and the Hasse invariant. The discriminant of $y \in Y_n(k/k, 1)$ is the square class $\det y k^{*2}$. The Hasse invariant is defined as follows. There exists $g \in \mathrm{GL}_n(k)$ such that $y \star g = \mathrm{diag}(a_1, \dots, a_n)$ is diagonal and the Hasse invariant is defined by

$$\mathrm{Hasse}_k(y) = \prod_{1 \leq i < j \leq n} (a_i, a_j)_k$$

where $(\cdot, \cdot)_k$ is the quadratic Hilbert symbol on $k^* \times k^*$. We have that $y_1, y_2 \in Y_n(k/k, 1)$ are in the same $\mathrm{GL}_n(k)$ -orbit if and only if $\det y_1 k^{*2} = \det y_2 k^{*2}$ and $\mathrm{Hasse}_k(y_1) = \mathrm{Hasse}_k(y_2)$.

For $n = 1$ we have that $Y_1(k/k, 1) = k^*$ and the $\mathrm{GL}_1(k) = k^*$ -orbits are the square classes in k^*/k^{*2} . For $n = 2$ there is a unique orbit of discriminant $-k^{*2}$ (it has Hasse invariant one) and two orbits for every other discriminant. For $n \geq 3$ there are two orbits for every discriminant class.

The stabilizer $G(y, k/k)$ is the orthogonal group $\mathrm{O}(y, k)$ associated to an element $y = {}^t y$ in $\mathrm{GL}_n(k)$.

- (3) **The unitary case:** K/k is a quadratic extension. The space $Y_n(K/k, \varepsilon)$ is the space of ε -hermitian matrices in $\mathrm{GL}_n(K)$ with respect to K/k . We have that $y_1, y_2 \in Y_n(K/k, \varepsilon)$ are in the same $\mathrm{GL}_n(K)$ -orbit if and only if $y_1 N(K/k) = y_2 N(K/k)$. By local class field theory $N(K/k)$ is an index two subgroup of k^* and

therefore $Y_n(K/k, \varepsilon)$ consists of two $\mathrm{GL}_n(K)$ -orbits. The stabilizer $G(y, K/k)$ is the unitary group $U(y, K/k)$ associated to the element $y = \varepsilon {}^t y^\rho$ in $\mathrm{GL}_n(K)$.

Let $y \in Y_n(K/k, \varepsilon)$. In the symplectic case, every element of $G(y, K/k)$ has determinant one. In either the unitary or orthogonal case \det maps $G(y, K/k)$ to $(K/k)_1$. We observe below that this map is in fact surjective. We apply the well known fact that in both cases every $\mathrm{GL}_n(K)$ -orbit in $Y_n(K/k, \varepsilon)$ contains a diagonal element.

Observation 1. *In the orthogonal and unitary cases for every $y \in Y_n(K/k, \varepsilon)$ and $a \in (K/k)_1$ there exists $h \in G(y, K/k)$ such that $\det h = a$. Furthermore, in the orthogonal case there exists such an h that satisfies $h^2 = I_n$.*

Indeed, $y \star g$ is diagonal for some $g \in \mathrm{GL}_n(K)$ and we can take $h = g \operatorname{diag}(a, I_{n-1}) g^{-1}$.

4.2. Classical Galois pairs-The set up. Fix a quadratic extension of non Archimedean local fields E/F of characteristic zero and let $\sigma(x) = \bar{x}$ be the associated Galois action. In particular, $E = F[\iota]$ for some $\iota \in E \setminus F$ such that $\iota^2 \in F$.

We would like to consider distinction for unitary, special orthogonal and symplectic Galois pairs. In order to unify notation let $F' = F$ and τ be the identity on F' in the symplectic or orthogonal case while F' is a quadratic extension of F such that $E \cap F' = F$ and τ is the Galois action associated to F'/F in the unitary case. Let $E' = EF'$ and note that σ (resp. τ) has a unique extension to an automorphisms of E' over F' (resp. over E) that, by abuse of notation, we denote by σ (resp. τ) and that E'/F is a Klein extension (that is, a Galois extension such that $\operatorname{Gal}(E'/F)$ is isomorphic to the Klein group $(\mathbb{Z}/2\mathbb{Z})^2$). As a consequence, for $x \in E'$ we can write \bar{x}^τ for $\sigma(\tau(x)) = \tau(\sigma(x))$ with no ambiguity.

4.2.1. The space of ε -hermitian forms. Fix $n \in \mathbb{N}$ and $\varepsilon \in \{\pm 1\}$ with the convention that if $\varepsilon = -1$ then $F' = F$ and n is even and set $Y_n(E) = Y_n(E'/E, \varepsilon)$ and $Y_n(F) = Y_n(F'/F, \varepsilon)$.

Remark 4. Note that if F'/F is a quadratic extension and $j \in F'$ is such that $\tau(j) = -j$ then $y \mapsto jy$ is a stabilizer preserving bijection between $Y_n(E'/E, 1)$ and $Y_n(E'/E, -1)$ and therefore in the unitary case it is enough to consider $\varepsilon = 1$.

4.2.2. The general Galois symmetric space. We consider another symmetric space

$$Z_n(E) = \{g \in \mathrm{GL}_n(E') : g\bar{g} = I_n\}$$

with the left $\mathrm{GL}_n(E')$ -action $g \cdot z = gz\bar{g}^{-1}$ by twisted conjugation. Since the first cohomology set $H^1(\operatorname{Gal}(E'/F), \mathrm{GL}_n(E'))$ is trivial, we have that $Z_n(E) = \mathrm{GL}_n(E') \cdot I_n$ is a unique $\mathrm{GL}_n(E')$ -orbit.

4.2.3. The symmetric space associated to a classical Galois pair. Fix $j \in Y_n(F)$ and let G^j be the algebraic group, defined over F , of isometries of the ε -hermitian form j . That is

$$G^j(F) = \{g \in \mathrm{GL}_n(F') : {}^t g^\tau j g = j\}$$

and

$$G^j(E) = \{g \in \mathrm{GL}_n(E') : {}^t g^\tau j g = j\}.$$

Consider the $G^j(E)$ -space

$$X^j = X^j(E/F) = \{x \in G^j(E) : x\bar{x} = I_n\}$$

with the $G^j(E)$ action

$$g \cdot x = gx\bar{g}^{-1}, g \in G^j(E), x \in X^j.$$

Note that X^j is a $G^j(E)$ -invariant subspace of $Z_n(E)$.

4.3. A classification of orbits. We parameterize $G^j(E)$ -orbits in X^j in terms of certain $\mathrm{GL}_n(F')$ -orbits in $Y_n(F)$ by the following lemma.

Lemma 3. *For $x \in X^j$ let $z \in \mathrm{GL}_n(E')$ be such that $x = z \cdot I_n$ (see §4.2.2) and let $y = j \star z \in Y_n(E)$. Then, in fact, $y \in Y_n(F)$ and the assignment $G^j(E) \cdot x \mapsto y \star \mathrm{GL}_n(F')$ is well defined and defines a bijection between the $G^j(E)$ -orbits in X^j and the $\mathrm{GL}_n(F')$ -orbits in $j \star \mathrm{GL}_n(E') \cap Y_n(F)$.*

Proof. Note first that for $z \in \mathrm{GL}_n(E')$ we have that $z \cdot I_n = \bar{z}z^{-1} \in G^j(E)$ if and only if $j \star z = {}^t z^\tau jz \in Y_n(F)$. Moreover, for $z, z' \in \mathrm{GL}_n(E')$ we have that $z \cdot I_n = z' \cdot I_n$ if and only if $z^{-1}z' \in \mathrm{GL}_n(F')$ if and only if $j \star z$ and $j \star z'$ belong to the same $\mathrm{GL}_n(F')$ -orbit. Hence the formula $p(z \cdot I_n) = (j \star z) \star \mathrm{GL}_n(F')$ well defines a map p from X^j to the $\mathrm{GL}_n(F')$ -orbits in $j \star \mathrm{GL}_n(E') \cap Y_n(F)$. For $x = z \cdot I_n$ and $g \in G^j(E)$ we have $g \cdot x = (gz) \cdot I_n$ and $j \star z = j \star (gz)$ and therefore $p(x) = p(g \cdot x)$. Consequently, the map p descends to a map \bar{p} from the $G^j(E)$ -orbits in X^j to the $\mathrm{GL}_n(F')$ -orbits in $j \star \mathrm{GL}_n(E') \cap Y_n(F)$.

Assume now that $z, z' \in \mathrm{GL}_n(E')$ are such that $j \star z = j \star z' \in Y_n(F)$. Then $z'z^{-1} \in G^j(E)$ and therefore $z \cdot I_n$ and $z' \cdot I_n$ are in the same $G^j(E)$ -orbit in X^j . Consequently, the formula $q(j \star z) = G^j(E) \cdot (z \cdot I_n)$ well defines a map q from $j \star \mathrm{GL}_n(E') \cap Y_n(F)$ to the $G^j(E)$ -orbits in X^j . It is again straightforward that q is fixed on $\mathrm{GL}_n(F')$ -orbits and therefore descends to a map \bar{q} from the $\mathrm{GL}_n(F')$ -orbits in $j \star \mathrm{GL}_n(E') \cap Y_n(F)$ to the $G^j(E)$ -orbits in X^j . It is also straight forward to verify that \bar{p} and \bar{q} are inverses of one another and the lemma follows. \square

We indicate the relation between orbits in X^j for $G^j(E)$ and for its intersection with $\mathrm{SL}_n(E')$. Of course in the symplectic case $G^j(E)$ is a subgroup of $\mathrm{SL}_n(E')$, the following lemma is relevant to the case $\varepsilon = 1$.

Let $SG^j(E) = G^j(E) \cap \mathrm{SL}_n(E')$. Next we observe that determinant is the only invariant of an $SG^j(E)$ -orbit in a $G^j(E)$ -orbit in X^j .

Lemma 4. *For every $x \in X^j$ we have $SG^j(E) \cdot x = \{x' \in G^j(E) \cdot x : \det x' = \det x\}$. In particular, in the orthogonal case $SG^j(E) \cdot x = G^j(E) \cdot x$.*

Proof. Clearly, all elements of $SG^j(E) \cdot x$ have the same determinant $\det x$ and the inclusion \subseteq follows. For the other inclusion, note that if $g \in G^j(E)$ is such that $\det(g \cdot x) = \det x$ then $\det g \in F' \cap (E'/E)_1 = (F'/F)_1$. It is therefore enough to show that for any $a \in (F'/F)_1$ there exists $h \in G^j(E)$ with $\det h = a$ such that $h \cdot x = x$. Indeed, for $a = \det g^{-1}$ and such h we have $g \cdot x = (gh) \cdot x$ and $gh \in SG^j(E)$.

Now let $z \in \mathrm{GL}_n(E')$ be such that $x = z \cdot I_n$ (see §4.2.2) and recall that $y = j \star z \in Y_n(F)$ (see Lemma 3). Applying Observation 1 let $h' \in G^y(F)$ be such that $\det h' = a$ and set $h = zh'z^{-1} \in G^j(E)$. Then

$$h \cdot x = (zh') \cdot I_n = z \cdot I_n = x.$$

□

We record separately consequences of Lemmas 3 and 4 in the three cases we consider.

4.3.1. *The symplectic case.* Recall that in the symplectic case $Y_n(F) = j \star \mathrm{GL}_n(F)$ is a unique $\mathrm{GL}_n(F)$ -orbit. As an immediate consequence of Lemma 3 we obtain

Corollary 3. *In the symplectic case $X^j = G^j(E) \cdot I_n$ is a unique $G^j(E)$ -orbit.* □

4.3.2. *The unitary case.* We begin with the following simple observation.

Lemma 5. *Let L/F be a Klein extension of local fields and let E be a quadratic extension of F in L . Then $F^* \subseteq N(L/E)$.*

Proof. Let E_1 and E_2 be the two different quadratic extensions of F contained in L and different from E . Restriction to E_i defines an isomorphism from $\mathrm{Gal}(L/E)$ to $\mathrm{Gal}(E_i/F)$ and therefore $N_{L/E}(E_i^*) = N(E_i/F)$ and in particular $N(E_i/F)$ is a subgroup of $N(L/E)$, $i = 1, 2$. By local class field theory $N(E_i/F)$, $i = 1, 2$ are two different subgroups of F^* of index two and therefore their product equals F^* . The lemma follows. □

Let

$$(4) \quad \Gamma = \{a^{1-\sigma} : a \in (E'/E)_1\}$$

and note that Γ is a subgroup of $(E'/E)_1 \cap (E'/F')_1$. Note further that \det maps X^j to $(E'/E)_1 \cap (E'/F')_1$.

Observation 2. *In the unitary case, for every $a \in (E'/E)_1 \cap (E'/F')_1$ there exists $x \in X^j$ such that $\det x = a$.*

Indeed, if j is diagonal then $\mathrm{diag}(a, I_{n-1}) \in X^j$ for every $a \in (E'/E)_1 \cap (E'/F')_1$. Otherwise, there exists $g \in \mathrm{GL}_n(F')$ such that $j \star g$ is diagonal and we then have $X^j = gX^{j \star g}g^{-1}$ so that $\det(X^j) = \det(X^{j \star g})$.

Corollary 4. *In the unitary case $Y_n(F) \subseteq j \star \mathrm{GL}_n(E')$. In particular, X^j consists of exactly two $G^j(E)$ -orbits. Namely, Γ is of index two in $(E'/E)_1 \cap (E'/F')_1$ and the two $G^j(E)$ -orbits in X^j are $\{x \in X^j : \det x \in \Gamma\} = G^j(E) \cdot I_n$ and $\{x \in X^j : \det x \notin \Gamma\}$.*

Proof. Consider first the case $n = 1$. Note that $Y_1(F) = F^* \subseteq Y_1(E) = E^*$ and $j \star \mathrm{GL}_1(E') = jN(E'/E)$ so that

$$j \star \mathrm{GL}_1(E') \cap Y_1(F) = jN(E'/E) \cap F^* = j(N(E'/E) \cap F^*).$$

It follows from Lemma 5 that $j \star \mathrm{GL}_1(E') \cap Y_1(F) = jF^*$. Note further that $\mathrm{GL}_1(F')$ acts on $j \star \mathrm{GL}_1(E') \cap Y_1(F)$ by $y \star g = yN_{F'/F}(g)$ and since $N(F'/F)$ is of index two in F^* it follows that $j \star \mathrm{GL}_1(E') \cap Y_1(F)$ consists of two $\mathrm{GL}_1(F')$ -orbits.

As a consequence of Lemma 3 we obtain that X^j consists of two $G^j(E)$ -orbits. Note that $j \in F^*$ and $G^j(E) = (E'/E)_1$ acts on $X^j = (E'/E)_1 \cap (E'/F')_1$ by $g \cdot x = g^{1-\sigma}x$. It follows that, indeed, Γ has index two in $(E'/E)_1 \cap (E'/F')_1$.

We now consider a general n . We first show that $j \star \mathrm{GL}_n(E') \cap Y_n(F)$ contains two different $\mathrm{GL}_n(F')$ -orbits. After replacing j with an appropriate element of $j \star \mathrm{GL}_n(F')$, we may assume without loss of generality that j is diagonal. Now let $a \in (E')^*$ be such that $N_{E'/E}(a) = aa^\tau \in F^* \setminus N(F'/F)$ be given by Lemma 5. Then $y = j \star \mathrm{diag}(a, I_{n-1}) \in j \star \mathrm{GL}_n(E') \cap Y_n(F)$ but $y \notin j \star \mathrm{GL}_n(F')$. Since $Y_n(F)$ consists of two $\mathrm{GL}_n(F')$ -orbits it follows that $Y_n(F) = j \star \mathrm{GL}_n(E') \cap Y_n(F)$. By Lemma 3 we get that X^j consists of two G^j -orbits.

It follows from Observations 1 (resp. 2) that $\det(G^j \cdot x) = \Gamma \det x$ (resp. $\det(X^j) = (E'/E)_1 \cap (E'/F')_1$). The lemma follows. \square

4.3.3. *The orthogonal case.* Consider the orthogonal case and let $SX^j = X^j \cap SG^j(E)$. Note that SX^j (and its complement $X^j \setminus SX^j$) are $G^j(E)$ -invariant spaces since the $G^j(E)$ -action preserves determinant. We observe the following refinement of Lemma 3.

Lemma 6. *Let $\mathrm{GL}_n^D(E) = \{g \in \mathrm{GL}_n(E) : \det g \in D\}$ for a subset D of E . Consider the orthogonal case.*

- (1) *The bijection defined by Lemma 3 restricts to a bijection between $SG^j(E)$ -orbits in SX^j and $\mathrm{GL}_n(F)$ -orbits in $j \star \mathrm{GL}_n^F(E) \cap Y_n(F)$ and to a bijection between $SG^j(E)$ -orbits in $X^j \setminus SX^j$ and $\mathrm{GL}_n(F)$ -orbits in $j \star \mathrm{GL}_n^{iF}(E) \cap Y_n(F)$.*
- (2) *Each of SX^j and $X^j \setminus SX^j$ contains at most two $SG^j(E)$ -orbits. For $n = 2$ if $\det j \in -F^{*2}$ then SX^j is a unique $SG^j(E)$ -orbit and if $\det j \in -i^2F^{*2}$ then $X^j \setminus SX^j$ is a unique $SG^j(E)$ -orbit.*

Proof. Note first that $a \in E^*$ satisfies $a^2 \in F^*$ if and only if $a \in F^* \sqcup iF^*$. As a consequence $E^{*2} \cap F^* = F^{*2} \sqcup i^2F^{*2}$ and therefore comparing determinants we obtain the disjoint union

$$j \star \mathrm{GL}_n(E) \cap Y_n(F) = (j \star \mathrm{GL}_n^F(E) \cap Y_n(F)) \sqcup (j \star \mathrm{GL}_n^{iF}(E) \cap Y_n(F)).$$

Let $x \in X^j$ and $z \in \mathrm{GL}_n(E)$ be such that $x = z \cdot I_n$. Recall that $G^j(E) \cdot x$ corresponds to $(j \star z) \star \mathrm{GL}_n(F)$ under the bijection of Lemma 3. Note that if $x \in SX^j$ then $1 = \det x = \det z \bar{z}^{-1}$ and therefore $z \in \mathrm{GL}_n^F(E)$ and if $x \in X^j \setminus SX^j$ then $-1 = \det x = \det z \bar{z}^{-1}$ and therefore $z \in \mathrm{GL}_n^{iF}(E)$. By Lemma 4 every $SG^j(E)$ -orbit is a $G^j(E)$ -orbit. The first part of the lemma follows.

Note further that every element in $j \star \mathrm{GL}_n^F(E) \cap Y_n(F)$ has the fixed discriminant $\det j F^{*2}$ and every element of $j \star \mathrm{GL}_n^{iF}(E) \cap Y_n(F)$ has the fixed discriminant $i^2 \det j F^{*2}$. The second part of the lemma follows from the first part and the classification of $\mathrm{GL}_n(F)$ -orbits in $Y_n(F)$. \square

Proposition 5. (1) *The space SX^j is a unique $SG^j(E)$ -orbit if and only if there is a unique $\mathrm{GL}_n(F)$ -orbit in $Y_n(F)$ with discriminant $\det j F^{*2}$. That is, if and only if either $n = 1$ or $n = 2$ and $\det j \in -F^{*2}$.*

- (2) The space $X^j \setminus SX^j$ is a unique $SG^j(E)$ -orbit if and only if there is a unique $GL_n(F)$ -orbit in $Y_n(F)$ with discriminant $i^2 \det j F^{*2}$. That is, if and only if either $n = 1$ or $n = 2$ and $\det j \in -i^2 F^{*2}$.
- (3) If either $n \geq 3$ or $n = 2$ and $\det j \notin -F^{*2}$ (resp. $n \geq 3$ or $n = 2$ and $\det j \notin -i^2 F^{*2}$) then SX^j (resp. $X^j \setminus SX^j$) consists of exactly two orbits.

Proof. By Lemma 6(1) the number of $SG^j(E)$ -orbits in SX^j equals the number of $GL_n(F)$ -orbits in $j \star GL_n^F(E) \cap Y_n(F)$ and the number of $SG^j(E)$ -orbits in $X^j \setminus SX^j$ equals the number of $GL_n(F)$ -orbits in $j \star GL_n^F(E) \cap Y_n(F)$. Note that

$$GL_n^{iF}(E) = \text{diag}(i, I_{n-1})GL_n^F(E) = GL_n^F(E) \text{diag}(i, I_{n-1}).$$

Replacing j by an appropriate element of $j \star GL_n(F)$ we assume without loss of generality that j is diagonal. When this is the case

$$j \star GL_n^{iF}(E) \cap Y_n(F) = (j \text{diag}(i^2, I_{n-1})) \star GL_n^F(E) \cap Y_n(F).$$

As a consequence, the second part of the proposition follows from the first and Lemma 6(1). The third part follows from the first two and Lemma 6(2). It remains to prove the first part.

When $n = 1$ we have that $SG^j(E) = \{\pm 1\}$ acts trivially on $SX^j = \{1\}$ and the statement is obvious. Consider the case $n = 2$. If $\det j \in -F^{*2}$ the statement follows from Lemma 6(2). Assume that $\det j \notin -F^{*2}$. Without loss of generality assume that j is diagonal and write $j = \text{diag}(a, b)$. Let K be a quadratic extension of F such that $-a^{-1}b = u^2$ for some $u \in K^*$ and ρ the Galois action of K/F . Note that for $\eta = \begin{pmatrix} u & -u \\ 1 & 1 \end{pmatrix} \in GL_2(K)$ we have

$$j \star \eta = \begin{pmatrix} 0 & 2b \\ 2b & 0 \end{pmatrix} \quad \text{and} \quad \eta^{-1}\eta^\rho = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Consider first the case that $-ab \in E^{*2}$ so that $K = E$ and $\rho = \sigma$. Then

$$SG^j(E) = \eta SG^{j \star \eta}(E) \eta^{-1} = \eta \{t_a : a \in E^*\} \eta^{-1}$$

where $t_a = \text{diag}(a, a^{-1})$, $a \in E^*$. For $a \in E^*$ and $g = \eta t_a \eta^{-1}$ we have

$$g\bar{g} = \eta t_{a\bar{a}^{-1}} \eta^{-1}.$$

Hence

$$SX^j = \eta \{t_a : a \in F^*\} \eta^{-1}.$$

Let $a \in E^*$ and $b \in F^*$ and write $g = \eta t_a \eta^{-1} \in SG^j(E)$ and $x = \eta t_b \eta^{-1} \in SX^j$. Then

$$g \cdot x = \eta t_{a\bar{a}b} \eta^{-1}$$

and since $N(E/F)$ is of index two in F^* it follows that SX^j consists of two different $SG^j(E)$ -orbits.

Assume now that $-ab \notin E^{*2}$ and let $L = KE$. Then L/F is a Klein extension. By abuse of notation we continue to denote by ρ the Galois action of L/E and by σ the Galois action of L/K . Note that $\eta^\sigma = \eta$.

It follows from the previous case (with L/E replacing E/F) that

$$SG^j(L) = \eta \{t_a : a \in L^*\} \eta^{-1}.$$

For $a \in L^*$ we have

$$(\eta t_a \eta^{-1})^\rho = \eta t_{a^{-\rho}} \eta^{-1}$$

and therefore

$$SG^j(E) = SG^j(L) \cap \mathrm{GL}_2(E) = \eta \{t_a : a \in (L/E)_1\} \eta^{-1}.$$

For $a \in (L/E)_1$ and $g = \eta t_a \eta^{-1}$ we have

$$g\bar{g} = \eta t_{aa^\sigma} \eta^{-1}$$

and therefore

$$SX^j = \eta \{t_a : a \in (L/E)_1 \cap (L/K)_1\} \eta^{-1}.$$

For $a \in (L/E)_1$ and $b \in (L/E)_1 \cap (L/K)_1$ set $g = \eta t_a \eta^{-1} \in SG^j(E)$ and $x = \eta t_b \eta^{-1} \in SX^j(E)$. It follows that

$$g \cdot x = \eta t_{a^{1-\sigma} b} \eta^{-1}.$$

It follows from Corollary 4 that $\{a^{1-\sigma} : a \in (L/E)_1\}$ is of index two in $(L/E)_1 \cap (L/K)_1$ and therefore SX^j consists of exactly two $SG^j(E)$ -orbits.

Next we show that for $n = 3$, the space $j \star \mathrm{GL}_3^F(E) \cap Y_3(F)$ contains two different $\mathrm{GL}_3(F)$ -orbits. Note that for $a \in F^*$ the map $y \mapsto ay$ defines a $\mathrm{GL}_3(F)$ -equivariant bijection between $j \star \mathrm{GL}_3^F(E) \cap Y_3(F)$ and $(aj) \star \mathrm{GL}_3^F(E) \cap Y_3(F)$. We may therefore assume without loss of generality that $\det j \in -F^{*2}$. Let $u \in F^* \setminus N(E/F)$ and recall that $(u, u^2)_F = -1$. It follows that $y_1 = \mathrm{diag}(1, 1, -1)$ and $y_2 = \mathrm{diag}(u^2, u, -u^2)$ both have discriminant $-F^{*2}$, are not in the same $\mathrm{GL}_3(F)$ -orbit but are in the same $\mathrm{GL}_3(E)$ -orbit. There is therefore $z \in \mathrm{GL}_3(E)$ such that $y_2 = y_1 \star z$ and comparing determinants it follows that $z \in \mathrm{GL}_3^F(E)$. Since $j \star \mathrm{GL}_3(F) = y_i \star \mathrm{GL}_3(F)$ for some $i \in \{1, 2\}$ our claim follows.

Next we show that for $n > 3$ the space $j \star \mathrm{GL}_n^F(E) \cap Y_n(F)$ contains two different $\mathrm{GL}_n(F)$ -orbits. We may again assume without loss of generality that j is diagonal. Write $j = \mathrm{diag}(j_1, j_2)$ where $j_1 \in Y_3(F)$. It follows from the $n = 3$ case that there exists $z \in \mathrm{GL}_3^F(E)$ such that $j_1 \star z \notin j_1 \star \mathrm{GL}_3(F)$. It follows from Witt's cancellation theorem that $j \star \mathrm{diag}(z, I_{n-3}) \notin j \star \mathrm{GL}_n(F)$ while clearly $\mathrm{diag}(z, I_{n-3}) \in \mathrm{GL}_n^F(E)$. All together this shows that for $n \geq 3$ it is always the case that $j \star \mathrm{GL}_n^F(E) \cap Y_n(F)$ contains two different $\mathrm{GL}_n(F)$ -orbits. The proposition now follows from Lemma 6(1). \square

4.3.4. Invariants for the $SG^j(E)$ -orbits in X^j in the orthogonal case. Continue to consider the orthogonal case. It will be convenient to introduce invariants that classify the $SG^j(E)$ -orbits in X^j based on the above results. Let $x \in X^j$, $z \in \mathrm{GL}_n(E)$ such that $x = z \cdot I_n$ and $y = j \star z \in Y_n(F)$. Note that

$$\det y F^{*2} = \begin{cases} \det j F^{*2} & x \in SX^j \\ i^2 \det j F^{*2} & x \in X^j \setminus SX^j \end{cases}$$

and $\mathrm{Hasse}_F(y)$ are invariants of $SG^j(E) \cdot x$ that uniquely determine it. Let

$$\partial^j(x) = \det y F^{*2} \quad \text{and} \quad \hbar^j(x) = \mathrm{Hasse}_F(y).$$

For $x, x' \in X^j$ we have $SG^j(E) \cdot x = SG^j(E) \cdot x'$ if and only if $(\partial^j(x), \hbar^j(x)) = (\partial^j(x'), \hbar^j(x'))$.

5. DISTINCTION FOR CLASSICAL GALOIS PAIRS-STATEMENT OF THE MAIN RESULT

Let $j \in Y_{n_0}(F)$ be anisotropic (that is, for $0 \neq v \in (F')^{n_0}$ we also have ${}^t v^\tau j v \neq 0$). This implies that in the symplectic case $n_0 = 0$, in the unitary case $n_0 \in \{0, 1, 2\}$ and in the orthogonal case $n_0 \in \{0, 1, 2, 3, 4\}$. For every $n \in \mathbb{Z}_{\geq 0}$ let $w_n = (\delta_{i, n+1-j}) \in \mathrm{GL}_n(F)$,

$$j[n] = \begin{pmatrix} & & w_n \\ & j & \\ \varepsilon w_n & & \end{pmatrix}, \quad G_n = G^{j[n]}(E), \quad X_n = X^{j[n]} \quad \text{and} \quad H_n = G^{j[n]}(F).$$

The group G_0 is compact. In the orthogonal case let $G_n^\circ = G_n \cap \mathrm{SL}_{n_0+2n}(E)$ be the special orthogonal group. In order to maintain a unified notation, we set $G_n^\circ = G_n$ in the unitary and symplectic cases. Thus G_n° is a connected reductive p -adic group.

Fix $n \in \mathbb{N}$ and let $G = G_n$, $X = X_n$ and $H = H_n$. For any subset S of G let $S^\circ = S \cap G_n^\circ$. Let $N = n_0 + 2n$ so that $j[n] \in Y_N(F)$ and G is a subgroup of $\mathrm{GL}_N(E')$.

5.1. The standard parabolic subgroups of G° . For a decomposition $\alpha = (N_1, \dots, N_k)$ of N let $Q_\alpha = L_\alpha \rtimes V_\alpha$ be the standard parabolic subgroup of $\mathrm{GL}_N(E')$ of type α with standard Levi subgroup L_α and unipotent radical V_α . That is, Q_α is the unique parabolic subgroup containing all upper-triangular matrices in $\mathrm{GL}_N(E')$ with Levi subgroup

$$L_\alpha = \{\mathrm{diag}(g_1, \dots, g_k) : g_i \in \mathrm{GL}_{N_i}(E'), \quad i = 1, \dots, k\}.$$

For the definition of a parabolic subgroup of G and the following analysis (particularly in the orthogonal case when G is not connected) we refer to [MgVW87, CHAPITRE 1, III 2].

The standard minimal parabolic subgroup of G is $P_0 = M_0 \rtimes U_0$ with Levi part

$$M_0 = \{\mathrm{diag}(a_1, \dots, a_n, h, a_n^{-\tau}, \dots, a_1^{-\tau}) : a_1, \dots, a_n \in (E')^*, \quad h \in G_0\}$$

and unipotent radical $U_0 = V_{(1^{(n)}, n_0, 1^{(n)})} \cap G$ consisting of block upper triangular unipotent matrices. (Here $1^{(n)}$ is the n -tuple $(1, \dots, 1)$.)

Let A be the set of all tuples α of non-negative integers of the form $\alpha = (n_1, \dots, n_k; r)$ where $k, r \in \mathbb{Z}_{\geq 0}$, $n_i \in \mathbb{N}$, $1 \leq i \leq k$ and $n = n_1 + \dots + n_k + r$. When $r = 0$ we will simply write $\alpha = (n_1, \dots, n_k) \in A$ and denote by A_0 the subset of all tuples in A with $r = 0$.

For $m \in \mathbb{N}$ and $g \in \mathrm{GL}_m(E')$ let $g^* = w_m {}^t g^{-\tau} w_m$. For $\alpha = (n_1, \dots, n_k; r) \in A$ let $\iota = \iota_\alpha : \mathrm{GL}_{n_1}(E') \times \dots \times \mathrm{GL}_{n_k}(E') \times G_r \rightarrow G$ be the imbedding

$$\iota(g_1, \dots, g_k; h) = \mathrm{diag}(g_1, \dots, g_k, h, g_k^*, \dots, g_1^*).$$

When $\alpha \in A_0$ we simply write $\iota(g_1, \dots, g_k) = \mathrm{diag}(g_1, \dots, g_k, g_k^*, \dots, g_1^*)$. Denote by M_α the image of ι_α and let

$$U_\alpha = V_{(n_1, \dots, n_k, n_0+2r, n_k, \dots, n_1)} \cap G$$

and $P_\alpha = M_\alpha \rtimes U_\alpha$. The groups P_α , $\alpha \in A$ are parabolic subgroups of G containing P_0 . The map $\alpha \mapsto P_\alpha$ is a bijection from A to a complete set of representatives of the G -conjugacy classes of parabolic subgroups of G . A parabolic subgroup of G is called standard if it is of the form P_α for some $\alpha \in A$.

Excluding the split even orthogonal case (that is, when $F' = F$, $\varepsilon = 1$ and $n_0 = 0$) the map $\alpha \mapsto P_\alpha^\circ$ is a bijection from A to the parabolic subgroups of G° containing P_0° . In order to describe the standard parabolic subgroups of G° in the split even orthogonal case we introduce some further notation.

Consider the split even orthogonal case and let $\kappa = \kappa_n = \iota(I_{n-1}, w_2)$. We set $\iota'_\alpha = \text{Ad}(\kappa) \circ \iota_\alpha$ and let $M'_\alpha = \kappa M_\alpha \kappa$ be the image of ι'_α . When α is clear from the context we also write $\iota' = \iota'_\alpha$. We further write $U'_\alpha = \kappa U_\alpha \kappa$ and $P'_\alpha = \kappa P_\alpha \kappa = M'_\alpha \rtimes U'_\alpha$. Then P'_α is a parabolic subgroup of G containing P_0 . Note that for $\alpha = (n_1, \dots, n_k; r) \in A$ we have $P_\alpha = P'_\alpha$ except for α such that $r = 0$ and $n_k \neq 1$. Furthermore, for $\alpha, \beta \in A$ we have that the two sets $\{P_\beta, P'_\beta\}$ and $\{P_\alpha, P'_\alpha\}$ intersect if and only if $\beta = \alpha$.

A parabolic subgroup of G° is called standard if it contains P_0 . Every standard parabolic subgroup of G° is either of the form P_α° or $(P'_\alpha)^\circ$. However, there are some repetitions. We have $P_\alpha^\circ = (P'_\alpha)^\circ$ except for α such that $r = 0$ and $n_k \neq 1$. Furthermore, if $n_1 + \dots + n_m + 1 = n$ we also have $P_{(n_1, \dots, n_m, 1)}^\circ = P_{(n_1, \dots, n_m; 1)}^\circ$. These cases provide the only possible repetitions.

In order to unify notation, back to the general case, let A° equal A unless we are in the split even orthogonal case in which case we define A° to be the disjoint union of

$$\{\alpha = (n_1, \dots, n_k; r) \in A : r > 1 \text{ or } r = 0, n_k = 1\}$$

and

$$\{[\alpha, i] : \alpha = (n_1, \dots, n_k) \in A_0, n_k \neq 1, i \in \{\pm 1\}\}.$$

For such $[\alpha, i]$ we write

$$P_{[\alpha, i]} = \begin{cases} P_\alpha & i = 1 \\ P'_\alpha & i = -1. \end{cases}, \quad M_{[\alpha, i]} = \begin{cases} M_\alpha & i = 1 \\ M'_\alpha & i = -1. \end{cases} \quad \text{and} \quad U_{[\alpha, i]} = \begin{cases} U_\alpha & i = 1 \\ U'_\alpha & i = -1. \end{cases}$$

The map $a \mapsto P_a^\circ$ is a bijection from A° to the set of standard parabolic subgroups of G° .

5.2. Explication of Corollary 2 for classical symmetric pairs. Let $a \in A^\circ$ and $P = M \rtimes U$ with $M = M_a$ and $U = U_a$. That is, $P^\circ = M^\circ \rtimes U$ is a standard parabolic subgroup of G° . If $a \in A \cap A^\circ$ let $\alpha = a$. Otherwise, write $a = [\alpha, \epsilon]$ where $\alpha = (n_1, \dots, n_k) \in A_0$ and $\epsilon \in \{\pm 1\}$ and set $r = 0$. If either $a \in A \cap A^\circ$ or $\epsilon = 1$ denote by ι_a° the restriction of ι_α to $\text{GL}_{n_1}(E') \times \dots \times \text{GL}_{n_k}(E') \times G_r^\circ$. Otherwise, let $\iota_a^\circ = \iota'_\alpha$. Thus,

$$\iota_a^\circ : \text{GL}_{n_1}(E') \times \dots \times \text{GL}_{n_k}(E') \times G_r^\circ \rightarrow M^\circ$$

is an isomorphism.

Let π_i be a representation of $\text{GL}_{n_i}(E')$, $i \in [1, k]$ and π_0 a representation of G_r° . (If $n_0 = r = 0$, by π_0 we will always mean the trivial representation of the trivial group). Let $\pi = (\pi_1 \otimes \dots \otimes \pi_k \otimes \pi_0) \circ \iota_a^\circ$ be the associated representation of M° . We apply the following standard notation for parabolic induction

$$\pi_1 \times \dots \times \pi_k \rtimes \pi_0 = I_{P^\circ}^{G^\circ}(\pi).$$

Our main result in this section is the characterization of distinction for classical symmetric pairs of representations parabolically induced from cuspidal.

For $\rho \in S_k$ and a subset \mathfrak{c} of $[1, k]$ let

- $I(\rho\mathbf{c}) = \{i \in \mathbf{c} : \rho(i) = i\}$;
- $o(\mathbf{c}) = o_M(\mathbf{c})$ be the number of $i \in \mathbf{c}$ such that n_i is odd;
- $N(\rho\mathbf{c}) = N_M(\rho\mathbf{c}) = \sum_{i \in I(\rho\mathbf{c})} n_i$.

Theorem 2. *With the above notation, let $x \in X^\circ$, π_i an irreducible cuspidal representation of $\mathrm{GL}_{n_i}(E')$, $i \in [1, k]$ and π_0 a cuspidal representation of G_r° . The representation*

$$\pi_1 \times \cdots \times \pi_k \rtimes \pi_0$$

of G° is G_x° -distinguished if and only if there exist an involution $\rho \in S_k$ and a subset \mathbf{c} of $[1, k]$ such that $\rho(\mathbf{c}) = \mathbf{c}$ and $n_{\rho(i)} = n_i$, $i \in [1, k]$, $y_i \in Y_{n_i}(E'/(E')^{\sigma^\tau}, \varepsilon)$, $i \in I(\rho\mathbf{c})$ and $z \in X_r$ such that

$$\begin{array}{ll} \pi_{\rho(i)} \simeq \bar{\pi}_i^\vee & i \notin \mathbf{c}, \rho(i) \neq i \\ \pi_i \text{ is } \mathrm{GL}_{n_i}(F')\text{-distinguished} & i \notin \mathbf{c}, \rho(i) = i \\ \pi_{\rho(i)} \simeq \pi_i^{\tau\sigma} & i \in \mathbf{c}, \rho(i) \neq i \\ \pi_i \text{ is } \mathrm{U}(y_i, E'/(E')^{\sigma^\tau})\text{-distinguished} & i \in \mathbf{c}, \rho(i) = i \\ \pi_0 \text{ is } (G_r^\circ)_z\text{-distinguished} & \end{array}$$

and furthermore

- in the unitary case $(-1)^{o(\mathbf{c})} \det z \prod_{i \in I(\rho\mathbf{c})} \det y_i^{\tau-1} \in \det x \Gamma$ (see (4)). (Here if $n_0 = r = 0$ replace $\det z$ by one.)
- in the orthogonal case the following condition is satisfied:

$$(-1)^{ro(\mathbf{c}) + \binom{N(\rho\mathbf{c})}{2}} (2 \det j)^{o(\mathbf{c})} \prod_{j \in I(w)} \det y_j \in N(E/F)$$

if and only if

$$\frac{\hbar^{j[r]}(z)}{\hbar^{j[r]}(I_{n_0+2r})} = \frac{\hbar^{j[n]}(x)}{\hbar^{j[n]}(I_N)}.$$

(See §4.3.4 for the definition of the invariant $\hbar^{j[r]}$. Here if $n_0 = 0$ replace $\det j$ by one and if $n_0 + 2r = 0$ replace the left hand side of the equality by one.) We further remark that

$$\hbar^{j[n]}(I_N) = \mathrm{Hasse}_F(j[n]) = (\det j, -1)_F^n (-1, -1)_F^{\binom{n}{2}} \mathrm{Hasse}_F(j).$$

5.3. A necessary condition for distinction. As a consequence of Theorem 2 we obtain a necessary condition for distinction that is easier to formulate and is useful for applications.

We continue to assume that $n_1 + \cdots + n_k + r = n$. For $w = \rho\mathbf{c} \in \mathfrak{W}_k$ and a tuple $\pi = (\pi_1, \dots, \pi_k; \pi_0)$ such that π_0 is a representation of G_r° and π_i is a representation of $\mathrm{GL}_{n_i}(E')$ let

$$(5) \quad w\pi = (\pi'_1, \dots, \pi'_k; \pi_0) \quad \text{where} \quad \pi'_i = \begin{cases} \pi_{\rho^{-1}(i)} & i \notin \mathbf{c} \\ \pi_{\rho^{-1}(i)}^* & i \in \mathbf{c}. \end{cases}$$

Corollary 5. *Let $x \in X^\circ$, π_i an irreducible cuspidal representation of $\mathrm{GL}_{n_i}(E')$, $i \in [1, k]$ and π_0 a cuspidal representation of G_r° . If the representation*

$$\pi_1 \times \cdots \times \pi_k \rtimes \pi_0$$

of G° is G_x° -distinguished and $w = \rho\mathbf{c} \in \mathfrak{W}_k^\circ[2, M]$ is given by Theorem 2 then

$$w(\pi_1, \dots, \pi_k; \pi_0) = (\bar{\pi}_1^\vee, \dots, \bar{\pi}_k^\vee; \pi_0).$$

Proof. Write $w\pi = (\pi'_1, \dots, \pi'_k; \pi_0)$ and recall that for an irreducible representation ν of $\mathrm{GL}_m(E')$ we have $\nu^* \simeq (\nu^\vee)^\tau$ by [GK75]. The identity $\pi'_i = \bar{\pi}_i^\vee$ is therefore clear whenever $\rho(i) \neq i$. For $i \notin \mathbf{c}$ such that $\rho(i) = i$ it further follows from [Fli91, Proposition 12] and for $i \in \mathbf{c}$ such that $\rho(i) = i$ it further follows from [FLO12, Theorem 6.1(1)]. \square

6. DISTINCTION FOR CLASSICAL GALOIS PAIRS-PROOF OF THE MAIN RESULT

6.1. A reduction step. We begin the proof of the theorem with a reduction to the case where P is a standard parabolic subgroup of G , that is, either $a \in A$ or $\epsilon = 1$ in the above notation. This step is only relevant to the split even orthogonal case.

Consider the split even orthogonal case and let $\alpha = (n_1, \dots, n_k) \in A_0$. Recall that $\kappa = \iota(I_{n-1}; w_2)$ and for a representation π of G° let π^κ be the representation of G° on the space of π defined by $\pi^\kappa(g) = \pi(\kappa g \kappa^{-1})$. More generally, for a representation π of M_α denote by π^κ the representation of $M'_\alpha = \kappa M_\alpha \kappa^{-1}$ defined similarly.

Lemma 7. (1) For a representation π of G° we have

$$\mathrm{Hom}_{H^\circ}(\pi, \mathbf{1}) = \mathrm{Hom}_{H^\circ}(\pi^\kappa, \mathbf{1}).$$

(2) Let $\alpha \in A_0$ and π a representation of M_α . Then

$$\mathrm{Hom}_{H^\circ}(I_{P_\alpha}^{G^\circ}(\pi), \mathbf{1}) \simeq \mathrm{Hom}_{H^\circ}(I_{P'_\alpha}^{G^\circ}(\pi^\kappa), \mathbf{1}).$$

Proof. The first part is immediate from the fact that κ normalizes H° . The second part follows from the first part and the observation that $I_{P'_\alpha}^{G^\circ}(\pi^\kappa) \simeq I_{P_\alpha}^{G^\circ}(\pi)^\kappa$. Indeed, for $\varphi \in I_{P_\alpha}^{G^\circ}(\pi)$ let $\varphi^\kappa(g) = \varphi(\kappa g \kappa^{-1})$, $g \in G^\circ$. Then $\varphi \mapsto \varphi^\kappa$ realizes this equivalence of representations. \square

We continue with some further notation and preliminaries.

6.2. Admissible orbits. For a standard parabolic subgroup $P = M \rtimes U$ of G we recall some facts on the P° -orbits in X° following [Off17, Section 3]. For $x \in X^\circ$ the P° -orbit $P^\circ \cdot x$ is contained in the Bruhat cell $P^\circ x P^\circ$. By the Bruhat decomposition it corresponds to a double coset in $W_{M^\circ} \backslash W_{G^\circ} / W_{M^\circ}$. Let w be the element of minimal length in this double coset, then w is an involution. We say that $P^\circ \cdot x$ is M° -admissible if w normalizes M° . Intersection with $N_{G^\circ}(M^\circ)$ defines a bijection from M° -admissible P° -orbits in X° to M° -orbits in $X \cap N_{G^\circ}(M^\circ)$. In the sequel we study M° -orbits in $X \cap N_{G^\circ}(M^\circ)$.

6.3. The signed permutation group. Let \mathfrak{W}_k be the signed permutation group on k elements realized as $\mathfrak{W}_k = S_k \rtimes \Xi_k$ where $\Xi_k \simeq (\mathbb{Z}/2\mathbb{Z})^k$ is the group of subsets of the integer interval $[1, k]$ with the operation of symmetric difference and the group S_k of permutations of $[1, k]$ acts naturally on Ξ_k (In \mathfrak{W}_k we have $\rho\mathbf{c}\rho^{-1} = \rho(\mathbf{c})$, $\rho \in S_k$, $\mathbf{c} \in \Xi_k$. That is, conjugation of an element \mathbf{c} in Ξ_k by the permutation ρ is the image of the set \mathbf{c} under

the permutation ρ). Denote by $S_k[2]$ the set of involutions in S_k and by $\mathfrak{W}_k[2]$ the set of involutions in \mathfrak{W}_k . Note that

$$\mathfrak{W}_k[2] = \{\rho\mathbf{c} : \rho \in S_k[2], \mathbf{c} \in \Xi_k, \rho(\mathbf{c}) = \mathbf{c}\}.$$

When we write $w = \rho\mathbf{c} \in \mathfrak{W}_k$ we always mean that $\rho \in S_k$ and $\mathbf{c} \in \Xi_k$.

6.4. The split even orthogonal case. Some of the arguments in the sequel will require further justification in the case where $F' = F$, $n_0 = 0$ and $\varepsilon = 1$ henceforth the split even orthogonal case.

In the split even orthogonal case we define the Weyl group W_G of the non-connected group G as $W_G = N_G(T)/T$. Clearly W_{G° is a subgroup of W_G . Note that $T = \{\iota(a_1, \dots, a_n) : a_i \in F^*, i \in [1, n]\}$ is a maximal split torus of G° contained in M_0 and let $e_i \in X^*(T)$ be defined by $e_i(\iota(a_1, \dots, a_n)) = a_i$, $i \in [1, n]$. The split connected group G° has a root system

$$R(T, G^\circ) = \{\pm(e_i \pm e_j) : 1 \leq i \neq j \leq n\}$$

of type D_n with a basis of simple roots

$$\Delta_0^\circ = \{e_i - e_{i+1} : i \in [1, n-1]\} \sqcup \{e_{n-1} + e_n\}.$$

The set

$$R(T, G)^1 = R(T, G^\circ) \sqcup \{\pm 2e_i : i \in [1, n]\}$$

is a root system of type C_n with basis

$$\Delta_0 = (\Delta_0^\circ \setminus \{e_{n-1} + e_n\}) \sqcup \{2e_n\}.$$

It is easy to see that the action of $N_G(T)$ on T by conjugation identifies W_G as the Weyl group of the root system $R(T, G)$ and in particular, defines an isomorphism

$$W_G \simeq \mathfrak{W}_n.$$

In particular, W_G admits a length function with respect to the elementary reflections associated with the simple roots in Δ_0 .

For the rest of this section let $\alpha = (n_1, \dots, n_k; r) \in A \cap A^\circ$, $M = M_\alpha$, $U = U_\alpha$ and $P = M \ltimes U$.

6.5. The set $W_G(M)$ and elementary symmetries. Let $\{s_\alpha : \alpha \in \Delta_{M^\circ}\}$ be the set of elementary symmetries in $W_{G^\circ}(M^\circ)$ and $\ell = \ell_{M^\circ} : W_{G^\circ}(M^\circ) \rightarrow \mathbb{Z}_{\geq 0}$ the length function defined in [MW95, I.1.7].

For the sake of unified notation set $W_G(M) = W_{G^\circ}(M^\circ)$, $\ell_M = \ell_{M^\circ}$, $\Delta_P = \Delta_{P^\circ}$ and $R(A_M, G) = R(A_{M^\circ}, G^\circ)$ except if $r = 0$ in the split even orthogonal case.

Assume that $r = 0$ in the split even orthogonal case and note that $M = M^\circ$. Let $W_G(M)$ be the subset of elements $w \in W_G$ such that w has minimal length in wW_M and wMw^{-1} is a standard Levi subgroup of G (that is, of the form M_β for some $\beta \in A$). Let $R(A_M, G)$ be the set of non-zero restrictions to A_M of the roots in $R(T, G)$ and let Δ_P be the non-zero restrictions to A_M of the roots in Δ_0 . For $\alpha \in R(A_M, G)$ we write $\alpha > 0$ if

¹the notation $R(T, G)$ is somewhat artificial, note that there are no root subgroups for the roots $2e_i$

and recall that $o(\mathbf{c}) = o_M(\mathbf{c})$ is the number of $i \in \mathbf{c}$ such that n_i is odd. Thus, $t_{\mathbf{c},M} \in G^\circ$ except if $o_M(\mathbf{c})$ is odd and $r = 0$ in the split even orthogonal case in which case $t_{\mathbf{c},M} \in G \setminus G^\circ$. Finally, for $w = \rho\mathbf{c} \in \mathfrak{W}_k$ let

$$t_w = t_{w,M} = t_{\rho,M}t_{\mathbf{c},M}.$$

Note that $t_w \in N_G(T) \cap H$ and in fact $t_w \in G^\circ$ except if $o_M(\mathbf{c})$ is odd and $r = 0$ in the split even orthogonal case. Furthermore, t_w represents an element in $W_G(M)$ (that is, $t_w M_0 \in W_G(M)$) and for $g_i \in \mathrm{GL}_{n_i}(E')$, $i \in [1, k]$, $h \in G_r$ and $m = \iota(g_1, \dots, g_k; h) \in M$ we have

$$(6) \quad t_w m t_w^{-1} = \iota(g'_1, \dots, g'_k; h') \quad \text{where} \quad g'_i = \begin{cases} g_{\rho^{-1}(i)} & i \notin \mathbf{c} \\ g_{\rho^{-1}(i)}^* & i \in \mathbf{c} \end{cases} \quad \text{and} \quad h' = \eta_r^{o(\mathbf{c})} h \eta_r^{-o(\mathbf{c})}.$$

In particular, $t_w M t_w^{-1} = M_{w\alpha}$ where $w\alpha = (n_{\rho^{-1}(1)}, \dots, n_{\rho^{-1}(k)}; r)$. Note further that for $w_1, w_2 \in \mathfrak{W}_k$, $t_{w_1} = t_{w_1, M_{w_2\alpha}}$, $t_{w_2} = t_{w_2, M}$ and $t_{w_1 w_2} = t_{w_1 w_2, M}$ we have

$$t_{w_1} t_{w_2} m (t_{w_1} t_{w_2})^{-1} = t_{w_1 w_2} m t_{w_1 w_2}^{-1}, \quad m \in M.$$

That is, $t_{w_1 w_2}^{-1} t_{w_1} t_{w_2}$ lies in the center of M and in particular in M_0 .

Let s_1, \dots, s_k be the standard simple reflections in \mathfrak{W}_k , that is, $s_i = (i, i+1) \in S_k$, $i \in [1, k-1]$ and $s_k = \{k\} \in \Xi_k$. Note that t_{s_1}, \dots, t_{s_k} are representatives of the elementary symmetries in $W_G(M)$. It is now a simple consequence of the results in [MW95, I.1.7 and I.1.8] and the discussion in §6.5 that the map $w \mapsto t_w M_0 : \mathfrak{W}_k \rightarrow W_G(M)$ is bijective. We denote by $j_M : W_G(M) \rightarrow \mathfrak{W}_k$ its inverse. For $w \in W_G(M)$ and $w' \in W_G(M')$ where $M' = w M w^{-1}$ we have

$$j_M(w'w) = j_{M'}(w')j_M(w).$$

6.6.1. In this paragraph assume that $r = 0$ in the even orthogonal case. Note that neither of $W_G(M)$ and $W_{G^\circ}(M^\circ)$ is necessarily contained in the other. In fact, it is easy to see that

$$W_G(M) \cap W_{G^\circ}(M^\circ) = \{w \in W_G(M) : j_M(w) = \rho\mathbf{c} \text{ where } o(\mathbf{c}) \text{ is even}\}.$$

Furthermore, clearly $W_{G^\circ}(M^\circ, M^\circ) \subseteq W_G(M) \cap W_{G^\circ}(M^\circ)$ and we conclude that

$$W_{G^\circ}(M^\circ, M^\circ) = \{w \in W_G(M) : j_M(w) = \rho\mathbf{c} \text{ where } o(\mathbf{c}) \text{ is even and } n_{\rho(i)} = n_i, i \in [1, k]\}.$$

In particular, $t_{j_M(w)} \in G^\circ$, $w \in W_{G^\circ}(M^\circ, M^\circ)$.

6.7. **Orbits and stabilizers in $X \cap N_{G^\circ}(M^\circ)$.** Let

$$\mathfrak{W}_k[2, M] = \{w = \rho\mathbf{c} \in \mathfrak{W}_k[2] : n_{\rho(i)} = n_i, i \in [1, k]\}.$$

We begin with two lemmas that consist of simple computations that allow us to compute M° -orbits and stabilizers in $X \cap N_{G^\circ}(M^\circ)$.

Lemma 8. *Let $w = \rho\mathbf{c} \in \mathfrak{W}_k[2, M]$ and $m = \iota(g_1, \dots, g_k; h) \in M$ where $g_i \in \mathrm{GL}_{n_i}(E')$, $i \in [1, k]$ and $h \in G_r$. Then*

$$t_w m \overline{t_w m} = \iota(a_1, \dots, a_k; b) \quad \text{where} \quad a_i = \begin{cases} g_{\rho(i)} \bar{g}_i & i \notin \mathbf{c} \\ \varepsilon g_{\rho(i)}^* \bar{g}_i & i \in \mathbf{c} \end{cases} \quad \text{and} \quad b = \eta_r^{o(\mathbf{c})} h \eta_r^{o(\mathbf{c})} \bar{h}.$$

In particular, $t_w m \in X$ if and only if

$$\begin{aligned} g_{\rho(i)} \bar{g}_i &= I_{n_i} & i \notin \mathfrak{c}, \rho(i) \neq i \\ g_i &\in Z_{n_i}(E) & i \notin \mathfrak{c}, \rho(i) = i \\ g_{\rho(i)}^* \bar{g}_i &= \varepsilon I_{n_i} & i \in \mathfrak{c}, \rho(i) \neq i \\ w_{n_i} \bar{g}_i &\in Y_{n_i}(E'/(E')^{\sigma\tau}, \varepsilon) & i \in \mathfrak{c}, \rho(i) = i \\ \eta_r^{\circ(\mathfrak{c})} h &\in X_r. \end{aligned}$$

Proof. Note that $t_w \in N_G(M)$ and recall that $\bar{t}_w = t_w$. It follows from the above discussion that t_w^2 lies in the center of M and therefore $t_w m \bar{t}_w m = t_w^2 (t_w m t_w^{-1}) \bar{m}$. In fact, it is easy to compute explicitly that

$$t_w^2 = \iota(u_1, \dots, u_k; I_{n_0+2r}) \quad \text{where} \quad u_i = \begin{cases} I_{n_i} & i \notin \mathfrak{c} \\ \varepsilon I_{n_i} & i \in \mathfrak{c}. \end{cases}$$

Combined with (6) the lemma follows. \square

Lemma 9. *Let $w = \rho\mathfrak{c} \in \mathfrak{W}_k[2, M]$ and $m = \iota(g_1, \dots, g_k; h)$, $x = \iota(x_1, \dots, x_k, y) \in M$ where $g_i, x_i \in \mathrm{GL}_{n_i}(E')$, $i \in [1, k]$ and $h, y \in G_r$. Then*

$$m t_w x \bar{m}^{-1} = t_w \iota(a_1, \dots, a_k; b) \quad \text{where} \quad a_i = \begin{cases} g_{\rho(i)} x_i \bar{g}_i^{-1} & i \notin \mathfrak{c} \\ g_{\rho(i)}^* x_i \bar{g}_i^{-1} & i \in \mathfrak{c} \end{cases} \quad \text{and} \quad b = \eta_r^{\circ(\mathfrak{c})} h \eta_r^{-\circ(\mathfrak{c})} y \bar{h}^{-1}.$$

Proof. Since t_w^2 is in the center of M we have $m t_w = t_w^2 m t_w^{-1}$ and therefore $m t_w x \bar{m}^{-1} = t_w (t_w m t_w^{-1}) x \bar{m}^{-1}$. The lemma is therefore straightforward from (6). \square

If $r = 0$ in the split even orthogonal case let

$$\mathfrak{W}_k^\circ[2, M] = \{w = \rho\mathfrak{c} \in \mathfrak{W}_k[2, M] : \circ(\mathfrak{c}) \text{ is even}\}.$$

For the sake of unified notation set $\mathfrak{W}_k^\circ[2, M] = \mathfrak{W}_k[2, M]$ otherwise.

Recall that for $w = \rho\mathfrak{c} \in \mathfrak{W}_k[2]$ we have

$$I(w) = \{i \in \mathfrak{c} : \rho(i) = i\}.$$

For $w = \rho\mathfrak{c} \in \mathfrak{W}_k^\circ[2, M]$, $y_i \in Y_{n_i}(E'/(E')^{\sigma\tau}, \varepsilon)$, $i \in I(w)$ and $z \in X_r$ let

$$(7) \quad x_w(\{y_i\}_{i \in I(w)}, z) = t_w \iota(u_1, \dots, u_k; \eta_r^{\circ(\mathfrak{c})} z) \quad \text{where} \quad u_i = \begin{cases} I_{n_i} & i \notin I(w) \\ w_{n_i} \bar{y}_i & i \in I(w). \end{cases}$$

Proposition 6. (1) *The disjoint M° -orbit decomposition of $X \cap N_{G^\circ}(M^\circ)$ is given by*

$$X \cap N_{G^\circ}(M^\circ) = \sqcup_{w \in \mathfrak{W}_k^\circ[2, M]} \sqcup_{\{y_i\}_{i \in I(w)}, z} M^\circ \cdot x_w(\{y_i\}_{i \in I(w)}, z)$$

where y_i ranges over a choice of representatives for the two $\mathrm{GL}_{n_i}(E')$ -orbits in $Y_{n_i}(E'/(E')^{\sigma\tau}, \varepsilon)$, $i \in I(w)$ and z ranges over a set of representatives for the (at most two) G_r° -orbits in

$$X_r \cap \eta_r^{\circ(\mathfrak{c})} G_r^\circ = \begin{cases} X_r \setminus X_r^\circ & \circ(\mathfrak{c}) \text{ is odd and either } n_0 > 0 \text{ or } r > 1 \text{ in the orthogonal case} \\ X_r^\circ & \text{otherwise.} \end{cases}$$

(2) In particular, for $w \in \mathfrak{W}_k^\circ[2, M]$ the number of M° -orbits in $X \cap t_w M^\circ$ is $2^{|I(w)| + \delta_{w, M}}$ where $\delta_{w, M} \in \{0, 1\}$ is determined as follows:

- In the symplectic case $\delta_{w, M} = 0$.
- In the unitary case $\delta_{w, M} = \begin{cases} 1 & n_0 + 2r > 0 \\ 0 & n_0 = r = 0. \end{cases}$
- In the orthogonal case $\delta_{w, M} = 0$ if
 - $n_0 \in \{0, 1\}$ and $r = 0$ or
 - $n_0 = 2$, $r = 0$ and $\det j$ is in $-F^{*2}$ if $\mathfrak{o}(\mathfrak{c})$ is even and in $-i^2 F^{*2}$ if $\mathfrak{o}(\mathfrak{c})$ is odd.

Otherwise, $\delta_{w, M} = 1$.

(3) The stabilizer of $x_w(\{y_i\}_{i \in I(w)}, z)$ in M° consists of elements $\iota(g_1, \dots, g_k; h)$ such that

$$\begin{aligned} g_{\rho(i)} &= \bar{g}_i \in \mathrm{GL}_{n_i}(E') & i \notin \mathfrak{c}, \rho(i) \neq i \\ g_i &\in \mathrm{GL}_{n_i}(F') & i \notin \mathfrak{c}, \rho(i) = i \\ g_{\rho(i)}^* &= \bar{g}_i \in \mathrm{GL}_{n_i}(E') & i \in \mathfrak{c}, \rho(i) \neq i \\ \bar{g}_i &\in \mathrm{U}(y_i, (E'/(E')^{\sigma\tau})) & i \in \mathfrak{c}, \rho(i) = i \\ h &\in (G_r^\circ)_z. \end{aligned}$$

Proof. Recall that the map $w \mapsto wM^\circ : W_{G^\circ}(M^\circ, M^\circ) \rightarrow N_{G^\circ}(M^\circ)/M^\circ$ is a group isomorphism. It follows from the discussion in §6.2 that if $x \in X \cap N_{G^\circ}(M^\circ)$ then $xM = wM$ where $w \in W_{G^\circ}(M^\circ, M^\circ)$ is an involution. Furthermore, $W_{G^\circ}(M^\circ, M^\circ) \subseteq W_G(M) \cap W_{G^\circ}(M^\circ)$ and $t_{j_M(w)} \in G^\circ$ for $w \in W_{G^\circ}(M^\circ, M^\circ)$ (see §6.6.1 if $r = 0$ in the split orthogonal case). Note that j_M maps the set of involutions in $W_{G^\circ}(M^\circ, M^\circ)$ to $\mathfrak{W}_k^\circ[2, M]$. We therefore have the disjoint union

$$X \cap N_{G^\circ}(M^\circ) = \sqcup_{w \in \mathfrak{W}_k^\circ[2, M]} X \cap t_w M^\circ.$$

The disjoint decomposition

$$X \cap t_w M^\circ = \sqcup_{\{y_i\}_{i \in I(w)}, z} M^\circ \cdot x_w(\{y_i\}_{i \in I(w)}, z)$$

as well as the explication of the stabilizer $M_{x_w(\{y_i\}_{i \in I(w)}, z)}^\circ$ are immediate from Lemmas 8 and 9 and the facts that $Y_m(E'/(E')^{\sigma\tau}, \varepsilon)$ consists of two $\mathrm{GL}_m(E')$ -orbits (see the unitary case in §4.1) and $Z_m(E) = \mathrm{GL}_m(E') \cdot I_m$ (see §4.2.2). The formula for the number of M° -orbits in $X \cap t_w M^\circ$ further follows from the classification of G_r° -orbits in X_r° and in $X_r \setminus X_r^\circ$ (see Corollary 3 in the symplectic case, Corollary 4 in the unitary case and Proposition 5 in the orthogonal case). □

Finally it will be relevant for us to determine, for $x \in X^\circ$, when is the representative (7) in $G^\circ \cdot x$.

Lemma 10. *Let $x \in X^\circ$, $w = \rho\mathfrak{c} \in \mathfrak{W}_k^\circ[2, M]$, $y_i \in Y_{n_i}(E'/(E')^{\sigma\tau}, \varepsilon)$, $i \in I(w)$ and $z \in X_r \cap \eta_r^{\mathfrak{o}(\mathfrak{c})} G_r^\circ$. Set $x_w = x_w(\{y_i\}_{i \in I(w)}, z) \in X^\circ$.*

(1) In the symplectic case $x_w \in G^\circ \cdot x$.

(2) In the unitary case $x_w \in G^\circ \cdot x$ if and only if

$$(-1)^{o(c)} \det z \prod_{i \in I(w)} \det y_i^{\tau-1} \in \det x \Gamma.$$

(Here, if $n_0 = r = 0$ replace $\det z$ by one. See (4) for the definition of Γ .)

(3) In the orthogonal case

- for $x_1, x_2 \in X^\circ$ we have $G^\circ \cdot x_1 = G^\circ \cdot x_2$ if and only if $\hbar^{j[n]}(x_1) = \hbar^{j[n]}(x_2)$;
- $x_w \in G^\circ \cdot x$ if and only if the following condition is satisfied:

$$(-1)^{r o(c) + \binom{N(\rho^c)}{2}} (2 \det j)^{o(c)} \prod_{j \in I(w)} \det y_j \in N(E/F)$$

if and only if

$$\frac{\hbar^{j[r]}(z)}{\hbar^{j[r]}(I_{n_0+2r})} = \frac{\hbar^{j[n]}(x)}{\hbar^{j[n]}(I_N)}.$$

(See §4.3.4 for the definition of the invariant $\hbar^{j[r]}$. Here if $n_0 = 0$ replace $\det j$ by one and if $n_0 + 2r = 0$ replace the left hand side of the equality by one.)

Proof. In the symplectic case the result of the lemma is immediate from Corollary 3. In the unitary case it is immediate from Corollary 4 and the observation that

$$\det x_w(\{y_i\}_{i \in I(w)}; z) = (-1)^{o(c)} \det z \prod_{i \in I(w)} \det y_i^{\tau-1}.$$

The orthogonal case is more elaborate and requires a careful computation of invariants. We point out all the ingredients of our computation.

It follows from Lemma 6 (1) that $\partial^{j[n]}(x_1) = \partial^{j[n]}(x_2)$. The first part immediately follows (see §4.3.4). For the second part, it follows that $x_w \in G^\circ \cdot x$ if and only if $\hbar^{j[n]}(x) = \hbar^{j[n]}(x_w)$. Let $z_w \in \mathrm{GL}_N(E)$ be such that $x_w = z_w \cdot I_N$ and $y_w = j[n] \star z_w$. By definition $\hbar(x_w) = \mathrm{Hasse}_F(y_w)$.

We make the following observations. Let $\zeta_m = \begin{pmatrix} I_m & u I_m \\ I_m & -u I_m \end{pmatrix}$ and $v_m = w_{2m} \star \zeta_m$. We observe that

$$\zeta_m \cdot I_{2m} = \begin{pmatrix} I_m & \\ & I_m \end{pmatrix} \quad \text{and} \quad v_m = \mathrm{diag}(2w_m, -2v^2 w_m).$$

Note further that

$$w_{4m} \star \mathrm{diag}(\zeta_{2m}, \zeta_{2m}) = \begin{pmatrix} & \eta_m \\ \eta_m & \end{pmatrix}$$

and that $\eta_{2m}, \begin{pmatrix} & \eta_m \\ \eta_m & \end{pmatrix} \in w_{4m} \star \mathrm{GL}_{4m}(F) = I_{4m} \star \mathrm{GL}_{4m}(F)$.

For $y \in Y_m(E/F, 1)$ let $x(y) = \begin{pmatrix} & y^{-1} w_m \\ w_m \bar{y} & \end{pmatrix} \in X^{w_{2m}}$, $z(y) = \begin{pmatrix} I_m & w_m \\ w_m \bar{y} & -w_m \bar{y} w_m \end{pmatrix} \in \mathrm{GL}_{2m}(E)$ and $\eta(y) = w_{2m} \star z(y)$. Note that $z(y) \cdot I_{2m} = x(y)$ and therefore $\eta(y) \in Y_{2m}(F/F, 1)$. If we assume further that $y \in Y_m(F/F, 1) = Y_m(E/F, 1) \cap \mathrm{GL}_m(F)$ then

$$\eta(y) = \mathrm{diag}(2y, -2v^2 w_m y w_m).$$

We can deduce that

$$\frac{\det \eta(y)}{\det w_{2m}} \in v^{2m} F^{\star 2}$$

and

$$\frac{\text{Hasse}_F(\eta(y))}{\text{Hasse}_F(w_{2m})} = (\det y, i^2)_F (2, i^2)_F^m (-1, i^2)_F^{\binom{m}{2}}.$$

Indeed, note that $\eta(y) \star \text{GL}_{2m}(F)$ (hence $\det \eta(y) F^{*2}$ and $\text{Hasse}_F(\eta(y))$) depends only on $y \star \text{GL}_m(E)$ and therefore we can replace y with $\text{diag}(I_{m-1}, \det y)$. In this case $\eta(y)$ is diagonal and its invariants are easy to compute. We further recall that $\det w_{2m} = (-1)^m$.

Based on the above observations we have that $\text{diag}(u_1, \dots, u_k, u) \in y_w \star \text{GL}_N(F)$ where

$$u_i = \begin{cases} \eta(y_i) & i \in I(w) \\ I_{2n_i} & \text{otherwise} \end{cases} \quad \text{and} \quad u = j[r] \star \zeta \text{ where } z = \zeta \cdot I_{n_0+2r}.$$

It therefore follows that

$$\frac{\text{Hasse}_F(y_w)}{\text{Hasse}_F(j[n])} = \frac{\hbar^{j[r]}(z)}{\text{Hasse}_F(j[r])} \left[\prod_{i \in I(w)} \frac{(\partial^{j[r]}(z), \det \eta(y_i))_F \text{Hasse}_F(\eta(y_i))}{(\det j[r], \det w_{2n_i})_F \text{Hasse}_F(w_{2n_i})} \right] \times \prod_{\{i, j \in I(w): i < j\}} \frac{(\det \eta(y_i), \det \eta(y_j))_F}{(\det w_{2n_i}, \det w_{2n_j})_F}.$$

Let $o(w) = o_M(w)$ be the number of $i \in I(w)$ such that n_i is odd. If $n_0 + 2r > 0$ then $\det \eta_r = -1$ and therefore $\det z = (-1)^{o(w)}$. It follows (see §4.3.4) that

$$\frac{\partial^{j[r]}(z)}{\det j[r]} \in i^{2o(w)} F^{*2}.$$

It follows from the above computations that

$$\prod_{\{i, j \in I(w): i < j\}} \frac{(\det \eta(y_i), \det \eta(y_j))_F}{(\det w_{2n_i}, \det w_{2n_j})_F} = (-1, i^2)_F^{\binom{o(w)}{2}}$$

and

$$\prod_{i \in I(w)} \frac{\text{Hasse}_F(\eta(y_i))}{\text{Hasse}_F(w_{2n_i})} = (2, i^2)_F^{o(w)} \prod_{j \in I(w)} (-1, i^2)_F^{\binom{n_j}{2}} (\det y_j, i^2)_F.$$

Recall that $N(w) = N_M(w) = \sum_{i \in I(w)} n_i$ and note that

$$\sum_{j \in I(w)} \binom{n_j}{2} \equiv \binom{N(w)}{2} + \binom{o(w)}{2} \pmod{2}.$$

It follows that

$$\prod_{\{i, j \in I(w): i < j\}} \frac{(\det \eta(y_i), \det \eta(y_j))_F}{(\det w_{2n_i}, \det w_{2n_j})_F} \times \prod_{i \in I(w)} \frac{\text{Hasse}_F(\eta(y_i))}{\text{Hasse}_F(w_{2n_i})} = (2, i^2)_F^{o(w)} (-1, i^2)_F^{\binom{N(w)}{2}} \left(\prod_{j \in I(w)} \det y_j, i^2 \right)_F.$$

Recall further that $\det j[r] = (-1)^r \det j$. Hence

$$\prod_{i \in I(w)} \frac{(\partial^{j[r]}(z), \det \eta(y_i))}{(\det j[r], \det w_{2n_i})_F} = (-1, i^2)_F^{ro(w)} (\det j, i^2)_F^{o(w)}.$$

We note further that $o(w) \equiv o(\mathfrak{c}) \pmod{2}$. We conclude that

$$\frac{\text{Hasse}_F(y_w)}{\text{Hasse}_F(J[n])} = ((-1)^r 2 \det J, \iota^2)_F^{o(\mathfrak{c})} (-1, \iota^2)_F^{\binom{N(w)}{2}} \left(\prod_{j \in I(w)} \det y_j, \iota^2 \right)_F \frac{\hbar^{j[r]}(z)}{\text{Hasse}_F(J[r])}.$$

Recall that from the definition of the quadratic Hilbert symbol we have $(a, \iota^2)_F = 1$ if and only if $a \in N(E/F)$. The lemma follows. \square

6.8. The contribution of an admissible orbit. Let

$$\Delta(\text{GL}_m(E')) = \{(g, g) : g \in \text{GL}_m(E')\}$$

be the image of the diagonal imbedding of $\text{GL}_m(E')$ in $\text{GL}_m(E') \times \text{GL}_m(E')$. The following is an immediate consequence of the explication of the stabilizers in Proposition 6 (3).

Corollary 6. *Let $\alpha = (n_1, \dots, n_k; r) \in A$ and $M = M_\alpha$. Let π_i be a representation of $\text{GL}_{n_i}(E')$, $i \in [1, k]$ and π_0 a representation of G_r° . Let $w = \rho\mathfrak{c} \in \mathfrak{W}_k^\circ[2, M]$, $y_i \in Y_{n_i}(E'/(E')^{\sigma\tau}, \varepsilon)$, $i \in I(w)$, $z \in X_r \cap \eta_r^{o(\mathfrak{c})} G_r^\circ$ and $x_w(\{y_i\}_{i \in I(w)}, z) \in X^\circ$ be defined as in (7). The representation $(\pi_1 \otimes \dots \otimes \pi_k \otimes \pi_0) \circ \iota_\alpha$ of M° is $M_{x_w(\{y_i\}_{i \in I(w)}, z)}^\circ$ -distinguished if and only if*

$$\begin{aligned} \pi_i \otimes \bar{\pi}_{\rho(i)} & \text{ is } \Delta(\text{GL}_{n_i}(E'))\text{-distinguished} & i \notin \mathfrak{c}, \rho(i) \neq i \\ \pi_i & \text{ is } \text{GL}_{n_i}(E')\text{-distinguished} & i \notin \mathfrak{c}, \rho(i) = i \\ \pi_i \otimes \bar{\pi}_{\rho(i)}^* & \text{ is } \Delta(\text{GL}_{n_i}(E'))\text{-distinguished} & i \in \mathfrak{c}, \rho(i) \neq i \\ \pi_i & \text{ is } \text{U}(y_i, E'/(E')^{\sigma\tau})\text{-distinguished} & i \in \mathfrak{c}, \rho(i) = i \\ \pi_0 & \text{ is } (G_r^\circ)_z\text{-distinguished.} \end{aligned}$$

\square

Remark 5. For irreducible representations π_1 and π_2 of $\text{GL}_m(E')$ we clearly have that $\pi_1 \otimes \pi_2$ is $\Delta(\text{GL}_{n_i}(E'))$ -distinguished if and only if $\pi_2 \simeq \pi_1^\vee$. Furthermore, for an irreducible representation π of $\text{GL}_m(E')$ we have $\pi^* \simeq (\pi^\tau)^\vee$ (see [GK75]). As a consequence, in the above lemma (and in its notation) if $\rho(i) \neq i$ and π_i and $\pi_{\rho(i)}$ are irreducible then $\pi_i \otimes \bar{\pi}_{\rho(i)}$ is $\Delta(\text{GL}_{n_i}(E'))$ -distinguished if and only if $\pi_{\rho(i)} \simeq \bar{\pi}_i^\vee$ and $\pi_i \otimes \bar{\pi}_{\rho(i)}^*$ is $\Delta(\text{GL}_{n_i}(E'))$ -distinguished if and only if $\pi_{\rho(i)} \simeq \pi_i^{\sigma\tau}$.

6.9. Completion of the proof of Theorem 2. It follows from Lemma 7 that we may assume without loss of generality that $P = P_\alpha$ for some $\alpha = (n_1, \dots, n_k; r) \in A$ (with $r \neq 1$ in the split even orthogonal case). As a consequence of Corollary 2 we have that $\pi_1 \times \dots \times \pi_k \rtimes \pi_0$ is G_x° -distinguished if and only if there exists $x' \in G \cdot x \cap N_{G^\circ}(M^\circ)$ such that $(\pi_1 \otimes \dots \otimes \pi_k \otimes \pi_0) \circ \iota_\alpha$ is $M_{x'}^\circ$ -distinguished. Note that for a representation v of M° and $m \in M^\circ$ we have that $\ell \mapsto \ell \circ v(m)$ is an isomorphism from $\text{Hom}_{M^\circ}, (v, \mathbf{1})$ to $\text{Hom}_{M_{x'}^\circ}, (v, \mathbf{1})$ and in particular, v is $M_{x'}^\circ$ -distinguished if and only if it is $M_{m \cdot x'}^\circ$ -distinguished. The theorem is therefore now immediate from Proposition 6(1), Lemma 10, Corollary 6 and Remark 5. \square

7. DISTINCTION AND CONJUGATE-SELF DUALITY IN THE SPLIT ODD ORTHOGONAL CASE

It was recently proved by Beuzart-Plessis [BP] that for any quasi-split reductive group \mathbb{H} defined over F and irreducible, generic representation π of $\mathbb{H}(E)$ that is $\mathbb{H}(F)$ -distinguished we have $\pi^\sigma = \pi^\vee$.

When \mathbb{H} is the split odd special orthogonal group we apply our previous results in order to prove a global analogue as well as several local generalizations of this result. In particular, we show that in the above statement $\mathbb{H}(F)$ -distinction can be replaced by $\mathbb{H}'(F)$ -distinction where \mathbb{H}' is a non-quasi-split inner form of \mathbb{H} such that $\mathbb{H}'(E) = \mathbb{H}(E)$.

Let $\mathrm{SO}_{2n+1} = \mathrm{SO}(w_{2n+1})$ be the split odd special orthogonal group. Over any field k we have

$$\mathrm{SO}_{2n+1}(k) = \{g \in \mathrm{SL}_{2n+1}(k) : {}^t g w_{2n+1} g = w_{2n+1}\}.$$

The following is well known. Since we do not know a proper reference we include the argument.

- Observation 3.** (1) For a p -adic field F and an irreducible, generic representation π of $\mathrm{SO}_{2n+1}(F)$ we have $\pi^\vee \simeq \pi$.
 (2) For a number field k and an irreducible, generic, cuspidal automorphic representation Π of $\mathrm{SO}_{2n+1}(\mathbb{A}_k)$ we have $\Pi^\vee = \Pi$.

Proof. Consider first the local statement and assume further that π is unramified. Then, π is fully induced from a character χ of the standard Borel subgroup. Since χ^{-1} is in the Weyl orbit of χ it follows that π is selfdual.

In the global setting, this implies that Π_v is selfdual for almost all places v of k . The global statement is therefore immediate from the rigidity theorem of Jiang and Soudry [JS03, Theorem 5.3].

Now a simple globalization argument such as [Sha91, Proposition 5.1] implies that every irreducible, generic and cuspidal representation π of $\mathrm{SO}_{2n+1}(F)$ is selfdual.

For the general local case, in terms of its cuspidal support, π^\vee is a subquotient of a representation induced from an irreducible, cuspidal and generic representation of a standard Levi subgroup. That is, there exists a decomposition $n = n_1 + \cdots + n_k + r$, irreducible cuspidal representations π_i of $\mathrm{GL}_{n_i}(F)$, $i \in [1, k]$ and an irreducible, generic and cuspidal representation π_0 of G_r° such that π^\vee is a subquotient of $\pi_1 \times \cdots \times \pi_k \rtimes \pi_0$. It follows from the cuspidal case that π_0 is selfdual. Thus both π and π^\vee are subquotients of $\pi_1^\vee \times \cdots \times \pi_k^\vee \rtimes \pi_0^\vee$. However, by a well known result of Rodier [Rod73], $\pi_1^\vee \times \cdots \times \pi_k^\vee \rtimes \pi_0^\vee$ has a unique irreducible, generic subquotient and it follows that π is selfdual. \square

Let \mathbb{G} be a semi-simple algebraic group and \mathbb{H} a reductive subgroup of \mathbb{G} both defined over a number field k . It follows from [AGR93, Proposition 1] that the period integral

$$\int_{\mathbb{H}(k) \backslash \mathbb{H}(\mathbb{A}_k)} \phi(h) dh$$

is absolutely convergent for every cuspidal automorphic form ϕ on $\mathbb{G}(\mathbb{A}_k)$. A cuspidal automorphic representation Π of $\mathbb{G}(\mathbb{A}_k)$ is called \mathbb{H} -distinguished if this period integral is non-vanishing for some cusp form ϕ in the space of Π .

Theorem 3. *Let K/k be a quadratic extension of number fields with Galois action σ , $\mathbb{G} = \text{Res}_{K/k}(\text{SO}_{2n+1})$, $\mathbb{X} = \{x \in \mathbb{G} : x^\sigma = x^{-1}\}$ and $g \cdot x = gxg^{-\sigma}$ the action of \mathbb{G} on \mathbb{X} by twisted conjugation. Denote by \mathbb{G}_x the stabilizer of $x \in \mathbb{X}(k)$ in \mathbb{G} , an algebraic group defined over k . Let Π be an irreducible cuspidal generic representation of $\mathbb{G}(\mathbb{A}_K) = \text{SO}_{2n+1}(\mathbb{A}_K)$ that is \mathbb{G}_x -distinguished. Then $\Pi^\sigma \simeq \Pi^\vee$.*

Proof. Write $\Pi = \otimes_v \Pi_v$ where the restricted tensor product is over all places v of k . Then Π_v is $\mathbb{G}_x(k_v)$ -distinguished for all places v of F .

If v is split in K (that is $K_v \simeq k_v \times k_v$) then we may identify $\mathbb{G}(K_v)$ with $\text{SO}_{2n+1}(k_v) \times \text{SO}_{2n+1}(k_v)$ and σ acts on it by interchanging the coordinates. Then $x_v = (x_0, x_0^{-1})$ for some $x_0 \in \text{SO}_{2n+1}(k_v)$ and

$$\mathbb{G}_x(k_v) = \{(g, x_0^{-1}gx_0) : g \in \text{SO}_{2n+1}(k_v)\}.$$

Write $\Pi_v = \pi_1 \otimes \pi_2$ for irreducible representations π_1 and π_2 of $\text{SO}_{2n+1}(k_v)$. Since Π_v is $\mathbb{G}_x(k_v)$ -distinguished we see that $\pi_2 \simeq \pi_1^\vee$, that is, $\Pi_v^\sigma \simeq \Pi_v^\vee$.

Since GL_{2n+1} has trivial Galois cohomology, there exists $z \in \text{GL}_{2n+1}(K)$ such that $x = zz^{-\sigma}$. Note that $y = {}^tzw_{2n+1}z$ is a symmetric matrix in $\text{GL}_{2n+1}(k)$. Since $\det x = 1$ we have $\det y k^{*2} = \det w_{2n+1} k^{*2}$. Since the Hasse invariant of y is one at almost all places it follows that y and w_{2n+1} are in the same $\text{GL}_{2n+1}(k_v)$ -orbit for almost all places v of k . It follows from Lemma 3 that $\mathbb{G}_x(k_v)$ is $\text{SO}_{2n+1}(K_v)$ -conjugate to $\text{SO}_{2n+1}(k_v)$ for almost all places v of k that are inert in K . Let v be a finite and inert place of k such that Π_v is unramified. Since Π is generic we can write $\Pi_v \simeq \chi_1 \times \cdots \times \chi_n \rtimes \mathbf{1}_0$ as irreducibly induced from a character of the Borel subgroup. Thus, $\Pi_v \simeq \chi'_{\rho(1)} \times \cdots \times \chi'_{\rho(n)} \rtimes \mathbf{1}_0$ whenever $\chi'_i \in \{\chi_i, \chi_i^{-1}\}$, $i \in [1, n]$ and $\rho \in S_n$. It now follows from Corollary 5 that $\Pi_v^\sigma = \Pi_v^\vee$.

We conclude that $\Pi_v^\sigma = \Pi_v^\vee$ for almost all places v of k and the theorem now follows from the rigidity theorem of Jiang and Soudry [JS03, Theorem 5.3]. \square

For the rest of this section we go back to the notation of §5 in the split odd orthogonal setting. Thus $G^\circ = \text{SO}_{2n+1}(E)$ and $X^\circ = \{x \in G^\circ : x^{-1} = x^\sigma\}$. Since $X = X^\circ \sqcup (-X^\circ)$ we will use without further mention the fact that $G_x^\circ = G_{-x}^\circ$, $x \in X$.

A representation π of $\text{GL}_m(E)$ is called essentially tempered if $|\det|^{-e} \otimes \pi$ is tempered for some real number e . The number e is uniquely determined by π and we write $e(\pi) = e$. Every irreducible and cuspidal representation of $\text{GL}_m(E)$ is essentially tempered.

Theorem 4. *Let π be an irreducible, generic representation of G° . If π is G_x -distinguished for some $x \in X$ then $\pi^\sigma \simeq \pi^\vee$.*

Proof. If π is cuspidal this is immediate from Theorem 3 and the globalization of distinguished, generic representations in [BP]. We apply the cuspidal case in order to complete the proof of the theorem for a general π .

In terms of its cuspidal support we realize π as a quotient of a representation induced from an irreducible, generic and cuspidal representation of a standard Levi subgroup of G° . That is, there exists a decomposition $n = n_1 + \cdots + n_k + r$, irreducible cuspidal representations π_i of $\text{GL}_{n_i}(E)$, $i \in [1, k]$ and an irreducible, generic and cuspidal representation π_0 of G_r°

such that π^\vee is a quotient of $I = \pi_1 \times \cdots \times \pi_k \rtimes \pi_0$. Hence I is also G_x° -distinguished. Furthermore, π is the unique generic irreducible subquotient of I by [Rod73].

By Theorem 2 the representation π_0 is $(G_r^\circ)_z$ -distinguished for some $z \in X_r$ and by the cuspidal case of this theorem we conclude that $\pi_0^\sigma \simeq \pi_0^\vee$. It further follows from Corollary 5 that $(\pi_1, \dots, \pi_k) = ((\pi'_{\rho(1)})^\sigma, \dots, (\pi'_{\rho(k)})^\sigma)$ for some involution $\rho \in S_k$ and some choice $\pi'_i \in \{\pi_i, \pi_i^\vee\}$, $i \in [1, k]$. Since $(\pi'_{\rho(1)})^\sigma \times \cdots \times (\pi'_{\rho(k)})^\sigma \rtimes (\pi_0^\vee)^\sigma$ has the same decomposition series as $(I^\vee)^\sigma$ we conclude that I and $(I^\vee)^\sigma$ have the same decomposition series. Thus, $(\pi^\vee)^\sigma$ is a generic subquotient of I and therefore $\pi^\sigma = \pi^\vee$. \square

Recall that by generalized injectivity [Han20], if a standard module is generic and reducible then its Langlands quotient is not generic. We now extend our result and show that for a certain family of irreducible, degenerate representations distinction implies Galois invariance.

Proposition 7. *Let I be a standard module of G° of the form $I = \pi_1 \times \cdots \times \pi_s \rtimes T$ where π_i is irreducible cuspidal for $i \in [1, s]$ and such that $e(\pi_1) \geq \cdots \geq e(\pi_s) > 0$ and T is irreducible, generic and has unitary cuspidal support (in particular, T is tempered). If either I or I^\vee is G_x° -distinguished for some $x \in X$ then $I^\sigma = I$. Consequently, if the Langlands quotient of I is G_x° -distinguished for some $x \in X$ then it is Galois invariant.*

Proof. The representation T is the unique generic direct summand of a semi-simple representation of the form $\pi_{s+1} \times \cdots \times \pi_k \rtimes \pi_0$ where π_0 is an irreducible, generic and cuspidal representation of G_r° for some r and π_i is irreducible cuspidal for $i \in [s+1, k]$. Therefore I is a quotient of $\Pi = \pi_1 \times \cdots \times \pi_k \rtimes \pi_0$ and I^\vee is a quotient of Π^\vee .

If I (resp. I^\vee) is G_x° -distinguished then so is Π (resp. Π^\vee). It follows from Theorem 2 that π_0 (resp. π_0^\vee) is $(G_r^\circ)_z$ -distinguished for some $z \in X_r$ and from Theorem 4 and Observation 3 that $\pi_0^\sigma = \pi_0$. It further follows from Corollary 5, the inequalities $e(\pi_1) \geq \cdots \geq e(\pi_s) > 0$ and the equalities $e(\pi_{s+1}) = \cdots = e(\pi_k) = 0$ that we have the following equalities of multi-sets of cuspidal representations

$$\{\pi_1, \dots, \pi_s\} = \{\pi_1^\sigma, \dots, \pi_s^\sigma\} \quad \text{and} \quad \{\pi'_{s+1}, \dots, \pi'_k\} = \{\pi_{s+1}^\sigma, \dots, \pi_k^\sigma\}$$

for some choice of $\pi'_i \in \{\pi_i, \pi_i^\vee\}$, $i \in [s+1, k]$. By well known results on standard modules for the general linear group over E the first identity implies the equality of the induced representations

$$\pi_1 \times \cdots \times \pi_s = \pi_1^\sigma \times \cdots \times \pi_s^\sigma$$

while the second identity combined with the fact that π_0 is Galois invariant imply the equality of the semi-simple representations

$$\pi_{s+1} \times \cdots \times \pi_k \rtimes \pi_0 = \pi_{s+1}^\sigma \times \cdots \times \pi_k^\sigma \rtimes \pi_0^\sigma.$$

Since T^σ is a generic summand of the right hand side we conclude that $T = T^\sigma$. All together, this implies that I is Galois invariant.

The second part of the proposition follows from the first part and the facts that Galois invariance of I implies Galois invariance of its Langlands quotient and G_x° -distinction of the Langlands quotient implies G_x° -distinction of I . \square

Remark 6. The proof of Proposition 7 uses Theorem 4 only in the case where π is cuspidal. In fact, the proposition can be used as an alternative way to reduce Theorem 4 to the cuspidal case as follows. Let π be as in Theorem 4. We freely use the notation of the proof of the proposition. We may realize π^\vee as the unique generic irreducible subquotient of a representation of the form Π . We may now define T to be the unique generic summand of the semi-simple representation $\pi_{s+1} \times \cdots \times \pi_k \times \pi_0$ so the standard module I is defined as in the proposition. Thus, π^\vee is the unique generic irreducible subquotient of I and by generalized injectivity [Han20], π^\vee is a subrepresentation of I . Consequently, π is a quotient of I^\vee and therefore I^\vee is G_x° -distinguished. It therefore follows from the proposition that I is Galois invariant. Hence $(\pi^\vee)^\sigma$ is an irreducible generic subquotient of I and therefore equals π^\vee . It follows that π is Galois invariant.

8. ON DISTINCTION AND LOCAL LANGLANDS CORRESPONDENCE

In this section we identify a certain family of irreducible representations π of the classical group G° (in the notation of §5), parabolically induced from cuspidal, such that their functorial transfer to $\mathrm{GL}_N(E')$ is also irreducibly induced from cuspidal and Galois distinction of π implies Galois distinction of its transfer. We only consider quasi-split classical groups and the family of representations will contain, in particular, all generic principal series representations (that is, irreducibly induced from a character of the Borel subgroup).

The results are simple applications of Theorem 2, however, we make the relation to Prasad's conjectures in [Pra] explicit and this requires lengthy and rather technical computations.

8.1. On Galois distinction for GL_N . For $m = m_1 + \cdots + m_t$ and representations π_i of $\mathrm{GL}_{m_i}(E')$, $i \in [1, t]$ we denote by $\pi_1 \times \cdots \times \pi_t$ the representation of $\mathrm{GL}_m(E')$ parabolically induced from $\pi_1 \otimes \cdots \otimes \pi_t$ (with the usual normalized induction). If π_i is $\mathrm{GL}_{m_i}(F')$ -distinguished for all $i \in [1, t]$ then $\pi_1 \times \cdots \times \pi_t$ is $\mathrm{GL}_m(F')$ -distinguished by a closed orbit argument (see e.g. [Fli92, Proposition 26]). Furthermore, for any representation π of $\mathrm{GL}_m(E')$ of finite length, the representation $\pi^\vee \times \pi^\sigma$ of $\mathrm{GL}_{2m}(E')$ is $\mathrm{GL}_{2m}(F')$ -distinguished by an open orbit argument (see e.g. [Gur15, Proposition 2.3]).

For a representation π of $\mathrm{GL}_m(E')$ and a character χ of $(E')^*$ let $\pi\chi = \pi \otimes (\chi \circ \det)$. Let $\eta_{E'/F'}$ be the unique character of $(F')^*$ with kernel, the index two subgroup $N(E'/F')$, and let η be any extension of $\eta_{E'/F'}$ to a character of $(E')^*$. In particular, $\eta^\sigma = \eta^{-1}$. Clearly, π is $\mathrm{GL}_m(F')$ -distinguished if and only if $\pi\eta$ is $(\mathrm{GL}_m(F'), \eta_{E'/F'} \circ \det)$ -distinguished. Since, with the above notation, $(\pi_1 \times \cdots \times \pi_t)\eta \simeq (\pi_1\eta) \times \cdots \times (\pi_t\eta)$, the closed orbit argument also implies that if π_i is $(\mathrm{GL}_{m_i}(F'), \eta_{E'/F'} \circ \det)$ -distinguished for all $i \in [1, t]$ then $\pi_1 \times \cdots \times \pi_t$ is $(\mathrm{GL}_m(F'), \eta_{E'/F'} \circ \det)$ -distinguished. Similarly, $(\pi^\vee \times \pi^\sigma)\eta \simeq (\pi\eta^{-1})^\vee \times (\pi\eta^{-1})^\sigma$ and therefore, the open orbit argument also implies that for any representation π of $\mathrm{GL}_m(E')$ of finite length, the representation $\pi^\vee \times \pi^\sigma$ of $\mathrm{GL}_{2m}(E')$ is $(\mathrm{GL}_{2m}(F'), \eta_{E'/F'})$ -distinguished. In the next lemma we freely use both the open and the closed orbit argument.

We make the following simple observation that, in conjunction with Theorem 2, will allow us to relate between Galois distinction for a classical group and a general linear group.

Lemma 11. *Let $N = 2(n_1 + \cdots + n_k) + r$, Π_0 a representation of $\mathrm{GL}_r(E')$, π_i an irreducible representation of $\mathrm{GL}_{n_i}(E')$, $i \in [1, k]$ and χ be either the trivial character of $(F')^*$ or $\eta_{E'/F'}$. Assume that*

(1) *the representation*

$$\Pi = \pi_1 \times \cdots \times \pi_k \times \Pi_0 \times (\pi_k^\vee)^\tau \times \cdots \times (\pi_1^\vee)^\tau$$

of $\mathrm{GL}_N(E')$ is irreducible,

(2) *Π_0 is $(\mathrm{GL}_r(F'), \chi \circ \det)$ -distinguished and*

(3) *there exist an involution $\rho \in S_k$ and a subset \mathfrak{c} of $[1, k]$ such that $\rho(\mathfrak{c}) = \mathfrak{c}$ and $n_{\rho(i)} = n_i$, $i \in [1, k]$ and $y_i \in Y_{n_i}(E'/(E')^{\sigma\tau}, \varepsilon)$ for any $i \in \mathfrak{c}$ such that $\rho(i) = i$ such that*

$$\begin{array}{ll} \pi_{\rho(i)} \simeq (\pi_i^\vee)^\sigma & i \notin \mathfrak{c}, \rho(i) \neq i \\ \pi_i \text{ is } (\mathrm{GL}_{n_i}(F'), \chi \circ \det)\text{-distinguished} & i \notin \mathfrak{c}, \rho(i) = i \\ \pi_{\rho(i)} \simeq \pi_i^{\tau\sigma} & i \in \mathfrak{c}, \rho(i) \neq i \\ \pi_i \text{ is } \mathrm{U}(y_i, E'/(E')^{\sigma\tau})\text{-distinguished} & i \in \mathfrak{c}, \rho(i) = i. \end{array}$$

Then Π is $(\mathrm{GL}_N(F'), \chi \circ \det)$ -distinguished.

Proof. Since Π is irreducible we may freely permute its factors. For $i \notin \mathfrak{c}$ such that $\rho(i) \neq i$ we have that

$$\pi_i \times \pi_{\rho(i)} \simeq \pi_i \times (\pi_i^\vee)^\sigma \quad \text{and} \quad (\pi_i^\vee)^\tau \times (\pi_{\rho(i)}^\vee)^\tau \simeq (\pi_i^\vee)^\tau \times \pi_i^{\sigma\tau}.$$

For $i \in \mathfrak{c}$ such that $\rho(i) \neq i$ we have

$$\pi_i \times (\pi_{\rho(i)}^\vee)^\tau \simeq \pi_i \times (\pi_i^\vee)^\sigma.$$

For $i \in \mathfrak{c}$ such that $\rho(i) = i$ it follows from [FLO12, Theorem 6.1(1)] that $\pi_i \simeq \pi_i^{\sigma\tau}$ and therefore that

$$\pi_i \times (\pi_i^\vee)^\tau \simeq \pi_i^{\sigma\tau} \times (\pi_i^\vee)^\tau.$$

By the open orbit argument, it follows that Π can be expressed as a product of representations each of the appropriate $\mathrm{GL}_m(E')$ that is $(\mathrm{GL}_m(F'), \chi \circ \det)$ -distinguished. Therefore, Π is $(\mathrm{GL}_N(F'), \chi \circ \det)$ -distinguished by the closed orbit argument. \square

8.2. L-groups and transfer maps. Recall (see §5) that $G^\circ = \mathbb{H}^\circ(E)$ where \mathbb{H}° is the connected component of the algebraic group $\mathbb{H} = \{g \in \mathrm{GL}_N : {}^t g^\tau j[n]g = j[n]\}$ defined over F . We denote by W_k the Weil group of the non-archimedean local field k and by $W'_k = W_k \times \mathrm{SL}_2(\mathbb{C})$ the Weil-Deligne group.

Let $\mathbb{G}^\circ = \mathbb{H}_E^\circ$ be the base change of \mathbb{H}° to E . Assume for the rest of this work that \mathbb{G}° is quasi-split. The Langlands dual group (L-group) of \mathbb{G}° is $\widehat{\mathbb{G}^\circ}(\mathbb{C}) \rtimes W_E$ with connected component $\widehat{\mathbb{G}^\circ}(\mathbb{C})$. For many applications it suffices to keep track of the action of W_E on $\widehat{\mathbb{G}^\circ}(\mathbb{C})$ and it is convenient to consider a simplified version of the L-group. For instance, if \mathbb{G}° is split then W_E acts trivially on $\widehat{\mathbb{G}^\circ}(\mathbb{C})$ and we only keep track of the connected component.

We denote by ${}^L G^\circ$ the L-group of G° defined as follows. In the symplectic, the odd orthogonal and the split even orthogonal cases, we take ${}^L G^\circ = \widehat{\mathbb{G}^\circ}(\mathbb{C})$. Explicitly, $\widehat{\mathbb{G}^\circ}(\mathbb{C})$

equals $\mathrm{SO}_{2n+1}(\mathbb{C})$, $\mathrm{Sp}_{2n}(\mathbb{C})$ or $\mathrm{SO}_{2n}(\mathbb{C})$ in the symplectic, odd orthogonal or split even orthogonal case respectively. In the non-split but quasi-split even orthogonal case, \mathbb{G}° splits over a quadratic extension Q/E and we set

$${}^L G^\circ = \mathrm{SO}_{2n+2}(\mathbb{C}) \rtimes \mathrm{Gal}(Q/E).$$

Sending the Galois involution $\theta_{Q/E}$ of Q/E to $s = \mathrm{diag}(I_n, w_2, I_n)$ we obtain an identification

$${}^L G^\circ = \mathrm{O}_{2n+2}(\mathbb{C})$$

with the split orthogonal group defined with respect to w_{2n+2} . Hence in all these cases we have a natural imbedding of ${}^L G^\circ$ into $\mathrm{GL}_m(\mathbb{C})$ where $m = m(n)$ equals $2n + 1$ in the symplectic case, $2n$ in the split orthogonal case and $2n + 2$ in the quasi-split non-split orthogonal case.

In the unitary case

$${}^L G^\circ = \mathrm{GL}_N(\mathbb{C}) \rtimes \mathrm{Gal}(E'/E)$$

where the Galois involution τ of E'/E acts on $\mathrm{GL}_N(\mathbb{C})$ by

$$\tau g \tau^{-1} = J_N^t g^{-1} J_N^{-1}$$

where $J_N = \mathrm{diag}(1, -1, \dots, (-1)^{N-1})w_N$. In this case we set $m = m(n) = N$.

We denote by $\Phi(G^\circ)$ the set of L -parameters for G° . These are the $\widehat{\mathbb{G}^\circ}(\mathbb{C})$ -conjugacy classes $[\phi]$ of (admissible) L -homomorphisms $\phi : W'_E \rightarrow {}^L G^\circ$. Similarly, we denote by $\Phi(\mathrm{GL}_m(E'))$ the set of L -parameters of $\mathrm{GL}_m(E')$. That is, the $\mathrm{GL}_m(\mathbb{C})$ -conjugacy classes $[\phi]$ of L -homomorphisms $\phi : W'_{E'} \rightarrow \mathrm{GL}_m(\mathbb{C})$.

Note that in all cases an L -homomorphism $\phi : W'_E \rightarrow {}^L G^\circ$ for G° maps $W'_{E'}$ into $\mathrm{GL}_m(\mathbb{C})$ and therefore defines an L -homomorphism for $\mathrm{GL}_m(E')$. Denote by $I : \Phi(G^\circ) \rightarrow \Phi(\mathrm{GL}_m(E'))$ the map defined by

$$I([\phi]) = [\phi|_{W'_{E'}}].$$

The local Langlands correspondence is a conjectural classification of the irreducible representations of a local reductive group in terms of, so called, enhanced L -parameters. The work of Arthur [Art13] followed by that of Mok [Mok15] establish the local Langlands correspondence for tempered representations of the quasi-split classical groups G° that we consider. In fact, we will need a much weaker version of their result as formulated in [Kal16, Conjecture A]. In conjunction with [ABPS14] or [SZ18] it provides a finite to one map from the set of isomorphism classes of irreducible representations of G° to the set $\Phi(G^\circ)$ of L -parameters of G° with certain natural properties. The fibers of this map form the L -packets for G° . Together with the local Langlands correspondence for general linear groups ([Hen00],[HT01]) this defines a transfer of irreducible representations from G° to $\mathrm{GL}_m(E')$ along the map I . We denote by $T(\pi)$ the transfer of an irreducible representation π of G° . It is the representation of $\mathrm{GL}_m(E')$ with parameter $I([\phi])$ where $[\phi]$ is the parameter of π .

It is known by the work of Arthur, Mok and others (see [Ato17]), that whenever one fixes a non-degenerate character ψ of the unipotent radical of the Borel subgroup of G° , each tempered L -packet contains a unique ψ -generic representation.

8.3. Transfer of induced generic representations. Throughout this section fix a decomposition $n = n_1 + \cdots + n_k + r$ and representations π_0 of G_r° and π_i of $\mathrm{GL}_{n_i}(E')$, $i \in [1, k]$. In the split even orthogonal case, if $r = 0$ we also fix a sign in order to choose one of the two standard parabolic subgroups of type (n_1, \dots, n_k) (see §5.1). Consider the representations

$$\pi = \pi_1 \times \cdots \times \pi_k \rtimes \pi_0$$

of G° and

$$\Pi = \pi_1 \times \cdots \times \pi_k \times T(\pi_0) \times (\pi_k^\vee)^\tau \times \cdots \times (\pi_1^\vee)^\tau.$$

of $\mathrm{GL}_m(E')$ where $m = m(n)$ and the representation $T(\pi_0)$ of $\mathrm{GL}_{m(r)}(E')$ is the transfer of π_0 defined by the local Langlands correspondence.

We will say that an irreducible representation of G° is generic if it has a Whittaker model with respect to a non-degenerate character of the standard maximal unipotent subgroup of G° . By the standard module theorem [Mui01, Theorem 1.1] every irreducible generic representation of G° is of the form π above where π_0 is an irreducible generic tempered representation of G_r° and π_1, \dots, π_k are irreducible essentially square-integrable representations such that $e(\pi_1) \geq \cdots \geq e(\pi_k) > 0$. The standard parabolic subgroup of G° and the data $(\pi_1, \dots, \pi_k; \pi_0)$ are uniquely determined by π . When this is the case, we will show that Π is also irreducible and generic and in fact $\Pi = T(\pi)$.

Lemma 12. *Assume that π_0 is tempered, π_1, \dots, π_k are essentially square-integrable, $e(\pi_1) \geq \cdots \geq e(\pi_k) \geq 0$ and both π and Π are irreducible. Then $\Pi = T(\pi)$.*

Proof. Let $M^\circ \simeq \mathrm{GL}_{n_1}(E') \times \cdots \times \mathrm{GL}_{n_k}(E') \times G_r^\circ$ be the standard Levi subgroup of the standard parabolic subgroup of G° fixed in the discussion above from which π is induced. Denote by ${}^L M^\circ$ the relevant standard Levi subgroup of ${}^L G^\circ$ (cf. [Bor79, §3]) corresponding to M° . Let M' be the standard Levi subgroup of $\mathrm{GL}_{m(n)}(\mathbb{C})$ of type $(n_1, \dots, n_k, m(r), n_k, \dots, n_1)$. The connected component of ${}^L M^\circ$ equals $M' \cap G^\circ$.

By compatibility of the local Langlands correspondence with parabolic induction for tempered representations (as in [Kal16, Conjecture A]) and with the Langlands classification ([ABPS14] or [SZ18]), the L-parameter $[\phi]$ of π is represented by some L-homomorphism

$$\phi : W'_E \rightarrow {}^L M^\circ$$

with image in ${}^L M^\circ$. Then ϕ maps W'_E into the connected component of ${}^L M^\circ$ and in particular into $M' = \mathrm{diag}(\mathrm{GL}_{n_1}(\mathbb{C}), \dots, \mathrm{GL}_{n_k}(\mathbb{C}), \mathrm{GL}_{m(r)}(\mathbb{C}), \mathrm{GL}_{n_k}(\mathbb{C}), \dots, \mathrm{GL}_{n_1}(\mathbb{C}))$ and from the compatibility condition defining an L-homomorphism one checks that

$$I[\phi] = [\phi|_{W'_E}] = [\mathrm{diag}(\phi_1, \dots, \phi_k, (\phi_0)|_{W'_E}, \phi'_k, \dots, \phi'_1)]$$

where $[\phi_i]$ is the L-parameter of π_i , $i \in [0, k]$ and $[\phi'_i]$ the L-parameter of $(\pi_i^\vee)^\tau$, $i \in [1, k]$. That is, $[\phi]$ is the L-parameter for Π . □

Lemma 13. *Assume that π_0 is generic tempered, π_1, \dots, π_k are essentially square-integrable, $e(\pi_1) \geq \cdots \geq e(\pi_k) \geq 0$ and π is irreducible. Then Π is irreducible and generic. In particular $\Pi = T(\pi)$.*

Proof. Note that $T(\pi_0)$ is tempered and in particular unitary. If $e(\pi_1) = 0$ then Π is induced from a unitary representation and is therefore irreducible by a theorem of Bernstein [Ber84]. Otherwise, let l be maximal such that $e(\pi_l) > 0$. Replacing π_0 by $\pi_{l+1} \times \cdots \times \pi_k \rtimes \pi_0$ and k by l we may assume without loss of generality that $l = k$, i.e., that $e(\pi_k) > 0$.

Since π is irreducible, we also have that $\pi'_i \rtimes \pi_0$, $\pi'_i \times \pi'_j$ are irreducible for all $i, j \in [1, k]$ where $\pi'_i \in \{\pi_i, (\pi_i^\vee)^\tau\}$ for $i \in [1, k]$. Since $T(\pi_0)$ is tempered it is of the form

$$T(\pi_0) \simeq \delta_1 \times \cdots \times \delta_t$$

for a unique multi-set $\{\delta_1, \dots, \delta_t\}$ of square-integrable representations. Since the standard module $\pi_i \rtimes \pi_0$ is irreducible it follows from [Mui01, Lemma 1.1 and Theorem 1.1] that

$$L(1, \pi_i^\vee, \pi_0) = L(1, \pi_i^\vee, T(\pi_0)) = \prod_{j=1}^t L(1, \pi_i^\vee, \delta_j) \neq \infty$$

where $L(s, \sigma_1, \sigma_2)$ denotes the local Rankin-Selberg L -function of irreducible representations σ_i of $\mathrm{GL}_{m_i}(E')$, $i = 1, 2$. The first equality can be viewed as a definition of the left hand side and the second follows from [JPSS83, Proposition 8.4]. It follows from [MO18, Lemma 2.3], in its terminology, that δ_j does not precede π_i and since $e(\pi_i) > 0 = e(\delta_j)$ it is also the case that π_i does not precede δ_j . It therefore follows from [Zel80, Theorem 9.7] that $\pi_i \times \delta_j$ is irreducible for every $i \in [1, k]$ and $j \in [1, t]$. It follows from [GGP12, Theorem 8.1] that $(T(\pi_0)^\vee)^\tau \simeq T(\pi_0)$ and consequently $\{\delta_1, \dots, \delta_t\}$ is stable under $\delta \mapsto (\delta^\vee)^\tau$. Hence $\pi_i \times (\delta_j^\vee)^\tau$ and therefore also its conjugate dual $(\pi_i^\vee)^\tau \times \delta_j$ are irreducible. It now follows from [Zel80, Theorem 9.7] that Π is irreducible and generic. The equality $\Pi = T(\pi)$ follows from Lemma 12. □

8.4. A consequence of a conjecture of Prasad. We state a consequence of a conjecture of Prasad for distinguished representations in the context of a Galois pair. The conjecture is formulated in terms of the opposition group $\mathbf{Y}^{\mathrm{op}} = \mathbf{Y}^{\mathrm{op}, E/F}$ defined in [Pra, §5] and a character $\omega_{\mathbf{Y}} = \omega_{\mathbf{Y}, E/F}$ of $\mathbf{Y}(F)$ of order at most two constructed in [Pra, Proposition 6.4] for any connected reductive group \mathbf{Y} defined and quasi-split over F with respect to the quadratic extension E/F .

We begin by explicating \mathbf{Y}^{op} and $\omega_{\mathbf{Y}}$ for groups \mathbf{Y} relevant to this work.

8.4.1. A table for \mathbf{Y}^{op} and $\omega_{\mathbf{Y}, E/F}$. We denote by $U_{m, K/F}$ the quasi-split unitary group defined over F associated with a quadratic extension K/F and defined by w_m . Let $\mathrm{wsn} : U_{m, K/F}(F) \rightarrow K^*/F^*$ be the composition of $\det : U_{m, K/F}(F) \rightarrow (K/F)_1$ with the inverse of the isomorphism $zF^* \mapsto zz^{-\tau}$ from K^*/F^* to $U_{1, K/F}(F)$. (In [Wal59], Wall interpreted this map as a spinor norm analogue.) For the purpose of the table below, K/F will be a quadratic extension such that $K \cap E = F$ so that KE/F is a Klein extension and contains a third quadratic extension of F , that we denote by K' , different than K and E .

We denote by SO_m any special orthogonal group defined and quasi-split over F of the form $\{g \in \mathrm{SL}_m : {}^t g y g = y\}$ for some $y = {}^t y \in \mathrm{GL}_m(F)$. (With this convention SO_2 stands for a rank one torus that is either split over F (isomorphic to GL_1) or splits over a

quadratic extension L of F (isomorphic to $U_{1,L/F}$). We denote by $\text{sn} : \text{SO}_m(F) \rightarrow F^*/F^{*2}$ the spinor norm map.

We also denote by $\eta_{E/F}$ the quadratic character of F^* with kernel $N(E/F)$. We have

\mathbf{Y}	$\omega_{\mathbf{Y}}$	\mathbf{Y}^{op}
GL_m	$\eta_{E/F}^{m-1} \circ \det$	$U_{m,E/F}$
$U_{m,E/F}$	$\mathbf{1}$	GL_m
Sp_{2m}	$\mathbf{1}$	Sp_{2m}
SO_m	$\eta_{E/F}^m \circ \text{sn}$	SO_m
$U_{m,K/F}$	$\eta_{EK/K}^{m-1} \circ \text{wsn}$	$U_{m,K'/F}$

In the sequel, this table is sometimes used without further mention. For the sake of completeness we provide the details of the computation in Appendix A.

8.4.2. *Prasad's conjecture and transfer of distinction for quasi-split classical groups.* Let \mathbf{Y} be an F -quasi-split group. Since $(\mathbf{Y}^{\text{op}})_E \simeq \mathbf{Y}_E$ the L -group ${}^L\mathbf{Y}(E)$ of \mathbf{Y}_E is a subgroup of the L -group ${}^L\mathbf{Y}^{\text{op}}(F)$ of \mathbf{Y}^{op} with the same connected component and for any L -homomorphism $\phi : W'_F \rightarrow {}^L\mathbf{Y}^{\text{op}}(F)$ we have $\phi(W'_E) \subseteq {}^L\mathbf{Y}(E)$. This defines a basechange map

$$\mathbf{BC}_F^E : \Phi(\mathbf{Y}^{\text{op}}(F)) \mapsto \Phi(\mathbf{Y}(E)), \quad \mathbf{BC}_F^E([\phi]) = [\phi|_{W'_E}]$$

from L -parameters of $\mathbf{Y}^{\text{op}}(F)$ to L -parameters of $\mathbf{Y}(E)$.

We say that a representation π of $\mathbf{Y}(E)$ is E/F -generic if it is (U, ψ) -generic for a σ -stable maximal unipotent subgroup U of $\mathbf{Y}(E)$ and a non-degenerate character ψ of U satisfying $\psi^\sigma = \psi^{-1}$ (which is the same as $\psi|_{U^\sigma} \equiv \mathbf{1}$). The following is a part of [Pra, Conjecture 13.3].

Conjecture 1 (Dipendra Prasad). *Assume that the Langlands correspondence is known for $\mathbf{Y}(E)$ and $\mathbf{Y}^{\text{op}}(F)$.*

- (1) *Let π be a $(\mathbf{Y}(F), \omega_{\mathbf{Y},E/F})$ -distinguished irreducible representation of $\mathbf{Y}(E)$. Then the parameter of the L -packet of π belongs to $\mathbf{BC}_F^E(\Phi(\mathbf{Y}^{\text{op}}(F)))$.*
- (2) *Conversely, if the L -packet of an E/F -generic representation π of $\mathbf{Y}(E)$ corresponds to a parameter in $\mathbf{BC}_F^E(\Phi(\mathbf{Y}^{\text{op}}(F)))$ then π is $(\mathbf{Y}(F), \omega_{\mathbf{Y},E/F})$ -distinguished.*

Remark 7. This conjecture is now a theorem when $\mathbf{Y} = \text{GL}_n$. It follows from the results in [Kab04, AKT04, AR05, Hen10, Mat11, Gur15].

Fix a quadratic extension K of F contained in E' (so that E'/K is also quadratic) and let $U_n(E'/K) = U_{n,E'/K}(K)$. We will need the following characterization of the image of

$$\mathbf{BC}_K^{E'} : \Phi(U_n(E'/K)) \rightarrow \Phi(\text{GL}_n(E')).$$

We say that an L -homomorphism

$$\phi_{E'} : W'_{E'} \rightarrow \text{GL}_n(\mathbb{C})$$

(or the L -parameter $[\phi]$) is E'/K -conjugate of sign $\epsilon \in \{\pm 1\}$ (we also say E'/K -conjugate orthogonal if the sign is 1 and E'/K -conjugate symplectic if the sign is -1) if there exists a non-degenerate bilinear form

$$B : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}$$

such that

$$(8) \quad B(\phi_{E'}(w)x, \phi_{E'}(sws^{-1})y) = B(x, y) \quad \text{and} \quad B(x, \phi_{E'}(s^2)y) = \epsilon B(y, x)$$

for $w \in W'_{E'}$, $x, y \in \mathbb{C}^n$ where s is a choice of an element of $W'_K \setminus W'_{E'}$.

Theorem 5. ([GGP12, Theorem 8.1]) *An L -homomorphism $\phi_{E'}$ as above is E'/K -conjugate of sign $(-1)^{n-1}$ if and only if $[\phi_{E'}] \in \mathbf{BC}_K^{E'}(\Phi(\mathbf{U}_n(E'/K)))$.*

We will also need the following slightly more precise result.

Theorem 6. ([Kab04, AKT04, AR05, Hen10, Mat11]) *Let π be a generic irreducible representation of $\mathrm{GL}_n(E')$ with L -parameter $[\phi_\pi]$. Then ϕ_π is E'/K -conjugate symplectic if and only if π is $(\mathrm{GL}_n(K), \eta_{E'/K} \circ \det)$ -distinguished and E'/K -conjugate orthogonal if and only if π is $\mathrm{GL}_n(K)$ -distinguished.*

We are now in a position to show that the first part of Conjecture 1 for the generic spectrum implies a relation between distinction in the classical group and distinction of the transfer.

Proposition 8. *Let $\mathbf{Y} = \mathbb{H}^\circ$ be a quasi-split classical F -group of F -rank n , defined as in §8.2 and assume that Conjecture 1 (1) holds for \mathbf{Y} . If π is an irreducible, generic and $(\mathbf{Y}(F), \omega_{\mathbf{Y}, E/F})$ -distinguished representation of $\mathbf{Y}(E)$ then the representation $T(\pi)$ of $\mathrm{GL}_m(E')$ (for the appropriate $m = m(n)$) is $(\mathrm{GL}_m(F'), \chi)$ -distinguished where χ is the trivial character in the even orthogonal case and equals $\omega_{\mathrm{GL}_m, E'/F'}$ otherwise.*

Proof. Let $[\phi] \in \Phi(\mathbf{Y}(E))$ be the parameter of π so that $I([\phi]) \in \Phi(\mathrm{GL}_m(E'))$ is the parameter of $T(\pi)$. It follows from Lemma 13 that $T(\pi)$ is generic and therefore, by Theorem 6, it suffices to show that $[\phi']$ is E'/F' -conjugate of sign ϵ where ϵ equals one in the even orthogonal case and $(-1)^{m-1}$ otherwise.

In the unitary case $m = n$, $\mathbf{Y}(E) = \mathbf{U}_n(E'/E)$, $\mathbf{Y}^{\mathrm{op}}(F) = \mathbf{U}_n(F''/F')$ where $F'' = (E')^{\sigma\tau}$ and

$$I = \mathbf{BC}_E^{E'} : \Phi(\mathbf{U}_n(E'/E)) \rightarrow \Phi(\mathrm{GL}_n(E'))$$

is the corresponding basechange map. This implies the commutativity of the diagram

$$\begin{array}{ccc} \Phi(\mathbf{U}_n(E'/E)) & \xrightarrow{I} & \Phi(\mathrm{GL}_n(E')) \\ \mathbf{BC}_F^E \uparrow & & \mathbf{BC}_{F'}^{E'} \uparrow \\ \Phi(\mathbf{U}_n(F''/F)) & \xrightarrow{\mathbf{BC}_F^{F'}} & \Phi(\mathbf{U}_n(E'/F')). \end{array}$$

(Both compositions from $\Phi(\mathbf{U}_n(F''/F))$ to $\Phi(\mathrm{GL}_n(E'))$ are defined by restriction to $W'_{E'}$.) By assumption, $[\phi] \in \Phi(\mathbf{U}_n(E'/E))$ lies in the image of \mathbf{BC}_F^E and therefore $I([\phi])$ lies in the image of $\mathbf{BC}_{F'}^{E'}$ and by Theorem 5 it is E'/F' -conjugate of sign $(-1)^{n-1}$.

Consider now the symplectic or orthogonal case. Again, by assumption $[\phi]$ lies in the image of $\mathbf{BC}_F^E : \Phi(\mathbf{Y}^{\text{op}}(F)) \rightarrow \Phi(\mathbf{Y}(E))$. Let $\phi' : W'_F \rightarrow {}^L\mathbf{Y}^{\text{op}}(F)$ be such that

$$[\phi] = I(\mathbf{BC}_F^E([\phi]')) = [\phi'_{|W'_E}].$$

Note that there exists a non-degenerate bilinear form $B : \mathbb{C}^N \times \mathbb{C}^N \rightarrow \mathbb{C}^N$ such that $B(y, x) = \epsilon B(x, y)$, $x, y \in \mathbb{C}^N$ and the image of ϕ' preserves B . Let $s \in W'_F \setminus W'_E$ and define the bilinear form $B' : \mathbb{C}^N \times \mathbb{C}^N \rightarrow \mathbb{C}^N$ by

$$B'(x, y) = B(\phi'(s)x, y).$$

It is now a straightforward verification that B' and ϕ satisfy the conditions (8) and since I is defined via the inclusion ${}^L\mathbf{Y}(E) \subseteq \text{GL}_N(\mathbb{C})$ it follows that $I([\phi])$ is E/F -conjugate of sign ϵ . \square

Remark 8. In the odd orthogonal case we could alternatively argue as in the unitary case using the commutative diagram

$$\begin{array}{ccc} \Phi(\text{SO}_{2n+1}(E)) & \xrightarrow{I} & \Phi(\text{GL}_{2n}(E)) \\ \mathbf{BC}_F^E \uparrow & & \mathbf{BC}_F^E \uparrow \\ \Phi(\text{SO}_{2n+1}(F)) & \xrightarrow{I_F} & \Phi(\text{U}_{2n}(E/F)) \end{array}$$

where I_F is induced by the inclusion $\text{GL}_{2n}(\mathbb{C})^\sigma \subseteq {}^L\text{U}_{2n}(E/F)$ as $\text{GL}_{2n}(\mathbb{C})^\sigma$ is conjugate to $\text{Sp}_{2n}(\mathbb{C})$ in $\text{GL}_{2n}(\mathbb{C})$.

Corollary 7. *Assume that $-1 \in N(E/F)$. In the notation of Proposition 8 assume that Conjecture 1 (1) holds for \mathbf{Y} . If π is a generic irreducible representation of $\mathbf{Y}(E)$ that is $\mathbf{Y}(F)$ -distinguished then $T(\pi)$ is $\text{GL}_{m(n)}(F')$ -distinguished.*

Proof. Taking the formula for $\omega_{\mathbf{Y}, E/F}$ into account, the corollary is exactly the statement of Proposition 8 except in the even unitary and odd orthogonal cases. Consider these two cases now. Let ω be the quadratic character of $\mathbf{Y}(E)$ that extends the character $\omega_{\mathbf{Y}, E/F}$ of $\mathbf{Y}(F)$ given by Lemma 15 (see Remark 10). The representation $\pi \otimes \omega$ is $(\mathbf{Y}(F), \omega_{\mathbf{Y}, E/F})$ -distinguished and since in both cases $m(n)$ is even it follows from Proposition 8 that $T(\pi \otimes \omega)$ is $(\text{GL}_{m(n)}(F'), \eta_{E'/F'} \circ \det)$ -distinguished. By Lemma 16 we have $T(\pi \otimes \omega) = T(\pi) \otimes (\eta_{E'/F'} \circ \det)$ and it therefore follows that $T(\pi)$ is $\text{GL}_{m(n)}(F')$ -distinguished as required. \square

8.5. Transfer of distinction for principal series. Let $\mathbb{H}^\circ = \mathbb{H}_n^\circ$ be a quasi-split classical F -group of F -rank n , defined as in §8.2, and set $H^\circ = H_n^\circ = \mathbb{H}^\circ(F)$ and $G^\circ = G_n^\circ = \mathbb{H}^\circ(E)$.

Our goal in this section is to show that for certain irreducible representations π of G° induced from a cuspidal representation of a Levi subgroup of the form M_α° with $\alpha \in A_0$ (see §5.1), i.e., contained in the Siegel Levi, the implication of Proposition 8 can be proved unconditionally.

We need some further preparation. The following will be used in the non-split quasi-split even orthogonal case. In this case $j \in Y_2(F/F, 1)$ is such that $\text{SO}(j)$ is anisotropic and

splits over a unique quadratic extension $K = F[\sqrt{-\det j}]$ of F . Let α be a character of $\mathrm{SO}(j, E)$.

If $K = E$ then $\mathrm{SO}(j, E) \simeq E^*$. Let λ_α be the character of E^* corresponding to α under this identification. It is a straightforward observation that

$$T(\alpha) = \lambda_\alpha \times \lambda_\alpha^{-1}.$$

Otherwise, let $L = KE$ and $\mathrm{SO}_2(L/E) = \mathrm{SO}(j, E)$. In particular, $\mathrm{SO}_2(L/E) \simeq (L/E)_1 \simeq L^*/E^*$ where the right isomorphism is via Hilbert 90. We denote by λ_α the corresponding character of L^* trivial on E^* . Let $\phi : W'_L \rightarrow \mathbb{C}^*$ be the L -parameter of λ_α . The two dimensional representation $\mathrm{Ind}_{W'_L}^{W'_E}(\phi)$ is therefore a L -homomorphism corresponding to a representation $\pi(\alpha)$ of $\mathrm{GL}_2(E)$. In fact, the representation $\pi(\alpha)$ has a model as the space of α -coinvariants in the Weil representation (see [Mat08, Définition 2.1.1]). According to our convention, ${}^L\mathrm{SO}_2(L/E) = \mathrm{O}_2(\mathbb{C})$ and for the L -homomorphism $\phi_\alpha : W'_E \rightarrow \mathrm{O}_2(\mathbb{C})$ corresponding to α it is easy to see that composing with the inclusion $\mathrm{O}_2(\mathbb{C}) \subseteq \mathrm{GL}_2(\mathbb{C})$ we get that $I(\phi_\alpha) \simeq \mathrm{Ind}_{W'_L}^{W'_E}(\phi)$. That is, $T(\alpha) = \pi(\alpha)$.

Proposition 9. *Let α be a character of G_0° that is $(G_0^\circ)_z$ -distinguished for some $z \in X_0$. Then the representation $T(\alpha)$ of $\mathrm{GL}_{m(0)}(E')$ is $\mathrm{GL}_{m(0)}(F')$ -distinguished.*

Proof. The lemma is trivially true if $m(0) = 0$ (both α and $T(\alpha)$ equal the trivial character of the trivial group). Otherwise, either $m(0) = 1$ in the odd unitary case or $m(0) = 2$ in the quasi-split non split even orthogonal case.

In the first case $G_0^\circ = (E'/E)_1$ and $(G_0^\circ)_z = (F'/F)_1$ is independent of $z \in X_0$. Thus α is a character of $(E'/E)_1$ that is trivial on $(F'/F)_1$ and $T(\alpha)(a) = \alpha(a^{1-\tau})$, $a \in (E')^*$. Since $a^{1-\tau} \in (F'/F)_1$ for $a \in (F')^*$ we conclude that $T(\alpha)$ is indeed $(G_0^\circ)_z$ -distinguished.

In the second case $\mathbb{H}_0^\circ = \mathrm{SO}(j)$. If $-\det j \in E^{*2}$ then as discussed above $T(\alpha) = \lambda_\alpha \times \lambda_\alpha^{-1}$ and $G_0^\circ \simeq E^*$. Under this isomorphism the group $(G_0^\circ)_z$ identifies with F^* if $z \in X_0^\circ$ and with $(E/F)_1$ otherwise. In the first case this means that λ_α (and hence also λ_α^{-1}) is trivial on F^* and in the second case that $\lambda_\alpha^\sigma = \lambda_\alpha$. Consequently, $T(\alpha)$ is $\mathrm{GL}_2(F)$ -distinguished by a closed orbit argument in the first case and by an open orbit argument in the second (see §8.1). If $-\det j \notin E^{*2}$ set $K = F[\sqrt{-\det j}]$ and $L = KE$ and let K' be the third quadratic extension of F contained in L that is different from E and K . As observed above $T(\alpha) = \pi(\alpha)$ and $G_0^\circ \simeq (L/E)_1$. In fact, the full orthogonal group G_0 is isomorphic to the semi-direct product $(L/E)_1 \rtimes \mathbb{Z}_2$ where the non-trivial element of \mathbb{Z}_2 acts on $(L/E)_1$ by the Galois action of L/E . Under this isomorphism $(G_0^\circ)_z$ identifies with $(K/F)_1$ if $z \in X_0^\circ$ and with $(K'/F)_1$ otherwise. In other words, with the above notation, λ_α is trivial on either K^* or $(K')^*$. The fact that for such α the representation $\pi(\alpha)$ is $\mathrm{GL}_2(F)$ -distinguished is proved in [Mat08, Théorème 2.3.3].

□

Let $n = n_1 + \dots + n_k$ and $M^\circ = M_{(n_1, \dots, n_k)}^\circ$ be the corresponding standard Levi subgroup of G° . For $x = x_w(\{y_i\}_{i \in I(w)}, z) \in G^\circ \cdot I_N$ defined by (7) let $\eta \in G^\circ$ be such that $\eta \cdot I_N = x$. Note that $M_x^\circ = M^\circ \cap \eta H^\circ \eta^{-1}$ and therefore the character $\omega_{\mathbb{H}^\circ, E/F} \circ \mathrm{Ad}(\eta^{-1})$ of $\eta H^\circ \eta^{-1}$ restricts to a character of M_x° .

Lemma 14. *Consider either the even unitary or odd orthogonal case. With the above notation let $g_i \in \mathrm{GL}_{n_i}(E')$, $i = 1, \dots, k$ be such that $m = \iota(g_1, \dots, g_k) \in M_x$. We have*

$$\omega_{\mathbb{H}^\circ, E/F}(\eta^{-1}m\eta) = \prod_{i \notin \mathfrak{c}, \rho(i)=i} \eta_{E'/F'}(\det g_i).$$

Proof. Note that in both cases G_0° is the trivial group. In the even unitary case we have

$$\omega_{U_{2n}, F'/F, E/F} = \eta_{E'/F'} \circ \alpha^{-1} \circ \det$$

where $\alpha : (F')^*/F^* \rightarrow (F'/F)_1$ is the isomorphism $aF^* \mapsto a^{1-\tau}$. It follows from Proposition 6 (3) that

$$\det(\eta^{-1}m\eta) = \det m = (aa^\sigma)^{1-\tau} b^{1-\tau} c^{1-\tau}$$

where $a = \prod_{i < \rho(i)} \det g_i \in (E')^*$, $b = \prod_{i \in \mathfrak{c}, \rho(i)=i} \det g_i \in (E'/(E')^{\sigma\tau})_1$ and $c = \prod_{i \notin \mathfrak{c}, \rho(i)=i} \det g_i \in (F')^*$. We have

$$\eta_{E'/F'}(\alpha^{-1}((aa^\sigma)^{1-\tau})) = \eta_{E'/F'}(N_{E'/F'}(a)) = 1$$

and

$$\eta_{E'/F'}(\alpha^{-1}(c^{1-\tau})) = \eta_{E'/F'}(c).$$

It remains to see that $\eta_{E'/F'}(\alpha^{-1}(b^{1-\tau})) = 1$. Let $d \in (E')^*$ be such that $b = d^{1-\sigma\tau}$ (by Hilbert 90). Then $b^{1-\tau} = (dd^\sigma)^{1-\tau}$ and therefore

$$\eta_{E'/F'}(\alpha^{-1}(b^{1-\tau})) = \eta_{E'/F'}(N_{E'/F'}(d)) = 1.$$

The even unitary case follows. In the odd orthogonal case

$$\omega_{\mathrm{SO}_{2n+1}, E/F} = \eta_{E/F} \circ \mathrm{sn}.$$

It follows from Proposition 6 (3) that $m = \iota(g_1, \dots, g_k) \mapsto (g_i)_{i \leq \rho(i)}$ defines an isomorphism

$$M_x \simeq \left[\prod_{i < \rho(i)} \mathrm{GL}_{n_i}(E) \right] \times \left[\prod_{i \notin \mathfrak{c}, \rho(i)=i} \mathrm{GL}_{n_i}(F) \right] \times \left[\prod_{i \in \mathfrak{c}, \rho(i)=i} \mathrm{U}(y_i, E/F) \right].$$

Let $m_i = \iota(h_1, \dots, h_k)$ where $h_j = I_{n_j}$ unless $j \in \{i, \rho(i)\}$ in which case $h_j = g_j$. Then clearly $m = \prod_{i \leq \rho(i)} m_i$ with all m_i 's commuting. Set also $m^\circ = \prod_{i \in \mathfrak{c}, \rho(i)=i} m_i$. We compute separately $\eta_{E/F} \circ \mathrm{sn}(\eta^{-1}m_i\eta)$ for each i such that either $i \notin \mathfrak{c}$ or $\rho(i) \neq i$ as well as $\eta_{E/F} \circ \mathrm{sn}(\eta^{-1}m^\circ\eta)$.

The computation for either $i \notin \mathfrak{c}$ or $\rho(i) \neq i$ is implicitly based on finding η explicitly. More precisely, let

$$x_i = \begin{cases} I_{2n_i} & i \notin \mathfrak{c}, \rho(i) = i \\ \iota \begin{pmatrix} I_{n_i} & \\ & I_{n_i} \end{pmatrix} & i \notin \mathfrak{c}, \rho(i) \neq i \\ \begin{pmatrix} I_{2n_i} & \\ & I_{2n_i} \end{pmatrix} & i \in \mathfrak{c}, \rho(i) \neq i \end{cases} \quad \text{and} \quad \ell_i = \begin{cases} \iota(g_i) & i \notin \mathfrak{c}, \rho(i) = i \\ \iota(g_i, g_i^\sigma) & i \notin \mathfrak{c}, \rho(i) \neq i \\ \iota(g_i, \bar{g}_i^*) & i \in \mathfrak{c}, \rho(i) \neq i. \end{cases}$$

Then $\eta_{E/F} \circ \mathrm{sn}(\eta^{-1}m_i\eta) = \eta_{E/F} \circ \mathrm{sn}(\eta_i^{-1}\ell_i\eta_i)$ where $\eta_i \in \mathrm{SO}_{2k_i}(E)$ is such that $\eta_i \cdot I_{2k_i} = x_i$ and k_i equals n_i if $\rho(i) = i$ and $2n_i$ otherwise. Below we provide an explicit such η_i whenever either $i \notin \mathfrak{c}$ or $\rho(i) \neq i$. This shows that $x \in \mathrm{SO}_{2n+1}(E) \cdot x'$ where $x' = x_{I(w)}(\{y_i\}_{i \in I(w)}, z)$

(here we think of the set of indices $I(w)$ as an element of \mathfrak{M}_k), that is, $x' \in \mathrm{SO}_{2n+1}(E) \cdot I_{2n+1}$. Similarly, set

$$x^\circ = \begin{pmatrix} & & I_t \\ & (-1)^t & \\ I_t & & \end{pmatrix} \iota(w_t \bar{y}) \in X_{2t+1}^\circ$$

where $i_1 < \dots < i_j$ are such that $I(w) = \{i_1, \dots, i_j\}$, $t = n_{i_1} + \dots + n_{i_j}$ and $y = \mathrm{diag}(y_{i_1}, \dots, y_{i_j})$. Applying Lemma 10 to both x' and x° we conclude that $x^\circ = \eta^\circ \cdot I_{2t+1}$ for some $\eta^\circ \in \mathrm{SO}_{2t+1}(E)$ and we then have

$$\eta_{E/F} \circ \mathrm{sn}(\eta^{-1} m^\circ \eta) = \eta_{E/F} \circ \mathrm{sn}((\eta^\circ)^{-1} \iota(g_{i_1}, \dots, g_{i_j}) \eta^\circ).$$

We also note that $\iota(g_{i_1}, \dots, g_{i_j}) \in U(y, E/F)$.

Furthermore, our explicit computations of the spinor norm are based on the fact that any $h \in \mathrm{SO}_m(F)$ can be written as an even product $h = s_{v_1} \dots s_{v_{2k}}$ of orthogonal reflections with respect to anisotropic vectors $v_i \in F^m$ for the quadratic form $q(v) = {}^t v w_m v$, $v \in F^m$ and that in such a situation

$$\mathrm{sn}(h) = q(v_1) \dots q(v_{2k}) F^{*2} \in F^*/F^{*2}.$$

Based on this we have

$$(9) \quad \mathrm{sn}(\iota(h)) = \det h F^{*2}, \quad h \in \mathrm{GL}_m(F).$$

Indeed, any homomorphism from $\mathrm{GL}_m(F)$ to F^*/F^{*2} factors through determinant and therefore $\mathrm{sn}(\iota(h)) = \chi(\det h)$ for some homomorphism $\chi : F^* \rightarrow F^*/F^{*2}$. In order to show that χ is the natural projection it suffices to show that $\mathrm{sn}(\iota(\mathrm{diag}(I_{m-1}, t))) = t F^{*2}$, $t \in F^*$ and this reduces to an $\mathrm{SO}_2(F)$ computation of $\mathrm{sn}(\mathrm{diag}(t, t^{-1}))$. Indeed,

$$(10) \quad \mathrm{diag}(t, t^{-1}) = s_{v(t)} s_{v(1)}$$

where ${}^t v(t) = (t, 1)$ so that $q(v(t)) = 2t$.

If $i \notin \mathfrak{c}$ and $\rho(i) = i$ then we may choose $\eta_i = I_{2n_i}$ and

$$\eta_{E/F}(\mathrm{sn}(\eta^{-1} m_i \eta)) = \eta_{E/F}(\mathrm{sn}(\iota(g_i))) = \eta_{E/F}(\det g_i)$$

by (9).

If $i \notin \mathfrak{c}$ and $i < \rho(i)$ then

$$\eta_{E/F}(\mathrm{sn}(\eta^{-1} m_i \eta)) = \eta_{E/F}(\mathrm{sn}(\iota(\zeta_{n_i}^{-1} \mathrm{diag}(g_i, g_i^\sigma) \zeta_{n_i})))$$

where $\zeta_m = \begin{pmatrix} I_m & \iota I_m \\ I_m & -\iota I_m \end{pmatrix}$ and the right hand side is computed in $\mathrm{SO}_{2n_i}(F)$. (Recall from the proof of Lemma 10 that $\zeta_m \cdot I_{2m} = \begin{pmatrix} I_m \\ I_m \end{pmatrix}$.) Note that $\zeta_{n_i}^{-1} \mathrm{diag}(g_i, g_i^\sigma) \zeta_{n_i} \in \mathrm{GL}_{2n_i}(F)$ and by (9) we see that

$$\eta_{E/F}(\mathrm{sn}(\eta^{-1} m_i \eta)) = \eta_{E/F}(N_{E/F}(\det g_i)) = 1.$$

For the next step we introduce the matrix

$$\xi_m = \begin{pmatrix} \frac{1}{2} I_m & & \iota I_m & \\ & -\frac{1}{2i} I_m & & I_m \\ \frac{1}{2} I_m & & -\iota I_m & \\ & \frac{1}{2i} I_m & & I_m \end{pmatrix} \in \mathrm{SO}_{4m}(E)$$

that satisfies

$$\xi_m \cdot I_{4m} = \begin{pmatrix} & I_{2m} \\ I_{2m} & \end{pmatrix}.$$

If $i \in \mathfrak{c}$ and $i < \rho(i)$ then

$$\eta_{E/F}(\mathrm{sn}(\eta^{-1}m_i\eta)) = \eta_{E/F}(\mathrm{sn}(\xi_{n_i}^{-1}\iota(g_i, \bar{g}_i^*)\xi_{n_i})).$$

By the argument used in the previous steps, in order to show that this equals to one for every $g_i \in \mathrm{GL}_{n_i}(E)$ it suffices to consider $g_i = \mathrm{diag}(I_{n_i-1}, t)$ for all $t \in E^*$. This reduces us to the computation in $\mathrm{SO}_4(F)$ of $\eta_{E/F}(\mathrm{sn}(\xi_1^{-1}\iota(t, t^{-\sigma})\xi_1))$. Note that for $v = {}^t(0, 1, 1, 0)$ the orthogonal reflection $s_v = \mathrm{diag}(1, -w_2, 1)$ and clearly $\mathrm{sn}(\xi_1^{-1}\iota(t, t^{-\sigma})\xi_1) = \mathrm{sn}(s_v\xi_1^{-1}\iota(t, t^{-\sigma})\xi_1s_v)$. Furthermore, $s_v\xi_1^{-1}\iota(t, t^{-\sigma})\xi_1s_v = \iota(a(t))$ where

$$a(t) = \begin{pmatrix} \frac{1}{2}(t+t^\sigma) & \iota(t^\sigma - t) \\ \frac{1}{4i}(t^\sigma - t) & \frac{1}{2}(t+t^\sigma) \end{pmatrix}.$$

Since $\det a(t) = N_{E/F}(t)$ it now follows from (9) that $\eta_{E/F}(\mathrm{sn}(\iota(a(t)))) = 1$ as required.

To conclude the lemma it remains to show that for $y \in Y_m(E/F, 1)$, such that

$$x(y) = \begin{pmatrix} & I_m \\ (-1)^m & \\ I_m & \end{pmatrix} \iota(w_m \bar{y}) \in \mathrm{SO}_{2m+1}(E) \cdot I_{2m+1}$$

and $g \in \mathrm{U}(y, E/F)$ we have

$$(11) \quad \eta_{E/F}(\mathrm{sn}(\eta(y)^{-1}\iota(g)\eta(y))) = 1$$

where $\eta(y) \in \mathrm{SO}_{2m+1}(E)$ is such that $\eta(y) \cdot I_{2m+1} = x(y)$. Indeed, we only need to show this for y of the form $\mathrm{diag}(y_{i_1}, \dots, y_{i_j})$ and g of the form $\mathrm{diag}(g_{i_1}, \dots, g_{i_j})$ as above. However, it is more convenient to show this more general statement.

Assume first that $-1 \in N(E/F)$. Then there is a quadratic character ω of E^* that extends $\eta_{E/F}$ (see Remark 10). Now denoting by sn_E the spinor norm on $\mathrm{SO}_{2m+1}(E)$ the character $\omega \circ \mathrm{sn}_E$ of $\mathrm{SO}_{2m+1}(E)$ is well defined and extends $\eta_{E/F} \circ \mathrm{sn}$. Note further that ω is trivial on $(E/F)_1$. Indeed, ω is trivial on both $N(E/F)$ and E^{*2} and by Hilbert 90 the group $(E/F)_1$ is contained in the product $N(E/F)E^{*2}$ ($z^{\sigma-1} = N_{E/F}(z)z^{-2}$, $z \in E^*$). Therefore,

$$\eta_{E/F}(\mathrm{sn}(\eta(y)^{-1}\iota(g)\eta(y))) = \omega(\mathrm{sn}_E(\eta(y)^{-1}\iota(g)\eta(y))) = \omega(\mathrm{sn}_E(\iota(g))).$$

By (9) this equals $\omega(\det g)$ that equals 1 since $\det g \in (E/F)_1$.

Assume now that $-1 \notin N(E/F)$. Note that if $\eta(y) \cdot I_{2m+1} = x(y)$ and $h \in \mathrm{GL}_m(E)$ then $(\iota(h)\eta(y)) \cdot I_{2m+1} = x({}^t h^{-\sigma} y h^{-1}) = x(y \star h^{-1})$ and $U(y \star h^{-1}, E/F) = hU(y, E/F)h^{-1}$. With our assumption it therefore suffices to show (11) for $y = \mathrm{diag}(a, I_{m-1})$ with $a = \pm 1$ (see §4.1). Fix the sign a and let $x = x(y)$ and $\eta = \eta(y)$. Let e_1, \dots, e_{2m+1} be the standard basis of F^{2m+1} and $v_i = \eta^{-1}e_i$, $i = 1, \dots, 2m+1$. Note that $x^2 = I_{2m+1}$ and $x e_1 = e_{2m+1}$ except when $a = -1$ and $m = 1$ where $x e_1 = -e_{2m+1}$. Consequently,

$$v_1^\sigma = \eta^{-\sigma} e_1 = \eta^{-1} x e_1 = \begin{cases} -v_{2m+1} & a = -1 \text{ and } m = 1 \\ v_{2m+1} & \text{otherwise.} \end{cases}$$

It follows from Lemma 10 (or by direct computation) that for $m = 2$ and $a = 1$ we have $x(y) = w_5 \notin \mathrm{SO}_5(F) \cdot I_5$. We may therefore exclude this case. In all other cases, we claim that $[\mathrm{U}(y, E/F), \mathrm{U}(y, E/F)] = \mathrm{SU}(y, E/F)$. Indeed, for $m = 1$ this is clear and for all other cases (excluding $m = 2$ and $a = 1$) y admits an isotropic vector. It therefore follows from [Die48, Théorème 5] that every proper normal subgroup of $\mathrm{SU}(y, E/F)$ is central. It is easy to see that $[\mathrm{U}(y, E/F), \mathrm{U}(y, E/F)]$ is not central. Consequently, there is a homomorphism $\chi : (E/F)_1 \rightarrow F^*/F^{*2}$ such that $\mathrm{sn} \circ \mathrm{Ad}(\eta^{-1}) \circ \iota = \chi \circ \det$. It therefore suffices to show (11) for $g = \mathrm{diag}(u, I_{m-1}) \in U(y, E/F)$ with $u \in (E/F)_1$. Note that for such g as in (10) we have

$$\iota(g) = s_{ue_1+e_{2m+1}} s_{e_1+e_{2m+1}}.$$

Applying Hilbert 90, let $z \in E^*$ be such that $z^{\sigma^{-1}} = u$. Then $ue_1 + e_{2m+1}$ and $\sigma(z)e_1 + ze_{2m+1}$ are co-linear in E^{2m+1} . Furthermore we have

$$(v_1 + v_{2m+1})^\sigma = \begin{cases} -(v_1 + v_{2m+1}) & a = -1 \text{ and } m = 1 \\ v_1 + v_{2m+1} & \text{otherwise.} \end{cases}$$

and similarly

$$(\sigma(z)v_1 + zv_{2m+1})^\sigma = \begin{cases} -(\sigma(z)v_1 + zv_{2m+1}) & a = -1 \text{ and } m = 1 \\ \sigma(z)v_1 + zv_{2m+1} & \text{otherwise.} \end{cases}$$

It follows that both $w = \alpha(v_1 + v_{2m+1})$ and $w' = \alpha(\sigma(z)v_1 + zv_{2m+1})$ are in F^{2m+1} where

$$\alpha = \begin{cases} i & a = -1 \text{ and } m = 1 \\ 1 & \text{otherwise.} \end{cases}$$

We deduce that

$$\eta^{-1}\iota(g)\eta = \eta^{-1}s_{\sigma(z)e_1+ze_{2m+1}}s_{e_1+e_{2m+1}}\eta = s_{\sigma(z)v_1+zv_{2m+1}}s_{v_1+v_{2m+1}} = s_{w'}s_w$$

and since $q = q \circ \mathrm{Ad}(\eta^{-1})$ we have

$$\mathrm{sn}(\eta^{-1}\iota(g)\eta) = q(w)q(w')F^{*2} = q(\alpha(\sigma(z)e_1+ze_{2m+1}))q(\alpha(e_1+e_{2m+1}))F^{*2} = 4\alpha^4 N_{E/F}(z)F^{*2}.$$

Since $4\alpha^4 \in F^{*2}$, we obtain

$$\eta_{E/F}(\mathrm{sn}(\eta^{-1}\iota(g)\eta)) = \eta_{E/F}(N_{E/F}(z)) = 1$$

as required. \square

For a certain class of generic irreducible representations π of G° we are now in a position to prove unconditionally the relation between distinction of π and of its transfer $T(\pi)$ that Proposition 8 shows follows more generally from Conjecture 1. In particular, π could be any irreducible principal series.

Theorem 7. *In the notation of §8.3 suppose that π_i is cuspidal for $i = 0, \dots, k$ (in particular, π_0 is the trivial character of the trivial group except in the odd unitary or quasi-split but non-split even orthogonal cases where π_0 is a character of G_0°) and π is irreducible. (In particular, π could be any irreducible generic principal series representation of G° .)*

- (1) If π is H° -distinguished then its transfer $T(\pi)$ is $\mathrm{GL}_{m(n)}(F')$ -distinguished.
- (2) If π is $(H^\circ, \omega_{\mathbb{H}^\circ, E/F})$ -distinguished then its transfer $T(\pi)$ is $(\mathrm{GL}_{m(n)}(F'), \chi)$ -distinguished where χ is the trivial character in the even orthogonal case and equals $\omega_{\mathrm{GL}_{m(n)}, E'/F'}$ otherwise.

Proof. Since π is irreducible, applying an appropriate Weyl element we assume without loss of generality that $e(\pi_1) \geq \cdots \geq e(\pi_k) \geq 0$. It follows from Lemma 13 that

$$T(\pi) = \pi_1 \times \cdots \times \pi_k \times T(\pi_0) \times (\pi_k^\vee)^\tau \times \cdots \times (\pi_1^\vee)^\tau.$$

If π is H° -distinguished then it follows from Theorem 2 and (in the odd unitary or quasi-split but non split even orthogonal case) Proposition 9 that $T(\pi)$ satisfies the conditions for Lemma 11 and we deduce that $T(\pi)$ is $\mathrm{GL}_{m(n)}(F')$ -distinguished. Note that we only applied the necessary condition for distinction of Theorem 2.

The second part of the theorem is only different from the first in the odd orthogonal and in the even unitary cases. Assume we are in one of those cases and note that G_0° is now the trivial group. Suppose that π is $(H^\circ, \omega_{\mathbb{H}^\circ, E/F})$ -distinguished. It follows from [Off17, Corollary 5.2 and Corollary 6.9], the orbits and stabilizers analysis done in Section 6.7, Lemma 14 and the fact that $v^* = (v^\tau)^\vee$ for an irreducible representation v of $\mathrm{GL}_m(E')$ (see [GK75]), that there exist an involution $\rho \in S_k$ and a subset \mathfrak{c} of $[1, k]$ such that $\rho(\mathfrak{c}) = \mathfrak{c}$ and $n_{\rho(i)} = n_i$, $i \in [1, k]$ and $y_i \in Y_{n_i}(E'/(E')^{\sigma\tau}, 1)$ for all $i \in \mathfrak{c}$ such that $\rho(i) = i$ such that

$$\begin{array}{ll} \pi_{\rho(i)} \simeq \bar{\pi}_i^\vee & i \notin \mathfrak{c}, \rho(i) \neq i \\ \pi_i \text{ is } (\mathrm{GL}_{n_i}(F'), \eta_{E'/F'} \circ \det)\text{-distinguished} & i \notin \mathfrak{c}, \rho(i) = i \\ \pi_{\rho(i)} \simeq \pi_i^{\tau\sigma} & i \in \mathfrak{c}, \rho(i) \neq i \\ \pi_i \text{ is } \mathrm{U}(y_i, E'/(E')^{\sigma\tau})\text{-distinguished} & i \in \mathfrak{c}, \rho(i) = i. \end{array}$$

It now follows from Lemma 11 that $T(\pi)$ is $(\mathrm{GL}_{m(n)}(F'), \eta_{E'/F'} \circ \det)$ -distinguished. Since in both cases $m(n)$ is even the theorem follows. \square

It is now tempting to suggest that a global analogue of this problem also holds. Let K/k be a quadratic extension of number fields and \mathbb{H}° be a k -quasisplit classical group amongst the ones considered in this paper. Recall from [CPSS11, Theorem 1.1] that for every globally generic cuspidal automorphic representation π of $\mathbb{H}^\circ(\mathbb{A}_K)$ the functorial lift $T(\pi)$ exists and is an automorphic representations of $\mathrm{GL}_{m(n)}(\mathbb{A}_K)$. Furthermore, the definition of Prasad's character $\omega_{\mathbb{H}^\circ, K/k}$ makes sense globally (see A.2 in Appendix A).

Conjecture 2. *Let π be an irreducible globally generic cuspidal automorphic representation of $\mathbb{H}^\circ(\mathbb{A}_K)$ such that $T(\pi)$ is a cuspidal automorphic representation of $\mathrm{GL}_{m(n)}(\mathbb{A}_K)$.*

- (1) If π is $\mathbb{H}^\circ(\mathbb{A}_k)$ -distinguished, then $T(\pi)$ is $\mathrm{GL}_{m(n)}(\mathbb{A}_k)$ -distinguished.
- (2) If π is $(\mathbb{H}^\circ(\mathbb{A}_k), \omega_{\mathbb{H}^\circ, K/k})$ -distinguished, then $T(\pi)$ is $(\mathrm{GL}_{m(n)}(\mathbb{A}_k), \chi_{\mathrm{GL}_{m(n)}, K/k})$ -distinguished where $\chi_{\mathrm{GL}_{m(n)}, K/k}$ is the trivial character in the even orthogonal case and equals $\omega_{\mathrm{GL}_{m(n)}, K/k}$ otherwise.

APPENDIX A. EXPLICIT COMPUTATIONS OF PRASAD'S QUADRATIC CHARACTER AND OPPOSITION GROUP

A.1. Prasad's quadratic character. Let \mathbf{Y} be a connected reductive group defined and quasi-split over F . We recall the construction of a quadratic character $\omega_{\mathbf{Y}, E/F}$ of $Y = \mathbf{Y}(F)$ in [Pra, Proposition 6.4] associated to \mathbf{Y} and the quadratic extension E/F .

The character $\omega_{\mathbf{Y}, E/F}$ factors through the natural projection $Y \mapsto Y^{\text{ad}} = \mathbf{Y}^{\text{ad}}(F)$ to the adjoint group and it suffices to define it for adjoint groups.

Denote by \mathbf{Y}^{sc} the simply connected cover of \mathbf{Y}^{ad} , by \mathbf{Z} its center and by $p : \mathbf{Y}^{\text{sc}} \rightarrow \mathbf{Y}^{\text{ad}}$ the natural projection. Then we have an exact sequence of algebraic F -groups

$$(12) \quad 1 \rightarrow \mathbf{Z} \rightarrow \mathbf{Y}^{\text{sc}} \rightarrow \mathbf{Y}^{\text{ad}} \rightarrow 1$$

that gives rise to an exact sequence of local groups

$$1 \rightarrow \mathbf{Z}(F) \rightarrow \mathbf{Y}^{\text{sc}}(F) \xrightarrow{p} \mathbf{Y}^{\text{ad}}(F) \xrightarrow{\delta_{p,F}} H^1(F, \mathbf{Z}).$$

(The Galois cohomology group $H^1(F, \mathbf{Z})$ and the connecting homomorphism $\delta_{p,F}$ are defined for example in [Ser94].)

Let $\mathbf{B}^{\text{ad}} = \mathbf{T}^{\text{ad}} \ltimes \mathbf{U}$ be a Borel subgroup of \mathbf{Y}^{ad} with a maximal torus \mathbf{T}^{ad} and unipotent radical \mathbf{U} defined over F . Then $p^{-1}(\mathbf{B}^{\text{ad}})$ is a Borel subgroup of \mathbf{Y}^{sc} and $\mathbf{Z} \subseteq \mathbf{T}^{\text{sc}} = p^{-1}(\mathbf{T}^{\text{ad}})$. Thus p identifies \mathbf{U} with its pre-image in \mathbf{Y}^{sc} that we still denote by \mathbf{U} . It is known that the half sum (with multiplicities) ρ of the positive roots of \mathbf{T}^{sc} (with respect to \mathbf{U}) is a weight (i.e. an algebraic character) of \mathbf{T}^{sc} defined over F and that 2ρ is a weight of \mathbf{T}^{ad} , that is, contains \mathbf{Z} in its kernel. By restriction ρ induces a character $\rho|_{\mathbf{Z}} \rightarrow \{\pm 1\}$. By the functorial properties of Galois cohomology ρ gives rise to a homomorphism

$$[\rho]_F : H^1(F, \mathbf{Z}) \rightarrow H^1(F, \{\pm 1\}) \simeq F^*/F^{*2}.$$

Let $\eta_{E/F}$ be the quadratic character of F^* with kernel $(E/F)_1$, the norm one subgroup. Then

$$\omega_{\mathbf{Y}^{\text{ad}}, E/F} = \eta_{E/F} \circ [\rho]_F \circ \delta_{p,F} : Y^{\text{ad}} \rightarrow \{\pm 1\}$$

is the character defined by Prasad for Y^{ad} . Composing with the natural map from Y to Y^{ad} we obtain Prasad's character $\omega_{\mathbf{Y}, E/F} : Y \rightarrow \{\pm 1\}$.

Remark 9. For semi-simple \mathbf{Y} the character $\omega_{\mathbf{Y}, E/F}$ can be defined more directly since the projection $p : \mathbf{Y}^{\text{sc}} \rightarrow \mathbf{Y}^{\text{ad}}$ factors through \mathbf{Y} . Let $p' : \mathbf{Y}^{\text{sc}} \rightarrow \mathbf{Y}$ be the corresponding projection (so that p is the composition of p' with the natural projection $\mathbf{Y} \rightarrow \mathbf{Y}^{\text{ad}}$). Then the short exact sequence

$$1 \rightarrow \mathbf{Z}' \rightarrow \mathbf{Y}^{\text{sc}} \xrightarrow{p'} \mathbf{Y} \rightarrow 1$$

gives rise to the connecting homomorphism $\delta_{p',F} : \mathbf{Y}(F) \rightarrow H^1(F, \mathbf{Z}')$. The kernel \mathbf{Z}' of p' is a subgroup of \mathbf{Z} so that $\rho|_{\mathbf{Z}'} : \mathbf{Z}' \rightarrow \{\pm 1\}$ defines the map $[\rho]_F : H^1(F, \mathbf{Z}') \rightarrow H^1(F, \{\pm 1\})$ and we have

$$\omega_{\mathbf{Y}, E/F} = \eta_{E/F} \circ [\rho]_F \circ \delta_{p',F}.$$

Taking both F -points and E -points in Equation (12) and composing with $[\rho]_E$ and $[\rho]_F$ respectively we obtain the commutative diagram

$$\begin{array}{ccccccccc}
1 & \longrightarrow & \mathbf{Z}(E) & \longrightarrow & \mathbf{Y}^{\text{sc}}(E) & \longrightarrow & \mathbf{Y}^{\text{ad}}(E) & \xrightarrow{\delta_{p,E}} & H^1(E, \mathbf{Z}) & \xrightarrow{[\rho]_E} & H^1(E, \{\pm 1\}) & \xrightarrow{\sim} & E^*/E^{*2} \\
& & \uparrow i_{F,E} & & \uparrow i_{F,E} & & \uparrow i_{F,E} & & \uparrow [i_{F,E}] & & \uparrow [i_{F,E}] & & \uparrow i \\
1 & \longrightarrow & \mathbf{Z}(F) & \longrightarrow & \mathbf{Y}^{\text{sc}}(F) & \longrightarrow & \mathbf{Y}^{\text{ad}}(F) & \xrightarrow{\delta_{p,F}} & H^1(F, \mathbf{Z}) & \xrightarrow{[\rho]_F} & H^1(F, \{\pm 1\}) & \xrightarrow{\sim} & F^*/F^{*2}
\end{array}$$

where the vertical arrows $i_{F,E}$ are the natural embeddings of F -points into E -points of an algebraic group defined over F and $[i_{F,E}]$ are the maps induced by restriction of cocycles from the absolute Galois group of F to that of E . We arrive at the following statement.

Lemma 15. *We have the following equality of characters of $\mathbf{Y}^{\text{ad}}(F)$*

$$[\rho]_E \circ \delta_{p,E} \circ i_{F,E} = [i_{F,E}] \circ [\rho]_F \circ \delta_{p,F}.$$

In particular, if $\eta_{E/F}$ extends to a quadratic character η of E^ then*

$$\omega_{E,\eta}^{\text{ad}} := \eta \circ [\rho]_E \circ \delta_{p,E}$$

is a character of $\mathbf{Y}^{\text{ad}}(E)$ that extends $\omega_{\mathbf{Y}^{\text{ad}},E/F}$ and hence, composition with the natural projection $\mathbf{Y}(E) \rightarrow \mathbf{Y}^{\text{ad}}(E)$ gives rise to a character $\omega_{E,\eta}$ of $\mathbf{Y}(E)$ that extends $\omega_{\mathbf{Y},E/F}$.

Proof. The lemma is immediate from the observation that the map $[i_{F,E}] : H^1(F, \{\pm 1\}) \rightarrow H^1(E, \{\pm 1\})$ identifies with the natural map $i : F^*/F^{*2} \rightarrow E^*/E^{*2}$ induced by the inclusion $F^* \subseteq E^*$. \square

Remark 10. There exists a quadratic character η of E^* that extends $\eta_{E/F}$ if and only if $-1 \in N(E/F)$. Indeed, such η exists if and only if $\eta_{E/F}$ considered as a character of F^*/F^{*2} is trivial on the kernel of the natural map $aF^{*2} \mapsto aE^{*2} : F^*/F^{*2} \rightarrow E^*/E^{*2}$ and if $a \in E \setminus F$ is such that $a^2 \in F$ then this kernel is $F^{*2} \sqcup a^2F^{*2}$ while $-a^2 \in N(E/F)$. When this is the case, we can choose a quadratic extension L/E such that $\eta_{L/E}$ extends $\eta_{E/F}$. Note that in this case we have

$$\omega_{E,\eta_{L/E}} = \omega_{\mathbf{Y}_E,L/E}$$

where \mathbf{Y}_E is the base change of \mathbf{Y} to E and the left hand side is defined in the lemma.

We compute $\omega_{\mathbf{Y},E/F}$ when \mathbf{Y} is either GL_n or the quasi-split classical group \mathbb{H}° as in §8.2. The method that we use was explained to us by Beuzart-Plessis and we thank him for his help. We also mention that he obtained a different proof of Prasad's character computation in the case of unitary groups, similar to that of the general linear and odd orthogonal case.

- Assume that $\mathbf{Y} = \text{GL}_n$ so that $\mathbf{Y}^{\text{ad}} = \text{PGL}_n$ and $\mathbf{Y}^{\text{sc}} = \text{SL}_n$. Recall that $\text{End}(\mathbb{G}_m) = \mathbb{Z}$ where we identify $n \in \mathbb{Z}$ with the n th power endomorphism of the multiplicative group \mathbb{G}_m . Let μ_n be the algebraic subgroup of \mathbb{G}_m defined by the exact sequence

$$1 \longrightarrow \mu_n \longrightarrow \mathbb{G}_m \xrightarrow{n} \mathbb{G}_m \longrightarrow 1.$$

Consider the following two commutative diagrams

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mu_n & \longrightarrow & \mathbb{G}_m & \xrightarrow{n} & \mathbb{G}_m \longrightarrow 1 \\
 & & \downarrow \binom{n}{2} & & \downarrow \binom{n}{2} & & \downarrow n-1 \\
 1 & \longrightarrow & \mu_2 & \longrightarrow & \mathbb{G}_m & \xrightarrow{2} & \mathbb{G}_m \longrightarrow 1
 \end{array}$$

and

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mu_n & \longrightarrow & \mathrm{SL}_n & \longrightarrow & \mathrm{PGL}_n \longrightarrow 1 \\
 & & \parallel & & \uparrow p_2 & & \uparrow s \\
 1 & \longrightarrow & \mu_n & \longrightarrow & \mathbb{G}_m \times \mathrm{SL}_n & \longrightarrow & \mathrm{GL}_n \longrightarrow 1 \\
 & & \downarrow -1 & & \downarrow p_1 & & \downarrow \det \\
 1 & \longrightarrow & \mu_n & \longrightarrow & \mathbb{G}_m & \xrightarrow{n} & \mathbb{G}_m \longrightarrow 1
 \end{array}$$

where $s : \mathrm{GL}_n \rightarrow \mathrm{PGL}_n$ is the natural projection and p_i the projection to the i th coordinate, $i = 1, 2$. Note that $\mathbf{Z} = \mu_n$ is the center of SL_n and $\rho|_{\mathbf{Z}} = \binom{n}{2}$. By passing to F -points and applying the long exact sequence of [Ser94, Proposition 43] we, in particular, get from the first diagram the commutative diagram

$$\begin{array}{ccc}
 F^* & \xrightarrow{\delta_n} & H^1(F, \mu_n) \\
 \downarrow n-1 & & \downarrow [\binom{n}{2}] \\
 F^* & \xrightarrow{\delta_2} & H^1(F, \mu_2) \xrightarrow{\sim} F^*/(F^*)^2
 \end{array}$$

and from the second diagram the commutative diagram

$$\begin{array}{ccc}
 \mathrm{PGL}_n(F) & \xrightarrow{\delta} & H^1(F, \mu_n) \\
 \uparrow s & & \parallel \\
 \mathrm{GL}_n(F) & \xrightarrow{\delta'} & H^1(F, \mu_n) \\
 \downarrow \det & & \downarrow [-1] \\
 F^* & \xrightarrow{\delta_n} & H^1(F, \mu_n)
 \end{array}$$

where $\delta_n, \delta_2, \delta$ and δ' are the corresponding connecting homomorphisms. Note that

$$\omega_{\mathrm{GL}_n, E/F} = \eta_{E/F} \circ \left[\binom{n}{2} \right]_F \circ \delta \circ s.$$

Putting the two last diagrams together we obtain

$$\left[- \binom{n}{2} \right]_F \circ \delta \circ s = \delta_2 \circ (n-1) \circ \det.$$

Since δ_2 induces the natural projection $F^* \rightarrow F^*/F^{*2}$ and $[-1]_F$ the identity map on F^*/F^{*2} we deduce that

$$\omega_{\mathrm{GL}_n, E/F} = \eta_{E/F}^{n-1} \circ \det.$$

- In the symplectic case we have that $\omega_{\mathrm{Sp}_{2n}, E/F}$ is trivial. Indeed, if $\mathbf{Y} = \mathrm{Sp}_{2n}$ then $\mathbf{Y} = \mathbf{Y}^{\mathrm{ad}} = \mathbf{Y}^{\mathrm{sc}}$ and $\mathbf{Z} = \{1\}$. In fact, $\mathrm{Sp}_{2n}(F)$ has no non-trivial characters.
- Consider the quasi-split orthogonal case. Thus, $\mathbf{Y} = \mathrm{SO}(j[n])$ where in the odd case we may take $j[n] = w_{2n+1}$ and in the even case, either $j[n] = w_{2n}$ (the split case) or j is anisotropic of size two (the quasi-split non-split case). If $n = 0$ then \mathbf{Y} is anisotropic, hence ρ is trivial and therefore so is $\omega_{\mathbf{Y}, E/F}$. Assume now that $n \geq 1$. Then \mathbf{Y} is semi-simple and $\mathbf{Y}^{\mathrm{sc}} = \mathrm{Spin}(j[n])$ is the associated spin group. The short exact sequence

$$1 \rightarrow \mu_2 \rightarrow \mathrm{Spin}(j[n]) \rightarrow \mathrm{SO}(j[n]) \rightarrow 1$$

gives rise to the connecting homomorphism

$$\delta_{\mathrm{SO}} : \mathrm{SO}(j[n], F) \rightarrow H^1(F, \mu_2) \simeq F^*/(F^*)^2.$$

By Remark 9 we have $\omega_{\mathrm{SO}(j[n]), E/F} = \eta_{E/F} \circ [\rho]_F \circ \delta_{\mathrm{SO}}$. One verifies that in the even case ρ is a weight of the diagonal maximal split torus while in the odd case it is not. In particular, in the even case $[\rho]_F$ is the trivial map so that $\omega_{\mathrm{SO}(j[n]), E/F}$ is trivial and in the odd case $[\rho]_F$ is the identity map so that $\omega_{\mathrm{SO}(j[n]), E/F} = \eta_{E/F} \circ \delta_{\mathrm{SO}}$.

The connecting homomorphism δ_{SO} is known to be the spinor norm sn . For convenience we provide the argument that is analogous to the GL_n computation. Denote by $\tilde{\mathrm{sn}} : \mathrm{GSpin}(j[n]) \rightarrow \mathbb{G}_m$ the spinor norm on $\mathrm{GSpin}(j[n])$, so that if

$$p : \mathrm{GSpin}(j[n]) \rightarrow \mathrm{SO}(j[n])$$

is the natural projection with kernel \mathbb{G}_m , the composed maps

$$\mathrm{GSpin}(j[n], F) \xrightarrow{p} \mathrm{SO}(j[n]) \xrightarrow{\mathrm{sn}} F^*/F^{*2}$$

and

$$\mathrm{GSpin}(j[n], F) \xrightarrow{\tilde{\mathrm{sn}}} F^* \rightarrow F^*/F^{*2}$$

are equal. Consider the commutative diagram

$$\begin{array}{ccccccccc} 1 & \longrightarrow & \mu_2 & \longrightarrow & \mathrm{Spin}(j[n]) & \longrightarrow & \mathrm{SO}(j[n]) & \longrightarrow & 1 \\ & & \parallel & & \uparrow p_2 & & \uparrow p & & \\ 1 & \longrightarrow & \mu_2 & \longrightarrow & \mathbb{G}_m \times \mathrm{Spin}(j[n]) & \longrightarrow & \mathrm{GSpin}(j[n]) & \longrightarrow & 1 \\ & & \parallel & & \downarrow p_1 & & \downarrow \tilde{\mathrm{sn}} & & \\ 1 & \longrightarrow & \mu_2 & \longrightarrow & \mathbb{G}_m & \xrightarrow{2} & \mathbb{G}_m & \longrightarrow & 1. \end{array}$$

Taking F -points we get the following commutative diagram

$$\begin{array}{ccc}
 \mathrm{SO}(j[n], F) & \xrightarrow{\delta_{\mathrm{SO}}} & H^1(F, \mu_2) \simeq F^*/(F^*)^2 \\
 \uparrow p & & \parallel \\
 \mathrm{GSpin}(j[n], F) & \xrightarrow{\delta_{\mathrm{GSpin}}} & H^1(F, \mu_2) \simeq F^*/(F^*)^2 \\
 \downarrow \tilde{\mathrm{sn}} & & \parallel \\
 F^* & \xrightarrow{\delta_2} & H^1(F, \mu_2) \simeq F^*/F^{*2}.
 \end{array}$$

Since p is surjective on F -points (see e.g. [Sch85, Chapter 9, Theorem 3.3]) and δ_2 induces the natural projection from F^* to F^*/F^{*2} we deduce that $\delta_{\mathrm{SO}} = \mathrm{sn}$. All together we conclude that for all $n \geq 0$ we have

$$\omega_{\mathrm{SO}(j[n]), E/F} = \eta_{E/F}^{n_0} \circ \mathrm{sn}$$

where $n_0 \in \{0, 1, 2\}$ is the size of j .

- Let $\mathbf{Y} = \mathrm{U}_{n, K/F} = \mathrm{U}_{K/F}(w_n)$ be the F -quasi-split unitary group with respect to a quadratic extension K/F and τ the Galois involution associated to K/F . Thus, $\mathbf{Y}^{\mathrm{sc}} = \mathrm{SU}_{n, K/F}$ and $\mathbf{Y}^{\mathrm{ad}} = \mathrm{PU}_{n, K/F}$. Let $m : \mathrm{U}_{n, K/F} \rightarrow \mathrm{PU}_{n, K/F}$ be the natural projection. Its restriction to $\mathrm{SU}_{n, K/F}$ gives rise to the short exact sequence

$$1 \rightarrow \mu'_n \rightarrow \mathrm{SU}_{n, K/F} \rightarrow \mathrm{PU}_{n, K/F} \rightarrow 1$$

where the center μ'_n of $\mathrm{SU}_{n, K/F}$ is its intersection with the center $\mathrm{Res}_{K/F}(\mu_n)$ of $\mathrm{Res}_{K/F}(\mathrm{SL}_n)$. By taking F -points we get the connecting homomorphism $\delta : \mathrm{PU}_{n, K/F}(F) \rightarrow H^1(F, \mu'_n)$ and

$$\omega_{\mathrm{U}_{n, K/F}, E/F} = \eta_{E/F} \circ [\rho]_F \circ \delta \circ m.$$

Note that if n is odd then the only homomorphism $\mu'_n \rightarrow \mu_2$ is the trivial one and it follows that $[\rho]_F$ and therefore also $\omega_{\mathrm{U}_{n, K/F}, E/F}$ is trivial.

We assume from now on that n is even. Then one verifies that

$$\rho(z) = z^{-\frac{n}{2}}, \quad z \in \mu'_n(F).$$

We now interpret the connecting homomorphism δ . Denote by

$$\mathrm{wsn} : \mathrm{U}_{n, K/F}(F) = \mathrm{U}_n(K/F) \rightarrow K^*/F^*$$

the composition of the determinant

$$\det : \mathrm{U}_{n, K/F}(F) \rightarrow \mathrm{U}_{1, K/F}(F)$$

with the inverse of the isomorphism $zF^* \mapsto zz^{-\tau}$ from K^*/F^* to $\mathrm{U}_{1, K/F}(F)$. (In [Wal59], Wall interpreted this map as a spinor norm analogue.) We show that

$$(13) \quad [\rho]_F \circ \delta \circ m = N_{K/F} \circ \mathrm{wsn}.$$

The character $N_{K/F} \circ \mathrm{wsn}$ naturally appears in another manner, which we now describe. Let $\delta_K \in K \setminus F$ be such that $\delta_K^2 \in F$. We denote by O_{2n}^q the orthogonal

group defined over F with respect to the quadratic form q on F^{2n} associated to the bilinear form

$$b : \left(\begin{pmatrix} X_1 \\ X_2 \end{pmatrix}, \begin{pmatrix} Y_1 \\ Y_2 \end{pmatrix} \right) \mapsto \text{Tr}_{K/F}(({}^t X_1 - \delta_K {}^t X_2)w_n(Y_1 + \delta_K Y_2)), \quad X_i, Y_i \in F^n, \quad i = 1, 2.$$

Since we assume that n is even, one checks that the quadratic space (F^{2n}, q) is a sum of hyperbolic planes hence O_{2n}^q is F -split (for n odd it is quasi-split and splits over K). There is now an obvious natural imbedding

$$i : \text{U}_{n,K/F} \rightarrow \text{SO}_{2n}^q,$$

defined over F , and by [Sch85, Chapter 10, Theorem 1.5] we have

$$N_{K/F} \circ \text{wsn} = \text{sn} \circ i.$$

We will make use of this equality to prove equation (13). Consider the commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mu'_n & \xrightarrow{i_1} & \text{U}_{1,K/F} \times \text{SU}_{n,K/F} & \xrightarrow{s_1} & \text{U}_{n,K/F} \longrightarrow 1 \\ & & \parallel & & \downarrow p_2 & & \downarrow m \\ 1 & \longrightarrow & \mu'_n & \longrightarrow & \text{SU}_{n,K/F} & \longrightarrow & \text{PU}_{n,K/F} \longrightarrow 1 \end{array}$$

where $i_1(\mu) = (\mu^{-1}, \mu I_n)$ and $s_1(z, u) = zu$, that implies the following one

$$\begin{array}{ccc} \text{U}_{n,K/F}(F) & \xrightarrow{\delta_{s_1,F}} & H^1(F, \mu'_n) \\ \downarrow m & & \parallel \\ \text{PU}_{n,K/F}(F) & \xrightarrow{\delta} & H^1(F, \mu'_n) \end{array}$$

so that

$$\delta \circ m = \delta_{s_1,F}.$$

Let

$$\overline{\text{U}_{n,K/F}}^2 = \{(z, u) \in \text{U}_{1,K/F} \times \text{U}_{n,K/F}, \det(u) = z^2\}$$

and $s_2 : \overline{\text{U}_{n,K/F}}^2 \rightarrow \text{U}_{n,K/F}$ the projection to the second coordinate

$$s_2(z, u) = u.$$

The map $i_2 : \mu \mapsto (\mu, I_n)$ identifies $\mu'_2 = \mu_2$ with the kernel of s_2 . Let $p_{n,2} : \text{U}_{1,K/F} \times \text{SU}_{n,K/F} \rightarrow \overline{\text{U}_{n,K/F}}^2$ be given by

$$p_{n,2}(z, u) = (z^{n/2}, zu) = (\rho(z)^{-1}, zu).$$

This gives the commutative diagram

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mu'_n & \xrightarrow{i_1} & \mathrm{U}_{1,K/F} \times \mathrm{SU}_{n,K/F} & \xrightarrow{s_1} & \mathrm{U}_{n,K/F} \longrightarrow 1 \\
 & & \rho \downarrow & & \downarrow p_{n,2} & & \parallel \\
 1 & \longrightarrow & \mu'_2 & \xrightarrow{i_2} & \overline{\mathrm{U}}_{n,K/F}^2 & \xrightarrow{s_2} & \mathrm{U}_{n,K/F} \longrightarrow 1.
 \end{array}$$

By taking F -points we obtain the commutative diagram

$$\begin{array}{ccc}
 \mathrm{U}_{n,K/F}(F) & \xrightarrow{\delta_{s_1,F}} & H^1(F, \mu'_n) \\
 \parallel & & \downarrow [\rho]_F \\
 \mathrm{U}_{n,K/F}(F) & \xrightarrow{\delta_{s_2,F}} & H^1(F, \mu'_2)
 \end{array}$$

so that

$$\delta_{s_2,F} = [\rho]_F \circ \delta_{s_1,F}.$$

To prove (13) it remains to show that

$$(14) \quad \delta_{s_2,F} = \mathrm{sn} \circ i.$$

To this end we recall the exact sequence

$$1 \rightarrow \mu_2 \xrightarrow{u} \mathrm{Spin}_{2n}^q \xrightarrow{v} \mathrm{SO}_{2n}^q \rightarrow 1$$

and show that there exists a map

$$j : \overline{\mathrm{U}}_{n,K/F}^2 \rightarrow \mathrm{Spin}_{2n}^q$$

defined over F such that the following diagram is commutative

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mu'_2 & \xrightarrow{i_2} & \overline{\mathrm{U}}_{n,F'/F}^2 & \xrightarrow{s_2} & \mathrm{U}_{n,F'/F} \longrightarrow 1 \\
 & & \parallel & & \downarrow j & & \downarrow i \\
 1 & \longrightarrow & \mu_2 & \xrightarrow{u} & \mathrm{Spin}_{2n} & \xrightarrow{v} & \mathrm{SO}_{2n} \longrightarrow 1.
 \end{array}$$

Taking F -points and recalling from the orthogonal case that the spinor norm sn is the connecting homomorphism associated with the bottom row we arrive at (14) and complete our proof.

To show that such j exists we first make the following simple observation. There exists at most one map $j : \overline{\mathrm{U}}_{n,K/F}^2 \rightarrow \mathrm{Spin}_{2n}^q$ such that $v \circ j = i \circ s_2$. Indeed, if j' is another such map then $u \mapsto j'(u)j(u)^{-1} : \overline{\mathrm{U}}_{n,K/F}^2 \rightarrow \mu_2$ must be trivial since $\overline{\mathrm{U}}_{n,K/F}^2$ is connected. It is therefore enough to find such a j defined over K . Indeed, if the above diagram commutes for j it also does for j^τ . Over K we may identify

$$\mathrm{U}_{n,K/F} \simeq \mathrm{GL}_n,$$

and

$$\overline{\mathrm{U}}_{n,K/F}^2 \simeq \{(z, g) \in \mathbb{G}_m \times \mathrm{GL}_n, \det(g) = z^2\} =: \overline{\mathrm{GL}}_n^2$$

and with these identifications $s_2(z, g) = g$ and $i : \mathrm{GL}_n \rightarrow \mathrm{SO}_{2n}^q$ is an imbedding defined over K . It now suffices to show that there exists $j : \overline{\mathrm{GL}_n}^2 \rightarrow \mathrm{Spin}_{2n}^q$ defined over K such that $v \circ j = s_2 \circ i$ and $j(-1, I_n) = u(-1)$. In fact, we claim that if there exists j such that $v \circ j = s_2 \circ i$ then $j(-1, I_n) = u(-1)$ is automatic. Indeed, otherwise j defines a map $\bar{j} : \mathrm{GL}_n \rightarrow \mathrm{Spin}_{2n}^q$ such that $v \circ \bar{j} = i$. Note that GL_n , Spin_{2n}^q and SO_{2n}^q are all of rank n . Since i is injective it maps any maximal torus of GL_n to a maximal torus \mathbf{T} of SO_{2n}^q . Furthermore, \bar{j} must also be injective and therefore $\bar{j}(\mathbf{T})$ is a maximal torus of Spin_{2n}^q . Clearly $\bar{j}(\mathbf{T})$ is a subgroup of $v^{-1}(\mathbf{T})$. It is a well known fact on simply connected covers that $v^{-1}(\mathbf{T})$ is a maximal torus of Spin_{2n}^q and therefore $v^{-1}(\mathbf{T}) = \bar{j}(\mathbf{T})$. Since $u(\mu_2) \subseteq v^{-1}(\mathbf{T})$ is in the kernel of v this contradicts the injectivity of i .

Note that we may further identify $\overline{\mathrm{GL}_n}^2$ as the semidirect product $\mathbb{G}_m \ltimes \mathrm{SL}_n$ where the action of \mathbb{G}_m on SL_n is given by

$$zg z^{-1} = \begin{pmatrix} z^2 & \\ & I_{n-1} \end{pmatrix} g \begin{pmatrix} z^{-2} & \\ & I_{n-1} \end{pmatrix}, \quad z \in \mathbb{G}_m, g \in \mathrm{SL}_n.$$

Indeed $(z, g) \mapsto (z, \mathrm{diag}(z^2, I_{n-1})g) : \mathbb{G}_m \ltimes \mathrm{SL}_n \rightarrow \overline{\mathrm{GL}_n}^2$ is an isomorphism. It now suffices to show that there is a map $j : \mathbb{G}_m \ltimes \mathrm{SL}_n \rightarrow \mathrm{Spin}_{2n}^q$ defined over K such that

$$j(z, g) = i(\mathrm{diag}(z^2, I_{n-1})g).$$

Let $j_1 = i|_{\mathrm{SL}_n} : \mathrm{SL}_n \rightarrow \mathrm{SO}_{2n}^q$ and $j_2 : \mathbb{G}_m \rightarrow \mathrm{SO}_{2n}^q$ be given by $j_2(z) = i(\mathrm{diag}(z^2, I_{n-1}))$. We claim that it suffices to lift j_1 and j_2 to Spin_{2n}^q . Indeed, if $\bar{j}_1 : \mathrm{SL}_n \rightarrow \mathrm{Spin}_{2n}^q$ and $\bar{j}_2 : \mathbb{G}_m \rightarrow \mathrm{Spin}_{2n}^q$ are K -morphisms such that $v \circ \bar{j}_t = j_t$, $t = 1, 2$ then we claim that

$$j(z, g) = \bar{j}_2(z) \bar{j}_1(g)$$

is automatically a K -morphism. Indeed, we need to verify that

$$\bar{j}_2(z) \bar{j}_1(g) \bar{j}_2(z)^{-1} = \bar{j}_2 \begin{pmatrix} z^2 & \\ & I_{n-1} \end{pmatrix} g \begin{pmatrix} z^{-2} & \\ & I_{n-1} \end{pmatrix}.$$

Fix z and let $f, f' : \mathrm{SL}_n \rightarrow \mathrm{Spin}_{2n}^q$ be defined by the left and right hand sides of the above equation respectively. Then $v \circ f = v \circ f'$ and therefore $g \mapsto f'(g) f(g)^{-1} : \mathrm{SL}_n \rightarrow \mu_2$ must be trivial as SL_n is connected. Thus, $f' = f$ as required.

To complete the proof we observe that j_1 and j_2 both lift to Spin_{2n}^q . The map j_1 has a lifting since SL_n is simply connected. Indeed, let \mathbf{Y} be the image of j_1 in SO_{2n}^q and $\mathbf{Y}' = v^{-1}(\mathbf{Y})^\circ$ (the connected component of the pre-image of \mathbf{Y} in Spin_{2n}^q). Then j_1 is an isomorphism of SL_n with \mathbf{Y} and $v : \mathbf{Y}' \rightarrow \mathbf{Y}$ is an isogeny (with kernel contained in μ_2). Since SL_n and therefore also \mathbf{Y} is simply connected, $v|_{\mathbf{Y}'}$ is in fact an isomorphism and therefore $\bar{j}_1 = (v|_{\mathbf{Y}'})^{-1} \circ j_1 : \mathrm{SL}_n \rightarrow \mathrm{Spin}_{2n}^q$ is such that $j_1 = v \circ \bar{j}_1$. Let $\mathbf{Z} \simeq \mathbb{G}_m$ be the image of j_2 in SO_{2n}^q and $\mathbf{Z}' = v^{-1}(\mathbf{Z})^\circ \simeq \mathbb{G}_m$ be the connected component of its pre-image in Spin_{2n}^q . If \mathbf{Z}' contains the kernel $u(\mu_2)$ of v then $v|_{\mathbf{Z}'} = w^2$ is the square of an isomorphism $w : \mathbf{Z}' \rightarrow \mathbf{Z}$ (any endomorphism of \mathbb{G}_m with kernel μ_2 is either the square map 2 or its composition with the inverse map). We can then define the desired lifting $\bar{j}_2(z) = w^{-1}(i(\mathrm{diag}(z, I_{n-1})))$ such that $v \circ \bar{j}_2 = j_2$. Otherwise, $v|_{\mathbf{Z}'} : \mathbf{Z}' \rightarrow \mathbf{Z}$ is an isomorphism and we can define $\bar{j}_2 = (v|_{\mathbf{Z}'})^{-1} \circ j_2$. This completes our computation. (In

fact, the second case cannot occur, it would result in a map $\bar{j} : \mathrm{GL}_n \rightarrow \mathrm{Spin}_{2n}^q$ such that $v \circ \bar{j} = i$ and as observed before this contradicts the fact that i is injective.)

All together we proved the formula

$$\omega_{\mathrm{U}_{n,K/F},E/F} = \eta_{E/F}^{n-1} \circ N_{K/F} \circ \mathrm{wsn}.$$

Remark 11. In particular, $\omega_{\mathrm{U}_{n,E/F},E/F}$ is always the trivial character since $\eta_{E/F} \circ N_{E/F}$ is the trivial character of E^* . For $K \neq E$ we have

$$\omega_{\mathrm{U}_{n,K/F},E/F} = \eta_{EK/K}^{n-1} \circ \mathrm{wsn}.$$

Indeed, in order to see that $\eta_{EK/K} = \eta_{E/F} \circ N_{K/F}$ we observe that on the one hand the right hand side is clearly trivial on $N(EK/K)$ and on the other hand it is not the trivial character since by local class field theory $N(K/F)$ and $N(E/F)$ are two different subgroups of F^* of index two. Note further that $\eta_{EK/K} \circ \mathrm{wsn}$ is well defined since $F^* \subseteq N(EK/K)$ by Lemma 5.

We describe another interpretation of $\omega_{\mathbf{Y},E/F}$ provided in [Pra] (see also [BP18, §5.2]). If \mathbf{Y} is quasi-split then according to [LL15] the Langlands morphism

$$\alpha_{\mathbf{Y}} : H^1(W_F, Z(\widehat{\mathbf{Y}}(\mathbb{C}))) \rightarrow \mathrm{Hom}(Y, \mathbb{C}^\times)$$

is bijective. The class

$$\chi_{\mathbf{Y},E/F} := \alpha_{\mathbf{Y}}^{-1}(\omega_{\mathbf{Y},E/F}) : W_F \rightarrow Z(\widehat{\mathbf{Y}}(\mathbb{C}))$$

is represented by the unique cocycle $\chi_{\mathbf{Y},E/F}$ that maps W_E to the identity and the non-trivial class in W_F/W_E to the image of $-I_2$ by any principal SL_2 -morphism from $\mathrm{SL}_2(\mathbb{C})$ to $\widehat{\mathbf{Y}}(\mathbb{C})$. In fact, $\chi_{\mathbf{Y},E/F}$ is a quadratic character of W_F and in the sequel we view $\chi_{\mathbf{Y},E/F}$ as a character of W'_F trivial on $\mathrm{SL}_2(\mathbb{C})$ so that it makes sense to twist an L -homomorphism of \mathbf{Y} by $\chi_{\mathbf{Y},E/F}$. The following lemma follows easily.

Lemma 16. *Suppose that $-1 \in N(E/F)$ and let L/E be as in Remark 10, so that $L' = LE'$ is a quadratic extension of E' . Let \mathbf{Y} be either SO_{2n+1} or $\mathrm{U}_{2n,E/F}$ and let N be either $2n+1$ or $2n$ respectively. For $[\phi] \in \Phi(\mathbf{Y}(E))$ we have*

$$I([\chi_{\mathbf{Y}_{E,L/E}} \phi]) = [\chi_{\mathrm{GL}_{N,L'/E'}} \phi'] \quad \text{where} \quad [\phi'] = I([\phi]).$$

Proof. In both cases, the principal SL_2 -morphism from $\mathrm{SL}_2(\mathbb{C})$ to $\widehat{\mathbf{Y}}(\mathbb{C})$ is the irreducible $2n$ -dimensional representation (it has image in $\mathrm{Sp}_{2n}(\mathbb{C})$ according to [GW09, Lemma 3.2.15]).

If $\mathbf{Y} = \mathrm{SO}_{2n+1}$ then $E' = E$ and $L' = L$, and $\chi_{\mathbf{Y}_{E,L/E}}$ identifies with the character of $\mathrm{Gal}(L/E)$ sending the non-trivial Galois element $\sigma_{L/E}$ to $-I_{2n}$ in the center of $\mathrm{Sp}_{2n}(\mathbb{C})$, whereas $\chi_{\mathrm{GL}_{N,L/E}}$ identifies with the character of $\mathrm{Gal}(L/E)$ sending $\sigma_{L/E}$ to $-I_{2n}$ in the center of $\mathrm{GL}_{2n}(\mathbb{C})$. As I is induced by composing L -homomorphisms with the natural inclusion of dual groups the result follows.

If $\mathbf{Y} = \mathrm{U}_{2n,E/F}$ then $I[\phi] = [\phi|_{W_{E'}}]$. Now $\chi_{\mathbf{Y}_{E,L/E}}$ identifies with the character of $\mathrm{Gal}(L/E)$ sending $\sigma_{L/E}$ to $-I_{2n}$ in the center of $\mathrm{GL}_{2n}(\mathbb{C})$, whereas $\chi_{\mathrm{GL}_{2n,L'/E'}}$ identifies with the character of $\mathrm{Gal}(L'/E')$ sending $\sigma_{L'/E'}$ to $-I_{2n}$ in the center of $\mathrm{GL}_{2n}(\mathbb{C})$, and the result follows. \square

A.2. Prasad's global character. Let K/k be a quadratic extension of number fields, and let $\eta_{K/k}$ be the quadratic automorphic character of \mathbb{A}_k^\times attached to K/k by global class field theory. It decomposes as $\eta_{K/k} = \prod_v \eta_{K_v/k_v}$ where the product is over all places v of k and η_{K_v/k_v} is the trivial character if v splits in K .

Denote by \mathbf{Y} a k -quasi-split group. Thanks to Section A.1 for each place v of k the natural map $\mathbf{Y}(k_v) \xrightarrow{p_v} \mathbf{Y}^{\text{ad}}(k_v)$ gives rise to the local homomorphism

$$N_v := [\rho]_{k_v} \circ \delta_{p,k_v} \circ p_v : \mathbf{Y}(k_v) \rightarrow k_v^*/k_v^{*2},$$

and with the obvious notation the exact sequence (12) also gives rise to the homomorphism

$$N_k := [\rho]_k \circ \delta_{p,k} \circ p_k : \mathbf{Y}(k) \rightarrow k^*/k^{*2}$$

that satisfies $j_{k_v} \circ N_k = N_v \circ i_{k_v}$ where $i_{k_v} : \mathbf{Y}(k) \rightarrow \mathbf{Y}(k_v)$ is the natural inclusion and $j_{k_v} : k^*/k^{*2} \rightarrow k_v^*/k_v^{*2}$ the natural homomorphism. The local homomorphisms N_v allows one to define the global homomorphism

$$N = \prod_v N_v : \mathbf{Y}(\mathbb{A}_k) \rightarrow \mathbb{A}_k^\times / \mathbb{A}_k^{\times 2}.$$

Note that $N|_{\mathbf{Y}(k)} = N_k$. We set

$$\omega_{\mathbf{Y},K/k} := \eta_{K/k} \circ N.$$

It is a smooth character of $\mathbf{Y}(\mathbb{A}_k)$, which is in fact automorphic since $\eta_{K/k}$ is and $N|_{\mathbf{Y}(k)}$ equals N_k composed with the natural diagonal map from k^*/k^{*2} to $\mathbb{A}_k^\times / \mathbb{A}_k^{\times 2}$. Prasad's global character is related to Prasad's local characters by the following formula:

$$\omega_{\mathbf{Y},K/k} = \prod_v \omega_{\mathbf{Y},K_v/k_v}$$

where $\omega_{\mathbf{Y},K_v/k_v}$ is the trivial character of $\mathbf{Y}(k_v)$ if v splits in K .

A.3. The opposition group. Following [Pra, §5] we define the opposition group \mathbf{Y}^{op} for any F -quasi-split reductive group \mathbf{Y} . We recall that \mathbf{Y} is a form of a unique F -split group \mathbf{Y}_s , and that any quasi-split form of \mathbf{Y}_s corresponds to an element in $\text{Hom}(\text{Gal}(\bar{F}/F), \text{Out}(\mathbf{Y}_s))$. Denote by $\mu_{\mathbf{Y}} : \text{Gal}(\bar{F}/F) \rightarrow \text{Out}(\mathbf{Y}_s)$ the homomorphism corresponding to \mathbf{Y} .

The group $\text{Out}(\mathbf{Y}_s)$ of outer automorphisms of \mathbf{Y}_s identifies with the group of pinned automorphisms of \mathbf{Y}_s associated with a quadruple $(\mathbf{Y}_s, \mathbf{B}, \mathbf{T}, (\mathbf{X}_\alpha)_{\alpha \in \Delta})$ where \mathbf{T} is a maximal torus inside a Borel subgroup \mathbf{B} of \mathbf{Y}_s , Δ is a basis of simple roots of \mathbf{Y}_s with respect to (\mathbf{B}, \mathbf{T}) and $(\mathbf{X}_\alpha)_{\alpha \in \Delta}$ is a pinning, all defined over F (see for example [Bor79]). Denote by w_0 the longest element in the Weyl group of \mathbf{Y}_s associated to \mathbf{T} . Then $\text{Aut}(\mathbf{H}_s, \mathbf{B}, \mathbf{T}, (\mathbf{X}_\alpha)_{\alpha \in \Delta})$ contains a unique element sending \mathbf{X}_α to $\mathbf{X}_{-w_0(\alpha)}$, $\alpha \in \Delta$, known as the *Chevalley involution* of \mathbf{Y}_s , which corresponds to an involution i_c in the center $Z(\text{Out}(\mathbf{Y}_s))$ of $\text{Out}(\mathbf{Y}_s)$ independent of the choice of data $(\mathbf{B}, \mathbf{T}, (\mathbf{X}_\alpha)_{\alpha \in \Delta})$. There is a unique element

$$\mu_{E/F} \in \text{Hom}(\text{Gal}(E/F), Z(\text{Out}(\mathbf{Y}_s))) \hookrightarrow \text{Hom}(\text{Gal}(\bar{F}/F), \text{Out}(\mathbf{Y}_s))$$

such that $\mu_{E/F}(\sigma) = i_c$. Then

$$\mu_{\mathbf{Y}^{\text{op}}} := \mu_{E/F} \mu_{\mathbf{Y}} \in \text{Hom}(\text{Gal}(\bar{F}/F), \text{Out}(\mathbf{Y}_s))$$

corresponds to a unique F -quasi-split form of \mathbf{Y}_s that we denote by \mathbf{Y}^{op} . Taking F -points we call $Y^{\text{op}} = \mathbf{Y}^{\text{op}}(F)$ the opposition group of Y . Note that \mathbf{Y} and \mathbf{Y}^{op} are isomorphic over E . It further follows from the definition that $(\mathbf{Y}^{\text{op}})^{\text{op}} = \mathbf{Y}$.

We now compute the opposition groups in the cases of interest to us.

- If $\mathbf{Y} = \text{GL}_n$ then $\mathbf{Y} = \mathbf{Y}_s$ and $\mu_{\mathbf{Y}}$ is the trivial character. Furthermore, the Chevalley involution of GL_n corresponds to $i_c \in Z(\text{Out}(\text{GL}_n))$ the class of the automorphism $g \mapsto w_n {}^t g^{-1} w_n^{-1}$. It is now easily verified that $\mathbf{Y}^{\text{op}} = \text{U}_{n,E/F}$.
- If $\mathbf{Y} = \text{Sp}_{2n}$ or $\mathbf{Y} = \text{SO}(j[n])$ (either the quasi-split orthogonal or symplectic case) then the Chevalley involution of \mathbf{Y}_s is trivial and therefore $\mathbf{Y} = \mathbf{Y}^{\text{op}}$.
- If $\mathbf{Y} = \text{U}_{n,F'/F}$ then $\mathbf{Y}_s = \text{GL}_n$ and $\mu_{\mathbf{Y}}$ factors through $\text{Gal}(F'/F)$ and sends τ to i_c . Hence $\mu_{\mathbf{Y}^{\text{op}}} = \mu_{E/F} \mu_{\mathbf{Y}}$ factors through $\text{Gal}(E'/F)$ and furthermore maps $\sigma\tau$ to i_c^2 hence lies in the kernel of $\mu_{\mathbf{Y}^{\text{op}}}$. Thus, $\mu_{\mathbf{Y}^{\text{op}}}$ factors through $\text{Gal}(F''/F)$ where $F'' = (E')^{\sigma\tau}$, and maps the non-trivial involution of F''/F (the restriction of either σ or τ) to i_c . That is,

$$\mathbf{Y}^{\text{op}} = \text{U}_{n,F''/F}.$$

REFERENCES

- [ABPS14] Anne-Marie Aubert, Paul Baum, Roger Plymen, and Maarten Solleveld, *On the local Langlands correspondence for non-tempered representations*, Münster J. Math. **7** (2014), no. 1, 27–50. MR 3271238 [43](#), [44](#)
- [AGR93] Avner Ash, David Ginzburg, and Steven Rallis, *Vanishing periods of cusp forms over modular symbols*, Math. Ann. **296** (1993), no. 4, 709–723. MR 1233493 (94f:11044) [38](#)
- [AKT04] U. K. Anandavardhanan, Anthony C. Kable, and R. Tandon, *Distinguished representations and poles of twisted tensor L -functions*, Proc. Amer. Math. Soc. **132** (2004), no. 10, 2875–2883. MR 2063106 [46](#), [47](#)
- [AR05] U. K. Anandavardhanan and C. S. Rajan, *Distinguished representations, base change, and reducibility for unitary groups*, Int. Math. Res. Not. (2005), no. 14, 841–854. MR 2146859 [46](#), [47](#)
- [Art13] James Arthur, *The endoscopic classification of representations*, American Mathematical Society Colloquium Publications, vol. 61, American Mathematical Society, Providence, RI, 2013, Orthogonal and symplectic groups. MR 3135650 [43](#)
- [Ato17] Hiraku Atobe, *On the uniqueness of generic representations in an L -packet*, Int. Math. Res. Not. IMRN (2017), no. 23, 7051–7068. MR 3801418 [43](#)
- [BD08] Philippe Blanc and Patrick Delorme, *Vecteurs distributions H -invariants de représentations induites, pour un espace symétrique réductif p -adique G/H* , Ann. Inst. Fourier (Grenoble) **58** (2008), no. 1, 213–261. MR 2401221 (2009e:22015) [10](#), [11](#)
- [Ber84] Joseph N. Bernstein, *P -invariant distributions on $\text{GL}(N)$ and the classification of unitary representations of $\text{GL}(N)$ (non-Archimedean case)*, Lie group representations, II (College Park, Md., 1982/1983), Lecture Notes in Math., vol. 1041, Springer, Berlin, 1984, pp. 50–102. MR 748505 (86b:22028) [45](#)
- [BM19] Paul Broussous and Nadir Matringe, *Multiplicity One for Pairs of Prasad–Takloo-Bighash Type*, International Mathematics Research Notices (2019), rnz254. [18](#)
- [Bor79] A. Borel, *Automorphic L -functions*, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 27–61. MR MR546608 (81m:10056) [44](#), [64](#)

- [BP] Raphaël Beuzart-Plessis, *On distinguished generic representations*, Preprint. [4](#), [38](#), [39](#)
- [BP18] ———, *On distinguished square-integrable representations for Galois pairs and a conjecture of Prasad*, *Invent. Math.* **214** (2018), no. 1, 437–521. MR 3858402 [63](#)
- [CPSS11] J. W. Cogdell, I. I. Piatetski-Shapiro, and F. Shahidi, *Functoriality for the quasisplit classical groups*, *On certain L-functions*, *Clay Math. Proc.*, vol. 13, Amer. Math. Soc., Providence, RI, 2011, pp. 117–140. MR 2767514 (2012f:22036) [54](#)
- [Die48] Jean Dieudonné, *Sur les groupes classiques*, *Actualités Sci. Ind.*, no. 1040 = *Publ. Inst. Math. Univ. Strasbourg (N.S.)* no. 1 (1945), Hermann et Cie., Paris, 1948. MR 0024439 [53](#)
- [Fli91] Yuval Z. Flicker, *On distinguished representations*, *J. Reine Angew. Math.* **418** (1991), 139–172. MR 1111204 [29](#)
- [Fli92] ———, *Distinguished representations and a Fourier summation formula*, *Bull. Soc. Math. France* **120** (1992), no. 4, 413–465. MR 1194271 [41](#)
- [FLO12] Brooke Feigon, Erez Lapid, and Omer Offen, *On representations distinguished by unitary groups*, *Publ. Math. Inst. Hautes Études Sci.* (2012), 185–323. MR 2930996 [18](#), [29](#), [42](#)
- [GGP12] Wee Teck Gan, Benedict H. Gross, and Dipendra Prasad, *Symplectic local root numbers, central critical L values, and restriction problems in the representation theory of classical groups*, *Astérisque* (2012), no. 346, 1–109, *Sur les conjectures de Gross et Prasad. I.* MR 3202556 [45](#), [47](#)
- [GK75] I. M. Gel'fand and D. A. Kajdan, *Representations of the group $GL(n, K)$ where K is a local field*, *Lie groups and their representations (Proc. Summer School, Bolyai János Math. Soc., Budapest, 1971)*, Halsted, New York, 1975, pp. 95–118. MR 0404534 (53 #8334) [29](#), [37](#), [54](#)
- [GO16] Maxim Gurevich and Omer Offen, *A criterion for integrability of matrix coefficients with respect to a symmetric space*, *J. Funct. Anal.* **270** (2016), no. 12, 4478–4512. MR 3490774 [13](#)
- [Gur15] Maxim Gurevich, *On a local conjecture of Jacquet, ladder representations and standard modules*, *Math. Z.* **281** (2015), no. 3-4, 1111–1127. MR 3421655 [18](#), [41](#), [46](#)
- [GW09] Roe Goodman and Nolan R. Wallach, *Symmetry, representations, and invariants*, *Graduate Texts in Mathematics*, vol. 255, Springer, Dordrecht, 2009. MR 2522486 (2011a:20119) [63](#)
- [Han20] Marcela Hanzer, *Generalized injectivity conjecture for classical p -adic groups II*, *J. Pure Appl. Algebra* **224** (2020), no. 1, 149–168. MR 3986415 [40](#), [41](#)
- [Hen00] Guy Henniart, *Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique*, *Invent. Math.* **139** (2000), no. 2, 439–455. MR 1738446 [43](#)
- [Hen10] ———, *Correspondance de Langlands et fonctions L des carrés extérieur et symétrique*, *Int. Math. Res. Not. IMRN* (2010), no. 4, 633–673. MR 2595008 [46](#), [47](#)
- [HT01] Michael Harris and Richard Taylor, *The geometry and cohomology of some simple Shimura varieties*, *Annals of Mathematics Studies*, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich. MR MR1876802 (2002m:11050) [43](#)
- [HW93] A. G. Helminck and S. P. Wang, *On rationality properties of involutions of reductive groups*, *Adv. Math.* **99** (1993), no. 1, 26–96. MR 1215304 (94d:20051) [7](#), [11](#), [14](#)
- [Jac95] Hervé Jacquet, *The continuous spectrum of the relative trace formula for $GL(3)$ over a quadratic extension*, *Israel J. Math.* **89** (1995), no. 1-3, 1–59. MR 1324453 (96a:22029) [3](#), [10](#)
- [JLR99] Hervé Jacquet, Erez Lapid, and Jonathan Rogawski, *Periods of automorphic forms*, *J. Amer. Math. Soc.* **12** (1999), no. 1, 173–240. MR 1625060 (99c:11056) [3](#), [10](#)
- [JPSS83] H. Jacquet, I. I. Piatetskii-Shapiro, and J. A. Shalika, *Rankin-Selberg convolutions*, *Amer. J. Math.* **105** (1983), no. 2, 367–464. MR 701565 (85g:11044) [45](#)
- [JS03] Dihua Jiang and David Soudry, *The local converse theorem for $SO(2n + 1)$ and applications*, *Ann. of Math. (2)* **157** (2003), no. 3, 743–806. MR 1983781 (2005b:11193) [4](#), [38](#), [39](#)
- [Kab04] Anthony C. Kable, *Asai L-functions and Jacquet's conjecture*, *Amer. J. Math.* **126** (2004), no. 4, 789–820. MR 2075482 (2005g:11083) [46](#), [47](#)

- [Kal16] Tasho Kaletha, *The local Langlands conjectures for non-quasi-split groups*, Families of automorphic forms and the trace formula, Simons Symp., Springer, [Cham], 2016, pp. 217–257. MR 3675168 [43](#), [44](#)
- [Lag08] Nathalie Lagier, *Terme constant de fonctions sur un espace symétrique réductif p -adique*, J. Funct. Anal. **254** (2008), no. 4, 1088–1145. MR MR2381204 (2009d:22013) [11](#), [16](#)
- [LL15] Jean-Pierre Labesse and Erez Lapid, *Characters of G over local and global fields*, 2015, appendix to E. Lapid, Z. Mao, A conjecture on Whittaker-Fourier coefficients of cusp forms, J. Number Theory 146 (2015), 448–505. [63](#)
- [LR03] Erez M. Lapid and Jonathan D. Rogawski, *Periods of Eisenstein series: the Galois case*, Duke Math. J. **120** (2003), no. 1, 153–226. MR 2010737 [3](#), [10](#), [11](#), [12](#), [15](#)
- [Mat08] Nadir Matringe, *Distinction et fonctions L d’Asai pour les représentations génériques des groupes linéaires généraux sur les corps p -adiques*, PhD Thesis, UNIVERSITÉ PARIS. DIDEROT - Paris 7 (2008). [49](#)
- [Mat11] ———, *Distinguished generic representations of $GL(n)$ over p -adic fields*, Int. Math. Res. Not. IMRN (2011), no. 1, 74–95. MR 2755483 (2012f:22032) [46](#), [47](#)
- [Mat17a] ———, *Distinction of the Steinberg representation for inner forms of $GL(n)$* , Math. Z. **287** (2017), no. 3–4, 881–895. MR 3719517 [18](#)
- [Mat17b] Nadir Matringe, *Gamma factors of intertwining periods and distinction for inner forms of $gl(n)$* , 2017. [18](#)
- [MgVW87] Colette Mœglin, Marie-France Vignéras, and Jean-Loup Waldspurger, *Correspondances de Howe sur un corps p -adique*, Lecture Notes in Mathematics, vol. 1291, Springer-Verlag, Berlin, 1987. MR 1041060 [26](#)
- [MO18] Nadir Matringe and Omer Offen, *Gamma factors, root numbers, and distinction*, Canad. J. Math. **70** (2018), no. 3, 683–701. MR 3785415 [45](#)
- [Mok15] Chung Pang Mok, *Endoscopic classification of representations of quasi-split unitary groups*, Mem. Amer. Math. Soc. **235** (2015), no. 1108, vi+248. MR 3338302 [43](#)
- [Mui01] Goran Muić, *A proof of Casselman-Shahidi’s conjecture for quasi-split classical groups*, Canad. Math. Bull. **44** (2001), no. 3, 298–312. MR 1847492 (2002f:22015) [44](#), [45](#)
- [MW95] C. Mœglin and J.-L. Waldspurger, *Spectral decomposition and Eisenstein series*, Cambridge Tracts in Mathematics, vol. 113, Cambridge University Press, Cambridge, 1995, Une paraphrase de l’Écriture [A paraphrase of Scripture]. MR 1361168 (97d:11083) [5](#), [6](#), [30](#), [31](#), [32](#)
- [Off06] Omer Offen, *On symplectic periods of the discrete spectrum of GL_{2n}* , Israel J. Math. **154** (2006), 253–298. MR 2254544 [3](#)
- [Off17] ———, *On parabolic induction associated with a p -adic symmetric space*, J. Number Theory **170** (2017), 211–227. MR 3541705 [3](#), [11](#), [12](#), [13](#), [14](#), [15](#), [17](#), [18](#), [29](#), [54](#)
- [Pra] Dipendra Prasad, *A relative langlands correspondence and geometry of parameter spaces*, Preprint. [3](#), [41](#), [45](#), [46](#), [55](#), [63](#), [64](#)
- [Rod73] François Rodier, *Whittaker models for admissible representations of reductive p -adic split groups*, Harmonic analysis on homogeneous spaces (Proc. Sympos. Pure Math., Vol. XXVI, Williams Coll., Williamstown, Mass., 1972), Amer. Math. Soc., Providence, R.I., 1973, pp. 425–430. MR 0354942 (50 #7419) [38](#), [40](#)
- [Sch85] Winfried Scharlau, *Quadratic and Hermitian forms*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 270, Springer-Verlag, Berlin, 1985. MR MR770063 (86k:11022) [59](#), [60](#)
- [Ser94] Jean-Pierre Serre, *Cohomologie galoisienne*, fifth ed., Lecture Notes in Mathematics, vol. 5, Springer-Verlag, Berlin, 1994. MR 1324577 (96b:12010) [55](#), [57](#)
- [Sha91] Freydoon Shahidi, *Langlands’ conjecture on Plancherel measures for p -adic groups*, Harmonic analysis on reductive groups (Brunswick, ME, 1989), Progr. Math., vol. 101, Birkhäuser Boston, Boston, MA, 1991, pp. 277–295. MR 1168488 (93h:22033) [38](#)

- [SZ18] Allan J. Silberger and Ernst-Wilhelm Zink, *Langlands classification for L -parameters*, *J. Algebra* **511** (2018), 299–357. MR 3834776 [43](#), [44](#)
- [SX] Miyu Suzuki and Hang Xue, *Linear intertwining periods and epsilon dichotomy for linear models*, Preprint. <https://www.math.arizona.edu/~xuehang/> [18](#)
- [Wal59] G. E. Wall, *The structure of a unitary factor group*, *Inst. Hautes Études Sci. Publ. Math.* (1959), no. 1, 23 pp. (1959). MR MR0104764 (21 #3517) [45](#), [59](#)
- [Zel80] A. V. Zelevinsky, *Induced representations of reductive p -adic groups. II. On irreducible representations of $GL(n)$* , *Ann. Sci. École Norm. Sup. (4)* **13** (1980), no. 2, 165–210. MR 584084 (83g:22012) [45](#)

Email address: nadirmatringe@outlook.fr

UNIVERSITÉ DE POITIERS, POITIERS, FRANCE

Email address: offen@brandeis.edu

BRANDEIS UNIVERSITY, WALTHAM, MA