

# A NOTE ON THE ASYMPTOTICS FOR INCOMPLETE BETAFUNCTIONS

JAN-CHRISTOPH SCHLAGE-PUCHTA

ABSTRACT. We determine the asymptotic behaviour of certain incomplete Betafunctions.

## 1. INTRODUCTION AND RESULTS

For integers  $k, \ell \geq 1$  define

$$P_{k,\ell} = \frac{(k+\ell)!}{(k-1)! \ell!} \int_{k/(k+\ell)}^1 t^{k-1} (1-t)^\ell dt.$$

By repeated partial integration we obtain the representation

$$P_{k,\ell} = \left(\frac{\ell}{\ell+k}\right)^{k+\ell} \sum_{\nu=0}^{k-1} \binom{k+\ell}{\nu} \left(\frac{k}{\ell}\right)^\nu.$$

Vietoris[5] showed that  $P_{k,\ell} \leq \frac{1}{2}$ . Alzer and Kwong [1] proved that  $P_{k,\ell} \geq \frac{1}{4}$  holds for all  $k, \ell$ . Interest in bounds for  $P_{k,\ell}$  stems from application to statistics, see [4].

In this note we consider the asymptotic behaviour of  $P_{k,\ell}$ . We prove the following.

**Theorem 1.** *We have*

$$(1) \quad P_{k,\ell} = e^{-k} \sum_{\nu=0}^{k-1} \frac{k^\nu}{\nu!} + \mathcal{O}\left(\frac{k^2}{\ell}\right),$$

$$(2) \quad P_{k,\ell} = 1 - e^{-\ell} \sum_{\nu=0}^{\ell} \frac{\ell^\nu}{\nu!} + \mathcal{O}\left(\frac{\ell^2}{k}\right)$$

and

$$(3) \quad P_{k,\ell} = \frac{1}{2} + \mathcal{O}\left(\frac{1}{\sqrt{\min(k, \ell)}}\right).$$

Note that the first two estimates are good if one of the parameters  $k, \ell$  is rather small, whereas the third one gives information in the general case.

Comparing (1) and (2) with Vietoris' result that  $P_{k,\ell} \leq \frac{1}{2}$  we see that

$$\sum_{\nu=0}^{\ell-1} \frac{\ell^\nu}{\nu!} < \frac{1}{2} e^\ell < \sum_{\nu=0}^{\ell} \frac{\ell^\nu}{\nu!}.$$

Note that equality is impossible as  $e$  is transcendental. A more precise version of this inequality has been asked by Ramanujan (Question 294) and was answered by

Karamata [2]. For a detailed discussion of this result, Uhlmann's inequalities [4] and Vietoris bound we refer the reader to the historical notes by Vietoris [6].

From Theorem 1 we deduce the following.

**Corollary 1.** *The only accumulation points of the set  $\{P_{k,\ell} : k, \ell \in \mathbb{N}\}$  are  $\frac{1}{2}$  and the real numbers  $e^{-k} \sum_{\nu=1}^{k-1} \frac{k^\nu}{\nu!}$  and  $1 - e^{-\ell} \sum_{\nu=1}^{\ell} \frac{\ell^\nu}{\nu!}$ .*

*Proof.* Suppose that  $(k_n), (\ell_n)$  are integer sequences such that the pairs  $(k_n, \ell_n)$  are all distinct and that the sequence  $(P_{k_n, \ell_n})$  converges. If both  $k_n, \ell_n$  tend to infinity, then by (3) we have that  $P_{k_n, \ell_n}$  tends to  $\frac{1}{2}$ . If one of these sequences does not tend to infinity, then we can pass to a subsequence and assume that  $k_n$  or  $\ell_n$  is constant. Then our claim follows from (1) or (2).  $\square$

Together with Vietoris bound we obtain the following.

**Corollary 2.** *We have*

$$\lim_{\ell \rightarrow \infty} P_{k,\ell} \geq \frac{1}{2} - \frac{1}{\sqrt{2\pi k}}.$$

*Proof.* We have

$$\lim_{\ell \rightarrow \infty} (P_{k,\ell} + P_{\ell,k}) = 1 - \frac{k^k}{e^k k!},$$

using Stirling's formula in the form  $k! = \left(\frac{k}{e}\right)^k \sqrt{2\pi k} e^{\theta/12k}$  with  $0 \leq \theta \leq 1$ , and  $P_{\ell,k} \leq \frac{1}{2}$ , we conclude

$$\lim_{\ell \rightarrow \infty} P_{k,\ell} \geq \frac{1}{2} - \frac{k^k}{e^k k!} \geq \frac{1}{2} - \frac{1}{\sqrt{2\pi k}}.$$

$\square$

## 2. PRELIMINARY ESTIMATES

For the proof of (3) we use another representation of  $P_{k,\ell}$ , which is due to Raab [3].

**Theorem 2.** *We have  $P_{k,\ell} = U_{k,\ell} V_{k,\ell}$ , where*

$$U_{k,\ell} = \exp \int_0^\infty \frac{1}{t} \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \left( e^{-(k+\ell)t} - e^{-kt} - e^{-\ell t} \right) dt,$$

$$V_{k,\ell} = \frac{\sqrt{\ell}}{2\pi} \sum_{\nu=1}^{\infty} \frac{c_\nu(k/\ell)}{\sqrt{\nu(\nu+\ell)}},$$

and

$$c_\nu(x) = \exp \int_0^\infty \frac{1}{t} \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \left( e^{-\nu(1+x)t} - e^{-\nu xt} - e^{-\nu t} \right) dt.$$

We first compute the occurring integrals.

**Lemma 1.** *We have for  $x \geq 1$*

$$\int_0^\infty \frac{1}{t} \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) e^{-xt} dt = \frac{1}{12x} + \mathcal{O}\left(\frac{1}{x^2}\right)$$

and

$$\int_0^\infty \frac{1}{t} \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) e^{-xt} dt = \frac{1}{2} \log x + C + \mathcal{O}(x)$$

for  $0 < x \leq 1$ , where  $C$  is some constant.

*Proof.* The series expansion of  $e^x$  yields for  $t \rightarrow 0$

$$\frac{1}{t} \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) = \frac{1}{t^2} \left( \frac{1}{1 + t/2 + t^2/6 + \mathcal{O}(t^3)} - 1 + \frac{t}{2} \right) = 1/12 + \mathcal{O}(t).$$

For  $t \rightarrow \infty$  this expression tends to 0, in particular, it is bounded for all positive  $t$ . Hence for  $x \rightarrow \infty$  the integral in question becomes

$$\frac{1}{12} \int_0^\infty e^{-xt} dt + \mathcal{O} \left( \int_0^\infty t e^{-xt} dt \right) = \frac{1}{12x} + \mathcal{O} \left( \frac{1}{x^2} \right).$$

For  $x \rightarrow 0$  we use  $e^{-xt} = 1 - xt + \mathcal{O}(x^2 t^2)$  and obtain

$$\int_0^1 \frac{1}{t} \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) e^{-xt} dt = \int_0^1 \frac{1}{t} \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) dt + \mathcal{O}(x) = C_1 + \mathcal{O}(x)$$

and

$$\int_1^\infty \frac{1}{t(e^t - 1)} e^{-xt} dt = \int_1^\infty \frac{1}{t(e^t - 1)} dt - x \int_1^\infty \frac{1}{e^t - 1} dt + \mathcal{O}(x^2).$$

As a function of  $x$ , the integral  $\int_1^\infty \frac{e^{-xt}}{t^2} dt$  defines a function that is differentiable from the right in 0 and has bounded second derivative in  $(0, \infty)$ , hence, for  $x \geq 0$  this integral is  $1 + \mathcal{O}(x)$ .

Finally

$$\begin{aligned} \int_1^\infty \frac{e^{-xt}}{2t} dt &= \int_x^\infty \frac{e^{-s}}{2s} ds = \\ &= \int_x^1 \frac{dt}{2t} + \int_0^1 \frac{e^{-t} - 1}{2t} dt + \int_1^\infty \frac{e^{-t}}{2t} dt + \int_0^x \frac{1 - e^{-t}}{2t} dt \\ &= -\frac{1}{2} \log x + C_3 + \mathcal{O}(x). \end{aligned}$$

Combining these estimates our claim follows.  $\square$

From this we obtain

**Lemma 2.** *We have*

$$U_{k,\ell} = 1 - \frac{1}{12} \left( \frac{1}{k} + \frac{1}{\ell} - \frac{1}{k+\ell} \right) + \mathcal{O} \left( \frac{1}{\min(k, \ell)^2} \right).$$

*Proof.* From Lemma 1 we obtain

$$U_{k,\ell} = \exp \left( \frac{1}{12} \left( \frac{1}{k+\ell} - \frac{1}{k} - \frac{1}{\ell} \right) + \mathcal{O} \left( \frac{1}{\min(k, \ell)^2} \right) \right),$$

inserting the Taylor series for exp and using the fact that  $k, \ell \geq 1$  our claim follows.  $\square$

Next we compute  $c_\nu(x)$ .

**Lemma 3.** *If  $\nu x \geq 1$ , then*

$$c_\nu(x) = 1 + \frac{1}{12\nu(1+x)} - \frac{1}{12\nu x} + \frac{1}{12\nu} + \mathcal{O} \left( \frac{1}{\nu^2 \min(1, x^2)} \right).$$

*If  $\nu x \leq 1$ , then*

$$c_\nu(x) = K\sqrt{\nu x} + \mathcal{O} \left( \sqrt{\frac{x}{\nu}} + (\nu x)^{3/2} \right)$$

*for some constant  $K$ .*

*Proof.* If  $\nu x > 1$ , then we apply Lemma 1 to obtain

$$\begin{aligned} c_\nu(x) &= \exp\left(\frac{1}{12\nu(x+1)} - \frac{1}{12\nu x} + \frac{1}{12\nu} + \mathcal{O}\left(\frac{1}{\nu^2 \min(1, x^2)}\right)\right) \\ &= 1 + \frac{1}{12\nu(x+1)} - \frac{1}{12\nu x} + \frac{1}{12\nu} + \mathcal{O}\left(\frac{1}{\nu^2 \min(1, x^2)}\right). \end{aligned}$$

If  $\nu x < 1$ , then  $\nu(x+1) \geq 1$ , and we obtain

$$\begin{aligned} c_\nu(x) &= \exp\left(\frac{1}{12\nu(x+1)} + \frac{1}{2} \log(\nu x) + C + \frac{1}{12\nu} + \mathcal{O}\left(\frac{1}{\nu^2} + \nu x\right)\right) \\ &= K\sqrt{\nu x} + \mathcal{O}\left(\sqrt{\frac{x}{\nu}} + (\nu x)^{3/2}\right). \end{aligned}$$

□

We now compute  $V_{k,\ell}$ .

**Lemma 4.** *We have*

$$V_{k,\ell} = \frac{1}{2} + \mathcal{O}\left(\frac{1}{\sqrt{k}} + \frac{1}{\sqrt{\ell}}\right)$$

*Proof.* We have

$$\sum_{\nu \leq \ell/k} \frac{c_\nu(k/\ell)}{\sqrt{\nu(\nu+\ell)}} \ll \sum_{\nu \leq \ell/k} \frac{\sqrt{\nu k/\ell}}{\sqrt{\nu(\nu+\ell)}} \ll \frac{1}{\sqrt{\ell k}}$$

thus,

$$\begin{aligned} V_{k,\ell} &= \frac{\sqrt{\ell}}{2\pi} \sum_{\nu > \ell/k} \frac{1}{\sqrt{\nu(\nu+\ell)}} \left(1 + \frac{1}{12\nu \frac{k+\ell}{\ell}} - \frac{1}{12\nu \frac{k}{\ell}} + \frac{1}{12\nu} + \mathcal{O}\left(\frac{1}{\nu^2 \min(1, k^2/\ell^2)}\right)\right) + \mathcal{O}\left(\frac{1}{\sqrt{k}}\right) \\ &= \frac{\sqrt{\ell}}{2\pi} \sum_{\nu > \ell/k} \frac{1}{\sqrt{\nu(\nu+\ell)}} \left(1 - \frac{1}{12\nu \frac{k}{\ell}} + \mathcal{O}\left(\frac{1}{\nu^2 \min(1, k^2/\ell^2)}\right)\right) + \mathcal{O}\left(\frac{1}{\sqrt{k}} + \sum_{\nu > \ell/k} \frac{1}{\nu^{3/2} \sqrt{\ell}}\right) \\ &= \frac{\sqrt{\ell}}{2\pi} \sum_{\nu > \ell/k} \frac{1}{\sqrt{\nu(\nu+\ell)}} \left(1 - \frac{1}{12\nu \frac{k}{\ell}} + \mathcal{O}\left(\frac{1}{\nu^2 \min(1, k^2/\ell^2)}\right)\right) + \mathcal{O}\left(\frac{1}{\sqrt{k}} + \frac{1}{\sqrt{\ell}}\right) \end{aligned}$$

If  $k > \ell$  we obtain

$$\begin{aligned} V_{k,\ell} &= \frac{\sqrt{\ell}}{2\pi} \sum_{\nu=1}^{\infty} \frac{1}{\sqrt{\nu(\nu+\ell)}} + \mathcal{O}\left(\frac{1}{\sqrt{k}} + \frac{1}{\sqrt{\ell}} + \sqrt{\ell} \sum_{\nu=1}^{\infty} \frac{1}{\nu^{3/2}(\nu+\ell)}\right) \\ &= \frac{1}{2\pi} \int_0^{\infty} \frac{dt}{\sqrt{t(t+1)}} + \mathcal{O}\left(\frac{1}{\sqrt{k}} + \frac{1}{\sqrt{\ell}}\right) \\ &= \frac{1}{2} + \mathcal{O}\left(\frac{1}{\sqrt{\ell}}\right). \end{aligned}$$

We have

$$\frac{\sqrt{\ell}}{2\pi} \sum_{\nu > \ell} \frac{1}{\sqrt{\nu(\nu+\ell)}} \left(1 - \frac{1}{12\nu \frac{k}{\ell}} + \mathcal{O}\left(\frac{1}{\nu^2 \min(1, k^2/\ell^2)}\right)\right) = \frac{\sqrt{\ell}}{2\pi} \sum_{\nu > \ell} \frac{1}{\sqrt{\nu(\nu+\ell)}} + \mathcal{O}\left(\frac{1}{\sqrt{k}}\right),$$

and for  $\ell/k \leq N \leq \ell$

$$\sum_{N \leq \nu \leq 2N} \frac{1}{\nu^{3/2}(\nu + \ell)} \leq \frac{1}{\sqrt{N}\ell}$$

as well as

$$\sum_{N \leq \nu \leq 2N} \frac{1}{\nu^{5/2}(\nu + \ell)} \leq \frac{1}{N^{3/2}\ell}$$

and therefore

$$\sqrt{\ell} \sum_{\nu \geq \ell/k} \frac{1}{\sqrt{\nu}(\nu + \ell)} \cdot \frac{1}{\nu^{k/\ell}} = \frac{\ell^{3/2}}{k} \sum_{\nu \geq \ell/k} \frac{1}{\nu^{3/2}(\nu + \ell)} = \mathcal{O}\left(\frac{1}{\sqrt{k}}\right)$$

and

$$\sqrt{\ell} \sum_{\nu \geq \ell/k} \frac{1}{\sqrt{\nu}(\nu + \ell)} \cdot \frac{1}{\nu^2 \min(k^2/\ell^2)} = \frac{\ell^{5/2}}{k^2} \sum_{\nu \geq \ell/k} \frac{1}{\nu^{5/2}(\nu + \ell)} = \mathcal{O}\left(\frac{1}{\sqrt{k}}\right).$$

Using these estimates we obtain

$$\begin{aligned} \frac{\sqrt{\ell}}{2\pi} \sum_{\ell/k \leq \nu \leq \ell} \frac{1}{\sqrt{\nu}(\nu + \ell)} \left(1 - \frac{1}{12\nu^{k/\ell}} + \mathcal{O}\left(\frac{1}{\nu^2 \min(1, k^2/\ell^2)}\right)\right) = \\ \frac{\sqrt{\ell}}{2\pi} \sum_{\ell/k \leq \nu \leq \ell} \frac{1}{\sqrt{\nu}(\nu + \ell)} + \mathcal{O}\left(\frac{1}{\sqrt{k}}\right). \end{aligned}$$

For  $k \leq \ell$  we obtain

$$\begin{aligned} V_{k,\ell} &= \frac{\sqrt{\ell}}{2\pi} \sum_{\nu \geq \ell/k} \frac{1}{\sqrt{\nu}(\nu + \ell)} + \mathcal{O}\left(\frac{1}{\sqrt{k}}\right) \\ &= \frac{1}{2\pi} \int_{1/k}^{\infty} \frac{dt}{\sqrt{t}(t+1)} + \mathcal{O}\left(\frac{1}{\sqrt{k}}\right) \\ &= \frac{1}{2\pi} \int_0^{\infty} \frac{dt}{\sqrt{t}(t+1)} + \mathcal{O}\left(\frac{1}{\sqrt{k}}\right) \\ &= \frac{1}{2} + \mathcal{O}\left(\frac{1}{\sqrt{k}}\right), \end{aligned}$$

and the proof is complete.  $\square$

## 3. PROOF OF THEOREM 1

We first prove (1). Note that this inequality is only interesting if the error term is  $o(1)$ , in particular we may assume that  $k^2 < \ell$ . Under this assumption we have

$$\begin{aligned}
P_{k,\ell} &= \left(\frac{\ell}{\ell+k}\right)^{k+\ell} \sum_{\nu=0}^{k-1} \binom{k+\ell}{\nu} \left(\frac{k}{\ell}\right)^\nu \\
&= \sum_{\nu=0}^{k-1} \left(\frac{\ell}{\ell+k}\right)^{k+\ell-\nu} \frac{k^\nu}{\nu!} \exp\left(\sum_{\mu=1}^{\nu} \log \frac{k+\ell-\mu}{k+\ell}\right) \\
&= \sum_{\nu=0}^{k-1} \left(\frac{\ell}{\ell+k}\right)^{k+\ell-\nu} \frac{k^\nu}{\nu!} \exp\left(-\sum_{\mu=1}^{\nu} \frac{\mu}{k+\ell} + \mathcal{O}\left(\frac{\mu^2}{(k+\ell)^2}\right)\right) \\
&= \sum_{\nu=0}^{k-1} \left(\frac{\ell}{\ell+k}\right)^{k+\ell-\nu} \frac{k^\nu}{\nu!} \exp\left(\mathcal{O}\left(\frac{\nu^2}{\ell}\right)\right) \\
&= \sum_{\nu=0}^{k-1} \left(\frac{\ell}{\ell+k}\right)^{k+\ell} \frac{k^\nu}{\nu!} \exp\left(\mathcal{O}\left(\frac{k^2}{\ell}\right)\right) \\
&= \exp\left((k+\ell) \log \frac{\ell}{\ell+k}\right) \sum_{\nu=0}^{k-1} \frac{k^\nu}{\nu!} \exp\left(\mathcal{O}\left(\frac{k^2}{\ell}\right)\right) \\
&= \exp\left(-k + \mathcal{O}\left(\frac{k^2}{\ell}\right)\right) \sum_{\nu=0}^{k-1} \frac{k^\nu}{\nu!} \\
&= e^{-k} \sum_{\nu=0}^{k-1} \frac{k^\nu}{\nu!} + \mathcal{O}\left(\frac{k^2}{\ell}\right).
\end{aligned}$$

The proof of (2) is quite similar. Analogously to the previous case we may assume that  $\ell^2 < k$ .

$$\begin{aligned}
P_{k,\ell} &= \left(\frac{\ell}{\ell+k}\right)^{k+\ell} \left( \left(\frac{\ell+k}{\ell}\right)^{k+\ell} - \sum_{\nu=k}^{k+\ell} \binom{k+\ell}{\nu} \left(\frac{k}{\ell}\right)^\nu \right) \\
&= 1 - \left(\frac{\ell}{\ell+k}\right)^{k+\ell} \sum_{\mu=0}^{\ell} \binom{k+\ell}{\mu} \left(\frac{k}{\ell}\right)^{k+\ell-\mu} \\
&= 1 - \left(\frac{k}{\ell+k}\right)^{k+\ell} \sum_{\mu=0}^{\ell} \binom{k+\ell}{\mu} \left(\frac{\ell}{k}\right)^\mu \\
&= 1 - \exp\left(-\ell + \mathcal{O}\left(\frac{\ell^2}{k}\right)\right) \sum_{\mu=0}^{\ell} \frac{\ell^\mu}{\mu!} \left(\frac{k+\ell}{k}\right)^\mu \\
&= 1 - e^{-\ell} \sum_{\mu=0}^{\ell} \frac{\ell^\mu}{\mu!}.
\end{aligned}$$

Finally (3) follows from Theorem 2, Lemma 2 and Lemma 4.

## REFERENCES

- [1] H. Alzer, M. K. Kwong, Inequalities for combinatorial sums, *Arch. Math.* **108** (2017), 601–607.
- [2] J. Karamata, Sur quelques problèmes posés par Ramanujan, *J. Indian Math. Soc. (N.S.)* **24** (1960), 343–365.
- [3] W. Raab, Die Ungleichungen von Vietoris, *Monatsh. Math.* **98** (1984), 311–322.
- [4] W. Uhlmann, *Statistische Qualitätskontrolle: Eine Einführung* Leitfäden der angewandten Mathematik und Mechanik Band 7. B. G. Teubner Verlagsgesellschaft, Stuttgart 1966.
- [5] L. Vietoris, Über gewisse die unvollständige Betafunktion betreffende Ungleichungen, *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* **191** (1982), 85–92.
- [6] L. Vietoris, Geschichtliches über gewisse Ungleichungen, *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* **193** (1984), 319–321.