

On the distribution of non-primitive lattice points in the plane

by Nikolay Moshchevitin ¹

1. Introduction.

In [2] Erdős proved two following easy and beautiful theorems.

Theorem A. *For every $\varepsilon > 0$ there exist arbitrarily large positive integer x and $y \geq x$ such that*

$$\text{g.c.d.}(x + i, y + j) > 1$$

for all pairs i, j with

$$0 \leq i, j \leq (1 - \varepsilon) \left(\frac{\log x}{\log \log x} \right)^{1/2}.$$

Theorem B. *For a certain positive constant c for any positive integers $x \leq y$ there exist a pair of integers i, j with*

$$0 \leq i, j \leq c \frac{\log x}{\log \log x}$$

such that

$$\text{g.c.d.}(x + i, y + j) = 1.$$

Applying Theorem B in [3] Chalk and Erdős proved the following result on coprime inhomogeneous approximation to a real number.

Theorem C. *For any given $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and any real number η , there exists an absolute constant C such that*

$$q |q\alpha - \eta - r| \leq C \left(\frac{\log q}{\log \log q} \right)^2$$

is satisfied by infinitely many coprime integers (q, r) , $q \geq 1$.

In the last years the problem on inhomogeneous approximation with coprime numbers again became of interest (see [4] and the bibliography therein). In particular, Jitomirskaya and Liu proved the following result.

Theorem D. *For any constant C , there exists $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and η such that the inequality*

$$q |q\alpha - \eta - r| \leq C$$

only has finitely many coprime integer solutions (q, r) , $q \geq 1$.

¹This work is supported by the Russian Science Foundation under grant 19-11-00001.

In the present paper we improve on the result by Jitomirskaya and Liu. Combining the original argument of Theorem A by Erdős with the construction from [4] we show that there exist uncountably many $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $\eta \in \mathbb{R}$ such that

$$\inf_{(q,r) \in \mathbb{Z}^2, q > 100, (q,r)=1} q \frac{\log \log q}{\sqrt{\log q}} |q\alpha - \eta - r| > 0.$$

In Section 2 below we describe generalized Erdős' construction. In Sections 3, 4 we explain how to construct α and η . In Section 5 we give the exact formulation of our result (Theorem 1) and complete the proof.

2. Constructions with prime numbers.

Let k be a non-negative integer. We define the sets of integer points Ω_k in the following way. We put $\Omega_0 = \{(0, 0)\}$ and define

$$\Omega_k = \{(i, j) \in \mathbb{Z}^2 : \max(1, |i|) \cdot \max(1, |j|) \leq k\}.$$

Let ω_k be the cardinality of the set Ω_k . It is clear that $\omega_0 = 1, \omega_1 = 5, \omega_2 = 13$ and by simple counting argument we have

$$\omega_k = 4k \log k + O(k), \quad k \rightarrow \infty. \quad (1)$$

We arrange the sequence

$$p_1 = 2, p_2 = 3, p_3 = 5, p_4 = 7, \dots$$

of all prime numbers in the following way. Let $p_{0,0} = 2$ and if we have a permutation of the first ω_k prime numbers indexed as

$$p_{i,j}, \quad (i, j) \in \Omega_k$$

we define next $\omega_{k+1} - \omega_k$ numbers

$$p_{i,j}, \quad (i, j) \in \Omega_{k+1} \setminus \Omega_k$$

as an arbitrary fixed permutation of prime numbers

$$p_s, \quad \omega_k < s \leq \omega_{k+1}.$$

We put

$$P_k = \prod_{(i,j) \in \Omega_k} p_{i,j} = \prod_{s=1}^{\omega_k} p_s. \quad (2)$$

By Prime Number Theorem and (1) we have

$$\log P_k \sim 4k (\log k)^2, \quad k \rightarrow \infty. \quad (3)$$

Define

$$Q_{k+1} = \frac{P_{k+1}}{P_k}, \quad \text{g.c.d.}(P_k, Q_{k+1}) = 1. \quad (4)$$

For every k we define numbers

$$A_i^{[k]} = \prod_{s: (i,s) \in \Omega_k} p_{i,s}, \quad -k \leq i \leq k$$

and

$$B_j^{[k]} = \prod_{s: (s,j) \in \Omega_k} p_{s,j}, \quad -k \leq j \leq k.$$

It is clear that

$$A_i^{[k]} | A_i^{[k+1]}, \quad B_j^{[k]} | B_j^{[k+1]}, \quad \prod_{i=-k}^k A_i^{[k]} = \prod_{j=-k}^k B_j^{[k]} = P_k,$$

$$\text{g.c.d.}(A_{i_1}^{[k]}, A_{i_2}^{[k]}) = \text{g.c.d.}(B_{j_1}^{[k]}, B_{j_2}^{[k]}) = 1, \quad i_1 \neq i_2, \quad j_1 \neq j_2 \quad (5)$$

and

$$\text{g.c.d.}(A_i^{[k_1]}, B_j^{[k_2]}) = \begin{cases} p_{i,j}, & \max(1, |i|) \cdot \max(1, |j|) \leq \min(k_1, k_2) \\ 1, & \max(1, |i|) \cdot \max(1, |j|) > \min(k_1, k_2) \end{cases} \quad (6)$$

for any k_1, k_2 .

We define X_k, Y_k from the conditions

$$X_k + i \equiv 0 \pmod{A_i^{[k]}}, \quad -k \leq i \leq k,$$

$$Y_k + j \equiv 0 \pmod{B_j^{[k]}}, \quad -k \leq j \leq k,$$

$$P_k \leq X_k, Y_k \leq 2P_k - 1$$

by Chinese Remainder Theorem. It is clear that

$$\text{g.c.d.}(X_k + i, Y_k + j) \neq 1 \quad \text{for} \quad \max(1, |i|) \cdot \max(1, |j|) \leq k \quad (7)$$

and

$$X_{k+1} \equiv X_k \pmod{P_k}, \quad Y_{k+1} \equiv Y_k \pmod{P_k}. \quad (8)$$

We should take into account that

$$P_0 = X_0 = Y_0 = A_0^{[0]} = B_0^{[0]} = 2.$$

Also for non-collinear integer vectors $\mathbf{f}, \mathbf{g} \in \mathbb{Z}^2$ we need to consider affine lattices $\Lambda_k(\mathbf{f}, \mathbf{g})$ defined for even values of k as

$$\Lambda_{2l}(\mathbf{f}, \mathbf{g}) = \{\mathbf{z} = x\mathbf{f} + y\mathbf{g} : x \equiv X_l \pmod{P_l}, y \equiv Y_l \pmod{P_l}\}.$$

and for odd values of k as

$$\Lambda_{2l+1}(\mathbf{f}, \mathbf{g}) = \{\mathbf{z} = x\mathbf{f} + y\mathbf{g} : x \equiv Y_{l+1} \pmod{P_{l+1}}, y \equiv X_l \pmod{P_l}\}.$$

3. Continued fractions and integer points.

Now we consider irrational α defined by its continued fraction expansion

$$\alpha = [a_0; a_1, a_2, a_3, \dots, a_k, \dots], \quad a_0 \in \mathbb{Z}, \quad a_j \in \mathbb{Z}_+, \quad j = 1, 2, 3, \dots \quad (9)$$

We suppose in our construction that

$$a_{k+1} \equiv 0 \pmod{P_{[k/2]}} \quad (10)$$

and

$$a_{k+1} \geq W_k \cdot P_{[k/2]+1} \quad \text{with integer } W_k \rightarrow \infty, \quad k \rightarrow \infty. \quad (11)$$

Let

$$\frac{u_k}{v_k} = [a_0; a_1, a_2, a_3, \dots, a_k], \quad \text{g.c.d.}(v_k, u_k) = 1$$

be convergent fractions to α . Define primitive vectors

$$\mathbf{e}_k = (v_k, u_k) \in \mathbb{Z}^2, \quad k = 0, 1, 2, 3, \dots$$

It is clear that

$$\mathbf{e}_{k+1} = a_{k+1}\mathbf{e}_k + \mathbf{e}_{k-1}. \quad (12)$$

It would be convenient to define the lattice

$$\mathfrak{L}_k = \Lambda_k(\mathbf{e}_{k-1}, \mathbf{e}_k),$$

so for $l = [k/2]$ we have

$$\mathfrak{L}_{2l} = \{\mathbf{z} = x\mathbf{e}_{2l-1} + y\mathbf{e}_{2l} : x \equiv X_l \pmod{P_l}, \quad y \equiv Y_l \pmod{P_l}\},$$

$$\mathfrak{L}_{2l+1} = \{\mathbf{z} = x\mathbf{e}_{2l} + y\mathbf{e}_{2l+1} : x \equiv Y_{l+1} \pmod{P_{l+1}}, \quad y \equiv X_l \pmod{P_l}\}.$$

Now we define inductively integer points

$$\mathfrak{z}_k = (b_k, c_k) = \mathfrak{x}_k\mathbf{e}_{k-1} + \mathfrak{y}_k\mathbf{e}_k \in \mathfrak{L}_k, \quad k = 0, 1, 2, 3, \dots \quad (13)$$

Put $\mathfrak{z}_0 = (0, 0)$. Then, if $\mathfrak{z}_k \in \mathfrak{L}_k$ is defined, we define $\mathfrak{z}_{k+1} \in \mathfrak{L}_{k+1}$ by the following procedure. We take $l = [k/2]$ and define

$$\mathfrak{z}_{k+1} = \mathfrak{z}_k + P_l\lambda_{k+1}\mathbf{e}_k = \mathfrak{x}_k\mathbf{e}_{k-1} + (\mathfrak{y}_k + P_l\lambda_{k+1})\mathbf{e}_k, \quad (14)$$

so

$$b_{k+1} = b_k + P_l\lambda_{k+1}v_k \quad \text{and} \quad c_{k+1} = c_k + P_l\lambda_{k+1}u_k. \quad (15)$$

The value of $\lambda_{k+1} \in \mathbb{Z}_+$ will be defined later in (21).

Taking into account equality (12), by (13,14) we can write

$$\mathfrak{y}_{k+1} = \mathfrak{x}_k \equiv \begin{cases} X_l \pmod{P_l}, & \text{if } k \text{ even} \\ Y_{l+1} \pmod{P_{l+1}}, & \text{if } k \text{ odd} \end{cases}. \quad (16)$$

and

$$\mathfrak{X}_{k+1} = \mathfrak{Y}_k + P_l \lambda_{k+1} - \mathfrak{X}_k a_{k+1}, \quad (17)$$

By (10) we have $\mathfrak{X}_{k+1} \equiv \mathfrak{Y}_k \pmod{P_l}$. Since $\mathfrak{Z}_k \in \mathfrak{L}_k$, by (8) we see that

$$\mathfrak{X}_{k+1} \equiv \begin{cases} Y_l \equiv Y_{l+1}, & \text{if } k \text{ even} \\ X_l \equiv X_{l+1}, & \text{if } k \text{ odd} \end{cases} \pmod{P_l}. \quad (18)$$

On the other hand $\text{g.c.d.}(P_l, Q_{l+1}) = 1$ (see (4)). So we can find $\lambda_{k+1}^* \in \{1, 2, \dots, Q_{l+1}\}$ such that for every $\lambda_{k+1} \equiv \lambda_{k+1}^* \pmod{Q_{l+1}}$ one has

$$\mathfrak{X}_{k+1} \equiv \begin{cases} Y_{l+1}, & \text{if } k \text{ even} \\ X_{l+1}, & \text{if } k \text{ odd} \end{cases} \pmod{Q_{l+1}}. \quad (19)$$

As $P_{l+1} = P_l Q_{l+1}$ equalities (18,19) imply

$$\mathfrak{X}_{k+1} \equiv \begin{cases} Y_{l+1}, & \text{if } k \text{ even} \\ X_{l+1}, & \text{if } k \text{ odd} \end{cases} \pmod{P_{l+1}}. \quad (20)$$

We see that equalities (16) and (20) ensure $\mathfrak{Z}_{k+1} \in \mathfrak{L}_{k+1}$.

Finally we take λ_{k+1} of the form $\lambda_{k+1} = \lambda_{k+1}^* + \lambda Q_{l+1}$, $\lambda \in \mathbb{Z}$ to satisfy

$$\left| \lambda_{k+1} - \frac{a_{k+1}}{2P_l} \right| \leq \frac{Q_{l+1}}{2}, \quad (21)$$

so λ is the nearest integer to $\frac{a_{k+1}}{2P_{l+1}} - \frac{\lambda_{k+1}^*}{Q_{l+1}}$.

We have defined $\mathfrak{Z}_{k+1} = (b_{k+1}, c_{k+1}) \in \mathfrak{L}_{k+1}$ of the form (14).

Here we should not that from (15) and (21) it follows that

$$\left| b_{k+1} - \left(b_k + \frac{v_{k+1}}{2} \right) \right| \leq P_{l+1} v_k$$

and by induction we have

$$\left| b_{k+1} - \frac{v_{k+1}}{2} \right| \leq 2P_{l+1} v_k. \quad (22)$$

4. The value of η .

From (15) we see that

$$b_k = \sum_{j=1}^k P_{[(j-1)/2]} \lambda_j v_{j-1}, \quad c_k = \sum_{j=1}^k P_{[(j-1)/2]} \lambda_j u_{j-1}.$$

Now we define

$$\eta = \lim_{k \rightarrow \infty} (b_k \alpha - c_k) = \sum_{j=1}^{\infty} P_{[(j-1)/2]} \lambda_j (v_{j-1} \alpha - u_{j-1}). \quad (23)$$

Of course, we should explain why the series in (23) converges. We use the estimate

$$\frac{1}{v_j + v_{j-1}} \leq |v_{j-1}\alpha - u_{j-1}| \leq \frac{1}{v_j},$$

By means of the upper bound here we see that

$$|P_{[(j-1)/2]+1}\lambda_j(v_{j-1}\alpha - u_{j-1})| \leq \frac{P_{[(j-1)/2]}\lambda_j}{v_j} \leq \frac{1}{2v_{j-1}} + \frac{P_{[(j-1)/2]}}{2v_j} \leq \left(\frac{1}{2} + \frac{1}{2W_j}\right) \frac{1}{v_{j-1}},$$

by (21) and (11,12), as $\frac{v_j}{v_{j-1}} \geq a_j \geq W_j P_{[(j-1)/2]}$. So the series from (23) converges absolutely since $v_j \geq \left(\frac{1+\sqrt{5}}{2}\right)^{j-1}$. Moreover, it is easy to see that the absolute values of the summands from series (23) decrease monotonically to zero. Futhermore, as $P_{[(j-1)/2]}\lambda_j$ is positive and $u_{j-1} - v_{j-1}\alpha$ changes its sign we see that

$$|\eta - (b_k\alpha - c_k)| < \left(\frac{1}{2} + \frac{1}{2W_k}\right) \frac{1}{v_k}. \quad (24)$$

The lower bound for the approximation is given by

$$\begin{aligned} |\eta - (b_k\alpha - c_k)| &> P_{[k/2]}\lambda_{k+1}|u_k - v_k\alpha| - P_{[(k+1)/2]}\lambda_{k+2}|u_{k+1} - v_{k+1}\alpha| \geq \\ &\geq \frac{P_{[k/2]}\lambda_{k+1}}{v_{k+1} + v_k} - \frac{P_{[(k+1)/2]}\lambda_{k+2}}{v_{k+2}} \geq \left(\frac{1}{2} - \frac{2}{W_k}\right) \frac{1}{v_k} \end{aligned} \quad (25)$$

(for the second summand we used here (24)). In addition by (12) and (11) we see that

$$|\eta - (b_k\alpha - c_k)| > |\eta - (b_{k+1}\alpha - c_{k+1})|.$$

We should note that the differences

$$\eta - (b_k\alpha - c_k) \quad \text{and} \quad \eta - (b_{k+1}\alpha - c_{k+1})$$

have different signs meanwhile the differences

$$\eta - (b_k\alpha - c_k) \quad \text{and} \quad \eta - ((b_{k+1} - v_k) - (c_{k+1} - u_k)\alpha)$$

have the same sign.

5. The main result.

Now we are able to give the exact formulation of our main result.

Theorem 1. *Suppose that the partial quotients of the continued fraction expansion (9) for α satisfy (10) and (11). Suppose that with some positive γ the inequality*

$$a_k \leq \exp(\gamma k(\log k)^2) \quad (26)$$

holds. Let η be defined by (23). Then for any primitive point $(q, r) \in \mathbb{Z}^2$ with $\text{g.c.d.}(q, r) = 1$ and $q > 100$ one has

$$q|q\alpha - \eta - r| > c \frac{\sqrt{\log q}}{\log \log q}, \quad (27)$$

with certain positive c .

The proof of Theorem 1 is close to original argument from [4], as well as to the argument by Worley [5] (see also [1]).

We suppose that constants in symbols \gg and \asymp below depend on γ .

Remark 1. It is clear that for any α under conditions (10,11,26) it is possible to find uncountably many η satisfying the conclusion of Theorem 1.

Remark 2. We can choose a_k satisfying (10,11) and (26) because of (3). From the inequalities $v_k \leq 2^k \prod_{j=1}^k a_k$ and (26) we deduce that

$$k \gg \frac{\sqrt{\log v_k}}{\log \log v_k}.$$

Moreover, from (22) it follows that $b_k \asymp v_k$ and so by (11) and (26) we have

$$k \gg \frac{\sqrt{\log b_k}}{\log \log b_k} \asymp \frac{\sqrt{\log b_{k+1}}}{\log \log b_{k+1}}. \quad (28)$$

Proof of Theorem 1.

From (28) we see that it is enough to prove the inequality

$$q |q\alpha - \eta - r| \gg k \quad (29)$$

for every primitive point $\mathbf{z} = (q, r) \in \mathbb{Z}^2$ satisfying

$$b_k \leq q < b_{k+1}. \quad (30)$$

Each point $\mathbf{z} = (q, r) \in \mathbb{Z}^2$ can be written in the form

$$\mathbf{z} = \mathfrak{Z}_k + x\mathbf{e}_{k-1} + y\mathbf{e}_k \quad (31)$$

and in another form

$$\mathbf{z} = \mathfrak{Z}_{k+1} + x'\mathbf{e}_k + y'\mathbf{e}_{k+1} \quad (32)$$

with integer x, y, x', y' .

We should note that if an integer point $\mathbf{z} = (q, r)$ can be represented in the form

$$\mathbf{z} = \mathfrak{Z}_k + x\mathbf{e}_{k-1} + y\mathbf{e}_k, \quad \text{with} \quad \max(1, |x|) \cdot \max(1, |y|) \leq \frac{k}{2}, \quad (33)$$

then by (7) and by the condition $\mathfrak{Z}_k \in \mathfrak{L}_k$ we see that for all \mathbf{z} of the form (33) we have $(\mathfrak{X}_k + x, \mathfrak{Y}_k + y) \neq 1$ and so $(q, r) \neq 1$.

Now we consider integer points $\mathbf{z} = (q, r)$ under the condition (30) and such that simultaneously

$$\max(1, |x|) \cdot \max(1, |y|) > \frac{k}{2} \quad \text{in (31), and} \quad \max(1, |x'|) \cdot \max(1, |y'|) > \frac{k}{2} \quad \text{in (32)}. \quad (34)$$

For these points we shall prove that (29) holds and this will complete the proof of Theorem 1.

Put $l = \lfloor \frac{k}{2} \rfloor$ and consider two cases:

case 1⁰: $\mathbf{z} = (q, r) = \mathfrak{Z}_k + y\mathbf{e}_k$ with $l \leq y \leq P_k \lambda_{k+1} - l$;

and

case 2⁰: $\mathbf{z} = (q, r) = \mathfrak{Z}_k + x\mathbf{e}_{k-1} + y\mathbf{e}_k$ with $x \neq 0$.

As $\mathfrak{Z}_{k+1} = (b_{k+1}, c_{k+1}) = \mathfrak{Z}_k + P_k \lambda_{k+1} \mathbf{e}_k$, in **case 1⁰** inequality (30) is satisfied automatically. In **case 2⁰** we should assume (30).

In **case 1⁰**,

$$q = b_k + yv_k, \quad \|\eta - q\alpha\| = \eta - ((b_k + yv_k)\alpha - (c_k + yu_k))$$

and the differences

$$\eta - ((b_k + lv_k)\alpha - (c_k + lu_k)) \quad \text{and} \quad \eta - ((b_{k+1} - lv_k)\alpha - (c_k + lu_k))$$

have the same sign (see the very end of Section 4).

We are interested in getting lower bound for approximation for primitive points \mathbf{z} , and it is enough to consider only those \mathbf{z} for which (34) holds. So we can assume that

$$b_k + lv_k \leq q \leq b_{k+1} - lv_k.$$

For

$$\mathbf{z} = (q, r) = (b_k + lv_k, c_k + lu_k)$$

we have $q = b_k + lv_k > lv_k$ and

$$q|\eta - q\alpha - r| \geq lv_k(|\eta - (b_k\alpha + c_k)| - l|v_k\alpha - u_k|) \geq \frac{l}{5} - \frac{l^2 v_k}{v_{k+1}} \geq \frac{l}{10} - \frac{l^2}{a_{k+1}} \geq \frac{l}{20}$$

as $|v_k\alpha - u_k| \leq \frac{1}{v_{k+1}}$ and by (25) and (11).

For

$$\mathbf{z} = (q, r) = (b_{k+1} - lv_k, c_{k+1} - lu_k)$$

we have $q = b_{k+1} - lv_k \geq b_{k+1}/2 \geq v_{k+1}/8$ (we use (22)) and

$$q|\eta - q\alpha - r| \geq \frac{v_{k+1}}{8}(l|v_k\alpha - u_k| - |\eta - (b_{k+1}\alpha + c_{k+1})|) \geq \frac{l}{10} - \frac{1}{8} \geq \frac{l}{20},$$

for $l \geq 3$, as we have inequalities $|v_k\alpha - u_k| \geq \frac{1}{v_{k+1} + v_k}$ and (24).

Now we see that for $\mathbf{z} = (q, r) = (c_k + yu_k, b_k + yv_k)$ the inequality

$$q|\eta - q\alpha - r| = (b_k + yv_k)|\eta - ((c_k + yu_k) - (b_k + yv_k)\alpha)| \geq \frac{l}{20}$$

is true for any intermediate value of y satisfying $l \leq y \leq P_k \lambda_{k+1} - l$ by convexity argument. Now we take into account (28) and (30). In **case 1⁰** for primitive \mathbf{z} inequality (29) is proven.

The proof in **case 2⁰** is based on the same ideas. In this case we have $x \neq 0$ and for any $\mathbf{z} = (q, r)$ under consideration from the lower bound (25) we deduce the inequality

$$|\eta - q\alpha - r| \geq \frac{|x|}{3v_k}. \tag{35}$$

By (22) and (30)

$$q \geq b_k \geq v_k/4. \quad (36)$$

If $|x| \geq l$, then by (36) we see that

$$q |\eta - q\alpha - r| \geq \frac{l}{12}.$$

Suppose $0 \neq |x| < l$. As we deal with primitive \mathbf{z} we should not consider \mathbf{z} which satisfy (33). So $y > \frac{l}{|x|}$ and instead of (36) we have

$$q \geq yv_k - b_k - lv_{k-1} \geq \frac{yv_k}{4} > \frac{lv_k}{4|x|}. \quad (37)$$

Now inequalities (35,37) give us

$$q |\eta - q\alpha - r| \geq \frac{l}{12}.$$

In **case 2** inequality (29) is proven also.

Theorem 1 is proven.

The author thanks Oleg German and Vasili Neckrasov for fruitful discussions.

References

- [1] A. Dujella, *Continued fractions and RSA with small secret exponent*, Tatra Mt. Math. Publ. **29** (2004), 101-112.
- [2] P. Erdős, *On an elementary problem in number theory*, Canadian Math. Bull. 1, (1958), 5-8.
- [3] J. Chalk, P. Erdős, *On the distribution of primitive lattice points in the plane*, Canad. Math. Bull. 2 (1959) 91-96.
- [4] Svetlana Jitomirskaya, Wencai Liu, *Inhomogeneous Diophantine approximation in the coprime setting*, Advances in Mathematics 355 (2019), 06773.
- [5] R.T. Worley, *Estimating $|\alpha - p/q|$* , Austral. Math. Soc. (Series A) **31** (1981), 202-206.

author:

Nikolay G. Moshchevitin
Steklov Mathematical Institute,
ul. Gubkina 8, Moscow, 119991
Russia