

# HYPONELLIPTIC ENTROPY DISSIPATION FOR STOCHASTIC DIFFERENTIAL EQUATIONS

QI FENG\* AND WUCHEN LI†

**ABSTRACT.** We study the Lyapunov convergence analysis for degenerate and non-reversible stochastic differential equations (SDEs). We apply the Lyapunov method towards the Fokker-Planck equation, in which the Lyapunov functional is chosen as a weighted relative Fisher information functional. We derive a structure condition under which the Lyapunov constant is obtained explicitly. Given the positive Lyapunov constant, we prove the exponential convergence result for the probability density function of SDEs towards its invariant distribution in  $L_1$  norm. Two examples are presented, including underdamped Langevin dynamics with variable diffusions and 3 oscillator chain model with nearest-neighbor couplings.

## 1. INTRODUCTION

Consider an Itô stochastic differential equation (SDE)

$$dX_t = b(X_t)dt + \sqrt{2}a(X_t)dB_t, \quad (1)$$

where  $X_t \in \mathbb{R}^{n+m}$  is a stochastic process with dimensions  $m, n \in \mathbb{Z}_+$ ,  $b \in \mathbb{R}^{n+m}$  is a smooth drift vector field,  $a \in \mathbb{R}^{(n+m) \times n}$  is a smooth diffusion matrix, and  $B_t$  is a standard Brownian motion in  $\mathbb{R}^n$ .

We study long time dynamical behaviors of SDE (1). How fast does the probability density function of SDE (1) converge to its invariant distribution? The convergence behavior of SDE (1) plays essential roles in both theory and applications. Theoretically, it is a core problem in probability [7, 25], non-equilibrium dynamics [16, 26, 29, 40, 48], differential geometry [10, 11] and ergodic theory [8, 32, 36, 43]. In applications, the convergence rate is useful in studying long time behaviors of molecular dynamics [17, 33, 45]. Nowadays, the convergence rate is important in estimating the speed of Markov-Chain-Monte-Carlo methods (MCMC). This is an emerging issue in AI (artificial intelligence) and Bayesian sampling algorithms [17, 19, 15, 45].

In current studies, when diffusion matrix  $a$  is non-degenerate and drift vector field  $b$  is a gradient direction, SDE (1) can be viewed as a reversible diffusion process, whose

---

\* Department of Mathematics, University of Michigan, Ann Arbor, MI 48109. Email: qif@umich.edu.

† Department of Mathematics, University of South Carolina, Columbia, SC 29208. Email: wuchen@mailbox.sc.edu.

†Wuchen Li is supported by a start-up funding in University of South Carolina.

**Key words:** Hypocoellipticity; Information Gamma calculus; Entropy dissipation; Underdamped Langevin dynamics; 3 oscillator chain model.

**MSC:** 37L45, 58J65, 60D05, 60H10.

convergence rate has been studied in various ways. A typical way is the entropy method, a.k.a. the Lyapunov method in probability density space; see [3]. In the derivation of entropy method, one important step is known as a Bakry-Émery Gamma calculus. Using it, one can derive the convergence rate of SDE (1) by a Bochner’s formula. However, the Bochner’s type formula often fails when matrix  $a$  is a degenerate matrix function of  $X_t$ . Here SDE (1) is a degenerate diffusion process with a possible irreversible drift vector field. The current entropy method [48] can not be applied directly.

In this paper, we derive a convergence analysis for SDE (1) by a modified entropy dissipation method. Here we design a weighted relative Fisher information functional as a Lyapunov functional. By studying the dissipation of weighted Fisher information functional, we introduce a structure condition for deriving an “information Bochner type formula”; see Assumption 1. We formulate an algebraic tensor/Hessian matrix, namely “Ricci curvature type tensor”; see details in Theorem 3. The smallest eigenvalue of Hessian matrix characterizes the convergence behavior of SDE (1) in term of  $L_1$  distance. As examples, we present explicit convergence rates for variable diffusion coefficient underdamped Langevin dynamics and 3 oscillator chain model.

In literature, there are many studies for convergence behaviors of degenerate SDEs, [2, 13, 17, 20, 27, 29, 30, 44, 15, 39, 46, 48]. Previous works have established the convergence rates under various metrics, e.g.  $H^1$  or  $L^2$  metric. In particular, [6] has applied a modified entropy dissipation method into the degenerate drift-diffusion process. They mainly work with a constant viscosity. Besides, the Gamma  $z$  calculus has been invented in [11], and it has been applied to study underdamped Langevin dynamics [10, 12]. A variable vector field  $z$  is introduced to address the degenerate SDE (8). The Gamma  $z$  method establishes the convergence in  $H^1$  metric, which requires an iterative symmetry condition between matrix functions  $a$  and  $z$ . Compared to previous methods, we derive a structure condition for [22]; see Assumption 1. And we derive explicit “Hessian matrices” to handle both non-gradient drift vectors and degenerate diffusion matrices. Technically speaking, we apply the second-order calculus in probability density space embedded with an optimal transport type metric [31, 34, 49]. Our computation extends the second-order derivative of Kullback–Leibler divergence in sub-Riemannian density manifold [22, 23].

We organize the paper below. In section 2, we present the main result of this paper. We present a matrix, whose smallest eigenvalue characterizes the convergence behavior of SDE (8). We apply the main results towards underdamped Langevin dynamics with variable diffusion matrix in section 3, and 3 oscillator chain model in section 4. All proofs are presented in section 5. In particular, we derive an “information Bochner type formula”.

## 2. MAIN RESULT

In this section, we present the main result of this paper.

**2.1. Settings.** Assume that the collections of vector fields  $\{b = a_0, a_1, \dots, a_n\}$  satisfy the following weak Hörmander condition [14]:

$$\text{Span}\left\{a_1(x), \dots, a_n(x), [a_{i_1}, \dots, [a_{i_{k-1}}, a_{i_k}] \dots](x), 0 \leq i_1, \dots, i_k \leq n, k \geq 2\right\} = \mathbb{R}^{n+m},$$

where  $[\cdot, \cdot]$  represents a Lie bracket operation. The weak Hörmander condition means that the Lie algebra generated by the vector fields  $\{b, a_1, \dots, a_n\}$  is of full rank at every point  $x \in \mathbb{R}^{n+m}$ . This condition [28, 14] ensures the existence of smooth probability density function of SDE (1). Consider the Fokker-Planck equation of SDE (1)

$$\partial_t p(t, x) = -\nabla_x \cdot (p(t, x)b(t, x)) + \sum_{i=1}^{n+m} \sum_{j=1}^{n+m} \frac{\partial^2}{\partial x_i \partial x_j} \left( (a(x)a(x)^\top)_{ij} p(t, x) \right), \quad (2)$$

where  $p(t, x)$  is the probability density function of SDE (1) with a smooth initial condition

$$p_0(x) = p(0, x), \quad \int p_0(x) dx = 1, \quad p_0(x) \geq 0.$$

Here  $\int$  denotes the integration over the entire spatial domain  $\mathbb{R}^{n+m}$ . We further assume that  $p(t, x)$  is smooth w.r.t both time and spatial variables  $t, x$ , respectively, and stays positive for all  $t \in [0, \infty)$ . We assume that the SDE (1) has a unique invariant distribution  $\pi > 0$ , which satisfies the equilibrium of equation (2):

$$-\nabla_x \cdot (\pi(x)b(x)) + \sum_{i=1}^{n+m} \sum_{j=1}^{n+m} \frac{\partial^2}{\partial x_i \partial x_j} \left( (a(x)a(x)^\top)_{ij} \pi(x) \right) = 0.$$

We also assume that  $\pi \in C^2(\mathbb{R}^{n+m})$  and has an analytical formula. In this paper, we study the convergence behavior of  $p(t, x)$  towards the invariant distribution  $\pi(x)$ .

**Decomposition.** The convergence result is based on the following decomposition/reformulation of Fokker-Planck equation (2). Consider

$$\partial_t p(t, x) = \nabla \cdot (p(t, x)a(x)a(x)^\top \nabla \log \frac{p(t, x)}{\pi(x)}) + \nabla \cdot (p(t, x)\gamma(x)), \quad (3)$$

where  $\gamma \in C^\infty(\mathbb{R}^{n+m})$  is a vector field satisfying

$$\gamma(x) := a(x)a(x)^\top \nabla \log \pi(x) - b(x) + \left( \sum_{j=1}^{n+m} \frac{\partial}{\partial x_j} (a(x)a(x)^\top)_{ij} \right)_{1 \leq i \leq n+m}.$$

It is a straightforward calculation to check that equation (3) is equivalent to equation (2). Under this decomposition, we notice the fact that

$$\nabla \cdot (\pi(x)\gamma(x)) = 0.$$

We observe that the term  $\nabla \cdot (paa^\top \nabla \log \frac{p}{\pi})$  corresponds to the “reversible” and “hypoelliptic” (degeneracy) component of SDE (1)’s Kolmogorov forward operator, while the term  $\nabla \cdot (p\gamma)$  represents the “irreversible” component in Fokker-Planck equation (2).

In later on proofs, we develop a Lyapunov analysis to study the dynamical behavior of SDE (1). Following decomposition (3), we develop a exponential convergence result for equation (2). Here the convergence rate explicitly addresses both degenerate and irreversible components of equation (2).

**2.2. Structure condition and Hessian matrix.** We present two assumptions for the main result. Firstly, denote two smooth matrix functions as

$$a \in \mathbb{R}^{(n+m) \times n}, \quad \text{and} \quad z \in \mathbb{R}^{(n+m) \times m}.$$

For a vector field  $U \in C^\infty(\mathbb{R}^{n+m})$ , denote

$$\begin{aligned} a_k^\top U &= \sum_{k'=1}^n a_{kk'}^\top U_{k'}, & (a^\top \nabla) \circ (a^\top U) &= \sum_{k=1}^n a_k^\top \nabla (a_k^\top U), \\ a^\top U &= \sum_{k=1}^n a_k^\top U, & a_k^\top \nabla a_i^\top U &= \sum_{k',i'=1}^n a_{kk'}^\top \frac{\partial a_{ii'}^\top}{\partial x_{k'}} U_{i'}, \end{aligned}$$

where  $a^\top$  is the transpose of  $a$ , i.e.  $a_{kk'}^\top = a_{k'k}$ .

**Assumption 1.** For any  $i \in \{1, \dots, n\}$  and  $k \in \{1, \dots, m\}$ , we assume

$$z_k^\top \nabla a_i^\top \in \text{Span}\{a_1^\top, \dots, a_n^\top\}. \quad (4)$$

In other words, for  $\hat{i} \in \{1, \dots, n+m\}$  and  $k' \in \{1, \dots, n+m\}$ , we assume that there exists functions  $\lambda_l^{k'i}$ , for  $l \in \{1, \dots, n\}$ , such that

$$z_k^\top \nabla a_i^\top := \sum_{\hat{i}, k'=1}^{n+m} z_{k\hat{i}}^\top \frac{\partial}{\partial x_{k'}} a_{\hat{i}\hat{i}}^\top = \sum_{\hat{i}, k'=1}^{n+m} \sum_{l=1}^n z_{k\hat{i}}^\top \lambda_l^{k'i} a_{\hat{i}l}^\top. \quad (5)$$

Secondly, we define the following matrix functions.

**Definition 1** (Hessian matrix). For  $U \in C^\infty(\mathbb{R}^{n+m})$ , a constant  $\beta \in \mathbb{R}$  and a matrix function  $z \in \mathbb{R}^{(n+m) \times (n+m)}$ , define a matrix function  $\mathfrak{R}: \mathbb{R}^{n+m} \rightarrow \mathbb{R}^{(n+m) \times (n+m)}$  as

$$\begin{aligned} \mathfrak{R}(U, U) &= (\mathfrak{R}_a + \mathfrak{R}_z + \mathfrak{R}_\pi)(U, U) - \Lambda_1^\top \Lambda_1 - \Lambda_2^\top \Lambda_2 + D^\top D + E^\top E \\ &\quad + (\beta \mathfrak{R}_{\mathcal{L}_a} + (1 - \beta) \mathfrak{R}_{\gamma_a} + \mathfrak{R}_{\gamma_z})(U, U). \end{aligned}$$

We denote  $\mathfrak{R}(x) \in \mathbb{R}^{n+m}$ , such that  $U^\top \mathfrak{R}(x) U = \mathfrak{R}(U, U)$ . We also define the following bi-linear forms.

$$\begin{aligned} \mathfrak{R}_a(U, U) &= \sum_{i,k=1}^n a_i^\top \nabla a_i^\top \nabla a_k^\top U (a_k^\top U) + \sum_{i,k=1}^n a_i^\top a_i^\top \nabla^2 a_k^\top U (a_k^\top U) \\ &\quad - \sum_{i,k=1}^n a_k^\top \nabla a_i^\top \nabla a_i^\top U (a_k^\top U) - \sum_{i,k=1}^n a_k^\top a_i^\top \nabla^2 a_i^\top U (a_k^\top U) \\ &\quad + \sum_{i=1}^n \sum_{\hat{k}=1}^{n+m} \left[ (a a^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} a_i^\top U - a_i^\top \nabla (a a^\top \nabla \log \pi)_{\hat{k}} U_{\hat{k}} \right] a_i^\top U \\ &\quad + \nabla a \circ \left( \sum_{k=1}^n [a^\top \nabla a_k^\top U - a_k^\top \nabla a^\top U] a_k^\top U \right) - \langle (a^\top \nabla^2 a \circ (a^\top U)), a^\top U \rangle_{\mathbb{R}^n}, \end{aligned}$$

$$\begin{aligned}
\mathfrak{R}_z(\mathbf{U}, \mathbf{U}) &= \sum_{i=1}^n \sum_{k=1}^m a_i^\top \nabla a_i^\top \nabla z_k^\top \mathbf{U} (z_k^\top \mathbf{U}) + \sum_{i,k=1}^n a_i^\top a_i^\top \nabla^2 z_k^\top \mathbf{U} (z_k^\top \mathbf{U}) \\
&\quad - \sum_{i=1}^n \sum_{k=1}^m z_k^\top \nabla a_i^\top \nabla a_i^\top \mathbf{U} (z_k^\top \mathbf{U}) - \sum_{i,k=1}^n z_k^\top a_i^\top \nabla^2 a_i^\top \mathbf{U} (z_k^\top \mathbf{U}) \\
&\quad + \sum_{k=1}^m \sum_{\hat{k}=1}^{n+m} \left[ (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} z_k^\top \mathbf{U} - z_k^\top \nabla (aa^\top \nabla \log \pi)_{\hat{k}} \mathbf{U}_{\hat{k}} \right] z_k^\top \mathbf{U} \\
&\quad + \nabla a \circ \left( \sum_{k=1}^m \left[ a^\top \nabla z_k^\top \mathbf{U} - z_k^\top \nabla a^\top \mathbf{U} \right] z_k^\top \mathbf{U} \right) - \langle (z^\top \nabla^2 a \circ (a^\top \mathbf{U})), z^\top \mathbf{U} \rangle_{\mathbb{R}^m},
\end{aligned}$$

$$\begin{aligned}
\mathfrak{R}_\pi(\mathbf{U}, \mathbf{U}) &= 2 \sum_{k=1}^m \sum_{i=1}^n \left[ \nabla z_k^\top z_k^\top \nabla a_i^\top \mathbf{U} a_i^\top \mathbf{U} + z_k^\top \nabla z_k^\top \nabla a_i^\top \mathbf{U} a_i^\top \mathbf{U} + z_k^\top z_k^\top \nabla^2 a_i^\top \mathbf{U} a_i^\top \mathbf{U} \right] \\
&\quad + 2 \sum_{k=1}^m \sum_{i=1}^n \left[ (z_k^\top \nabla a_i^\top \mathbf{U})^2 + (z^\top \nabla \log \pi)_k \left[ z_k^\top \nabla a_i^\top \mathbf{U} a_i^\top \mathbf{U} \right] \right] \\
&\quad - 2 \sum_{j=1}^m \sum_{l=1}^n \left[ \nabla a_l^\top a_l^\top \nabla z_j^\top \mathbf{U} z_j^\top \mathbf{U} + a_l^\top \nabla a_l^\top \nabla z_j^\top \mathbf{U} z_j^\top \mathbf{U} + a_l^\top a_l^\top \nabla^2 z_j^\top \mathbf{U} z_j^\top \mathbf{U} \right] \\
&\quad - 2 \sum_{j=1}^m \sum_{l=1}^n \left[ (a_l^\top \nabla z_j^\top \mathbf{U})^2 + (a^\top \nabla \log \pi)_l \left[ a_l^\top \nabla z_j^\top \mathbf{U} z_j^\top \mathbf{U} \right] \right],
\end{aligned}$$

$$\mathfrak{R}_{\mathcal{I}_a}(\mathbf{U}, \mathbf{U}) = \langle \mathbf{U}, \gamma \rangle \left[ (\nabla a) \circ (a^\top \mathbf{U}) + \sum_{i=1}^n a_i^\top \nabla a_i^\top \mathbf{U} + \langle aa^\top \nabla \log \pi, \mathbf{U} \rangle \right] - \sum_{\hat{i}=1}^n \sum_{k=1}^{n+m} \gamma_k \nabla_{x_k} a_{\hat{i}}^\top \mathbf{U} (a_{\hat{i}}^\top \mathbf{U}),$$

$$\mathfrak{R}_{\gamma_a}(\mathbf{U}, \mathbf{U}) = \frac{1}{2} \sum_{\hat{k}=1}^{n+m} \gamma_{\hat{k}} \langle \mathbf{U}, \nabla_{\hat{k}} (aa^\top \mathbf{U}) \rangle - \langle \nabla \gamma \mathbf{U}, aa^\top \mathbf{U} \rangle_{\mathbb{R}^{n+m}},$$

$$\mathfrak{R}_{\gamma_z}(\mathbf{U}, \mathbf{U}) = \frac{1}{2} \sum_{\hat{k}=1}^{n+m} \gamma_{\hat{k}} \langle \mathbf{U}, \nabla_{\hat{k}} (zz^\top \mathbf{U}) \rangle - \langle \nabla \gamma \mathbf{U}, zz^\top \mathbf{U} \rangle_{\mathbb{R}^{n+m}}.$$

We define vectors  $\mathbf{D} \in \mathbb{R}^{n^2 \times 1}$ , and  $\mathbf{E} \in \mathbb{R}^{(n \times m) \times 1}$  as below,

$$\mathbf{D}_{ik} = \sum_{\hat{i}, \hat{k}=1}^{n+m} a_{\hat{i}\hat{i}}^\top \partial_{x_i} a_{\hat{k}\hat{k}}^\top \mathbf{U}_{\hat{k}}, \quad \mathbf{E}_{ik} = \sum_{\hat{i}, \hat{k}=1}^{n+m} a_{\hat{i}\hat{i}}^\top \partial_{x_i} z_{\hat{k}\hat{k}}^\top \mathbf{U}_{\hat{k}}. \quad (6)$$

For  $\beta \in \mathbb{R}$ , the vectors  $\Lambda_1 \in \mathbb{R}^{n^2 \times 1}$  and  $\Lambda_2 \in \mathbb{R}^{(n \times m) \times 1}$  are defined as, for  $i, l \in \{1, \dots, n\}$ ,

$$\begin{aligned}
(\Lambda_1)_{il} &= \sum_{k=1}^n \left[ \sum_{i'=1}^{n+m} a_{ii'}^\top \lambda_l^{i'k} - \sum_{k'=1}^{n+m} a_{kk'}^\top \lambda_l^{k'i} \right] a_k^\top \mathbf{U} + \sum_{k=1}^m \left( \sum_{i'=1}^{n+m} a_{ii'}^\top \omega_l^{i'k} - \sum_{k'=1}^{n+m} z_{kk'}^\top \lambda_l^{k'i} \right) z_k^\top \mathbf{U} \\
&\quad - \sum_{k=1}^m \sum_{i'=1}^{n+m} a_{ii'}^\top \omega_l^{i'k} z_k^\top \mathbf{U} - \frac{\beta}{2} \alpha_l (a_i^\top \mathbf{U}) + \frac{\beta}{2} \langle \mathbf{U}, \gamma \rangle \mathbf{1}_{\{i=l\}} + \mathbf{D}_{il},
\end{aligned}$$

and for  $i \in \{1, \dots, n\}$ ,  $l \in \{1, \dots, m\}$ ,

$$\begin{aligned} (\Lambda_2)_{il} &= \sum_{k=1}^n \left[ \sum_{i'=1}^{n+m} a_{ii'}^\top \lambda_{l+n}^{i'k} - \sum_{k'=1}^{n+m} a_{kk'}^\top \lambda_{l+n}^{k'i} \right] a_k^\top \mathbf{U} + \sum_{k=1}^m \left( \sum_{i'=1}^{n+m} a_{ii'}^\top \omega_{l+n}^{i'k} - \sum_{k'=1}^{n+m} z_{kk'}^\top \lambda_{l+n}^{k'i} \right) z_k^\top \mathbf{U} \\ &\quad + \sum_{k=1}^n \sum_{k'=1}^{n+m} z_{lk'}^\top \lambda_i^{k'k} a_k^\top \mathbf{U} + z_l^\top \nabla a_i^\top \mathbf{U} - \sum_{k=1}^m \sum_{i'=1}^{n+m} a_{ii'}^\top \omega_{l+n}^{i'k} z_k^\top \mathbf{U} - a_i^\top \nabla z_l^\top \mathbf{U} - \frac{\beta}{2} \alpha_{l+n} (a_i^\top \mathbf{U}) + \mathbf{E}_{il}. \end{aligned}$$

For each indices  $i, k, \hat{k}$ , assume that there exists smooth functions  $\lambda_l^{i'k}$ ,  $\omega_l^{i'k}$  and  $\alpha_l$  for  $l = 1, \dots, n+m$ ,

$$\nabla_{i'} a_{k\hat{k}}^\top = \sum_{l=1}^n \lambda_l^{i'k} a_{l\hat{k}}^\top + \sum_{l=1}^m \lambda_{l+n}^{i'k} z_{l\hat{k}}^\top, \quad \nabla_{i'} z_{k\hat{k}}^\top = \sum_{l=1}^n \omega_l^{i'k} a_{l\hat{k}}^\top + \sum_{l=1}^m \omega_{l+n}^{i'k} z_{l\hat{k}}^\top,$$

and  $\gamma_{\hat{k}} = \sum_{l=1}^n \alpha_l a_{l\hat{k}}^\top + \sum_{l=1}^m \alpha_{l+n} z_{l\hat{k}}^\top$ . For a vector function  $\gamma \in \mathbb{R}^{n+m}$ , we define  $\nabla \gamma \in \mathbb{R}^{(n+m) \times (n+m)}$  with  $(\nabla \gamma)_{ij} = \nabla_i \gamma_j$ .

We next introduce the following eigenvalue condition.

**Assumption 2** (Hessian matrix condition). *Suppose that there exists a constant  $\lambda > 0$ , a smooth matrix function  $z \in C^\infty(\mathbb{R}^{(n+m) \times m})$  and a scalar  $\beta \in \mathbb{R}$ , such that*

$$\mathfrak{A}(x) \succeq \lambda (a(x)a(x)^\top + z(x)z(x)^\top), \quad \text{for any } x \in \mathbb{R}^{n+m}. \quad (7)$$

**2.3. Entropy dissipation.** We are now ready to present the convergence result for SDE (1). Denote a functional

$$\mathcal{I}_{a,z}(p||\pi) = \int \left( \nabla \log \frac{p(x)}{\pi(x)}, (a(x)a(x)^\top + z(x)z(x)^\top) \nabla \log \frac{p(x)}{\pi(x)} \right) p(x) dx,$$

which measures the difference between functions  $p$  and  $\pi$ . Here we call functional  $\mathcal{I}_{a,z}$  the *weighted relative Fisher information*.

In the next theorem, we establish the convergence of SDEs by the dissipation estimation of functional  $\mathcal{I}_{a,z}$ .

**Theorem 1** (Weighted Fisher information dissipation). *Suppose that Assumption 1, 2 hold. Then*

$$\mathcal{I}_{a,z}(p||\pi) \leq e^{-2\lambda t} \mathcal{I}_{a,z}(p_0||\pi),$$

where  $p = p(t, x)$  is the solution of Fokker-Planck equation (3) and  $p_0$  is a given smooth initial distribution.

Using Theorem 1, we demonstrate several convergence results in other functionals, such as the  $L_1$  distance.

**Corollary 2** (Functionals decay). *Assume condition (7) holds. Then there are following decay results.*

(i) *KL divergence decay:*

$$D_{\text{KL}}(p||\pi) \leq \frac{1}{2\lambda} e^{-2\lambda t} \mathcal{I}_{a,z}(p_0||\pi),$$

where

$$D_{\text{KL}}(p\|\pi) := \int p(t, x) \log \frac{p(t, x)}{\pi(x)} dx,$$

is the Kullback–Leibler (KL) divergence between functions  $p$  and  $\pi$ .

(ii)  $L_1$  distance decay:

$$\int \|p(t, x) - \pi(x)\| dx \leq \sqrt{\frac{1}{\lambda} \mathcal{I}_{a, z}(p_0\|\pi)} e^{-\lambda t}.$$

*Remark 1* (Structure conditions and Hessian matrix conditions). Assumption 1 can be easily verified for a class of degenerate SDEs. Two examples are given below.

- (i) If  $a$  is a constant matrix, then  $\nabla a = 0$ , and Assumption 1 is satisfied;
- (ii) If  $a$  is a matrix function by

$$a = a(x_1, \dots, x_n), \quad z \in \text{span}\{e_{n+1}, \dots, e_{n+m}\},$$

where  $e_i$  is the  $i$ -th Euclidean basis function, then Assumption 1 holds.

We can also verify Assumption 2 in several examples, including Langevin dynamics with variable diffusion matrix (Section 3). It can also be applied to study 3 oscillator chains models [20, 26] (Section 4), where the invariant Gibbs measure has an explicit formula. In fact, these two assumptions have been checked for drift-diffusion processes on Lie group induced sub-Riemannian manifolds, including Heisenberg group, Displacement group and Martinet flat sub-Riemannian structure. See [24, Examples 4.1, 4.2, 4.3].

*Remark 2* (Perturbed gradient flows in sub-Riemannian density manifold). We remark that equation (3) can be viewed as a perturbed gradient flow in probability density space embedded with the sub-Riemannian Wasserstein metric [22, 23]. See related studies for gradient flows in probability density space in [1]. Here an energy function is chosen as the KL divergence while a perturbation is given by the vector field  $\gamma$ . Under this viewpoint, our decomposition allows us to study the dynamical behavior of SDE (1) in three components. In later on proofs, we show that the Gamma two operator studies the reversible component in SDE (1); the generalized Gamma  $z$  operator overcomes the degenerate diffusion matrix function  $a$ ; and the irreversible Gamma operator handles the non-gradient drift vector field  $b$ .

*Remark 3* (Optimal convergence rate). We notice that the convergence bound  $\lambda$  depends on the choice of parameter  $\beta \in \mathbb{R}$  and the selection of vector field  $z(x)$ . In examples, we will demonstrate that the optimal choice of them will obtain better bounds for underdamped Langevin dynamics. See details in section 3.

### 3. EXAMPLE I: UNDERDAMPED LANGEVIN DYNAMICS

In this section, we apply Theorem 1 to prove several explicit convergence rates for variable coefficient irreversible degenerate SDEs.

Consider a underdamped Langevin dynamics with variable diffusion coefficients:

$$\begin{cases} dx_t = v_t dt \\ dv_t = (-r(x_t)v_t - \nabla_x U(x_t))dt + \sqrt{2r(x_t)}dB_t, \end{cases} \quad (8)$$

where  $(x_t, v_t) \in \mathbb{R}^2$  is a two dimensional stochastic process,  $B_t$  is a standard Brownian motion in  $\mathbb{R}$ , and  $r \in \mathbb{R}_+$  is a given positive smooth diffusion coefficient function. Equation (8) often arises in molecular dynamics and Bayesian sampling algorithms; see motivations in [45]. We can check the following facts about SDE (8).

**Proposition 1.** *Assume that  $\int e^{-U(x)} dx < \infty$ . Then the invariant distribution of SDE (8) satisfies*

$$\pi(x, v) = \frac{1}{Z} e^{-H(x, v)}, \quad (9)$$

where  $H(x, v) = \frac{\|v\|^2}{2} + U(x)$  and  $Z = \int_{\mathbb{R}^2} e^{-H(x, v)} dx dv < +\infty$  is a normalization constant.

**Proof** Denote the drift vector field and the diffusion coefficient as

$$b(x, v) = \begin{pmatrix} v \\ -r(x)v - \nabla_x U(x) \end{pmatrix}, \quad a = \begin{pmatrix} 0 \\ \sqrt{r(x)} \end{pmatrix}. \quad (10)$$

Here we only need to prove  $-\nabla \cdot (\pi(x, v)b(x, v)) + \nabla_{vv}^2 (r(x)\pi(x, v)) = 0$ . This equality holds by a direct calculation. Here the above equality can be formulated into

$$-\pi \nabla \cdot \begin{pmatrix} v \\ -\nabla_x U \end{pmatrix} - \begin{pmatrix} \nabla_x \pi \\ \nabla_v \pi \end{pmatrix} \cdot \begin{pmatrix} v \\ -\nabla_x U \end{pmatrix} + r \nabla_v \cdot (\pi v) + r \nabla_{vv}^2 \pi = 0.$$

We observe that  $r \nabla_v \cdot (\pi v) + r \nabla_{vv}^2 \pi = r \nabla_v \cdot (\pi \nabla_v \log \frac{\pi}{e^{-\frac{v^2}{2}}}) = 0$ . And we know that

$$\nabla \cdot \begin{pmatrix} v \\ -\nabla_x U \end{pmatrix} = 0, \quad \nabla_x \pi \cdot v + \nabla_v \pi \cdot (-\nabla_x U) = 0.$$

This finishes the proof.

We are now ready to prove the convergence behavior of SDE (8).

**Notations:** Denote  $X(t) = (x(t), v(t))^\top$ . Denote a vector field  $\gamma \in \mathbb{R}^2$  as

$$\gamma(x, v) = \begin{pmatrix} 0 \\ -rv \end{pmatrix} - \begin{pmatrix} v \\ -rv - \nabla_x U \end{pmatrix} = \begin{pmatrix} -v \\ \nabla_x U \end{pmatrix}.$$

Consider a constant vector  $z = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}$ .

**Proposition 2.** *For any  $\beta \in \mathbb{R}$ , define a matrix function  $\mathfrak{R} \in \mathbb{R}^{2 \times 2}$  as*

$$\mathfrak{R} = \mathfrak{R}_a + \mathfrak{R}_z + \mathfrak{R}_\pi - \mathfrak{M}_\Lambda + \beta \mathfrak{R}_{\mathcal{I}_a} + (1 - \beta) \mathfrak{R}_{\gamma_a} + \mathfrak{R}_{\gamma_z},$$



where

$$\begin{aligned}
\mathfrak{R}_a &= \begin{pmatrix} 0 & 0 \\ 0 & -\frac{\partial^2 \log \pi}{\partial v^2} |a_{12}^\top|^4 \end{pmatrix}, \quad \mathfrak{R}_\pi = \begin{pmatrix} 0 & 0 \\ 0 & C_\pi \end{pmatrix}, \\
\mathfrak{R}_z &= \frac{1}{2} \left[ \begin{pmatrix} 0 \\ -z_1^\top \nabla ((a_{12}^\top)^2 \frac{\partial \log \pi}{\partial v}) \end{pmatrix} z_1^\top + z_1 \begin{pmatrix} 0 & -z_1^\top \nabla ((a_{12}^\top)^2 \frac{\partial \log \pi}{\partial v}) \end{pmatrix} \right], \\
\mathfrak{R}_{\mathcal{L}_a} &= \frac{1}{2} \left[ \gamma (aa^\top \nabla \log \pi)^\top + (aa^\top \nabla \log \pi) \gamma^\top \right] - \begin{pmatrix} 0 & 0 \\ 0 & \gamma_1 \frac{\partial}{\partial x} a_{12}^\top a_{12}^\top \end{pmatrix}, \\
\mathfrak{R}_{\gamma_a} &= \frac{1}{2} \gamma_1 \nabla_1 (aa^\top) - \frac{1}{2} \left[ (\nabla \gamma)^\top aa^\top + aa^\top \nabla \gamma \right], \\
\mathfrak{R}_{\gamma_z} &= -\frac{1}{2} \left[ (\nabla \gamma)^\top z z^\top + z z^\top \nabla \gamma \right], \quad \mathfrak{M}_\Lambda = \frac{1}{(a_{12}^\top)^2} \mathsf{K}^\top (aa^\top + z z^\top)^{-1} \mathsf{K},
\end{aligned}$$

with

$$\begin{aligned}
C_\pi &= 2 \left[ z_1^\top z_1^\top \nabla^2 a_{12}^\top a_{12}^\top + (z_1^\top \nabla a_{12}^\top)^2 + (z_1^\top \nabla \log \pi) [z_1^\top \nabla a_{12}^\top a_{12}^\top] \right], \\
\mathsf{K} &= \begin{pmatrix} 0 & 2z_1^2 \partial_x [a_{12}^\top] a_{12}^\top - \frac{1}{2} \beta \gamma_1 (a_{12}^\top)^2 \\ -z_1^2 \partial_x [a_{12}^\top] a_{12}^\top + \frac{1}{2} \beta \gamma_1 (a_{12}^\top)^2 & z_1 z_2 \partial_x [a_{12}^\top] a_{12}^\top \end{pmatrix}.
\end{aligned}$$

If there exists a constant  $\lambda > 0$ , such that

$$\mathfrak{R} \succeq \lambda (aa^\top + z z^\top),$$

then entropy dissipation results in Theorem 1 and Corollary 2 hold. In other words,

$$\int_{\Omega} \|p(t, x, v) - \pi(x, v)\| dx = O(e^{-\lambda t}).$$

We present several examples to demonstrate the convergence rate of underdamped Langevin dynamics (8).

**Example 1** (Constant diffusion). Consider  $r(x) = 1$ . From Theorem 2,

$$\begin{aligned}
\mathfrak{R} &= \begin{pmatrix} 0 & 0 \\ 0 & -\frac{\partial^2 \log \pi}{\partial v^2} \end{pmatrix} + \frac{1}{2} \left[ \begin{pmatrix} 0 \\ -z_1^\top \nabla \left( \frac{\partial \log \pi}{\partial v} \right) \end{pmatrix} z_1^\top + z \begin{pmatrix} 0 & -z_1^\top \nabla \left( \frac{\partial \log \pi}{\partial v} \right) \end{pmatrix} \right] - \frac{1}{2} \left[ (\nabla \gamma)^\top z z^\top + z z^\top \nabla \gamma \right] \\
&\quad - \frac{1-\beta}{2} \left[ (\nabla \gamma)^\top aa^\top + aa^\top \nabla \gamma \right] + \frac{\beta}{2} \left[ \gamma (aa^\top \nabla \log \pi)^\top + (aa^\top \nabla \log \pi) \gamma^\top \right] \\
&\quad + \frac{\beta^2}{(a_{12}^\top)^2} \begin{pmatrix} 0 & -\frac{\gamma_1}{2} \\ \frac{\gamma_1}{2} & 0 \end{pmatrix}^\top (aa^\top + z z^\top)^{-1} \begin{pmatrix} 0 & -\frac{\gamma_1}{2} \\ \frac{\gamma_1}{2} & 0 \end{pmatrix}.
\end{aligned}$$

We remark that if  $\beta = 0$ , then  $\mathfrak{R}$  leads to the tensor in [6, Formula (7.7)]. Our formulation adds an additional constant  $\beta \in \mathbb{R}$  in the convergence rate. For example, we consider  $U(x) = \frac{x^2}{2}$ . Let  $\lambda$  be the smallest eigenvalue function of  $(aa^\top + z z^\top)^{-1} \mathfrak{R}$  for a given direction  $z = \begin{pmatrix} 1 \\ 0.1 \end{pmatrix} \in \mathbb{R}^2$ . We next plot the smallest eigenvalue function  $\lambda$  on a spatial domain  $[-1, 1]^2$ . As in Figure 2, we observe that chosen a suitable constant  $\beta \in \mathbb{R}$  can improve the convergence rate for a given local spatial region. In details, if  $\beta = 0$ , we observe that  $\lambda(x) = 0.0975$  for all  $x$ . If  $\beta = 0.1$ , we know that  $\lambda(x) = 0.1$  when  $x = [0, 0]$ .




FIGURE 1. Illustration of convergence rates  $\lambda$  with different choices of  $\beta$ .  
Left:  $\beta = 0$ ; Right  $\beta = 0.1$ .

**Example 2** (Variable diffusion). Consider  $U(x) = \frac{x^{2.5}-x}{2.5*1.5}$  with a variable temperature function  $r(x) = (\nabla_x^2 U(x))^{-1}$ . For simplicity of discussion, we study SDE (8) defined on a spatial domain  $[0.5, 1]^2$  with periodic boundary conditions. In this case, we can establish the convergence rate of SDE (8). Similarly, we plot the smallest eigenvalue function  $\lambda = \lambda_{\min}((aa^\top + zz^\top)^{-1}\mathfrak{R})$  on a spatial domain  $[0.5, 1]^2$ . Here we choose  $z = \begin{pmatrix} 1 \\ 0.1 \end{pmatrix}$ . Clearly, we can find a region, in which the solution of Fokker-Planck equation converges to the equilibrium with an positive exponential rate.




FIGURE 2. Illustration of convergence rates  $\lambda$  with different choices of  $\beta$ .  
Left:  $\beta = 0$ ; Right:  $\beta = 0.6$ .

**Proposition 3** (Sufficient conditions). In Example 1, let  $\beta = 0$ ,  $d = 1$ ,  $r(x) = r \in \mathbb{R}_+$ , then

$$\mathfrak{R} = \begin{pmatrix} z_1 z_2 & \frac{1}{2}[z_1 z_2 + z_2^2 - \partial_{xx}^2 U z_1^2 + r] \\ \frac{1}{2}[z_1 z_2 + z_2^2 - \partial_{xx}^2 U z_1^2 + r] & r^2 + z_2^2 - \partial_{xx}^2 U z_1 z_2 \end{pmatrix}_{2 \times 2}.$$

Assume that  $\underline{\lambda} \leq \partial_{xx}^2 U \leq \bar{\lambda}$ , and there exist constants  $z_2 \in (0, \frac{1+\sqrt{1+4r}}{2})$  and  $\delta > 0$ , such that  $\underline{\lambda}, \bar{\lambda}$  satisfies the following conditions,

$$2r\underline{\lambda} - \bar{\lambda}^2 > r^2 - \delta, \quad [2(z_2 - z_2^2)]\underline{\lambda} + (4r^2 - 2r)z_2 + 2z_2^3 - z_2^4 - (2r + 1)z_2^2 > \delta. \quad (11)$$

Then there exists  $\lambda > 0$ , such that

$$\mathfrak{R} \succeq \lambda(aa^\top + zz^\top).$$

**Proof** It is sufficient to prove  $\det(\mathfrak{R}) > 0$  for  $z_1 z_2 > 0$ , which is equivalent to

$$z_1 z_2 (r^2 + z_2^2 - \partial_{xx}^2 U z_1 z_2) - \frac{1}{4} (z_1 z_2 + z_2^2 - \partial_{xx}^2 U z_1^2 + r)^2 > 0.$$

By a direct computation, it is equivalent to the following inequality:

$$-z_1^4 (\partial_{xx}^2 U)^2 + [2(r + z_1 z_2 - z_2^2) z_1^2] \partial_{xx}^2 U + (4r^2 - 2r) z_1 z_2 + 2z_1 z_2^3 - r^2 - (z_1^2 + 2r) z_2^2 - z_2^4 > 0.$$

According to the assumption of  $\partial_{xx}^2 U$ , it is sufficient to prove the following conditions:

$$\begin{cases} z_1 z_2 > 0, & (r + z_1 z_2 - z_2^2) > 0; \\ -z_1^4 \bar{\lambda}^2 + [2(r + z_1 z_2 - z_2^2) z_1^2] \underline{\lambda} + (4r^2 - 2r) z_1 z_2 + 2z_1 z_2^3 - r^2 - (z_1^2 + 2r) z_2^2 - z_2^4 > 0. \end{cases} \quad (12)$$

Let  $z_1 = 1$ , then (11) implies (12). Let  $r + z_2 - z_2^2 > 0$ , for a small constant  $\delta > 0$ , there exists  $z_2 > 0$  such that  $[2(z_2 - z_2^2)]\underline{\lambda} + (4r^2 - 2r)z_2 + 2z_2^3 - z_2^4 - (2r + 1)z_2^2 > \delta$ , which completes the proof.

*Remark 4.* In general, the condition (12) is more general than (11), which holds for a larger class of potential function  $U$ . From simplicity of checking, we provide a simplified condition (11). For example, let  $\underline{\lambda} = \bar{\lambda} = 0.9$ ,  $r = 1$ ,  $\delta = 0.02$ ,  $z_2 = 0.3$ , condition (11) holds.

#### 4. EXAMPLE II: 3-OSCILLATORS WITH NEAREST-NEIGHBOR COUPLINGS

In this section, we apply Theorem 1 to prove  $L_1$  convergence rates for 3-oscillators with nearest-neighbor couplings [26].

Consider

$$\begin{cases} dq_j = p_j dt, & j = 0, 1, 2, \\ dp_0 = -\xi_0 p_0 dt - V_1'(q_0) dt - V_2'(q_0 - q_1) dt + \sqrt{2\xi_0 T} dB_0, \\ dp_1 = -V_1'(q_1) dt - V_2'(q_1 - q_0) dt - V_2'(q_1 - q_2) dt, \\ dp_2 = -\xi_2 p_2 dt - V_1'(q_2) dt - V_2'(q_2 - q_1) dt + \sqrt{2\xi_2 T} dB_2. \end{cases} \quad (13)$$

For simplicity of discussion, we abuse the notation slightly. Denote  $q \in \mathbb{T}^3$  as the state variable and  $p \in \mathbb{T}^3$  as the moment variable. Here  $\mathbb{T}^3$  is a 3 dimensional torus. The dynamics (13) is associated with the Hamiltonian function

$$H(p, q) = \sum_{i=0}^2 \left( \frac{p_i^2}{2} + V_1(q_i) \right) + \sum_{i=1}^2 V_2(q_i - q_{i-1}),$$

where function  $V_2$  is a smooth interaction potential between neighboring oscillators, and function  $V_1$  is a smooth pinning potential. In this paper, we consider  $\xi_0 = \xi_2$ . We assume that there exists a unique invariant measure  $\pi$ , where

$$\pi(p, q) = \frac{1}{Z} e^{-\frac{H(p, q)}{T}},$$

with a normalization constant  $Z = \int_{\mathbb{T}^3 \times \mathbb{T}^3} e^{-\frac{H(p, q)}{T}} dq dp < \infty$ .

*Remark 5.* If  $\xi_0 \neq \xi_2$ , there is no analytical formulation for the invariant density  $\pi$ . See related studies [20, 21, 26]. In this example, we focus on the case when  $\pi$  is explicitly known.

**Notations:** Denote  $X(t) = (q_0(t), q_1(t), q_2(t), p_0(t), p_1(t), p_2(t))^\top$  and  $x = (q_0, q_1, q_2, p_0, p_1, p_2)^\top$ . In this notation, equation (13) can be written as

$$dX_t = b(X_t)dt + \sqrt{2}adB_t,$$

where we denote  $B_t = (B_0(t), B_2(t))^\top$ ,

$$b(X) = \begin{pmatrix} p_0 & p_1 & p_2 & -\xi p_0 - \partial_{q_0} H & -\partial_{q_1} H & -\xi p_2 - \partial_{q_2} H \end{pmatrix}^\top,$$

and

$$a = \begin{pmatrix} 0 & 0 & 0 & \sqrt{\xi T} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sqrt{\xi T} \end{pmatrix}^\top.$$

Denote  $l_3 = \text{Diag}(1, 1, 1)_{3 \times 3}$  as the diagonal identity matrix. We select the matrix  $z$  as

$$z = \begin{pmatrix} z_1 l_3 & 0_{3 \times 1} \\ z_2 l_3 & (\vec{z}_3)_{3 \times 1} \end{pmatrix}_{6 \times 4},$$

where we denote  $\vec{z}_3 = (z_{31} \ z_{32}(p_0, p_2) \ z_{33})^\top$  for constants  $z_{31}, z_{33} \in \mathbb{R}$ , and function  $z_{32}(p_0, p_2) : \mathbb{R}^2 \rightarrow \mathbb{R}$ . We observe that

$$aa^\top = \begin{pmatrix} 0 & 0 \\ 0 & (\xi T) l_3^O \end{pmatrix}_{6 \times 6}, \quad zz^\top = \begin{pmatrix} z_1^2 l_3 & z_1 z_2 l_3 \\ z_1 z_2 l_3 & z_2^2 l_3 + \vec{z}_3 \vec{z}_3^\top \end{pmatrix}_{6 \times 6},$$

where we denote  $l_3^O = \text{Diag}(1, 0, 1)_{3 \times 3}$ , and

$$\begin{aligned} \gamma(x) &= a(x)a(x)^\top \nabla \log \pi(x) - b(x) + \left( \sum_{j=1}^6 \frac{\partial}{\partial x_j} (a(x)a(x)^\top)_{ij} \right)_{1 \leq i \leq 6} \\ &= \begin{pmatrix} -p_0 & -p_1 & -p_2 & \partial_{q_0} H & \partial_{q_1} H & \partial_{q_2} H \end{pmatrix}^\top. \end{aligned}$$

It is easy to verify  $\nabla \cdot (\pi \gamma) = 0$ . We have the following matrix function  $\mathfrak{R}$  for SDE (13).

**Proposition 4.** Define a matrix function  $\mathfrak{R} \in \mathbb{R}^{6 \times 6}$  as

$$\mathfrak{R} = \begin{pmatrix} R_1 & R_2 \\ R_2 & R_3 \end{pmatrix}_{6 \times 6},$$

with

$$\begin{aligned} R_1 &= z_1 z_2 l_3, \quad R_2 = \frac{1}{2} (z_1 z_2 \xi + \xi T) l_3^O + \frac{1}{2} (z_2^2 l_3 + \vec{z}_3 \vec{z}_3^\top - z_1^2 L), \\ R_3 &= ((\xi T)^2 + z_2^2 \xi) l_3^O - z_1 z_2 L + \frac{1}{2} (\vec{z}_3 \vec{z}_3^\top + \vec{z}_3 (\vec{z}_3)^\top) + l_\pi, \end{aligned}$$

where we denote

$$\begin{aligned} \mathbf{l}_3^O &= \text{Diag}(1, 0, 1)_{3 \times 3}, \quad \vec{z}_3 = (z_{31}\xi \quad S_1 \quad z_{33}\xi)^\top, \\ \mathbf{l}_\pi &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & -2z_{32}S_1 - 2\xi T[|\partial_{p_0} z_{45}^\top|^2 + |\partial_{p_2} z_{45}^\top|^2] & 0 \\ 0 & 0 & 0 \end{pmatrix}_{3 \times 3}, \quad L = (\nabla_{q_i q_j} H)_{0 \leq i, j \leq 2}, \\ S_1 &= \xi T \partial_{p_0 p_0} z_{32} + \xi T \partial_{p_2 p_2} z_{32} + (\xi T \partial_{p_0} \log \pi) \partial_{p_0} z_{32} + (\xi T \partial_{p_2} \log \pi) \partial_{p_2} z_{32}. \end{aligned}$$

If there exists a constant  $\lambda > 0$ , such that

$$\mathfrak{R} \succeq \lambda(aa^\top + zz^\top),$$

then entropy dissipation results in Theorem 1 and Corollary 2 hold.

To simplify the computation, we choose the following parameters,

$$\xi = T = 1, \quad z_{31} = z_{33} = 0, \quad z_{32}(p_0, p_2) = N - \varepsilon p_0^2/2, \quad z_1 = 1, \quad (14)$$

with a fixed constant  $N > 0$  and a fixed scaling constant  $\varepsilon > 0$ .

**Proposition 5** (Sufficient conditions). *Assume that  $(\nabla_{q_i q_j} H)_{0 \leq i, j \leq 2}$  satisfies the following conditions, for  $\delta_1, \delta_2 > 0$ ,  $z_2 \in (0, \min\{\frac{1+\sqrt{5}}{2}, N\})$ , and a small constant  $\varepsilon > 0$ ,*

$$\begin{cases} \underline{\lambda} \mathbf{l}_3 \preceq (\nabla_{q_i q_j} H)_{0 \leq i, j \leq 2} \preceq \bar{\lambda} \mathbf{l}_3; \\ 2\underline{\lambda} - \bar{\lambda}^2 \geq 1 - \delta_1, \quad -(z_2^2 + N^2)^2 + 2(N^2 - z_2^2)\underline{\lambda} - \bar{\lambda}^2 > 0; \\ 2z_2 + 2z_2^3 - z_2^4 - 3z_2^2 + 2(z_2 - z_2^2)\underline{\lambda} > \delta_1 \end{cases} \quad (15)$$

Here we denote  $0 \leq \underline{\lambda} \leq \bar{\lambda}$  as the lower and upper bound eigenvalues of the matrix  $\nabla_{q q} H$ . Then there exists a constant  $\lambda > 0$ , such that

$$\mathfrak{R} \succeq \lambda(aa^\top + zz^\top).$$

*Remark 6.* The condition of  $\nabla_{q q} H$  given in the above proposition implies the condition for  $\nabla_{q q} V_1$  and  $\nabla_{q q} V_2$ .

**Proof** It is sufficient to prove that  $\mathfrak{R}$  defined in Proposition 4 is positive definite. Applying the Schur complement for symmetric matrix function [47][Appendix A.5], the following conditions are equivalent,

- (1)  $\mathfrak{R} \succeq 0$  ( $\mathfrak{R}$  is positive semidefinite).
- (2)  $\mathbf{R}_1 \succeq 0$ ,  $(\mathbf{l}_3 - \mathbf{R}_1 \mathbf{R}_1^{-1}) \mathbf{R}_2 = 0$ ,  $\mathbf{R}_3 - \mathbf{R}_2 \mathbf{R}_1^{-1} \mathbf{R}_2 \succeq 0$ .

Applying the parameters in (14), let  $z_2 > 0$ , we just need to show  $R_3 - R_2 R_1^{-1} R_2 \succeq 0$ . In other words,

$$\begin{aligned}
& R_3 - R_2 R_1^{-1} R_2 \\
&= \begin{pmatrix} 1 + z_2^2 & 0 & 0 \\ 0 & \varepsilon(1 - p_0^2)(N - \varepsilon p_0^2/2) - 2\varepsilon^2 p_0^2 & 0 \\ 0 & 0 & 1 + z_2^2 \end{pmatrix} - z_2 L \\
&\quad - \frac{1}{z_2} \left( \begin{pmatrix} \frac{1}{2}z_2^2 + \frac{1}{2}z_2 + \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2}z_2^2 + \frac{1}{2}(N - \varepsilon p_0^2/2)^2 & 0 \\ 0 & 0 & \frac{1}{2}z_2^2 + \frac{1}{2}z_2 + \frac{1}{2} \end{pmatrix} - \frac{1}{2}L \right)^2 \\
&:= \text{Diag}(C_1, C_2, C_3) - z_2 \mathbf{O} \cdot \text{Diag}(\lambda_1, \lambda_2, \lambda_3)_{3 \times 3} \cdot \mathbf{O}^\top \\
&\quad - \frac{1}{4z_2} \left( \text{Diag}(C_4, C_5, C_6) - \mathbf{O} \cdot \text{Diag}(\lambda_1, \lambda_2, \lambda_3)_{3 \times 3} \cdot \mathbf{O}^\top \right)^2,
\end{aligned}$$

where

$$\begin{aligned}
C_1 &= C_3 = 1 + z_2^2, & C_2 &= \varepsilon(1 - p_0^2)(N - \varepsilon p_0^2/2) - 2\varepsilon^2 p_0^2, \\
C_4 &= C_6 = 1 + z_2^2 + z_2, & C_5 &= z_2^2 + (N - \varepsilon p_0^2/2)^2,
\end{aligned}$$

and we denote the eigenvalue decomposition of matrix  $L$  as

$$L = \mathbf{O} \cdot \text{Diag}(\lambda_1, \lambda_2, \lambda_3)_{3 \times 3} \cdot \mathbf{O}^\top.$$

Here we denote  $\mathbf{O} \in \mathbb{R}^{3 \times 3}$  as the orthogonal matrix with  $\mathbf{O}^{-1} = \mathbf{O}^\top$  and  $\lambda_1 \leq \lambda_2 \leq \lambda_3$  as the eigenvalues of matrix  $L$ . Thus

$$\begin{aligned}
& R_3 - R_2 R_1^{-1} R_2 \\
&= \mathbf{O} \cdot \text{Diag} \left( \left( C_i - z_2 \lambda_i - \frac{1}{4z_2} (C_{i+3}^2 - 2C_{i+3} \lambda_i + \lambda_i^2) \right)_{1 \leq i \leq 3} \right)_{3 \times 3} \cdot \mathbf{O}^\top \\
&= \mathbf{O} \cdot \text{Diag} \left( \left( C_i - \frac{C_{i+3}^2}{4z_2} + \left( \frac{C_{i+3}}{2z_2} - z_2 \right) \lambda_i - \frac{\lambda_i^2}{4z_2} \right)_{1 \leq i \leq 3} \right)_{3 \times 3} \cdot \mathbf{O}^\top
\end{aligned}$$

It is sufficient to find  $z_2$ ,  $N$ , and  $\varepsilon$  such that the following inequalities hold:

$$\left( C_i - \frac{C_{i+3}^2}{4z_2} + \left( \frac{C_{i+3}}{2z_2} - z_2 \right) \lambda_i - \frac{\lambda_i^2}{4z_2} \right) > 0, \quad 1 \leq i \leq 3.$$

Since  $z_2 > 0$ , it is equivalent to prove

$$\begin{cases} z_2(1 + z_2^2) - \frac{(1+z_2+z_2^2)^2}{4} + \left( \frac{1+z_2+z_2^2}{2} - z_2 \right) \lambda_i - \frac{\lambda_i^2}{4} > 0, \\ z_2 \varepsilon(1 - p_0^2)(N - \varepsilon p_0^2/2) - 2\varepsilon^2 p_0^2 - \frac{(z_2^2 + (N - \varepsilon p_0^2/2)^2)^2}{4} + \left( \frac{z_2^2 + (N - \varepsilon p_0^2/2)^2}{2} - z_2 \right) \lambda_i - \frac{\lambda_i^2}{4} > 0. \end{cases} \quad (16)$$

The first inequality in (16) holds under the following conditions. Suppose  $1 + z_2 - z_2^2 > 0$ ,

$$2\lambda - \bar{\lambda}^2 \geq 1 - \delta_1, \quad 2z_2 + 2z_2^3 - z_2^4 - 3z_2^2 + 2(z_2 - z_2^2)\lambda > \delta_1.$$

For  $p_0 \in \mathbb{T}^1$ , we select  $\varepsilon$  to be very small such that

$$-\frac{(z_2^2 + N^2)^2}{4} + (N^2 - z_2^2) \frac{\lambda}{2} - \frac{\bar{\lambda}^2}{4} > 0.$$

The proof is finished. ■

*Remark 7.* Consider  $\underline{\lambda} = 0.6$ ,  $\bar{\lambda} = 0.65$ ,  $N = 1$ ,  $z_2 = 0.2$ ,  $\delta_1 = 0.225$ ,  $\varepsilon = 10^{-7}$ . All conditions in (15) are satisfied.

## 5. PROOF

In this section, we present the main proof of this paper.

**5.1. Outline of main result.** Before showing all detailed proofs behind Theorem 1, we outline the main ideas by the following three steps. We postpone all derivations in subsection 5.3.

Step 1: We first compute the dissipation of the weighted Fisher information functional.

$$\frac{d}{dt} \mathcal{I}_{a,z}(p|\pi) = -2 \int \Gamma_{2,a,z,\gamma}(f, f) p dx,$$

where  $f = \log \frac{p}{\pi}$  and

$$\Gamma_{2,a,z,\gamma}(f, f) = \tilde{\Gamma}_2(f, f) + \tilde{\Gamma}_2^{z,\pi}(f, f) + \Gamma_{\mathcal{I}_{a,z}}(f, f).$$

The definition of the above operators will be introduced shortly in Definition 2.

Step 2: We next decompose the weak form of information Gamma calculus in details. This is to derive the information Bochner's formula (Theorem 3):

$$\int \Gamma_{2,a,z,\gamma}(f, f) p dx = \int \left[ \|\mathfrak{Hess}_\beta f\|_{\mathbb{F}}^2 + \mathfrak{R}(\nabla f, \nabla f) \right] p dx.$$

Step 3: From the information Bochner's formula, we establish the convergence result. In other words, if  $\mathfrak{R} \succeq \lambda(aa^\top + zz^\top)$ , then

$$\frac{d}{dt} \mathcal{I}_{a,z}(p|\pi) \leq -2\lambda \mathcal{I}_{a,z}(p|\pi).$$

From the Gronwall's inequality, we can prove the weighted Fisher information decay in Theorem 1. This decay result establishes other functional decays in Corollary 2.

We comment that Step 1 is based on a global in space computation, which comes from the second-order calculus of entropy functional in sub-Riemannian density manifold. Here we carefully handle the non-communicative operators based on  $\Gamma_1$ ,  $\Gamma_1^z$  and vector field  $\gamma$ . Step 2 is a local in space calculation, which can be viewed as a generalization of Bochner's formula. Step 3 follows directly from the modified Lyapunov method.

**5.2. Information Gamma calculus.** We next introduce the needed tensors for the main proof. We call these tensors as information Gamma calculus. Denote the following operators of SDE (1) below. For any  $f \in C^\infty(\mathbb{R}^{n+m})$ , the generator of SDE (1) satisfies

$$Lf = \tilde{L}f - \langle \gamma, \nabla f \rangle,$$

where

$$\tilde{L}f = \nabla \cdot (aa^\top \nabla f) + \langle aa^\top \nabla \log \pi, \nabla f \rangle, \quad (17)$$

is the reversible component of Kolmogorov backward operator. For a given matrix function  $a \in \mathbb{R}^{(n+m) \times n}$ , we construct a matrix function  $z \in \mathbb{R}^{(n+m) \times m}$ , which is to handle the degenerate component of SDE (1). We also denote a  $z$ -direction generator as

$$\tilde{L}_z f = \nabla \cdot (zz^\top \nabla f) + \langle zz^\top \nabla \log \pi, \nabla f \rangle.$$

Define Gamma one bilinear forms  $\Gamma_1, \Gamma_1^z: C^\infty(\mathbb{R}^{n+m}) \times C^\infty(\mathbb{R}^{n+m}) \rightarrow C^\infty(\mathbb{R}^{n+m})$  as

$$\Gamma_1(f, f) = \langle a^\top \nabla f, a^\top \nabla f \rangle_{\mathbb{R}^n}, \quad \Gamma_1^z(f, f) = \langle z^\top \nabla f, z^\top \nabla f \rangle_{\mathbb{R}^m}. \quad (18)$$

We are ready to define the *information Gamma operators* for SDE (1). In fact, it is a derivative calculation based on the reversible Kolmogorov backward operator  $\tilde{L}$ , the non-reversible vector field  $\gamma$  and the invariant distribution  $\pi$ .

**Definition 2** (Information Gamma operators). *Define the following three bi-linear forms:*

$$\tilde{\Gamma}_2, \tilde{\Gamma}_2^{z, \pi}, \Gamma_{\mathcal{I}_{a, z}}: C^\infty(\mathbb{R}^{n+m}) \times C^\infty(\mathbb{R}^{n+m}) \rightarrow C^\infty(\mathbb{R}^{n+m}).$$

(i) *Gamma two operator:*

$$\tilde{\Gamma}_2(f, f) = \frac{1}{2} \tilde{L} \Gamma_1(f, f) - \Gamma_1(\tilde{L} f, f).$$

(ii) *Generalized Gamma  $z$  operator:*

$$\begin{aligned} \tilde{\Gamma}_2^{z, \pi}(f, f) &= \frac{1}{2} \tilde{L} \Gamma_1^z(f, f) - \Gamma_1^z(\tilde{L} f, f) \\ &\quad + \mathbf{div}_z^\pi \left( \Gamma_{1, \nabla(aa^\top)}(f, f) \right) - \mathbf{div}_a^\pi \left( \Gamma_{1, \nabla(zz^\top)}(f, f) \right). \end{aligned}$$

Here  $\mathbf{div}_a^\pi$ ,  $\mathbf{div}_z^\pi$  are divergence operators defined by

$$\mathbf{div}_a^\pi(F) = \frac{1}{\pi} \nabla \cdot (\pi a a^\top F), \quad \mathbf{div}_z^\pi(F) = \frac{1}{\pi} \nabla \cdot (\pi z z^\top F),$$

for any smooth vector field  $F \in \mathbb{R}^{n+m}$ , and  $\Gamma_{1, \nabla(aa^\top)}$ ,  $\Gamma_{1, \nabla(zz^\top)}$  are vector Gamma one bilinear forms defined by

$$\begin{aligned} \Gamma_{1, \nabla(aa^\top)}(f, f) &= \langle \nabla f, \nabla(aa^\top) \nabla f \rangle = \left\langle \nabla f, \frac{\partial}{\partial x_k} (aa^\top) \nabla f \right\rangle_{k=1}^{n+m}, \\ \Gamma_{1, \nabla(zz^\top)}(f, f) &= \langle \nabla f, \nabla(zz^\top) \nabla f \rangle = \left\langle \nabla f, \frac{\partial}{\partial x_k} (zz^\top) \nabla f \right\rangle_{k=1}^{n+m}. \end{aligned}$$

(iii) *Irreversible Gamma operator:*

$$\Gamma_{\mathcal{I}_{a, z}}(f, f) = (\tilde{L} f + \tilde{L}_z f) \langle \nabla f, \gamma \rangle - \frac{1}{2} \langle \nabla (\Gamma_1(f, f) + \Gamma_1^z(f, f)), \gamma \rangle.$$

Under Assumption 1, we establish the following *information Bochner's formula*.

**Theorem 3** (Information Bochner's formula). *If Assumption 1 is satisfied, then the following decomposition holds. For any  $f = \log \frac{\rho}{\pi} \in C^\infty(\mathbb{R}^{n+m}, \mathbb{R})$  and any  $\beta \in \mathbb{R}$ ,*

$$\int \left[ \tilde{\Gamma}_2(f, f) + \tilde{\Gamma}_2^{z, \pi}(f, f) + \Gamma_{\mathcal{I}_{a, z}}(f, f) \right] p dx = \int \left[ \|\mathfrak{Hess}_\beta f\|_{\mathbb{F}}^2 + \mathfrak{R}(\nabla f, \nabla f) \right] p dx.$$

Here we denote

$$\|\mathfrak{Hess}_\beta f\|_{\mathbb{F}}^2 = [\mathbf{QX} + \Lambda_1]^\top [\mathbf{QX} + \Lambda_1] + [\mathbf{PX} + \Lambda_2]^\top [\mathbf{PX} + \Lambda_2],$$



where  $\mathfrak{R}$ ,  $\Lambda_1$ ,  $\Lambda_2$  are defined in 1. And we define matrices  $\mathbf{Q}$  and  $\mathbf{P}$  by

$$\mathbf{Q} = a^\top \otimes a^\top \in \mathbb{R}^{n^2 \times (n+m)^2}, \quad \mathbf{P} = a^\top \otimes z^\top \in \mathbb{R}^{(nm) \times (n+m)^2}, \quad (19)$$

with  $Q_{ik\hat{i}\hat{k}} = a_{i\hat{i}}^\top a_{k\hat{k}}^\top$  and  $P_{ik\hat{i}\hat{k}} = a_{i\hat{i}}^\top z_{k\hat{k}}^\top$ . More precisely, for each row (resp. column) of  $\mathbf{Q}$ , the row (resp. column) indices of  $Q_{ik\hat{i}\hat{k}}$  follows  $\sum_{i=1}^n \sum_{k=1}^n$  (resp.  $\sum_{\hat{i}=1}^{n+1} \sum_{\hat{k}=1}^{n+m}$ ). For any smooth function  $f : \mathbb{R}^{n+m} \rightarrow \mathbb{R}$ , we define  $\mathbf{X} \in \mathbb{R}^{(n+m)^2 \times 1}$  by the vectorization of the Hessian matrix of function  $f$  with

$$X_{\hat{i}\hat{k}} = \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}}, \quad \text{for } \hat{i}, \hat{k} = 1, \dots, n+m.$$

*Remark 8* (Connections with Bakry–Émery conditions). We remark that our result extends the classical diffusion hypercontractivity, namely Bakry–Émery Gamma calculus [7].

- (i) If  $\gamma = 0$  and  $m = 0$ , then information Gamma calculus recovers the Bakry–Émery Gamma 2 calculus. It is designed for reversible diffusions on a Riemannian manifold;
- (ii) If  $\gamma = 0$  and  $m \neq 0$ , and there exists a communicative restriction between  $a$  and  $z$ , i.e.  $\Gamma_1(f, \Gamma_1^z(f, f)) = \Gamma_1^z(f, \Gamma_1(f, f))$ , then information Gamma calculus leads to Baudoin–Garofalo Gamma  $z$  calculus [11]. This communicative restriction has been removed by [22], where the generalized Gamma  $z$  calculus term  $\mathbf{div}_z^\pi \left( \Gamma_{1, \nabla(aa^\top)}(f, f) \right) - \mathbf{div}_a^\pi \left( \Gamma_{1, \nabla(zz^\top)}(f, f) \right)$  overcomes the communicative restriction in proving  $L_1$  norm decay;
- (iii) If  $\beta = 0$  and  $m = 0$ , then the tensor of information Gamma calculus leads to the Arnold–Carlen–Ju tensor [4, 5];
- (iv) If  $a, z$  are constants and  $\beta = 0$ , then the derived tensor leads to the ones given in Arnold–Erb tensor [6];
- (v) If  $\beta = 1$ ,  $m = 0$  and  $a = \mathbb{I}$ , then the information Gamma calculus formulates the ones proposed in [23].

*Remark 9* (Optimal Bochner’s formula). We remark that the proposed information Bochner’s formula can be asymmetric, which could be formulated into a more symmetric way. In other words, denote a quadratic matrix function by

$$\mathcal{F}(Y) = [QY + \Lambda_1]^\top [QY + \Lambda_1] + [PY + \Lambda_2]^\top [PY + \Lambda_2],$$

where  $Y \in \mathbb{R}^{(n+m)^2 \times 1}$  is a vectorization of a symmetric matrix. We next consider a symmetric matrix optimization problem by

$$Y^* = \arg \min_Y \mathcal{F}(Y).$$

Then our Bochner’s formula forms

$$\begin{aligned} & \|\mathfrak{Hess}_\beta f\|_{\mathbb{F}}^2 + \mathfrak{R}(\nabla f, \nabla f) \\ &= \mathcal{F}(X) - \mathcal{F}(Y^*) + \mathcal{F}(Y^*) + \mathfrak{R}(\nabla f, \nabla f). \end{aligned}$$

Here we may call the term  $\mathcal{F}(X) - \mathcal{F}(Y^*)$  the “squared Hessian formula” while name the term  $\mathcal{F}(Y^*) + \mathfrak{R}(\nabla f, \nabla f) + \mathfrak{R}_{\mathcal{I}}(\nabla f, \nabla f)$  the “Ricci curvature type tensor”. This optimal choice of Bochner’s formula are left for future works. See Remark 11 in the Appendix for an example of the optimal Bochner’s formula.

### 5.3. Proof of Step 1.

**Proposition 6.** *The following equality holds.*

$$\frac{d}{dt}\mathcal{I}_{a,z}(p|\pi) = -2 \int \Gamma_{2,a,z,\gamma}(\log \frac{p}{\pi}, \log \frac{p}{\pi})pdx,$$

where  $p$  is the solution of Fokker-Planck equation (2).

**Proof** The proof combines the calculations in [22, 23]. We first present the outline of the proof here. Denote

$$\mathcal{I}_{a,z}(p|\pi) = \mathcal{I}_a(p|\pi) + \mathcal{I}_z(p|\pi),$$

where

$$\mathcal{I}_a(p|\pi) = \int (\nabla \log \frac{p}{\pi}, aa^\top \nabla \log \frac{p}{\pi})pdx, \quad \mathcal{I}_z(p|\pi) = \int (\nabla \log \frac{p}{\pi}, zz^\top \nabla \log \frac{p}{\pi})pdx.$$

We observe the following facts.

**Claim 0:**

$$\frac{d}{dt}\mathcal{I}_a(p|\pi) = -2 \int (\tilde{\Gamma}_2 + \Gamma_{\mathcal{I}_a})(\log \frac{p}{\pi}, \log \frac{p}{\pi})pdx,$$

and

$$\frac{d}{dt}\mathcal{I}_z(p|\pi) = -2 \int (\tilde{\Gamma}_2^{z,\pi} + \Gamma_{\mathcal{I}_z})(\log \frac{p}{\pi}, \log \frac{p}{\pi})pdx.$$

From the Claim 0, we obtain

$$\frac{d}{dt}\mathcal{I}_{a,z}(p|\pi) = \frac{d}{dt}\mathcal{I}_a(p|\pi) + \frac{d}{dt}\mathcal{I}_z(p|\pi) = -2 \int \Gamma_{2,a,z,\gamma}(\log \frac{p}{\pi}, \log \frac{p}{\pi})pdx,$$

which finishes the proof. Next, we prove Claim 0 in details. Denote that  $p$  solves the Fokker-Planck equation (2) and let  $\tilde{L}^*$  be the forward Kolmogorov operator.

Firstly, the dissipation of  $\mathcal{I}_a(p||\pi)$  follows

$$\begin{aligned}
& \frac{d}{dt} \mathcal{I}_a(p||\pi) \\
&= \frac{d}{dt} \int \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) p dx \\
&= \int \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p + 2\Gamma_1(\partial_t \log \frac{p}{\pi}, \log \frac{p}{\pi}) p dx \\
&= \int \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p + 2\Gamma_1(\frac{\partial_t p}{p}, \log \frac{p}{\pi}) p dx \\
&= \int \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p - 2 \frac{\nabla \cdot (p a a^\top \nabla \log \frac{p}{\pi})}{p} \partial_t p dx \\
&= \int \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p - 2 \left( \langle \nabla \log p, a a^\top \nabla \log \frac{p}{\pi} \rangle + \nabla \cdot (a a^\top \nabla \log \frac{p}{\pi}) \right) \partial_t p dx \\
&= \int \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p - 2 \left( \langle \nabla \log \frac{p}{\pi}, a a^\top \nabla \log \frac{p}{\pi} \rangle + \tilde{L} \log \frac{p}{\pi} \right) \partial_t p dx \\
&= -2 \int \left\{ \frac{1}{2} \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p + \tilde{L} \log \frac{p}{\pi} \partial_t p \right\} dx \\
&= -2 \int \left\{ \frac{1}{2} \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \nabla \cdot (p \gamma) + \tilde{L} \log \frac{p}{\pi} \nabla \cdot (p \gamma) \right. \\
&\quad \left. + \frac{1}{2} \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \tilde{L}^* p + \tilde{L} \log \frac{p}{\pi} \tilde{L}^* p \right\} dx \\
&= -2 \int \left\{ -\frac{1}{2} \langle \nabla \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}), \gamma \rangle + \tilde{L} \log \frac{p}{\pi} \langle \nabla \log \frac{p}{\pi}, \gamma \rangle \right. \\
&\quad \left. + \frac{1}{2} \tilde{L} \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) - \Gamma_1(\tilde{L} \log \frac{p}{\pi}, \log \frac{p}{\pi}) \right\} p dx \\
&= -2 \int \left\{ \Gamma_{\mathcal{I}_a}(\log \frac{p}{\pi}, \log \frac{p}{\pi}) + \tilde{\Gamma}_2(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \right\} p dx.
\end{aligned}$$

In above derivations, we apply the following fact:

$$\nabla \cdot (p \gamma) = p \left( \langle \nabla \log p, \gamma \rangle + \nabla \cdot \gamma \right) = p \langle \nabla \log \frac{p}{\pi}, \gamma \rangle,$$

where we use the fact of  $\pi$  by  $\nabla \cdot (\pi \gamma) = \pi \left( \langle \nabla \log \pi, \gamma \rangle + \nabla \cdot \gamma \right) = 0$ .

Secondly, we compute the dissipation of  $\mathcal{I}_z(p||\pi)$ . Denote

$$\tilde{L}_z = (\nabla \log \pi, z z^\top \nabla) + \nabla \cdot (z z^\top \nabla).$$

Hence

$$\begin{aligned}
& \frac{d}{dt} \mathcal{I}_z(p|\pi) \\
&= \frac{d}{dt} \int \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) p dx \\
&= \int \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p + 2\Gamma_1^z(\partial_t \log \frac{p}{\pi}, \log \frac{p}{\pi}) p dx \\
&= \int \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p + 2\Gamma_1^z(\frac{\partial_t p}{p}, \log \frac{p}{\pi}) p dx \\
&= \int \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p - 2 \frac{\nabla \cdot (p z z^\top \nabla \log \frac{p}{\pi})}{p} \partial_t p dx \\
&= \int \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p - 2 \left( \langle \nabla \log p, z z^\top \nabla \log \frac{p}{\pi} \rangle + \nabla \cdot (z z^\top \nabla \log \frac{p}{\pi}) \right) \partial_t p dx \\
&= \int \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p - 2 \left( \langle \nabla \log \frac{p}{\pi}, z z^\top \nabla \log \frac{p}{\pi} \rangle + \tilde{L}_z \log \frac{p}{\pi} \right) \partial_t p dx \\
&= -2 \int \left\{ \frac{1}{2} \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \partial_t p + \tilde{L}_z \log \frac{p}{\pi} \partial_t p \right\} dx \\
&= -2 \int \left\{ \frac{1}{2} \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \nabla \cdot (p \gamma) + \tilde{L}_z \log \frac{p}{\pi} \nabla \cdot (p \gamma) \right. \\
&\quad \left. + \frac{1}{2} \Gamma_1^z(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \tilde{L}^* p + \tilde{L}_z \log \frac{p}{\pi} \tilde{L}^* p \right\} dx \\
&= -2 \int \left\{ \Gamma_{\mathcal{I}_z}(\log \frac{p}{\pi}, \log \frac{p}{\pi}) + \frac{1}{2} \tilde{L}_z \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) - \Gamma_1(\tilde{L}_z \log \frac{p}{\pi}, \log \frac{p}{\pi}) \right\} p dx.
\end{aligned}$$

We notice that the above formula can not be used to derive the Bochner's formula. Since the related third derivative tensors can not be eliminated. To handle this issue, we present the following equality by

$$\int \left\{ \frac{1}{2} \tilde{L}_z \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) - \Gamma_1(\tilde{L}_z \log \frac{p}{\pi}, \log \frac{p}{\pi}) \right\} p dx = \int \tilde{\Gamma}_2^{z, \pi}(\log \frac{p}{\pi}, \log \frac{p}{\pi}) p dx. \quad (20)$$

The identity (20) has been derived in [22, Proposition 5.11]. For completeness of this paper, we also present its derivation here. Notice that

$$\begin{aligned}
& \frac{1}{2} \tilde{L}_z \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) - \Gamma_1(\tilde{L}_z \log \frac{p}{\pi}, \log \frac{p}{\pi}) \\
&= \frac{1}{2} \tilde{L}_z \Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}) - \Gamma_1^z(\tilde{L} \log \frac{p}{\pi}, \log \frac{p}{\pi}) + \Gamma_1^z(\tilde{L} \log \frac{p}{\pi}, \log \frac{p}{\pi}) - \Gamma_1(\tilde{L}_z \log \frac{p}{\pi}, \log \frac{p}{\pi}).
\end{aligned}$$

We next claim that

$$\begin{aligned}
& \int \left\{ \Gamma_1^z(\tilde{L} \log \frac{p}{\pi}, \log \frac{p}{\pi}) - \Gamma_1(\tilde{L}_z \log \frac{p}{\pi}, \log \frac{p}{\pi}) \right\} p dx \\
&= \int \left\{ \mathbf{div}^\pi \left( \Gamma_{1, \nabla(aa^\top)}(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \right) - \mathbf{div}_a^\pi \left( \Gamma_{1, \nabla(zz^\top)}(\log \frac{p}{\pi}, \log \frac{p}{\pi}) \right) \right\} p dx.
\end{aligned}$$

We derive it as follows. Firstly,

$$\begin{aligned}
& \int \Gamma_1^z(\tilde{L} \log \frac{p}{\pi}, \log \frac{p}{\pi}) p dx \\
&= \int \Gamma_1^z \left( \langle \nabla \log \pi, aa^\top \nabla \log \frac{p}{\pi} \rangle + \nabla \cdot (aa^\top \nabla \log \frac{p}{\pi}), \log \frac{p}{\pi} \right) p dx \\
&= \int \Gamma_1^z \left( \Gamma_1(\log \pi, \log \frac{p}{\pi}), \log \frac{p}{\pi} \right) p + \Gamma_1^z \left( \nabla \cdot (aa^\top \nabla \log \frac{p}{\pi}), \log \frac{p}{\pi} \right) p dx \\
&= \int \Gamma_1^z \left( \Gamma_1(\log \pi, \log \frac{p}{\pi}), \log \frac{p}{\pi} \right) p dx - \int \nabla \cdot (aa^\top \nabla \log \frac{p}{\pi}) \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) dx.
\end{aligned}$$

We notice that the second term in the above formula can be written below.

$$\begin{aligned}
& - \int \nabla \cdot (aa^\top \nabla \log \frac{p}{\pi}) \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) dx \\
&= - \int \nabla \cdot \left( \frac{\pi}{p} aa^\top \nabla \frac{p}{\pi} \right) \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) dx \\
&= - \int \left( \pi \langle \nabla \frac{1}{p}, aa^\top \nabla \frac{p}{\pi} \rangle + \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \right) \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) dx \\
&= - \int \pi \langle \nabla \frac{1}{p}, aa^\top \nabla \frac{p}{\pi} \rangle \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) dx - \int \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \nabla \cdot (pzz^\top \nabla \frac{p}{\pi}) dx \\
&= \int \frac{\pi}{p^2} \langle \nabla p, aa^\top \nabla \frac{p}{\pi} \rangle \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) - \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \nabla \cdot (pzz^\top \nabla \frac{p}{\pi}) dx \\
&= \int \langle \nabla \log p, aa^\top \nabla \log \frac{p}{\pi} \rangle \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) - \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \nabla \cdot (pzz^\top \nabla \frac{p}{\pi}) dx \\
&= \int \left\{ \langle \nabla \log \frac{p}{\pi}, aa^\top \nabla \log \frac{p}{\pi} \rangle \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) + \langle \nabla \log \pi, aa^\top \nabla \log \frac{p}{\pi} \rangle \nabla \cdot (pzz^\top \nabla \log \frac{p}{\pi}) \right. \\
&\quad \left. - \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \nabla \cdot (pzz^\top \nabla \frac{p}{\pi}) \right\} dx \\
&= \int \left\{ - \Gamma_1^z(\Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}), \log \frac{p}{\pi}) - \Gamma_1^z(\Gamma_1(\log \pi, \log \frac{p}{\pi}), \log \frac{p}{\pi}) p \right. \\
&\quad \left. - \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \nabla \cdot (pzz^\top \nabla \frac{p}{\pi}) \right\} dx \\
&= \int \left\{ - \Gamma_1^z(\Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}), \log \frac{p}{\pi}) p - \Gamma_1^z(\Gamma_1(\log \pi, \log \frac{p}{\pi}), \log \frac{p}{\pi}) p \right. \\
&\quad \left. - \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \nabla \cdot (pzz^\top \nabla \frac{p}{\pi}) \right\} dx.
\end{aligned}$$

We observe that

$$\begin{aligned}
& - \int \Gamma_1^z(\Gamma_1(\log \frac{p}{\pi}, \log \frac{p}{\pi}), \log \frac{p}{\pi}) p dx \\
= & - \int \nabla^2 \log \frac{p}{\pi} (aa^\top \nabla \log \frac{p}{\pi}, zz^\top \nabla \log \frac{p}{\pi}) p dx \\
& - \int \left( \Gamma_{1, \nabla(aa^\top)}(\nabla \log \frac{p}{\pi}, \nabla \log \frac{p}{\pi}) zz^\top \nabla \log \frac{p}{\pi} \right) p dx \\
= & - \int \nabla^2 \log \frac{p}{\pi} (aa^\top \nabla \log \frac{p}{\pi}, zz^\top \nabla \log \frac{p}{\pi}) p dx \\
& - \int \left( \Gamma_{1, \nabla(aa^\top)}(\nabla \log \frac{p}{\pi}, \nabla \log \frac{p}{\pi}) zz^\top \nabla \frac{p}{\pi} \right) \pi dx \\
= & - \int \nabla^2 \log \frac{p}{\pi} (aa^\top \nabla \log \frac{p}{\pi}, zz^\top \nabla \log \frac{p}{\pi}) p dx \\
& + \int \frac{1}{\pi} \nabla \cdot \left( zz^\top \Gamma_{1, \nabla(aa^\top)}(\nabla \log \frac{p}{\pi}, \nabla \log \frac{p}{\pi}) \right) dx.
\end{aligned}$$

Combining the above three formulas, we have

$$\begin{aligned}
& \int \Gamma_1^z(\tilde{L} \log \frac{p}{\pi}, \log \frac{p}{\pi}) p dx \\
= & \int \frac{1}{\pi} \nabla \cdot \left( zz^\top \Gamma_{1, \nabla(aa^\top)}(\nabla \log \frac{p}{\pi}, \nabla \log \frac{p}{\pi}) \right) dx \\
& - \int \nabla^2 \log \frac{p}{\pi} (aa^\top \nabla \log \frac{p}{\pi}, zz^\top \nabla \log \frac{p}{\pi}) p dx - \int \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \nabla \cdot (\pi zz^\top \nabla \frac{p}{\pi}) dx.
\end{aligned}$$

Secondly, by switching the order of  $a$  and  $z$ , we have

$$\begin{aligned}
& \int \Gamma_1(\tilde{L}_z \log \frac{p}{\pi}, \log \frac{p}{\pi}) p dx \\
= & \int \frac{1}{\pi} \nabla \cdot \left( aa^\top \Gamma_{1, \nabla(zz^\top)}(\nabla \log \frac{p}{\pi}, \nabla \log \frac{p}{\pi}) \right) dx \\
& - \int \nabla^2 \log \frac{p}{\pi} (aa^\top \nabla \log \frac{p}{\pi}, zz^\top \nabla \log \frac{p}{\pi}) p dx - \int \frac{1}{p} \nabla \cdot (\pi aa^\top \nabla \frac{p}{\pi}) \nabla \cdot (\pi zz^\top \nabla \frac{p}{\pi}) dx.
\end{aligned}$$

By subtracting the above two items, we derive the information Gamma  $z$  operator. ■

**5.4. Proof of Step 2.** We first introduce the following definition.

**Definition 3.** For any vector field  $\mathbf{U} \in \mathcal{C}^\infty(\mathbb{R}^{n+m})$ , we define vectors  $\mathbf{C}, \mathbf{F}, \mathbf{G}, \mathbf{V}^a \in \mathbb{R}^{(n+m)^2 \times 1}$ ,  $\mathbf{D} \in \mathbb{R}^{n^2 \times 1}$  and  $\mathbf{E} \in \mathbb{R}^{(n \times m) \times 1}$  as below. For  $\hat{i}, \hat{k} = 1, \dots, n+m$ ,

$$\begin{aligned} \mathbf{C}_{\hat{i}\hat{k}} &= \sum_{i,k=1}^n \left( a_{\hat{i}\hat{i}}^\top a_i^\top \nabla a_{\hat{k}\hat{k}}^\top - a_{\hat{i}\hat{k}}^\top a_k^\top \nabla a_{\hat{i}\hat{i}}^\top \right) a_k^\top \mathbf{U}, & \mathbf{D}_{ik} &= a_i^\top \nabla a_k^\top \mathbf{U}, \\ \mathbf{F}_{\hat{i}\hat{k}} &= \sum_{i=1}^n \sum_{k=1}^m \left( a_{\hat{i}\hat{i}}^\top a_i^\top \nabla z_{\hat{k}\hat{k}}^\top - z_k^\top \nabla a_{\hat{i}\hat{i}}^\top a_{\hat{k}\hat{k}}^\top \right) z_k^\top \mathbf{U}, & \mathbf{E}_{ik} &= a_i^\top \nabla z_k^\top \mathbf{U}, \\ \mathbf{G}_{\hat{i}\hat{k}} &= \sum_{i=1}^n \sum_{k=1}^m \left[ \left( z_{\hat{k}\hat{k}}^\top z_k^\top \nabla a_{\hat{i}\hat{i}}^\top a_i^\top \mathbf{U} + a_{\hat{i}\hat{i}}^\top z_{\hat{k}\hat{k}}^\top z_k^\top \nabla a_i^\top \mathbf{U} \right) - \left( a_{\hat{i}\hat{i}}^\top a_i^\top \nabla z_{\hat{k}\hat{k}}^\top z_k^\top \mathbf{U} + z_{\hat{k}\hat{k}}^\top a_{\hat{i}\hat{i}}^\top a_i^\top \nabla z_k^\top \mathbf{U} \right) \right], \\ \mathbf{V}_{\hat{i}\hat{k}}^a &= -\frac{1}{2} \sum_{i=1}^n \gamma_k a_{\hat{i}\hat{i}}^\top (a_i^\top \mathbf{U}) + \frac{1}{2} \sum_{k=1}^m a_{\hat{k}\hat{k}}^\top a_{\hat{k}\hat{i}}^\top \langle \mathbf{U}, \gamma \rangle. \end{aligned}$$

We further divide all needed calculations into the following steps. Let  $\mathbf{U} = \nabla f$  in Definition 3. We have the following facts.

**Proposition 7.** Denote  $f = \log \frac{p}{\pi}$ . Then

$$\int \Gamma_{\mathcal{I}_a}(f, f) p dx = \int \mathfrak{R}_{\gamma_a}(\nabla f, \nabla f) p dx, \quad \int \Gamma_{\mathcal{I}_z}(f, f) p dx = \int \mathfrak{R}_{\gamma_z}(\nabla f, \nabla f) p dx.$$

**Proof** We show the proof of the case  $\int [\Gamma_{\mathcal{I}_a}(f, f) + \Gamma_{\mathcal{I}_z}(f, f)] p dx$  below. For  $f = \log \frac{p}{\pi}$ , we have

$$\begin{aligned} & \int \Gamma_{\mathcal{I}_{a,z}}(f, f) p dx \\ &= \int \left[ (\tilde{L}f + \tilde{L}_z f) \langle \nabla f, \gamma \rangle - \frac{1}{2} \langle \nabla (\Gamma_1(f, f) + \Gamma_1^z(f, f)), \gamma \rangle \right] p dx \\ &= \int \left[ \nabla \cdot ((aa^\top + zz^\top) \nabla f) \langle \nabla f, \gamma \rangle + \langle \nabla f, \gamma \rangle \langle \nabla f, (aa^\top + zz^\top) \nabla \log \pi \rangle \right] p dx \\ & \quad + \int \frac{1}{2} \nabla \cdot (\gamma p) \langle \nabla f, (aa^\top + zz^\top) \nabla f \rangle dx \\ &= \int \left[ \nabla \cdot ((aa^\top + zz^\top) \nabla f) \langle \nabla f, \gamma \rangle + \langle \nabla f, \gamma \rangle \langle \nabla f, (aa^\top + zz^\top) \nabla \log \pi \rangle \right] p dx \\ & \quad + \int \frac{1}{2} \langle \nabla f, \gamma \rangle \langle \nabla f, (aa^\top + zz^\top) \nabla f \rangle p dx, \end{aligned}$$

where we use the fact  $\nabla \cdot (p\gamma) = \frac{\nabla \cdot (p\gamma)}{p} p = (\langle \nabla \log p, \gamma \rangle + \nabla \cdot \gamma) p = \langle \nabla f, \gamma \rangle p$ . Notice that

$$\begin{aligned} & \int \nabla \cdot ((aa^\top + zz^\top) \nabla f) \langle \nabla f, \gamma \rangle p dx \\ &= - \int \left[ \langle (aa^\top + zz^\top) \nabla f, \nabla \log p \rangle \langle \nabla f, \gamma \rangle + \langle (aa^\top + zz^\top) \nabla f, \nabla^2 f \gamma \rangle \right] p dx \\ & \quad - \int \langle (aa^\top + zz^\top) \nabla f, \nabla \gamma \nabla f \rangle p dx. \end{aligned}$$

Combining the above two terms, we have

$$\begin{aligned}
& \int \Gamma_{\mathcal{I}_{a,z}}(f, f) p dx \\
&= \int -\frac{1}{2} \langle \nabla f, \gamma \rangle \langle \nabla f, (aa^\top + zz^\top) \nabla f \rangle p \\
&\quad - \int \left[ \langle (aa^\top + zz^\top) \nabla f, \nabla \gamma \nabla f \rangle + \langle (aa^\top + zz^\top) \nabla f, \nabla^2 f \gamma \rangle \right] p dx \\
&= \int -\frac{1}{2} \langle \nabla p, \gamma \rangle \langle \nabla f, (aa^\top + zz^\top) \nabla f \rangle + \int \frac{1}{2} \langle \nabla \log \pi, \gamma \rangle \langle \nabla f, (aa^\top + zz^\top) \nabla f \rangle p dx \\
&\quad - \int \left[ \langle (aa^\top + zz^\top) \nabla f, \nabla \gamma \nabla f \rangle + \langle (aa^\top + zz^\top) \nabla f, \nabla^2 f \gamma \rangle \right] p dx.
\end{aligned}$$

In other words,

$$\begin{aligned}
& \int \Gamma_{\mathcal{I}_{a,z}}(f, f) p dx \\
&= \int \frac{1}{2} \nabla \cdot (\gamma \langle \nabla f, (aa^\top + zz^\top) \nabla f \rangle) p + \int \frac{1}{2} \langle \nabla \log \pi, \gamma \rangle \langle \nabla f, (aa^\top + zz^\top) \nabla f \rangle p dx \\
&\quad - \int \left[ \langle (aa^\top + zz^\top) \nabla f, \nabla \gamma \nabla f \rangle + \langle (aa^\top + zz^\top) \nabla f, \nabla^2 f \gamma \rangle \right] p dx \\
&= \frac{1}{2} \int \langle \gamma, \langle \nabla f, \nabla (aa^\top + zz^\top) \nabla f \rangle \rangle p dx - \int \langle \nabla \gamma \nabla f, (aa^\top + zz^\top) \nabla f \rangle p dx,
\end{aligned}$$

where we use the fact  $\frac{\nabla \cdot (\pi \gamma)}{\pi} = \langle \nabla \log \pi, \gamma \rangle + \nabla \cdot \gamma = 0$ . The proof is completed.  $\blacksquare$

**Proposition 8.** For any  $f \in C^\infty(\mathbb{R}^{n+m})$ , we have

$$\Gamma_{\mathcal{I}_a}(f, f) = \mathfrak{R}_{\mathcal{I}_a}(\nabla f, \nabla f) + 2V_a^\top X.$$

**Proof** From our definition,

$$\Gamma_{\mathcal{I}_a}(f, f) = \langle \nabla f, \gamma(x) \rangle \tilde{L}f - \frac{1}{2} \langle \nabla \Gamma_1(f, f), \gamma(x) \rangle.$$

We look at the first term by

$$\begin{aligned}
\langle \nabla f, \gamma \rangle \tilde{L}f &= \langle \nabla f, \gamma \rangle \Delta_a f + \langle \nabla f, \gamma \rangle \langle aa^\top \nabla \log \pi, \nabla f \rangle \\
&= \langle \nabla f, \gamma \rangle \left[ (\nabla a) \circ (a^\top \nabla f) + \sum_{i=1}^n a_i^\top \nabla a_i^\top \nabla f + \sum_{i=1}^n a_i^\top a_i^\top \nabla^2 f \right] \\
&\quad + \langle \nabla f, \gamma \rangle \langle aa^\top \nabla \log \pi, \nabla f \rangle,
\end{aligned}$$

where we use the fact that

$$\begin{aligned}
\Delta_a f &= \nabla \cdot (aa^\top \nabla f) \\
&= (\nabla a) \circ (a^\top \nabla f) + a^\top \nabla \circ (a^\top \nabla f) \\
&= (\nabla a) \circ (a^\top \nabla f) + \sum_{i=1}^n a_i^\top \nabla a_i^\top \nabla f + \sum_{i=1}^n a_i^\top a_i^\top \nabla^2 f.
\end{aligned}$$



We further reformulate the cross terms as below,

$$\langle \nabla f, \gamma \rangle \sum_{i=1}^n a_i^\top a_i^\top \nabla^2 f = \langle \nabla f, \gamma \rangle \sum_{i=1}^n a_{ii}^\top a_{ik}^\top \frac{\partial^2 f}{\partial x_i \partial x_k}.$$

As for the second term, we have

$$\begin{aligned} \frac{1}{2} \gamma \nabla \Gamma_1(f, f) &= \frac{1}{2} \sum_{k=1}^{n+m} \gamma_k \frac{\partial}{\partial x_k} (\langle a^\top \nabla f, a^\top \nabla f \rangle_{\mathbb{R}^n}) \\ &= \sum_{k,i=1}^{n+m} \sum_{i=1}^n \left( \gamma_k \frac{\partial a_{ii}^\top}{\partial x_k} \frac{\partial f}{\partial x_i} + \gamma_k a_{ii}^\top \frac{\partial^2 f}{\partial x_k \partial x_i} \right) (a_i^\top \nabla f), \\ &= \sum_{i=1}^n \sum_{k=1}^{n+m} \gamma_k \nabla_{x_k} a_i^\top \nabla f (a_i^\top \nabla f) + \sum_{i=1}^n \sum_{k=1}^{n+m} \gamma_k a_i^\top \nabla^2 f (a_i^\top \nabla f). \end{aligned}$$

Combining the above terms and taking into account the Definition 21, we derive  $\Gamma_{\mathcal{I}_a}(f, f)$ .

■

**Proposition 9.** *For any  $f \in C^\infty(\mathbb{R}^{n+m})$ , we have*

$$\tilde{\Gamma}_2(f, f) = (\mathbf{QX} + \mathbf{D})^\top (\mathbf{QX} + \mathbf{D}) + 2\mathbf{C}^\top \mathbf{X} + \mathfrak{R}_a(\nabla f, \nabla f),$$

$$\tilde{\Gamma}_2^z(f, f) = (\mathbf{PX} + \mathbf{E})^\top (\mathbf{PX} + \mathbf{E}) + 2\mathbf{F}^\top \mathbf{X} + \mathfrak{R}_z(\nabla f, \nabla f),$$

$$\mathbf{div}_z^\pi(\Gamma_{\nabla(aa^\top)} f, f) - \mathbf{div}_a^\pi(\Gamma_{\nabla(zz^\top)} f, f) = \mathfrak{R}_\pi(\nabla f, \nabla f) + 2\mathbf{G}^\top \mathbf{X}.$$

The detailed derivations of the above propositions are given in the following lemmas.

The proof of Proposition 9 is proved in [22][Lemma 3.8, Lemma 3.10, Lemma 3.17, Lemma 3.18, and Lemma 3.25]. We address the main differences in the current paper and give a self-contained proof. For notation convenience, we recall the operator  $\tilde{L}$  as below,

$$\tilde{L}f = \nabla \cdot (aa^\top \nabla f) + \langle aa^\top \nabla \log \pi, \nabla f \rangle,$$

and we denote

$$\tilde{L}f = \Delta_a f + \langle aa^\top \nabla \log \pi, \nabla f \rangle, \quad (21)$$

with

$$\Delta_a f = \nabla \cdot (aa^\top \nabla f), \quad (22)$$

and

$$\Gamma_{2,a} = \frac{1}{2} \Delta_a \Gamma_1(f, f) - \Gamma_1(\Delta_a f, f). \quad (23)$$

5.4.1. *Derivation of  $\tilde{\Gamma}_2(f, f)$ .*

**Lemma 4.**

$$\tilde{\Gamma}_2(f, f) = \Gamma_{2,a}(f, f) + \sum_{i=1}^n \sum_{\hat{k}=1}^{n+m} \left[ (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} a_i^\top \nabla f - a_i^\top \nabla (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} f \right] a_i^\top \nabla f.$$

**Proof** [Proof of Lemma 4] From our definition, one can get

$$\tilde{\Gamma}_2(f, f) = \Gamma_{2,a}(f, f) + \frac{1}{2} \langle aa^\top \nabla \log \pi, \nabla \Gamma_1(f, f) \rangle - \Gamma_1(\langle aa^\top \nabla \log \pi, \nabla f \rangle, f),$$

where we take  $b = aa^\top \nabla \log \pi$ , we get

$$\begin{aligned} \frac{1}{2} b \nabla \Gamma_1(f, f) &= \frac{1}{2} \sum_{\hat{k}=1}^{n+m} b_{\hat{k}} \frac{\partial}{\partial x_{\hat{k}}} (\langle a^T \nabla f, a^T \nabla f \rangle_{\mathbb{R}^n}) \\ &= \sum_{\hat{k}, \hat{i}=1}^{n+m} \sum_{i=1}^n (b_{\hat{k}} \frac{\partial a_{i\hat{i}}^T}{\partial x_{\hat{k}}} \frac{\partial f}{\partial x_{\hat{i}}} + b_{\hat{k}} a_{i\hat{i}}^T \frac{\partial^2 f}{\partial x_{\hat{k}} \partial x_{\hat{i}}}) (a^T \nabla f)_i, \end{aligned}$$

and

$$\begin{aligned} -\Gamma_1(b \nabla f, f) &= -\langle a^T \nabla (b \nabla f), a^T \nabla f \rangle_{\mathbb{R}^n} \\ &= -\sum_{i=1}^n \sum_{\hat{k}, \hat{i}=1}^{n+m} (a_{i\hat{i}}^\top \frac{\partial b_{\hat{k}}}{\partial x_{\hat{i}}} \frac{\partial f}{\partial x_{\hat{k}}} + a_{i\hat{i}}^\top b_{\hat{k}} \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}}) (a^T \nabla f)_i. \end{aligned}$$

Summing over all the terms, we have

$$\begin{aligned} \tilde{\Gamma}_2(f, f) &= \Gamma_{2,a}(f, f) + \sum_{\hat{k}, \hat{i}=1}^{n+m} \sum_{i=1}^n (b_{\hat{k}} \frac{\partial a_{i\hat{i}}^T}{\partial x_{\hat{k}}} \frac{\partial f}{\partial x_{\hat{i}}} - a_{i\hat{i}}^\top \frac{\partial b_{\hat{k}}}{\partial x_{\hat{i}}} \frac{\partial f}{\partial x_{\hat{k}}}) (a^T \nabla f)_i \\ &= \Gamma_{2,a}(f, f) + \sum_{i=1}^n \sum_{\hat{k}=1}^{n+m} \left[ (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} a_i^\top \nabla f - a_i^\top \nabla (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} f \right] a_i^\top \nabla f. \end{aligned}$$

■

**Lemma 5.**

$$\begin{aligned} &\frac{1}{2} \Delta_a \Gamma_1(f, f) - \Gamma_1(\Delta_a f, f) \\ &= \frac{1}{2} (a^\top \nabla \circ (a^\top \nabla |a^\top \nabla f|^2)) - \langle a^\top \nabla ([(a^\top \nabla) \circ (a^\top \nabla f)]), a^\top \nabla f \rangle_{\mathbb{R}^n} \\ &\quad + \nabla a \circ \left( \sum_{k=1}^n [a^\top \nabla a_k^\top \nabla f - a_k^\top \nabla a^\top \nabla f] a_k^\top \nabla f \right) - \langle (a^\top \nabla^2 a \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n}. \end{aligned} \tag{24}$$

**Proof** [Proof of Lemma 5] According to our definition, we have

$$\begin{aligned} \Delta_a \Gamma_1(f, f) &= \nabla \cdot (aa^\top |a^\top \nabla f|^2) \\ &= \nabla a \circ (a^\top \nabla |a^\top \nabla f|^2) + (a^\top \nabla) \circ (a^\top \nabla |a^\top \nabla f|^2). \end{aligned}$$

Similarly, we have

$$\nabla \cdot (aa^\top \nabla f) = \nabla a \circ (a^\top \nabla f) + (a^\top \nabla) \circ (a^\top \nabla f),$$

which gives us

$$\begin{aligned} \Gamma_1(\Delta_a f, f) &= \langle a^\top \nabla(\nabla \cdot (aa^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n}, \\ &= \langle a^\top \nabla(\nabla a \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n} + \langle a^\top \nabla((a^\top \nabla) \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n} \\ &= \langle (a^\top \nabla^2 a \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n} + \langle (\nabla a \circ (a^\top \nabla(a^\top \nabla f))), a^\top \nabla f \rangle_{\mathbb{R}^n} \\ &\quad + \langle a^\top \nabla((a^\top \nabla) \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n}. \end{aligned}$$

Combining the above terms, we get

$$\begin{aligned} &\frac{1}{2} \Delta_a \Gamma_1(f, f) - \Gamma_1, a(\Delta_a f, f) \\ &= \frac{1}{2} (a^\top \nabla \circ (a^\top \nabla |a^\top \nabla f|^2)) - \langle a^\top \nabla((a^\top \nabla) \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n} + \underbrace{\frac{1}{2} \nabla a \circ (a^\top \nabla |a^\top \nabla f|^2)}_{\mathbf{I}} \\ &\quad - \langle (a^\top \nabla^2 a \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n} - \langle (\nabla a \circ (a^\top \nabla(a^\top \nabla f))), a^\top \nabla f \rangle_{\mathbb{R}^n} \cdots \mathbf{II} \end{aligned}$$

Recalling the fact  $a_{ii}^\top = a_{ii}$ , we have

$$\begin{aligned} \mathbf{I} &= \frac{1}{2} \nabla a \circ (a^\top \nabla |a^\top \nabla f|^2) \\ &= \nabla a \circ \left( \sum_{k=1}^n a^\top \nabla (a_k^\top \nabla f) a_k^\top \nabla f \right) \\ &= \nabla a \circ \left( \sum_{k=1}^n a^\top \nabla a_k^\top \nabla f a_k^\top \nabla f \right) + \nabla a \circ \left( \sum_{k=1}^n a^\top a_k^\top \nabla^2 f a_k^\top \nabla f \right), \end{aligned}$$

and

$$\begin{aligned} \mathbf{II} &= -\langle (a^\top \nabla^2 a \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n} - \sum_{k=1}^n \langle (\nabla a \circ (a_k^\top \nabla(a^\top \nabla f))), a_k^\top \nabla f \rangle \\ &= -\langle (a^\top \nabla^2 a \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n} - \sum_{k=1}^n \langle (\nabla a \circ (a_k^\top a^\top \nabla^2 f)), a_k^\top \nabla f \rangle \\ &\quad - \sum_{k=1}^n \langle (\nabla a \circ (a_k^\top \nabla a^\top \nabla f)), a_k^\top \nabla f \rangle. \end{aligned}$$

Subtracting the above two terms, we end up with

$$\mathbf{I} - \mathbf{II} = \nabla a \circ \left( \sum_{k=1}^n [a^\top \nabla a_k^\top \nabla f - a_k^\top \nabla a^\top \nabla f] a_k^\top \nabla f \right) - \langle (a^\top \nabla^2 a \circ (a^\top \nabla f)), a^\top \nabla f \rangle_{\mathbb{R}^n}.$$

Now, we eventually derive the the following step

$$\begin{aligned}
& \frac{1}{2}\Delta_a\Gamma_1(f, f) - \Gamma_1(\Delta_a f, f) \\
= & \frac{1}{2}(a^\top\nabla \circ (a^\top\nabla|a^\top\nabla f|^2)) - \langle a^\top\nabla((a^\top\nabla) \circ (a^\top\nabla f)), a^\top\nabla f \rangle_{\mathbb{R}^n} \\
& + \nabla a \circ \left( \sum_{k=1}^n [a^\top\nabla a_k^\top\nabla f - a_k^\top\nabla a^\top\nabla f] a_k^\top\nabla f \right) - \langle (a^\top\nabla^2 a \circ (a^\top\nabla f)), a^\top\nabla f \rangle_{\mathbb{R}^n}.
\end{aligned}$$

The proof is completed. ■

**Lemma 6.**

$$\begin{aligned}
& \frac{1}{2}(a^\top\nabla \circ (a^\top\nabla|a^\top\nabla f|^2)) - \langle a^\top\nabla((a^\top\nabla) \circ (a^\top\nabla f)), a^\top\nabla f \rangle_{\mathbb{R}^n} \\
= & [\mathbf{Q}\mathbf{X} + \mathbf{D}]^\top[\mathbf{Q}\mathbf{X} + \mathbf{D}] + 2\mathbf{C}^\top\mathbf{X} \\
& + \sum_{i,k=1}^n a_i^\top\nabla a_i^\top\nabla a_k^\top\nabla f(a_k^\top\nabla f) + \sum_{i,k=1}^n a_i^\top a_i^\top\nabla^2 a_k^\top\nabla f(a_k^\top\nabla f) \\
& - \sum_{i,k=1}^n a_k^\top\nabla a_i^\top\nabla a_i^\top\nabla f(a_k^\top\nabla f) - \sum_{i,k=1}^n a_k^\top a_i^\top\nabla^2 a_i^\top\nabla f(a_k^\top\nabla f),
\end{aligned}$$

where the vectors  $\mathbf{C}$ ,  $\mathbf{D}$  and the matrix  $\mathbf{Q}$  are defined in Definition 3 and formula (19).

**Proof** [Proof of Lemma 6]

$$\begin{aligned}
& \frac{1}{2}(a^\top\nabla \circ (a^\top\nabla|a^\top\nabla f|^2)) - \langle a^\top\nabla([(a^\top\nabla) \circ (a^\top\nabla f)]), a^\top\nabla f \rangle_{\mathbb{R}^n} \\
= & \frac{1}{2} \sum_{i,k=1}^n (a_i^\top\nabla)(a_i^\top\nabla)|a_k^\top\nabla f|^2 - \sum_{i,k=1}^n (a_k^\top\nabla)[(a_i^\top\nabla)(a_i^\top\nabla f)](a_k^\top\nabla f) \\
= & \sum_{i,k=1}^n a_i^\top\nabla[(a_i^\top\nabla)(a_k^\top\nabla f)(a_k^\top\nabla f)] - \sum_{i,k=1}^n (a_k^\top\nabla)[(a_i^\top\nabla)(a_i^\top\nabla f)](a_k^\top\nabla f) \\
= & \sum_{i,k=1}^n |(a_i^\top\nabla)(a_k^\top\nabla f)|^2 + \sum_{i,k=1}^n \langle (a_i^\top\nabla)(a_i^\top\nabla)(a_k^\top\nabla f)(a_k^\top\nabla f) \rangle - \sum_{i,k=1}^n (a_k^\top\nabla)[(a_i^\top\nabla)(a_i^\top\nabla f)](a_k^\top\nabla f) \\
= & \mathbf{T}_1 + \mathbf{T}_2 - \mathbf{T}_3.
\end{aligned}$$

We then expand  $\mathbf{T}_1$ ,  $\mathbf{T}_2$  and  $\mathbf{T}_3$ .

$$\begin{aligned}
\mathbf{T}_1 &= \sum_{i,k=1}^n |(a_i^\top\nabla)(a_k^\top\nabla f)|^2 = \sum_{i,k=1}^n |a_i^\top a_k^\top\nabla^2 f + a_i^\top\nabla a_k^\top\nabla f|^2 \\
&= \sum_{i,k=1}^n \left[ |a_i^\top a_k^\top\nabla^2 f|^2 + 2(a_i^\top a_k^\top\nabla^2 f)(a_i^\top\nabla a_k^\top\nabla f) + |a_i^\top\nabla a_k^\top\nabla f|^2 \right] \\
&= [\mathbf{Q}\mathbf{X} + \mathbf{D}]^\top[\mathbf{Q}\mathbf{X} + \mathbf{D}],
\end{aligned}$$

$$\begin{aligned}
\mathbf{T}_2 &= \sum_{i,k=1}^n \langle (a_i^\top \nabla)(a_i^\top \nabla)(a_k^\top \nabla f)(a_k^\top \nabla f) \rangle \\
&= \sum_{i,k=1}^n a_i^\top a_i^\top a_k^\top \nabla^3 f(a_k^\top \nabla f) + \sum_{i,k=1}^n a_i^\top \nabla a_i^\top \nabla a_k^\top \nabla f(a_k^\top \nabla f) + \sum_{i,k=1}^n a_i^\top a_i^\top \nabla^2 a_k^\top \nabla f(a_k^\top \nabla f) \\
&\quad + 2 \sum_{i,k=1}^n a_i^\top a_i^\top \nabla a_k^\top \nabla^2 f(a_k^\top \nabla f) + \sum_{i,k=1}^n a_i^\top \nabla a_i^\top a_k^\top \nabla^2 f(a_k^\top \nabla f),
\end{aligned}$$

and

$$\begin{aligned}
\mathbf{T}_3 &= \sum_{i,k=1}^n (a_k^\top \nabla)[(a_i^\top \nabla)(a_i^\top \nabla f)](a_k^\top \nabla f) \\
&= \sum_{i,k=1}^n a_k^\top a_i^\top a_i^\top \nabla^3 f(a_k^\top \nabla f) + \sum_{i,k=1}^n a_k^\top \nabla a_i^\top \nabla a_i^\top \nabla f(a_k^\top \nabla f) + \sum_{i,k=1}^n a_k^\top a_i^\top \nabla^2 a_i^\top \nabla f(a_k^\top \nabla f) \\
&\quad + \sum_{i,k=1}^n a_k^\top a_i^\top \nabla a_i^\top \nabla^2 f(a_k^\top \nabla f) + 2 \sum_{i,k=1}^n a_k^\top \nabla a_i^\top a_i^\top \nabla^2 f(a_k^\top \nabla f).
\end{aligned}$$

Combing the above three terms, we have the following equality.

$$\begin{aligned}
&\mathbf{T}_1 + \mathbf{T}_2 - \mathbf{T}_3 \\
&= \sum_{i,k=1}^n a_i^\top \nabla a_i^\top \nabla a_k^\top \nabla f(a_k^\top \nabla f) + \sum_{i,k=1}^n a_i^\top a_i^\top \nabla^2 a_k^\top \nabla f(a_k^\top \nabla f) \\
&\quad - \sum_{i,k=1}^n a_k^\top \nabla a_i^\top \nabla a_i^\top \nabla f(a_k^\top \nabla f) - \sum_{i,k=1}^n a_k^\top a_i^\top \nabla^2 a_i^\top \nabla f(a_k^\top \nabla f) \\
&\quad + [\mathbf{QX} + \mathbf{D}]^\top [\mathbf{QX} + \mathbf{D}] + 2 \sum_{i,k=1}^n a_i^\top a_i^\top \nabla a_k^\top \nabla^2 f(a_k^\top \nabla f) - 2 \sum_{i,k=1}^n a_k^\top \nabla a_i^\top a_i^\top \nabla^2 f(a_k^\top \nabla f).
\end{aligned}$$

Furthermore, we investigate the last two terms in the above formula,

$$\begin{aligned}
&2 \sum_{i,k=1}^n a_i^\top a_i^\top \nabla a_k^\top \nabla^2 f(a_k^\top \nabla f) - 2 \sum_{i,k=1}^n a_k^\top \nabla a_i^\top a_i^\top \nabla^2 f(a_k^\top \nabla f) \\
&= 2 \sum_{\hat{i}, \hat{k}=1}^{n+m} \left[ \sum_{i,k=1}^n \sum_{i'=1}^{n+m} \left( \langle a_{\hat{i}\hat{i}}^\top a_{i'i'}^\top \left( \frac{\partial a_{k\hat{k}}^\top}{\partial x_{i'}} \right), (a^\top \nabla)_k f \rangle - \langle a_{k\hat{i}'}^\top a_{i\hat{k}}^\top \frac{\partial a_{\hat{i}\hat{i}}^\top}{\partial x_{i'}}, (a^\top \nabla)_k f \rangle \right) \right] \left( \frac{\partial}{\partial x_{\hat{i}}} \frac{\partial f}{\partial x_{\hat{k}}} \right) \\
&= \sum_{\hat{i}, \hat{k}=1}^n 2\mathbf{C}_{\hat{i}\hat{k}} \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}} = 2\mathbf{C}^\top \mathbf{X},
\end{aligned}$$

which completes the proof.  $\blacksquare$

5.4.2. *Derivation of  $\tilde{\Gamma}_{2,z}(f, f)$  calculus.* Similar to the proof of  $\tilde{\Gamma}_2(f, f)$ , we directly give the key lemmas, and skip the routine computations.

**Lemma 7.**

$$\begin{aligned}\tilde{\Gamma}_2^z(f, f) &= \frac{1}{2}\Delta_a\Gamma_{1,z} - \Gamma_{1,z}(\Delta_a f, f) \\ &\quad + \sum_{i=1}^n \sum_{\hat{k}=1}^{n+m} \left[ (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} z_i^\top \nabla f - z_i^\top \nabla (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} f \right] z_i^\top \nabla f.\end{aligned}$$

**Proof** [Proof of Lemma 7] From our definition, one can get

$$\tilde{\Gamma}_{2,z}(f, f) = \frac{1}{2}\Delta_a\Gamma_{1,z} - \Gamma_{1,z}(\Delta_a f, f) + \frac{1}{2}\langle aa^\top \nabla \log \pi, \nabla \Gamma_{1,z}(f, f) \rangle - \Gamma_{1,z}(\langle aa^\top \nabla \log \pi, \nabla f \rangle, f),$$

where we take  $b = aa^\top \nabla \log \pi$ , we get

$$\begin{aligned}\frac{1}{2}b \nabla \Gamma_{1,z}(f, f) &= \frac{1}{2} \sum_{\hat{k}=1}^{n+m} b_{\hat{k}} \frac{\partial}{\partial x_{\hat{k}}} (\langle z^T \nabla f, z^T \nabla f \rangle_{\mathbb{R}^m}) \\ &= \sum_{\hat{k}, \hat{i}=1}^{n+m} \sum_{i=1}^m (b_{\hat{k}} \frac{\partial z_{\hat{i}}^T}{\partial x_{\hat{k}}} \frac{\partial f}{\partial x_i} + b_{\hat{k}} z_{\hat{i}}^T \frac{\partial^2 f}{\partial x_{\hat{k}} \partial x_i}) (z^T \nabla f)_i,\end{aligned}$$

and

$$\begin{aligned}-\Gamma_{1,z}(b \nabla f, f) &= -\langle z^T \nabla (b \nabla f), z^T \nabla f \rangle_{\mathbb{R}^m} \\ &= -\sum_{i=1}^m \sum_{\hat{k}, \hat{i}=1}^{n+m} (z_{\hat{i}}^\top \frac{\partial b_{\hat{k}}}{\partial x_i} \frac{\partial f}{\partial x_{\hat{k}}} + z_{\hat{i}}^\top b_{\hat{k}} \frac{\partial^2 f}{\partial x_i \partial x_{\hat{k}}}) (a^T \nabla f)_i.\end{aligned}$$

Summing over all the terms, we have

$$\begin{aligned}\tilde{\Gamma}_2(f, f) &= \frac{1}{2}\Delta_a\Gamma_{1,z} - \Gamma_{1,z}(\Delta_a f, f) \\ &\quad + \sum_{i=1}^m \sum_{\hat{k}=1}^{n+m} \left[ (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} z_i^\top \nabla f - z_i^\top \nabla (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} f \right] z_i^\top \nabla f.\end{aligned}$$

■

**Lemma 8.**

$$\begin{aligned}&\frac{1}{2}\Delta_a\Gamma_{1,z} - \Gamma_{1,z}(\Delta_a f, f) \tag{25} \\ &= \frac{1}{2}(a^\top \nabla \circ (a^\top \nabla |z^\top \nabla f|^2)) - \langle z^\top \nabla ((a^\top \nabla) \circ (a^\top \nabla f)), z^\top \nabla f \rangle_{\mathbb{R}^m} \\ &\quad + \nabla a \circ \left( \sum_{k=1}^m [a^\top \nabla z_k^\top \nabla f - z_k^\top \nabla a^\top \nabla f] z_k^\top \nabla f \right) - \langle (z^\top \nabla^2 a \circ (a^\top \nabla f)), z^\top \nabla f \rangle_{\mathbb{R}^m}.\end{aligned}$$

We further investigate the extra term explicitly in Lemma 8.

**Lemma 9.**

$$\begin{aligned}
& \frac{1}{2}(a^\top \nabla \circ (a^\top \nabla |z^\top \nabla f|^2)) - \langle z^\top \nabla ((a^\top \nabla) \circ (a^\top \nabla f)), z^\top \nabla f \rangle_{\mathbb{R}^m} \\
&= [\mathbf{P}\mathbf{X} + \mathbf{E}]^\top [\mathbf{P}\mathbf{X} + \mathbf{E}] + 2\mathbf{F}^\top \mathbf{X} \\
&+ \sum_{i=1}^n \sum_{k=1}^m a_i^\top \nabla a_i^\top \nabla z_k^\top \nabla f (z_k^\top \nabla f) + \sum_{i=1}^n \sum_{k=1}^m a_i^\top a_i^\top \nabla^2 z_k^\top \nabla f (z_k^\top \nabla f) \\
&- \sum_{i=1}^n \sum_{k=1}^m z_k^\top \nabla a_i^\top \nabla a_i^\top \nabla f (z_k^\top \nabla f) - \sum_{i=1}^n \sum_{k=1}^m z_k^\top a_i^\top \nabla^2 a_i^\top \nabla f (z_k^\top \nabla f),
\end{aligned}$$

where vectors  $\mathbf{F}$ ,  $\mathbf{E}$ , and the matrix  $\mathbf{P}$  are defined in Definition 3 and formula (19).

*Remark 10.* The proof of the identity

$$\mathbf{div}_z^\pi(\Gamma_{\nabla(aa^\top)} f, f) - \mathbf{div}_a^\pi(\Gamma_{\nabla(zz^\top)} f, f) = \mathfrak{R}_\pi(\nabla f, \nabla f) + 2\mathbf{G}^\top \mathbf{X},$$

is the same as in [22][Lemma 3.25]. We skip the proof here.

Here Lemma 8 and Lemma 9 follow similar proofs in [22]. We omitted them here.

**5.5. Proof of Theorem 3.** Combining all above propositions, we are now ready to prove an *information Bochner's formula*.

**Proof** [Proof of Theorem 3] We show the proof in the following three steps.

**Step A:** According to the above three propositions, we have

$$\begin{aligned}
& \int \left[ \tilde{\Gamma}_2(f, f) + \tilde{\Gamma}_2^{z, \pi}(f, f) + \Gamma_{\mathcal{I}_{a, z}}(f, f) \right] p dx \\
&= \int \left[ \tilde{\Gamma}_2(f, f) + \tilde{\Gamma}_2^{z, \pi}(f, f) + (\beta + (1 - \beta))\Gamma_{\mathcal{I}_a}(f, f) + \Gamma_{\mathcal{I}_z}(f, f) \right] p dx \\
&= \int \left[ (\mathbf{Q}\mathbf{X} + \mathbf{D})^\top (\mathbf{Q}\mathbf{X} + \mathbf{D}) + 2\mathbf{C}^\top \mathbf{X} + \mathfrak{R}_a(\nabla f, \nabla f) \right] p dx \\
&+ \int \left[ (\mathbf{P}\mathbf{X} + \mathbf{E})^\top (\mathbf{P}\mathbf{X} + \mathbf{E}) + 2\mathbf{F}^\top \mathbf{X} + \mathfrak{R}_z(\nabla f, \nabla f) + \mathfrak{R}_\pi(\nabla f, \nabla f) + 2\mathbf{G}^\top \mathbf{X} \right] p dx \\
&+ \int \left[ \beta(\mathfrak{R}_{\mathcal{I}_a}(\nabla f, \nabla f) + 2\mathbf{V}_a^\top \mathbf{X}) + (1 - \beta)\mathfrak{R}_{\gamma_a}(\nabla f, \nabla f) + \mathfrak{R}_{\gamma_z}(\nabla f, \nabla f) \right] p dx.
\end{aligned}$$

**Step B:** Under Assumption 1, we show that there exists vectors  $\Lambda_1, \Lambda_2 \in \mathbb{R}^{(n+m)^2 \times 1}$  such that, for any  $\beta \in \mathbb{R}^1$ , we have

$$(\mathbf{Q}^\top \Lambda_1 + \mathbf{P}^\top \Lambda_2)^\top \mathbf{X} = (\mathbf{F} + \mathbf{C} + \mathbf{G} + \beta \mathbf{V}^a + \mathbf{Q}^\top \mathbf{D} + \mathbf{P}^\top \mathbf{E})^\top \mathbf{X}. \quad (26)$$

The main idea is to complete squares for all the second order terms listed below

$$(\mathbf{Q}\mathbf{X} + \mathbf{D})^\top (\mathbf{Q}\mathbf{X} + \mathbf{D}) + (\mathbf{P}\mathbf{X} + \mathbf{E})^\top (\mathbf{P}\mathbf{X} + \mathbf{E}) + 2\mathbf{C}^\top \mathbf{X} + 2\mathbf{F}^\top \mathbf{X} + 2\mathbf{G}^\top \mathbf{X} + 2\mathbf{V}_a^\top \mathbf{X},$$

where the leading terms are given by

$$\mathbf{X}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{X} = \sum_{i, k=1}^n |a_i^\top a_k^\top \nabla^2 f|^2, \quad \mathbf{X}^\top \mathbf{P}^\top \mathbf{P} \mathbf{X} = \sum_{i=1}^n \sum_{j=1}^m |a_i^\top z_j^\top \nabla^2 f|^2.$$

It is enough to show that for  $\mathbf{C}^\top \mathbf{X}$ , we have

**Claim 1:**

$$\mathbf{C}^\top \mathbf{X} = \sum_{i,k=1}^n (a_i^\top a_k^\top \nabla^2 f) \tilde{\mathbf{C}}_1(\mathbf{U})_{ik} + \sum_{i=1}^n \sum_{k=1}^m (a_i^\top z_k^\top \nabla^2 f) \tilde{\mathbf{C}}_2(\mathbf{U})_{ik}. \quad (27)$$

We show the details for  $\mathbf{C}^\top \mathbf{X}$  as below, and skip the details for the similar terms in  $\mathbf{F}^\top \mathbf{X}, \mathbf{G}^\top \mathbf{X}, \mathbf{V}^a \mathbf{X}$ . We have

$$\mathbf{C}^\top \mathbf{X} = \sum_{\hat{i}, \hat{k}=1}^{n+m} \mathbf{C}_{\hat{i}\hat{k}} \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}} = \sum_{\hat{i}, \hat{k}=1}^{n+m} \sum_{i,k=1}^n \left( a_{\hat{i}\hat{i}}^\top a_i^\top \nabla a_{k\hat{k}}^\top - a_{\hat{i}\hat{k}}^\top a_k^\top \nabla a_{\hat{i}\hat{i}}^\top \right) a_k^\top \mathbf{U} \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}}.$$

We look at the first term,  $a_i^\top \nabla a_{k\hat{k}}^\top = \sum_{i'=1}^{n+m} a_{i i'}^\top \nabla_{i'} a_{k\hat{k}}^\top$ , and apply Definition 1. Hence we derive the following expression by

$$\begin{aligned} & \sum_{\hat{i}, \hat{k}=1}^{n+m} \sum_{i,k=1}^n \left( a_{\hat{i}\hat{i}}^\top a_i^\top \nabla a_{k\hat{k}}^\top \right) a_k^\top \mathbf{U} \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}} \\ &= \sum_{\hat{i}, \hat{k}=1}^{n+m} \sum_{i,k,l=1}^n \left( a_{\hat{i}\hat{i}}^\top a_{i i'}^\top \lambda_l^{i'k} a_{l\hat{k}}^\top \right) a_k^\top \mathbf{U} \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}} + \sum_{\hat{i}, \hat{k}=1}^{n+m} \sum_{i,k=1}^n \sum_{l=1}^m \left( a_{\hat{i}\hat{i}}^\top a_{i i'}^\top \lambda_{l+n}^{i'k} z_{l\hat{k}}^\top \right) a_k^\top \mathbf{U} \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}} \\ &= \sum_{i,l=1}^n (a_i^\top a_l^\top \nabla^2 f) \left( \sum_{k=1}^n \sum_{i'=1}^{n+m} a_{i i'}^\top \lambda_l^{i'k} a_k^\top \mathbf{U} \right) + \sum_{i=1}^n \sum_{l=1}^m (a_i^\top z_l^\top \nabla^2 f) \left( \sum_{k=1}^n \sum_{i'=1}^{n+m} a_{i i'}^\top \lambda_{l+n}^{i'k} a_k^\top \mathbf{U} \right). \end{aligned}$$

We have the similar expansion for the second term of  $\mathbf{C}^\top \mathbf{X}$ . Thus

$$\begin{aligned} \tilde{\mathbf{C}}_1(\mathbf{U})_{il} &= \sum_{k=1}^n \left[ \sum_{i'=1}^{n+m} a_{i i'}^\top \lambda_l^{i'k} - \sum_{k'=1}^{n+m} a_{k k'}^\top \lambda_l^{k'i} \right] a_k^\top \mathbf{U}, \\ \tilde{\mathbf{C}}_2(\mathbf{U})_{il} &= \sum_{k=1}^n \left[ \sum_{i'=1}^{n+m} a_{i i'}^\top \lambda_{l+n}^{i'k} - \sum_{k'=1}^{n+m} a_{k k'}^\top \lambda_{l+n}^{k'i} \right] a_k^\top \mathbf{U}, \end{aligned}$$

which proves Claim 1 in (27). It is by a straightforward observation. We can show the same property for vectors  $\mathbf{F}$ ,  $\mathbf{V}^a$  and the last three terms of vector  $\mathbf{G}$ . We now just need to further look at the first term of vector  $\mathbf{G}$  as defined in Definition 1,

$$\begin{aligned} & \text{First Term of } \mathbf{G}^\top \mathbf{X} \\ &= \sum_{i=1}^n \sum_{k=1}^m \left( z_{k\hat{k}}^\top z_k^\top \nabla a_{\hat{i}\hat{i}}^\top a_i^\top \mathbf{U} \right) \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}} \\ &= \sum_{i,l=1}^n \sum_{k=1}^m \sum_{k'=1}^{n+m} \left( z_{k\hat{k}}^\top z_{k k'}^\top \lambda_l^{k'i} a_{l\hat{i}}^\top a_i^\top \mathbf{U} \right) \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}} + \sum_{i=1}^n \sum_{k,l=1}^m \sum_{k'=1}^{n+m} \left( z_{k\hat{k}}^\top z_{k k'}^\top \lambda_{l+n}^{k'i} z_{l\hat{i}}^\top a_i^\top \mathbf{U} \right) \frac{\partial^2 f}{\partial x_{\hat{i}} \partial x_{\hat{k}}} \\ &= \sum_{k=1}^m \sum_{l=1}^n (z_k^\top a_l^\top \nabla^2 f) \left( \sum_{i=1}^n \sum_{k'=1}^{n+m} z_{k k'}^\top \lambda_l^{k'i} a_i^\top \mathbf{U} \right) + \sum_{k,l=1}^m (z_k^\top z_l^\top \nabla^2 f) \left( \sum_{i=1}^n \sum_{k'=1}^{n+m} z_{k k'}^\top \lambda_{l+n}^{k'i} a_i^\top \mathbf{U} \right). \end{aligned}$$

If Assumption 1 holds, we get  $z_k^\top \nabla a_{\hat{i}\hat{i}}^\top = \sum_{k'=1}^{n+m} \sum_{l=1}^n z_{k k'}^\top \lambda_l^{k'i} a_{l\hat{i}}^\top$ , which means  $\lambda_{l+n}^{k'i} = 0$ , for  $l = 1, \dots, m$ . Thus we conclude the result by the following claim.



**Claim 2:**

$$\begin{aligned} [C + F + G^T + \beta V^a]X &= \sum_{i,k=1}^n (a_i^T a_k^T \nabla^2 f) \tilde{\Lambda}_1(U)_{ik} + \sum_{i=1}^n \sum_{k=1}^m (a_i^T z_k^T \nabla^2 f) \tilde{\Lambda}_2(U)_{ik} \\ &= \tilde{\Lambda}_1(U)^T QX + \tilde{\Lambda}_2(U)^T PX, \end{aligned}$$

where

$$\begin{aligned} \tilde{\Lambda}_1(U) &= \tilde{C}_1(U) + \tilde{F}_1(U) + \tilde{G}_1(U) + \beta \tilde{V}_1^a(U) \in \mathbb{R}^{n^2 \times 1}, \\ \tilde{\Lambda}_2(U) &= \tilde{C}_2(U) + \tilde{F}_2(U) + \tilde{G}_2(U) + \beta \tilde{V}_2^a(U) \in \mathbb{R}^{(n \times m) \times 1}, \end{aligned}$$

and

$$\begin{aligned} \tilde{F}_1(U)_{il} &= \sum_{k=1}^m \left( \sum_{i'=1}^{n+m} a_{ii'}^T \omega_l^{i'k} - \sum_{k'=1}^{n+m} z_{kk'}^T \lambda_l^{k'i} \right) z_k^T U, \\ \tilde{F}_2(U)_{il} &= \sum_{k=1}^m \left( \sum_{i'=1}^{n+m} a_{ii'}^T \omega_{l+n}^{i'k} - \sum_{k'=1}^{n+m} z_{kk'}^T \lambda_{l+n}^{k'i} \right) z_k^T U, \quad \tilde{G}_1(U)_{il} = - \sum_{k=1}^m \sum_{i'=1}^{n+m} a_{ii'}^T \omega_l^{i'k} z_k^T U, \\ \tilde{G}_2(U)_{lk} &= \sum_{i=1}^n \sum_{k'=1}^{n+m} z_{kk'}^T \lambda_l^{k'i} a_i^T U + z_k^T \nabla a_l^T U - \sum_{i=1}^m \sum_{i'=1}^{n+m} a_{li'}^T \omega_{k+n}^{i'i} z_i^T U - a_l^T \nabla z_k^T U, \\ \tilde{V}_1^a(U)_{il} &= -\frac{1}{2} \alpha_l (a_i^T U) + \frac{1}{2} \langle U, \gamma \rangle \mathbf{1}_{\{i=l\}}, \quad \tilde{V}_2^a(U)_{il} = -\frac{1}{2} \alpha_{l+n} (a_i^T U). \end{aligned}$$

We thus have

$$\begin{aligned} & (QX + D)^T (QX + D) + (PX + E)^T (PX + E) + 2C^T X + 2F^T X + 2G^T X + 2V_a^T X \\ &= [QX + \tilde{\Lambda}_1(U) + D]^T [QX + \tilde{\Lambda}_1(U) + D] + [PX + \tilde{\Lambda}_2(U) + E]^T [PX + \tilde{\Lambda}_2(U) + E] \\ &\quad - [\tilde{\Lambda}_1(U) + D]^T [\tilde{\Lambda}_1(U) + D] - [\tilde{\Lambda}_2(U) + E]^T [\tilde{\Lambda}_2(U) + E] + D^T D + E^T E \\ &= [QX + \Lambda_1]^T [QX + \Lambda_1] + [PX + \Lambda_2]^T [PX + \Lambda_2] - \Lambda_1^T \Lambda_1 - \Lambda_2^T \Lambda_2 + D^T D + E^T E, \end{aligned}$$

where

$$\Lambda_1 = \tilde{\Lambda}_1 + D \in \mathbb{R}^{n^2 \times 1}, \quad \Lambda_2 = \tilde{\Lambda}_2 + E \in \mathbb{R}^{(n \times m) \times 1}.$$

**Step C:** Given the condition (26) in **Step B**, we complete the proof by

$$\begin{aligned} & \int \left[ \tilde{\Gamma}_2(f, f) + \tilde{\Gamma}_2^{z, \pi}(f, f) + \Gamma_{\mathcal{I}_{a,z}}(f, f) \right] p dx \\ &= \int \left[ [QX + \Lambda_1]^T [QX + \Lambda_1] + [PX + \Lambda_2]^T [PX + \Lambda_2] + \mathfrak{R}(\nabla f, \nabla f) \right] p dx. \end{aligned}$$

■

**5.6. Proof of Step 3: entropy dissipation.** We last prove Step 3. It is a modified Lyapunov method in probability density space, which is a standard approach; see also [6, 23].

We first derive the following dissipation result.

**Proposition 10.** *Along with Fokker-Planck equation (2), the following equality holds*

$$\frac{d}{dt}D_{\text{KL}}(p\|\pi) = -\mathcal{I}_a(p\|\pi).$$

**Proof** The proof is based on a direct calculation. Notice that

$$\begin{aligned} \frac{d}{dt}D_{\text{KL}}(p\|\pi) &= \int \partial_t p \log \frac{p}{\pi} + p \partial_t \log \frac{p}{\pi} dx \\ &= \int \nabla \cdot (p\gamma) \log \frac{p}{\pi} + \nabla \cdot (paa^\top \nabla \log \frac{p}{\pi}) dx + \int p \frac{\partial_t p}{p} dx \\ &= - \int \langle \nabla \log \frac{p}{\pi}, \gamma \rangle p dx - \int \langle \nabla \log \frac{p}{\pi}, aa^\top \nabla \log \frac{p}{\pi} \rangle p dx. \end{aligned}$$

And

$$\begin{aligned} - \int \langle \nabla \log \frac{p}{\pi}, \gamma \rangle p dx &= \int -\langle \nabla p, \gamma \rangle + \langle \nabla \log \pi, \gamma \rangle dx \\ &= \int (\nabla \cdot \gamma + \langle \nabla \log \pi, \gamma \rangle) p dx \\ &= \int \frac{1}{\pi} \nabla \cdot (\pi \gamma) p dx \\ &= 0. \end{aligned}$$

Combining the above facts, we finish the proof.

We next show the following facts to conclude all results.

**Proof** [Proof of Theorem 1 and Corollary 2] From the information Gamma calculus and condition (7), we know that

$$\frac{d}{dt}\mathcal{I}_{a,z}(p\|\pi) \leq -2\lambda\mathcal{I}_{a,z}(p\|\pi).$$

From the Gronwall's inequality, we have

$$\mathcal{I}_{a,z}(p\|\pi) \leq e^{-2\lambda t}\mathcal{I}_{a,z}(p_0\|\pi).$$

This finishes the proof of Corollary 1. We next prove Corollary 2. Notice that

$$\begin{aligned} -D_{\text{KL}}(p\|\pi) &= \int_t^\infty \frac{d}{ds}D_{\text{KL}}(p_s\|\pi) ds \\ &= - \int_t^\infty \mathcal{I}_a(p_s\|\pi) ds \\ &\geq - \int_t^\infty \mathcal{I}_{a,z}(p_s\|\pi) ds \\ &\geq \frac{1}{2\lambda} \int_t^\infty \frac{d}{ds}\mathcal{I}_{a,z}(p_s\|\pi) ds \\ &= - \frac{1}{2\lambda}\mathcal{I}_{a,z}(p\|\pi), \end{aligned}$$

where we use the fact that  $p_\infty = \pi$ . Hence we prove the log-Sobolev inequality for  $aa^\top + zz^\top$  with a bound given by a constant  $\lambda$ . Using this log-Sobolev inequality for  $aa^\top + zz^\top$ , one

can prove (i), (ii) directly. Notice that

$$D_{\text{KL}}(p\|\pi) \leq \frac{1}{2\lambda} \mathcal{I}_{a,z}(p\|\pi) \leq \frac{1}{2\lambda} e^{-2\lambda t} \mathcal{I}_{a,z}(p_0\|\pi).$$

Following an inequality between KL divergence and  $L_1$  distance, i.e.,

$$\int \|p(t, x) - \pi(x)\| dx \leq \sqrt{2D_{\text{KL}}(p\|\pi)},$$

we can show the exponential decay of the solution in term of  $L_1$  distances. This follows from the exponential decay of KL divergence. We finish the proof.

#### REFERENCES

- [1] L. Ambrosio, N. Gigli and G. Savaré. *Gradient Flows in Metric Spaces and in the Space of Probability Measures*, Lectures in Mathematics ETH Zurich, Birkhauser Verlag, Basel, 2nd ed. 2008.
- [2] S. Armstrong and J. Mourrat. Variational methods for the kinetic Fokker-Planck equation. *arXiv:1902.04037*, 2019.
- [3] A. Arnold, P. Markowich, G. Toscani, A. Unterreiter. On convex Sobolev inequalities and the rate of convergence to equilibrium for Fokker-Planck type equations. *Communications in Partial Differential Equations*, 26, 43–100, 2001.
- [4] A. Arnold, E. Carlen. A generalized Bakry-Émery condition for non-symmetric diffusions. *EQUADIFF 99 Proceedings of the International Conference on Differential Equations, Berlin 1999, World Scientific Publishing*, 732-734, 2000.
- [5] A. Arnold, E. Carlen, Q. Ju. Large-time behavior of non-symmetric Fokker-Planck type equations. *Communications on Stochastic Analysis*, 4-1, 2008.
- [6] A. Arnold and J. Erb. Sharp entropy decay for hypocoercive and non-symmetric Fokker-Planck equations with linear drift. *arXiv preprint arXiv:1409.5425*, 2014.
- [7] D. Bakry and M. Émery. Diffusions hypercontractives. *Séminaire de probabilités de Strasbourg*, 19:177–206, 1985.
- [8] D. Bakry, P. Cattiaux and A. Guillin. Rate of convergence for ergodic continuous Markov processes: Lyapunov versus Poincaré. *Journal of Functional Analysis*, 254(3), 727-759, 2008.
- [9] D. Bakry, I. Gentil and M. Ledoux. Analysis and geometry of Markov diffusion operators, Vol 348, 2013. *Springer Science & Business Media*,
- [10] F. Baudoin. Bakry-Émery meet Villani. *Journal of functional analysis*, 273(7), 2275-2291, 2017.
- [11] F. Baudoin and N. Garofalo. Curvature-dimension inequalities and Ricci lower bounds for sub-Riemannian manifolds with transverse symmetries. *Journal of the EMS, Vol. 19, Issue 1*, 2017.
- [12] F. Baudoin, M. Gordina, and D. P. Herzog. Gamma calculus beyond Villani and explicit convergence estimates for Langevin dynamics with singular potentials. *arXiv:1907.03092*, 2019.
- [13] E. Bernard, M. Fathi, A. Levitt, and G. Stoltz. Hypocoercivity with Schur complements. *arXiv preprint arXiv:2003.00726*, 2020.
- [14] J. M. Bismut. Martingales, the Malliavin calculus and hypoellipticity under general Hörmander’s conditions. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 56, 4, 469-505, 1981, Springer.
- [15] Y. Cao, J. Lu, and L. Wang. On explicit L2-convergence rate estimate for underdamped Langevin dynamics. *arXiv:1908.04746*, 2019.
- [16] E. Carlen. Superadditivity of Fisher’s information and logarithmic Sobolev inequalities. *Journal of Functional Analysis*, vol. 101, no. 1, pp. 194–211, 1991.
- [17] X. Cheng, N. Chatterji, P. Bartlett, and M. Jordan. Underdamped Langevin MCMC: A non-asymptotic analysis. *Proceedings of the 31st Conference On Learning Theory*, PMLR 75:300-323, 2018.
- [18] R. Duan. Hypocoercivity of linear degenerately dissipative kinetic equations. *Nonlinearity*, Volume 24, Number 8, 2011.
- [19] A.B. Duncan, G.A. Pavliotis, and K.C. Zygalakis. Nonreversible Langevin Samplers: Splitting Schemes, Analysis and Implementation. *arXiv: 1701.04247*, 2017.

- [20] J.P. Eckmann and M. Hairer. Spectral properties of hypoelliptic operators. *Comm. Math. Phys.*, 235(2):233–253, 2003.
- [21] J.-P. Eckmann, C.-A. Pillet and L. Rey-Bellet. Non-Equilibrium Statistical Mechanics of Anharmonic Chains Coupled to Two Heat Baths at Different Temperatures. *Comm. Math. Phys.*, 201, 657–697, 1999.
- [22] Q. Feng and W. Li. Generalized Gamma  $z$  calculus via sub-Riemannian density manifold. *arXiv:1910.07480*, 2020.
- [23] Q. Feng and W. Li. Entropy dissipation via Information Gamma calculus: Non-reversible stochastic differential equations. *arXiv:1910.07480*, 2020.
- [24] Q. Feng and W. Li. Sub-Riemannian Ricci curvature via generalized Gamma  $z$  calculus. *arXiv:2004.01863*, 2020.
- [25] L. Gross. Logarithmic Sobolev inequalities. *American Journal of Mathematics*, 97(4), 1061–1083, 1975.
- [26] M. Hairer and J. Mattingly. Slow energy dissipation in anharmonic oscillator chains. *Communications on Pure and Applied Mathematics*, Vol. LXII, 0999–1032, 2009.
- [27] F. Hérau and F. Nier. Isotropic hypoellipticity and trend to equilibrium for the Fokker-Planck equation with a high-degree potential. *Arch. Ration. Mech. Anal.*, 171(2):151–218, 2004.
- [28] L. Hörmander. Hypoelliptic second order differential equations. *Acta Math.*, 119, (1967), 147–171.
- [29] A. Iacubucci, S. Olla, and G. Stoltz. Convergence rates for nonequilibrium Langevin dynamics. *Annales mathématiques du Québec*, 43, 1, 73–98, 2019.
- [30] S. Gadat and L. Miclo. Spectral decompositions and  $L^2$ -operator norms of toy hypocoercive semigroups, *Kinetic and Related Models*, 2, 317–372, 2013.
- [31] J. D. Lafferty. The density manifold and configuration space quantization. *Transactions of the American Mathematical Society*, 305(2):699–741, 1988.
- [32] F. Ledrappier and L.S. Young. Entropy Formula for Random Transformations. *Probab. Th. Rel. Fields*, 80, 217–240, 1988.
- [33] B. Leimkuhler and M. Sachs. Efficient Numerical Algorithms for the Generalized Langevin Equation. *arXiv preprint: arXiv:2012.04245*.
- [34] W. Li. Transport information geometry: Riemannian calculus on probability simplex. *arXiv:1803.06360*, 2018.
- [35] P. A. Markowich, C. Villani. On The Trend To Equilibrium For The Fokker-Planck Equation: An Interplay Between Physics And Functional Analysis. *Physics and Functional Analysis*, Matematica Contemporanea, 1999.
- [36] J. C. Mattingly, A.M. Stuart, D.J. Higham. Ergodicity for SDEs and approximations: locally Lipschitz vector fields and degenerate noise. *Stochastic Processes and their Applications*, Volume 101, Issue 2, Pages 185–232, 2002.
- [37] A. Menegaki. Quantitative Rates of Convergence to Non-equilibrium Steady State for a Weakly Anharmonic Chain of Oscillators. *Journal of Statistical Physics*, vol. 181, no. 1, Oct., pp. 53+, 2020.
- [38] P. Monmarché. Almost sure contraction for diffusions on  $\mathbb{R}^d$ . Application to generalised Langevin diffusions. *arXiv:2009.10828v4*, 2020.
- [39] C. Mouhot, and L. Neumann. Quantitative perturbative study of convergence to equilibrium for collisional kinetic models in the torus. *Nonlinearity*, 969–998, 19, 2006.
- [40] E. Nelson. *Quantum Fluctuations*. Princeton series in physics. Princeton University Press, Princeton, N.J, 1985.
- [41] F. Otto and C. Villani. Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality. *Journal of Functional Analysis*, 173 (2): 361–400, 2000.
- [42] G. A. Pavliotis. Stochastic processes and applications: diffusion processes, the Fokker-Planck and Langevin equations. *Springer, New York*, Vol. 60, 2014.
- [43] L. Rey-Bellet. Ergodic properties of Markov processes. Open quantum systems. II, *Lecture Notes in Math.*, Vol. 1881, Springer, Berlin, 2006, pp. 1–39.
- [44] L. Rey-Bellet, and L.E. Thomas. Asymptotic behavior of thermal nonequilibrium steady states for a driven chain of anharmonic oscillators. *Comm. Math. Phys.*, 215(1):1–24, 2000.
- [45] M. Sachs, B. Leimkuhler, and V. Danos. Langevin Dynamics with Variable Coefficients and Nonconservative Forces: From Stationary States to Numerical Methods. *Entropy*, 19(12), 647, 2017.

- [46] D. Talay. Stochastic Hamiltonian systems: exponential convergence to the invariant measure, and discretization by the implicit Euler scheme. *Markov Process and Related Fields*, 8(2):163–198. 2002.
- [47] L. Vandenberghe and S. Boyd. Convex optimization. *Volume 1. Cambridge University Press Cambridge*, 2004.
- [48] C. Villani. *Hypocoercivity*, Memoirs of the American Mathematical Society, 2009.
- [49] C. Villani. *Optimal Transport: Old and New*, 2009.

## APPENDIX: PROOF OF EXAMPLES.

In appendix, we provide the detailed proofs for all examples.

**Proof of Example I: underdamped Langevin dynamics.** For underdamped Langevin dynamics with variable coefficients defined in (10), we first observe that

$$aa^\top = \begin{pmatrix} 0 & 0 \\ 0 & r(x) \end{pmatrix}, \quad \left( \sum_{j=1}^2 \frac{\partial}{\partial x_j} (aa^\top)_{ij} \right)_{1 \leq i \leq 2} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad aa^\top \nabla \log \pi = \begin{pmatrix} 0 \\ (a_{12}^\top)^2 \frac{\partial \log \pi}{\partial v} \end{pmatrix}.$$

By routine computations, we have the following proposition.

**Proposition 11.** *For any  $\beta \in \mathbb{R}$ , and any smooth function  $f \in C^\infty(\mathbb{R}^2)$ , we have*

$$\begin{aligned} & \int \left[ \tilde{\Gamma}_2(f, f) + \tilde{\Gamma}_2^{z, \pi}(f, f) + \Gamma_{\mathcal{I}_{a, z}}(f, f) \right] p dx \\ &= \int \left[ \|\mathfrak{H}\text{ess}_\beta f\|_{\mathbb{F}}^2 + \mathfrak{R}(\nabla f, \nabla f) + \mathfrak{R}_{\mathcal{I}}(\nabla f, \nabla f) \right] p dx, \end{aligned}$$

where

$$\|\mathfrak{H}\text{ess}_\beta f\|_{\mathbb{F}}^2 = [\mathbf{X} + \Lambda]^\top (\mathbf{Q}^\top \mathbf{Q} + \mathbf{P}^\top \mathbf{P}) [\mathbf{X} + \Lambda],$$

with

$$\begin{aligned} \Lambda &= (0, 0, \Lambda_3, \Lambda_4), \quad \mathbf{Q} = \begin{pmatrix} 0 & 0 & 0 & (a_{12}^\top)^2 \end{pmatrix}, \quad \mathbf{P} = \begin{pmatrix} 0 & 0 & a_{12}^\top z_1 & a_{12}^\top z_2 \end{pmatrix}, \\ \begin{pmatrix} \Lambda_3 \\ \Lambda_4 \end{pmatrix} &= \frac{1}{(a_{12}^\top)^2} (aa^\top + zz^\top)^{-1} \mathbf{K} \nabla f. \end{aligned}$$

And the curvature tensor  $\mathcal{R}$  is presented in Theorem 2.

*Remark 11.* Here we use a different notion of the Hessian of function  $f$ , which gives us the other equivalent formulation of the tensor  $\mathfrak{R}$ . The key observation is the following relation

$$\begin{aligned} & [\mathbf{Q}\mathbf{X} + \hat{\Lambda}_1]^\top [\mathbf{Q}\mathbf{X} + \hat{\Lambda}_1] + [\mathbf{P}\mathbf{X} + \hat{\Lambda}_2]^\top [\mathbf{P}\mathbf{X} + \hat{\Lambda}_2] - \hat{\Lambda}_1^\top \hat{\Lambda}_1 - \hat{\Lambda}_2^\top \hat{\Lambda}_2 \\ &= [\mathbf{X} + \Lambda_1]^\top \mathbf{Q}^\top \mathbf{Q} [\mathbf{X} + \Lambda_1] + [\mathbf{X} + \Lambda_2]^\top \mathbf{P}^\top \mathbf{P} [\mathbf{X} + \Lambda_2] - \Lambda_1^\top \mathbf{Q}^\top \mathbf{Q} \Lambda_1 - \Lambda_2^\top \mathbf{P}^\top \mathbf{P} \Lambda_2, \end{aligned}$$

where  $\hat{\Lambda}_1^\top = \Lambda_1^\top \mathbf{Q}^\top$ ,  $\hat{\Lambda}_2^\top = \Lambda_2^\top \mathbf{P}^\top$ . Shortly, we show that the vectors  $\Lambda_1 \in \mathbb{R}^{(n+m)^2 \times 1}$  and  $\Lambda_2 \in \mathbb{R}^{(n+m)^2 \times 1}$  exist. And we compute them explicitly for the variable coefficient underdamped Langevin dynamics.

**Proof** [Proof of Proposition 11] We demonstrate explicit examples with constant matrix  $z$ , i.e.  $z^\top = (z_1 \ z_2)$ , with  $z_1$  and  $z_2$  being constants. In particular, we have used the notation below  $z_1^\top = (z_{11}^\top \ z_{12}^\top) = (z_1 \ z_2)$ , and  $\partial_1 f = \partial_x f$ ,  $\partial_2 f = \partial_v f$ .

**Step 1:** We first have the following simplified quantities:

$$\begin{aligned}
\mathfrak{R}_a(\nabla f, \nabla f) &= -(a_{12}^\top)^2 \frac{\partial^2 \log \pi}{\partial v^2} |a_{12}^\top \partial_2 f|^2, \quad \mathfrak{R}_z(\nabla f, \nabla f) = -z_1^\top \nabla \left( (a_{12}^\top)^2 \frac{\partial \log \pi}{\partial v} \right) \partial_2 f z_1^\top \nabla f, \\
\mathfrak{R}_\pi(\nabla f, \nabla f) &= 2 \left[ z_1^\top z_1^\top \nabla^2 a_1^\top \nabla f a_1^\top \nabla f + (z_1^\top \nabla a_1^\top \nabla f)^2 + (z^\top \nabla \log \pi)_1 \left[ z_1^\top \nabla a_1^\top \nabla f a_1^\top \nabla f \right] \right], \\
\mathfrak{R}_{\mathcal{I}_a}(\nabla f, \nabla f) &= \langle \nabla f, \gamma \rangle \langle a a^\top \nabla \log \pi, \nabla f \rangle - \gamma_1 \frac{\partial}{\partial x} a_{12}^\top \partial_2 f a_{12}^\top \partial_2 f, \\
\mathfrak{R}_{\gamma_a}(\nabla f, \nabla f) &= \frac{1}{2} \sum_{k=1}^2 \gamma_k \langle \nabla f, \nabla_k (a a^\top) \nabla f \rangle - \left\langle \begin{pmatrix} \nabla_1 \gamma_1 & \nabla_1 \gamma_2 \\ \nabla_2 \gamma_1 & \nabla_2 \gamma_2 \end{pmatrix} \begin{pmatrix} \partial_1 f \\ \partial_2 f \end{pmatrix}, a a^\top \nabla f \right\rangle, \\
\mathfrak{R}_{\gamma_z}(\nabla f, \nabla f) &= - \left\langle \begin{pmatrix} \nabla_1 \gamma_1 & \nabla_1 \gamma_2 \\ \nabla_2 \gamma_1 & \nabla_2 \gamma_2 \end{pmatrix} \begin{pmatrix} \partial_1 f \\ \partial_2 f \end{pmatrix}, z z^\top \nabla f \right\rangle.
\end{aligned}$$

And

$$\begin{aligned}
\mathbf{E} &= \mathbf{D} = 0, \quad \mathbf{C} = (0, 0, 0, 0)^\top, \quad \mathbf{F} = \left( 0, 0, 0, \mathbf{F}_{22} \right)^\top, \quad \mathbf{F}_{22} = -z_{11}^\top \partial_x a_{12}^\top a_{12}^\top z_1^\top \nabla f, \\
\mathbf{G} &= \left( 0, 0, \mathbf{G}_{21}, \mathbf{G}_{22} \right)^\top, \quad \mathbf{G}_{21} = 2(z_{11}^\top)^2 \partial_x a_{12}^\top a_{12}^\top \partial_2 f, \quad \mathbf{G}_{22} = 2z_{11}^\top z_{12}^\top \partial_x a_{12}^\top a_{12}^\top \partial_2 f, \\
\mathbf{V}^a &= (0, 0, \mathbf{V}_{21}^a, \mathbf{V}_{22}^a), \quad \mathbf{V}_{21}^a = -\frac{1}{2} \gamma_1 a_{12}^\top a_{12}^\top \partial_2 f, \quad \mathbf{V}_{22}^a = -\frac{1}{2} \gamma_2 a_{12}^\top a_{12}^\top \partial_2 f + \frac{1}{2} a_{12}^\top a_{12}^\top \langle \nabla f, \gamma \rangle.
\end{aligned}$$

**Step 2:** We now find the vector  $\Lambda$  in Remark 11. By Definition 1, we have

$$P^\top P = (a_{12}^\top)^2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & z_1^2 & z_1 z_2 \\ 0 & 0 & z_1 z_2 & z_2^2 \end{pmatrix}, \quad \text{and} \quad Q^\top Q = (a_{12}^\top)^2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & (a_{12}^\top)^2 \end{pmatrix},$$

and  $Q^\top D = (0, 0, 0, 0)^\top$ ,  $P^\top E = (0, 0, z_{11}^\top a_{12}^\top \mathbf{E}, z_{12}^\top a_{12}^\top \mathbf{E})^\top$ . For simplicity, taking  $\Lambda_1 = \Lambda_2 = \Lambda = (0, 0, \Lambda_3, \Lambda_4)$ , we have

$$(Q^\top Q \Lambda + P^\top P \Lambda)^\top X = (F + C + G + \beta V^a + Q^\top D + P^\top E)^\top X.$$

By matching the coefficients of  $\frac{\partial^2 f}{\partial x \partial v}$  and  $\frac{\partial^2 f}{\partial v \partial v}$  for both sides of the above equation, we have

$$(a_{12}^\top)^2 \begin{pmatrix} z_1^2 & z_1 z_2 \\ z_1 z_2 & z_2^2 + (a_{12}^\top)^2 \end{pmatrix} \begin{pmatrix} \Lambda_3 \\ \Lambda_4 \end{pmatrix} = \begin{pmatrix} \mathbf{K}_1 \\ \mathbf{K}_2 \end{pmatrix},$$

with

$$\mathbf{K}_1 =: \mathbf{F}_{21} + \mathbf{G}_{21} + \beta \mathbf{V}_{21}^a, \quad \mathbf{K}_2 =: \mathbf{F}_{22} + \mathbf{G}_{22} + \beta \mathbf{V}_{22}^a.$$

Notice that

$$\begin{aligned}
\mathbf{K}_1 &= \left( 2(z_{11}^\top)^2 \partial_x a_{12}^\top a_{12}^\top - \frac{\beta}{2} \gamma_1 a_{12}^\top a_{12}^\top \right) \partial_2 f := \mathbf{K}_{12} \partial_2 f, \\
\mathbf{K}_2 &= [-z_1^2 \partial_x [a_{12}^\top] a_{12}^\top + \frac{\beta}{2} (a_{12}^\top)^2 \gamma_1] \partial_1 f + z_1 z_2 \partial_x [a_{12}^\top] a_{12}^\top \partial_2 f := \mathbf{K}_{21} \partial_1 f + \mathbf{K}_{22} \partial_2 f.
\end{aligned}$$

Based on the above computation, we have

$$\begin{pmatrix} \Lambda_3 \\ \Lambda_4 \end{pmatrix} = \frac{1}{(a_{12}^\top)^2} \begin{pmatrix} z_1^2 & z_1 z_2 \\ z_1 z_2 & z_2^2 + (a_{12}^\top)^2 \end{pmatrix}^{-1} \begin{pmatrix} \mathbf{K}_{11} & \mathbf{K}_{12} \\ \mathbf{K}_{21} & \mathbf{K}_{22} \end{pmatrix} \nabla f = \frac{1}{(a_{12}^\top)^2} (a a^\top + z z^\top)^{-1} \mathbf{K} \nabla f.$$

Since  $aa^\top + zz^\top$  is symmetric, we have the following relation

$$\Lambda^\top(Q^\top Q + P^\top P)\Lambda = (\nabla f)^\top \mathfrak{M}_\Lambda \nabla f,$$

with the matrix  $\mathfrak{M}_\Lambda$  defined by

$$\mathfrak{M}_\Lambda = \frac{1}{(a_{12}^\top)^2} (\mathbf{K})^\top (aa^\top + zz^\top)^{-1} \mathbf{K}.$$

**Step 3:** We next transfer all the bilinear forms into its corresponding symmetric matrix forms. In particular, for each bi-linear form  $\mathfrak{R}(\nabla f, \nabla f)$ , we keep the convention

$$\mathfrak{R}(\nabla f, \nabla f) = (\nabla f)^\top \mathfrak{R} \nabla f.$$

We focus on the symmetric matrix form  $\mathfrak{R}$ :

$$\mathfrak{R}_a(\nabla f, \nabla f) = -(a_{12}^\top)^2 \frac{\partial^2 \log \pi}{\partial v^2} |a_{12}^\top \partial_2 f|^2 = (\nabla f)^\top \begin{pmatrix} 0 & 0 \\ 0 & -\frac{\partial^2 \log \pi}{\partial v^2} |a_{12}^\top|^4 \end{pmatrix} \nabla f,$$

$$\begin{aligned} \mathfrak{R}_z(\nabla f, \nabla f) &= -z_1^\top \nabla \left( (a_{12}^\top)^2 \frac{\partial \log \pi}{\partial v} \right) \partial_2 f z_1^\top \nabla f \\ &= \frac{1}{2} (\nabla f)^\top \left[ \begin{pmatrix} 0 & 0 \\ -z_1^\top \nabla \left( (a_{12}^\top)^2 \frac{\partial \log \pi}{\partial v} \right) & z_1^\top \nabla \left( (a_{12}^\top)^2 \frac{\partial \log \pi}{\partial v} \right) \end{pmatrix} \right] \nabla f, \end{aligned}$$

$$\begin{aligned} \mathfrak{R}_\pi(\nabla f, \nabla f) &= 2 \left[ z_1^\top z_1^\top \nabla^2 a_1^\top \nabla f a_1^\top \nabla f + (z_1^\top \nabla a_1^\top \nabla f)^2 + (z^\top \nabla \log \pi)_1 \left[ z_1^\top \nabla a_1^\top \nabla f a_1^\top \nabla f \right] \right] \\ &= (\nabla f)^\top \begin{pmatrix} 0 & 0 \\ 0 & C_\pi \end{pmatrix} \nabla f, \end{aligned}$$

with

$$C_\pi = 2 \left[ z_1^\top z_1^\top \nabla^2 a_{12}^\top a_{12}^\top + (z_1^\top \nabla a_{12}^\top)^2 + (z^\top \nabla \log \pi)_1 [z_1^\top \nabla a_{12}^\top a_{12}^\top] \right]. \quad (28)$$

We also have

$$\begin{aligned} \mathfrak{R}_{\mathcal{I}_a}(\nabla f, \nabla f) &= \langle \nabla f, \gamma \rangle \langle aa^\top \nabla \log \pi, \nabla f \rangle - \gamma_1 \frac{\partial}{\partial x} a_{12}^\top \partial_2 f a_{12}^\top \partial_2 f \\ &= (\nabla f)^\top \left[ \frac{\gamma (aa^\top \nabla \log \pi)^\top + (aa^\top \nabla \log \pi) \gamma^\top}{2} - \begin{pmatrix} 0 & 0 \\ 0 & \gamma_1 \frac{\partial}{\partial x} a_{12}^\top a_{12}^\top \end{pmatrix} \right] \nabla f, \end{aligned}$$

and

$$\begin{aligned} \mathfrak{R}_{\gamma_a}(\nabla f, \nabla f) &= \frac{1}{2} \langle \nabla f, \sum_{k=1}^2 \gamma_k \nabla_k (aa^\top) \nabla f \rangle - \langle \nabla \gamma \nabla f, aa^\top \nabla f \rangle \\ &= (\nabla f)^\top \left[ \frac{1}{2} \sum_{k=1}^2 \gamma_k \nabla_k (aa^\top) - \frac{1}{2} [(\nabla \gamma)^\top aa^\top + aa^\top \nabla \gamma] \right] \nabla f. \end{aligned}$$

The last equality follows from the fact that  $aa^\top$  is a diagonal matrix. Hence  $\sum_{k=1}^2 \gamma_k \nabla_k(aa^\top)$  is also a diagonal matrix. Similarly, we derive

$$\begin{aligned} \mathfrak{R}_{\gamma_z}(\nabla f, \nabla f) &= -\left\langle \begin{pmatrix} \nabla_1 \gamma_1 & \nabla_1 \gamma_2 \\ \nabla_2 \gamma_1 & \nabla_2 \gamma_2 \end{pmatrix} \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}, zz^\top \nabla f \right\rangle, \\ &= (\nabla f)^\top \frac{1}{2} \left[ -(\nabla \gamma)^\top zz^\top - zz^\top \nabla \gamma \right] \nabla f. \end{aligned}$$

The proof is completed. ■

### Proof of example II: 3 oscillator chain model.

**Proof** [Proof of Proposition 4] Following Theorem 3, consider a constant matrix  $a$  and a matrix function  $z$ ,

$$z = \begin{pmatrix} z_1 \mathbf{1}_3 & \mathbf{0}_{3 \times 1} \\ z_2 \mathbf{1}_3 & (\vec{z}_3)_{3 \times 1} \end{pmatrix}_{6 \times 4},$$

with  $\vec{z}_3 = (z_{31} \ z_{32}(p_0, p_2) \ z_{33})^\top$  and constants  $z_1, z_2$ . We compute the matrix tensor  $\mathfrak{R}$  for  $\beta = 0$ . By abusing the notations, we denote  $z_i^\top$  as the  $i$ -th row vector of the transpose matrix  $z^\top$ . We first find vectors  $\Lambda_1$  and  $\Lambda_2$ . For vectors  $C, F, G, V^a \in \mathbb{R}^{(6 \times 6) \times 1}$ ,  $D \in \mathbb{R}^{(2 \times 2) \times 1}$  and  $E \in \mathbb{R}^{(2 \times 4) \times 1}$  with  $U \in \mathbb{R}^6$ , we have

$$\begin{aligned} E_{ik} &= a_i^\top \nabla z_k^\top U, \quad E_{14,1} = a_{14}^\top \partial_{v_0} z_{45}^\top U_5, \quad E_{24,1} = a_{26}^\top \partial_{p_2} z_{45}^\top U_5; \\ C_{\hat{ik}} &= \sum_{i,k=1}^2 \left( a_{ii}^\top a_i^\top \nabla a_{kk}^\top - a_{ik}^\top a_k^\top \nabla a_{ii}^\top \right) a_k^\top U = 0, \quad D_{ik} = a_i^\top \nabla a_k^\top U = 0; \\ F_{\hat{ik}} &= \sum_{i=1}^2 \sum_{k=1}^4 \left( a_{ii}^\top a_i^\top \nabla z_{kk}^\top \right) z_k^\top U, \quad F_{45,1} = |a_{14}^\top|^2 \partial_{p_0} z_{45}^\top z_4^\top U, \quad F_{65,1} = |a_{26}^\top|^2 \partial_{p_2} z_{45}^\top z_4^\top U; \\ G_{\hat{ik}} &= \sum_{i=1}^2 \sum_{k=1}^4 \left[ - \left( a_{ii}^\top a_i^\top \nabla z_{kk}^\top z_k^\top U + z_{kk}^\top a_{ii}^\top a_i^\top \nabla z_k^\top U \right) \right], \\ G_{45,1} &= -|a_{14}^\top|^2 \partial_{p_0} z_{45}^\top z_4^\top U - |a_{14}^\top|^2 \partial_{p_0} z_{45}^\top z_{45}^\top U_5, \\ G_{65,1} &= -|a_{26}^\top|^2 \partial_{p_2} z_{45}^\top z_4^\top U - |a_{26}^\top|^2 z_{45}^\top \partial_{p_2} z_{45}^\top U_5; \\ (P^\top E)_{45,1} &= a_{14}^\top z_{45}^\top E_{14,1} = |a_{14}^\top|^2 z_{45}^\top \partial_{p_0} z_{45}^\top U_5, \quad (P^\top E)_{65,1} = a_{26}^\top z_{45}^\top E_{24,1} = |a_{26}^\top|^2 z_{45}^\top \partial_{p_2} z_{45}^\top U_5. \end{aligned}$$

By a direct computation, we observe that  $F + G + C + Q^\top D + P^\top E = \vec{0}_{36 \times 1}$ . We thus get  $\Lambda_1 = \Lambda_2 = \mathbf{0} \in \mathbb{R}^6$ . We next compute the following matrix tensors. For simplicity, we directly omit the zero terms based on Definition 1 and matrix functions  $a, z$ .

$$\begin{aligned} \mathfrak{R}_a(U, U) &= - \sum_{i=1}^2 \sum_{\hat{k}=1}^6 \left[ a_i^\top \nabla (aa^\top \nabla \log \pi)_{\hat{k}} U_{\hat{k}} \right] a_i^\top U \\ &= -a_{14}^\top \partial_{p_0} (|a_{14}^\top|^2 \partial_{p_0} \log \pi) U_4 a_{14}^\top U_4 - a_{26}^\top \partial_{p_2} (|a_{26}^\top|^2 \partial_{p_2} \log \pi) U_6 a_{26}^\top U_6 \\ &= U^\top \begin{pmatrix} 0 & 0 \\ 0 & (\xi T)^2 \mathbf{1}_3 \end{pmatrix} U. \end{aligned}$$



$$\begin{aligned}
& \mathfrak{R}_z(\mathbf{U}, \mathbf{U}) \\
&= \sum_{i,k=1}^2 a_i^\top a_i^\top \nabla^2 z_k^\top \mathbf{U} (z_k^\top \mathbf{U}) \\
&+ \sum_{k=1}^4 \sum_{\hat{k}=1}^6 \left[ (aa^\top \nabla \log \pi)_{\hat{k}} \nabla_{\hat{k}} z_k^\top \mathbf{U} - z_k^\top \nabla (aa^\top \nabla \log \pi)_{\hat{k}} \mathbf{U}_{\hat{k}} \right] z_k^\top \mathbf{U} \\
&= |a_{14}^\top|^2 \partial_{p_0 p_0} z_4^\top \mathbf{U} z_4^\top \mathbf{U} + |a_{26}^\top|^2 \partial_{p_2 p_2} z_4^\top \mathbf{U} z_4^\top \mathbf{U} \\
&+ (|a_{14}^\top|^2 \partial_{p_0} \log \pi) \partial_{p_0} z_4^\top \mathbf{U} z_4^\top \mathbf{U} + (|a_{26}^\top|^2 \partial_{p_2} \log \pi) \partial_{p_2} z_4^\top \mathbf{U} z_4^\top \mathbf{U} \\
&- z_1^\top \nabla ((a_{14}^\top)^2 \partial_{p_0} \log \pi \mathbf{U}_4) z_1^\top \mathbf{U} - z_3^\top \nabla ((a_{26}^\top)^2 \partial_{p_2} \log \pi \mathbf{U}_6) z_3^\top \mathbf{U} \\
&- z_4^\top \nabla [(a_{14}^\top)^2 \partial_{p_0} \log \pi \mathbf{U}_4 + (a_{26}^\top)^2 \partial_{p_2} \log \pi \mathbf{U}_6] z_4^\top \mathbf{U} \\
&= |a_{14}^\top|^2 \partial_{p_0 p_0} z_{45}^\top \mathbf{U} z_{45}^\top \mathbf{U} + |a_{26}^\top|^2 \partial_{p_2 p_2} z_{45}^\top \mathbf{U} z_{45}^\top \mathbf{U} \\
&+ (|a_{14}^\top|^2 \partial_{p_0} \log \pi) \partial_{p_0} z_{45}^\top \mathbf{U} z_{45}^\top \mathbf{U} + (|a_{26}^\top|^2 \partial_{p_2} \log \pi) \partial_{p_2} z_{45}^\top \mathbf{U} z_{45}^\top \mathbf{U} \\
&- z_1^\top \nabla ((a_{14}^\top)^2 \partial_{p_0} \log \pi \mathbf{U}_4) z_1^\top \mathbf{U} - z_3^\top \nabla ((a_{26}^\top)^2 \partial_{p_2} \log \pi \mathbf{U}_6) z_3^\top \mathbf{U} \\
&- z_4^\top \nabla [(a_{14}^\top)^2 \partial_{p_0} \log \pi \mathbf{U}_4 + (a_{26}^\top)^2 \partial_{p_2} \log \pi \mathbf{U}_6] z_4^\top \mathbf{U} \\
&= \frac{1}{2} \mathbf{U}^\top \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -z_1^\top \nabla ((a_{14}^\top)^2 \partial_{p_0} \log \pi) & 0 & 0 & -z_4^\top \nabla ((a_{14}^\top)^2 \partial_{p_0} \log \pi) \\ 0 & 0 & 0 & \mathbf{S}_1 \\ 0 & 0 & -z_3^\top \nabla ((a_{26}^\top)^2 \partial_{p_2} \log \pi) & -z_4^\top \nabla ((a_{26}^\top)^2 \partial_{p_2} \log \pi) \end{pmatrix} z^\top \right. \\
&\quad \left. + z \begin{pmatrix} 0 & 0 & 0 & -z_1^\top \nabla ((a_{14}^\top)^2 \partial_{p_0} \log \pi) & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -z_3^\top \nabla ((a_{26}^\top)^2 \partial_{p_2} \log \pi) \\ 0 & 0 & 0 & -z_4^\top \nabla ((a_{14}^\top)^2 \partial_{p_0} \log \pi) & \mathbf{S}_1 & -z_4^\top \nabla ((a_{26}^\top)^2 \partial_{p_2} \log \pi) \end{pmatrix} \right\} \mathbf{U},
\end{aligned}$$

where we denote

$$\begin{aligned}
\mathbf{S}_1 &= |a_{14}^\top|^2 \partial_{p_0 p_0} z_{45}^\top + |a_{26}^\top|^2 \partial_{p_2 p_2} z_{45}^\top + (|a_{14}^\top|^2 \partial_{p_0} \log \pi) \partial_{p_0} z_{45}^\top + (a_{26}^\top)^2 \partial_{p_2} \log \pi \partial_{p_2} z_{45}^\top \\
&= \xi T \partial_{p_0 p_0} z_{45}^\top + \xi T \partial_{p_2 p_2} z_{45}^\top + (\xi T \partial_{p_0} \log \pi) \partial_{p_0} z_{45}^\top + (\xi T \partial_{p_2} \log \pi) \partial_{p_2} z_{45}^\top.
\end{aligned}$$

According to our definition, for  $z_{44}^\top = z_{31}$ ,  $z_{45}^\top = z_{32}(p_0, p_2)$ ,  $z_{46}^\top = z_{33}$ , we have

$$\begin{aligned}
\mathfrak{R}_z(\mathbf{U}, \mathbf{U}) &= \frac{1}{2} \mathbf{U}^\top \left[ \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ z_2 \xi & 0 & 0 & z_{31} \xi \\ 0 & 0 & 0 & \mathbf{S}_1 \\ 0 & 0 & z_2 \xi & z_{33} \xi \end{pmatrix} z^\top + z \begin{pmatrix} 0 & 0 & 0 & z_2 \xi & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & z_2 \xi \\ 0 & 0 & 0 & z_{31} \xi & \mathbf{S}_1 & z_{33} \xi \end{pmatrix} \right] \mathbf{U} \\
&= \begin{pmatrix} 0 & \frac{1}{2} z_1 z_2 \xi \mathbf{I}_3^O \\ \frac{1}{2} z_1 z_2 \xi \mathbf{I}_3^O & \frac{1}{2} (\vec{z}_3 \vec{z}_3^\top + \vec{z}_3 (\vec{z}_3^\top)) \end{pmatrix}_{6 \times 6},
\end{aligned}$$

where we denote  $\vec{z}_3^S = (z_{31}\xi \quad S_1 \quad z_{33}\xi)^T$ . Furthermore, we have  $\mathfrak{R}_{\gamma_a}(\mathbf{U}, \mathbf{U}) = -\langle \nabla\gamma\mathbf{U}, aa^T\mathbf{U} \rangle_{\mathbb{R}^6} = -\mathbf{U}^T \frac{1}{2} [(\nabla\gamma)^T aa^T + aa^T \nabla\gamma] \mathbf{U}$ . From our definition for  $(\nabla\gamma)_{ij} = \nabla_i \gamma_j$ , we have

$$\nabla\gamma = \begin{pmatrix} 0 & \nabla_{qq}H \\ -\nabla_{pp}H & 0 \end{pmatrix} = \begin{pmatrix} 0 & L \\ -I_3 & 0 \end{pmatrix}.$$

Here we denote  $\nabla_{qq}H$  ( $\nabla_{pp}H$  resp.) the Hessian in the  $q$  ( $p$  resp.) direction, i.e.  $L = (\nabla_{q_i q_j} H)_{0 \leq i, j \leq 2}$ . Plugging them in  $\nabla\gamma$ , we have

$$\mathfrak{R}_{\gamma_a} = \frac{1}{2} \begin{pmatrix} 0 & \xi T I_3^O \\ \xi T I_3^O & 0 \end{pmatrix}_{6 \times 6}.$$

Similarly,  $\mathfrak{R}_{\gamma_z}(\mathbf{U}, \mathbf{U}) = -\frac{1}{2} [(\nabla\gamma)^T z z^T + z z^T \nabla\gamma]$ . And we get

$$\mathfrak{R}_{\gamma_z} = \begin{pmatrix} \frac{1}{2}(z_2^2 I_3 + \vec{z}_3^S \vec{z}_3^{S^T} - z_1^2 L) & \\ & -z_1 z_2 L \end{pmatrix}_{6 \times 6}.$$

Lastly, we compute the matrix tensor  $\mathfrak{R}_\pi$ .

$$\begin{aligned} \mathfrak{R}_\pi(\mathbf{U}, \mathbf{U}) &= -2 \sum_{j=1}^4 \sum_{l=1}^2 \left[ a_l^T a_l^T \nabla^2 z_j^T \mathbf{U} z_j^T \mathbf{U} \right] \\ &\quad - 2 \sum_{j=1}^4 \sum_{l=1}^2 \left[ (a_l^T \nabla z_j^T \mathbf{U})^2 + (a^T \nabla \log \pi)_l \left[ a_l^T \nabla z_j^T \mathbf{U} z_j^T \mathbf{U} \right] \right], \\ &= -2|a_{14}^T|^2 \partial_{p_0 p_0} z_{45}^T \mathbf{U}_5 z_4^T \mathbf{U} - 2|a_{26}^T|^2 \partial_{p_2 p_2} z_{45}^T \mathbf{U}_5 z_4^T \mathbf{U} \\ &\quad - 2(a_{14}^T \partial_{p_0} z_{45}^T \mathbf{U}_5)^2 - 2(a_{26}^T \partial_{p_2} z_{45}^T \mathbf{U}_5)^2 \\ &\quad - 2a_{14}^T \partial_{p_0} \log \pi a_{14}^T \partial_{p_0} z_{45}^T \mathbf{U}_5 z_4^T \mathbf{U} - 2a_{26}^T \partial_{p_2} \log \pi a_{26}^T \partial_{p_2} z_{45}^T \mathbf{U}_5 z_4^T \mathbf{U} \\ &= -2\xi T [\partial_{p_0 p_0} z_{45}^T + \partial_{p_2 p_2} z_{45}^T + \partial_{p_0} \log \pi \partial_{p_0} z_{45}^T + \partial_{p_2} \log \pi \partial_{p_2} z_{45}^T] \mathbf{U}_5 z_4^T \mathbf{U} \\ &\quad - 2(a_{14}^T \partial_{p_0} z_{45}^T \mathbf{U}_5)^2 - 2(a_{26}^T \partial_{p_2} z_{45}^T \mathbf{U}_5)^2 \\ &= -2S_1 \mathbf{U}_5 z_4^T \mathbf{U} - 2\xi T (\partial_{p_0} z_{45}^T \mathbf{U}_5)^2 - 2\xi T (\partial_{p_2} z_{45}^T \mathbf{U}_5)^2 \\ &= -\mathbf{U}^T \left[ \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ S_1 \\ 0 \end{pmatrix} z_4^T + z_4 \begin{pmatrix} 0 & 0 & 0 & 0 & S_1 & 0 \end{pmatrix} \right] \mathbf{U} \\ &\quad - \mathbf{U}^T \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2\xi T [|\partial_{p_0} z_{45}^T|^2 + |\partial_{p_2} z_{45}^T|^2] & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \mathbf{U}, \end{aligned}$$

which directly gives

$$\mathfrak{A}_\pi = \begin{pmatrix} 0 & 0 \\ 0 & \mathfrak{l}_\pi \end{pmatrix}_{6 \times 6}, \quad \text{with } \mathfrak{l}_\pi = \begin{pmatrix} 0 & 0 & 0 \\ 0 & -2z_{32}\mathcal{S}_1 - 2\xi T[|\partial_{v_0} z_{45}^\top|^2 + |\partial_{v_2} z_{45}^\top|^2] & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Summing over all above matrices, we finish the proof. ■