

FIELDS INTERPRETABLE IN REAL CLOSED VALUED FIELDS AND SOME EXPANSIONS

ASSAF HASSON AND YA'ACOV PETERZIL

ABSTRACT. Let $\mathcal{M} = \langle K; O \rangle$ be a real closed valued field and let k be its residue field. We prove that every interpretable field in \mathcal{M} is definably isomorphic to either K , $K(\sqrt{-1})$, k , or $k(\sqrt{-1})$. The same result holds when K is a model of T , for T a power bounded o-minimal expansion of a real closed field, and O is a T -convex subring.

The proof is direct and does not make use of known results about elimination of imaginaries in valued fields.

1. INTRODUCTION

The goal of this work is to classify fields interpretable (i.e. fields which are given as definable quotients) in real closed valued fields or, more generally, in expansions of power bounded o-minimal structures by a proper T -convex subring. This type of classification of definable quotients originates in Poizat's influential model theoretic consideration of the Borel-Tits theorem, [23] A similar study of interpretable groups and fields, in the setting of (pure) algebraically closed valued fields, was carried out By Hrushovski and Rideau-Kikuchi in [10] using different techniques.

We prove (see Section 3 for the definition of T_{conv}):

Theorem. *Let K be a real closed valued field, or more generally a model of T_{conv} , where T is an o-minimal power bounded expansion of a real closed field. Let k be its residue field. If \mathcal{F} is a field interpretable in K then \mathcal{F} is definably isomorphic to either K , $K(\sqrt{-1})$, k , or $k(\sqrt{-1})$.*

This result extends a similar theorem of Bays and the second author, [1], proved for *definable* fields in real closed valued fields. The analogous problem of definable fields in \mathbb{Q}_p was addressed in [22], but its generalisation to interpretable fields remains open.

The study of imaginaries in valued fields was suggested by Holly in [9] and first studied in depth and in full generality, in the setting of algebraically closed valued fields, by Haskell, Hrushovski and Macpherson in [5] and [6]. Elimination of imaginaries in real closed valued fields analogous to that of [5] was proved by Mellor in [18]. Some results analogous to [6] for real closed valued fields were obtained by Ealy, Haskell and Marikova in [3].

The work in [10] uses the theorems on elimination of imaginaries, stable domination and other structural results on algebraically closed valued fields, and accomplishes considerably more than the classification of interpretable fields. We adopt a different approach circumventing elimination of imaginaries, and avoiding almost completely the so called geometric sorts. In fact, our main result covers expansions of real closed valued fields by analytic functions where no elimination of imaginaries results are currently available (see [7]). Our proof is based on the analysis of one dimensional (equivalently dp-minimal) subsets of the

Date: February 3, 2022.

The second author was partially supported by Israel Science Foundation grant number 290/19 .

interpreted field \mathcal{F} , and as such it borrows ideas from Johnson's work on fields of finite dp-rank (see for example [12] and [11]), as well as [21].

The outline of the proof is as follows: We identify \mathcal{F} with X/E for a definable $X \subseteq K^n$ and a definable equivalence relation E . We then find a one-dimensional definable set $J \subseteq X$ intersecting infinitely many E -classes. After possibly shrinking J we endow it with the structure of a weakly o-minimal structure \mathcal{I} . Its universe $I = J/E$ is our basic building block (see [12] for the similar notion of a *quasi-minimal sets* or [21] for the notion of \mathbb{G} -minimal set). Next, we show that, after possibly shrinking J further, \mathcal{I} can be definably embedded in one of four weakly o-minimal structures: $\langle K; \langle \cdot \rangle \rangle$, $\langle k; \langle \cdot \rangle \rangle$, $\langle K/O; \langle \cdot \rangle \rangle$ or $\langle \Gamma; \langle \cdot \rangle \rangle$ (the value group). By analysing definable functions in K/O and showing that they are locally "affine" with respect to the additive structure of K/O , we eliminate the possibility that I could be embedded into K/O . As for Γ , we consider two cases: If T is power-bounded then the o-minimal K -induced structure on Γ cannot interpret a field (see [28]), and in fact we show that I cannot be embedded in Γ . In the exponential case we show that $\langle \Gamma, + \rangle$ is definably isomorphic to $\langle K/O, + \rangle$ and therefore, by the above, I cannot be embedded into Γ . Thus, regardless of whether T is power bounded or exponential, we are left with the first two possibilities, of K and k . Using the notion of infinitesimals we prove in these two cases that the field is definably isomorphic to a definable field in K_{conv} or in k . The final result follows from the work on definable fields in o-minimal structures ([20]).

We remark that given the o-minimal expansion of K in a signature \mathcal{L} , as in the statement of Theorem 1, it suffices to prove the result for some $\langle K; O \rangle \prec \langle K'; O \rangle$. Therefore, throughout, **we tacitly assume that $\langle K; O \rangle$ is $(|\mathcal{L}| + \aleph_0)^+$ -saturated** as will be all structures considered below.

As a corollary to our main theorem, and using the work of Hempel and Palacin [8], we obtain a theorem about definable division rings:

Corollary 1.1. *Let K be a real closed valued field, or more generally a model of T_{conv} , where T is an o-minimal power bounded expansion of a real closed field. If D is a division ring interpretable in K then D is definably isomorphic to either K , $K(\sqrt{-1})$, or the quaternions over K , or to k , $k(\sqrt{-1})$, or the quaternions over k .*

Proof. By [8, Theorem 2.9], D is a finite extension of its center, the field \mathcal{F} . By the theorem above, \mathcal{F} is definably isomorphic to a field which is definable in $\langle K; +, \cdot \rangle$, or in $\langle k; +, \cdot \rangle$. Thus D , as a finite extension, is definably isomorphic to a division ring definable in these. The result follows from [20, Theorem 1.1] on division rings in o-minimal structures. \square

Remark. (1) Weakly o-minimal structures are dp-minimal (namely, have dp-rank 1), thus if \mathcal{F} is an interpretable field in a real closed valued field (or in a model of T_{conv}) then $\text{dp-rk}(\mathcal{F}) < \omega$. It is therefore possible that the algebraic classification of \mathcal{F} , into a real or an algebraically closed field of characteristic 0, follows from the seminal work of Johnson on fields of finite dp-rank (see for example [13]). Note however that our theorem gives the additional connection between \mathcal{F} and the fields K and k .

(2) A weakly o-minimal theory is distal [25, Chapter 9], as is T^{eq} (see Exercise 9.12 there). Thus an interpretable field in T_{conv} is definable in a distal structure. As was pointed out by Chernikov, such fields cannot have characteristic $p > 0$ due to the combinatorial regularity in [2]. So our field \mathcal{F} must have characteristic zero.

However, our arguments do not make use of either of the above algebraic facts.

- (3) As the two comments above suggest, there are various model theoretic frameworks which fit our setting. The field \mathcal{F} has finite dp-rank. It is also definable in a distal structure. Finally, the structure $\langle K; O \rangle$, as well as the induced structures on K/O , k and Γ are all weakly o-minimal expansions of a groups thus they are all uniform dp-minimal structures in the sense of Simon and Walsberg, [26]. While no direct application of their paper appears in this final version we found the results there to be very helpful during our work on this project.
- (4) We expect that a similar direct proof could be given for the theorem of Hrushovski and Rideau-Kikuchi, on interpretable fields in algebraically closed valued fields.
- (5) An earlier version of the article claimed the result, wrongly, for T the theory of an arbitrary o-minimal expansion of a real closed field. We thank Yatir Halevi for noticing the gap.

1.1. Acknowledgments. A second article, in which similar methods are applied to analyze interpretable fields in P-minimal fields, was written jointly with Yatir Halevi, see [4], and we plan to combine both articles into one joint paper.

We thank the model theory group in Düsseldorf for their careful reading of the current article. Their helpful comments will be incorporated into the final version of the paper.

2. DIMENSION IN WEAKLY O-MINIMAL THEORIES

Throughout this section \mathcal{M} denotes a model of a weakly o-minimal theory.

We first collect some useful facts concerning dimension in weakly o-minimal structures. The set M is equipped with the order topology and M^n with the product topology. By an *open box* in M^n we mean a cartesian product of convex subsets of M (since M is weakly o-minimal the end points might not be in M).

We recall that if \mathcal{M} is a weakly o-minimal structure and $S \subseteq M^r$ is a definable set then $\dim(S) = l$ if l is the maximal natural number such that the projection of S on l of the coordinates has non empty interior.

We remind that definable functions in weakly o-minimal structures may be locally constant without being piecewise continuous, so that dcl need not, in general, satisfy the exchange principle. Though dcl satisfies exchange in the main sorts of T_{conv} (because it coincides with $\text{dcl}_{\mathcal{L}}$), it is not the case for K/O .

The following can be inferred readily from [24, Theorem 3.8] and the subsequent comment.

Fact 2.1. *If $Y \subseteq M^n$ is definable then $\text{dp-rk}(Y) = \dim(Y)$.*

For a a tuple in M^n and $A \subseteq M$ we let $\dim(a/A)$ be the minimal dimension of an A -definable subset of M^n containing a . Given $X \subseteq M^n$ definable over A , we say that a is *generic in X over A* if $a \in X$ and $\dim(a/A) = \dim(X)$. Our standing saturation assumption assures that generics over small parameter sets always exist.

Corollary 2.2. *If $a \in M^n$ and $A \subseteq M$ then $\text{dp-rk}(a/A) = \dim(a/A)$.*

Proof. By [24, Corollary 3.5] in dp-minimal structures, $\text{dp-rk}()$ is local, namely $\text{dp-rk}(a/A)$ is the minimal dp-rank of an A -definable set containing a . By definition, the same is true

for $\dim(a/A)$, so we can use Fact 2.1. \square

By the sub-additivity of the dp-rank, [14], we have:

Fact 2.3. $\dim(ab/A) \leq \dim(a/A) + \dim(b/Aa)$.

We shall use several times the fact that a definable subset of M^n in weakly o-minimal theories has a finite decomposition into definable cells, each homeomorphic, via an appropriate projection to an open set in some M^k (see [16, Theorem 4.11]). Also, we shall use the fact, [16, Theorem 4.7], that dimension of definable sets is preserved under definable bijection.

From now on we only use the weakly o-minimal dimension (and not dp-rank), providing – for the sake of completeness – self contained proofs of the properties we need.

Lemma 2.4. *Assume that $X \subseteq M^n$ is definable over A and that there is an externally defined set W (i.e. W is definable in an elementary extension \mathcal{N}) such that $X(M) \subseteq W$. If $\dim(W) \leq s$ then $\dim(X) \leq s$.*

Proof. By weak o-minimality, there is a cell decomposition C_1, \dots, C_l for W and projections $\pi_i : C_i \rightarrow M^{t_i}$ such that $\pi_i(C_i)$ is open and $\pi_i|_{C_i}$ is a homeomorphism. So $\dim(C_i) = t_i$. It follows that we can find N -definable sets $W_1, \dots, W_r \subseteq W$ such that every W_i has a finite-to-one projection onto s of the coordinates. Without loss of generality, W is one of these W_i and hence there is a finite-to-one projection $\pi : W \rightarrow N^s$. Since $X(M) \subseteq W$ the restriction of π to $X(M)$ is also finite-to-one. Hence $\dim(X) \leq s$. \square

We also need the following:

Lemma 2.5. *Let (J, \preceq) be a definable infinite linearly ordered set without a maximum element. Let $\{X_b : b \in J\}$ be a definable family of subsets of M^n such that for $b_1 \preceq b_2$, $X_{b_1} \subseteq X_{b_2}$. Assume that for every $b \in J$ we have $\dim(X_b) \leq m$. Let $U = \bigcup_{b \in J} X_b$. Then $\dim(U) \leq m$.*

Proof. Let $b^* \in J(N)$ be an element in an elementary extension \mathcal{N} , such that b^* is greater than all elements of $J(M)$ (since J has no maximum such a b^* exist). Since dimension is definable in parameters, $\dim(X_{b^*}) \leq m$. For every $b \in M$, $X_b(M) \subseteq X_{b^*}$ therefore $U(M) \subseteq X_{b^*}$. It follows from Lemma 2.4 that $\dim(U) \leq m$. \square

Lemma 2.6. *Let c_1, \dots, c_m be each a tuple of elements from M , of possibly different lengths, and let $C \subseteq M$ be an A -definable convex set.*

Given any initial segment $I \subseteq C$ (possibly defined over additional parameters), there exists $b \in I$ such that for every $i = 1, \dots, m$, $\dim(bc_i/A) = 1 + \dim(c_i/A)$.

Proof. For each $i = 1, \dots, m$ we fix W_i , definable over A , such that $\dim(c_i/A) = \dim W_i$. Consider the type $p(x)$:

$$\{x \in I\} \cup \bigcup_{i=1}^m \{\neg X(x, c_i) : \dim(X) \leq \dim(c_i/A) \text{ and } X \subseteq M \times W_i \text{ definable over } A\}.$$

Since C is definable over A , we may take above only those sets X such that for every $w \in W_1$, $X(M, w)$ is a nonempty subset of C .

If p is consistent then its realization will be the desired b . So, we assume towards contradiction that p is inconsistent.

It follows that there are X_1, \dots, X_r as above, all A -definable, and for each $j = 1, \dots, r$, there is $i(j) \in \{1, \dots, m\}$, such that

$$\forall x \left(x \in I \rightarrow \bigvee_{j=1}^r X_i(x, c_{i(j)}) \right).$$

In addition, for each $j = 1, \dots, r$, $\dim(X_j) \leq \dim(W_{i(j)})$.

By weak o-minimality, each set $X_j(M, c_{i(j)})$ is a finite union of convex subsets of C , and since I is a non-empty initial segment of C , the above implies that for some fixed j , the set $X_j(M, c_{i(j)})$ contains an initial segment of C . For simplicity, we write X for X_j and assume that $i(j) = 1$. Namely, we assume that $X(M, c_1)$ contains an initial segment of C . This is an A -definable property of c_1 , therefore we may assume, after possibly shrinking W_1 , that for every $w \in W_1$, the set $X(M, w)$ contains an initial segment of C . Moreover, by choosing in each $X(M, w)$ the first convex component, we may assume that for every $w \in W_1$, the set $X(M, w)$ contains a single convex set, which is an initial segment of C .

For $b \in C$, let $X_b = \{w \in W_1 : \langle b, w \rangle \in X\}$. Notice that $\bigcup_{b \in C} X_b = W_1$. Our assumptions imply that for $b_1 \leq b_2 \in C$, we have $X_{b_2} \subseteq X_{b_1}$. It now follows from lemma 2.5 (with the ordering $<$ reversed) that for some $b_0 \in C$ we have $\dim(X_{b_0}) = \dim(W_1)$.

Also, our assumptions imply that for every $b < b_0$ in C and $w \in X_{b_0}$ we have $\langle \langle b, w \rangle \in X$, namely $C^{<b_0} \times X_{b_0} \subseteq X$. Thus, $\dim(X) \geq 1 + \dim X_{b_0} > \dim W_1$. But by our assumption, $\dim(X) \leq \dim(c_1/A) = \dim(W_1)$. This is a contradiction and the lemma is proved. \square

As a corollary we obtain:

Corollary 2.7. *Assume that \mathcal{M} has definable Skolem functions, and let $X \subseteq M^n$ be an A -definable set. Let c_1, \dots, c_m be tuples from M (of possibly different lengths). Then there exists $b \in X$ such that for each $i = 1, \dots, m$,*

$$\dim(bc_i/A) = \dim(X) + \dim(c_i/A).$$

Proof. By cell decomposition we may assume that X is an open subset of M^n . We use induction on n .

The case $n = 1$ follows from Lemma 2.6. For a general n , note first that X contains an A -definable open box $I_1 \times \dots \times I_n$. Indeed, since X is open it contains an open box and by the existence of Skolem functions we can find such a box defined over A . Applying induction, we can find $a' = \langle a_2, \dots, a_n \rangle \in I_2 \times \dots \times I_n$ such that for each $i = 1, \dots, m$, $\dim(a'c_i/A) = (n-1) + \dim(c_i/A)$.

We now let $c'_i = \langle a', c_i \rangle$, for $i = 1, \dots, m$, and using again the case $n = 1$, find $a_1 \in I_1$ such that, for all i $\dim(a_1c'_i/A) = n + \dim(c_i/A)$. The tuple $\langle a_1 a' \rangle$ is the desired b . \square

In the above existence of Skolem functions somewhat simplifies the proof, but is not, in fact, needed.

2.1. Dimension and domination in quotients by convex equivalence relations. We establish here several properties of quotients of one-dimensional sets by convex equivalence relations. Recall:

Definition 2.8. An equivalence relation E on a linearly ordered set $(J, <)$ is convex if all its equivalence classes are.

If \mathcal{M} is weakly o-minimal, E a definable convex equivalence relation with infinitely many classes then M/E is linearly ordered and itself weakly o-minimal. If in addition $Th(\mathcal{M})$ is weakly o-minimal then so is $Th(\mathcal{M}/E)$. It follows, in particular (see next section), that if $\langle K; O \rangle \models T_{conv}$ then the theories of K , K/O , Γ and k are weakly o-minimal.

Let \mathcal{M} be a weakly o-minimal structure, E a definable convex equivalence relation on M , and let $\pi : M \rightarrow M/E$ be the quotient map. For simplicity we also denote by E the equivalence relation on M^n given by E^n and let $\pi : M^n \rightarrow (M/E)^n$ be the associated quotient map. We prove here a domination result for such quotients.

Lemma 2.9. *Let $I_1, \dots, I_n \subseteq M$ be open convex sets such that each I_j/E is infinite, and $X \subseteq B = I_1 \times \dots \times I_n$. If $\pi(X) = \pi(B)$ then $\dim(\pi(B \setminus X)) < n$.*

Proof. Let us say that a definable set X crosses an E -class $[a]$ if both X and X^c intersect $[a]$.

We proceed by induction on n . The result for $n = 1$ follows from weak o-minimality of K/E : Indeed, if $X \subseteq I$ is definable then, since X consists of at most r -convex sets and each E -class is convex, then X crosses at most $2r$ -many classes. Thus, since X intersects all E -classes, $\pi(I_1 \setminus X)$ is finite.

Assume now that $X \subseteq M^n$ and $\pi(X) = \pi(B)$. Let $Y = B \setminus X$, and assume towards a contradiction that $\dim(\pi(Y)) = n$. This implies that $\pi(Y)$ has nonempty interior, and therefore contains a box. The pre-image of such a box is itself a box (since E is convex, and we may assume that its domain is an open box). Thus there exists an open box $B_1 \subseteq Y$ such that $\pi(B_1)$ is an open subset of $\pi(X)$. By our assumption, we also have $\pi(B_1 \cap X) = \pi(B_1) = \pi(B_1 \cap Y)$.

Thus we may assume, by replacing B with B_1 , that $\pi(X) = \pi(Y) = \pi(B)$. To simplify notation let $B' = I_1 \times \dots \times I_{n-1}$. We define

$$X^* = \{ \langle a, b \rangle \in B' \times I_n : X_a \text{ crosses } [b] \}.$$

By the case $n = 1$, for each $a \in B'$, X_a crosses at most finitely many E -classes in I_n . It follows that $\pi(X^*) \subseteq \pi(B)$ projects finite-to-one into $\pi(B')$, and thus $\dim(\pi(X^*)) < n$. Hence, there exists an open box $R \subseteq \pi(B)$ which is disjoint from $\pi(X^*)$. By replacing B with $\pi^{-1}(R)$ we may assume that $X^* = \emptyset$. Namely, for every $\langle a, b \rangle \in B' \times I_n$, either X_a contains $[b]$ or it is disjoint from $[b]$.

For every $b \in I_n$, we let $X^b = \{ a \in B' : \langle a, b \rangle \in X \}$. We claim that for each $b \in I_n$, $\pi(X^b) = \pi(B')$. Indeed, if $a \in B'$ then there is $\langle a', b' \rangle \in X$ such that $\langle a', b' \rangle E \langle a, b \rangle$ (since $\pi(X) = \pi(B)$). By our assumption, $X_{a'}$ does not cross $[b']$, therefore $[b']$ is contained in $X_{a'}$. We have bEb' , hence $\langle a', b \rangle \in X$. It follows that $\pi(a) = \pi(a') \in \pi(X)$. Our induction assumption thus implies that $\dim(B' \setminus X^b) < n - 1$.

We also claim that for $b_1 E b_2$, we have $X^{b_1} = X^{b_2}$ (and hence also $B' \setminus X^{b_1} = B' \setminus X^{b_2}$). Indeed, if $a \in X^{b_1}$ then $X_a \cap [b_1] \neq \emptyset$ and therefore $[b_1] \subseteq X_a$, so $b_2 \in X_a$, namely $a \in X^{b_2}$. The opposite inclusion follows.

Thus, for every $b \in I_n$, we have $\pi(B \setminus X)^{\pi(b)} = \pi(B' \setminus X^b)$, which has dimension $< n - 1$. It follows that for every $c \in \pi(I_n)$, $\dim(\pi(B \setminus X)^c) < n - 1$. By the sub-additivity of dimension, $\dim(\pi(B \setminus X)) < n$, contradicting our assumption. \square

The following could be viewed as domination of types in M^n by generic types in M^n/E .

Proposition 2.10. *In the above setting, assume that $p \vdash (M/E)^n$ is a complete generic type over $A \subseteq M$. Then $q = \pi^{-1}(p)$ is a complete type over A .*

Proof. If q is not complete then there is an A -definable set $X \subseteq M^n$ such that $q \vdash \pi(X)$ and $q \vdash \pi(X^c)$. But then $\dim(\pi(X)) = \dim(\pi(X^c)) = n$, contradicting Lemma 2.9. \square

3. THE THEORY T_{conv}

Let T be an o-minimal expansion of a real closed field in a signature \mathcal{L} . The theory T_{conv} was introduced by Lowenberg and v.d. Dries in [31]: Given $K \models T$, a T -convex subring of K is a subring $O \subseteq K$ with the property that $f(O) \subseteq O$ for any \emptyset -definable continuous function $f : K \rightarrow K$. Since O is convex it is a valuation ring, its maximal ideal is denoted μ , the value group is Γ and the residue field is k . The theory T_{conv} , in the language $\mathcal{L}_{conv} := \mathcal{L} \cup \{O\}$, is the extension of T by the axioms saying that O is a proper valuation ring which is T -convex. Here are some results from [31] and [28]:

Fact 3.1. (1) *The theory T_{conv} has quantifier elimination relative to T , it is complete and weakly o-minimal. ([31, 3.10, 3.13, 3.14].*

Let $T_{conv,c}$ be the extension of T_{conv} by the formula $c \notin O$, for a new constant c . Then:

- (2) *The theory $T_{conv,c}$ has definable Skolem functions ([28, Remark 2.4]).*
- (3) *In models of $T_{conv,c}$ the definable closure in \mathcal{L} and in \mathcal{L}_{conv} are the same (see [28, Lemma 2.6]). Moreover, every \mathcal{L}_{conv} -definable function coincides with an \mathcal{L} -definable function around generic points of the domain.*

From now on, we add a constant symbol c interpreted as a positive element outside O , and still denote the new language \mathcal{L}_{conv} . Throughout the text we will be using the above results without further reference.

Definition 3.2. Let \mathcal{M} be an o-minimal expansion of real closed field K . A *power function* on K is a definable endomorphism of the ordered multiplicative group $K^{>0}$.

\mathcal{M} is called *power bounded* (generalizing ‘‘polynomially bounded’’) if every definable function of one variable is eventually bounded by some power function. An o-minimal theory T is called *power bounded* if every model of T is power bounded.

By Miller’s [19], \mathcal{M} is power bounded if and only if it is not exponential, namely one cannot define a (necessarily monotone) isomorphism of $(K, +)$ and $(K^{>0}, \cdot)$.

The structure of the residue field and the value group of T_{conv} is described by the work of v.d. Dries :

Fact 3.3 (Theorem A, Theorem B). [28]] *Let T be the theory of an o-minimal expansion of a real closed field K and O a T -convex subring of R . Then:*

- (1) *The residue field with its induced structure can be given a structure of a T -model, and it is stably embedded as such. In particular it is o-minimal, so by [20], any field interpretable in the residue field is definably isomorphic to the residue field itself or to its algebraic closure.*
- (2) *If, in addition, T is power bounded, then the value group is, up to a change of signature, an ordered vector space over the field of exponents of T and is stably embedded as such.*

We shall also use the following result from J. Tyne’s PhD thesis [27, p.94]:

Fact 3.4. *Let T be power bounded and o-minimal and $K \text{conv} \models T_{\text{conv}}$. Then every $\mathcal{L}_{\text{conv}}$ -definable subset of K is already definable in the valued field, namely it is a boolean combination of balls and intervals.*

To sum up the last facts in few words, if we denote k_{ind} the induced structure on k (in the signature \mathcal{L}_{ind}) then k_{ind} is o-minimal, and up to a change of signature elementarily equivalent to R . In case T is power bounded, the value group, too, is an o-minimal vector space with no additional structure and the 1-variable definable subsets of K are the same as in the underlying real closed valued field.

From now on, we denote the induced structure on k by k_{ind} and its language by \mathcal{L}_{ind} .

We finally note:

Lemma 3.5. *Let T be an exponential o-minimal expansion of a real closed field, T_{conv} its expansion by a T -convex valuation ring. If $K \models T$ then $\langle K/O; <, + \rangle$ is in definably isomorphic with $\langle \Gamma; <, + \rangle$.*

Proof. We first note that $\exp(O) = O_{>0} \setminus \mu$. Indeed, the right-to-left inclusion follows from the fact that O is T -convex, and $\exp(x) > x$. For the converse, assume for contradiction that $a = \exp(b) \in \mu$ for some $b \in O$. Then $a^{-1} = \exp(-b) \notin O$, contradicting the fact that O is T -convex.

Since $O \setminus \mu = O^\times$ it follows that \exp induces a definable isomorphism between $\langle K/O; <, + \rangle$ and $\langle K_{>0}/O_{>0}^\times; <, + \rangle$, which is isomorphic to $\langle \Gamma; <, + \rangle$. \square

For example, in this case the residue field is internal to the value group, and the latter's theory (relative to T) is not known (see [28, Clause 6.5]).

It will be convenient to define in $\mathcal{L}_{\text{conv}}$ the relation $v(x) > v(y)$ if $x/y \in O \wedge y/x \notin O$. Since $O = \{x : v(x) \geq v(1)\}$, quantifier elimination for T_{conv} in $\mathcal{L}_{\text{conv}}$ is equivalent to quantifier elimination for \mathcal{L} expanded by the above binary predicate, and we will use the two languages interchangeably.

The following could probably be read off [28], but we need the explicit formulation below. It is a consequence of quantifier elimination relative to T :

Lemma 3.6. *Let $S \subseteq K^n$ be $\mathcal{L}_{\text{conv}}$ -definable, $a \in S$ generic over the parameters defining S (in the weakly o-minimal sense). Then there exists an open box $B \ni a$ (defined possibly over new parameters) such that $B \cap S$ is \mathcal{L} -definable.*

Proof. We first note that the associated valuation induces the same topology on K as the order topology. Quantifier elimination implies that any definable set is of the form

$$\bigcup_i \left(\bar{T}_i(\bar{x}) \cap \left(\bigcap_j (t_{i,j}(\bar{x}) \in O)^{\pm 1} \right) \right)$$

where \bar{T}_i are \mathcal{L} -definable sets, $t_{i,j}(\bar{x})$ are \mathcal{L} -terms and $(x \in O)^{-1}$ means $x \notin O$. It suffices to verify the lemma for sets of the form

$$\bar{T}(\bar{x}) \cap \left(\bigcap_j (t_j(\bar{x}) \in O)^{\pm 1} \right)$$

The set O is clopen in K and since a is generic with respect to the o-minimal structure, each $t_{i,j}$ defines a continuous function at a . It follows that a is an interior point of both

the preimage of O and its complement under each t_j . Thus there is an open box $B \subseteq K^n$ containing a such that B is contained in the set $\bigcap_j (t_j(\bar{x}) \in O)^{\pm 1}$. We have $T \cap B = S \cap B$, so it is \mathcal{L} -definable. \square

4. DEFINABLE QUOTIENTS OF K

In this section we classify, locally, definable quotients of K itself. The first lemma will eventually allow us to reduce much of the work from quotients of the form K^n/E to quotients of the form $(K/E)^n$.

Lemma 4.1. *Let \mathcal{M} be a weakly o-minimal structure with definable Skolem functions. Let $S = X/E$, for some definable $X \subseteq K^n$ and E a definable equivalence relation. If S is infinite then there exists an infinite $S' \subseteq S$ (definable over additional parameters) such that S' is in definable bijection with J/E' for some interval $J \subseteq M$ and E' a definable equivalence relation on M .*

Proof. We use induction on n , where the case $n = 1$ is trivial. Let $W := \pi(X)$ be the projection of X onto the last $n - 1$ coordinates (assuming, without loss of generality, this projection has infinite image). If for some $w \in W$ the fibre $\pi^{-1}(w)$ meets infinitely many E -classes then take $S' = X^w = \{x \in M : \langle x, w \rangle \in X\}$ and E' the relation on S' obtained from $\pi^{-1}(w)$.

So we may assume that $|\pi^{-1}(w)/E| < \infty$ for all $w \in \pi(X)$. Since \mathcal{M} has definable Skolem function we can find, uniformly for each $w \in W$ a finite set of representatives $Y(w)$ of $\pi^{-1}(w)/E$. More precisely, there exists a definable set $Z \subseteq K^n$ such that $\pi(Z) = \pi(X)$ and for each $w \in \pi(X)$, $Z \cap \pi^{-1}(w)$ has exactly one representative for each E -class intersecting $\pi^{-1}(w)$.

By weak o-minimality, there exists a cell decomposition Z_1, \dots, Z_l of Z such that each Y_i is definably homeomorphic to an open cell $C_i \subseteq M^{\dim(Z_i)}$. Assume, without loss, that $|Z_1/E| = \infty$. If $\pi : Z_1 \rightarrow C_1$ is the definable homeomorphism let $E_1 := \pi(E|Z_1)$ and we conclude by induction. \square

We now fix an infinite field $\langle \mathcal{F}; +, \cdot \rangle$ interpretable in K . We fix an interval $J \subseteq K$ and a definable equivalence relation E on J as provided by the above lemma. As we show next, after possibly shrinking J , the set J/E can be definably embedded into one of four weakly o-minimal structures.

Theorem 4.2. *Let $K \models T_{conv}$, where T is power bounded, and let E be a definable equivalence relation on $J \subseteq K$ with infinitely many classes. Then there exists a definable ordering on J/E such that $\langle J/E, < \rangle$, with the induced structure, is weakly o-minimal, and there exists an $<$ -interval $I \subseteq J/E$ which is in definable bijection with an interval in K , in Γ , in k or in K/O .*

Proof. By weak o-minimality each E -equivalence class is a finite union of convex sets. So by replacing E with the equivalence relation E' , choosing the first component in each E -class, we get a convex equivalence relation with $D/E' = J/E$ for some definable $D \subseteq J$. So we assume that $E = E'$. Thus J/E is linearly ordered and weakly o-minimal.

If E is the identity on an infinite set we can find a convex set $I \subseteq J$ where this is true, and then $I/E \hookrightarrow K$, so we are done. So we assume that this is not the case. It follows that for all but finitely many classes, $\sup([x])$ is not an element of K (for otherwise the set $\exists y(x = \sup[y])$ is an infinite discrete set in K). For the same reason $\inf([x])$ is not an element of K for all but finitely many of the E -classes. By shrinking J we may assume that for all $x \in J$, the class $[x]$ is a bounded convex set without supremum or infimum.

For a valational ball B (either open or closed) write $x < B$ if $x < y$ for all $y \in B$. We write $x \leq B$ if $x < B \vee x \in B$. Recall (see, e.g., [18]) that a *disc cut* is a set of the form $x \square B$ where $\square \in \{<, \leq\}$ and B is a definable ball (either open or closed) or a point in K . By Tyne's work, Fact 3.4, every \mathcal{L}_{conv} -definable subset of K is a finite boolean combination of disc cuts. Since the E -classes are convex it follows from our assumptions that they are intersections of two disc cuts where B is actually a ball (and not a point).

Let us call a set of the form $B_1 \square_1 x \square_2 B_2$, where $\square_i \in \{<, \leq\}$, a *ball-interval*. The *form* of a ball interval is the data specifying which of the balls B_i is open, and which of the inequalities is weak.

Below we use valational radius for balls, namely for $\gamma \in \Gamma$, an open ball of radius γ is a set of the form $\{x \in K : v(x - x_0) > \gamma\}$ and a closed ball is of the form $\{x \in K : v(x - x_0) \geq \gamma\}$.

Claim. If C is a ball interval $B_1 \square_1 x \square_2 B_2$ then B_1 and B_2 are uniquely determined by C . More explicitly, B_1 and B_2 are definable over a code for C .

Proof. We deal first with the case where \square_1 is \leq . If B_1 is a ball of radius γ (either closed or open) then

$$\gamma = \inf\{\delta : (\forall x \in C \exists y \in C)(y < x \wedge B_\delta(y) \subseteq C)\}.$$

And $B_1 = B_\gamma(y)$ for any small enough $y \in C$. We can recognise whether B_1 is open or closed by verifying whether γ is attained as a minimum in the above formula or not. B_2 is defined similarly when \square_2 is \leq .

The case of strong inequalities follows from the previous case by replacing I with its complement. □claim.

The claim allows us to definably assign to each class $[x]$ the left and right “end-balls” of the ball-interval, $B_1[x]$ and $B_2[x]$, respectively. There are finitely many possible forms of ball-intervals, hence we may further assume that all equivalence classes $[x] \subseteq J$ have the same form. Note that if $[x] \neq [y]$ then $B_1[x] \neq B_1[y]$, for otherwise $[x] \cap [y] \neq \emptyset$. So the function $[x] \mapsto B_1[x]$ is a definable injection into the family of open (closed) balls. If for some $\gamma \in \Gamma$ the fibre of the function sending $[x]$ to the valational radius of $B_1[x]$ is infinite then, depending on whether $B_1[x]$ is open or closed, we get an injection of some interval in J/E into K/O or k .

If all fibres of this function are finite we get a finite-to-one map from an interval in J/E into Γ . The linear order on J/E implies the existence of an injection from an interval in J/E into Γ , finishing the proof of Theorem 4.2. □

5. DEFINABLE FUNCTIONS IN K/O

Combining Lemma 4.1 with Theorem 4.2 we conclude that there exists an infinite definable I in our interpretable field \mathcal{F} such that I is in definable bijection with an interval in one of the four weakly o-minimal structures K , Γ , k or K/O . While the definable sets in K , Γ and k are well understood, the situation is less clear in K/O . Our goal in this section is to show that definable functions in K/O are locally affine with respect to the additive structure of K/O . The “polynomial boundedness” assumption is not used in this section.

We start with the following general lemma.

Lemma 5.1. *Assume that $p \vdash K^n$ is a partial \mathcal{L} -type and that $H : K^n \rightarrow K$ is a partial \mathcal{L} -definable function such that $p \vdash \text{dom}(H)$.*

Assume also that $C \subseteq K$ is an $\mathcal{L}_{\text{conv}}$ -definable convex set which is bounded above (or below) and for every $\alpha \models p$, $H(\alpha) \in C$. Then there exists $c \in C$ such that for every $\alpha \models p$ we have $H(\alpha) < c$ (or $H(\alpha) > c$).

Proof. We are using the saturation of $\mathcal{M}_{\text{conv}}$, and assume that C is bounded above. For every \mathcal{L} -definable set $X \subseteq K^n$, for which $p \vdash X$, let $s(X)$ be the supremum of $H(W)$, which exists by o-minimality and the boundedness of C . We claim that for some such X we have $s(X) \in C$ (and then we may take $c = s(X)$). Indeed, if not then for every X in p there exists $\alpha \in X$ such that $s(\alpha) > C$. But then, by saturation we may find $\alpha \models p$ such that $H(\alpha) > C$, contradicting our assumption. \square

Definition 5.2. A subset of K is called *long* if it contains infinitely many cosets of O . A subset of K^n is *long* if it contains a cartesian n -product of long subsets of K . A type $p \vdash K^n$ is long if every set in p is long.

Definition 5.3. We say that a (partial) function $F : K^n \rightarrow K$ *descends to K/O* if whenever $a - b \in O^n$ also $F(a) - F(b) \in O$.

Example 5.4. If $a \in O$ then the linear function $x \mapsto a \cdot x$ descends to an endomorphism of $\langle K/O, + \rangle$. In the case that $a \in \mu$, the map descends to an endomorphism of K/O with infinite kernel. Thus we obtain a definable locally constant, surjective endomorphism on K/O .

For a K -differentiable $F : K^n \rightarrow K$, and $i = 1, \dots, n$, we let F_{x_i} denote the partial derivative with respect to x_i , and let $\nabla F = (F_{x_1}, \dots, F_{x_n})$.

We are going to use several times the following (see Exercise 3.2.19(2)): *If $F : K^n \rightarrow K$ is an \mathcal{L} -definable partial function and $p \vdash \text{dom}(F)$ is a generic type then the restriction of F to $p(\mathcal{M})$ is monotone in each coordinate separately.*

Lemma 5.5. *Assume that $F : K^n \rightarrow K$ is an $\mathcal{L}(A)$ -definable (partial) function which descends to K/O . Assume that $p \vdash \text{dom}(F)$ is a partial long $\mathcal{L}_{\text{conv}}(A)$ -type which implies a complete generic $\mathcal{L}(A)$ -type in K^n ,*

Then for every $a \models p$, and $i = 1, \dots, n$, we have $F_{x_i}(a) \in O$.

Proof. Because p implies a generic type in K^n the function F is smooth at every $a \models p$. Assume towards a contradiction that the conclusion of the lemma does not hold. Without loss of generality, assume that $F_{x_1}(a) > O$ for some $a = (a_1, \dots, a_n) \models p$.

Using monotonicity of F_{x_1} in the first coordinate, we can find some $\alpha > O$ and an interval $J \subseteq K$ of length greater than 1 such that every element in $J \times \{a_2, \dots, a_n\}$ still realizes p , and for every $x \in J$, $F_{x_1}(x, a_2, \dots, a_n) > \alpha$. By Lagrange's Mean Value Theorem for o-minimal structures, for $x, x+1 \in J$, $F(x+1, a_2, \dots, a_n) - F(x, a_2, \dots, a_n) > \alpha$, contradicting the assumption that F descends to K/O . \square

Lemma 5.6. *Assume that $G : K^n \rightarrow K$ is an $\mathcal{L}(A)$ -definable partial function, and $p \vdash \text{dom}(G)$ is a long partial $\mathcal{L}_{\text{conv}}(A)$ -type which implies a complete generic $\mathcal{L}(A)$ -type in K^n . Assume also that for every $a \models p$, $G(a) \in O$. Then for all $a \models p$ and all $i \in \{1, \dots, n\}$, we have $G_{x_i}(a) \in \mu$.*

Proof. We prove the result for $i = 1$. By Lemma 5.1, there is $c \in O$ such that for all $a \models p$ we have $|G(a)| \leq c$. Assume towards contradiction that for some $a = (a_1, \dots, a_n) \models p$ we have $|G_{x_1}(a)| > \mu$. Then we can find a positive $\alpha \in O \setminus \mu$, such that $|G_{x_1}(a)| > \alpha$.

Using the monotonicity of G_{x_1} in the first coordinate, we may assume that for every positive $b \in O$, we have $G_{x_1}(a_1 + b, a_2, \dots, a_n) > \alpha$. Pick $b > 2c/\alpha \in O$. Then, by Lagrange's mean value theorem in o-minimal structures,

$$|G(a_1 + b, a_2, \dots, a_n) - G(a_1, \dots, a_n)| > |G_{x_1}(\xi, a_2, \dots, a_n)|(2c/\alpha)$$

for $\xi \in (a_1, a_1 + b)$. By our assumptions, $|G_{x_1}(\xi, a_2, \dots, a_n)| > \alpha$. It follows that $|G(a_1 + b, a_2, \dots, a_n) - G(a_1, \dots, a_n)| > 2c$, in contradiction to the fact that $|G(x)| \leq c$ for all $x \models p$. \square

We conclude:

Lemma 5.7. *Assume that $F : K^n \rightarrow K$ is a partial $\mathcal{L}(A)$ -definable function which descends to K/O . Let $p \vdash \text{dom}(F)$ be a long $\mathcal{L}_{conv}(A)$ -type which implies a complete generic $\mathcal{L}(A)$ -type. Then there are $a_1, \dots, a_n \in O$, $b \in K$ and a long box $B \subseteq p(K)$ such that for all $x = \langle x_1, \dots, x_n \rangle \in B$,*

$$F(x) - \sum_{i=1}^n a_i(x_i - \alpha_i) - b \in \mu,$$

Proof. By Lemma 5.5, each F_{x_i} takes values in O , on the type p . By Lemma 5.6, applied to each of the functions F_{x_i} , we have $F_{x_i, x_j}(a) \in \mu$ for all $a \models p$ and $i, j = 1, \dots, n$. By Lemma 5.1, there is some fixed positive $\beta \in \mu$ such that $|F_{x_i, x_j}(a)| < \beta$ for all $a \models p$.

Fix $\alpha \models p$ and let $(a_1, \dots, a_n) = \nabla F(\alpha)$.

By [30, Lemma 7.2.9], for all $x \models p$,

$$(*) \quad \left| F(x) - F(\alpha) - \sum_{i=1}^n a_i(x_i - \alpha_i) \right| \leq |x - \alpha| \cdot \max_{y \in [\alpha, x]} |\nabla F(x) - \nabla F(\alpha)|,$$

where $[\alpha, x]$ is the closed line segment in K^n connecting α and x and $|\nabla F(x)|$ is the operator norm.

Fix an $x \models p$ and for $i = 1, \dots, n$, and $t \in [0, 1]$, consider the function

$$g_i(t) = F_{x_i}((1-t)\alpha + tx).$$

The derivative of $\frac{dg_i}{dt}$ is, by the chain rule,

$$(F_{x_1, x_i}(y), \dots, F_{x_n, x_i}(y)) \cdot (x - \alpha).$$

Applying Lagrange's Theorem to g_i and substituting, we get $\xi \in (0, 1)$ such that

$$|F_{x_i}(x) - F_{x_i}(\alpha)| = |g_i(1) - g_i(0)| = |g_i'(\xi)| = |(F_{x_1, x_i}(y), \dots, F_{x_n, x_i}(y)) \cdot (x - \alpha)|,$$

where $y = (1 - \xi)\alpha + \xi x$.

Since each $|F_{x_j, x_i}(y)| < \beta$ we have $|F_{x_i}(x) - F_{x_i}(\alpha)| \leq \beta|x - \alpha|$ for all $i = 1, \dots, n$. Thus $|\nabla F(x) - \nabla F(\alpha)| \leq n\beta|x - \alpha|$, so by (*),

$$|F(x) - F(\alpha) - \sum_{i=1}^n a_i(x_i - \alpha_i)| \leq n\beta|x - \alpha|^2.$$

Since p is long and $\beta \in \mu$, we can find a long box $B \subseteq p(\mathcal{M})$, centered at α such that for every $x \in B$, we have $|x - \alpha| \leq (1/\beta)^{1/4}$. Thus, for $x \in B$ we have

$$|F(x) - F(\alpha) - \sum_{i=1}^n a_i(x_i - \alpha_i)| \leq n\sqrt{\beta} \in \mu.$$

If we now let $b = F(\alpha)$ then we have the desired result. \square

In fact, for the next corollary it would have been sufficient to show above that $F(x) - F(\alpha) - \sum_{i=1}^n a_i(x_i - \alpha_i) + b \in O$.

Corollary 5.8. *Assume that $f : (K/O)^n \rightarrow K/O$ is a (partial) \emptyset -definable function whose domain is open. Then for every generic $c \in \text{dom}(f)$ there exists an open box $R \subseteq \text{dom}(f)$ centered at c , definable endomorphisms of $\langle K/O; + \rangle$ denoted by ℓ_1, \dots, ℓ_n , and $d \in K/O$, such that for every $y = \langle y_1, \dots, y_n \rangle \in R$ we have*

$$f(y) = \sum_{i=1}^n \ell_i(y) + d.$$

Proof. Let $p = \pi^{-1}(tp_{K/O}(c/\emptyset))$. It is clearly a long type and by Proposition 2.10, it is in fact a complete \mathcal{L}_{conv} -type which implies a generic \mathcal{L} -type. Using definable Skolem functions we may lift f to an \mathcal{L} -definable partial $F : K^n \rightarrow K$, which descends to K/O . Namely, for every $x \in \text{dom}(F)$, we have $F(x) \in f(x + O^n)$. We have $p \vdash \text{dom}(F)$.

Applying Lemma 5.7, we can find a long box $B \subseteq K^n$ centered at $\alpha \models p$, and $a_1, \dots, a_n \in O, b \in K$, such that for every $x \in B$ we have

$$F(x) - b - \sum_{i=1}^n a_i(x_i - \alpha_i) \in \mu.$$

Each function $x \mapsto a_i \cdot x$ descends to an endomorphism ℓ_i of $\langle K/O; + \rangle$, so if we let R be the image of B in K/O and $e = b + O$, then we have for all $y \in R$,

$$f(y) = \sum_{i=1}^n \ell_i(y_i - \alpha_i) + e = \sum_{i=1}^n \ell_i(y_i) + d,$$

for $d = e - \sum_{i=1}^n \ell_i(\alpha_i)$. \square

We end this section by commenting that in the *o-minimal setting* the local affiness of definable functions provided by Corollary 5.8, would imply that the structure is linear (in the sense of [15]), and thus does not interpret a field. This is not true for K/O :

Proposition 5.9. *The field k is definably isomorphic to a field interpretable (in induced structure on) K/O .*

Proof. Fix some t such that $v(t) < 0$. Consider the balls $tO \supseteq t\mu$. Since $O \subseteq t\mu$, the subgroups tO/O and $t\mu/O$ are definable subgroups of K/O . Thus, we get that the quotient $\langle tO/t\mu; + \rangle$ is interpretable in K/O . The former is definably isomorphic to $\langle k; + \rangle$, hence we can definably endow it with multiplication which makes it isomorphic to $\langle k; +, \cdot \rangle$. \square

As we shall show in Proposition 6.4, no field is *definable* in K/O .

6. SUBSETS OF \mathcal{F} THAT ARE STRONGLY INTERNAL TO I

As we saw in Theorem 4.2, if T is power bounded then there exists an infinite \mathcal{L}_{conv} -definable $I \subseteq \mathcal{F}$ such that I is in definable bijection with an interval in one of the structures K, Γ, k or K/O . Once we obtain such a set I we can drop the assumption that T is power bounded. In all four cases, the induced structure on I is weakly o-minimal. Let us be more precise about this:

Remark 6.1. Recall that given an \aleph_1 -saturated structure \mathcal{M} a definable set $S \subseteq M^n$ is *stably embedded* if for any definable $X \subseteq M^{nk}$ the set $X \cap S^k$ is definable using only

parameters from S . Thus the \mathcal{M} -induced structure on S does not depend on the model or on the choice of parameters.

However, intervals in a weakly o-minimal structure need not be stably embedded so that the notion of “the structure induced on I ” is not well defined. For $A \subseteq K$ we let \mathcal{I}_A denote the structure obtained by taking the traces on I^n , $n \in \mathbb{N}$, of all A -definable sets. The structure \mathcal{I}_K has a weakly o-minimal theory. So the same is true of any ordered reduct of the full K -induced structure on I and in particular of reducts of the form \mathcal{I}_A . We assume that \mathcal{F} , I , and an embedding of I into K , k , Γ or K/O are all definable over \emptyset , and initially work in \mathcal{I}_\emptyset . We will, throughout the course of the proof, add small sets of parameters (not necessarily coming from I), tacitly expanding the structure \mathcal{I}_\emptyset in into one of the form \mathcal{I}_A . We denote the resulted structure by \mathcal{I} .

Definition 6.2. A definable $Y \subseteq \mathcal{F}$ is *strongly internal to I* (or *strongly I -internal*) if there exists a definable injection, possibly over additional parameters, from Y into I^r for some r .

We now fix $Y \subseteq \mathcal{F}$ which is strongly internal to I and has maximal dimension, denoted by n , in the sense of I (equivalently, maximal dp-rank).

We identify Y with its image in I^r and view it with the induced I -topology. We assume for simplicity that Y is definable over \emptyset . Unless otherwise stated we use $+$, $-$, \cdot , $()^{-1}$ for the field operations of \mathcal{F} .

Proposition 6.3. Let $\langle b, c, d \rangle \in I \times Y^2$ be such that $\dim(b, c, d/\emptyset) = 2n + 1$. Then there exists an initial segment $J \subseteq I^{>b}$ and a definable $S \subseteq Y^2$, all defined over an additional parameter set B , such that $\dim(c, d/B) = 2n$, $\langle c, d \rangle \in S$, (in particular, $\dim(S) = 2n$), and such that for every $\langle x, y, z \rangle \in J \times S \subseteq J \times Y^2$, we have $(x - b)y + z \in Y$.

Proof. For $\langle x, y, z \rangle \in I \times Y \times Y$ consider the function $f_y(x, z) = xy - z$.

Claim The point $\langle b, d \rangle \subseteq I \times Y \subseteq I^{1+r}$ is not an isolated point of the set $f_c^{-1}(f_c(b, d))$.

Proof. Assume $\langle b, d \rangle$ were isolated in $f_c^{-1}(f_c(b, d))$. By the weak o-minimality of $Th(\mathcal{I})$, there is a number r such that for every $d \in \text{Im}(f_c)$, the set $f_c^{-1}(d)$ has at most r -many isolated points. Because I is linearly ordered, each of these points is in $\text{dcl}(c, d)$. Thus, there is a c -definable (partial) function $\sigma : \mathcal{F} \rightarrow I \times Y$, such that for every $w \in \text{dom}(\sigma)$, $\sigma(w)$ is an isolated point of $f_c^{-1}(w)$, and if $w = f_c(b, d)$ then $\sigma(w) = \langle b, d \rangle$. In particular, $f_c \circ \sigma(w) = w$, so σ is injective.

The image of σ is a c -definable set T containing $\langle b, d \rangle$ on which f_c is injective. It follows that $\dim(T) = n + 1$. Because I and Y were strongly I -internal then so is $f_c(T)$, and we have $\dim(f_c(T)) = n + 1$. This contradicts the maximality of dimension of Y . \square

Thus $\langle b, d \rangle$ is a cluster point of $f_c^{-1}(bc - d)$. It now follows that there exists an open interval $J = (b, b') \subseteq I$ (or (b', b)) such that for every $x \in J$ there is $z \in Y$ with $xc - z = bc - d$. In particular, the map $x \mapsto xc - (bc - d)$ sends J injectively into Y . We now apply Lemma 2.6 and obtain b_1 , $b < b_1 < b'$, such that $\dim(b_1cd/b) = 1 + 2n$. In particular, $\dim(cd/b_1b) = 2n$.

Our assumption is that for every x in the interval $J_1 = (b, b_1)$, we have $(x - b)c + d \in Y$. This is a first order property of c, d over the parameters bb_1 . Thus there is a bb_1 -definable set $S \subseteq Y^2$, with $\langle c, d \rangle \in S$, such that for every $\langle y, z \rangle \in S$ and $x \in J_1$, we have $(x - b)y + z \in Y$. It follows that $\dim(S) = 2n$ and $\langle c, d \rangle$ is a generic point in S over bb_1 . We now replace J with J_1 . \square

Proposition 6.4. The set I is neither a subset of K/O nor a subset of Γ .

Proof. Assume for a contradiction that I is either a subset of K/O or of Γ , and denote these by V . In both cases V has an underlying ordered group structures (which is in fact an ordered \mathbb{Q} -vector space). In order to distinguish it from the additive structure of the field we denote here the additive structure on V by $\langle V; \oplus \rangle$. We still use $+$, \cdot to denote the operations of \mathcal{F} .

Let $\langle b, c, d \rangle \in I \times Y^2$ be such that $\dim(b, c, d/\emptyset) = 2n + 1$. We fix $J \subseteq I$ and $S \subseteq Y^2$ as provided by Proposition 6.3. For simplicity we assume that all these sets are definable over \emptyset . For $\langle x, y, z \rangle \in J \times S$, we let $F(x, y, z) = (x - b)y + z$. Our choice of S and J assures that F takes values in Y .

Using cell decomposition in either K/O or Γ we may assume that S is an open subset of I^s .

Claim. There exists an additive (with respect to \oplus) function $L : V^{1+2n} \rightarrow V^s$ and $e \in V$, such that $F(x, y, z) = L(x, y, z) + e$ on a definable subset of $J \times S$ of the same dimension.

Proof. The claim follows from Corollary 5.8 if V is K/O . If T is exponential then by Lemma 3.5, Γ is definably isomorphic to K/O , so the claim holds for Γ as well. If T is power bounded then by Fact 3.3, Γ with its induced structure is an o-minimal vector space over the field of definable exponents. In that case the result follows from quantifier elimination for ordered vector spaces. \square_{Claim}

For simplicity of notation we still denote the set provided by the last claim as $J \times S$.

By the definition of the function F in the field \mathcal{F} , for every $\langle y_1, z_1 \rangle \neq \langle y_2, z_2 \rangle$, there is at most one x such that $F(x, y_1, z_1) = F(x, y_2, z_2)$.

Fix $x_0 \in J \subseteq V$ such that $x_0 \neq 0_V$. The map $\langle y, z \rangle \mapsto F(x_0, y, z)$ sends $Z_1 \times Z_2$ into Y . Since $\dim(Z_1 \times Z_2) = 2n > n$, it follows that the map cannot be injective and hence there are $\langle y_1, z_1 \rangle \neq \langle y_2, z_2 \rangle$ for which $F(x_0, y_1, z_1) = F(x_0, y_2, z_2)$.

It follows that $L(x_0, y_1, z_1) = L(x_0, y_2, z_2)$, hence $L(x_0, y_1, z_1) \ominus L(x_0, y_2, z_2) = 0$. The function $T : x \mapsto L(x, y_1, z_1) \ominus L(x, y_2, z_2) : V \rightarrow V^s$ is an additive function from V into V^s , and $x_0 \in \ker(T)$. It follows that $\mathbb{Z}x_0 \subseteq \ker(T)$, so by weak o-minimality, $\ker(T)$ is a convex subgroup of (V, \oplus) . Thus T is locally constant. This implies that $F(x, y_1, z_1) = F(x, y_2, z_2)$ for infinitely many x , a contradiction. \square

From now on we assume that $I \subseteq K$ or $I \subseteq k$.

7. FIELDS INTERPRETABLE IN T_{conv}

We fix Y a definable subset of \mathcal{F} strongly internal to I , a subset of K or k , and of maximal \mathcal{I} -dimension. As in the previous section, we identify Y with a subset of I^s , and equip it with the topology induced from the weakly o-minimal topology of I^s . For simplicity we assume that I and Y are \emptyset -definable. As before, the argument below does not require T to be power bounded.

7.1. Infinitesimal neighborhoods. We introduce the notion of infinitesimals with respect to the ambient weakly o-minimal structures $(K, \text{ or } k)$:

Given $a \in I$, the *infinitesimal neighborhood of a* (with respect to the ambient weakly o-minimal structure \mathcal{I}) is the partial type over \mathcal{M} consisting of all \mathcal{M} -definable \mathcal{I} -open sets containing a . Equivalently, it is the collection $\{c < x < d : c < a < d, c, d \in I\}$. We

denote it by $\nu(a)$. For $a = (a_1, \dots, a_n) \in I^n$, we let $\nu(a) = \nu(a_1) \times \dots \times \nu(a_n)$. Note that $\nu(a)$ is a partial type in the language \langle (in either K or k).

For a definable $Y \subseteq I^s$, and $a \in Y$, we let $\nu_Y(a) = Y \cap \nu(a)$. In the case where $I \subseteq k$, Y is definable in the induced o-minimal language on k and therefore the type $\nu_Y(a)$ is given by \mathcal{L}_{ind} -formulas. As we now note, this is also the case when $I \subseteq K$:

Remark 7.1. Assume that $I \subseteq K$, $Y \subseteq I^s$ and a is generic in Y . By Lemma 3.6, there is an \mathcal{L} -definable set T and an open box $B \ni a$ such that $B \cap Y$ is \mathcal{L} -definable. Thus, the type $\nu_Y(a)$ is given by \mathcal{L} -formulas.

Finally, note that if $f : I^r \rightarrow I^s$ is an \mathcal{I} -definable function continuous at $a \in I^r$, then $f(\nu(a)) \subseteq \nu(f(a))$.

7.2. The infinitesimal subgroup of $\langle \mathcal{F}, + \rangle$ associated to Y . Throughout $+$, $-$, \cdot and $()^{-1}$ denote the operations in \mathcal{F} .

Lemma 7.2. *If d is generic in Y then $\nu_Y(d) - d$ is a type definable subgroup of $(\mathcal{F}, +)$. Moreover, for every d_1, d_2 both generic in Y over \emptyset , we have $\nu_Y(d_1) - d_1 = \nu_Y(d_2) - d_2$.*

Proof. We fix any $b \in I$ and $c \in Y$ such that $\dim(b, c, d) = 2n + 1$ and apply Proposition 6.3. It follows that the b -definable set

$$\{\langle x, y, z \rangle \in I \times Y \times Y : (x - b)y + z \in Y\}$$

has dimension $2n + 1$ and contains a set of the form $J \times S$ with J an interval (b, b') , $\dim(S) = 2 \dim Y$ and $\langle c, d \rangle$ generic in S over bb' .

Apply Lemma 2.6 to obtain $b_1 \in J$ such that $\dim(b_1, c, d/bb') = 2n + 1$. Let $Y' = (b_1 - b)Y$ and $c' = (b_1 - b)c$. Note that Y' is strongly internal to I and that c' is interdefinable with c over b_1, b . Therefore $\dim(c', d/b_1, b) = 2n$, i.e., $\langle c', d \rangle$ is generic in $Y' \times Y$ over $b_1 b$. Since \mathcal{I} is weakly o-minimal it follows (see [16, Theorem 4.8, Theorem 4.11]) that the function $(y, z) \mapsto y + z$, from $Y' \times Y \subseteq I^{2s}$ into $Y \subseteq I^s$, is I -continuous at $\langle c', d \rangle$. In particular, it sends the types $\nu_{Y'}(c') \times \nu_Y(d)$ into $\nu_Y(c' + d)$.

We now work in an elementary extension of \mathcal{M} and realize the various infinitesimal types there.

The function $y + z$ is injective in each coordinate and therefore for every $y \in \nu_{Y'}(c')$ the map $z \mapsto y + z$ induces a bijection between $\nu_Y(d)$ and $\nu_Y(c' + d)$. Similarly, for every $z \in \nu_Y(d)$, the function $x \mapsto x + z$ induces a bijection of $\nu_Y(c')$ and $\nu_Y(c' + d)$. Finally, for every $w \in \nu_Y(c' + d)$ the function $y \mapsto w - y$ induces a bijection between $\nu_{Y'}(c')$ and $\nu_Y(d)$.

This shows that for every $y \in \nu_{Y'}(c')$ and $w \in \nu_Y(c' + d)$ there exists $z \in \nu_Y(d)$ such that $w = y + z$. Similarly we obtain that $\nu_{Y'}(c') - c' = \nu_Y(d) - d = \nu_Y(c' + d) - (c' + d)$ and they are all subgroups of $(\mathcal{F}, +)$. E.g., if $x, y \in \nu(d) - d$ then $x + c' \in \nu_Y(c')$, $y + d \in \nu_Y(d)$ and therefore $(x + y + c' + d) \in \nu_Y(c' + d)$, so that $x + y \in \nu_Y(c' + d) = \nu_Y(d)$. Closure of $\nu_Y(d)$ under inverses is proved similarly.

In particular, the above shows that whenever $\langle c, d \rangle$ is generic in $Y \times Y$ then $\nu_Y(c) - c = \nu_Y(d) - d$. Since either $Y \subseteq K^s$ or $Y \subseteq k^s$ we know that Y is definable in a weakly o-minimal structure with definable Skolem functions. Therefore, given any $d_1, d_2 \in Y$, each generic in Y , we can find, using Corollary 2.7 $c \in Y$ such that both $\langle c, d_1 \rangle$ and $\langle c, d_2 \rangle$ are generic in $Y \times Y$. It follows that $\nu_Y(d_1) - d_1 = \nu_Y(c) - c = \nu_Y(d_2) - d_2$. \square

We may thus associate to every strongly I -internal Y of maximal dimension a type-definable subgroup ν_Y of $(\mathcal{F}, +)$, which is any of $\nu_Y(c) - c$ for c generic in Y . We now conclude:

Lemma 7.3. *For every Y_1, Y_2 strongly internal to I of maximal dimension, we have $\nu_{Y_1} = \nu_{Y_2}$, as partial types in \mathcal{F} .*

Proof. Let Y be the disjoint union of Y_1 and Y_2 (both embedded in some I^r). Then Y is strongly internal to I and thus for every generic of Y_1 or of Y_2 can be viewed as a generic of Y . The result follows from Lemma 7.2. \square

We let $\nu_{\mathcal{I}}$ denote the type-definable subgroup ν_Y of $(\mathcal{F}, +)$ associated with (any) strongly I -internal set $Y \subseteq F$ of maximal dimension. We still work with a fixed such Y , but now replace it with $Y - a$ for a generic in Y , so that $\nu_{\mathcal{I}} = \nu_Y(0)$. Because $\nu_Y(0)$ is an additive subgroup of \mathcal{F} , there is a definable $Y' \subseteq Y$, with $\nu_Y(0) \vdash Y'$, such that $-Y' \subseteq Y$ and $Y' + Y' \subseteq Y$. We replace Y with $Y' \cap -Y'$, so now $-Y = Y$ and $Y + Y$ is still \mathcal{I} -internal.

Our next goal is to show that the field \mathcal{F} equals to the set $Y \cdot (Y \setminus \{0\})^{-1}$ (thus interpretable in \mathcal{I}) and that in fact, it can be made definable in the o-minimal language of either K or k (without the valuation). For that purpose we apply several times ideas from Marikova's work, [17]. We start with a direct analogue of Lemma 2.10 in her article.

Lemma 7.4. *Assume that $a \in Y$ is generic over the parameters defining Y . Then the functions $(x, y, z) \mapsto x - y + z$ and $(x, y, z) \mapsto -x + y - z$ are \mathcal{I} -continuous at (a, a, a) . Furthermore, if $I \subseteq K$ then the functions are \mathcal{L} -definable in a neighborhood of (a, a, a) .*

Proof. Fix $c \in Y$ such that $\langle c, a \rangle$ is generic in Y^2 . By our assumptions on Y , $c+a, c-a, -a$ and $-c$ are all in I^s . We view the map $x - y + z$ via a composition of 4 maps, and use repeatedly the weakly o-minimal fact that definable functions are continuous at generic points of their domains and, in the case that $I \subseteq K$, use Lemma 3.1(4) which says that \mathcal{L}_{conv} -definable functions coincide with \mathcal{L} -definable functions on neighborhoods of generic points of their domain.

$$\begin{aligned} \varphi_1 & : (x, y, z) \mapsto (c + x, -y, z) : Y^3 \rightarrow Y^3, \\ \varphi_2 & : (x, y, z) \mapsto (x + y, z) : Y^3 \rightarrow Y^2, \\ \varphi_3 & : (x, y) \mapsto (-c, x + y) : Y^2 \rightarrow Y \times Y, \\ \varphi_4 & : (x, y) \mapsto x + y : Y \times Y \rightarrow Y. \end{aligned}$$

Since a is generic in Y , the function $x \mapsto -x$ is continuous near a and since $\dim(c, a) = 2n$, the function $x \mapsto c + x$ is continuous near a . Thus, φ_1 is continuous near (a, a, a) and sends (a, a, a) to $(c + a, -a, a)$. We now have $(c + a, -a)$ generic in Y^2 thus φ_2 is continuous near $(c + a, -a, a)$ and sends it to (c, a) .

We proceed in the same way, and at each step use the fact that we work near a generic point, to conclude that φ_i is continuous locally. Finally, $x - y + z = \varphi_4 \varphi_3 \varphi_2 \varphi_1(x, y, z)$ is continuous near (a, a, a) .

In the case that I is a subset of K , the same argument shows that $x - y + z$ and $-x + y - z$ is \mathcal{L} -definable near (a, a, a) . Applying this result to the point $(-a, -a, -a)$ we get the same result for $-x + y - z$. \square

We now proceed with our proof:

Proposition 7.5. (1) *For every nonzero $c \in \mathcal{F}(\mathcal{M})$, the partial types $c \cdot \nu_{\mathcal{I}}$ and $\nu_{\mathcal{I}}$ are equal.*

(2) *The set of realizations of $\nu_{\mathcal{I}}$ in an elementary extension is a subring of \mathcal{F} .*

Proof. To see (1), notice that for every nonzero $c \in \mathcal{F}$, the set cY is also strongly internal, of the same dimension, hence by Lemma 7.3, $c\nu_Y = \nu_{cY} = \nu_Y$, (as partial types).

(2). We first argue in \mathcal{M} and fix a generic in Y . Since $a\nu_Y = \nu_Y$, there is, by compactness, a definable subset $Y_1 \subseteq Y$ such $aY_1 \subseteq Y$. We fix $b \in Y_1$ such that $\langle a, b \rangle$ is generic in Y^2 . Since $\nu_Y(a) - a$ does not depend on the choice of a it suffices to show that $(\nu_Y(a) - a)(\nu_Y(b) - b) = \nu_Y(ab) - ab$.

We now work in an elementary extension of \mathcal{M} , with realizations of the various partial types. For any $x \in \nu_Y(a)$ and $y \in \nu_Y(b)$ we have $(x - a)(y - b) = xy - xb - ay + ab$. Since $\langle a, b \rangle$ is generic in Y^2 , \mathcal{F} -multiplication is continuous near $\langle a, b \rangle$, hence $xb, ay, xy \in \nu_Y(ab)$. Since a, b are independent generic also ab is generic in Y . Thus by Lemma 7.4, $xy - xb + ab \in \nu_Y(-ab)$. Thus, $(x - a)(y - b) \in \nu_Y(-ab) + ab = \nu(\mathcal{I})$. \square

We can now conclude the proof of our main theorem:

Theorem 7.6. *The field \mathcal{F} is definably isomorphic to a definable field in k_{ind} or to an \mathcal{L} -definable field in K .*

Proof. Let $\nu_{\mathcal{I}} = \nu_Y(0)$ be as above. By Proposition 7.5, the partial type $\nu_{\mathcal{I}}$ is invariant under multiplication by scalars from \mathcal{F} . For every $c \in \mathcal{F}$, the set $cY \cap Y$ is infinite (as both sets contain $\nu_{\mathcal{I}}$). Take a nonzero $s \in cY \cap Y$ and then $s = cr$ for some nonzero $r \in Y$ and so $c = s(r^{-1})$. It follows that

$$(1) \quad \mathcal{F} = \{xy^{-1} : x, y \in Y, y \neq 0\} = Y(Y \setminus \{0\})^{-1}.$$

In fact, since $\nu_Y(a) - a = \nu_{\mathcal{I}}$ for every generic $a \in Y$, the above shows that for two generic elements $a, b \in Y$, we have $\mathcal{F} = (Y - a) \cdot ((Y \setminus \{b\}) - b)^{-1}$. Moreover, we may replace Y by any relatively open neighborhoods $U, V \subseteq Y$ of a and b respectively, and then $\mathcal{F} = (U - a)(V^* - b)^{-1}$, where $V^* = V \setminus \{b\}$.

Because Y is strongly internal to I , the above already implies that \mathcal{F} is interpretable in \mathcal{I} .

Our standing assumption is: $I \subseteq K$ or $I \subseteq k$. When $I \subseteq k$ we can use elimination of imaginaries in the o-minimal structure k_{ind} to deduce that \mathcal{F} is definable in k_{ind} . So we are left with the case $I \subseteq K$, and we want to prove that \mathcal{F} is definably isomorphic to an \mathcal{L} -definable field in K .

We fix $\langle a, b \rangle$ generic in Y^2 such that $a \cdot b \in Y$. By (1) and the comment right after it, for every K -neighborhoods $U \ni a$ and $V \ni b$, we have $(U - a)(V^* - b)^{-1} = \mathfrak{F}$. We define on $U \times V^*$ the following equivalence relation: $\langle x_1, y_1 \rangle \sim \langle x_2, y_2 \rangle$ if

$$(x_1 - a)(y_1 - b)^{-1} = (x_2 - a)(y_2 - b)^{-1} \Leftrightarrow (x_1 - a)(y_2 - b) = (x_2 - a)(y_1 - b).$$

Note that there is a definable bijection between $U \times V^* / \sim$ and \mathcal{F} , given by $[\langle x, y \rangle] \mapsto (x - a)(y - b)^{-1}$.

Claim 7.7. *There are \mathcal{L} -definable $U \ni a$ and $V \ni b$ such that \sim is \mathcal{L} -definable on $U \times V^*$, and $U \times V^* / \sim$ has an \mathcal{L} -definable set of representatives.*

Proof. Unravelling the definition of \sim , we obtain:

$$\langle x_1, y_1 \rangle \sim \langle x_2, y_2 \rangle \Leftrightarrow x_1y_2 - ay_2 - x_1b = x_2y_1 - ay_1 - x_2b.$$

The function $\langle x, y \rangle \mapsto xy$ is \mathcal{L} -definable near $\langle a, b \rangle$ because of genericity. Thus, $y \mapsto ay$ is \mathcal{L} -definable near b and $x \mapsto xb$ is \mathcal{L} -definable near a . Finally, by Lemma 7.4, the map

$\langle x, y, z \rangle \mapsto -x + y - z$ is \mathcal{L} -definable near the point (ab, ab, ab) . It follows that the function $\langle x, y \rangle \mapsto xy - ay - xb$ is \mathcal{L} -definable near $\langle a, b \rangle$, thus \sim is an \mathcal{L} -definable relation near $\langle a, b \rangle$.

We fix \mathcal{L} -definable relatively open neighbourhoods $U, V \subseteq Y$ of a and b , respectively, such that the restriction of \sim to $U \times V^*$ is \mathcal{L} -definable. We require further, using Lemma 7.4, that the functions $(x, y, z) \mapsto x - y + z$ and $(x, y, z) \mapsto -x + y - z$ are \mathcal{L} -definable on $U \cdot V$. Using definable choice in o-minimal structures we can thus find an \mathcal{L} -definable set of representatives, call it S , for \sim . \square

We have so far a definable bijection between \mathfrak{F} and S , an \mathcal{L} -definable set in K^n . Namely, \mathcal{F} is definably isomorphic to an \mathcal{L}_{conv} -definable field (as the field operations might still be definable in \mathcal{L}_{conv}).

Our goal is to show that \mathcal{F} is definably isomorphic to an \mathcal{L} -definable field. One approach for doing that is by noticing that the proof of the analogous result, [1, Theorem 4.2], in the case of real closed valued fields, works word-for-word for T_{conv} . However, for the sake of completeness we give a different, self-contained proof that we can endow S with \mathcal{L} -definable field operations making \mathcal{F} into a \mathcal{L} -definable field. We first need:

Claim 7.8. *Given $\langle x, y \rangle \in (U \setminus \{a\}) \times (V \setminus \{b\})$, for all Y -neighborhoods $U_1 \ni a$ and $V_1 \ni b$ sufficiently small, and for all $a_1 \in U_1 \setminus \{a\}$ and $b_1 \in V_1 \setminus \{b\}$ there are unique $x_1 \in U_1$ and unique $y_1 \in V_1$ such that*

$$\langle x, y \rangle \sim \langle a_1, y_1 \rangle \sim \langle x_1, b_1 \rangle.$$

Moreover, the two families of functions $F_{x,y}(b_1) = x_1$ and $G_{x,y}(a_1) = y_1$ are \mathcal{L} -definable, as $\langle x, y \rangle$ varies.

Proof. The uniqueness is clear from the definition. Let $c = (x - a)^{-1}(y - b)$. The sets $c(Y - a)$ and $(Y - b)$ both contain $\nu_{\mathcal{L}}$. Thus, for $a_1 \in \nu_Y(a)$ there exists $y_1 \in \nu_Y(b)$ such that $c(a_1 - a) = y_1 - b$. It now follows that $(x - a)(y - b)^{-1} = (a_1 - a)(y_1 - b)^{-1}$, so $\langle x, y \rangle \sim \langle a_1, y_1 \rangle$.

Similarly, there exists $x_1 \in \nu_Y(a)$ such that $\langle x, y \rangle \sim \langle x_1, b_1 \rangle$. The existence of U_1, V_1 follows by compactness. Because S is \mathcal{L} -definable, the map $\langle x, y, b_1 \rangle \mapsto x_1$ and the map $\langle x, y, a_1 \rangle \mapsto y_1$ are \mathcal{L} -definable on their appropriate domains. \square

We can now show that the field operations, as induced from \mathcal{F} on the \mathcal{L} -definable set S , are \mathcal{L} -definable. Given $U \ni a$ and $V \ni b$, we let $U^* = U \setminus \{a\}$ and $V^* = V \setminus \{b\}$.

The definability of addition.

For $\langle x_1, y_1 \rangle, \langle x_2, y_2 \rangle, \langle x_3, y_3 \rangle \in S$, let

$$M_+((x_1, y_1), (x_2, y_2)) = \langle x_3, y_3 \rangle \text{ if } (x_1 - a)(y_1 - b)^{-1} + (x_2 - a)(y_2 - b)^{-1} = (x_3 - a)(y_3 - b)^{-1}.$$

If $x_1 = a$ then $M_+((x_1, y_1), (x_2, y_2)) = \langle x_2, y_2 \rangle$. Similarly, $M_+((x_1, y_1), (a, y_2)) = \langle x_1, y_1 \rangle$. We thus assume that $x_1 \in U^*$ and $y_1 \in V^*$.

By Claim 7.8, we can replace $\langle x_2, y_2 \rangle$ with $\langle F_{x_2, y_2}(y_1), y_1 \rangle$ and $\langle x_3, y_3 \rangle$ with $\langle F_{x_3, y_3}(y_1), y_1 \rangle$, uniformly and \mathcal{L} -definably in the parameters. So we may assume that $y_1 = y_2 = y_3$. Setting $x'_i := F_{x_i, y_i}(y_1)$ we get

$$M_+((x_1, y_1), (x_2, y_1)) = \langle x_3, y_1 \rangle \Leftrightarrow (x'_1 - a) + (x'_2 - a) = (x_3 - a) \Leftrightarrow x'_1 - x'_3 + x_2 = a.$$

By the choice of U , we may apply Lemma 7.4, to get that this is an \mathcal{L} -definable relation.

The definability of multiplication.

For $\langle x_1, y_1 \rangle, \langle x_2, y_2 \rangle, \langle x_3, y_3 \rangle \in S$, let

$$M_\bullet((x_1, y_1), (x_2, y_2)) = \langle x_3, y_3 \rangle \text{ if } (x_1 - a)(y_1 - b)^{-1} \cdot (x_2 - a)(y_2 - b)^{-1} = (x_3 - a)(y_3 - b)^{-1}.$$

To see that M_\bullet is \mathcal{L} -definable, we first fix $a_0 \in U^*$ and $b_0 \in V^*$ such that $a_0 - a = b_0 - b$. Using F , as above, we can find x'_2 such that $\langle x'_2, y_3 \rangle \sim \langle x_2, y_2 \rangle$. Thus

$$M_\bullet((x_1, y_1), (x_2, y_2)) = \langle x_3, y_3 \rangle \Leftrightarrow (x_1 - a)(y_1 - b)^{-1} \cdot (x'_2 - a) = x_3 - a.$$

We can now change, \mathcal{L} -definably, $\langle x_1, y_1 \rangle$ in a similar way to an equivalent $\langle x_3, y'_1 \rangle$ so that

$$M_\bullet((x_1, y_1), (x_2, y_2)) = \langle x_3, y_3 \rangle \Leftrightarrow (y'_1 - b)^{-1} \cdot (x'_2 - a) = 1.$$

We now use the fact that $(a_0 - a)(b_0 - b)^{-1} = 1$ to conclude:

$$M_\bullet((x_1, y_1), (x_2, y_2)) = \langle x_3, y_3 \rangle \Leftrightarrow \langle x'_2, y'_1 \rangle \sim \langle a_0, b_0 \rangle.$$

Thus M_\bullet is \mathcal{L} -definable, concluding the proof of the theorem. \square

For the sake of clarity we now sum up everything done up until this point to get a complete proof of Theorem 1:

Corollary 7.9. *Let T be an o-minimal power bounded expansion of a field, T_{conv} a T -convex expansion of T . Any field \mathcal{F} which is interpretable in $K \models T_{conv}$ is definably isomorphic to one of K , $K(\sqrt{-1})$, k , or $k(\sqrt{-1})$*

Proof. Fix some interval $J \subseteq K$ and a definable equivalence relation E on J such that J/E is in definable bijection with an infinite subset of \mathcal{F} , as provided by Lemma 4.1. Shrink J further to obtain a definable bijection between the weakly o-minimal structure J/E and an infinite interval in K , in k , in Γ or in K/O (Theorem 4.2). Apply Proposition 6.4 to eliminate the two last cases. So we are reduced to the case where I is in definable bijection with either k or K and use Theorem 7.6 to get that \mathcal{F} definable in either the o-minimal structure k_{ind} or in the o-minimal structure K (restricted to \mathcal{L}). This allows us to use the main result of [20] to complete the proof. \square

REFERENCES

- [1] Martin Bays and Ya'acov Peterzil. Definability in the group of infinitesimals of a compact Lie group. *Confluentes Math.*, 11(2):3–23, 2019.
- [2] Artem Chernikov and Sergei Starchenko. Regularity lemma for distal structures. *J. Eur. Math. Soc. (JEMS)*, 20(10):2437–2466, 2018.
- [3] Clifton Ealy, Deirdre Haskell, and Jana Maříková. Residue field domination in real closed valued fields. *Notre Dame J. Form. Log.*, 60(3):333–351, 2019.
- [4] Yarit Halevi, Assaf Hasson and Ya'acov Peterzil. Fields interpretable in P -minimal fields. Math Arxiv, 2021
- [5] Deirdre Haskell, Ehud Hrushovski, and Dugald Macpherson. Definable sets in algebraically closed valued fields: elimination of imaginaries. *J. Reine Angew. Math.*, 597:175–236, 2006.
- [6] Deirdre Haskell, Ehud Hrushovski, and Dugald Macpherson. *Stable domination and independence in algebraically closed valued fields*, volume 30 of *Lecture Notes in Logic*. Association for Symbolic Logic, Chicago, IL, 2008.
- [7] Deirdre Haskell, Ehud Hrushovski, and Dugald Macpherson. Unexpected imaginaries in valued fields with analytic structure. *J. Symbolic Logic*, 78(2):523–542, 2013.
- [8] Nadja Hempel and Daniel Palacín. Division rings with ranks. *Proc. Amer. Math. Soc.*, 146(2):803–817, 2018.

- [9] Jan E. Holly. Prototypes for definable subsets of algebraically closed valued fields. *J. Symbolic Logic*, 62(4):1093–1141, 1997.
- [10] Ehud Hrushovski and Silvain Rideau-Kikuchi. Valued fields, metastable groups. *Selecta Math. (N.S.)*, 25(3):Paper No. 47, 58, 2019.
- [11] Will Johnson. Dp-finite fields ii: the canonical topology and its relation to henselianity, 2019.
- [12] Will Johnson. Dp-finite fields i: infinitesimals and positive characteristic, 2020.
- [13] Will Johnson. Dp-finite fields vi: the dp-finite shelah conjecture, 2020.
- [14] Itay Kaplan, Alf Onshuus, and Alexander Usvyatsov. Additivity of the dp-rank. *Trans. Amer. Math. Soc.*, 365(11):5783–5804, 2013.
- [15] James Loveys and Ya’acov Peterzil. Linear o-minimal structures. *Israel J. Math.*, 81(1-2):1–30, 1993.
- [16] Dugald Macpherson, David Marker, and Charles Steinhorn. Weakly o-minimal structures and real closed fields. *Trans. Amer. Math. Soc.*, 352(12):5435–5483 (electronic), 2000.
- [17] Jana Maříková. Type-definable and invariant groups in o-minimal structures. *J. Symbolic Logic*, 72(1):67–80, 2007.
- [18] T. Mellor. Imaginaries in real closed valued fields. *Ann. Pure Appl. Logic*, 139(1-3):230–279, 2006.
- [19] Chris Miller and Sergei Starchenko. A growth dichotomy for o-minimal expansions of ordered groups. *Trans. Amer. Math. Soc.*, 350(9):3505–3521, 1998.
- [20] Margarita Otero, Ya’acov Peterzil, and Anand Pillay. On groups and rings definable in o-minimal expansions of real closed fields. *Bull. London Math. Soc.*, 28(1):7–14, 1996.
- [21] Ya’acov Peterzil, Anand Pillay, and Sergei Starchenko. Simple algebraic and semialgebraic groups over real closed fields. *Trans. Amer. Math. Soc.*, 352(10):4421–4450, 2000.
- [22] Anand Pillay. On fields definable in \mathbf{Q}_p . *Arch. Math. Logic*, 29(1):1–7, 1989.
- [23] Bruno Poizat. MM. Borel, Tits, Zilber et le Général Nonsense. *J. Symbolic Logic*, 53(1):124–131, 1988.
- [24] Pierre Simon. Dp-minimality: invariant types and dp-rank. *J. Symb. Log.*, 79(4):1025–1045, 2014.
- [25] Pierre Simon. *A guide to NIP theories*, volume 44 of *Lecture Notes in Logic*. Association for Symbolic Logic, Chicago, IL; Cambridge Scientific Publishers, Cambridge, 2015.
- [26] Pierre Simon and Erik Walsberg. Tame topology over dp-minimal structures. *Notre Dame J. Form. Log.*, 60(1):61–76, 2019.
- [27] James Michael Tyne. *T-levels and T-convexity* Ph.D. Thesis, Univerisyt of Illinois at Urbana-Champaign, 2003.
- [28] Lou van den Dries. *T-convexity and tame extensions. II.* *J. Symbolic Logic*, 62(1):14–34, 1997.
- [29] Lou van den Dries. Correction to: “*T-convexity and tame extensions. II*” [*J. Symbolic Logic* **62** (1997), no. 1, 14–34; MR1450511 (98h:03048)]. *J. Symbolic Logic*, 63(4):1597, 1998.
- [30] Lou van den Dries. *Tame topology and o-minimal structures*, volume 248 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1998.
- [31] Lou van den Dries and Adam H. Lewenberg. *T-convexity and tame extensions.* *J. Symbolic Logic*, 60(1):74–102, 1995.

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, BE’ER-SHEVA 84105, ISRAEL

Email address: hassonas@math.bgu.ac.il

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA, ISRAEL

Email address: kobi@math.haifa.ac.il