

On zeros of the Riemann zeta function

Xiaolong Wu

Ex. Institute of Mathematics, Chinese Academy of Sciences

xwu622@comcast.net

January 20, 2021

Abstract

This paper shows that, in the critical strip, the Riemann zeta function $\zeta(s)$ have the same set of zeros as $F(s) := \int_0^\infty t^{s-1}(e^t + 1)^{-1}dt$, and then discusses the behavior of $F(s)$.

Introduction

Define $\zeta(s)$ as the Riemann zeta function, and define

$$F(s) := \int_0^\infty \frac{t^{s-1}}{e^t + 1} dt, \quad s \in \mathbb{C}. \quad (1)$$

Define the open critical strip by

$$\mathcal{S} := \{s \in \mathbb{C} : 0 < \mathcal{R}(s) < 1\} \quad (2)$$

where $\mathcal{R}(s)$ is the real part of s .

Theorem 1 shows that $\zeta(s)$ and $F(s)$ have the same set of zeros in \mathcal{S} . So we can study the zeros of $F(s)$ instead of $\zeta(s)$, and $F(s)$ looks simpler.

Write $s = a + ib$. It is obvious that

$$F(s) = \int_0^\infty \frac{t^{a-1} \cos(b \log t)}{e^t + 1} dt + i \int_0^\infty \frac{t^{a-1} \sin(b \log t)}{e^t + 1} dt =: F_1(a, b) + iF_2(a, b). \quad (3)$$

So, $F_2(a, b) \neq 0$ means $F(s) \neq 0$. This paper will work on F_2 . We split the integration range of $F_2(a, b)$ into two sub-ranges $[0, R]$ and $[R, \infty)$ for some real $R = e^{2K\pi}$ with integer $K \geq 0, 1 \leq R < \pi$.

Define

$$f(t, a) := \frac{t^{a-1}}{e^t + 1}, \quad g(t) := \frac{1}{e^t + 1}. \quad (4)$$

Theorem 2 shows that

$$\int_0^R f(t, a) \sin(b \log t) dt = - \sum_{n=0}^{\infty} \frac{g^{(n)}(0)b}{n!((n+a)^2 + b^2)} R^{n+a}. \quad (5)$$

Theorems 3 and 4 discuss properties of $g^{(n)}(0)$, especially, its relation with the Bernoulli numbers.

Theorem 5 shows that

$$\int_0^R f(t, a) \sin(b \log t) dt > \frac{R^a}{b(e^R + 1)} - \frac{0.47177R^a}{b^2}. \quad (6)$$

Theorems 6 to 10 study the behavior of

$$\int_R^{\infty} f(t, a) \sin(b \log t) dt.$$

In particular, Theorem 6 shows

$$\int_R^{\infty} f(t, a) \sin(b \log t) dt = \sum_{k=K}^{\infty} \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} h(t, a, b) \sin(b \log t) dt.$$

where

$$h(t, a, b) := \frac{t^{a-1}}{e^t + 1} - \frac{t^{a-1} e^{a\pi/b}}{e^{te^{\pi/b}} + 1}. \quad (7)$$

Theorem 7 shows, for $c = e^{\pi/b}$,

$$\int_R^{\infty} h(t, a, b) dt = \int_R^{cR} \frac{t^{a-1}}{e^t + 1} dt. \quad (8)$$

Theorem 10 shows, for some range of t, a, b ,

$$h(t, a, b) > 0. \quad (9)$$

Write the gamma function as

$$\Gamma(s) := \int_0^{\infty} \frac{t^{s-1}}{e^t} dt. \quad (10)$$

Main Content

Theorem 1. For $s \in \mathcal{S}$, $\zeta(s) = 0$ if and only if $F(s) = 0$.

Proof. It is well known that, ([Titchmarsh 1986] formula 2.4.1),

$$\zeta(s) = \frac{G(s)}{\Gamma(s)}, \quad (1.1)$$

where

$$G(s) := \int_0^\infty \frac{t^{s-1}}{e^t - 1} dt. \quad (1.2)$$

Since $\Gamma(s)$ has no zero nor pole in \mathcal{S} , $\zeta(s)$ has the same set of zeros as $G(s)$ in \mathcal{S} . It is easy to see, write $u = 2t$,

$$\begin{aligned} G(s) - F(s) &= \int_0^\infty \frac{t^{s-1}}{e^t - 1} dt - \int_0^\infty \frac{t^{s-1}}{e^t + 1} dt \\ &= \int_0^\infty \frac{2t^{s-1}}{e^{2t} - 1} dt = \int_0^\infty \frac{2(u/2)^{s-1}}{e^u - 1} d\frac{u}{2} \\ &= 2^{1-s} \int_0^\infty \frac{u^{s-1}}{e^u - 1} du = 2^{1-s} G(s). \end{aligned} \quad (1.3)$$

This means

$$(1 - 2^{1-s})G(s) = F(s). \quad (1.4)$$

Since $1 - 2^{1-s}$ has no zero nor pole in \mathcal{S} , $F(s)$ has the same set of zeros as $G(s)$, hence as $\zeta(s)$ in \mathcal{S} . \square

Theorem 2. Let $b \geq 100$ be a real, $K > 0$ be an integer, and $R = \exp(2K\pi/b)$ such that $1 \leq R < \pi$. Let $f(t, a)$ and $g(t)$ be defined as in (4).

Then

$$\int_0^R f(t, a) \sin(b \log t) dt = - \sum_{n=0}^{\infty} \frac{g^{(n)}(0)b}{n!((n+a)^2 + b^2)} R^{n+a}. \quad (2.1)$$

Proof. Since $g(t) = (e^t + 1)^{-1}$ has a pole at $t = \pi i$, the radius of convergence of series

$$g(t) = \sum_{n=0}^{\infty} \frac{g^{(n)}(0)}{n!} t^n \quad (2.2)$$

is π . We have

$$\begin{aligned}
\int_0^R \frac{t^{s-1}}{e^t + 1} dt &= \int_0^R t^{s-1} \sum_{n=0}^{\infty} \frac{g^{(n)}(0)}{n!} t^n dt \\
&= \sum_{n=0}^{\infty} \frac{g^{(n)}(0)}{n!} \int_0^R t^{n+s-1} dt \\
&= \sum_{n=0}^{\infty} \frac{g^{(n)}(0)}{n!(n+s)} R^{n+s} \\
&= \sum_{n=0}^{\infty} \frac{g^{(n)}(0)(n+a-ib)}{n!((n+a)^2+b^2)} R^{n+a} e^{ib \log R} \\
&= \sum_{n=0}^{\infty} \frac{g^{(n)}(0)(n+a)}{n!((n+a)^2+b^2)} R^{n+a} - i \sum_{n=0}^{\infty} \frac{g^{(n)}(0)b}{n!((n+a)^2+b^2)} R^{n+a}.
\end{aligned} \tag{2.3}$$

On the other hand, we have

$$\begin{aligned}
\int_0^R \frac{t^{s-1}}{e^t + 1} dt &= \int_0^R \frac{t^{a-1+bi}}{e^t + 1} dt \\
&= \int_0^R \frac{t^{a-1} e^{ib \log t}}{e^t + 1} dt \\
&= \int_0^R \frac{t^{a-1}}{e^t + 1} \cos(b \log t) dt + i \int_0^R \frac{t^{a-1}}{e^t + 1} \sin(b \log t) dt
\end{aligned} \tag{2.4}$$

Equate the imaginary part of (2.3) and (2.4), we get (2.1). \square

Theorem 3. Let $g(t) = (e^t + 1)^{-1}$ as in (4), then

$$g^{(n)}(0) = \frac{1 - 2^{n+1}}{n+1} B_{n+1}, \tag{3.1}$$

where B_n is the n -th Bernoulli number.

Proof. The Bernoulli numbers have a generating function:

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} t^n. \tag{3.2}$$

On the other hand, we have

$$\frac{t}{e^t + 1} = \sum_{n=0}^{\infty} \frac{g^{(n)}(0)}{n!} t^{n+1}. \tag{3.3}$$

Since

$$\frac{t}{e^t - 1} - \frac{t}{e^t + 1} = \frac{2t}{e^{2t} - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} (2t)^n, \quad (3.4)$$

we have

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{g^{(n)}(0)}{n!} t^{n+1} &= \frac{t}{e^t + 1} \\ &= \frac{t}{e^t - 1} - \frac{2t}{e^{2t} - 1} \\ &= \sum_{n=0}^{\infty} \frac{B_n}{n!} t^n - \sum_{n=0}^{\infty} \frac{B_n}{n!} (2t)^n \\ &= \sum_{n=0}^{\infty} \frac{B_n}{n!} (1 - 2^n) t^n \end{aligned} \quad (3.5)$$

Equate coefficients of t^{n+1} , we get

$$\frac{g^{(n)}(0)}{n!} = \frac{B_{n+1}}{(n+1)!} (1 - 2^{n+1}) \quad (3.6)$$

$$g^{(n)}(0) = \frac{1 - 2^{n+1}}{n+1} B_{n+1}. \quad (3.7)$$

□

Theorem 4. (1) $g^{(2m)}(0) = 0$, $\forall m \geq 1$.

(2) $g^{(4m+1)}(0) < 0$, $\forall m \geq 0$.

(3) $g^{(4m-1)}(0) > 0$, $\forall m \geq 1$.

(4)

$$\frac{g^{(2m-1)}(0)}{(2m-1)!} = (-1)^m (1 - 2^{-2m}) \zeta(2m) \frac{2}{\pi^{2m}}, \quad \forall m \geq 1. \quad (4.1)$$

(5)

$$-\frac{g^{(4m+1)}(0)}{(4m+1)!} \pi^2 < \frac{g^{(4m-1)}(0)}{(4m-1)!} < -1.00013814 \frac{g^{(4m+1)}(0)}{(4m+1)!} \pi^2, \quad \forall m \geq 2. \quad (4.2)$$

(6)

$$\frac{2}{\pi^{2m}} < \frac{|g^{(2m-1)}(0)|}{(2m-1)!} = (1 - 2^{-2m}) \zeta(2m) \frac{2}{\pi^{2m}}, \quad \forall m \geq 1. \quad (4.3)$$

(7)

$$g(t) = \frac{1}{2} - \frac{1}{2} \tanh \frac{t}{2}. \quad (4.4)$$

Proof. (1), (2) and (3) easily follow from well-known properties of the Bernoulli numbers.

(4) It is well known that

$$\zeta(2m) = \frac{(-1)^{m+1}(2\pi)^{2m}B_{2m}}{2(2m)!}, \quad \forall m \geq 1. \quad (4.5)$$

Hence by Theorem 3, we have

$$\begin{aligned} g^{(2m-1)}(0) &= \frac{1-2^{2m}}{2m}B_{2m} \\ &= \frac{1-2^{2m}}{2m} \cdot \frac{(-1)^{m+1}2(2m)!\zeta(2m)}{(2\pi)^{2m}} \\ &= \frac{(-1)^m 2(1-2^{-2m})(2m-1)!\zeta(2m)}{\pi^{2m}}, \quad \forall m \geq 1. \end{aligned} \quad (4.6)$$

(5) By equation (4.1) we have

$$\begin{aligned} \frac{g^{(4m-1)}(0)/(4m-1)!}{-g^{(4m+1)}(0)/(4m+1)!} &= \frac{(1-2^{-4m})\zeta(4m)2/\pi^{4m}}{(1-2^{-4m-2})\zeta(4m+2)2/\pi^{4m+2}} \\ &= \frac{(1-2^{-4m})\zeta(4m)}{(1-2^{-4m-2})\zeta(4m+2)}\pi^2, \quad \forall m \geq 1. \end{aligned} \quad (4.7)$$

The last fraction in (4.7) is a decreasing function in m and approaches 1 when $m \rightarrow \infty$. Hence, we have

$$1 < \frac{(1-2^{-4m})\zeta(4m)}{(1-2^{-4m-2})\zeta(4m+2)} < \frac{(1-2^{-8})\zeta(8)}{(1-2^{-10})\zeta(10)} < 1.00013814, \quad \forall m \geq 2. \quad (4.8)$$

(6) By [Ge 2012] Theorem 1.1,

$$\frac{2(2m)!}{\pi^{2m}(2^{2m}-1)} < |B_{2m}| = \zeta(2m)\frac{2(2m)!}{(2\pi)^{2m}}. \quad (4.9)$$

Substitute $B_{2m} = 2mg^{(2m-1)}(0)/(1-2^{2m})$, we get

$$\frac{2(2m)!}{\pi^{2m}(2^{2m}-1)} < \left| \frac{2mg^{(2m-1)}(0)}{1-2^{2m}} \right| = \zeta(2m)\frac{2(2m)!}{(2\pi)^{2m}}. \quad (4.10)$$

$$\frac{2}{\pi^{2m}} < \left| \frac{g^{(2m-1)}(0)}{(2m-1)!} \right| = (1-2^{-2m})\zeta(2m)\frac{2}{\pi^{2m}}. \quad (4.11)$$

(7)

$$\frac{1}{e^t+1} - \frac{1}{2} = -\frac{1}{2} \cdot \frac{e^t-1}{e^t+1} = -\frac{1}{2} \tanh \frac{t}{2}. \quad (4.12)$$

□

Theorem 5. Let $K > 0$ be the largest integer such that $R = e^{2K\pi/b} \leq 2$. Let $a \leq 0.1$ and $b \geq 100$. Then

$$-\sum_{n=0}^{\infty} \frac{g^{(n)}(0)b}{n!((n+a)^2 + b^2)} R^{n+a} > -\frac{R^a}{b(e^R + 1)} - \frac{0.47177R^a}{b^3}. \quad (5.1)$$

Proof. By maximality of K , we have

$$Re^{2\pi/b} = e^{2K\pi/b} e^{2\pi/b} = e^{2(K+1)\pi/b} > 2. \quad (5.2)$$

Hence

$$R > 2e^{-2\pi/b} > 2 \left(1 - \frac{2\pi}{b}\right) = 2 - \frac{4\pi}{b}. \quad (5.3)$$

$$\begin{aligned} & -\sum_{n=0}^{\infty} \frac{g^{(n)}(0)b}{n!((n+a)^2 + b^2)} R^{n+a} + \frac{R^a}{b(e^R + 1)} \\ &= -\sum_{n=0}^{\infty} \frac{g^{(n)}(0)b}{n!((n+a)^2 + b^2)} R^{n+a} + \frac{R^a}{b} \sum_{n=0}^{\infty} \frac{g^{(n)}(0)}{n!} R^n \\ &= \frac{R^a}{b} \sum_{n=0}^{\infty} \frac{g^{(n)}(0)R^n}{n!} \left(\frac{-b^2}{(n+a)^2 + b^2} + 1 \right) \\ &= \frac{R^a}{b} \sum_{n=0}^{\infty} \frac{g^{(n)}(0)R^n}{n!} \left(\frac{(n+a)^2}{(n+a)^2 + b^2} \right) \\ &= \frac{R^a}{b} \left(\frac{a^2}{2(a^2 + b^2)} - \frac{(1+a)^2 R}{4((1+a)^2 + b^2)} + \frac{(3+a)^2 R^3}{4((3+a)^2 + b^2)} - \frac{(5+a)^2 R^5}{4((5+a)^2 + b^2)} \right) \\ &\quad + \frac{R^a}{b} \sum_{n=7}^{\infty} \frac{g^{(n)}(0)R^n}{n!} \left(\frac{(n+a)^2}{(n+a)^2 + b^2} \right) \\ &> \frac{R^a}{b} \left(-\frac{0.76667}{b^2} + \sum_{m=2}^{\infty} c_m \right), \end{aligned} \quad (5.4)$$

where

$$\begin{aligned} c_m &:= \frac{g^{(4m-1)}(0)R^{4m-1}}{(4m-1)!} \left(\frac{(4m-1+a)^2}{(4m-1+a)^2 + b^2} \right) \\ &\quad + \frac{g^{(4m+1)}(0)R^{4m+1}}{(4m+1)!} \left(\frac{(4m+1+a)^2}{(4m+1+a)^2 + b^2} \right). \end{aligned} \quad (5.5)$$

By Theorem 4, we have $g^{(4m-1)}(0) > 0, g^{(4m+1)}(0)$ and

$$\frac{g^{(4m-1)}(0)}{(4m-1)!} > -\frac{g^{(4m+1)}(0)\pi^2}{(4m+1)!}, \quad (5.6)$$

$$-\frac{g^{(4m+1)}(0)}{(4m+1)!} > \frac{2}{\pi^{4m+2}}. \quad (5.7)$$

Hence

$$\begin{aligned} c_m &> -\frac{g^{(4m+1)}(0)R^{4m-1}}{(4m+1)!} \left(\frac{(4m-1+a)^2}{(4m-1+a)^2+b^2} - \frac{(4m+1+a)^2R^2}{(4m+1+a)^2+b^2} \right) \\ &> \frac{2R^{4m-1}}{\pi^{4m+2}} \left(\frac{(4m+1+a)^2}{(4m+1+a)^2+b^2} \left(\frac{7^2\pi^2}{9^2} - R^2 \right) \right) \\ &= \frac{2}{Rb^2} \left(\frac{R}{\pi} \right)^{4m} \left(\frac{7^2}{9^2} - \frac{R^2}{\pi^2} \right) \left(\frac{(4m+1+a)^2}{(4m+1+a)^2b^{-2}+1} \right) \end{aligned} \quad (5.8)$$

Then

$$\begin{aligned} \sum_{m=0}^{\infty} c_m &> \frac{2}{Rb^2} \left(\frac{7^2}{9^2} - \frac{R^2}{\pi^2} \right) \sum_{m=2}^{\infty} \left(\frac{R}{\pi} \right)^{4m} \left(\frac{(4m+1+a)^2}{(4m+1+a)^2b^{-2}+1} \right) \\ &> \frac{0.1996535370}{b^2} \sum_{m=2}^{\infty} \left(\frac{R}{\pi} \right)^{4m} \left(\frac{9^2}{9^2b^{-2}+1} \right) \\ &> \frac{16.04199633}{b^2} \sum_{m=2}^{\infty} \left(\frac{R}{\pi} \right)^{4m} \\ &= \frac{16.04199633}{b^2} \cdot \frac{(R/\pi)^8}{1-(R/\pi)^4} \\ &> \frac{0.29490227}{b^2} \end{aligned} \quad (5.9)$$

Hence

$$\begin{aligned} &-\sum_{n=0}^{\infty} \frac{g^{(n)}(0)b}{n!((n+a)^2+b^2)} R^{n+a} + \frac{R^a}{b(e^R+1)} \\ &> \frac{R^a}{b} \left(-\frac{0.76667}{b^2} + \frac{0.29490}{b^2} \right) > -\frac{0.47177R^a}{b^3}. \end{aligned} \quad (5.10)$$

□

Remark. Theorem 5 chooses $R \approx 2$ to make the series converge and make the calculation simple.

Theorem 6. Let a, b be reals such that $0 < a < 1, b \geq 100$. Let $K > 0$ be the largest integer such that $R = e^{2K\pi/b} \leq 2$. Then

$$\int_R^\infty f(t, a) \sin(b \log t) dt = \sum_{k=K}^\infty \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} h(t, a, b) \sin(b \log t) dt. \quad (6.1)$$

Proof. Write

$$\int_R^\infty f(t, a) \sin(b \log t) dt = \sum_{k=K}^\infty I_k, \quad (6.2)$$

where

$$I_k := \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} f(t, a) \sin(b \log t) dt. \quad (6.3)$$

Then split the range of I_k in to two sub-intervals according to the sign of $\sin(b \log t)$. That is

$$I_k = I_{k,1} + I_{k,2}, \quad (6.4)$$

where

$$I_{k,1} := \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} f(t, a) \sin(b \log t) dt, \quad (6.5)$$

$$I_{k,2} := \int_{\exp((2k+1)\pi/b)}^{\exp(2(k+1)\pi/b)} f(t, a) \sin(b \log t) dt. \quad (6.6)$$

By changing variable $u = te^{-\pi/b}$ in $I_{k,2}$, we get

$$\begin{aligned} I_{k,2} &= \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} f(ue^{\pi/b}, a) \sin(b \log(ue^{\pi/b})) d(ue^{\pi/b}) \\ &= e^{\pi/b} \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} f(ue^{\pi/b}, a) \sin(b \log u + \pi) du \\ &= -e^{\pi/b} \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} f(ue^{\pi/b}, a) \sin(b \log u) du. \end{aligned} \quad (6.7)$$

Hence

$$\begin{aligned} I_k &= \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} (f(t, a) - e^{\pi/b} f(te^{\pi/b}, a) \sin(b \log t)) dt \\ &= \int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} h(t, a, b) \sin(b \log t) dt \end{aligned} \quad (6.8)$$

□

Theorem 7. Let a, b, R be reals such that $0 < a < 1, b > 0, R \geq 1$. Let $c := e^{\pi/b}$. Then

$$\int_R^\infty h(t, a, b) dt = \int_R^{cR} \frac{t^{a-1}}{e^t + 1} dt. \quad (7.1)$$

$$(c-1) \frac{c^{a-1} R^a}{e^{cR} + 1} < \int_R^\infty h(t, a, b) dt < (c-1) \frac{R^a}{e^R + 1}. \quad (7.2)$$

Proof. Let $0 \leq \alpha < \beta$ be reals, $u = ct, t = c^{-1}u$. Then

$$\int_\alpha^\beta \frac{t^{a-1}}{e^{ct} + 1} dt = \int_{c\alpha}^{c\beta} \frac{c^{1-a} u^{a-1}}{e^u + 1} d(c^{-1}u) = c^{-a} \int_{c\alpha}^{c\beta} \frac{u^{a-1}}{e^u + 1} du. \quad (7.3)$$

Hence

$$\begin{aligned} \int_R^\infty h(t, a, b) dt &= \int_R^\infty \left(\frac{t^{a-1}}{e^t + 1} - \frac{t^{a-1} e^{a\pi/b}}{e^{te^{\pi/b}} + 1} \right) dt \\ &= \int_R^\infty \frac{t^{a-1}}{e^t + 1} dt - c^a \int_R^\infty \frac{t^{a-1}}{e^{ct} + 1} dt \\ &= \int_R^\infty \frac{t^{a-1}}{e^t + 1} dt - c^a \left(c^{-a} \int_{cR}^\infty \frac{t^{a-1}}{e^t + 1} dt \right) \\ &= \int_R^\infty \frac{t^{a-1}}{e^t + 1} dt - \int_{cR}^\infty \frac{t^{a-1}}{e^t + 1} dt \\ &= \int_R^{cR} \frac{t^{a-1}}{e^t + 1} dt \end{aligned} \quad (7.4)$$

Since $t^{a-1}/(e^t + 1)$ is decreasing in t , we have

$$\int_R^{cR} \frac{t^{a-1}}{e^t + 1} dt > (cR - R) \frac{(cR)^{a-1}}{e^{cR} + 1} = (c-1) \frac{c^{a-1} R^a}{e^{cR} + 1}, \quad (7.5)$$

$$\int_R^{cR} \frac{t^{a-1}}{e^t + 1} dt > (cR - R) \frac{R^{a-1}}{e^R + 1} = (c-1) \frac{R^a}{e^R + 1}. \quad (7.6)$$

□

Theorem 8. Let k be an integer, $b \geq 10$ be a real. Define the average

$$A := \frac{1}{e^{(2k+1)\pi/b} - e^{2k\pi/b}} \int_{e^{2k\pi/b}}^{e^{(2k+1)\pi/b}} \sin(b \log t) dt. \quad (8.1)$$

Then

$$\frac{2}{\pi} - \frac{2}{\pi b^2} < A < \frac{2}{\pi}. \quad (8.2)$$

Proof. Substitute $u = b \log t, t = e^{u/b}$, we get

$$\begin{aligned}
\int_{\exp(2k\pi/b)}^{\exp((2k+1)\pi/b)} \sin(b \log t) dt &= \frac{1}{b} \int_{2k\pi/b}^{(2k+1)\pi/b} e^{u/b} \sin u du \\
&= \frac{e^{u/b}}{b(b^{-2} + 1)} \left[\frac{1}{b} \sin u - \cos u \right]_{2k\pi}^{(2k+1)\pi} \\
&= \frac{e^{(2k+1)\pi/b} + e^{2k\pi/b}}{b(b^{-2} + 1)}. \tag{8.3}
\end{aligned}$$

So the average is

$$A = \frac{e^{(2k+1)\pi/b} + e^{2k\pi/b}}{b(b^{-2} + 1)(e^{(2k+1)\pi/b} - e^{2k\pi/b})} = \frac{1 + e^{-\pi/b}}{b(b^{-2} + 1)(1 - e^{-\pi/b})}. \tag{8.4}$$

Since

$$\begin{aligned}
1 + e^{-\pi/b} - \frac{2b}{\pi}(1 - e^{-\pi/b}) &= 1 + \sum_{n=0}^{\infty} \frac{(-\pi)^n}{n!b^n} + \frac{2b}{\pi} \sum_{n=1}^{\infty} \frac{(-\pi)^n}{n!b^n} \\
&= 2 + \sum_{n=1}^{\infty} \frac{(-\pi)^n}{n!b^n} - 2 + 2 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}\pi^n}{(n+1)!b^n} \\
&= \sum_{n=1}^{\infty} \frac{(-\pi)^n}{b^n} \left(\frac{1}{n!} - \frac{2}{(n+1)!} \right) \\
&= \sum_{n=1}^{\infty} \frac{(-\pi)^n}{b^n} \cdot \frac{n-1}{(n+1)!} \\
&= \frac{\pi^2}{3!b^2} - \dots, \tag{8.5}
\end{aligned}$$

we have

$$\begin{aligned}
\frac{2}{\pi} &< \frac{1 + e^{-\pi/b}}{b(1 - e^{-\pi/b})} \\
&< \frac{2}{\pi} + \frac{\pi^2}{3!b^2b(\pi/b - (\pi/b)^2/2)} \\
&= \frac{2}{\pi} + \frac{\pi}{3!b^2(1 - \pi/(2b))} \\
&\leq \frac{2}{\pi} + \frac{\pi}{3!b^2(1 - \pi/20)} \\
&< \frac{2}{\pi} + \frac{\pi}{3!b^2(5/6)} = \frac{2}{\pi} + \frac{\pi}{5b^2}, \quad \forall b \geq 10. \tag{8.6}
\end{aligned}$$

Hence

$$\begin{aligned}
A &< \frac{1}{b^{-2} + 1} \left(\frac{2}{\pi} + \frac{\pi}{5b^2} \right) \\
&< \left(1 - \frac{1}{b^2} + \frac{1}{b^4} \right) \left(\frac{2}{\pi} + \frac{\pi}{5b^2} \right) \\
&= \frac{2}{\pi} + \frac{\pi}{5b^2} - \frac{2}{\pi b^2} - \frac{\pi}{5b^4} + \frac{2}{\pi b^4} + \frac{\pi}{5b^6} \\
&= \frac{2}{\pi} - \frac{0.0083}{b^2} + \frac{0.0083}{b^4} + \frac{0.6283}{b^6} < \frac{2}{\pi}, \tag{8.7}
\end{aligned}$$

$$A > \frac{1}{b^{-2} + 1} \left(\frac{2}{\pi} \right) > \frac{2}{\pi} \left(1 - \frac{1}{b^2} \right). \tag{8.8}$$

□

Theorem 9. Let k be an integer, $b \geq 100$ be a real. Write $t_0 = \exp(2k\pi/b)$, $t_1 = \exp((2k+1)\pi/b)$. Let $h(t)$ be a continuous function and

$$M_1 = \min_{t_0 \leq t \leq t_1} h(t), \quad M_2 = \max_{t_0 \leq t \leq t_1} h(t). \tag{9.1}$$

Then

$$\left(\frac{2}{\pi} - \frac{2}{\pi b^2} \right) \int_{t_0}^{t_1} M_1 dt < \int_{t_0}^{t_1} h(t) \sin(b \log t) dt < \frac{2}{\pi} \int_{t_0}^{t_1} M_2 dt. \tag{9.2}$$

Proof. Since $\sin(b \log t) \geq 0$ in interval $[t_0, t_1]$, we have, by Theorem 8,

$$\begin{aligned}
\int_{t_0}^{t_1} h(t) \sin(b \log t) dt &\geq \int_{t_0}^{t_1} M_1 \sin(b \log t) dt \\
&> M_1 \left(\frac{2}{\pi} - \frac{2}{\pi b^2} \right) (t_1 - t_0) \\
&= \left(\frac{2}{\pi} - \frac{2}{\pi b^2} \right) \int_{t_0}^{t_1} M_1 dt. \tag{9.3}
\end{aligned}$$

The other inequality can be proved similarly. □

Theorem 10. Let t, a, b be reals such that $t \geq 1, 0 < a \leq e/(e+1) = 0.731 \dots$ and $b > 0$. We have

$$h(t, a, b) > 0. \tag{10.1}$$

Proof. We need only to prove that

$$\frac{1}{e^t + 1} - \frac{e^{a\pi/b}}{e^{te^{\pi/b}} + 1} > 0. \quad (10.2)$$

$$\begin{aligned} \frac{1}{e^t + 1} - \frac{e^{a\pi/b}}{e^{te^{\pi/b}} + 1} &= \frac{e^{te^{\pi/b}} + 1 - e^{t+a\pi/b} - e^{a\pi/b}}{(e^t + 1)(e^{te^{\pi/b}} + 1)} \\ &> \frac{e^{t(1+\pi/b)} + 1 - e^{t+a\pi/b} - e^{a\pi/b}}{(e^t + 1)(e^{te^{\pi/b}} + 1)} \\ &= \frac{e^t(e^{t\pi/b} - e^{a\pi/b}) - (e^{a\pi/b} - 1)}{(e^t + 1)(e^{te^{\pi/b}} + 1)} \\ &> \frac{e^t(t\pi/b - a\pi/b) - a\pi/b}{(e^t + 1)(e^{te^{\pi/b}} + 1)} \\ &= \frac{\pi}{b} \cdot \frac{te^t - ae^t - a}{(e^t + 1)(e^{te^{\pi/b}} + 1)} \end{aligned} \quad (10.3)$$

Since $te^t - ae^t - a > 0$ when $t \geq 1, a \leq e/(e+1) = 0.731 \dots$, (10.2) holds. \square

References

- [Ge 2012] Hua-feng Ge. *New Sharp Bounds for the Bernoulli Numbers and Refinement of Becker-Stark Inequalities*. Journal of Applied Mathematics Vol 2012, (2012)
- [Titchmarsh 1986] E.C. Titchmarsh and D.R. Heath Brown. *The Theory of the Riemann Zeta-function*. Oxford University, 2nd edition (1986)

Appendix

$g^{(n)}(0)$ for $n \leq 15$.

$$\begin{aligned} g^{(0)}(0) &= \frac{1}{2}, & g^{(1)}(0) &= -\frac{1}{4}, & g^{(3)}(0) &= \frac{1}{8}, \\ g^{(5)}(0) &= -\frac{1}{4}, & g^{(7)}(0) &= \frac{17}{16}, & g^{(9)}(0) &= -\frac{31}{4}, \\ g^{(11)}(0) &= \frac{691}{8}, & g^{(13)}(0) &= -\frac{5461}{4}, & g^{(15)}(0) &= \frac{929569}{32}. \end{aligned}$$