

# Role Coloring Bipartite Graphs

Sukanya Pandey<sup>1</sup> and Vibha Sahlot<sup>2</sup>

<sup>1</sup> Utrecht University, The Netherlands  
s.pandey1@uu.nl

<sup>2</sup> The Institute of Mathematical Sciences, HBNI, Chennai, India  
sahlotvibha@gmail.com

**Abstract.** A  $k$ -role coloring  $\alpha$  of a graph  $G$  is an assignment of  $k$  colors to the vertices of  $G$  such that if any two vertices are assigned the same color, then their neighborhood are assigned the same set of colors. That is, if  $\alpha(u) = \alpha(v)$  for a pair of vertices  $u$  and  $v$ , then the set of colors assigned to  $N(u)$  and  $N(v)$  are the same (where  $N(u)$  is the set of neighbors of  $u$ ). By definition, every graph on  $n$  vertices admits an  $n$ -role coloring.

While for every graph on  $n$  vertices, it is trivial to decide if it admits a 1-role coloring, determining whether a graph admits a  $k$ -role coloring is a notoriously hard problem for  $k \geq 2$ . In fact, it is known that  $k$ -ROLE COLORING is NP-complete for  $k \geq 2$  on arbitrary graphs.

There has been extensive research on the complexity of  $k$ -ROLE COLORING on various hereditary graph classes. Furthering this direction of research, we show that  $k$ -ROLE COLORING is NP-complete on bipartite graphs for  $k \geq 3$  (while it is trivial for  $k = 2$ ). We complement the hardness result by characterizing 3-role colorable bipartite chain graphs, leading to a polynomial time algorithm for 3-ROLE COLORING for this class of graphs. We further show that 2-ROLE COLORING is NP-complete for graphs that are  $d$  vertices or edges away from the class of bipartite graphs, even when  $d = 1$ .

**Keywords:** Role coloring · Bipartite Graphs · NP-hard · Bipartite Chain Graphs

## 1 Introduction

The notion of role coloring of graphs was introduced by Borgatti and Everett [9] in 1991 motivated by applications in sociology. The underlying principle was that in a social network, individuals play the same role if they relate in the same way to other individuals playing counterpart roles.

Formally, given a graph  $G(V, E)$ , role coloring is an assignment of color to its vertices such that if two vertices get the same color, then the set of colors assigned to their neighborhood is also the same. That is,  $k$ -role coloring is an assignment  $\alpha : V \rightarrow \{1, 2, \dots, k\}$  such that for all  $u, v \in V$ , if  $\alpha(u) = \alpha(v)$  then  $\alpha(N(u)) = \alpha(N(v))$  where  $\alpha(S)$ , for a set of vertices, is the set  $\{\alpha(v) | v \in S\}$ .

Corresponding to a  $k$ -role assignment  $\alpha$ , the role graph  $R(V_R, E_R)$  is defined as the graph with  $V_R = \{1, 2, \dots, k\}$  and  $E_R = \{(\alpha(u), \alpha(v)) | (u, v) \in E(G)\}$ .

Without loss of generality, we assume that each color in a role graph is assigned to some vertex in  $G$ . Then it is easy to see that when  $G$  is connected, the role graph  $R$  is also connected and  $|V_R| \leq |V|$ . Throughout the paper, we assume that the cardinality of the vertex set of  $G$  to be  $n$ . Also, for all  $v \in V(G)$ ,  $deg_G(v) \geq deg_R(\alpha(v))$ .

We define the problem  $k$ -ROLE COLORING as follows,

$k$ -ROLE COLORING

**Input:** An undirected graph  $G$  and an integer  $k$ .

**Question:** Does there exist a function  $\alpha : V(G) \rightarrow \{1, 2, \dots, k\}$  satisfying: if  $\alpha(u) = \alpha(v)$  then  $\alpha(N(u)) = \alpha(N(v))$  for all  $u, v \in V(G)$ ?

We can also define the problem in terms of role graphs. By  $N_G(u)$  and  $N_R(v)$ , we refer to the vertices in the neighborhood of the vertices  $u \in V$  and  $v \in V_R$ , respectively.

$R$ -ROLE COLORING

**Input:** Undirected graphs  $G(V, E)$  and  $R(V_R, E_R)$ .

**Question:** Does there exist a function  $\alpha : V \rightarrow V_R$  satisfying:  $\alpha(N_G(u)) = N_R(\alpha(u))$ , for all  $u \in V$ ?

This problem is equivalent to deciding if there exists a *locally surjective* homomorphism between the graphs  $G$  and  $R$  [4]. Notice that given a role graph  $R$  and a role coloring  $\alpha$  of  $G$ , we can find in polynomial time if  $\alpha$  agrees with  $R$ , that is if  $G$  is  $R$ -role colorable with respect to  $\alpha$ . Also, given a role coloring  $\alpha$  of  $G$ , we can find whether  $\alpha$  is a valid role coloring in polynomial time. Further, we can also find out the role graph  $R$  corresponding to  $\alpha$ .

**Previous Work:** Fiala et al [4] showed that  $R$ -ROLE COLORING is polynomial time solvable when  $R$  has no edges, or has a component isomorphic to a single loop-incident vertex, or it is simple and bipartite and has at least one component isomorphic to a  $K_2$ . For all other cases, they proved that  $R$ -ROLE COLORING is NP-complete. They used this result to show that  $k$ -ROLE COLORING is NP-complete for all  $k \geq 3$ . Prior to their work, it was shown by Roberts and Sheng [11] that  $k$ -ROLE COLORING is NP-complete for  $k = 2$ . Hence, the following dichotomy exists for  $k$ -ROLE COLORING: Given any  $n$  vertex graph  $G(V, E)$ ,  $k$ -ROLE COLORING is polynomial time solvable on  $G$  when  $k = 1$  or  $n$  and NP-complete otherwise.

Various studies have been conducted to find the complexity of computing role-colorings on hereditary graph classes. Sheng [12] gave a greedy algorithm to compute a 2-role coloring of chordal graphs in polynomial time. Later, van't Hof et al. [8] showed that  $k$ -ROLE COLORING is NP-complete on chordal graphs when  $k \geq 3$ . Following these results, Dourado [3] showed that on split graphs, a subclass of chordal graphs, 3-ROLE COLORING can be decided in linear time, whereas for  $k \geq 4$ ,  $k$ -ROLE COLORING is NP-complete on split graphs. The problem remains NP-complete on planar graphs for  $k \geq 2$  [10]. The only known classes of graphs for which  $k$ -ROLE COLORING can be decided in polynomial time

for any fixed  $k$ , are Cographs [10] and Proper Interval graphs [6].

**Our Contribution:** Continuing the line of research on complexity of ROLE COLORING on hereditary graph classes, we explore this problem when restricted to the class of bipartite graphs.  $R$ -ROLE COLORING is known to be NP-complete [4] on bipartite graphs when

- $|E_R| \geq 2$  and
- $R$  does not have a component isomorphic to a single loop-incident vertex, and
- $R$  is not bipartite with at least one component isomorphic to  $K_2$ .

We build on this result and focus on the more general problem of  $k$ -ROLE COLORING on bipartite graphs. We show that  $k$ -ROLE COLORING is NP-complete on the class of bipartite graphs for all  $k \geq 3$  (while it is trivial when  $k \leq 2$ ).

Note that the NP-completeness of  $k$ -ROLE COLORING does not follow from the NP-completeness of  $R$ -ROLE COLORING, where  $R$  is a graph on  $k$  vertices satisfying the characterization in [4]. To exemplify this, let us consider the two variants of the problem with respect to the class of split graphs. Let  $R_1$  be a graph isomorphic to a  $K_2$  with a self loop on exactly one end point, and  $R_2$  be isomorphic to a  $K_2$  with self loops incident on both the end points. It has been shown in [8] that  $R_2$ -ROLE COLORING is NP-complete on split graphs. However, 2-ROLE COLORING is trivially solvable in polynomial time on split graphs with the corresponding role graph being isomorphic to  $R_1$ .

Complementing our hardness result, we give a polynomial time algorithm for 3-ROLE COLORING on bipartite chain graphs. Bipartite chain graphs are bipartite graphs such that for the vertices in each partition, their neighborhoods can be ordered linearly with respect to inclusion. Equivalently, they are bipartite graphs that exclude  $2K_2$  as an induced subgraph. (A  $2K_2$  is a set of two disjoint edges).

Finally, we investigate the problem 2-ROLE COLORING on graphs that are  $d$  vertices (or edges) away from bipartite graphs. We show that the above problem is NP-complete even when  $d = 1$ . Henceforth, we shall refer to such a graph  $G$  as an “almost bipartite” graph. Parameterization by distance from a graph class was first studied for the vertex coloring problem in [1]. We draw inspiration from that and explore it with respect to ROLE COLORING.

**Organization of the paper:** The paper is organised as follows. We define the terminology and notations used in this paper in Section 2.

Section 3 gives the proof for NP-completeness of  $k$ -ROLE COLORING on bipartite graphs, for fixed  $k \geq 3$ . The proof is divided into four parts. We show separately the NP-completeness for each of the cases when  $k \in \{3, 4, 5\}$  and for the case when  $k > 5$ . In Section 4 we characterize bipartite chain graphs that are 3-role colorable, and in section 5, we show that 2-ROLE COLORING is NP-hard on “almost bipartite graphs”.

## 2 Preliminaries

In this section, we state the graph theoretic terminology and notation used in this paper. The set of consecutive integers from 1 to  $n$  is denoted by  $[n]$ . Let  $f : A \rightarrow B$  be a function. Then, for any non-empty set  $A' \subseteq A$ , by  $f(A')$ , we denote the set  $\{f(a) | a \in A'\}$ . The vertex set and the edge set of a graph  $G$  are denoted by  $V(G)$  and  $E(G)$  respectively. By  $|G|$  or  $|V|$  we mean  $|V(G)|$ . An edge between vertices  $u$  and  $v$  is denoted as  $uv$  or  $(u, v)$ . For a vertex  $v \in V(G)$ , its *neighborhood*  $N_G(v)$  is the set of all vertices adjacent to it and its *closed neighborhood*  $N_G[v]$  is the set  $N_G(v) \cup \{v\}$ . This notation is extended to subsets of vertices as  $N_G[S] = \bigcup_{v \in S} N_G[v]$  and  $N_G(S) = N_G[S] \setminus S$  where  $S \subseteq V(G)$ . The *degree* of a vertex  $v \in V(G)$ , denoted by  $d_G(v)$ , is the size of  $N_G(v)$ . A *pendant vertex* is a vertex  $v \in V(G)$  with  $d_G(v) = 1$ . A *pendant path* is an induced path attached to a vertex. A *path* in a simple graph is a sequence of distinct vertices with an edge between every pair of consecutive vertices. The *length of a path* is the number of vertices in a path. We denote a path on  $n$  vertices by  $P_n$ .

In HYPER-GRAPH 2-COLORING, we are given a hyper-graph  $H = (Q_H, S_H)$  where  $Q_H$  is a finite set of vertices and  $S_H$  is a collection of non-empty subsets of  $Q_H$  called hyper-edges. The goal is to determine whether the vertices in  $Q_H$  can be colored with two colors such that in each hyper-edge, there is a vertex of either color. The canonical incidence graph of a hyper-graph  $H = (Q_H, S_H)$  is a bipartite graph  $G(Q, S, E)$  with  $Q$  and  $S$  as the two parts of the bipartition, and for all  $q \in Q$  and  $s \in S$ ,  $(q, s) \in E(G)$  if and only if the vertex corresponding to  $q \in Q_H$  belongs to the hyper-edge  $s \in S_H$ . For further details on graphs, refer [2].

Following are some observations about the role graph that we use throughout the paper without explicitly referring to them.

**Observation 2.1** [4] If  $G$  is  $R$ -role assignable with role assignment  $\alpha$ , then  $\deg_G(u) \geq \deg_R(\alpha(u))$  for all vertices  $u \in V(G)$ .

**Observation 2.2** If  $G$  is a connected graph that is  $R$ -role assignable with the role assignment  $\alpha$ , then  $R$  must be connected.

*Proof.* Since  $G$  is connected, there is a path between every pair of vertices  $u, v \in V(G)$ . Let  $P = ux_1x_2\dots x_{l-2}v$  be a path of length  $l$  in  $G$ . By definition,  $\alpha(N_G(u)) = N_R(\alpha(u))$ . That is,  $\alpha(x_1) \in N_R(\alpha(u))$ . By induction on  $l$ , it can be shown that there exists a path between  $\alpha(u)$  and  $\alpha(v)$  in  $R$ . Since the choice of  $u$  and  $v$  was arbitrary, the claim is true for all pairs of vertices in  $R$ . Hence,  $R$  is connected.  $\square$

## 3 NP-completeness of $k$ -ROLE COLORING on bipartite graphs


In this section, we prove that  $k$ -ROLE COLORING is NP-complete on the class of bipartite graphs, when  $k \geq 3$ .

**Theorem 3.1**  $k$ -ROLE COLORING is NP-complete on bipartite graphs, for any fixed  $k \geq 3$ .

*Proof.* It is easy to see that  $k$ -ROLE COLORING is in NP. We prove the NP-hardness in four parts. We separately deal with the cases when  $k = 3$ ,  $k = 4$ ,  $k = 5$  and  $k > 5$ . In all the four cases, we give a reduction from HYPER-GRAPH 2-COLORING (which is known to be NP-complete [5]). For each case, the hyper-graph is denoted by  $H = (Q_H, S_H)$ , where  $Q_H$  is a finite set of vertices and  $S_H$  is a collection of non-empty subsets of  $Q_H$  called hyper-edges. The corresponding canonical incidence graph of a hyper-graph  $H = (Q_H, S_H)$  is a bipartite graph denoted as  $G(Q, S, E)$  with  $Q$  and  $S$  as the two parts of the bipartition, and for all  $q \in Q$  and  $s \in S$ ,  $(q, s) \in E(G)$  if and only if the vertex corresponding to  $q \in Q_H$  belongs to the hyper-edge  $s \in S_H$ .

**Lemma 3.1** 3-ROLE COLORING is NP-complete on connected bipartite graphs even with the restriction that the pendant vertices lie in at most one partition.

*Proof. Construction:* We assume that  $H$  has no hyper-edges with less than two vertices, else  $H$  would trivially be a no instance. Thus no vertex in  $S$  can be a pendant vertex in the canonical incidence graph  $G$  of  $H$ . To each vertex  $q \in Q$ , we add a path on two vertices, such that the vertex adjacent to  $q$  is labeled  $b_q$  and the other vertex adjacent to  $b_q$  is  $a_q$ . For each  $q \in Q$ , the sub-graph induced on  $\{q, b_q, a_q\}$  is a “pendant path” and has length 3. The new graph thus obtained is denoted by  $G'$  (refer to Figure 1).




**Fig. 1.** Construction in Lemma 3.1

We claim that  $H$  is a *yes* instance of HYPER-GRAPH-2-COLORING if and only if  $G'$  is a *yes* instance of 3-ROLE COLORING. First, assume that  $H$  is a *yes* instance and  $\beta : Q_H \rightarrow \{1, 2\}$  is a 2-coloring of the vertices in  $Q_H$ . Then, the role assignment  $\alpha : V(G') \rightarrow \{1, 2, 3\}$  can be defined as: for all  $q \in Q$ ,  $\alpha(q) = \beta(q)$ ;  $\alpha(b_q) = 2$  if  $\alpha(q) = 1$  and  $\alpha(b_q) = 1$ , otherwise.  $\alpha(s) = 3$  for all  $s \in S$ ,  $\alpha(a_q) = 2$ .  $G'$  is, therefore, a *yes* instance of 3-ROLE COLORING.

Conversely, suppose that  $G'$  is a *yes* instance of 3-ROLE COLORING with the 3-role coloring assignment  $\alpha$  and the corresponding role graph  $R$ . Without loss of generality, assume that for some  $q \in Q$ ,  $\alpha(a_q) = 1$ . Then,  $\alpha(b_q) \neq 1$ , or else the role graph would be disconnected (with one component isomorphic to a single vertex with a loop). This is a contradiction as  $G'$  is connected. So let  $\alpha(b_q) = 2$ . Now  $\alpha(q) \neq 1$  and  $\alpha(q) \neq 2$ , otherwise role graph would be disconnected. So, let  $\alpha(q) = 3$ . We prove that  $\alpha(S) = \{2\}$ . The neighbor of  $q$ , say  $s \in S$  could either be colored 2 or 3. Suppose that  $\alpha(s) = 3$ . This forces one of the neighbors of  $s$  to be colored 2. Since, no hyper-edge in  $S_H$  is singleton, let there be a neighbor  $q'$  of  $s$  such that  $\alpha(q') = 2$ . Consider the vertex  $b_{q'}$  adjacent to  $q'$ .  $\alpha(b_{q'}) \neq 1$  as that would force  $\alpha(a_{q'}) = 2$ . This is not possible because  $a_{q'}$  has no neighbor colored 3. Similarly,  $\alpha(b_{q'}) \neq 3$ . Therefore, in a valid role assignment of  $G'$ ,  $\alpha(s) \neq 3$ , for any  $s \in S$ . Also  $\alpha(s) \neq 1$  for  $s \in S$  as 3 is not adjacent to 1 in the role graph. Hence,  $\alpha(s) = 2$  for all  $s \in S$ . We define the coloring of the hypergraph  $\beta : Q_H \rightarrow \{2, 3\}$ , as follows as  $\beta(q) = \alpha(q)$ . This is a valid 2-coloring of  $H$ , as every  $s \in S$  is adjacent to at least one vertex colored 2 and at least one vertex colored 3. Hence,  $H$  is a *yes* instance of HYPERGRAPH-2-COLORING.  $\square$

**Lemma 3.2** 4-ROLE COLORING is NP-complete on bipartite graphs.

*Proof. Construction:* Let  $H$  be an instance of HYPER-GRAPH 2-COLORING. The incidence graph  $G(Q, S, E)$  is a connected bipartite graph, with the bipartition  $(Q, S)$ . To construct the graph  $G'$  from  $G$ , we add a pendant vertex  $p^s$  to every vertex  $s \in S$  (refer to figure 2). Let  $P^s = \{p^s | s \in S\}$ .



**Fig. 2.** Construction in Lemma 3.2

We now claim that  $H = (Q_H, S_H)$  is 2-colorable if and only if  $G'$  is 4-role colorable. Suppose that  $H$  has a 2-coloring,  $\beta : Q_H \rightarrow \{1, 2\}$ . Then the role

assignment for  $G'$  is defined as: For each  $q \in Q$ ,  $\alpha(q)=\beta(q)$ ; for all  $s \in S$ ,  $\alpha(s) = 3$  and  $\alpha(p^s) = 4$ . Corresponding to  $\alpha$ , the role graph  $R$  is defined as  $V(R) = \{1, 2, 3, 4\}$  and  $E(R) = \{(1, 3), (2, 3), (3, 4)\}$ . As  $H$  is a *yes* instance of HYPER-GRAPH 2-COLORING, every vertex  $s \in S$  colored 3 is adjacent to vertices colored 1 and 2 in  $Q$  and the vertex  $p^s$  colored 4. Thus,  $\alpha$  is consistent with  $R$ .

Conversely, suppose that  $G'$  has a 4-role coloring  $\alpha : V(G) \rightarrow \{1, 2, 3, 4\}$ , and the corresponding role graph is  $R$ .

**Case 1.**  $Q$  is monochromatic. Let  $\alpha(q) = 1$ , for each  $q \in Q$ .

**Claim** *No vertex in  $S$  is assigned a color same as that of the vertices in  $Q$ .*

*Proof.* Towards a contradiction, suppose that there exists  $s \in S$  colored 1. Then, 1 has a self-loop incident on it in the role graph  $R$ . If the pendant attached to  $s$  gets colored 1, then by connectedness of  $G'$ , all the vertices of  $G'$  get assigned the color 1. This is a contradiction to  $\alpha$  being a 4-role coloring of  $G'$ . Therefore, suppose that  $\alpha(p^s) = 2$ . So,  $N_R(2) = \{1\}$ . As  $q$  is a vertex colored 1, it must have another neighbor in  $S$  colored 2. Hence, there must exist a vertex  $s' \in S$  such that  $\alpha(s') = 2$ . This in turn would force  $\alpha(p^{s'})$  to be equal to 1, which is not possible, as  $deg_R(1) = 2 > deg_G(p^{s'})$ . Hence,  $S$  cannot have a vertex with the same color as that assigned to the vertices of  $Q$ .  $\square$

**Claim.** *None of the pendant vertices in  $P^s$  gets the color 1.*

*Proof.* Suppose that there exists a vertex  $p^s$  of  $P^s$  colored 1. By the previous claim, no vertex in  $S$  is colored 1. Suppose that  $\alpha(s) = 2$ . Since the vertices colored 1 are only adjacent to vertices colored 2,  $S$  must be monochromatic with the color 2, as  $Q$  is monochromatic with the color 1. Therefore,  $P^s$  must be monochromatic with the color 1 assigned to its vertices, a contradiction to  $\alpha$  being a 4-role coloring of  $G'$ . Therefore, it follows that  $1 \notin \alpha(P^s \cup S)$ .  $\square$

**Claim.**  *$P^s$  must be monochromatic.*

*Proof.* Towards a contradiction, suppose that there exist vertices  $p^s$  and  $p^{s'}$  colored 2 and 3, respectively. Then  $\alpha(s) \neq 2$  and  $\alpha(s') \neq 3$ , else every vertex colored 2 or 3 could only have neighbors of the same color. As every  $s \in S$  has a neighbor in  $Q$  colored 1, we know that is not possible. This implies that some vertices in  $S$  are adjacent to vertices colored 1 and 2, whereas the others have neighbors colored 1 and 3. Therefore, there is no color that can be assigned to the vertices of  $S$ , which is a contradiction to  $G'$  being 4-role colorable. Hence,  $P^s$  must be monochromatic.  $\square$

Let  $\alpha(P^s)=\{2\}$ . That forces  $S$  to be monochromatic with a third color, say 3, which again contradicts the assumption that  $G'$  is 4-role colorable. Therefore, when  $Q$  is monochromatic,  $G'$  is not 4-role colorable, which contradicts our assumption.

**Case 2:**  $Q$  is bi-chromatic. Let the colors assigned to vertices in  $Q$  by  $\alpha$  be 1 and 2.

**Claim.** *The vertices in  $S$  are not assigned the colors 1 and 2.*

*Proof.* We assume the contrary. Suppose that there exists  $s \in S$  such that  $\alpha(s) = 1$ . If  $\alpha(p^s) = 1$ , then 1 would only be adjacent to itself and the role graph would be disconnected. This is not possible as  $G'$  is connected. Suppose that  $\alpha(p^s) =$

2. Then  $N_R(2) = \{1\}$ . For the role graph to be connected, 1 must be adjacent to a third color, say 3. However, the neighbors of  $s$  in  $Q$  are colored either 1 or 2, which is a contradiction. Similarly, no  $s \in S$  gets colored 2.  $\square$

**Claim.** *The pendants in  $P^s$  are not assigned any color in  $\alpha(Q)$ .*

*Proof.* Suppose that there exists a pendant vertex  $p^s$  colored 1. Let its neighbor  $s$  in  $S$  have role 3 (from the previous claim we know that it can get neither role 1 nor role 2). Also, 3 must be adjacent to a different color, say 2, otherwise the role graph would be disconnected with 1 and 3 forming an induced  $K_2$ . This means that no pendant receives the color 3 as the degree of 3 is at least equal to 2. If there exists a pendant colored 2, then its neighbor must be colored 3 as well. So,  $N_R(1) = N_R(2) = \{3\}$  and  $N_R(3) = \{1, 2\}$ . All the vertices in  $Q$  are adjacent only to vertices in  $S$ . Hence,  $S$  is monochromatic with the role 3. This is a contradiction to  $\alpha$  being a 4-role assignment. This implies that  $1 \notin \alpha(P^s)$  and  $2 \notin \alpha(P^s)$ . This proves our claim that  $\alpha(Q) \cap \alpha(P^s) = \phi$ .  $\square$

Thus, the only possible role assignment in this case is :  $\alpha(Q) \in \{1, 2\}$ ,  $\alpha(S) = \{3\}$ , and  $\alpha(P^s) = \{4\}$ . So, every vertex in  $S$  is forced to be adjacent to at least one vertex of role 1 and at least one vertex of role 2.

Having dealt with the cases when  $Q$  is monochromatic and bi-chromatic respectively, we now show that  $|\alpha(Q)| = 2$ .

**Claim.** *If  $G'$  has a 4-role coloring,  $|\alpha(Q)| = 2$ .*

*Proof.* For  $|\alpha(Q)| = 1$ , we have already seen that no 4-role coloring exists for  $G'$ . Suppose that  $|\alpha(Q)| \geq 3$ . By the previous claims  $\alpha(Q) \cap \alpha(\{P^s \cup S\}) = \phi$ . Therefore, the set  $P^s \cup S$  is monochromatic. That is not possible because every vertex in  $S$  is adjacent to vertices of at least 2 different role colors, whereas the vertices in  $P^s$  can only be adjacent to one vertex. Therefore,  $|\alpha(Q)| = 2$ .  $\square$

**Claim.**  $H = (Q_H, S_H)$  is a yes instance of HYPERGRAPH 2-COLORING

*Proof.* Define the coloring of the vertices of  $Q_H$  as  $\beta(q) = \alpha(q)$  for all  $q \in Q$ . Since, each  $s \in S$  is adjacent to some vertex colored 1 and some vertex colored 2, the corresponding hyper-edges are all bi-chromatic.  $\square$


**Lemma 3.3** 5-ROLE COLORING is NP-complete on connected bipartite graphs.

*Proof. Construction:* The construction is mostly the same as in the proof of Lemma 3.2. To the bipartite graph  $G'(Q \cup P^s, S, E)$ , we add a vertex  $u$  and make it adjacent to all the vertices in  $Q$ . The resulting graph  $G''$  is bipartite, as  $u$  has no neighbors in  $S$  (refer to Figure 4).

We now show that  $H$  has a 2-coloring if and only if  $G''$  has a 5-role coloring. Suppose that  $H$  has a 2-coloring assignment,  $\beta : Q_H \rightarrow \{1, 2\}$ . Then, the role assignment for  $G''$  is defined as follows:  $\alpha(q) = \beta(q)$ , for all  $q \in Q$ ;  $\alpha(u) = 3$ ;  $\alpha(s) = 4$ , for all  $s \in S$ ; and  $\alpha(p^s) = 5$ , for all  $p^s \in P^s$ . The role graph  $R(V_R, E_R)$ , corresponding to the 5-role coloring  $\alpha$ , is defined as:  $V_R = \{1, 2, 3, 4, 5\}$  and  $E_R = \{(1, 3), (1, 4), (2, 3), (2, 4), (4, 5)\}$ .

Conversely, suppose that  $G''$  has a 5-role coloring,  $\alpha : V(G'') \rightarrow \{1, 2, 3, 4, 5\}$ . We first argue that the set  $Q \subseteq V(G)$  can not be monochromatic.

**Case 1.** Let  $Q$  be monochromatic. Without loss of generality, let  $\alpha(Q) = 1$ .



**Fig. 3.** Construction in Lemma 3.3

We claim that the vertex  $u$  must not be assigned the color 1. If  $\alpha(u) = 1$ , then every vertex colored 1 can only be adjacent to vertices colored 1. That would mean that the role-graph  $R$  is disconnected. This is not possible as  $G''$  is connected.

Without loss of generality, let  $\alpha(u)=2$ . So,  $N_R(2) = \{1\}$ . Now we prove that neither the vertices in  $S$  nor the pendants in  $P^s$  are assigned the colors 1 and 2.

Suppose that there exists  $s \in S$ , such that  $\alpha(s) = 1$ . Then  $s$  is adjacent to vertices colored 1 in  $Q$  as well as a pendant vertex  $p^s$ . Then,  $\alpha(p^s) = 2$ , as 1 is adjacent to 1 and 2, but all the other neighbors of  $s$  are colored 1. In fact, all the vertices in the set  $S$  must get colored 1 because  $N(Q) = S \cup \{u\}$  and  $u$  is colored 2. This forces every  $p^s$  to be colored 2. This is a contradiction to the hypothesis that  $\alpha$  is a 5-role coloring of  $G''$ .

Since each  $s \in S$  is adjacent to a pendant vertex,  $\alpha(s) \neq 2$ , because 1 is adjacent to both 1 and 2 but  $\deg_{G''}(p^s) = 1$ . So, the pendant vertices cannot be colored 1. Therefore, each element of  $S \cup P^s$  must be assigned one of the three colors, 3, 4 and 5.

We claim that  $P^s$  is monochromatic. Suppose that  $P^s$  has vertices with role 3 and 4. Then no vertex of  $S$  gets the colors 3 and 4 as these vertices are adjacent to vertices in  $Q$  colored 1, whereas the pendants are not. This leaves us with the role 5 for  $S$  but, some vertices of  $S$  are adjacent to pendants colored 3 whereas some others to pendants colored 4. Thus, there is no color that can be assigned to  $S$  which contradicts the assumption that  $G''$  has a 5-role coloring.

Hence, we assume that  $P^s$  is monochromatic with the role 3. However, this forces  $S$  to be monochromatic with the role 4. This again contradicts  $\alpha$  being a 5-role coloring. Therefore, when  $Q$  is monochromatic,  $G''$  does not have any

5-role coloring.

Case 2:  $Q$  is bi-chromatic. Without loss of generality, assume that  $\alpha(Q) = \{1, 2\}$ .

We claim that  $\alpha(u) \notin \{1, 2\}$ . Suppose that  $\alpha(u) = 1$ . Then,  $N_R(1) = \{1, 2\}$ . This implies that the vertices in  $Q$  colored 1 must be adjacent to some vertex colored 2. That is only possible if some vertices in  $S$  receive the color 2. Let  $Q_1$  be the set of vertices in  $Q$  colored 1 and  $S_2$  be the set of vertices in  $S$  colored 2. Then, the vertices in  $P^s \cap N(S_2)$  cannot be colored 1 ( $deg_R(1) = 2$ ). They cannot be colored 2 either, as 2 must be adjacent to some vertex colored 1. Suppose that these vertices get colored 3. Then, the vertices in  $Q \setminus Q_1$  must be adjacent to some vertex with role 3, as well. This would imply that some vertices in  $S$  get the role 3. However, this is not possible, as 3 is only adjacent to 2 but the pendants attached to these vertices cannot be colored 2. Therefore,  $\alpha(u) \neq 1$ . Similarly,  $\alpha(u) \neq 2$ .

Therefore, let  $\alpha(u) = 3$ .

Now we prove that  $\alpha(Q \cup \{u\}) \cap \alpha(S \cup P^s) = \phi$ .

Suppose that there exists  $s \in S$  colored 1. Then  $s$  must be adjacent to a vertex colored 3 as  $3 \in N_R(1)$ . However, the neighbors of  $s$  include vertices in  $Q$ , which are colored 1 or 2, as well as the pendant vertex  $p^s$ , which cannot be colored 3 because  $deg_R(3) = 2 > deg_G''(p^s)$ . Similarly, no vertex of  $S$  can be given the role 2 either. Suppose that some  $s' \in S$  gets the role 3, then as 3 is only adjacent to 1 and 2, the pendant  $p^{s'}$  must get assigned the colors 1 or 2. This would force all the vertices in  $S$  to be colored 3 and all the vertices of  $P^s$  to get the role  $\alpha(p^{s'})$ . This is a contradiction to  $\alpha$  being a 5-role coloring.

Now, the vertices in  $S \cup P^s$  can be colored with 4 and 5. Suppose that  $p^s$  gets the role 4. Then,  $\alpha(s) \neq 4$  because  $s$  is adjacent to vertices of role 1 and 2 whereas  $p^s$  is not. So,  $\alpha(s) = 5$ . This implies that no pendant vertex gets the role 5, so  $P^s$  is monochromatic with the role 4 and  $S$  is monochromatic with role 5. Also, each vertex of  $S$  must be adjacent to at least one vertex with role 1 and one vertex of role 2.

In this case, we assign colors to the vertices of the hyper-graph as  $\beta(q) = \alpha(q)$  for all  $q \in Q_H$ .

Now we prove that if  $G''$  is 5-role colorable, then  $|\alpha(Q)| = 2$ . We have seen that when  $Q$  is monochromatic, there is no possible 5-role assignment for  $G''$ . Suppose that  $|\alpha(S)| \geq 3$ . By the previous claim,  $\alpha(Q \cup \{u\}) \cap \alpha(S \cup P^s) = \phi$ . That would leave us with at most one role color for the vertices in  $S \cup P^s$ . However, we have also shown that  $S \cup P^s$  cannot be monochromatic. So,  $G''$  does not have a 5-role coloring, which is a contradiction. Hence,  $|\alpha(Q)| = 2$ .  $\square$

**Lemma 3.4** *k-ROLE COLORING is NP-complete on connected bipartite graphs, for any fixed  $k > 5$ .*

*Proof. Construction:* We use the same reduction as in proof of the Lemma 3.3. Consider the graph  $G''$ . In  $G''$ , we tweak the set  $P^s$  to include “pendant paths”

instead of vertices, defined as  $P^s = \{[p_1^s, \dots, p_{k-4}^s] : \text{for all } s \in S\}$ . Each path in  $P^s$  is attached to the corresponding  $s$  through the edge  $(p_{k-4}^s, s)$ .




Fig. 4. Construction in Lemma 3.4

We prove that  $H$  has a 2-coloring assignment if and only if  $G''$  has a  $k$ -role coloring. Suppose that the hyper-graph  $H$  has a 2-coloring,  $\beta : Q_H \rightarrow \{1, 2\}$ . Let the role assignment for  $G''$  be the following:  $\alpha(q) = \beta(q)$ , for each  $q \in Q$ ;  $\alpha(u) = 3$ ;  $\alpha(s) = 4$ , for all  $s \in S$ ;  $\alpha(p_{k-4}^s) = 5$ , for all  $s \in S$ ;  $\alpha(p_j^s) = k - j + 1$ , where  $1 \leq j \leq k - 5$ . This gives us a  $k$ -role coloring of  $G''$ ,  $\alpha$  with the corresponding role-graph  $R(V_R, E_R)$  defined as:  $V_R = \{1, 2, \dots, k\}$  and  $E_R = \{(1, 3), (2, 3), (1, 4), (2, 4), (4, 5), (j, j + 1)\}$ , for all  $5 \leq j \leq k - 1$ .

To prove the converse, we have already shown that  $Q$  cannot be monochromatic and that  $|\alpha(Q)| = 2$ . We have also shown that  $\alpha(Q \cup \{u\}) \cap \alpha(S \cup P^s) = \phi$ . Hence, the only thing left to show is that  $\alpha(Q \cup S \cup \{u\}) \cap \alpha(P^s) = \phi$ .

We prove that  $\alpha(Q \cup S \cup \{u\}) \cap \alpha(P^s) = \phi$ . To see this, consider the path attached to  $s$ . For convenience, we remove the superscript and denote the path as  $[p_1, \dots, p_{k-4}]$ . Towards a contradiction, suppose that  $\alpha(p_{k-4}) = \alpha(s) = 4$ . Then,  $N_R(4) = \{1, 2, 4\}$ . However, the degree of  $p_{k-4}$  is 2, hence it cannot be assigned a color with degree greater than or equal to 3.

Now  $\alpha(p_{k-4}) \neq 3$  as 3 is not adjacent to 4. Its color cannot be equal to 1 or 2 because that would force  $p_{k-3}$  to get role 3, contradicting  $G''$  having a  $k$ -role assignment,  $k > 5$ . So, let  $\alpha(p_{k-4}) = 5$ . Since, 1, 2 and 3 are not adjacent to 5,  $\alpha(p_{k-5}) \notin \{1, 2, 3\}$ . Since 4 is adjacent to 1, 2 and 5 in  $R$ , no vertex in the path can be assigned the role 4, as these vertices have degree equals two which is

less than the degree of 4 in the role graph. If  $\alpha(p_{k-5}) = 5$  then  $\alpha(p_{k-6})$  is forced to be 4, which is not possible. Hence,  $p_{k-5}$  must receive a fresh color, say 6. By induction on  $j$ , we can show that each  $p_{k-j}$  receives a fresh color  $j$ ,  $j \geq 4$ . This completes the proof.  $\square$

This completes the proof of Theorem 3.1  $\square$

## 4 3-role coloring bipartite chain graphs

**Definition.** ([7], [13]) A bipartite graph is a *chain graph* if and only if the neighborhood of the vertices of each partition can be linearly ordered with respect to inclusion. Equivalently, a bipartite graph is a chain graph if and only if it is  $2K_2$ -free.

A *universal vertex*<sup>3</sup> is a vertex in one partition of a bipartite graph that is adjacent to every vertex in the other partition.

Throughout this section, we assume  $G = (X, Y, E)$  is a bipartite chain graph.  $U_X$  and  $U_Y$  are the sets of universal vertices in the partitions  $X$  and  $Y$ , respectively. Likewise,  $P_X$  and  $P_Y$  are sets of pendant vertices in the partitions  $X$  and  $Y$ . Note that  $P_X$  and/or  $P_Y$  may be empty. We also assume  $|V| \geq 3$  or else,  $G$  is trivially a *no* instance. In the rest of this section, we aim to prove the following theorem.

**Theorem 4.1** *A bipartite chain graph  $G = (X, Y, E)$ , where  $|X \cup Y| \geq 3$ , is 3-role colorable if and only if any of the following is true:*

- $G$  is not connected
- $|X| = 1$
- $|X| = 2$  and both the vertices of  $X$  are universal vertices.
- $|X| = 2$ ,  $|Y| > 2$  and  $|Y \setminus P_Y| > 1$ .
- $|X| \geq 3$  and  $|Y| \geq 3$ .

Furthermore, given any bipartite chain graph we can check in polynomial time whether it is 3-Role colorable.

*Proof.* We proceed by considering the case when  $|X| \geq 3$ , and  $|Y| \geq 3$ . We separately handle the edge cases when  $|X| \leq 2$  or  $|Y| \leq 2$ .

**Case 1.** First, let us assume that  $G$  is connected and  $|X| \geq 3$ , and  $|Y| \geq 3$ .

As there are no isolated vertices in  $G$ , the linear ordering on the neighborhood of vertices in each partition ensures that the largest neighborhood contains all the vertices in the partition. Hence, each partition has at least one universal vertex. We demonstrate a 3-role coloring for each of the following sub-cases:

**Sub-case 1.**  $P_X = \phi = P_Y$

Let  $\alpha : V(G) \rightarrow \{1, 2, 3\}$  be a coloring assignment of  $V(G)$  where  $\alpha(u) = \{1\}$  for some  $u \in U_X$ ;  $\alpha(Y) = \{2\}$ ; and  $\alpha(w) = 3$  for all  $w \in G \setminus N[U_X]$ .

<sup>3</sup> Our definition of “universal vertex” is different from the standard definition.

**Claim.**  $\alpha$  is a valid role-coloring of  $G$ .

*Proof.* The vertices in  $Y$  form the neighborhood of  $U_X$ , and  $\alpha(Y) = \{2\}$ . Each vertex in  $Y$  has a neighbor of color 1. Since there are no pendants in  $G$ , each of them has at least one other neighbor in  $X$ . This ensures that each vertex colored 2 has a neighbor colored 3.  $\square$

**Sub-case 2.**  $P_X \neq \phi$ , and  $P_Y = \phi$

**Claim.** The unique neighbor of a pendant in  $P_X$  must be a universal vertex. Moreover,  $|U_Y| = 1$ .

*Proof.* Let  $v \in P_X$  and let  $u \in Y$  be its unique neighbor. If  $u$  is not a universal vertex, then there exists a vertex  $x$  in  $X$ , that is not adjacent to  $u$ . However, since no isolated vertices exist in  $G$ ,  $x$  must be adjacent to some other vertex in  $G$ , say  $y$ . Therefore,  $(x,y)$  and  $(u,v)$  induce a  $2K_2$  which contradicts the assumption that  $G$  is a bipartite chain graph. Suppose that  $|U_Y| \geq 2$ . Then every vertex in  $X$  would have degree at least 2. This contradicts our assumption that  $P_X \neq \phi$ .  $\square$

Now, we define the role coloring of vertices in  $G$  as follows:  $\alpha : V(G) \rightarrow \{1, 2, 3\}$ ,  $\alpha(v) = 1$ , for all  $v$  in  $Y$ ;  $\alpha(w) = 2$ , where  $w \in U_X$  and  $\alpha(X \setminus \{w\}) = 3$ . Every vertex colored 1 is adjacent to the vertex colored 2 (that is  $w$ ) and at least one vertex colored 3. Thus,  $\alpha$  is a valid role-coloring of  $G$ .

**Sub-case 3.** Pendants occur in both the partitions.

**Claim.**  $|U_X| = |U_Y| = 1$

*Proof.* Follows from the previous claim in Case 2.  $\square$

Let  $u$  and  $v$  be the universal vertices in  $X$  and  $Y$ , respectively. We first assume that  $N(u) \cup N(v)$  does not induce an independent set. Let  $\alpha : V \rightarrow \{1, 2, 3\}$  be a role coloring of  $G$ , as before.

In this case,  $\alpha(P_X \cup P_Y) = \{1\}$ ;  $\alpha(u) = \alpha(v) = 2$ ;  $\alpha(G \setminus (\{u\} \cup \{v\} \cup P_X \cup P_Y)) = \{3\}$ . Therefore, every vertex colored 1 is adjacent only to vertices colored 2. The two vertices colored 2 are adjacent to all the vertices in each partition and hence adjacent to both 1 and 3. Since the neighborhood of  $u$  and  $v$  do not form an independent set, at least one vertex in each partition is not a pendant. Hence, there are vertices colored 3 which are adjacent to each other and the universal vertices and  $\alpha$  is a valid role coloring of  $G$ .

Now, suppose that  $N(u) \cup N(v)$  is an independent set. Then,  $|N(u)| \geq 2$  and  $|N(v)| \geq 2$ , as each partition has at least 3 vertices. Define the role coloring as:  $\alpha(u) = \alpha(v) = 2$ . Color any one neighbor each of  $u$  and  $v$  with the color 1 and another neighbor with the color 3. The remaining vertices can be arbitrarily given the colors 1 or 3.  $\alpha$  is a valid role coloring of  $G$ , in this case too.  $\square$

**Case 2.** Now we deal with the case where either  $|X| \leq 2$  or  $|Y| \leq 2$ . We first consider the case where  $|X| = 2$  and  $X = \{u, v\}$ .

**Claim.**  $G$  is 3-role colorable if both the vertices in  $X$  are universal.

*Proof.* Here both  $u$  and  $v$  are universal vertices. Assign color 1 to  $u$ , color 2 to  $v$  and color 3 to vertices of  $Y$ . The role graph is defined as  $R = (\{1, 2, 3\}, \{(1, 3), (2, 3)\})$ .  $\square$

Now, suppose that  $u$  is not universal. If  $|Y| = 2$ , then  $G$  is a *no* instance as  $P_4$  is not 3-role colorable. Henceforth, we assume that  $|Y| > 2$ . Let  $Y = P_Y \cup T$  where  $T$  is the set of vertices of degree two.

**Claim.** If  $|U_X| = 1$  then  $G$  is 3-role colorable if and only if  $|T| \geq 2$ .

*Proof.* Assume that  $u \in X$  is universal and  $v$  is not. Then,  $P_Y$  is not empty. Let  $\alpha(a) = 1$ , for some  $a \in P_Y$ . We cannot assign 1 to  $u$ , due to the connectivity constraint of the role graph. Let  $\alpha(u) = 2$ . So  $(1, 2) \in E(R)$  where  $R$  is the role graph corresponding to  $\alpha$ .

We prove the forward direction, by proving the contrapositive. Suppose that  $|T| < 2$ .  $T \neq \phi$  because  $G$  is connected. Hence,  $|T| = 1$ . Let  $t \in T$ . If possible, let  $\alpha(v) = 1$ . It forces  $\alpha(t) = 2$  which forces some vertex in  $P_Y \setminus \{a\}$  to have color 2 as for  $\alpha$  to be consistent,  $u$  needs a neighbor colored 2. However, the color 2 cannot be assigned to a pendant, as  $\deg_G(u) \geq \deg_R(\alpha(u))$ ,  $\forall u \in V(G)$ .

So let  $\alpha(v) = 2$ . If  $\alpha(t) = 1$ , it forces all vertices in  $P_Y$  to get color 1. This contradicts the assumption that  $\alpha$  is a 3-role coloring assignment. If  $\alpha(t) = 2$ , it forces the role graph to have a single vertex labeled 2, with a self loop which contradicts the fact that  $(1, 2) \in E(R)$ . If  $\alpha(t) = 3$ , it forces role graph to have just an edge  $(2, 3)$  which is again a contradiction as  $(1, 2) \in E(R)$ .

Therefore, let  $\alpha(v) = 3$ . Then,  $\alpha(t) \neq 1$ , as  $t$  is adjacent to vertices colored 2 and 3, but the degree of 1 is equal to one in  $R$ . Also,  $\alpha(t) \neq 2$ , as  $t$  has no neighbor colored 1. If possible, let  $\alpha(t) = 3$ . This is not a valid role coloring as  $\alpha(v) = \alpha(t) = 3$  but  $\alpha(N(u)) = \{3\} \neq \{2, 3\} = \alpha(N(b))$ . Hence if  $|T = 1|$ , then  $G$  is a *no* instance.

Conversely, suppose that  $|T| \geq 2$ . Let  $t'$  also belong to  $T$ . We define the role coloring as:  $\alpha(v) = \alpha(u) = 2$ ,  $\alpha(t) = 1$ ,  $\alpha(t') = 3$ . Also, color all vertices in  $P_Y$  with color 1 and all the remaining vertices in  $T$  with the color 3. It is easy to verify that this coloring is consistent with the role graph  $R = (\{1, 2, 3\}, \{(1, 2), (2, 3)\})$ .  $\square$

Now consider the case when  $|X| = 1$ .

**Claim.**  $G$  is 3- role colorable if  $|X| = 1$ .

*Proof.* As  $G$  is connected, thus  $v \in X$  is universal. Let  $\alpha(v) = 1$  and for a vertex  $w \in Y$   $\alpha(w) = 2$ . Assign color 3 to the rest of vertices in  $Y$ . It is easy to check that this coloring satisfies the role graph  $R = (\{1, 2, 3\}, \{(1, 2), (1, 3)\})$ .  $\square$

We can argue symmetrically for  $Y$ .

Now we claim that a disconnected bipartite chain graph is always 3-role colorable if it has more than two vertices. Note that edges are present in at most one of the connected components of a disconnected bipartite chain graph, otherwise there will be a  $2K_2$ . So there are two cases, first when there is no edge in the bipartite chain graph. As we have assumed that there are at least three vertices, thus the graph is trivially 3-role colorable with an edgeless role graph. In the second case, one connected component has edges. In that case, color the connected component with edges using colors 1 and 2. This can be done as the graph is bipartite. Color rest of the isolated vertices with color 3. Note that this is a valid 3-role coloring with role graph  $(\{1, 2, 3\}, \{(1, 2)\})$ .

There are a finite number of cases for which  $G$  is 3-role colorable, and each case can be verified in  $\mathcal{O}(n + m)$  time. Hence, we can decide 3-ROLE COLORING on bipartite chain graphs in polynomial time.  $\square$

## 5 Almost bipartite graphs

An “almost bipartite” graph is a graph that has a set of at most  $d$  vertices or  $d$  edges,  $d$  being a constant, which on deletion yield a bipartite graph. Here, we prove that 2-ROLE COLORING is NP-complete on the class of almost bipartite graphs, even if  $d = 1$ .

**Theorem 5.1** 2-ROLE COLORING is NP-complete on the class of almost bipartite graphs.

*Proof.* We know from [4] that given a connected bipartite graph  $G$  and a role graph  $R$  isomorphic to an edge with a self loop incident on one of its end points, deciding if  $G$  is 2-role colorable, is NP-complete. Thus, we show a reduction from this problem to 2-ROLE COLORING on almost bipartite graphs with  $d = 1$ . Refer to Figure 5.




Fig. 5. Construction in Theorem 5.1

Let  $R = (V_R, E_R)$  be defined as  $V_R = \{1, 2\}$  and  $E_R = \{(1, 1), (1, 2)\}$ . Let  $G$  be a connected bipartite graph and let  $x \in V(G)$ . We construct  $G'$  from  $G$  by adding to it 3 vertices  $a, b$  and  $c$  and the edges  $(a, b), (a, c), (x, a), (x, b)$ .  $G'$  is not bipartite as it contains a triangle, namely,  $abx$ . It is 1 vertex (remove  $b$ ) and 1 edge (remove  $(x, b)$ ) away from the bipartite graph  $G$ .

Suppose that  $G$  is a *yes* instance of  $R$ -ROLE COLORING. Then, it has a role assignment  $\alpha$ , with the role graph being isomorphic to  $R$ . Then we can extend  $\alpha$  to the vertices of  $G'$  as follows: If  $\alpha(x) = 1$ , then color  $a$  with 1 and  $b$  and  $c$  with 2. If  $\alpha(x) = 2$  then color both  $a$  and  $b$  with 1 and  $c$  with 2. Thus,  $G'$  has an  $R$ -role coloring and therefore a valid 2-role coloring.

Conversely, suppose that  $G'$  has a 2-role coloring  $\alpha'$ . First assume that  $\alpha'(a) = 1$ . If both  $x$  and  $b$  received the color 1, then  $b$  would only be adjacent to vertices colored 1. Thus, the role graph would be disconnected, as 1 could only be adjacent to itself. This is not possible as  $G'$  is a connected graph.

Hence, at least one of them gets the color 2. Also,  $c$  gets color 2 as its degree is one, but the degree of 1 in  $R$  is two. Now, there are two possible subcases:

**Subcase 1:**  $\alpha'(b) = 1$  and  $\alpha'(x) = 2$ .

Any other neighbor  $y$  of  $x$  in  $G'$  must be colored 1. As per the assignment  $\alpha'$ ,  $y$  must have a neighbor colored 1 and another neighbor colored 2. Therefore,  $\alpha'$  restricted to  $G$  is an  $R$ -role coloring, and  $G$  is  $R$ -role colorable.

**Subcase 2:**  $\alpha'(b) = 2$  and  $\alpha'(x) = 1$

Suppose that all the other neighbors of  $x$  in  $G'$  are colored 2. Those neighbors would, in turn, be adjacent to vertices colored 1 in the partition of  $x$ . These vertices colored 1 must obey the coloring  $\alpha'$ , and have neighbors of both the colors 1 and 2. Once again, restricting  $\alpha'$  to  $G$ , gives an  $R$ -role coloring of  $G$ , showing that  $G$  is  $R$ -role colorable.

Symmetrically, if  $\alpha'(a) = 2$ , the role graph is again isomorphic to  $R$ . Hence, if  $G'$  has a 2-role coloring it has an  $R$ -role coloring and therefore  $G$  is  $R$ -role colorable.  $\square$

## 6 Conclusions

We have proved that  $k$ -ROLE COLORING, for a fixed  $k \geq 3$ , remains NP-complete even when restricted to the class of bipartite graphs. We have also shown that 2-ROLE COLORING, which is polynomial time solvable on bipartite graphs, becomes NP-complete on graphs that are a constant number of vertices or edges away from bipartite graphs. From the viewpoint of parameterized complexity, it means that 2-ROLE COLORING, when parameterized by the distance  $d$  from a graph class, is para-NP-hard on graphs that are  $d$ -away from bipartite graphs. To the best of our knowledge, this parameter has not been explored with respect to ROLE COLORING, before. This opens avenues for further research.

On a subclass of  $2K_2$ -free graphs, i.e, bipartite chain graphs, we have shown that 3-ROLE COLORING can be solved in polynomial time. However, the complexity of  $k$ -ROLE COLORING, for  $k=2$  and 3, is open on general  $2K_2$ -free graphs.

## Acknowledgements

The authors would like to thank Professor Venkatesh Raman for his invaluable comments.

## References

1. Cai, L.: Parameterized complexity of vertex colouring. *Discrete Applied Mathematics* **127**(3), 415 – 429 (2003)
2. Diestel, R.: *Graph Theory*. Springer-Verlag (2005)
3. Dourado, M.C.: Computing role assignments of split graphs. *Theor. Comput. Sci.* **635**, 74–84 (2016)
4. Fiala, J., Paulusma, D.: A complete complexity classification of the role assignment problem. *Theor. Comput. Sci.* **349**(1), 67–81 (2005)

5. Garey, M.R., Johnson, D.S.: *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W.H.Freeman and Company (1979)
6. Heggenes, P., van 't Hof, P., Paulusma, D.: Computing role assignments of proper interval graphs in polynomial time. *J. Discrete Algorithms* **14**, 173–188 (2012)
7. Heggenes, P., Kratsch, D.: Linear-time certifying recognition algorithms and forbidden induced subgraphs. *Nordic J. of Computing* **14**(1), 87–108 (jan 2007)
8. van't Hof, P., Paulusma, D., van Rooij, J.M.M.: Computing role assignments of chordal graphs. *Theor. Comput. Sci.* **411**(40-42), 3601–3613 (2010)
9. Martin G. Everett, S.B.: Role colouring a graph. *Mathematical Social Sciences* pp. 183–188 (1991)
10. Purcell, C., Rombach, M.P.: On the complexity of role colouring planar graphs, trees and cographs. *J. Discrete Algorithms* **35**, 1–8 (2015)
11. Roberts, F.S., Sheng, L.: How hard is it to determine if a graph has a 2-role assignment? *Networks* **37**, 67–73 (2001)
12. Sheng, L.: 2-role assignments on triangulated graphs. *Theor. Comput. Sci.* **304**(1-3), 201–214 (2003)
13. Yannakakis, M.: The complexity of the partial order dimension problem. *SIAM J. Algebraic Discrete Methods* **3**, 351–358 (1982)