

Instantons to the people: the power of one-form symmetries

Giulio Bonelli, Fran Globblek, Alessandro Tanzini

*International School of Advanced Studies (SISSA), via Bonomea 265,
34136 Trieste, Italy and INFN, Sezione di Trieste and
Institute for Geometry and Physics, IGAP, via Beirut 2, 34136 Trieste, Italy*

(Dated: February 3, 2021)

Abstract: We show that the non-perturbative dynamics of $\mathcal{N} = 2$ super Yang-Mills theories in a self-dual Ω -background and with an arbitrary simple gauge group is fully determined by studying renormalization group equations of vevs of surface operators generating one-form symmetries. The corresponding system of equations is a *non-autonomous* Toda chain, the time being the RG scale. We obtain new recurrence relations which provide a systematic algorithm computing multi-instanton corrections from the tree-level one-loop prepotential as the asymptotic boundary condition of the RGE. We exemplify by computing the E_6 and G_2 cases up to two-instantons.

In an ideal world the non-perturbative structure of gauge theories should be computed by quantum equations of motion determined by a symmetry principle. The presence of extended operators generating higher form symmetries in quantum field theory is a powerful tool to concretely realise such a programme. A perturbative analysis in a weakly coupled regime, if any, would supply appropriate asymptotic conditions. In this letter we present a class of theories where the full non-perturbative result is fixed in such a framework. These are $\mathcal{N} = 2$ super Yang-Mills theories in four dimensional self-dual Ω -background, which enjoy a one-form symmetry generated by surface operators [1]. We show that the renormalization group equation obeyed by the vacuum expectation value of such surface operators provides a recursion relation which fully determines, from the perturbative one-loop prepotential, all instanton contributions on the self-dual Ω -background or, equivalently, the all-genus topological string amplitudes on the relevant geometric background. Actually, partition functions with surface operators display a very clear resurgent structure led by the summation over the magnetic fluxes [2].

The system of equations we study is a *non-autonomous* twisted affine Toda chain of type $(\hat{G})^\vee$, where $(\hat{G})^\vee$ is the Langlands dual of the untwisted affine Kac-Moody algebra \hat{G} . Each node of the corresponding affine Dynkin diagram defines a surface operator, the associated τ -function being its vacuum expectation value. The time flow corresponds in the gauge theory to the renormalization group. The resulting recurrence relations constitute a new effective algorithm to determine instanton contributions for all classical groups G . Let us remark that the τ -functions we obtain provide the general solution at the canonical rays for the Jimbo-Miwa-Ueno isomonodromic deformation problem [3, 4] on the sphere with two-irregular punctures for all classical groups, which to the best of our knowledge was not known in the previous literature. The recursion relations we obtain are different from the blow-up equations of [5] further elaborated in [6]. Indeed the latter necessarily involve the knowledge of the partition function in different Ω -backgrounds. This

makes the recursion relations (and the results) coming from blow-up equations more involved and difficult to handle. However, we expect a relation between the two approaches to follow from blow-up relations in presence of surface defects. Indeed, the isomonodromic τ -function for the sphere with four regular punctures was obtained in a similar way from $SU(2)$ gauge theory with $N_f = 4$ in [7]. In this letter we summarise our results and refer to a subsequent longer paper for a fully detailed discussion.

The τ -functions are labeled by the simple roots of the affinization of the Lie algebra of the gauge group $\alpha \in \hat{\Delta}$, namely $\{\tau_\alpha\}_{\alpha \in \hat{\Delta}}$, and satisfy the equations

$$D^2(\tau_\beta) = -\frac{\beta^\vee \cdot \beta^\vee}{2} t^{1/h^\vee} \prod_{\beta \in \hat{\Delta}, \beta \neq \alpha} [\tau_\alpha]^{-\alpha \cdot \beta^\vee} \quad (1)$$

where $t := (\Lambda/\epsilon)^{2h^\vee}$ and the logarithmic Hirota derivative is given by $D^2(f) = f \partial_{\log t}^2 f - (\partial_{\log t} f)^2$. Given a simple root α , its coroot is as usual given by $\alpha^\vee = 2\alpha/(\alpha, \alpha)$, where (\cdot, \cdot) is the scalar product defined by the affine Cartan matrix. Eq. (1) is the de-autonomization of the τ -form of the standard Toda integrable system [8, 9] governing the classical Seiberg-Witten (SW) theory [10]. The de-autonomization is induced by coupling the theory to a self-dual Ω -background $(\epsilon_1, \epsilon_2) = (\epsilon, -\epsilon)$ [11]. In the autonomous limit $\epsilon \rightarrow 0$, τ -functions reduce to θ -functions on the classical SW curve [12], which were used to provide recursion relations on the coefficients of the SW prepotential in [13]. The gauge theory interpretation of these τ -functions is the v.e.v. of surface operators associated to the corresponding decomposition of the Lie algebra representation under which these are charged. We expect these equations and their generalizations to describe chiral ring relations in presence of a surface operator, which deserve further investigation. Higher chiral observables should generate the flows of the full *non-autonomous* Toda hierarchy. The actual form of equations (1) depends on the Dynkin diagram. For the classical groups A, B and D these reduce to bilinear equations which we solve via general recursion relations. For C, E, F and G the resulting equations are of higher

order and we study them case by case. The symmetries of the equations are given by the center of the group G , namely

$$\begin{array}{c|c|c|c|c|c|c|c|c} \mathfrak{g} & A_n & B_n & C_n & D_{2n} & D_{2n+1} & E_n & F_4 & G_2 \\ \hline Z(G) & \mathbb{Z}_{n+1} & \mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 \times \mathbb{Z}_2 & \mathbb{Z}_4 & \mathbb{Z}_{9-n} & 1 & 1 \end{array}$$

Moreover, the center is isomorphic to the coset of the affine coweight lattice by the affine coroot lattice, and coincides with the automorphism group of the affine Dynkin diagram. By a remark in [14], the coweights, and by extension the lattice cosets, corresponding to these nodes are the miniscule coweights, a representation of \mathfrak{g} being miniscule if all its weights form a single Weyl-orbit. This remark will be crucial while solving the τ -system.

The τ -functions corresponding to the affine nodes, that is the ones which can be removed from the Dynkin diagram leaving behind that of an irreducible simple Lie algebra, play a special rôle. Indeed, these are related to simple surface operators associated to elements of the center $Z(G)$, and are bounded by fractional 't Hooft lines. Such surface operators are the generators of the one-form symmetry of the corresponding gauge theory, [1]. Since their magnetic charge is defined modulo the magnetic root lattice, a natural Ansatz for their expectation value is

$$\tau_{\alpha_{\text{aff}}}(\sigma, \eta | \kappa_{\mathfrak{g}} t) = \sum_{\mathbf{n} \in Q_{\text{aff}}^{\vee}} e^{2\pi\sqrt{-1}\eta \cdot \mathbf{n}} t^{\frac{1}{2}(\sigma + \mathbf{n})^2} B(\sigma + \mathbf{n}|t) \quad (2)$$

where $B(\sigma|t) = B_0(\sigma) \sum_{i \geq 0} t^i Z_i(\sigma)$ with $Z_0(\sigma) \equiv 1$ and $Q_{\text{aff}}^{\vee} = \lambda_{\text{aff}}^{\vee} + Q^{\vee}$, Q^{\vee} being the co-root lattice and $(\lambda_{\text{aff}}^{\vee}, \alpha) = \delta_{\alpha_{\text{aff}}, \alpha}$ for any simple root α . The constant $\kappa_{\mathfrak{g}} = (-n_{\mathfrak{g}})^{r_{\mathfrak{g},s}}$, where $n_{\mathfrak{g}}$ is the ratio of the squares of long vs. short roots and $r_{\mathfrak{g},s}$ is the number of short simple roots. For simply laced, all roots are long and $\kappa_{\mathfrak{g}} = 1$.

We will now show how the term $t^{\frac{1}{2}\sigma^2} B(\sigma|t)$ in (2) is the full Nekrasov partition function in the self-dual Ω -background upon the identification $\sigma = \mathbf{a}/\epsilon$, where \mathbf{a} is the Cartan parameter. In the A_n case, (2) is known as the Kiev Ansatz. In the A_1 case, it was used to give the general solution of Painlevé III₃ equation in [15] and further analysed in [16].

Let us remark that the τ -function (2) displays a clear resurgent structure, with ‘‘instantons’’ given by the magnetic fluxes in the lattice summed with ‘‘resurgent’’ coefficients $B(\sigma|t)$ and trans-series parameter $e^{2\pi\sqrt{-1}\eta}$, see [17] for a similar analysis in the Painlevé III₃ case.

The Ansatz (2) is consistent with equations (1). Indeed, after eliminating the τ -functions associated to the non-affine nodes, the resulting equation is bilinear and therefore the Ansatz (2) reduces to a set of recursion relations for the coefficients $Z_i(\sigma)$. The variables η and σ are the integration constants of the second order differential equations (1) and correspond to the initial position and velocity of the de-autonomized Toda particle.

Let us set more precisely the boundary conditions which we impose to the solutions of equations (1). We consider the asymptotic behaviour of the solutions at $t \rightarrow 0$ and $\sigma \rightarrow \infty$ as


$$\log(B_0) \sim -\frac{1}{4} \sum_{\mathbf{r} \in R} (\mathbf{r} \cdot \sigma)^2 \log(\mathbf{r} \cdot \sigma)^2 \quad (3)$$

up to quadratic and log-terms [18]. We will show that the solution of (1) which satisfies the above asymptotic condition is such that

$$B_0(\sigma) = \mathcal{Z}_{1\text{-loop}}(\sigma) \equiv \prod_{\mathbf{r} \in R} \frac{1}{G(1 + \mathbf{r} \cdot \sigma)} \quad (4)$$

where $G(z)$ is the Barnes’ G-function and R is the adjoint representation of the group G . The expansion of the above function matches the one-loop gauge theory result upon the appropriate identification of the log-branch. This reads, in the gauge theory variables, as $\ln[\sqrt{-1}\mathbf{r} \cdot \mathbf{a}/\Lambda] \in \mathbb{R}$ and matches the canonical Stokes rays obtained in [19].

Let us first focus on the A_n case whose affine Dynkin diagram is



The root lattice is $Q = \{ \sum_{i=1}^{n+1} c_i e_i \mid \sum_{i=1}^{n+1} c_i = 0 \}$, and all the fundamental weights are miniscule, namely

$$\lambda_i = \frac{1}{n+1} (1^i, 0^{n+1-i}) - \frac{i}{n+1} (1^{n+1}),$$

where $(1^p, 0^{n+1-p})$ stands for a vector whose first p entries are 1 and the remaining entries vanish. We label the τ -functions as $\tau_{\alpha_j} \equiv \tau_j$. The τ -system is given by the closed chain of differential equations

$$D^2(\tau_j) = -t^{\frac{1}{n+1}} \tau_{j-1} \tau_{j+1}, \quad (5)$$

with $\tau_j = \tau_{n+1+j}$. Since all the nodes in this case are affine we can use the Kiev Ansatz (2). Then, all the τ -functions are determined by τ_0 as $\tau_j(\sigma|t) = \tau_0(\sigma + \lambda_j|t)$. It is therefore enough to solve the single equation

$$D^2(\tau_0(\sigma)) = -\tau_0(\sigma \pm e_1). \quad (6)$$

Here and in the following we use the notation $f(y \pm x) \equiv f(y+x)f(y-x)$. The Ansatz (2) for τ_0 reads

$$\tau_0(\sigma, \eta|t) = \sum_{\mathbf{n} \in Q, i \geq 0} e^{2\pi\sqrt{-1}\eta \cdot \mathbf{n}} t^{\frac{1}{2}(\sigma + \mathbf{n})^2 + i} B_0(\sigma + \mathbf{n}) Z_i(\sigma + \mathbf{n})$$

and by inserting it into (6) one gets after some simplifi-

cations

$$\begin{aligned}
& \sum_{\substack{\mathbf{n}_1, \mathbf{n}_2 \in Q \\ i_1, i_2 \geq 0}} e^{2\pi\sqrt{-1}(\mathbf{n}_1 + \mathbf{n}_2) \cdot \boldsymbol{\eta}} t^{\frac{1}{2}\mathbf{n}_1^2 + \frac{1}{2}\mathbf{n}_2^2 + i_1 + i_2 + \boldsymbol{\sigma} \cdot (\mathbf{n}_1 + \mathbf{n}_2)} \\
& \times \left(\frac{1}{2}\mathbf{n}_1^2 - \frac{1}{2}\mathbf{n}_2^2 + i_1 - i_2 + \boldsymbol{\sigma} \cdot (\mathbf{n}_1 - \mathbf{n}_2) \right)^2 \\
& \times B_0(\boldsymbol{\sigma} + \mathbf{n}_1) B_0(\boldsymbol{\sigma} + \mathbf{n}_2) Z_{i_1}(\boldsymbol{\sigma} + \mathbf{n}_1) Z_{i_2}(\boldsymbol{\sigma} + \mathbf{n}_2) \\
& = - \sum_{\substack{\mathbf{m}_1, \mathbf{m}_2 \in Q \\ j_1, j_2 \geq 0}} t^{1 + \frac{1}{2}\mathbf{m}_1^2 + \frac{1}{2}\mathbf{m}_2^2 + e_1 \cdot (\mathbf{m}_1 - \mathbf{m}_2) + j_1 + j_2 + \boldsymbol{\sigma} \cdot (\mathbf{m}_1 + \mathbf{m}_2)} \times \\
& e^{2\pi\sqrt{-1}(\mathbf{m}_1 + \mathbf{m}_2) \cdot \boldsymbol{\eta}} B_0(\boldsymbol{\sigma} + \mathbf{m}_1 + e_1) B_0(\boldsymbol{\sigma} + \mathbf{m}_2 - e_1) \times \\
& Z_{j_1}(\boldsymbol{\sigma} + \mathbf{m}_1 + e_1) Z_{j_2}(\boldsymbol{\sigma} + \mathbf{m}_2 - e_1) \quad (7)
\end{aligned}$$

Now we simply equate the exponents. To fix $B_0(\boldsymbol{\sigma})$, we look at the lowest order in t . This produces a quadratic constraint and $n+1$ linear constraints on the root lattice variables $(\mathbf{n}_1, \mathbf{n}_2)$ and $(\mathbf{m}_1, \mathbf{m}_2)$. Let us fix $p, q \in \{0, \dots, n+1\}$, $p \neq q$. Up to Weyl reflections, the only solution to the above mentioned constraints is given by $\mathbf{n}_1 = e_p - e_q$, $\mathbf{n}_2 = 0$ and $\mathbf{m}_1 = e_p - e_1$, $\mathbf{m}_2 = -e_q + e_1$, leading to

$$\begin{aligned}
& (1 + (e_p - e_q) \cdot \boldsymbol{\sigma})^2 B_0(\boldsymbol{\sigma} + e_p - e_q) B_0(\boldsymbol{\sigma}) = \\
& -B_0(\boldsymbol{\sigma} + e_p) B_0(\boldsymbol{\sigma} - e_q). \quad (8)
\end{aligned}$$

This is solved by (4) up to a function periodic on the root lattice, which is set to one by the asymptotic condition (3). The higher order terms in (7) provide the recursion relations

$$\begin{aligned}
k^2 Z_k(\boldsymbol{\sigma}) = & - \sum_{\substack{\mathbf{n}^2 + j_1 + j_2 = k \\ \mathbf{n} \in e_1 + Q, j_1, j_2 < k}} \frac{B_0(\boldsymbol{\sigma} \pm \mathbf{n})}{B_0(\boldsymbol{\sigma})^2} \times \\
& Z_{j_2}(\boldsymbol{\sigma} - \mathbf{n}) Z_{j_1}(\boldsymbol{\sigma} + \mathbf{n}) + \sum_{\substack{\mathbf{n}^2 + i_1 + i_2 = k \\ \mathbf{n} \in Q, i_1, i_2 < k}} (i_1 - i_2 + 2\mathbf{n} \cdot \boldsymbol{\sigma})^2 \\
& \times \frac{B_0(\boldsymbol{\sigma} \pm \mathbf{n})}{B_0(\boldsymbol{\sigma})^2} Z_{i_1}(\boldsymbol{\sigma} + \mathbf{n}) Z_{i_2}(\boldsymbol{\sigma} - \mathbf{n}),
\end{aligned}$$


where $B_0(\boldsymbol{\sigma})$ is given by (4). For $k=1$ we easily obtain

$$Z_1(\boldsymbol{\sigma}) = - \sum_{i=1}^{n+1} \frac{B_0(\boldsymbol{\sigma} \pm e_i)}{B_0(\boldsymbol{\sigma})^2} = (-1)^{n+1} \sum_{i=1}^{n+1} \frac{1}{\prod_{j \neq i} (\sigma_i - \sigma_j)^2}$$

and, upon abbreviating $\sigma_{ij} = \sigma_i - \sigma_j$, the next term

$$\begin{aligned}
Z_2(\boldsymbol{\sigma}) = & -\frac{1}{4} \sum_{i=1}^{n+1} \frac{B_0(\boldsymbol{\sigma} \pm e_i)}{B_0(\boldsymbol{\sigma})^2} [Z_1(\boldsymbol{\sigma} + e_i) + Z_1(\boldsymbol{\sigma} - e_i)] \\
& + \sum_{i < j}^{n+1} (\sigma_i - \sigma_j)^2 \frac{B_0(\boldsymbol{\sigma} \pm (e_i - e_j))}{B_0(\boldsymbol{\sigma})^2}
\end{aligned}$$


The above coincide with one and two instanton contributions to the $SU(n+1)$ Nekrasov partition function as computed from supersymmetric localization [20, 21]. Let us remark that the use of the τ -system (5) provides a completely independent tool to compute all instanton corrections just starting from the asymptotic behaviour (3). This procedure extends to all classical groups.



D_n is a simply laced root system, with the checkerboard lattice $Q = Q^\vee = \{\sum_{i=1}^n c_i e_i \mid \sum_{i=1}^n c_i \in 2\mathbb{Z}\}$. We consider $n > 4$. It has four miniscule weights, $\lambda_0 = (0^n)$, $\lambda_1 = (1, 0^{n-1})$, $\lambda_{n-1} = ((\frac{1}{2})^{n-1}, -\frac{1}{2})$, $\lambda_n = ((\frac{1}{2})^{n-1}, +\frac{1}{2})$. These correspond to the "legs" of the affine diagram. Whichever rank we consider, we always have the consistency conditions

$$D^2(\tau_0) = D^2(\tau_1), \quad D^2(\tau_{n-1}) = D^2(\tau_n) \quad (9)$$

which are also equal if $n=4$, due to the enhanced symmetry of D_4 .



B_n is non-simply laced. The coroot lattice is the checkerboard lattice $Q^\vee = \{\sum_{i=1}^n c_i e_i \mid \sum_{i=1}^n c_i \in 2\mathbb{Z}\}$, and the two miniscule weights are $\lambda_0^\vee = (0^n)$ and $\lambda_1^\vee = (1, 0^{n-1})$, corresponding to the "antennae" of the diagram. The τ -system coincides with that of D_{n+1} , with the modification that (i) there is no τ_{n+1} and (ii) that

$$D^2(\tau_{n-1}) = -2t^{\frac{1}{2n-1}} \tau_{n-2} \tau_n, \quad D^2(\tau_n) = -t^{\frac{1}{2n-1}} \tau_{n-1}^2.$$

For $n \geq 3$, the analysis proceeds as for D_n except we may only use the left antennae and consider the first equation in (9). Therefore, we have a unified approach for both D_n and B_n . Explicitly, inserting (2) and $\tau_1(\boldsymbol{\sigma}|t) = \tau_0(\boldsymbol{\sigma} + \lambda_1|t)$ into the first of (9) we get after some simplification a formula analogous to (7) leading to quadratic and linear constraints on the lattice labels. By repeating the analysis similarly to the previous case, the equation, analogous to (8), fixing B_0 is

$$\begin{aligned}
& (1 + (e_p + e_q) \cdot \boldsymbol{\sigma})^2 B_0(\boldsymbol{\sigma}) B_0(\boldsymbol{\sigma} + e_p + e_q) \\
& = ((e_p - e_q) \cdot \boldsymbol{\sigma})^2 B_0(\boldsymbol{\sigma} + e_p) B_0(\boldsymbol{\sigma} + e_q). \quad (10)
\end{aligned}$$

The two cases are distinguished by the corresponding different asymptotic conditions (3). Indeed, we have

$$\begin{aligned}
B_0^{[D_n]}(\boldsymbol{\sigma}) &= \prod_{i < j}^n \frac{1}{G(1 \pm \sigma_i \pm \sigma_j)} \\
B_0^{[B_n]}(\boldsymbol{\sigma}) &= \left(\prod_{k=1}^n \frac{1}{G(1 \pm \sigma_k)} \right) B_0^{[D_n]}(\boldsymbol{\sigma})
\end{aligned}$$

Also the recursion relations are the same, upon using the

appropriate root systems R :

$$k^2 Z_k(\boldsymbol{\sigma}) = \sum_{\substack{(\mathbf{n}-\boldsymbol{\lambda}_1)^2+j_1+j_2=k \\ \mathbf{n} \in \boldsymbol{\lambda}_1+Q, j_{1,2}<k}} Z_{j_1}(\boldsymbol{\sigma}+\mathbf{n})Z_{j_2}(\boldsymbol{\sigma}-\mathbf{n}) \\ (j_1-j_2+2\mathbf{n}\cdot\boldsymbol{\sigma})^2 \frac{B_0(\boldsymbol{\sigma}\pm\mathbf{n})}{B_0(\boldsymbol{\sigma})^2} - \sum_{\substack{\mathbf{n}^2+i_1+i_2=k \\ \mathbf{n} \in Q, i_{1,2}<k}} Z_{j_1}(\boldsymbol{\sigma}+\mathbf{n}) \\ \times Z_{j_2}(\boldsymbol{\sigma}-\mathbf{n}) (i_1-i_2+2\mathbf{n}\cdot\boldsymbol{\sigma})^2 \frac{B_0(\boldsymbol{\sigma}\pm\mathbf{n})}{B_0(\boldsymbol{\sigma})^2}$$

This result is in line with the contour integral formulae for the relevant Nekrasov partition functions. Indeed the poles in the D_n and B_n cases are the same, with different residues. From the above recursion relation we can compute the 1-instanton terms


$$Z_1(\boldsymbol{\sigma}) = \sum_{k=1}^n 4\sigma_k^2 \frac{B_0(\boldsymbol{\sigma}\pm e_k)}{B_0(\boldsymbol{\sigma})^2} = \begin{cases} \sum_{k=1}^n \frac{-1}{\prod_{j \neq k} (\sigma_k^2 - \sigma_j^2)^2}, & B_n \\ \sum_{k=1}^n \frac{4\sigma_k^2}{\prod_{j \neq k} (\sigma_k^2 - \sigma_j^2)^2}, & D_n \end{cases}$$

and the 2-instantons

$$Z_2(\boldsymbol{\sigma}) = \sum_{\alpha \in Q^\vee, \alpha^2=2} \frac{-1}{(\alpha \cdot \boldsymbol{\sigma})^2 ((\alpha \cdot \boldsymbol{\sigma})^2 - 1)^2 \prod_{\beta \cdot \alpha=1} (\beta \cdot \boldsymbol{\sigma})^2} \\ + \sum_{k=1}^n \frac{Z_1(\boldsymbol{\sigma}+e_k)(\sigma_k + \frac{1}{2})^2 + Z_1(\boldsymbol{\sigma}-e_k)(\sigma_k - \frac{1}{2})^2}{\prod_{\beta \cdot e_k=\pm 1} (\beta \cdot \boldsymbol{\sigma})^2}$$

and so on. These are easily compared to [22].

We now turn to the analysis of the other classical groups, which is more involved. Indeed, the τ -system reduces to higher order equations which produce more complicated recurrence relations to be solved by a case by case analysis. We performed explicit checks for C_3 , C_4 and C_5 up to two-instantons again in agreement with [22].



For the exceptional group E_6 we obtain the system

$$\tau_6 D^4(\tau_0) = \tau_0 D^4(\tau_6). \quad (11)$$

where we used the notation $D^{2n} := D^2 \circ D^{2n-2}$. The equations which specify B_0 can be written as follows. Choose the miniscule weight to be $\boldsymbol{\lambda} = (0^5, (-\frac{2}{3})^3)$. Let p_1, \dots, p_5 be a permutation of $\{1, \dots, 5\}$ and let $\boldsymbol{\delta} := ((\frac{1}{2})^8)$. Then one gets from the lowest order in (11)


$$(1 + \sigma_{p_1} + \sigma_{p_2})^2 (1 + \sigma_{p_1} + \sigma_{p_3})^2 (\sigma_{p_2} - \sigma_{p_3})^2 \times \\ B_0(\boldsymbol{\sigma}) B_0(\boldsymbol{\sigma} + e_{p_1} + e_{p_2}) B_0(\boldsymbol{\sigma} + e_{p_1} + e_{p_3}) = \\ ((\boldsymbol{\delta} - e_{p_2} - e_{p_3}) \cdot \boldsymbol{\sigma})^2 ((\boldsymbol{\delta} - e_{p_2} - e_{p_3} - e_{p_4} - e_{p_5}) \cdot \boldsymbol{\sigma})^2 \times \\ (\sigma_{p_4} + \sigma_{p_5})^2 B_0(\boldsymbol{\sigma} + \boldsymbol{\delta} + \boldsymbol{\lambda}) \times \\ B_0(\boldsymbol{\sigma} + \boldsymbol{\delta} + \boldsymbol{\lambda} - e_{p_4} - e_{p_5}) B_0(\boldsymbol{\sigma} + e_{p_1} - \boldsymbol{\lambda}/2)$$

The solution satisfying the asymptotic behaviour (3) is

$$B_0^{[E_6]} = \prod_{i < j=1}^5 \frac{1}{G(1 \pm \sigma_i \pm \sigma_j)} \prod_{\substack{\varepsilon_i = \pm 1 \\ \prod_{i=1}^8 \varepsilon_i = 1, \\ \varepsilon_6 = \varepsilon_7 = \varepsilon_8}} \frac{1}{G(1 + \frac{1}{2} \sum_{i=1}^8 \varepsilon_i \sigma_i)}$$

We also solved the recurrence relation arising from (11) up to two-instantons. For one-instanton, our results agree with the ones of [23], while the two instantons result is a too huge formula to be reported here. We remark that (11) represents a completely novel way of obtaining equivariant volumes of instanton moduli spaces for exceptional groups.

Unimodular algebras G_2, F_4, E_8 have no outer automorphisms and consequently all the τ -functions associated to different nodes are independent. Therefore, the equations on the τ -function associated to the affine node turn out to be more difficult to solve. Let us display them for the G_2 case.



In the normalization where its longest root has length 2, the G_2 coroot lattice is the span $Q^\vee = \mathbb{Z} \frac{1}{\sqrt{3}}(-2, 1, 1) \oplus \mathbb{Z} \sqrt{3}(1, -1, 0)$. We introduce $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ but all expressions should be restricted to $\sigma_1 + \sigma_2 + \sigma_3 = 0$. By eliminating τ_1 and τ_2 , the τ -system reduces to the single equation

$$D^2(\tau_0^{-1} D^4(\tau_0)) = 3t(D^2(\tau_0))^3. \quad (12)$$

The operator on the l.h.s. of (12) turns out to factorize as $D^2(\tau_0^{-1} D^4(\tau_0)) = \tilde{D}^4(\tau_0) \cdot D^2(\tau_0)$, where $\tilde{D}^4(\tau_0)$ is a fourth order operator in τ_0 and its derivatives. The trivial solution of $D^2(\tau_0) = 0$ is $\tau_0 = at^b$ which we discard being incompatible with (2). In the remainder we insert

$$\tau_0(\boldsymbol{\sigma}, \boldsymbol{\eta}|t) = \sum_{\mathbf{n} \in Q^\vee} e^{2\pi\sqrt{-1}\boldsymbol{\eta}\cdot\mathbf{n}} \left(-\frac{t}{3}\right)^{\frac{1}{2}(\boldsymbol{\sigma}+\mathbf{n})^2} B(\boldsymbol{\sigma}+\mathbf{n} | -\frac{t}{3})$$

and obtain, after a rescaling $t \mapsto -3t$,

$$\sum_{\substack{\{\mathbf{n}_k\} \in Q^\vee \\ \{i_k\} \in \mathbb{N}}} \prod_{k=1}^4 e^{2\pi\sqrt{-1}\boldsymbol{\eta}\cdot\mathbf{n}_k} t^{\frac{1}{2}(\boldsymbol{\sigma}+\mathbf{n}_k)^2 + i_k} B_0(\boldsymbol{\sigma}+\mathbf{n}_k) Z_{i_k}(\boldsymbol{\sigma}+\mathbf{n}_k) \\ \left(\frac{1}{4!} \prod_{k_1 < k_2} \left(\frac{1}{2} \mathbf{n}_{k_1}^2 + i_{k_1} - \frac{1}{2} \mathbf{n}_{k_2}^2 - i_{k_2} + (\mathbf{n}_{k_1} - \mathbf{n}_{k_2}) \cdot \boldsymbol{\sigma} \right)^2 \right. \\ \left. + \frac{9}{4} \left(\frac{1}{2} \mathbf{n}_1^2 + i_1 - \frac{1}{2} \mathbf{n}_2^2 - i_2 + (\mathbf{n}_1 - \mathbf{n}_2) \cdot \boldsymbol{\sigma} \right)^2 \right. \\ \left. \left(\frac{1}{2} \mathbf{n}_3^2 + i_3 - \frac{1}{2} \mathbf{n}_4^2 - i_4 + (\mathbf{n}_3 - \mathbf{n}_4) \cdot \boldsymbol{\sigma} \right)^2 \right) = 0. \quad (13)$$

The lowest order terms in (13), namely the coefficient of $t^{3+\boldsymbol{\sigma}\cdot(\frac{4}{\sqrt{3}}, -\frac{2}{\sqrt{3}}, -\frac{2}{\sqrt{3}})}$, gives a quartic relation which simpli-

fies to the following quadratic one

$$\begin{aligned} & \left(\frac{2\sigma_1 - \sigma_2 - \sigma_3}{\sqrt{3}} + 1 \right)^2 B_0(\boldsymbol{\sigma}) B_0 \left(\boldsymbol{\sigma} + \frac{1}{\sqrt{3}}(2, -1, -1) \right) \\ &= - \left(\frac{\sigma_2 - \sigma_3}{\sqrt{3}} \right)^2 \left(\frac{\sigma_1 + \sigma_2 - 2\sigma_3}{\sqrt{3}} \right)^2 \left(\frac{\sigma_1 - 2\sigma_2 + \sigma_3}{\sqrt{3}} \right)^2 \\ & \times \left(\frac{\sigma_1 + \sigma_2 - 2\sigma_3}{\sqrt{3}} + 1 \right)^2 \left(\frac{\sigma_1 - 2\sigma_2 + \sigma_3}{\sqrt{3}} + 1 \right)^2 \\ & \times B_0 \left(\boldsymbol{\sigma} + \frac{1}{\sqrt{3}}(1, -2, 1) \right) B_0 \left(\boldsymbol{\sigma} + \frac{1}{\sqrt{3}}(1, 1, -2) \right) \end{aligned}$$

By imposing (3), these are solved by $B_0^{[G_2]}(\boldsymbol{\sigma}) =$

$$\prod_{i < j}^3 \frac{1}{G(1 \pm \frac{1}{\sqrt{3}}(\sigma_i - \sigma_j))} \prod_{\substack{ijk \\ \text{cyclic}}}^3 \frac{1}{G(1 \pm \frac{1}{\sqrt{3}}(2\sigma_i - \sigma_j - \sigma_k))}$$

The 1-instanton contribution is obtained by considering the coefficient of the next order $t^{3+\boldsymbol{\sigma} \cdot (\sqrt{3}, 0, -\sqrt{3})}$ term: all $B_0(\boldsymbol{\sigma})$ factors drop out and we obtain just

$$Z_1(\boldsymbol{\sigma})^{[G_2]}|_{\sigma_3 = -\sigma_1 - \sigma_2} = -\frac{2}{3\sigma_1^2 \sigma_2^2 (\sigma_1 + \sigma_2)^2}$$

in agreement with [23]. The next order in t gives the 2-instanton term $Z_2(\boldsymbol{\sigma})^{[G_2]}|_{\sigma_3 = -\sigma_1 - \sigma_2} =$

$$\frac{3(9\sigma_1^4(6\sigma_2^2+1)+18\sigma_1^3(6\sigma_2^3+\sigma_2)+3\sigma_1^2(18\sigma_2^4+9\sigma_2^3-2)+6\sigma_1\sigma_2(3\sigma_2^2-1)+(1-3\sigma_2^2)^2)}{\sigma_1^2(1-3\sigma_1^2)^2\sigma_2^2(1-3\sigma_2^2)^2(\sigma_1+\sigma_2)^2(1-3(\sigma_1+\sigma_2)^2)^2}.$$

It should be possible to apply the approach proposed here to general class- \mathcal{S} theories [24] by studying the related isomonodromic deformation problem (for example for linear and circular quivers). It would be also interesting to extend the analysis to non-self-dual Ω -background, which should amount to quantum τ -systems, and its lift to five dimensional gauge theories on $\mathbb{R}^4 \times S^1$, which should correspond to q -difference τ -systems [25–27]. Finally, it would be great to apply similar ideas to models with less or no supersymmetry, trying to constrain their dynamics by the study of renormalization group equations in presence of surface operators.

Acknowledgements: We would like to thank M. Mariño and T. Nosaka for fruitful discussions. This research is partially supported by the INFN Research Projects GAST and ST&FI, by PRIN "Geometria delle varietà algebriche" and by PRIN "Non-perturbative Aspects Of Gauge Theories And Strings".

[1] Gaiotto, D., Kapustin, A., Seiberg, N., and Willett, B., *JHEP* **02**, 172 (2015), [arXiv:1412.5148 \[hep-th\]](#).

[2] For an introduction to resurgence in QFT, see for example M. Marino, *Instantons and Large N*, (2015) Cambridge University Press.

- [3] Jimbo, M., Miwa, T., and Ueno, a. K., *Physica* **D2**, 306 (1981).
- [4] Jimbo, M., Miwa, T., and Ueno, a. K., *Physica* **D2**, 407 (1982).
- [5] Nakajima, H. and Yoshioka, K., *Invent. Math.* **162**, 313 (2005), [arXiv:math/0306198 \[math.AG\]](#).
- [6] J. Kim, S.-S. Kim, K.-H. L. K. L. and Song, J., *JHEP* **11**, 092 (2019), [Erratum: *JHEP* 06, 124 (2020)], [arXiv:1908.11276 \[hep-th\]](#).
- [7] Nekrasov, N., (2020), [arXiv:2007.03646 \[hep-th\]](#).
- [8] Gorsky, A., Krichever, I., Marshakov, A., Mironov, A., and Morozov, A., *Phys. Lett.* **B355**, 466 (1995), [arXiv:hep-th/9505035 \[hep-th\]](#).
- [9] Martinec, E. J. and Warner, N. P., *Nucl. Phys. B* **459**, 97 (1996), [arXiv:hep-th/9509161](#).
- [10] Seiberg, N. and Witten, E., *Nucl. Phys.* **B426**, 19 (1994), [Erratum: *Nucl. Phys.* B430,485(1994)], [arXiv:hep-th/9407087 \[hep-th\]](#).
- [11] Bonelli, G., Lisovyy, O., Maruyoshi, K., Sciarappa, A., and Tanzini, A., *Letters in Mathematical Physics* **107**, 2359 (2017), [arXiv:1612.06235 \[hep-th\]](#).
- [12] Bonelli, G., Del Monte, F., Gavrylenko, P., and Tanzini, A., *Commun. Math. Phys.* **377**, 1381 (2020), [arXiv:1901.10497 \[hep-th\]](#).
- [13] Edelstein, J. D., Marino, M., and Mas, J., *Nucl. Phys. B* **541**, 671 (1999), [arXiv:hep-th/9805172](#).
- [14] Bourbaki [Lie gps Ch. VIII §7].
- [15] Its, A., Lisovyy, O., and Tykhyi, Y., (2014), [10.1093/imrn/rnu209](#), [arXiv:1403.1235 \[math-ph\]](#).
- [16] Mironov, A. and Morozov, A., *Phys. Lett. B* **773**, 34 (2017), [arXiv:1707.02443 \[hep-th\]](#).
- [17] Dunne, G. V., *J. Phys. A* **52**, 463001 (2019), [arXiv:1901.02076 \[hep-th\]](#).
- [18] Notice that if one chooses $\log(B_0) \sim \sum_{\mathbf{r} \in R} c_{\mathbf{r},m}(\mathbf{r} \cdot \boldsymbol{\sigma})^{2n} \log((\mathbf{r} \cdot \boldsymbol{\sigma})^2)^m$, then the equation itself dictates that $(n, m) = (1, 1)$ and $(2, 0)$.
- [19] Guest, M. A., Its, A. R., and Lin, C.-S., (2012), [arXiv:1209.2045 \[math.DG\]](#).
- [20] Nekrasov, N. A., in *International Congress of Mathematicians (ICM 2002) Beijing, China, August 20-28, 2002* (2003) [arXiv:hep-th/0306211 \[hep-th\]](#).
- [21] Nekrasov, N. and Okounkov, A., *Prog. Math.* **244**, 525 (2006), [arXiv:hep-th/0306238 \[hep-th\]](#).
- [22] Marino, M. and Wyllard, N., *JHEP* **05**, 021 (2004), [arXiv:hep-th/0404125](#).
- [23] Keller, C. A., Mekareeya, N., Song, J., and Tachikawa, Y., *JHEP* **03**, 045 (2012), [arXiv:1111.5624 \[hep-th\]](#).
- [24] Gaiotto, D., *JHEP* **08**, 034 (2012), [arXiv:0904.2715 \[hep-th\]](#).
- [25] Bonelli, G., Grassi, A., and Tanzini, A., *Lett. Math. Phys.* **109**, 1961 (2019), [arXiv:1710.11603 \[hep-th\]](#).
- [26] Bershtein, M., Gavrylenko, P., and Marshakov, A., (2018), [arXiv:1804.10145 \[math-ph\]](#).
- [27] Bonelli, G., Del Monte, F., and Tanzini, A., (2020), [arXiv:2007.11596 \[hep-th\]](#).