

Polynomial approximation of piecewise analytic functions on quasi-smooth arcs

Liudmyla Kryvonos

Abstract

For a function f that is piecewise analytic on a quasi-smooth arc \mathcal{L} and any $0 < \sigma < 1$ we construct a sequence of polynomials that converge at a rate e^{-n^σ} at each point of analyticity of f and are close to the best polynomial approximants on the whole \mathcal{L} . Moreover, we give examples when such polynomials can be constructed for $\sigma = 1$.

MSC: 30E10

Keywords: Polynomial approximation, Quasi-smooth arcs, Near-best approximation.

1 Introduction and main results

Let \mathcal{L} be a quasi-smooth arc on the complex plane \mathbf{C} , that is, for any $z, \zeta \in \mathcal{L}$ the length $|\mathcal{L}(z, \zeta)|$ of the subarc $\mathcal{L}(z, \zeta)$ of \mathcal{L} between points z, ζ satisfies

$$|\mathcal{L}(z, \zeta)| \leq c|z - \zeta|$$

for some $c = c(\mathcal{L}) \geq 1$.

Consider a piecewise analytic function f on \mathcal{L} belonging to $C^k(\mathcal{L})$, $k \geq 0$, that means f is k times continuously differentiable on \mathcal{L} and there exist points z_2, z_3, \dots, z_{m-1} such that f is analytic on $\mathcal{L} \setminus \{z_1, z_2, \dots, z_m\}$, (z_1, z_m – endpoints of \mathcal{L}), but is not analytic at points z_1, z_2, \dots, z_m . We call the z_i points of singularity of f .

The rate of the best uniform approximation of a function f by polynomials of degree at most $n \in \mathbb{N} := \{1, 2, \dots\}$ is denoted by

$$E_n(f) = E_n(f, \mathcal{L}) := \inf_{P_n: \deg P_n \leq n} \|f - P_n\|_{\mathcal{L}}. \quad (1.1)$$

Here $\|\cdot\|_{\mathcal{L}}$ means the supremum norm over \mathcal{L} . Also, let $p_n^*(f, z)$ be the (unique) polynomial minimizing the uniform norm in (1.1).

It is natural to expect the difference $f(z) - p_n^*(z)$ to converge faster at points of analyticity of f . But, it turns out, singularities of f adversely affect the behavior over the whole \mathcal{L} of a subsequence of the best polynomial approximants $p_n^*(f, z)$. This so-called "principle of contamination" manifests itself in density of extreme points of $f - p_n^*$, discussed by A. Kroo' and E.B. Saff in [8] and

accumulation of zeros of $p_n^*(f, z)$, showed by H.-P. Blatt and E.B. Saff in [6]. For more details, we refer the reader to [10].

Surprisingly, such behavior of zeros and extreme points need not hold for polynomials of "near-best" approximation, that is for polynomials P_n that satisfy

$$\|f - P_n\|_{\mathcal{L}} \leq CE_n(f), \quad n = 1, 2, \dots,$$

with a fixed $C > 1$. Hence, it is natural to seek "near-best" polynomials which would converge faster at points $z \in \mathcal{L} \setminus \{z_1, z_2, \dots, z_m\}$.

For the case of $\mathcal{L} = [-1, 1]$ and a piecewise analytic function f belonging to $C^k[-1, 1]$, E.B. Saff and V. Totik in [12] have proved that if non-negative numbers α, β satisfy $\alpha < 1$ and $\beta \geq \alpha$ or $\alpha = 1$ and $\beta > 1$, then there exist constants $c, C > 0$ and polynomials $P_n, n = 1, 2, \dots$, such that for every $x \in [-1, 1]$

$$|f(x) - P_n(x)| \leq CE_n(f)e^{-cn^\alpha d(x)^\beta}, \quad (1.2)$$

where $d(x)$ denotes the distance from x to the nearest singularity of f in $(-1, 1)$.

Accordingly, the question of constructing "near-best" polynomials arises when $[-1, 1]$ is replaced by an arbitrary quasi-smooth arc \mathcal{L} in \mathbf{C} . Polynomial approximation of functions on arcs is an important case of a more general problem of approximation of functions on an arbitrary continuum of the complex plane studied in the works of N.A. Shirokov [13], V.K. Dzjadyk and G.A. Alibekov [1], V.V. Andrievskii [3] and others (see, for example, [7]).

The behavior of "near-best" polynomials is well studied in the case of approximation on compact sets K with non-empty interior $Int(K)$. The following results demonstrate how the possible rate of convergence inside K depends on the geometry of K . V.V. Maimeskul have proved in [9] that if $\Omega := \overline{C} \setminus K$ satisfies the α -wedge condition with $0 < \alpha \leq 1$, then for any $\sigma < \alpha/2$ there exist "near-best" polynomials converging at a rate e^{-n^σ} in the interior of K . E.B. Saff and V. Totik in [11] show the possibility of geometric convergence of "near-best" polynomials inside K if the boundary of K is an analytic curve. Meantime, N.A. Shirokov and V. Totik in [14] discuss the rate of approximation by "near best" polynomials of a function f given on a compact set K with a generalized external angle smaller than π at some point $z_0 \in \partial K$. They showed that if f has a singularity at z_0 , then geometric convergence inside K , where f is analytic, is impossible. Taking into account these results, the most interesting case for us is when singularities of the function f occur at points where the angle between subarcs of \mathcal{L} is different from π . It turns out that for some such arcs there are no restrictions on the rate of convergence of "near-best" polynomials and it can be geometric at points where f is analytic, as opposed to the result for compact sets with non-empty interior. We formulate and prove this assertion in Theorem 2. Furthermore, the general case is given by the following

Theorem 1. *Let f be a piecewise analytic function on a quasi-smooth arc \mathcal{L} , i.e. there exist points $z_2, \dots, z_{m-1} \in \mathcal{L}$, such that they divide \mathcal{L} into $\mathcal{L}^1, \mathcal{L}^2, \dots, \mathcal{L}^{m-1}$ and*

$$f(z) = f_i(z), \quad z \in \mathcal{L}^i, \quad i = \overline{1, m-1}, \quad (1.3)$$

where $f_i(z)$ are analytic in some neighborhood of \mathcal{L}^i , respectively, and satisfy

$$f_{i-1}^{(r)}(z_i) = f_i^{(r)}(z_i), \quad f_{i-1}^{(k_i)}(z_i) \neq f_i^{(k_i)}(z_i) \quad (1.4)$$


for $r = \overline{0, k_i}$, $i = \overline{2, m-1}$. Then, for any $0 < \sigma < 1$, there exists a sequence $\{P_n\}_1^\infty$ of "near-best" polynomial approximants of f on \mathcal{L} , such that

$$\lim_{n \rightarrow \infty} \|f - P_n\|_E e^{n^\sigma} = 0 \quad (1.5)$$

holds for any compact set $E \subset \mathcal{L} \setminus \{z_2, \dots, z_{m-1}\}$.

On the complex plane consider lemniscates that are level lines of some complex polynomials. Namely, take $P(z) = P_N(z) := (z - a_1)(z - a_2)\dots(z - a_N)$, where $a_k = Re^{i\frac{2\pi(k-1)}{N}}$, $k = \overline{1, N}$ and $R > 0$ is a fixed number. Then $|P(z)| = R^N$ is an equation of a lemniscate. Note that the origin is a point of this lemniscate (since $|P(0)| = R^N$).

The lemniscate divides the plane into three parts, namely the curve itself, points $\{z : |P(z)| < R^N\}$ and $\{z : |P(z)| > R^N\}$. Consider an arc $\mathcal{L} = \mathcal{L}' \cup \mathcal{L}''$, where $\mathcal{L}', \mathcal{L}''$ may belong to different petals of the lemniscate, meet at the origin and satisfy $|P(z)| < R^N$, $z \in \mathcal{L} \setminus \{0\}$. An example for $N = 4$, $R = 1$ you can see below.



In particular, two line segments meeting at the origin at angle $0 < \varphi \leq \pi$ satisfy this property: if $\frac{2\pi}{m+1} < \varphi \leq \frac{2\pi}{m}$ for some integer m , it is enough to take R to be sufficiently large and $N = m$.

Let f be a piecewise analytic function on \mathcal{L} given by

$$f(z) = \begin{cases} f_1(z), & \text{if } z \in \mathcal{L}' \\ f_2(z), & \text{if } z \in \mathcal{L}'' \end{cases}$$

where f_1, f_2 are functions, analytic on \mathcal{L}' and \mathcal{L}'' correspondingly, satisfying

$$f_1^{(r)}(0) = f_2^{(r)}(0), \quad r = \overline{0, k}, \quad f_1^{(k+1)}(0) \neq f_2^{(k+1)}(0).$$

With these assumptions we prove the following result

Theorem 2. *Let \mathcal{L} and f be as above. Then there exist a constant $c > 0$ and a sequence of "near-best" polynomials $\{P_n\}_1^\infty$, such that*

$$\lim_{n \rightarrow \infty} \|f - P_n\|_E e^{cnd(E)} = 0,$$

where $d(E) > 0$ for any compact set $E \subset \mathcal{L} \setminus \{0\}$.

2 Auxiliary results

In this section we give some results which allow us to get estimates for the $E_n(f)$ and are needed for constructing "near-best" polynomials.

For $a > 0$ and $b > 0$ we will use the notation $a \preccurlyeq b$ if $a \leq cb$, with some constant $c > 0$. The expression $a \asymp b$ means $a \preccurlyeq b$ and $b \preccurlyeq a$.

Let \mathcal{L} be a quasi-smooth arc and $\Omega := \overline{\mathbb{C}} \setminus \mathcal{L}$. Consider a conformal mapping $\Phi : \Omega \rightarrow \Delta := \{\omega : |\omega| > 1\}$, normalized in such a way that $\Phi(\infty) = \infty$, $\Phi'(\infty) > 0$, and denote $\Psi := \Phi^{-1}$.

By $\tilde{\Omega}$ we denote compactification of the domain Ω by prime ends in the Caratheodory sense, and $\tilde{\mathcal{L}} := \tilde{\Omega} \setminus \Omega$. For the endpoints z_1, z_2 of \mathcal{L} and $u > 0, j = 1, 2$, let

$$\begin{aligned} \Phi(z_j) &:= \tau_j; \\ \Delta_1 &:= \{\tau : \tau \in \Delta, \arg \tau_1 < \arg \tau < \arg \tau_2\}; \\ \Delta_2 &:= \Delta \setminus \overline{\Delta_1}, \quad \tilde{\Omega}^j := \Psi(\overline{\Delta_j}), \quad \Omega^j := \Psi(\Delta_j); \\ \tilde{\mathcal{L}}^j &:= \tilde{\Omega}^j \cap \tilde{\mathcal{L}}; \\ \mathcal{L}_u^j &:= \{\zeta : \zeta \in \tilde{\Omega}^j, |\Phi(\zeta)| = 1 + u\}; \\ \rho_u^j(z) &:= \text{dist}(z, \mathcal{L}_u^j); \quad \rho_u^*(z) := \max_{j=1,2} \rho_u^j(z). \end{aligned}$$

Let z_0 be a point of \mathcal{L} , distinct from endpoints of the arc. Then point z_0 divides \mathcal{L} into two parts, \mathcal{L}' and \mathcal{L}'' . Consider the function

$$f(z) = \begin{cases} f_1(z), & \text{if } z \in \mathcal{L}' \\ f_2(z), & \text{if } z \in \mathcal{L}'' \end{cases} \quad (2.1)$$

where f_1, f_2 are functions, analytic on \mathcal{L}' and \mathcal{L}'' , i.e. analytic in some neighborhoods of \mathcal{L}' and \mathcal{L}'' correspondingly, and satisfying

$$f_1^{(r)}(z_0) = f_2^{(r)}(z_0), \quad r = \overline{0, k}, \quad f_1^{(k+1)}(z_0) \neq f_2^{(k+1)}(z_0). \quad (2.2)$$

By U we will denote an open circular neighborhood of the point z_0 , where both f_1, f_2 are analytic.

Let $Z_0^1, Z_0^2 \in \tilde{\mathcal{L}}$ be the prime ends, s.t. $|Z_0^j| = z_0, j = 1, 2$. Set

$$\tau_0^j := \Phi(Z_0^j), \quad j = 1, 2.$$

Points $\tau_1^j, j = 1, 2$ we define by

$$\tau_1^j = \lambda \tau_0^j,$$

with $\lambda > 1$ such that

$$\Gamma^1, \Gamma^2 \subset U,$$

where

$$\Gamma^j = \Gamma_0^j := \{\zeta : 1 < |\Phi(\zeta)| < \lambda, \arg \Phi(\zeta) = \arg \tau_0^j\}, \quad j = 1, 2. \quad (2.3)$$

The arcs Γ^1, Γ^2 are rectifiable (see [4, Chap. 5]), thus, can be oriented in such a way that for all $z \in \mathcal{L} \setminus \{z_0\}$ function f can be represented, by the Cauchy formula, as

$$f(z) = h_1(z) + h_2(z),$$

where

$$h_1(z) = \frac{1}{2\pi i} \int_{\Gamma^1 \cup \Gamma^2} \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} d\zeta, \quad (2.4)$$

and $h_2(z)$ is analytic for all $z \in \mathcal{L}$, therefore it can be approximated with a geometric rate on \mathcal{L} .

We will make use of the following lemma.

Lemma 1. *Let \mathcal{L} be a quasi-smooth arc. Then for any fixed non-negative integer k , a positive integer n and $\zeta \in \Gamma^1 \cup \Gamma^2$ there exists a polynomial kernel $K_n(\zeta, z)$ of the form $K_n(\zeta, z) = \sum_{j=0}^n a_j(\zeta) z^j$ with continuous in ζ coefficients $a_j(\zeta), j = \overline{0, n}$, satisfying for $z \in \mathcal{L}$ and ζ with $|\zeta - z_0| \geq \rho_{1/n}^*(z_0)$*

$$\left| \frac{1}{\zeta - z} - K_n(\zeta, z) \right| \leq c[\rho_{1/n}^*(z_0)]^{k+2} |\zeta - z_0|^{-(k+3)}, \quad (2.5)$$

where $c = c(\mathcal{L}) > 0$.

Proof. To show (2.5), we repeat word by word the proof for $k = 0$, ([4, Lemma 5.4]).

Let n be sufficiently large. For fixed m and r we consider the Dzyadyk polynomial kernel $K_{0,m,r,n}(\zeta, z)$ (see, e.g., [4, Chap. 3]). Then, for $r \geq 5$ and $z \in \mathcal{L}$, $\zeta \in \Gamma^j$, $j = 1, 2$,

$$\left| \frac{1}{\zeta - z} - K_{0,m,r,n}(\zeta, z) \right| \preccurlyeq \frac{1}{|\zeta - z|} \left| \frac{\tilde{\zeta} - \zeta}{\tilde{\zeta} - z} \right|^{rm}$$

where $\tilde{\zeta} := \tilde{\zeta}_{1/n}^j := \Psi[(1 + 1/n)\Phi(\zeta)]$.

Since

$$\left| \frac{\tilde{\zeta} - \zeta}{\tilde{\zeta} - z} \right| \preccurlyeq \left| \frac{\tilde{\zeta} - \zeta}{\tilde{\zeta} - z_0} \right| \preccurlyeq \left| \frac{\rho_{1/n}^j(z_0)}{\zeta - z_0} \right|^c \preccurlyeq \left| \frac{\rho_{1/n}^*(z_0)}{\zeta - z_0} \right|^c,$$

it is enough to take r and m such that $rmc \geq k + 2$, and set $K_n(\zeta, z) := K_{0,m,r,[\varepsilon n]}(\zeta, z)$, where $\varepsilon = \varepsilon(r, m) > 0$ is sufficiently small. \square

The next theorem is also a generalization of the case $k = 0$ in (2.2) and the proof essentially repeats the proof of [4, Theorem 5.2].

Theorem 3. *Let \mathcal{L} be a quasi-smooth arc, and let function f be given by (2.1), (2.2). Then*

$$c' [\rho_{1/n}^*(z_0)]^{k+1} \leq E_n(f, \mathcal{L}) \leq c'' [\rho_{1/n}^*(z_0)]^{k+1}, \quad (2.6)$$

where c', c'' don't depend on n .

Proof. First, we estimate $E_n(f, \mathcal{L})$ from above.

Without loss of generality, we can assume $z_0 = 0$ and n is sufficiently large. Let $d_n := \rho_{1/n}^*(0)$, $\gamma = \gamma_n := \{\zeta : \zeta \in \Gamma^1 \cup \Gamma^2, |\zeta| \geq d_n\}$,

$$P_n = \frac{1}{2\pi i} \int_{\gamma} (f_1(\zeta) - f_2(\zeta)) K_n(\zeta, z) d\zeta.$$

From (2.2), for all ζ in some neighborhood U of the point $z_0 = 0$

$$f_1(\zeta) = c_0 + c_1\zeta + \dots + c_k\zeta^k + c_{k+1}\zeta^{k+1} + \varphi_1(\zeta)\zeta^{k+2} \quad (2.7)$$

$$f_2(\zeta) = c_0 + c_1\zeta + \dots + c_k\zeta^k + \tilde{c}_{k+1}\zeta^{k+1} + \varphi_2(\zeta)\zeta^{k+2}, \quad (2.8)$$

where $c_{k+1} \neq \tilde{c}_{k+1}$ and $\varphi_1(\zeta), \varphi_2(\zeta)$ are functions, analytic in U .

Hence, there exists a constant C such that

$$|f_1(\zeta) - f_2(\zeta)| \leq C|\zeta^{k+1}|, \quad \zeta \in U. \quad (2.9)$$

By (2.5), (2.9), for all $z \in \mathcal{L}$

$$\begin{aligned} & \left| \frac{1}{2\pi i} \int_{\Gamma^1 \cup \Gamma^2} \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} d\zeta - P_n(z) \right| \\ & \leq \frac{1}{2\pi} \int_{\gamma} |f_1(\zeta) - f_2(\zeta)| \left| \frac{1}{\zeta - z} - K_n(z, \zeta) \right| |d\zeta| + \frac{1}{2\pi} \int_{(\Gamma^1 \cup \Gamma^2) \setminus \gamma} \left| \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} \right| |d\zeta| \end{aligned}$$

$$\leq \frac{C d_n^{k+2}}{2\pi} \int_{\gamma} \frac{|d\zeta|}{|\zeta|^2} + \frac{C}{2\pi} \int_{(\Gamma^1 \cup \Gamma^2) \setminus \gamma} \frac{|\zeta^{k+1}|}{|\zeta - z|} |d\zeta|. \quad (2.10)$$

Integration by parts of $\int_{\gamma} \frac{|d\zeta|}{|\zeta|^2}$ yields $\int_{\gamma} \frac{|d\zeta|}{|\zeta|^2} \asymp \frac{1}{d_n}$. Since $\text{dist}(\zeta, \mathcal{L}) \asymp |\zeta|$, (see [4, Chap. 5]), and $|(\Gamma^1 \cup \Gamma^2) \setminus \gamma| \asymp d_n$, it implies $\int_{(\Gamma^1 \cup \Gamma^2) \setminus \gamma} \frac{|\zeta^{k+1}|}{|\zeta - z|} |d\zeta| \asymp d_n^{k+1}$. Thus, combining with (2.10), we obtain the estimate from above in (2.6).

Now, we estimate $E_n(f, \mathcal{L})$ from below.

Let p_n^* be the polynomial of the best approximation, that is

$$|f(z) - p_n^*(z)| \leq E_n(f), \quad z \in \mathcal{L} \quad (2.11)$$

Without loss of generality we can assume that

$$E_n(f) \leq d_n = \rho_{1/n}^1(0).$$

Denote by $l_3 \subset \Omega^1$ any arc of a circle $\{\zeta : |\zeta| = d_n\}$, separating the prime end Z_0^1 from ∞ .

Let $z' \in \mathcal{L}'$ and $z'' \in \mathcal{L}''$ be the endpoints of the arc l_3 . Denote

$$l_1 := \mathcal{L}(0, z'), \quad l_2 := \mathcal{L}(0, z'').$$

Next, take a point z so that $z \in \Gamma^2$, $|z| = \varepsilon d_n$ (we'll choose the constant ε later). With a corresponding choice of orientation of arcs l_j , $j = 1, 2, 3$

$$I := \int_{l_1 \cup l_2} \frac{\tilde{f}(\zeta)}{(\zeta - z)^{k+2}} d\zeta = \int_{l_1 \cup l_2} \frac{\tilde{f}(\zeta) - \tilde{p}_n^*(\zeta)}{(\zeta - z)^{k+2}} d\zeta + \int_{l_3} \frac{\tilde{p}_n^*(\zeta)}{(\zeta - z)^{k+2}} d\zeta, \quad (2.12)$$

where $\tilde{f}(\zeta) = f(\zeta) - (c_0 + c_1\zeta + \dots + c_k\zeta^k)$ and $\tilde{p}_n^*(\zeta) = p_n^* - (c_0 + c_1\zeta + \dots + c_k\zeta^k)$.

Notice that $f(\zeta) - p_n^*(\zeta) = \tilde{f}(\zeta) - \tilde{p}_n^*(\zeta)$.

In the following estimates we use notations $a_i, \tilde{a}_i, \tilde{C}, \hat{C}, C_i$ for constants.

For the left hand side we have

$$\begin{aligned} & \left| \int_{l_1 \cup l_2} \frac{\tilde{f}(\zeta)}{(\zeta - z)^{k+2}} d\zeta \right| \\ &= \left| c_{k+1} \int_{l_1} \frac{\zeta^{k+1}}{(\zeta - z)^{k+2}} d\zeta + \tilde{c}_{k+1} \int_{l_2} \frac{\zeta^{k+1}}{(\zeta - z)^{k+2}} d\zeta \right. \\ & \quad \left. + \int_{l_1} \frac{\varphi_1(\zeta) \zeta^{k+2}}{(\zeta - z)^{k+2}} d\zeta + \int_{l_2} \frac{\varphi_2(\zeta) \zeta^{k+2}}{(\zeta - z)^{k+2}} d\zeta \right| \\ &= \left| c_{k+1} \log \frac{z}{z - z'} + \tilde{c}_{k+1} \log \frac{z - z''}{z} + \frac{a_1 z'^{k+1} + a_2 z''^k z + \dots + a_{k+1} z' z^k}{(z' - z)^{k+1}} \right. \\ & \quad \left. + \frac{\tilde{a}_1 z''^{k+1} + \tilde{a}_2 z''^k z + \dots + \tilde{a}_{k+1} z'' z^k}{(z'' - z)^{k+1}} + \tilde{C} + \int_{l_1} \frac{\varphi_1(\zeta) \zeta^{k+2}}{(\zeta - z)^{k+2}} d\zeta + \int_{l_2} \frac{\varphi_2(\zeta) \zeta^{k+2}}{(\zeta - z)^{k+2}} d\zeta \right| \\ & \geq \left| (\tilde{c}_{k+1} - c_{k+1}) \log \frac{z - z''}{z} + c_{k+1} \log \frac{z - z''}{z - z'} \right| - \frac{C_1 \varepsilon}{(1 - \varepsilon)^{k+1}} - \hat{C} \end{aligned}$$

$$\geq |\tilde{c}_{k+1} - c_{k+1}| \log \frac{1-\varepsilon}{\varepsilon} - \frac{C_1 \varepsilon}{(1-\varepsilon)^{k+1}} - C_2$$

Next, we estimate the right hand side of (2.12). By (2.11) and by the choice of z

$$\left| \int_{l_1 \cup l_2} \frac{\tilde{f}(\zeta) - \tilde{p}_n^*(\zeta)}{(\zeta - z)^{k+2}} d\zeta \right| \leq C_3 \frac{E_n}{\varepsilon^{k+1} d_n^{k+1}}.$$

To estimate the integral over l_3 notice that by (2.7) and (2.8)

$$|\tilde{f}(\zeta)| \leq c |\zeta|^{k+1}, \quad \zeta \in \mathcal{L}$$

for some constant c . Without loss of generality, we assume $c = 1$ (otherwise the arc l_3 must be considered with a radius $\frac{d_n}{c}$ instead). Since the estimate

$$|\tilde{p}_n^*(\zeta)| \leq |\tilde{p}_n^*(\zeta) - \tilde{f}(\zeta)| + |\tilde{f}(\zeta)| \leq d_n^{k+1} \left(1 + \left| \frac{\zeta}{d_n} \right|^{k+1} \right), \quad \zeta \in \mathcal{L}$$

holds, [4, Theorem 6.1] implies

$$|\tilde{p}_n^*(\zeta)| \leq C_4 d_n^{k+1}, \quad \zeta \in l_3.$$

The last inequality yields

$$\left| \int_{l_3} \frac{\tilde{p}_n^*(\zeta)}{(\zeta - z)^{k+2}} d\zeta \right| \leq \frac{2\pi C_4}{(1-\varepsilon)^{k+2}}.$$

Combining the estimates above, for some small but fixed ε we get

$$\begin{aligned} C_3 \frac{E_n}{\varepsilon^{k+1} d_n^{k+1}} &\geq |\tilde{c}_{k+1} - c_{k+1}| \log \frac{1-\varepsilon}{\varepsilon} - \frac{C_1 \varepsilon}{(1-\varepsilon)^{k+1}} - C_2 - \frac{2\pi C_4}{(1-\varepsilon)^{k+2}} \\ &\geq \frac{|\tilde{c}_{k+1} - c_{k+1}|}{2} \log \frac{1-\varepsilon}{\varepsilon}. \end{aligned}$$

Consequently, the estimate from below in (2.6) holds. \square

With reasoning completely similar, we obtain the following.

Theorem 4. *Let \mathcal{L} be a quasi-smooth arc, and let function f be given by (1.3), (1.4). Then*

$$c' [\rho_{1/n}^*(z_0)]^{k+1} \leq E_n(f, \mathcal{L}) \leq c'' [\rho_{1/n}^*(z_0)]^{k+1},$$

where $k := \min_{i=2, m-1} \{k_i\}$ and c', c'' don't depend on n .

3 Proof of Theorem 1

As it was mentioned above, f can be represented as

$$f(z) = \sum_{j=2}^{m-1} (h_1^j(z) + h_2^j(z)),$$

where $h_2^j(z)$ are analytic functions on \mathcal{L} and

$$h_1^j(z) = \frac{1}{2\pi i} \int_{\Gamma_j^1 \cup \Gamma_j^2} \frac{f_{j-1}(\zeta) - f_j(\zeta)}{\zeta - z} d\zeta,$$

with Γ_j^1, Γ_j^2 being the arcs given by (2.3), that correspond to the point z_j . Therefore, it's enough to construct polynomial approximants for $h_1^j(z)$ only.

To approximate the integral over Γ_j^i , $i = 1, 2$, consider a function $F_j^i : \mathcal{L} \cup \Gamma_j^i \rightarrow \mathcal{L}_{\varphi, j}^i$, such that F_j^i is one-to-one and satisfies

$$|F_j^i(z) - F_j^i(\zeta)| \leq c|z - \zeta|, \quad z, \zeta \in \mathcal{L} \cup \Gamma_j^i,$$

$$F_j^i(z_j) = 0,$$

$$F_j^i(\mathcal{L}(z_1, z_j)) = \mathcal{L}',$$

$$F_j^i(\mathcal{L}(z_j, z_m)) = \mathcal{L}'',$$

$$F_j^i(\Gamma_j^i) = \tilde{\Gamma},$$

where z_1, z_m are endpoints of \mathcal{L} , \mathcal{L}' is a line segment in $[0, \infty)$, \mathcal{L}'' is a line segment in the upper half plane that form an angle $\varphi > 0$ with \mathcal{L}' , (this angle will be determined below), and $\tilde{\Gamma}$ – a line segment at an angle $\frac{\varphi}{2}$ to the \mathcal{L}' .

Such a mapping $F_j^i \in Lip_1[\mathcal{L} \cup \Gamma_j^i]$ always exists, and to see this it is enough to note that \mathcal{L} and Γ_j^i are quasi-smooth and

$$dist(\zeta, \mathcal{L}) \asymp |\zeta - z_j|$$


holds for all $\zeta \in \Gamma_j^i$ (see [4, Chap. 5]).

By [2, Theorem 4] the function F_j^i can be approximated by polynomials $Q_n(z) := Q_{n,j}^i(z)$ with the rate $\frac{1}{n^\alpha}$, for some $\alpha > 0$, that is

$$|F_j^i(z) - Q_n(z)| \leq \frac{C}{n^\alpha}, \quad z \in \mathcal{L} \cup \Gamma_j^i, \quad (3.1)$$

where constant C does not depend on z and n .

For fixed $0 < \sigma < 1$ take an integer $k \geq 2$, such that $1 - \sigma > \frac{1}{1+k\alpha}$. Now, for $\varphi = \frac{2\pi}{k}$ consider corresponding mapping F_j^i and approximating polynomials Q_n .



Let

$$\widehat{P}_{n,j}^i(z, \zeta) = \frac{1 - \left(\frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right)^{\left[\frac{n^{1-\beta}}{2k} \right]}}{\zeta - z} + \left(\frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right)^{\left[\frac{n^{1-\beta}}{2k} \right]} K_{[\frac{n}{2}]}(z, \zeta).$$

It is not hard to see that $\widehat{P}_{n,j}^i(z, \zeta)$ is a polynomial in z of degree at most n . The idea of constructing such a polynomial is motivated by [5, p. 380].

We will show that for some choice of β and ζ_0 the term $\left| \frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right|^{\left[\frac{n^{1-\beta}}{2k} \right]}$ is bounded uniformly on \mathcal{L} by a constant that does not depend on n , and at points of analyticity of f it can be bounded by $q^{\left[\frac{n^{1-\beta}}{2k} \right]}$, for some $q < 1$.

For n sufficiently large the arc \mathcal{L} can be written as a disjoint union

$$\mathcal{L} = A_1 \cup A_2 \cup A_3,$$

where

$$A_1 := \left\{ z \in \mathcal{L} : |Q_n(z)| < \frac{C}{n^\alpha \sin \frac{\pi}{4k}} \right\}, \quad (3.2)$$

$$A_2 := \left\{ z \in \mathcal{L} : |Q_n(z)| \geq \frac{C}{n^\alpha \sin \frac{\pi}{4k}}, \text{dist}(Q_n(z), \mathcal{L}') \leq \frac{C}{n^\alpha} \right\}, \quad (3.3)$$

$$A_3 := \left\{ z \in \mathcal{L} : |Q_n(z)| \geq \frac{C}{n^\alpha \sin \frac{\pi}{4k}}, \text{dist}(Q_n(z), \mathcal{L}'') \leq \frac{C}{n^\alpha} \right\}, \quad (3.4)$$

where C is the constant from (3.1).
Points of A_1 satisfy

$$|Q_n^k(z)| < \frac{C^k}{n^{\alpha k} (\sin \frac{\pi}{4k})^k}. \quad (3.5)$$

For A_2 we have

$$|\sin(\arg Q_n(z))| \leq \frac{C}{n^\alpha |Q_n(z)|} \leq \sin \frac{\pi}{4k},$$

that implies

$$-\frac{\pi}{4} \leq \arg Q_n^k(z) \leq \frac{\pi}{4}. \quad (3.6)$$

Similarly, for A_3

$$|\sin(\frac{2\pi}{k} - \arg Q_n(z))| \leq \frac{C}{n^\alpha |Q_n(z)|} \leq \sin \frac{\pi}{4k},$$

that yields

$$-\frac{\pi}{4} \leq \arg Q_n^k(z) \leq \frac{\pi}{4}. \quad (3.7)$$

Γ_j^i can also be written as a disjoint union

$$\Gamma_j^i = B_1 \cup B_2,$$

where

$$B_1 := \{\zeta \in \Gamma_j^i : |Q_n(\zeta)| < \frac{C}{n^\alpha \sin \frac{\pi}{4k}}\}, \quad (3.8)$$

$$B_2 := \{\zeta \in \Gamma_j^i : |Q_n(\zeta)| \geq \frac{C}{n^\alpha \sin \frac{\pi}{4k}}\}. \quad (3.9)$$

Points of B_1 satisfy

$$|Q_n^k(\zeta)| < \frac{C^k}{n^{\alpha k} (\sin \frac{\pi}{4k})^k}. \quad (3.10)$$

For B_2 we have

$$\begin{aligned} |\sin(\frac{\pi}{k} - \arg Q_n(\zeta))| &\leq \frac{C}{n^\alpha |Q_n(\zeta)|} \leq \sin \frac{\pi}{4k}, \\ \pi - \frac{\pi}{4} &\leq \arg Q_n^k(\zeta) \leq \pi + \frac{\pi}{4}. \end{aligned} \quad (3.11)$$

Now, if we choose ζ_0 to be a point in $(0, \infty)$ with $\zeta_0 > \max\{|\mathcal{L}'|^k, |\mathcal{L}''|^k\}$, then (3.5), (3.6) and (3.7) imply $|Q_n^k(z) - \zeta_0| \leq \zeta_0 + \frac{C^k}{n^{\alpha k} (\sin \frac{\pi}{4k})^k}$, $z \in \mathcal{L}$. Also, by (3.10) and (3.11) the estimate $|Q_n^k(\zeta) - \zeta_0| \geq \zeta_0 - \frac{C^k}{n^{\alpha k} (\sin \frac{\pi}{4k})^k}$ holds for $\zeta \in \Gamma_j^i$.

According to these observations, we have

$$\left| \frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right|^{\left[\frac{n^{1-\beta}}{2k} \right]} \leq \left(1 + \frac{\tilde{C}}{n^{\alpha\beta k}} \right)^{\left[\frac{n^{1-\beta}}{2k} \right]} \quad (3.12)$$

where $\tilde{C} = \frac{2C^k}{\zeta_0 n^{\alpha\beta k} (\sin \frac{\pi}{4k})^k - C^k} \leq C^k$ for n large enough.

Let β be such that $1 - \sigma > \beta > \frac{1}{1+k\alpha}$, so that $1 - \beta < \alpha\beta k$ and $\sigma < 1 - \beta$. From (3.12) it follows

$$\left| \frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right|^{\left[\frac{n^{1-\beta}}{2k} \right]} \leq e^{\tilde{C}n^{-\alpha\beta k} n^{1-\beta}} \leq \hat{C}, \quad (3.13)$$

where \hat{C} does not depend on n .

Also, for all points z of a compact set $E \subset \mathcal{L} \setminus \{z_1, z_2, \dots, z_m\}$ and n sufficiently large the estimate

$$\left| \frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right|^{\left[\frac{n^{1-\beta}}{2k} \right]} \leq q^{\left[\frac{n^{1-\beta}}{2k} \right]}, \quad (3.14)$$

holds with some $q = q(E) < 1$.

Therefore, if we denote

$$d_n := \rho_{1/n}^*(z_j), \quad \gamma = \gamma_n := \{\zeta : \zeta \in \Gamma_j^i, |\zeta - z_j| \geq d_n\}$$

and consider polynomial

$$\begin{aligned} P_{n,j}^i(z) &= \frac{1}{2\pi i} \int_{\gamma} (f_{j-1}(\zeta) - f_j(\zeta)) \hat{P}_{n,j}^i(z, \zeta) d\zeta \\ &+ \frac{1}{2\pi i} \int_{\Gamma_j^i \setminus \gamma} (f_{j-1}(\zeta) - f_j(\zeta)) \frac{\left(1 - \left(\frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right)^{\left[\frac{n^{1-\beta}}{2k} \right]} \right)}{\zeta - z} d\zeta, \end{aligned}$$

by (3.13) and Theorem 4, for all $z \in \mathcal{L}$ we get

$$\begin{aligned} &\left| \frac{1}{2\pi i} \int_{\Gamma_j^i} \frac{f_{j-1}(\zeta) - f_j(\zeta)}{\zeta - z} d\zeta - P_{n,j}^i(z) \right| \\ &\leq \frac{1}{2\pi} \int_{\gamma} |f_{j-1}(\zeta) - f_j(\zeta)| \left| \frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right|^{\left[\frac{n^{1-\beta}}{2k} \right]} \left| \frac{1}{\zeta - z} - K_{[\frac{n}{2}]}(z, \zeta) \right| |d\zeta| \\ &+ \frac{1}{2\pi} \int_{\Gamma_j^i \setminus \gamma} \left| \frac{f_{j-1}(\zeta) - f_j(\zeta)}{\zeta - z} \right| \left| \frac{Q_{[n\beta]}^k(z) - \zeta_0}{Q_{[n\beta]}^k(\zeta) - \zeta_0} \right|^{\left[\frac{n^{1-\beta}}{2k} \right]} |d\zeta| \\ &\preceq d_n^{k_j+2} \int_{\gamma} \frac{|d\zeta|}{|\zeta|^2} + \int_{\Gamma_j^i \setminus \gamma} \left| \frac{\zeta^{k_j+1}}{\zeta - z} \right| |d\zeta| \preceq E_n(f, \mathcal{L}), \quad (3.15) \end{aligned}$$

where the last inequality follows by the reasoning, similar to the one we use in (2.10).

If $z \in E$, by (3.14), (2.10) and Theorem 4 we have

$$\left| \frac{1}{2\pi i} \int_{\Gamma_j^i} \frac{f_{j-1}(\zeta) - f_j(\zeta)}{\zeta - z} d\zeta - P_{n,j}^i(z) \right|$$

$$\begin{aligned}
&\leq \frac{q^{\lfloor \frac{n^{1-\beta}}{2k} \rfloor}}{2\pi} \int_{\gamma} |f_{j-1}(\zeta) - f_j(\zeta)| \left| \frac{1}{\zeta - z} - K_{\lfloor \frac{n}{2} \rfloor}(z, \zeta) \right| |d\zeta| \\
&\quad + \frac{q^{\lfloor \frac{n^{1-\beta}}{2k} \rfloor}}{2\pi} \int_{\Gamma_j^i \setminus \gamma} \left| \frac{f_{j-1}(\zeta) - f_j(\zeta)}{\zeta - z} \right| |d\zeta| \\
&\preceq E_n(f, \mathcal{L}) q^{\lfloor \frac{n^{1-\beta}}{2k} \rfloor} \preceq E_n(f, \mathcal{L}) e^{-\tilde{c}n^{1-\beta}}. \tag{3.16}
\end{aligned}$$

Let $P_n(z) = \sum_{j=2}^{m-1} (P_{n,j}^1(z) + P_{n,j}^2(z))$.

By (3.15), (3.16), polynomials $\{P_n\}$ are "near best" polynomials, approximating $\sum_{j=2}^{m-1} h_1^j(z)$ and satisfying (1.5). \square

4 Proof of Theorem 2

Since changing the R corresponds to scaling the lemniscate, we can always scale the picture and without loss of generality assume for simplicity $R = 1$.

As it was shown above, it's enough to approximate the function

$$h_1(z) = \frac{1}{2\pi i} \int_{\Gamma^1 \cup \Gamma^2} \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} d\zeta.$$

Here Γ^1 and Γ^2 we choose in such a way that $|P(\zeta)| > 1$ for all $\zeta \in (\Gamma^1 \cup \Gamma^2) \setminus \{0\}$. While the image of $\Gamma^1 \cup \Gamma^2$ under the mapping P belongs to the complement of the unit disc, the image of \mathcal{L} is inside the disc, that yields

$$\left| \frac{P(z)}{P(\zeta)} \right| \leq 1, \quad z \in \mathcal{L}, \quad \zeta \in \Gamma^1 \cup \Gamma^2 \tag{4.1}$$

Moreover, due to geometry of \mathcal{L} the equality in (4.1) occurs only if $\zeta = z = 0$.

Let

$$\widehat{P}_n(z, \zeta) = \frac{1 - \left(\frac{P(z)}{P(\zeta)}\right)^{\lfloor \frac{n}{2N} \rfloor}}{\zeta - z} + \left(\frac{P(z)}{P(\zeta)}\right)^{\lfloor \frac{n}{2N} \rfloor} K_{\lfloor \frac{n}{2} \rfloor}(z, \zeta). \tag{4.2}$$

One may check that $\widehat{P}_n(z, \zeta)$ is a polynomial in z of degree at most n .

Let $d_n := \rho_{1/n}^*(0)$, $\gamma = \gamma_n := \{\zeta : \zeta \in \Gamma^1 \cup \Gamma^2, |\zeta| \geq d_n\}$, and consider

$$\begin{aligned}
P_n(z) &= \frac{1}{2\pi i} \int_{\gamma} (f_1(\zeta) - f_2(\zeta)) \widehat{P}_n(z, \zeta) d\zeta \\
&\quad + \frac{1}{2\pi i} \int_{(\Gamma^1 \cup \Gamma^2) \setminus \gamma} (f_1(\zeta) - f_2(\zeta)) \left(\frac{1 - \left(\frac{P(z)}{P(\zeta)}\right)^{\lfloor \frac{n}{2N} \rfloor}}{\zeta - z} \right) d\zeta.
\end{aligned}$$

By virtue of Theorem 3, estimates (2.10) and (4.1), for all $z \in \mathcal{L}$

$$\left| \frac{1}{2\pi i} \int_{\Gamma^1 \cup \Gamma^2} \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} d\zeta - P_n(z) \right|$$

$$\begin{aligned}
&\leq \frac{1}{2\pi} \int_{\gamma} |f_1(\zeta) - f_2(\zeta)| \left| \frac{P(z)}{P(\zeta)} \right|^{\lfloor \frac{n}{2N} \rfloor} \left| \frac{1}{\zeta - z} - K_{\lfloor \frac{n}{2} \rfloor}(z, \zeta) \right| |d\zeta| \\
&\quad + \frac{1}{2\pi} \int_{(\Gamma^1 \cup \Gamma^2) \setminus \gamma} \left| \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} \right| \left| \frac{P(z)}{P(\zeta)} \right|^{\lfloor \frac{n}{2N} \rfloor} |d\zeta| \\
&\leq \frac{1}{2\pi} \int_{\gamma} |f_1(\zeta) - f_2(\zeta)| \left| \frac{1}{\zeta - z} - K_{\lfloor \frac{n}{2} \rfloor}(z, \zeta) \right| |d\zeta| + \frac{1}{2\pi} \int_{(\Gamma^1 \cup \Gamma^2) \setminus \gamma} \left| \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} \right| |d\zeta| \\
&\quad \preceq d_n^{k+2} \int_{\gamma} \frac{|d\zeta|}{|\zeta|^2} + \int_{(\Gamma^1 \cup \Gamma^2) \setminus \gamma} \left| \frac{\zeta^{k+1}}{\zeta - z} \right| |d\zeta| \\
&\quad \preceq [\rho_{1/n}^*(0)]^{k+1} \preceq E_n(f, \mathcal{L}).
\end{aligned}$$

If E is a compact set in $\mathcal{L} \setminus \{z_1, 0, z_2\}$, then for all $z \in E$

$$|P(z)| < q, \quad (4.3)$$

for some $q = q(E) < 1$.

Let

$$d(E) := \min_{z \in E} \{1 - |P(z)|\}.$$

By (4.3), $d(E) > 0$ for any compact set $E \subset \mathcal{L} \setminus \{z_1, 0, z_2\}$.

Therefore, for all $z \in E$

$$\left| \frac{P(z)}{P(\zeta)} \right|^{\lfloor \frac{n}{2N} \rfloor} \leq |P(z)|^{\lfloor \frac{n}{2N} \rfloor} \leq |1 - d(E)|^{\lfloor \frac{n}{2N} \rfloor} \leq e^{-cnd(E)},$$

where the constant $c > 0$ does not depend on n and E .

Hence, for $z \in E$

$$\begin{aligned}
&\left| \frac{1}{2\pi i} \int_{\Gamma^1 \cup \Gamma^2} \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} d\zeta - P_n(z) \right| \\
&\leq \frac{e^{-cnd(E)}}{2\pi} \int_{\gamma} |f_1(\zeta) - f_2(\zeta)| \left| \frac{1}{\zeta - z} - K_{\lfloor \frac{n}{2} \rfloor}(z, \zeta) \right| |d\zeta| \\
&\quad + \frac{e^{-cnd(E)}}{2\pi} \int_{(\Gamma^1 \cup \Gamma^2) \setminus \gamma} \left| \frac{f_1(\zeta) - f_2(\zeta)}{\zeta - z} \right| |d\zeta| \\
&\preceq [\rho_{1/n}^*(0)]^{k+1} e^{-cnd(E)} \preceq E_n(f, \mathcal{L}) e^{-cnd(E)}. \quad \square
\end{aligned}$$

Acknowledgment

The author would like to warmly thank Vladimir Andrievskii for guidance and many useful discussions.

References

- [1] G.A.Alibekov, V.K.Dzyadyk, *Uniform approximation of functions of a complex variable on closed sets with corners*, Mat. Sb. (N.S.), 75(117):4 (1968), 502–557; Math. USSR-Sb., 4:4 (1968), 463–517
- [2] V.V.Andrievskii, *Approximation characterization of classes of functions on continua of the complex plane*, Math. USSR-Sb., 53:1 (1986), 69–87.
- [3] V.V.Andrievskii, *Direct theorems of approximation theory on quasiconformal arcs*, Math. USSR-Izv., 16:2 (1981), 221–238
- [4] V.V.Andrievskii, V.I.Belyi, V.K.Dzjadyk, *Conformal Invariants in Constructive Theory of Functions of Complex Variable*, World Federation Publisher, Atlanta, GA, 1995.
- [5] V.V.Andrievskii, I.E.Pritsker, R.S.Varga, *Simultaneous approximation and interpolation of functions on continua in the complex plane*, J.Math.Pures Appl. 80, 4 (2001) 373-388.
- [6] H.-P.Blatt, E.B.Saff, *Behavior of zeros of polynomials of near best approximation*, J. Approx. Theory 46 No. 4 (1986). 323-344.
- [7] V.K. Dzyadyk, *On the theory of the approximation of functions on closed sets of the complex plane (apropos of a certain problem of S.M. Nikol'skii)*, Theory of functions and its applications, A collection of articles dedicated to Academician Sergei Mihailovic Nikol'skii on the occasion of his seventieth birthday, Trudy Mat. Inst. Steklov., 134, 1975, 63–114; Proc. Steklov Inst. Math., 134 (1977), 75–130
- [8] A.Kroo, E.B.Saff, *The density of extreme points in complex polynomial approximation*, Proc. Amer. Math. Soc., Vol.103, No.1 (May, 1988), pp.203-209.
- [9] V.V.Maimeskul, *Degree of approximation of analytic functions by nearly best polynomial approximants*, Constr. Approx. 11 (1995), 1–21.
- [10] E.B.Saff, *A principle of contamination in best polynomial approximation*, Approx. and Optim., Lecture Notes in Math., Vol. 1354, Springer-Verlag, Heidelberg, (1988), pp. 79-97.
- [11] E.B.Saff, V.Totik, *Behavior of Polynomials of Best Uniform Approximation*, Transactions of the American Mathematical Society, vol. 316, no. 2, 1989, pp. 567–593.
- [12] E.B.Saff, V.Totik, *Polynomial approximation of piecewise analytic functions*, J. London Math. Soc. (2) 39 (1989), no. 3, 487–498.
- [13] N.A.Shirokov, *On uniform approximation of functions on closed sets having a finite number of corner points with nonzero exterior angles*, Dokl. Akad. Nauk SSSR, 205:4 (1972), 798–800

- [14] N.A. Shirokov, V. Totik, *Polynomial Approximation on the Boundary and Strictly Inside*, *Constr. Approx.* (1994) 10: 145-152.