

Missing Mass of Rank-2 Markov Chains

Prafulla Chandra, Andrew Thangaraj
 Department of Elec Engg
 Indian Institute of Technology, Madras
 Chennai, India 600036
 {ee16d402, andrew}@ee.iitm.ac.in

Nived Rajaraman
 Department of Elec Engg and Comp Sci
 University of California, Berkeley
 Berkeley, CA 94720, USA
 nived.rajaraman@gmail.com

Abstract—Estimation of missing mass with the popular Good-Turing (GT) estimator is well-understood in the case where samples are independent and identically distributed (iid). In this article, we consider the same problem when the samples come from a stationary Markov chain with a rank-2 transition matrix, which is one of the simplest extensions of the iid case. We develop an upper bound on the absolute bias of the GT estimator in terms of the spectral gap of the chain and a tail bound on the occupancy of states. Borrowing tail bounds from known concentration results for Markov chains, we evaluate the bound using other parameters of the chain. The analysis, supported by simulations, suggests that, for rank-2 irreducible chains, the GT estimator has bias and mean-squared error falling with number of samples at a rate that depends loosely on the connectivity of the states in the chain.

I. INTRODUCTION

A. Preliminaries

Consider a Markov chain $X^n = (X_1, X_2, \dots, X_n)$ with the states $X_i \in \mathcal{X} = [K] \triangleq \{1, 2, \dots, K\}$ and

$$\Pr(X_i = x_i | X_{i-1} = x_{i-1}, \dots, X_1 = x_1) = \Pr(X_2 = x_i | X_1 = x_{i-1})$$

for $i = 2, \dots, n$ and all $x_i \in \mathcal{X}$. The transition probability matrix (t.p.m) of the Markov chain, denoted P , is the $K \times K$ matrix with (i, j) -th element $P_{ij} \triangleq \Pr(X_2 = j | X_1 = i)$. A distribution $\pi = [\pi_1, \dots, \pi_K]$ on $\mathcal{X} = [K]$ is said to be a stationary or invariant distribution of the Markov chain if $\pi P = \pi$ [1]. We will consider stationary Markov chains for which the distribution of X_1 (and all X_i) is a stationary distribution.

For $x \in \mathcal{X}$, let

$$F_x(X^n) \triangleq \sum_{i=1}^n I(X_i = x),$$

where $I(\cdot)$ is an indicator random variable, denote the number of occurrences of x in X^n . For a stationary Markov chain with state distribution π , the missing mass or the total probability of letters that did not appear in X^n , denoted $M_0(X^n, \pi)$, is defined as

$$M_0(X^n, \pi) \triangleq \sum_{x \in \mathcal{X}} \pi_x I(F_x(X^n) = 0).$$

For $l \geq 0$, let

$$\phi_l(X^n) \triangleq \sum_{x \in \mathcal{X}} I(F_x(X^n) = l)$$

denote the number of letters that have occurred l times in X^n . The popular and standard Good-Turing estimator [2] for $M_0(X^n, \pi)$ is defined as

$$G_0(X^n) \triangleq \frac{\phi_1(X^n)}{n}. \quad (1)$$

We will drop the arguments X^n, π whenever it is non-ambiguous.

II. PRIOR WORK AND PROBLEM SETTING

When the t.p.m has a rank of 1, the chain is *iid* and the bias of Good-Turing estimators falls as $1/n$ [3]. So, in the *iid* case, the number of letters occurring once (i.e. ϕ_1) acts as a good proxy for the missing letters in a sequence. However, the Markov case can be very different, in general. Consider a t.p.m with $P_{ii} = 1 - \eta$ and $P_{ij} = \eta/(K-1)$, $i \neq j$. For small η , every state is "sticky" and ϕ_1 appears to be inadequate to capture the mass of missing letters. As expected, a simulation of Good-Turing estimators for such sticky chains shows a non-vanishing bias.

Markov models occur naturally in language modeling [4] and several other applications in practice, where Good-Turing estimators and their modifications are routinely used. Therefore, it is interesting to analytically understand and study Good-Turing estimators for missing mass in Markov chains. Missing mass has been studied extensively in the *iid* case [3], [5]–[10], and estimation in Markov chains has been studied as well [11]–[14]. Recently, there has been interest in studying concentration and estimation of missing mass in Markov chains [15] [16].

The sticky chain example above has a full-rank t.p.m, while the *iid* case has a rank-1 t.p.m. This naturally motivates the study of Good-Turing estimators for chains with t.p.ms of other ranks. In particular, in this article, we consider chains with rank-2 t.p.ms, which are, in some sense, the simplest in the non-*iid* Markovian case.

Consider a Markov chain with a rank-2 t.p.m P , which we will call, loosely, as a rank-2 Markov chain. Since P has all entries in $[0, 1]$ with each row adding to 1 and it has rank 2, the eigenvalues of P will be $1, \lambda_2, 0, \dots, 0$, and $-1 \leq \lambda_2 \leq 1$ (by Perron-Frobenius theorem) [17]. The value of λ_2 determines several important properties of the chain. If $\lambda_2 = 1$, the chain is reducible. If $\lambda_2 = -1$, the chain is periodic with period 2. For $-1 < \lambda_2 < 1$, the chain is irreducible and aperiodic.

III. RESULTS

The main results of this article provide bounds on the absolute bias of Good-Turing estimators of missing mass $|E[G_0(X^n) - M_0(X^n, \pi)]|$ on rank-2 stationary Markov chains. To the best of our knowledge, these are perhaps the first such bounds to have appeared in the literature.

For the purpose of bounding, the bias is split into two significant components. The first component has contributions from states x for which π_x is low. This component is bounded in terms of the spectral gap

$$\beta(P) \triangleq 1 - \lambda_2 \in [0, 2]. \quad (2)$$

For bounding the remaining part of the bias, we use tail bounds on the occupancy $F_x(X^n)$ for states x with a high enough value of π_x .

The following theorem states the main starting point of our bounds on bias. We let $\bar{x} \triangleq 1 - x$.

Theorem 1. *Let X^n be a stationary Markov chain with state distribution π and a rank-2 t.p.m with spectral gap β . Let $1/n < \delta \leq \beta/5$. Then, there exist universal constants $c_1 > 0$ and $c_2 > 0$ such that*

$$|E[G_0(X^n) - M_0(X^n, \pi)]| \leq \frac{\delta}{\beta} \left(c_1 + \frac{c_2}{n\beta} \right) + 2 \max_{\substack{x: \pi_x > \delta, \\ P_{xx} \neq \pi_x}} \Pr(F_x(X^n) \leq 1) + O(1/n). \quad (3)$$

As $\beta \rightarrow 0$, the chain becomes reducible, and the bias bound grows unbounded as expected because consistent estimation is not possible for missing mass in reducible chains [15]. For example, consider the chain with t.p.m

$$\begin{bmatrix} (2/K)\mathbf{1}_{K/2 \times K/2} & \mathbf{0}_{K/2 \times K/2} \\ \mathbf{0}_{K/2 \times K/2} & (2/K)\mathbf{1}_{K/2 \times K/2} \end{bmatrix},$$

where $\mathbf{b}_{r \times c}$ denotes the $r \times c$ all- b matrix, with the uniform distribution on all states as π . This chain does not visit half the states making the estimation of missing mass inconsistent.

For a non-vanishing β , using Theorem 1 with a specific tail bound for $\Pr(F_x \leq 1)$ and a choice for δ , we obtain bounds that depend only on the chain's parameters. Tail bounds, obtained from concentration inequalities for Markov chains, typically use other parameters derived from the t.p.ms. The inequality of Kontorovich and Ramanan in [18] uses a parameter $\theta(P)$ defined as

$$\theta(P) \triangleq \sup_{x, x' \in \mathcal{X}} d_{\text{TV}}(P(\cdot|x), P(\cdot|x')), \quad (4)$$

where $d_{\text{TV}}(p, q) = \frac{1}{2} \sum_{x \in \mathcal{X}} |p_x - q_x|$ is the total variation (TV) distance between two distributions p and q on the same alphabet \mathcal{X} . We will call $\theta(P)$ or simply θ as the *maximum TV gap* of the chain.

Corollary 2. *For a rank-2 Markov chain with spectral gap β and maximum TV gap θ satisfying $\beta \geq \beta_0 \triangleq 5 \left(\frac{1}{\theta n^c} + \frac{1}{n} \right)$ for some $c \in (0, 0.5)$, we have*

$$|E[G_0(X^n) - M_0(X^n, \pi)]| \leq \frac{1}{\beta} \left(\frac{1}{n} + \frac{1}{\theta n^c} \right) \left(c_1 + \frac{c_2}{n\beta} \right) + 4e^{-0.5n^{(1-2c)}} + O(1/n). \quad (5)$$

In the above corollary, a tail bound from the concentration inequality of [18] is used for $\Pr(F_x \leq 1)$ in Theorem 1 with $\delta = \beta_0/5$. The bias bound is now expressed in terms of n , the spectral gap β and the maximum TV gap θ , and the rate of fall of absolute bias with n is bounded by how the two gap parameters β and θ vary with n .

If two rows of the t.p.m are disjoint in their non-zero positions, we get $\theta = 1$ and Corollary 2 does not apply. We will need to use other concentration inequalities in such cases.

The concentration inequality for Markov chains due to Naor, Rao and Regev in [19] uses a parameter $\lambda_\pi(P)$ defined as

$$\lambda_\pi(P) \triangleq \sup_{z: \sum_{i=1}^K \pi_i z_i^2 = 1} \sqrt{\sum_{i=1}^K \pi_i \left(\sum_{j=1}^K (P_{ij} - \pi_j) z_j \right)^2}, \quad (6)$$

which is the norm of $P - \mathbf{1}_{K \times 1} \pi$ in the Hilbert space $L_2(\pi)$ with $\|z\|_\pi^2 = \sum_i \pi_i z_i^2$. If P is orthonormally diagonalizable in $L_2(\pi)$, λ_π is equal to the absolute second eigenvalue $|\lambda_2|$ in the rank-2 case. However, in most other cases, the optimization in (6) needs to be evaluated to find λ_π , and the evaluation is usually feasible in the rank-2 case. We will call the parameter $\lambda_\pi(P)$ or simply λ_π as the *non-iid weighted norm* of the chain.

Corollary 3. *For a rank-2 Markov chain with spectral gap β and non-iid weighted norm λ_π satisfying $\beta \geq \beta_1 \triangleq 15 \sqrt{\frac{\ln n}{n \lambda_\pi}} + 5/n$, we have*

$$|E[G_0(X^n) - M_0(X^n, \pi)]| \leq \frac{1}{\beta} \left(c_1 + \frac{c_2}{n\beta} \right) \left(3 \sqrt{\frac{\ln n}{\lambda_\pi n}} + \frac{1}{n} \right) + O(1/n), \quad (7)$$

where $c_1, c_2 > 0$ are universal constants.

In the above corollary, a tail bound from the concentration inequality of [19] is used for $\Pr(F_x \leq 1)$ in Theorem 1 with $\delta = \beta_1/5$.

We will now consider a few specific types of rank-2 irreducible chains and use Corollaries 2 and 3 to bound the absolute bias of the Good-Turing estimator for missing mass in terms of β , θ and non-iid weighted norm λ_π .

Chains with $\theta = 0$ (IID): The iid case $P = \mathbf{1}_{K \times 1} \pi$ (i.e. each row equal to π) has $\beta = 1$ and $\theta = 0$ resulting in a $1/\sqrt{n}$ upper bound from Corollary 2, which, while capturing the fall of bias to zero, is poorer in rate than the well-known $1/n$ bound.

Non-iid chains: Consider the following $K \times K$ rank-2 t.p.ms.

$$P_1 = \begin{bmatrix} a\mathbf{1}_{\frac{K}{2} \times \frac{K-K_1}{2}} & a\mathbf{1}_{\frac{K}{2} \times \frac{K_1}{2}} & a\mathbf{1}_{\frac{K}{2} \times \frac{K_1}{2}} & \mathbf{0} \\ \mathbf{0} & a\mathbf{1}_{\frac{K}{2} \times \frac{K_1}{2}} & a\mathbf{1}_{\frac{K}{2} \times \frac{K_1}{2}} & a\mathbf{1}_{\frac{K}{2} \times \frac{K-K_1}{2}} \end{bmatrix},$$

$$P_2 = \begin{bmatrix} \frac{2}{K}\mathbf{1}_{\frac{K-K_1}{2} \times \frac{K}{2}} & \mathbf{0} \\ \mathbf{0} & \frac{2}{K}\mathbf{1}_{\frac{K_1}{2} \times \frac{K}{2}} \\ \frac{2}{K}\mathbf{1}_{\frac{K_1}{2} \times \frac{K}{2}} & \mathbf{0} \\ \mathbf{0} & \frac{2}{K}\mathbf{1}_{\frac{K-K_1}{2} \times \frac{K}{2}} \end{bmatrix},$$

$$P_3 = \begin{bmatrix} \frac{2}{K}\mathbf{1}_{\frac{K-K_1}{2} \times \frac{K}{2}} & \mathbf{0} \\ \frac{1}{K}\mathbf{1}_{K_1 \times \frac{K}{2}} & \frac{1}{K}\mathbf{1}_{K_1 \times \frac{K}{2}} \\ \mathbf{0} & \frac{2}{K}\mathbf{1}_{\frac{K-K_1}{2} \times \frac{K}{2}} \end{bmatrix},$$

where $a = 2/(K + K_1)$ and $\mathbf{0}$ is an all-zero matrix of appropriate size. In all of the three cases above, States 1 to $K/2$ and States $K/2 + 1$ to K are connected only through the K_1 states from $(K - K_1)/2 + 1$ to $(K + K_1)/2$. Motivated by the *iid* worst case, we let $K = n$ and consider $K_1 = \Theta(n^\kappa)$ for $\kappa \leq 1$ to control the connectivity in the chains. Table I lists the dominant terms in the parameters β , $\bar{\theta}$, $\bar{\lambda}_\pi$ and the bounds from Corollaries 2 and 3 for the t.p.ms P_1 , P_2 and P_3 (up to a multiplicative constant). The three chains considered

| $K = n$ $K_1 = n^\kappa$ | β | $\bar{\theta}$ | $\bar{\lambda}_\pi$ | Cor 2 $\kappa > 3/4$ | Cor 3 $\kappa > 2/3$ |
|-----------------------------|--------------|----------------|---------------------|-----------------------------|--------------------------------------|
| P_1 | n^κ/n | n^κ/n | n^κ/n | $\frac{1}{n^{2\kappa-3/2}}$ | $\sqrt{\frac{\ln n}{n^{3\kappa-2}}}$ |
| P_2 | n^κ/n | 0 | 0 | — | — |
| P_3 | n^κ/n | 0 | n^κ/n | — | $\sqrt{\frac{\ln n}{n^{3\kappa-2}}}$ |

TABLE I: Dominant terms of β , $\bar{\theta}$, $\bar{\lambda}_\pi$ and bounds (up to a multiplicative constant).

above illustrate different regimes of values for the parameters θ and λ . The corollaries apply when $\bar{\theta} \neq 0$ or $\bar{\lambda} \neq 0$ and when κ is large enough to satisfy the lower bound on the spectral gap β in the corollaries. The case of $\kappa = 1$ results in a bound tending to $1/\sqrt{n}$, and lower κ results in weaker bounds.

Fig. 1 shows simulation plots of the rate of fall of absolute bias and mean-squared error versus n for the three t.p.ms above with $\kappa = 1$ and $\kappa = 1/4$. We observe that both the absolute bias and the mean-squared error fall with n in all cases. For the highly connected case of $\kappa = 1$, the rate of fall in simulations is close to $1/n$. As κ decreases and connectivity reduces, the rate of fall of both absolute bias and MSE reduce. *Periodic case:* Consider the following $n \times n$ t.p.m

$$\begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 1 & 0 & 0 & \cdots & 0 \end{bmatrix}_{r \times r} \otimes \begin{bmatrix} r/n & \cdots & r/n \\ \vdots & \ddots & \vdots \\ r/n & \cdots & r/n \end{bmatrix}_{n/r \times n/r},$$




Fig. 1: Rate of fall of absolute bias ($|ME|$) and mean-squared error (MSE).

which is a Kronecker product of an $r \times r$ right-shift-by-1 permutation matrix and an $n/r \times n/r$ matrix with each entry equal to r/n . The rank of the above t.p.m is r , the chain is irreducible and periodic with all states having a period r , and the uniform distribution is the unique stationary distribution. For $r = 2$, we get the rank-2 irreducible, period-2 chain. Using a direct computation in a manner similar to the *iid* case, we obtain the following exact characterization of bias:

$$|E[G_0(X^n) - M_0(X^n, \pi)]| = \frac{r}{n} \left(1 - \frac{r}{n}\right)^{\frac{n}{r}-1}. \quad (8)$$

For sub-linear r , we see that bias tends to zero. However, for $r \propto n$, we get a bias that is constant.

In summary, we see that the bias of the Good-Turing estimator for missing mass in rank-2 Markov chains can be analytically bounded using the spectral gap (β in (2)) and other parameters such as the maximum TV gap (θ in (4)) or the non-iid weighted norm (λ_π in (6)) that control the concentration of state occupancies. If better tail bounds can be developed, the rate of fall of bias predicted from the bounds can be tightened further.

From examples such as the sticky chain and the periodic case above, we expect a constant bias for the Good-Turing estimator in certain higher rank Markov chains. Improvement in estimation methods for missing mass is needed for such higher rank chains.

IV. PROOF OF THEOREMS AND COROLLARIES

A. Theorem 1

We begin with the result of [15, Theorem 1], which states that for a Markov chain X^n with state distribution π ,

$$|E[G_0(X^n) - M_0(X^n, \pi)]| \leq \frac{1}{n} + \left| \sum_{x \in \mathcal{X}} \pi_x \bar{\pi}_x \Gamma_x \right|, \quad (9)$$

where

$$n\Gamma_x = \sum_{i=1}^n \Pr(F_x(X_{\sim i}^n)=0 | X_i=x) - \Pr(F_x(X_{\sim i}^n)=0 | X_i \neq x)$$

and $X_{\sim i}^n = (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n)$.

Lemma 4. For a rank-2 t.p.m P , we have the following:

- 1) For $x \in \mathcal{X}$ with $P_{xx} = \pi_x$, $\Gamma_x = 0$.
- 2) For δ satisfying $1/n < \delta \leq \beta/5$, there exist constants c_1 and c_2 such that

$$\sum_{\substack{x: \pi_x \leq \delta, \\ P_{xx} \neq \pi_x}} \pi_x \bar{\pi}_x |\Gamma_x| \leq \frac{\delta}{\beta} \left(c_1 + \frac{c_2}{n\beta} \right).$$

Proof. See Section V. \square

Using Part 1 of Lemma 4, it suffices to upper bound $\left| \sum_{x \in \mathcal{X}: P_{xx} \neq \pi_x} \pi_x \bar{\pi}_x \Gamma_x \right|$. In order to carry out this analysis, we show that π_x and Γ_x bear an inverse relationship. For this, we fix a threshold δ (to be decided later) and partition the set of states with $P_{xx} \neq \pi_x$ into two sets

$$\begin{aligned} A(\delta) &= \{x \in \mathcal{X} : \pi_x \leq \delta, P_{xx} \neq \pi_x\}, \\ \bar{A}(\delta) &= \{x \in \mathcal{X} : \pi_x > \delta, P_{xx} \neq \pi_x\}. \end{aligned}$$

Observe that

$$\begin{aligned} \pi_x \bar{\pi}_x \Gamma_x &= \frac{1}{n} \left(\sum_{i=1}^n \bar{\pi}_x \Pr(F_x(X_{\sim i}^n)=0, X_i=x) \right. \\ &\quad \left. - \pi_x \Pr(F_x(X_{\sim i}^n)=0, X_i \neq x) \right) \\ &= \frac{\bar{\pi}_x}{n} \left(\sum_{i=1}^n \Pr(F_x(X_{\sim i}^n)=0, X_i=x) \right) - \pi_x \Pr(F_x(X^n) = 0) \\ &= \frac{\bar{\pi}_x}{n} \Pr(F_x(X^n) = 1) - \pi_x \Pr(F_x(X^n) = 0). \end{aligned}$$

Summing over $x \in \bar{A}(\delta)$, we get

$$\begin{aligned} &\left| \sum_{x \in \bar{A}(\delta)} \pi_x \bar{\pi}_x \Gamma_x \right| \\ &= \left| \sum_{x \in \bar{A}(\delta)} \frac{\bar{\pi}_x}{n} \Pr(F_x(X^n) = 1) - \pi_x \Pr(F_x(X^n) = 0) \right| \\ &\leq \sum_{x \in \bar{A}(\delta)} \left(\frac{\bar{\pi}_x}{n} + \pi_x \right) \Pr(F_x(X^n) \leq 1) \\ &\leq 2 \max_{x \in \bar{A}(\delta)} \Pr(F_x(X^n) \leq 1), \end{aligned} \quad (10)$$

where, in the last inequality, we use $\bar{\pi}_x < 1$ and $|\bar{A}(\delta)| \leq 1/\delta < n$. Using Part 2 of Lemma 4 and (10), the proof is complete.

B. Corollaries 2 and 3

For any x , we have $E[F_x(X^n)/n] = \pi_x$. For large π_x , if $F_x(X^n)$ concentrates, we can bound the tail probability $\Pr(F_x(X^n) \leq 1)$ using concentration of $F_x(X^n)$ as follows:

$$\begin{aligned} \Pr(F_x(X^n) \leq 1) &= \Pr(\pi_x - F_x(X^n)/n \geq \pi_x - 1/n) \\ &\leq \Pr(|\pi_x - F_x(X^n)/n| \geq \pi_x - 1/n). \end{aligned} \quad (11)$$

Using the concentration results in [18] and [19], we obtain the following:

Lemma 5. 1) If the chain has maximum TV gap $\theta < 1$,

$$\Pr(|\pi_x - F_x(X^n)/n| \geq \epsilon) \leq 2e^{-0.5n\theta^2\epsilon^2}.$$

2) If the chain has non-iid weighted norm $\lambda_\pi < 1$,

$$\Pr(|\pi_x - F_x(X^n)/n| \geq \epsilon) \leq C \left(\frac{q}{\lambda_\pi n} \right)^{q/2} \pi_x \epsilon^{-q},$$

where C is a constant and $q \geq 2$.

Proof. See Section V. \square

1) *Proof of Corollary 2:* We consider the set $\bar{A}(\delta)$ with

$$\delta = \frac{1}{\theta n^c} + \frac{1}{n}, \quad c \in (0, 0.5). \quad (12)$$

For $x \in \bar{A}(\delta)$, we have $\pi_x - \frac{1}{n} > \frac{1}{\theta n^c}$. So, setting $\epsilon = \frac{1}{\theta n^c}$ in Part 1 of Lemma 5, we get

$$\Pr(F_x(X^n) \leq 1) \leq 2e^{-0.5n^{1-2c}}.$$

Using δ from (12) and the above bound for $\Pr(F_x(X^n) \leq 1)$ in Theorem 1, we get (5) and the proof for Corollary 2 is complete.

2) *Proof of Corollary 3:* We consider the set $\bar{A}(\delta)$ with

$$\delta = 3\sqrt{\frac{\ln n}{n\lambda_\pi}} + \frac{1}{n}. \quad (13)$$

For $x \in \bar{A}(\delta)$, we have $\pi_x - 1/n > 3\sqrt{\frac{\ln n}{n\lambda_\pi}}$. So, setting $\epsilon = 3\sqrt{\frac{\ln n}{n\lambda_\pi}}$ and $q = 3 \ln n$ in Part 2 of Lemma 5, we get

$$\max_{x \in \bar{A}(\delta)} \Pr(F_x(X^n) \leq 1) \leq C/n^{1.5}.$$

Using δ from (13) and the above bound for $\Pr(F_x(X^n) \leq 1)$ in Theorem 1, we get (7) and the proof for Corollary 3 is complete.

V. PROOFS OF LEMMAS

Consider a rank-2 $K \times K$ irreducible t.p.m P with stationary distribution $\pi = [\pi_1 \cdots \pi_K]^T$. There exist vectors $u = [u_1 \cdots u_K]^T$ and $v = [v_1 \cdots v_K]^T$ satisfying the following decompositions:

- 1) If P is diagonalizable with spectral gap $\beta \neq 1$, then

$$P = RDS \text{ with } SR = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

where $R = [\mathbf{1}_{K \times 1} \quad v]$, $D = \begin{bmatrix} 1 & 0 \\ 0 & \beta \end{bmatrix}$ and $S = \begin{bmatrix} \pi^T \\ u^T \end{bmatrix}$. Since P is a t.p.m, we have, for $1 \leq i, j \leq K$,

$$0 \leq \pi_j + (1 - \beta)v_i u_j \leq 1. \quad (14)$$

- 2) If P is non-diagonalizable with spectral gap $\beta = 1$, then

$$P = RS \text{ with } SR = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

Since P is a t.p.m, we have, for $1 \leq i, j \leq K$,

$$0 \leq \pi_j + v_i u_j \leq 1.$$

A. Lemma 4, Part 2 (see [20] for Part 1)

The quantity Γ_x , when $P_{xx} \neq \pi_x$, can be expressed in terms of some more spectral parameters for both the diagonalizable and the non-diagonalizable cases.

Lemma 6. *If P is diagonalizable and $x \in \mathcal{X}$ with $P_{xx} \neq \pi_x$,*

$$\begin{aligned} \Gamma_x &= \bar{\beta} v_x u_x \left[\left((\lambda_2^{\sim x})^{n-1} - (\lambda_1^{\sim x})^{n-1} \right) \frac{\Delta_x^{-1}}{\pi_x} \right. \\ &\quad - \left. \left((\lambda_1^{\sim x})^{n-1} + (\lambda_2^{\sim x})^{n-1} \right) \left(1 - \frac{2}{n} \right) \frac{\beta}{\Delta_x^2} \right. \\ &\quad \left. + \frac{2}{n} \frac{\beta}{\Delta_x^3} \lambda_1^{\sim x} \lambda_2^{\sim x} \left((\lambda_1^{\sim x})^{n-2} - (\lambda_2^{\sim x})^{n-2} \right) \right], \quad (15) \end{aligned}$$

where $\lambda_i^{\sim x} = 0.5(\bar{\pi}_x + \bar{\beta}(1 - v_x u_x) + (-1)^{i+1} \Delta_x)$, $i = 1, 2$, are the eigenvalues of the matrix $\left(I_{2 \times 2} - \begin{bmatrix} \pi_x \\ u_x \end{bmatrix} \begin{bmatrix} 1 & v_x \end{bmatrix} \right) D$,

and $\Delta_x^2 = \left[\beta - \pi_x + \bar{\beta} v_x u_x \right]^2 + 4\bar{\beta} \pi_x v_x u_x$.

Proof. See [20] for a proof. \square

Now, $|\Gamma_x|$ is upper-bounded by the sum of absolute value of all the terms in the RHS of (15). To simplify the bound, we use $|\lambda_i^{\sim x}| < 1$ and $1 - 2/n < 1$ to get

$$|\Gamma_x| \leq 2 \left(\frac{\Delta_x^{-1}}{\pi_x} + \frac{\beta}{\Delta_x^2} \left(1 + \frac{2\Delta_x^{-1}}{n} \right) \right) |\bar{\beta} v_x u_x|. \quad (16)$$

For $x \in A(\delta)$, we have $\pi_x \leq \delta$ and $\delta \leq \beta/5 \leq 2/5$. Using this and (14), it is possible to show that

Claim 1: $\Delta_x \geq \beta/3$ for $x \in A(\delta)$.

Using these bounds in (16), we get

$$|\Gamma_x| \leq \frac{2}{\beta} \left(7 + \frac{27}{n\beta} \right) |\bar{\beta} v_x u_x|, \quad x \in A(\delta). \quad (17)$$

Next we claim that

Claim 2: $\sum_{x \in A(\delta)} |\bar{\beta} v_x u_x| \leq 3$.

Using this and (17), we get

$$\sum_{x \in A(\delta)} \pi_x \bar{\pi}_x |\Gamma_x| \leq \frac{6\delta}{\beta} \left(7 + \frac{27}{n\beta} \right),$$

where we have bounded $\bar{\pi}_x \leq 1$ and $\pi_x < \delta$. This completes the proof for the diagonalizable case.

1) *Proofs of Claims:* **Claim 1:** Using $\pi_x \leq \delta \leq \beta/5$ for $x \in A(\delta)$ and $\bar{\beta} v_x u_x \geq -\pi_x$ from (14), we get

(i) $\beta - \pi_x + \bar{\beta} v_x u_x \geq \beta - 2\pi_x \geq 5\pi_x - 2\pi_x \geq 0$, and (ii) $4\pi_x \bar{\beta} v_x u_x \geq -4\pi_x^2$. This implies that,

$$\begin{aligned} \Delta_x^2 &= (\beta - \pi_x + \bar{\beta} v_x u_x)^2 + 4\pi_x \bar{\beta} v_x u_x \\ &\geq (\beta - 2\pi_x)^2 - 4\pi_x^2 \geq \beta^2/5 \geq \beta^2/9 \end{aligned} \quad (18)$$

and hence $\Delta_x \geq \beta/3$ for $x \in A(\delta)$.

Claim 2: To prove this, we partition $A(\delta)$ into two sets, $A_1(\delta) = \{x \in A(\delta) : \bar{\beta} v_x u_x \leq 0\}$ and its complement, $\bar{A}_1(\delta) = A(\delta) \setminus A_1(\delta)$.

- 1) For $x \in A_1(\delta)$, using $\bar{\beta} v_x u_x \leq 0$ and $\bar{\beta} v_x u_x \geq -\pi_x$ from (14), we get

$$0 \geq \sum_{x \in A_1(\delta)} \bar{\beta} v_x u_x \geq -\sum_{x \in A_1(\delta)} \pi_x \geq -1 \text{ which implies } \left| \sum_{x \in A_1(\delta)} \bar{\beta} v_x u_x \right| \leq 1.$$

- 2) On the other hand, to deal with the sum over $\bar{A}_1(\delta)$, note that $\sum_{x \in \mathcal{X}} v_x u_x = 1$ since the matrix product $SR = I$. Therefore,

$$\sum_{x \in \mathcal{X}: \bar{\beta} v_x u_x > 0} \bar{\beta} v_x u_x = \bar{\beta} - \sum_{x \in \mathcal{X}: \bar{\beta} v_x u_x \leq 0} \bar{\beta} v_x u_x$$

Since the summation on the right in the above equation is bounded above by 1 in absolute value (because of (14) again), we get $\left| \sum_{x \in \bar{A}_1(\delta)} \bar{\beta} v_x u_x \right| \leq \sum_{x \in \mathcal{X}: \bar{\beta} v_x u_x > 0} \bar{\beta} v_x u_x \leq 1 + \bar{\beta} \leq 2$.

Lemma 7. *If P is non-diagonalizable and $x \in \mathcal{X}$ with $P_{xx} \neq \pi_x$,*

$$\begin{aligned} \Gamma_x &= \frac{v_x u_x}{\Delta_x} \left[\left((\lambda_2^{\sim x})^{n-1} - (\lambda_1^{\sim x})^{n-1} \right) \frac{1}{\pi_x} \right. \\ &\quad - \left. \left((\lambda_1^{\sim x})^{n-1} + (\lambda_2^{\sim x})^{n-1} \right) \left(1 - \frac{2}{n} \right) \Delta_x^{-1} \right. \\ &\quad \left. + \frac{2}{n} \Delta_x^{-2} \lambda_1^{\sim x} \lambda_2^{\sim x} \left((\lambda_1^{\sim x})^{n-2} - (\lambda_2^{\sim x})^{n-2} \right) \right], \quad (19) \end{aligned}$$

where $\lambda_i^{\sim x} = 0.5(\bar{\pi}_x - v_x u_x) + (-1)^{i+1} \Delta_x$, $i = 1, 2$ are the eigenvalues of the matrix $\begin{bmatrix} \bar{\pi}_x & -\pi_x v_x \\ -u_x & -u_x v_x \end{bmatrix}$ with $\Delta_x^2 = (\bar{\pi}_x - v_x u_x)^2 + 4v_x u_x$.

Proof. Similar to that of Lemma 6. \square

Bounding $\sum_{x \in A} \pi_x \bar{\pi}_x |\Gamma_x|$ using a method similar to the diagonalizable case, completes the proof.

B. Lemma 5

Theorem 8 ([18]). Suppose that X^n is a Markov chain with state space \mathcal{X} and t.p.m P with $\theta(P) < 1$, and $\psi(X^n) : \mathcal{X}^n \rightarrow \mathbb{R}$ is a d -Lipschitz function with respect to the normalized Hamming metric, then

$$\Pr(|\psi(X^n) - E[\psi(X^n)]| \geq \epsilon) \leq e^{-n(1-\theta)^2 \epsilon^2 / 2d^2}. \quad (20)$$

Since $F_x(X^n)/n$ is 1-Lipschitz, Theorem 20 directly results in Part 1. For Part 2, we use the following theorem from [19].

Theorem 9 ([19]). Suppose that X^n is a stationary Markov chain with state space $\mathcal{X} = [K]$, stationary distribution π and t.p.m P such that $\lambda_\pi(P) < 1$. Then every $f : [K] \rightarrow \mathbb{R}$ satisfies the following inequality for every $n \in \mathbb{N}$ and $q \geq 2$,

$$\begin{aligned} & \left(E \left[\left| \frac{\sum_{i=1}^n f(X_i)}{n} - E[f(X_1)] \right|^q \right] \right)^{\frac{1}{q}} \\ & \lesssim \sqrt{\frac{q}{(1-\lambda_\pi)n}} \left(E[|f(X_1)|^q] \right)^{\frac{1}{q}}, \end{aligned} \quad (21)$$

where $D_1 \lesssim D_2$ implies there exists a universal constant $C > 0$ such that $D_1 \leq CD_2$.

Using Markov's inequality,

$$\begin{aligned} \Pr \left(\left| \pi_x - F_x(X^n)/n \right|^q \geq \epsilon^q \right) & \leq E \left[\left| \pi_x - F_x(X^n)/n \right|^q \right] \epsilon^{-q} \\ & \stackrel{(a)}{\leq} C \left(\frac{q}{\lambda_\pi n} \right)^{q/2} \pi_x \epsilon^{-q}, \end{aligned} \quad (22)$$

where (a) follows by setting $f(X_i) = I(X_i = x)$ in (21).

C. Proofs of Lemma 4 part 1, Lemma 6

We prove Lemma 6 and part 1 of Lemma 4 (for diagonalizable t.p.ms) using the Lemma below. (The proof of part 1 of Lemma 4 for non-diagonalizable t.p.ms follows a similar method and is hence omitted.)

Lemma 10. Let P be a rank-2 diagonalizable t.p.m. Let $E_{\sim m}^x$ denote the event $F_x(X_{\sim m}^n) = 0$.

- 1) For $x \in \mathcal{X}$ with $P_{xx} = \pi_x$,
 $\Pr(E_{\sim m}^x | X_m = x) = \Pr(E_{\sim m}^x | X_m \neq x) = (\bar{\pi}_x)^{n-1}$ for $m = 1$ to n .
- 2) For $x \in \mathcal{X}$ with $P_{xx} \neq \pi_x$,
 - a) for $m = 1$ to n ,

$$\begin{aligned} & \Pr(E_{\sim m}^x | X_m \neq x) \\ & = \frac{1}{2\bar{\pi}_x} \left[(\lambda_1^{\sim x})^n \left(\frac{s_x}{\Delta_x} + 1 \right) + (\lambda_2^{\sim x})^n \left(1 - \frac{s_x}{\Delta_x} \right) \right] \end{aligned} \quad (23)$$

b)

$$\begin{aligned} & P(E_{\sim 1}^x | X_1 = x) = P(E_{\sim n}^x | X_n = x) \\ & = (\lambda_1^{\sim x})^{n-1} \left(\frac{s_x + \Delta_x}{2\Delta_x} - \frac{(1-\beta)v_x u_x}{\Delta_x} \right) \\ & \quad + (\lambda_2^{\sim x})^{n-1} \left(\frac{\Delta_x - s_x}{2\Delta_x} + \frac{(1-\beta)v_x u_x}{\Delta_x} \right) \end{aligned} \quad (24)$$

c) for $m = 2$ to $n - 1$,

$$\begin{aligned} & \Pr(E_{\sim m}^x | X_m = x) \\ & = (\lambda_1^{\sim x})^{n-1} \left(\frac{s_x + \Delta_x}{2\Delta_x} - \frac{(1-\beta)v_x u_x}{\Delta_x} \right)^2 \\ & \quad + (\lambda_2^{\sim x})^{n-1} \left(\frac{\Delta_x - s_x}{2\Delta_x} + \frac{(1-\beta)v_x u_x}{\Delta_x} \right)^2 \\ & \quad + \frac{1}{\Delta_x^2} \beta(1-\beta)v_x u_x \left((\lambda_1^{\sim x})^{m-1} (\lambda_2^{\sim x})^{n-m} \right. \\ & \quad \left. + (\lambda_1^{\sim x})^{n-m} (\lambda_2^{\sim x})^{m-1} \right) \end{aligned} \quad (25)$$

where $s_x = 1 - \pi_x - (1-\beta)(1 - v_x u_x)$, $\lambda_1^{\sim x}$, $\lambda_2^{\sim x}$, and Δ_x are as defined in Lemma 6.

Recall that $n\Gamma_x = \sum_{m=1}^n \Pr(E_{\sim m}^x | X_m = x) - \Pr(E_{\sim m}^x | X_m \neq x)$.

Using part 1 of Lemma 10 in the above expression completes the proof of part 1 of Lemma 4. Using part 2 of Lemma 10 in the above expression for Γ_x and simplifying, we get (15). This completes the proof of Lemma 6.

D. Proof of Lemma 10

The probabilities $\Pr(E_{\sim m}^x | X_m = x)$ and $\Pr(E_{\sim m}^x | X_m \neq x)$ involve the events $E_{i,j}^x \triangleq (X_i \neq x, X_{i+1} \neq x, \dots, X_j \neq x)$ for suitable choices of i, j depending on m . To compute the probability of $E_{i,j}^x$ i.e. the chain X^n not visiting the state x consecutively for a certain number of steps, we use the matrix $P^{\sim x}$ which is obtained by replacing the column corresponding to state x in the t.p.m P with 0s. Since $P = RDS$, we have $P^{\sim x} = RDS^{\sim x}$, where $S^{\sim x}$ is obtained by replacing the column corresponding to state x in the matrix S with zeros. Since $SR = I$, we get $S^{\sim x}R = \begin{bmatrix} 1 - \pi_x & -\pi_x v_x \\ -u_x & 1 - u_x v_x \end{bmatrix}$.

We use the following Lemmas to prove Lemma 10. The first two Lemmas help us express $\Pr(E_{\sim m}^x | X_m \neq x)$ and $\Pr(E_{\sim m}^x | X_m = x)$ in terms of $S^{\sim x}$, R and D .

Lemma 11. For $m = 1, 2, \dots, n$

$$\begin{aligned} & \Pr(E_{\sim m}^x | X_m \neq x) \\ & = \left[1 \quad -\frac{(1-\beta)}{1-\pi_x} \pi_x v_x \right] (S^{\sim x} R D)^{n-2} \begin{bmatrix} 1 - \pi_x \\ -u_x \end{bmatrix} \end{aligned} \quad (26)$$

Proof. Appendix □

Lemma 12. 1) For $m \geq 2$,

$$\begin{aligned} & \Pr(E_{2:m}^x | X_1 = x) \\ & = \left[1 \quad (1-\beta)v_x \right] (S^{\sim x} R D)^{m-2} \begin{bmatrix} 1 - \pi_x \\ -u_x \end{bmatrix} \end{aligned} \quad (27)$$

2)

$$\Pr(E_{1:n-1}^x | X_n = x) = \Pr(E_{2:n}^x | X_1 = x) \quad (28)$$

For $m = 2, \dots, n-1$,

$$\Pr(E_{\sim m}^x | X_m = x) = \Pr(E_{2:m}^x | X_1 = x) \\ \Pr(E_{2:n-m+1}^x | X_1 = x) \quad (29)$$

Proof. Appendix \square

The next lemma gives an expression for $(S^{\sim x}RD)^l$ ($l \geq 1$) for various values of π_x, v_x and u_x .

Lemma 13. $(S^{\sim x}RD)^l$:

1) For $x \in \mathcal{X}$, with $v_x = u_x = 0$,

$$(S^{\sim x}RD)^l = \begin{bmatrix} \overline{\pi_x}^l & 0 \\ 0 & \beta^l \end{bmatrix}$$

2) For $x \in \mathcal{X}$, with $v_x = 0, u_x \neq 0$,

a) $\pi_x = \beta$:

$$(S^{\sim x}RD)^l = \begin{bmatrix} \overline{\pi_x}^l & 0 \\ -l u_x \overline{\pi_x}^{l-1} & \overline{\pi_x}^l \end{bmatrix}$$

b) $\pi_x \neq \beta$:

$$(S^{\sim x}RD)^l = \begin{bmatrix} 1 & 0 \\ \frac{u_x}{\pi_x - \beta} & 1 \end{bmatrix} \begin{bmatrix} \overline{\pi_x}^l & 0 \\ 0 & \beta^l \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -\frac{u_x}{\pi_x - \beta} & 1 \end{bmatrix} \quad (30)$$

3) For $x \in \mathcal{X}$, with $v_x \neq 0, u_x = 0$,

a) $\pi_x = \beta$:

$$(S^{\sim x}RD)^l = \begin{bmatrix} \overline{\pi_x}^l & -l \pi_x v_x \overline{\pi_x}^{l-1} \\ 0 & \overline{\pi_x}^l \end{bmatrix}$$

b) $\pi_x \neq \beta$:

$$(S^{\sim x}RD)^l = \begin{bmatrix} 1 & -\frac{\beta \pi_x v_x}{\pi_x - \beta} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \overline{\pi_x}^l & 0 \\ 0 & \beta^l \end{bmatrix} \begin{bmatrix} 1 & \frac{\beta \pi_x v_x}{\pi_x - \beta} \\ 0 & 1 \end{bmatrix} \quad (31)$$

4) For $x \in \mathcal{X}$, with $v_x \neq 0, u_x \neq 0$,

$$(S^{\sim x}RD)^l = V^{\sim x} \begin{bmatrix} (\lambda_1^{\sim x})^l & 0 \\ 0 & (\lambda_2^{\sim x})^l \end{bmatrix} (V^{\sim x})^{-1} \quad (32)$$

$$\text{where } V^{\sim x} = \begin{bmatrix} 1 & 1 \\ -\frac{\Delta_x - s_x}{2(1-\beta)\pi_x v_x} & \frac{\Delta_x + s_x}{2(1-\beta)\pi_x v_x} \end{bmatrix} \\ (V^{\sim x})^{-1} = \begin{bmatrix} \frac{\Delta_x + s_x}{2\Delta_x} & -\frac{(1-\beta)\pi_x v_x}{\Delta_x} \\ \frac{\Delta_x - s_x}{2\Delta_x} & \frac{(1-\beta)\pi_x v_x}{\Delta_x} \end{bmatrix}$$

with $s_x = 1 - \pi_x - (1 - \beta)(1 - v_x u_x)$, $\lambda_1^{\sim x}, \lambda_2^{\sim x}$ and Δ_x are as defined in Lemma 6.

For $x \in \mathcal{X}$ that fall in any of the first three categories in Lemma 13 (which makes $P_{xx} = \pi_x$), substituting the

corresponding form of $(S^{\sim x}RD)^l$ in (26), (27) and simplifying we get $\Pr(E_{\sim m}^x | X_m \neq x) = (1 - \pi_x)^{n-1}$ for $m = 1$ to n and $\Pr(E_{2:m}^x | X_1 = x) = (1 - \pi_x)^{m-1}$, for all $2 \leq m \leq n$. Using this in (29) and (28), we get $\Pr(E_{\sim m}^x | X_m = x) = (1 - \pi_x)^{n-1}$, for $m = 1$ to n . This completes the proof of the first part of Lemma 10.

For $x \in \mathcal{X}$ with $v_x \neq 0$ and $u_x \neq 0$, substituting the expression for $(S^{\sim x}RD)^l$ from (32) in (27) and simplifying we get

$$P(E_{2:m}^x | X_1 = x) = (\lambda_1^{\sim x})^{m-1} \left(\frac{s_x + \Delta_x}{2\Delta_x} - \frac{(1-\beta)v_x u_x}{\Delta_x} \right) \\ + (\lambda_2^{\sim x})^{m-1} \left(\frac{\Delta_x - s_x}{2\Delta_x} + \frac{(1-\beta)v_x u_x}{\Delta_x} \right) \quad (33)$$

for $m \geq 2$.

Using (33) with $m = n$ along with (28) completes the proof of (24). Using (33) in (29) and simplifying, we get (25). Lastly, substituting $(S^{\sim x}RD)^l$ from (32) in (26) and simplifying, we get (23) and the proof is complete.

APPENDIX

A. Proof of Lemma 11

To prove Lemma 11, we first note that

$$\Pr(E_{\sim m}^x | X_m \neq x) = \frac{\Pr(F_x(X_{\sim m}^n) = 0, X_m \neq x)}{\Pr(X_m \neq x)} \\ = \frac{\Pr(F_x(X^n) = 0)}{\Pr(X_m \neq x)}$$

We can write $\Pr(F_x(X^n) = 0)$ as

$$\Pr(F_x(X^n) = 0) = \sum_{y \in \mathcal{X}, y \neq x} \pi_y \Pr(E_{\sim 1}^x | X_1 = y).$$

Since $\Pr(E_{\sim 1}^x | X_1 = y)$ is the probability that the Markov chain does not visit the state x in the steps X_2 through X_n given that it starts in state y , $\Pr(E_{\sim 1}^x | X_1 = y)$ equals the row sum of $(P^{\sim x})^{n-1}$ taken along the row corresponding to the state y . Therefore

$$\Pr(F_x(X^n) = 0) = \sum_{y \in \mathcal{X}, y \neq x} \pi_y \sum_{z \in \mathcal{X}} [(P^{\sim x})^{n-1}]_{y,z} \\ \stackrel{(a)}{=} \sum_{y \in \mathcal{X}, y \neq x} \pi_y \sum_{z \in \mathcal{X}} [(RDS^{\sim x})^{n-1}]_{y,z} \\ = \sum_{y \in \mathcal{X}, y \neq x} \pi_y \sum_{z \in \mathcal{X}} \left[RD(S^{\sim x}RD)^{n-2} S^{\sim x} \right]_{y,z}$$

where we use $P^{\sim x} = RDS^{\sim x}$ to get (a). Noting that $(S^{\sim x}RD)^{n-2}, S^{\sim x}$ are matrices of dimensions 2×2 and $2 \times K$

respectively and $[1 \quad (1 - \beta)v_y]$ is the row of the matrix RD corresponding to the state y , we have

$$\begin{aligned} & \Pr(F_x(X^n) = 0) \\ & \stackrel{(b)}{=} \sum_{y \in \mathcal{X}, y \neq x} \pi_y [1 \quad (1 - \beta)v_y] (S^{\sim x} RD)^{n-2} \begin{bmatrix} 1 - \pi_x \\ -u_x \end{bmatrix} \\ & \stackrel{(c)}{=} [1 - \pi_x \quad -(1 - \beta)\pi_x v_x] (S^{\sim x} RD)^{n-2} \begin{bmatrix} 1 - \pi_x \\ -u_x \end{bmatrix} \end{aligned}$$

where we use $\pi^T \mathbf{1}_{K \times 1} = 1$, $u^T \mathbf{1}_{K \times 1} = 0$ (from $SR = I$) to sum up the rows of $S^{\sim x}$ to get (b) and use $\pi^T \mathbf{1}_{K \times 1} = 1$, $u^T v = 0$ (from $SR = I$) to get (c).

Using $\Pr(E_{\sim m}^x | X_m \neq x) = \frac{1}{1 - \pi_x} \Pr(F_x(X^n) = 0)$ for $m = 1, 2, \dots, n$ completes the proof.

B. Proof of Lemma 12

1) Using a method similar to the proof of Lemma 11, we get

$$\begin{aligned} & \Pr(E_{2:m}^x | X_1 = x) \\ & = \sum_{z \in \mathcal{X}} [(P^{\sim x})^{m-1}]_{x,z} = \sum_{z \in \mathcal{X}} [(RD S^{\sim x})^{m-1}]_{x,z} \\ & = [1 \quad (1 - \beta)v_x] \left(S^{\sim x} RD \right)^{m-2} \begin{bmatrix} 1 - \pi_x \\ -u_x \end{bmatrix} \end{aligned}$$

2) We begin with the result of [15, Lemma 8] which states that for a stationary Markov chain X^n ,

$$\Pr(X_j = x, E_{1:j-1}^x) = \Pr(X_1 = x, E_{2:j}^x), \text{ for } 2 \leq j \leq n \quad (34)$$

Since

$$\Pr(E_{1:n-1}^x | X_n = x) = \frac{\Pr(E_{1:n-1}^x, X_n = x)}{\Pr(X_n = x)}$$

using (34) and the stationarity of X^n , we get

$$\Pr(E_{1:n-1}^x | X_n = x) = \Pr(E_{2:n}^x | X_1 = x)$$

We now factorise $\Pr(E_{\sim m}^x | X_m = x)$ (for $m = 2, \dots, n - 1$) as shown below:

$$\begin{aligned} & \Pr(E_{\sim m}^x | X_m = x) \\ & \stackrel{(a)}{=} \Pr(E_{1:m-1}^x | X_m = x) \Pr(E_{m+1:n}^x | X_m = x) \\ & \stackrel{(b)}{=} \Pr(E_{2:m}^x | X_1 = x) \Pr(E_{m+1:n}^x | X_m = x) \\ & \stackrel{(c)}{=} \Pr(E_{2:m}^x | X_1 = x) \Pr(E_{2:n-m+1}^x | X_1 = x) \end{aligned}$$

where we use Markov property to get (a), use (28) with $n = m$ to get (b) and finally use the stationarity of X^n to get (c). This completes proof of (29).

C. Proof of Lemma 13

Since $SR = I$, we have $S^{\sim x} R = \begin{bmatrix} \overline{\pi}_x & -\pi_x v_x \\ -u_x & 1 - v_x u_x \end{bmatrix}$ and

$$S^{\sim x} RD = \begin{bmatrix} \overline{\pi}_x & -\overline{\beta} \pi_x v_x \\ -u_x & \overline{\beta}(1 - v_x u_x) \end{bmatrix} \quad (35)$$

1) We substitute $v_x = u_x = 0$ in (35) and raise the power on both sides to l .

2) a) Substituting $v_x = 0$, $\beta = \pi_x$ in (35), we get $S^{\sim x} RD = \begin{bmatrix} \overline{\pi}_x & 0 \\ -u_x & \overline{\pi}_x \end{bmatrix}$. Using induction on power l , we get $(S^{\sim x} RD)^l = \begin{bmatrix} \overline{\pi}_x^l & 0 \\ -l u_x \overline{\pi}_x^{l-1} & \overline{\pi}_x^l \end{bmatrix}$.

b) Substituting $v_x = 0$ in (35), we get $S^{\sim x} RD = \begin{bmatrix} \overline{\pi}_x & 0 \\ -u_x & \overline{\beta} \end{bmatrix}$. We observe that $\overline{\pi}_x, \overline{\beta}$ are the eigenvalues of $S^{\sim x} RD$ with $\begin{bmatrix} 1 \\ \frac{u_x}{\overline{\pi}_x - \overline{\beta}} \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ as their respective right eigenvectors resulting in the diagonalised form in (30).

3) a) Substituting $u_x = 0$, $\beta = \pi_x$ in (35), we get $S^{\sim x} RD = \begin{bmatrix} \overline{\pi}_x & -\overline{\pi}_x \pi_x v_x \\ 0 & \overline{\pi}_x \end{bmatrix}$. Using induction on power l , we get $(S^{\sim x} RD)^l = \begin{bmatrix} \overline{\pi}_x^l & -l \pi_x v_x \overline{\pi}_x^{l-1} \\ 0 & \overline{\pi}_x^l \end{bmatrix}$.

b) Substituting $u_x = 0$ in (35), we get $S^{\sim x} RD = \begin{bmatrix} \overline{\pi}_x & -\overline{\beta} \pi_x v_x \\ 0 & \overline{\beta} \end{bmatrix}$. We observe that $\overline{\pi}_x, \overline{\beta}$ are the eigenvalues of $S^{\sim x} RD$ with $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} -\frac{\overline{\beta} \pi_x v_x}{\overline{\pi}_x - \overline{\beta}} \\ 1 \end{bmatrix}$ as their respective right eigenvectors resulting in the diagonalised form in (31).

4) Solving $\det(S^{\sim x} RD - \lambda I) = 0$, we get $\lambda_i^{\sim x} = 0.5(\overline{\pi}_x + \overline{\beta}(1 - v_x u_x) + (-1)^{i+1} \Delta_x)$, $i = 1, 2$, as the eigenvalues of $S^{\sim x} RD$ with $\begin{bmatrix} 1 \\ -\frac{\Delta_x - s_x}{2(1 - \beta)\pi_x v_x} \end{bmatrix}$, $\begin{bmatrix} 1 \\ \frac{\Delta_x + s_x}{2(1 - \beta)\pi_x v_x} \end{bmatrix}$ as their respective right eigenvectors, where $\Delta_x^2 = (\beta - \pi_x + \overline{\beta} v_x u_x)^2 + 4\overline{\beta} \pi_x v_x u_x$, $s_x = 1 - \pi_x - (1 - \beta)(1 - v_x u_x)$, resulting in the diagonalised form in (32).

This completes the proof of Lemma 13.

REFERENCES

- [1] R. G. Gallager, "Finite state Markov chains," in *Discrete Stochastic Processes*. Springer, 1996, pp. 103–147.
- [2] I. J. Good, "The population frequencies of species and the estimation of population parameters," *Biometrika*, vol. 40, no. 3/4, pp. 237–264, 1953.
- [3] D. A. McAllester and R. E. Schapire, "On the convergence rate of Good-Turing estimators," in *Proceedings of the Thirteenth Annual Conference on Computational Learning Theory*, 2000, pp. 1–6.
- [4] S. F. Chen and J. Goodman, "An empirical study of smoothing techniques for language modeling," in *Proceedings of the 34th Annual Meeting on Association for Computational Linguistics*, ser. ACL '96, 1996, pp. 310–318.
- [5] D. McAllester and L. Ortiz, "Concentration inequalities for the missing mass and for histogram rule error," *J. Mach. Learn. Res.*, vol. 4, pp. 895–911, Dec. 2003.
- [6] D. Berend and A. Kontorovich, "On the concentration of the missing mass," *Electron. Commun. Probab.*, vol. 18, p. 7 pp., 2013.
- [7] A. Ben-Hamou, S. Boucheron, and M. I. Ohannessian, "Concentration inequalities in the infinite urn scheme for occupancy counts and the missing mass, with applications," *Bernoulli*, vol. 23, no. 1, pp. 249–287, 02 2017.

- [8] N. Rajaraman, A. Thangaraj, and A. T. Suresh, "Minimax risk for missing mass estimation," in *2017 IEEE International Symposium on Information Theory (ISIT)*, June 2017, pp. 3025–3029.
- [9] J. Acharya, Y. Bao, Y. Kang, and Z. Sun, "Improved bounds for minimax risk of estimating missing mass," in *2018 IEEE International Symposium on Information Theory (ISIT)*, June 2018, pp. 326–330.
- [10] P. Chandra and A. Thangaraj, "Concentration and tail bounds for missing mass," in *2019 IEEE International Symposium on Information Theory (ISIT)*, July 2019, pp. 1862–1866.
- [11] M. Asadi, R. P. Torghabeh, and N. P. Santhanam, "Stationary and transition probabilities in slow mixing, long memory Markov processes," *IEEE Transactions on Information Theory*, vol. 60, no. 9, pp. 5682–5701, 2014.
- [12] M. Falahatgar, A. Orlitsky, V. Pichapati, and A. T. Suresh, "Learning Markov distributions: Does estimation trump compression?" in *2016 IEEE International Symposium on Information Theory (ISIT)*, July 2016, pp. 2689–2693.
- [13] Y. Hao, A. Orlitsky, and V. Pichapati, "On learning Markov chains," in *Proceedings of the 32nd International Conference on Neural Information Processing Systems*, ser. NIPS'18. Red Hook, NY, USA: Curran Associates Inc., 2018, p. 646–655.
- [14] G. Wolfer and A. Kontorovich, "Minimax learning of ergodic Markov chains," in *Proceedings of the 30th International Conference on Algorithmic Learning Theory*, 2019, pp. 904–930.
- [15] P. Chandra, A. Thangaraj, and N. Rajaraman, "Missing mass of Markov chains," in *2020 IEEE International Symposium on Information Theory (ISIT)*, 2020, pp. 1207–1212.
- [16] M. Skórski, "Missing mass in Markov chains," *arXiv preprint, arXiv:2001.03603*, 2020.
- [17] S. U. Pillai, T. Suel, and S. Cha, "The perron-frobenius theorem: some of its applications," *IEEE Signal Processing Magazine*, vol. 22, no. 2, pp. 62–75, 2005.
- [18] L. A. Kontorovich and K. Ramanan, "Concentration inequalities for dependent random variables via the martingale method," *The Annals of Probability*, vol. 36, no. 6, p. 2126–2158, Nov 2008.
- [19] A. Naor, S. Rao, and O. Regev, "Concentration of Markov chains with bounded moments," *Ann. Inst. H. Poincaré Probab. Statist.*, vol. 56, no. 3, pp. 2270–2280, 08 2020.
- [20] P. Chandra, A. Thangaraj, and N. Rajaraman, "Missing mass of rank-2 Markov chains," *arXiv preprint, arXiv:2102.01938v1*, 2021. [Online]. Available: <https://arxiv.org/pdf/2102.01938.pdf>